



## LJMU Research Online

**Kirk, C, Clark, DR, Langan-Evans, C and Morton, JP**

**The physical demands of mixed martial arts: A narrative review using the ARMSS model to provide a hierarchy of evidence.**

<http://researchonline.ljmu.ac.uk/id/eprint/13541/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Kirk, C, Clark, DR, Langan-Evans, C and Morton, JP (2020) The physical demands of mixed martial arts: A narrative review using the ARMSS model to provide a hierarchy of evidence. Journal of Sports Sciences. ISSN 1466-447X**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

1 **The physical demands of mixed martial arts: A narrative review using the ARMSS model to provide a**  
2 **hierarchy of evidence**

3 Christopher Kirk<sup>1,2</sup>, David R Clark<sup>2</sup>, Carl Langan-Evans<sup>2</sup>, James P Morton<sup>2</sup>

4 <sup>1</sup>University of Derby, College of Life and Natural Sciences, Kedleston Rd., Derby, United Kingdom, DE22 1GB

5 <sup>2</sup>Liverpool John Moores University, Research Institute for Sport and Exercise Sciences, Tom Reilly Building,  
6 Liverpool, United Kingdom, L3 3AF

7 Corresponding author: Christopher Kirk Email: [C.Kirk@Derby.ac.uk](mailto:C.Kirk@Derby.ac.uk)

8 CK Orcid: 0000-0002-6207-027X DC Orcid: 0000-0002-6661-6137

9 DC Email: [D.R.Clark@LJMU.ac.uk](mailto:D.R.Clark@LJMU.ac.uk)

10 CL Email: [C.Langanevans@LJMU.ac.uk](mailto:C.Langanevans@LJMU.ac.uk)

11 JM Email: [J.P.Morton@LJMU.ac.uk](mailto:J.P.Morton@LJMU.ac.uk)

12

13 **Abstract**

14 The physical and physiological demands of mixed martial arts (MMA) training and competition is not yet well  
15 quantified. The Applied Research Model for the Sport Sciences (ARMSS) provides a framework through which  
16 to conduct sport science, determining pertinent questions to apply and test research findings in real world settings.  
17 The aim of this review was to evaluate MMA research within the context of ARMSS to critically analyse our  
18 understanding of the physical and physiological requirements of MMA training and competition. Research  
19 databases were searched, with 70 peer-reviewed articles being discussed in relation to the specific stage of the  
20 ARMSS in which their results best fit. MMA research was found to be mostly foundational and descriptive in  
21 nature and has generally not developed along systematic lines. The internal and external physiological loads and  
22 responses to training and competition have not been adequately identified. Therefore, it is not currently possible  
23 to state which variables are key predictors of success, or how coaches can optimally manipulate these variables.  
24 We propose that MMA research could be refocused to be conducted within ARMSS. Specifically, stage 2 studies  
25 describing the physical, physiological and technical demands of MMA training and competition, and stage 3  
26 studies determining the physiological predictors of competition performance should be initially prioritised.

27

28

29 **Key Words**

30 Combat sports; ARMSS; performance

31 **Word Count = 5,663**

32

33

34

## 35 **1 – Introduction**

36 The aim of scientific research is to provide a structured understanding of a particular field to develop solutions to  
37 encountered problems (Niiniluoto 1993). Though many disciplines have histories of achieving this within a formal  
38 framework, sports science has not generally proceeded in this manner. This has been suggested to inhibit the  
39 translation of science to practice (Bishop 2008). Research lacking robustness and/or applicability tends to be most  
40 evident, though not exclusively, in young sports where the knowledge base is under-developed. One such sport  
41 where this is the case is mixed martial arts (MMA). MMA is a body-mass regulated combat sport in which  
42 competitors engage each other using kicks, punches, elbows and knees along with grappling manoeuvres to  
43 overcome their opponent. Bouts are 3 x 5 minute rounds (5 x 5 minute rounds for championship bouts or bouts  
44 deemed to be of significance) for professionals and 3 x 3 minute rounds for amateurs (NJSAC 2002, James et al.  
45 2016, Kirk 2018a). A contest is won or lost when: i) the referee is given reason to believe either opponent is no  
46 longer intelligently defending themselves against strikes (deemed a technical knockout (TKO)), or is no longer  
47 physically able to intelligently defend themselves against strikes (deemed a knockout (KO)); ii) either opponent  
48 ‘submits’ by indicating they are no longer willing to continue the contest due to joint manipulation, choking or  
49 injury, or if a submission hold has rendered them unconscious or physically debilitated; iii) either opponent or  
50 their coaches indicates they are unable or unwilling to continue the contest, iv) if an appointed medical doctor  
51 determines that either participant is no longer fit to continue safely, or, v) all scheduled rounds of the bout are  
52 completed and judges decide the winner based on pre-set criteria (ABC 2018).

53

54 Though MMA has been compared to the ancient Greek sport of pankration (Buse 2006, Seidenberg 2011) it  
55 developed from Brazilian vale tudo (no holds barred) contests of the early to mid-20<sup>th</sup> Century (Gracie and Danaher  
56 2003). This format was imported to the USA in the 1990s by several organisations, most notably the Ultimate  
57 Fighting Championship (UFC) (Gracie and Danaher 2003, Souza-Junior et al. 2015). MMA became a codified  
58 sport with the 2002 adoption of the ‘unified rules’ (NJSAC 2002). Due to its recent establishment, research into  
59 this population of athletes is nascent, as discussed previously (Lenetsky and Harris 2012, Del Vecchio and  
60 Franchini 2013, Andreato and Branco 2016, Kirk 2018a). Presently, there are 4,000+ male and female  
61 professional competitors worldwide (Fightmatrix 2020), with the International MMA Federation (IMMAF)  
62 consisting of 85+ national governing bodies across five continents (IMMAF 2019). MMA has rapidly increased  
63 in popularity and participation, as highlighted by the 2016 sale of UFC for \$4 billion, just 16 years after being  
64 bought for \$2 million (Rovell and Okamoto 2016). As an indication of the use of sports science in MMA, UFC  
65 established performance institutes designed to provide training and support for its contracted athletes (UFC 2019).  
66 However, there is evidence that MMA coaches do not view scientific research as important to their role (Bujak et  
67 al. 2013). If this perspective is to be altered the work produced by researchers must be demonstrated to be of real-  
68 world value to the end user, allowing a symbiotic relationship to occur.

69

70 To improve the quality of data produced by sport science, Bishop (2008) proposed the Applied Research Model  
71 for the Sport Sciences (ARMSS). Consisting of eight interacting stages (Table 1) that guide the logical progression  
72 of research by building on the findings of each stage, ARMSS aims to ensure results are based firmly on  
73 cumulative knowledge. Hence, research is continuously tested and verified before being applied to improve  
74 performance or health. This approach may ensure sport science is more focused and integrative, which should

75 result in applicable findings avoiding stand-alone studies with little or no practitioner or researcher impact. Whilst  
76 other suggested models focus on improving research to practice transfer (Drust and Green 2013, Verhagen et al.  
77 2014, Eisenmann 2017), ARMSS provides a more detailed structure of the cumulative steps of the evidence  
78 gathering process. This structure allows a nuanced assessment of the current research base. Therefore, the aim of  
79 the present paper is to review our current understanding of the physiological and physical demands of MMA  
80 training and competition, as presented within a hierarchy of evidence provided by the ARMSS framework. This  
81 work is important as it provides coaches, applied practitioners and researchers an important and valuable  
82 summation of the performance related research completed during developmental stages of the sport. The findings  
83 provided here lay the foundations of future work to be performed to better understand the requirements and effects  
84 of MMA.

## 85 **2 – Categorising MMA research within the ARMSS model**

86 Studies included in this review were acquired between the months of March 2017 and November 2019 inclusively  
87 by searching peer-reviewed articles written in English (or providing an abstract in English) in the following  
88 databases and repositories: PubMed, OVID, Google Scholar, Researchgate.net, LibraryPlus and Academia.edu.  
89 Keywords used in searches included: MMA, mixed martial arts, training load, time motion analysis, S&C, strength  
90 and conditioning, performance, performance analysis, anthropometry, age, physiology. Finally, reference lists and  
91 bibliographies were searched for additional sources. The only inclusion criteria was that articles had to discuss  
92 performance and/or preparation related aspects of MMA from a physical or physiological standpoint. Studies  
93 discussing psychological, sociological or other non-physiological aspects were excluded.

94 The discussion that forms the body of this review is split into sections corresponding with each stage of ARMSS.  
95 Articles were assigned to ARMSS stages based on how closely their key findings fit the stage explanations  
96 provided by Bishop (2008). Articles which could fit into more than one stage have been placed according to the  
97 main conclusion provided by the article author(s). Review and placement of each article was initially undertaken  
98 by the lead author, then reviewed and agreed by the remaining authors. To ensure a coherent narrative, articles  
99 within each ARMSS stage were grouped into the following themes and discussed in turn:

- 100 • Characteristics of the MMA athlete (any article that describes the physical and/or physiological capabilities  
101 and/or capacities of MMA participants);
- 102 • Characteristics and/or physical/physiological effects of MMA training (any article that describes training  
103 methods used and/or the effects of training methods used by/suggested for MMA participants, inclusive of  
104 intervention and non-intervention based research);
- 105 • Characteristics of MMA performance (any article that describes the internal/external load or the  
106 technical/tactical aspects of official competition or simulated bouts, and the technical, physical or  
107 physiological predictors/characteristics of successful MMA performance);

108 These themes were chosen from common areas of sports science research and areas of concern around combat  
109 sport athlete preparation (Ruiz 2017). In total, 70 unique articles were found and included. Table 1 details the  
110 distribution of these articles in each ARMSS stage. 8 were retrieved that are related to MMA performance but do  
111 not fit in any ARMSS stage. Each article included in the review is summarised in Table 2. Whilst injuries,

112 concussions and body-mass regulation are recognised as key facets of MMA, they are outside the scope of this  
 113 review. The incidences, rates and severity of each of these areas are, however, in need of more focussed research  
 114 in order to support and protect the participants involved (Seidenberg 2011, Lockwood et al. 2018, Barley et al.  
 115 2019).

116

**Table 1 – Descriptors of each ARMSS(Bishop 2008) stage and the number of MMA related articles that have been reviewed and placed within each stage.**

Purpose	Stage	n = MMA related studies included in each stage
Description	<b>1 Defining the problem</b> Identification of types of real-world problems and issues that coaches and athletes face. Discussions with practitioners, reviews and meta-analyses.	<b>11</b>
	<b>2 Descriptive research (hypothesis development)</b> Studies that describe what is currently occurring in the field, cross-sectional studies and methodological studies.	<b>25</b>
	<b>3 Predictors of performance</b> Studies to identify which factors are most likely to affect performance. Should include replication studies.	<b>21</b>
Experimentation	<b>4 Experimental testing of predictors</b> Studies that determine which of the previously identified predictors are likely causal (controlled, randomised, double blind studies).	<b>5</b>
	<b>5 Determinants of key performance predictors</b> Studies to determine which intervention(s) most effect the previously identified causal predictors.	<b>0</b>
	<b>6 Efficacy studies</b> Studies to determine whether the previously highlighted interventions cause substantial positive or negative effects on performance in controlled conditions.	<b>0</b>
Implementation	<b>7 Barriers to uptake</b> Which real world factors may hinder the application of the research findings from the previous stages?	<b>0</b>
	<b>8 Implementation studies</b> Studies to determine how effective at improving performance the previously identified interventions are when used in a real-world, uncontrolled setting.	<b>0</b>

*Nb. 8 additional performance related publications do not fit into any stage of the ARMSS model.*

117

### 118 3 – ARMSS Stage 1: Defining the Problem

119 Stage 1 research is currently lacking in MMA, despite calls for work of this kind to be conducted (Lenetsky and  
 120 Harris 2012, Andreato and Branco 2016). Due to this lack of information, several authors have used data from  
 121 related combat sports to provide suggestions about the requirements of MMA competition and preparation  
 122 strategies (James et al. 2016). A lack of understanding about the potential positive and negative effects of training  
 123 methods amongst practitioners has been highlighted (Amtmann 2004, 2010). Whilst the details of each of these  
 124 studies are discussed in other sections of this review, this lack of foundational research presents an immediate  
 125 deviation from the ARMSS. As the aim of stage 1 is to define the nature of the event being studied and the  
 126 problems requiring attention(Bishop 2008), the absence of this understanding in MMA may have negative  
 127 implications for the applicability of much of the data discussed in this review.

128 **4 – ARMSS Stage 2: Descriptive Research**

129 **4.1 – Characteristics of the MMA Athlete**

130 It has been suggested via needs analyses of other combat sports that MMA participants would likely require a  
131 range of physiological capabilities, including high lower body power and anaerobic capacity (James et al. 2016,  
132 Lonergan et al. 2018), but these conclusions are based on limited evidence. Though most studies report stature  
133 and mass of participants, lean tissue and fat distribution data is limited to the studies reported in Table 3. Each of  
134 these used a cross-divisional cohort without delineating between mass divisions, which may mask important  
135 distinctions in performance and requirements. A greater but still limited number of papers outlined physical  
136 capabilities of MMA participants (example data can be viewed in Table 4). These data tend to suggest that MMA  
137 competitors have relatively low  $\dot{V}O_2\text{max}$ , hand grip strength and muscular endurance, alongside moderate relative  
138 strength, vertical jump and linear sprint speed, especially when compared to other combat and non-combat sports  
139 (Guidetti et al. 2002, Bosquet et al. 2007, Argus et al. 2012, Lockie et al. 2018). A review of anthropometry and  
140 physical capabilities demonstrated that MMA competitors tend to have low body fat, high hip and lower torso  
141 flexibility and muscular endurance, but moderate cardiovascular capacity (Spanias et al. 2019). The key finding,  
142 however, was that most studies use different testing protocols and methods, making conclusions about the  
143 population as a whole difficult.

144 Regarding laboratory measurements, electromyography (EMG) found evidence of a double activation peak of the  
145 latissimus dorsi, erector spinae, gluteals and rectus femoris muscles during punches and kicks onto a heavy bag  
146 (McGill et al. 2010). Though providing an interesting discussion of relative muscle activation during MMA related  
147 techniques, the importance or otherwise of these measurements to performance is not alluded to. Finally, the UFC  
148 Performance Institute published suggested normative data (not peer reviewed) for assessing an MMA athlete's  
149 performance in several physiological components, including reactive strength index (RSI), peak power output,  
150  $\dot{V}O_2\text{max}$  and rate of force development (RFD) (Institute 2018). This provides a useful resource for practitioners to  
151 understand the physical abilities of high-level competitors. A limitation of this document from a research  
152 viewpoint is that it does not provide specific means or methods for each of these indices, reducing the opportunity  
153 for comparison, benchmarking or replication.

154

155

156

157

158

159

160

**Table 3 – Example anthropometrical and body tissue distribution data of MMA competitors**

Study	n	Stature (cm)	Body mass (kg)	Body fat (kg)	Body fat (%)	BMI (kg·m <sup>-2</sup> )	Muscle mass (%)	Fat free mass (kg)
(Marinho et al. 2016)	8	177 ± 5	82.1 ± 9.6	11.8 ± 6.2	13.4 ± 5.6*	26 ± 3.3	69.6 ± 4.6	
(Schick et al. 2010)	11	175 ± 5	77.4 ± 11.4		11.7 ± 4*			
(Marinho et al. 2011)	13	176 ± 5	82.1 ± 10.9		11.9 ± 5.1*			
(de Oliveira et al. 2015)	18	172 ± 5	78.3 ± 6.9		15 ± 7.3**			
(LaRocca et al. 2019)	40					24.6 ± 3.3		
(Kasper et al. 2018)	1		80.2	11.7	15***			63.5
(Alm and Yu 2013)	5	180.4 ± 9	80.8 ± 11.1	9.9 ± 2	12.25 ± .5***		83.2 ± 0.8	

*Nb. BMI = body mass index; \* = estimated using skinfolds; \*\* = estimated using air displacement plethysmography; \*\*\*estimated using dual energy x-ray absorptiometry*

162

163

164

165

166

167

168

169

170

171

172

173

174

175

**Table 4 – Example physiological and physical assessment data for MMA competitors**

Study	n	$\dot{V}O_2\text{max}$ (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )	Hand grip strength (kg)	1RM squat (kg)	1RM relative bench press (kg·kg <sup>-1</sup> )	CMJ (cm)	Relative CMJ (cm·kg <sup>-1</sup> )	SJ peak power (W·kg <sup>-1</sup> )	SJ (cm)	Standing broad jump (cm)	10m sprint (s)	Upper body peak power (W·kg <sup>-1</sup> )
(Schick et al. 2010)	11	55.5 ± 7.3	45.8 ± 6.2		1.2 ± 0.1							
(Marinho et al. 2011)	13			73 ± 15						219 ± 0.25		
(Lovell et al. 2013)	1	Pre = 55, Post = 56.8 training program										Pre = 8.9, Post = 9.7
(Alm and Yu 2013)	5	Pre = 50 ± 6.5, Post = 48.5 ± 5.7 training program					Pre = 0.63 ± 0.4, Post = 0.59 ± 0.1 training program		Pre = 40.3 ± 3.8, Post = 36.2 ± 3.6			
(de Oliveira et al. 2015)	18	44.2 ± 6.7	R = 46 ± 9, L = 45 ± 8.5									
(Marinho et al. 2016)	8			69 ± 6	1 ± 0.2					219 ± 0.31		
(Kostiakiadis et al. 2018)	10	Pre = 41.5 ± 11.1*, Post = 46.2 ± 10.3* training program		Pre = 140.7 ± 22.5, Post = 167.1 ± 25.6		Pre = 33.1 ± 5.2, Post = 35.1 ± 3.8			Pre = 31 ± 4.6, Post = 31.6 ± 3.9			Pre = 1.95 ± 0.06, Post = 1.88 ± .05
(James, Beckman, et al. 2017)	29				HL = 1.21 ± 0.18, LL = 1.07 ± 0.2			HL = 44.45 ± 7.54, LL = 38.47 ± 6.74				

*Nb. 1RM = one repetition maximum; CMJ = countermovement jump; SJ = squat jump; \* = estimated; R = right hand; L = left hand; HL = pro/semi pro competitors with ≥50% wins; LL = pro/amateur competitors with ≤50% wins*

176

177

#### 178 4.2 – Assessment of MMA Training Loads

179 In terms of training load assessment, there are large gaps in our knowledge. Two attempts have used participant  
180 questionnaires to understand how competitors prepare for bouts (Amtmann 2004, 2010). Both instances found  
181 that training in MMA has wide variations in frequency and type (1 – 7 strength and conditioning sessions, 3 -12  
182 MMA sessions per week). Additionally, 83% of participants used resistance training, only 26% incorporated  
183 Olympic lifting derivatives and 53% used neck strengthening movements. It was concluded that most participants  
184 had poor strength and conditioning knowledge or a lack of support in this area. The only study reporting actual  
185 training load data is a case study of an elite participant (n=1) preparing for a professional world championship  
186 tournament (Jukic et al. 2017). This competitor experienced mean weekly sessional rating of perceived exertion  
187 (sRPE) of 3,653 AU during a 9-week training camp (range = 1,345 - 6,035 AU). Unfortunately, these data are  
188 from preparations for a 2006 defunct competition format (one-night, multi-bout tournament), so its utility for  
189 current performers is limited. A second more recent training case study does provide details as to the distances,

190 times and prescribed intensities of training for a male professional competitor and the measurable effects of this  
191 program(Tota et al. 2014). However, this article does not describe the specific content or training modes of the  
192 sessions themselves, nor how they were incorporated alongside the participant’s technical training. Torso-  
193 mounted accelerometry has been used to measure the external load of standing and grounded MMA strikes and  
194 takedowns. Punches, right sided kicks and double leg takedowns elicited greater external loads in training  
195 compared to the same techniques in simulated competition (Kirk et al. 2015a). How these loads affect the athlete’s  
196 overall readiness to perform or how this can be manipulated by the coaches is, however, unknown. The paucity  
197 of data regarding the current training practices and loads of MMA participants makes it extremely difficult to  
198 form conclusions regarding the efficacy of the methods used. This gap in our understanding of the sport’s  
199 preparatory practices is a key limitation of MMA research.

### 200 **4.3 – Physiological adaptations to MMA Training**

201 Alm and Yu (2013) observed a reduction in squat jump (SJ) and countermovement jump (CMJ) relative to body  
202 mass after 12 months of MMA training(Alm and Yu 2013). These results are limited, however, as it was  
203 conducted with 5 participants being tested in the first instance and only 4 being tested in the second instance due  
204 to injury and as such does not provide robust data for practitioners to work from. Furthermore, individual training  
205 content and volume were not reported. A case study (n = 1) reporting responses to an 8-week training period  
206 demonstrated increased  $\dot{V}O_2\text{max}$  on a cycling ergometer and increased relative upper body peak power(Lovell et  
207 al. 2013). This study did report training content, allowing more nuanced considerations to be made by researchers  
208 and trainers. This facet of detail was further developed by Kostikiadis et al. (2018) demonstrating significant  
209 improvements in physiological variables in this population through employment of a specific 4-week strength and  
210 conditioning program. Further details of this study are discussed in section 6.1.

211

### 212 **4.4 – Physical and physiological demands of MMA competition**

#### 213 **4.4.1 – External Load of MMA competition**

214 External load of MMA competition has been measured directly and by proxy in official competitions and  
215 simulated bouts. Torso mounted accelerometry measured accumulated player load ( $PL_{ACC} = 224.32 \pm 26.59$  AU)  
216 and accumulated player load per minute ( $PL_{ACC} \cdot \text{min}^{-1} = 14.91 \pm 1.78$  AU) in simulated bouts of 3 x 5 minute  
217 rounds (Kirk et al. 2015b). This demonstrates that MMA external loads may be comparable to those of an  
218 Australian rules football match (Mooney et al. 2013), but greater than those experienced by elite netball  
219 players(Cormack et al. 2014) and defenders in association football (Domene 2013). Given accelerometry displays  
220 high reliability in the measurement of external load of MMA (Hurst et al. 2014), this method shows promise for  
221 use within training and competition and should be explored further by researchers and practitioners alike.

222 The physical demands of MMA competition has also been measured by proxy through time motion analyses  
223 (TMA). Del Vecchio et al. (2011) determined that within a 5 minute round, there was 2-3 minutes of activity,  
224 which generally reduced as the round and the bout commenced. Work-to-rest ratio (W:R) was 9:1 – 6:1, when  
225 taking the 1 minute break between rounds into account. When excluding the 1 minute break, W:R was 1:2 – 1:4,

226 with the briefest category being high intensity standing ( $9 \pm 8$  s) and the longest being low intensity groundwork  
227 ( $21 \pm 19$  s). ‘Rest’ periods included low intensity combat phases (stable positions requiring little effort or  
228 movement) and pauses (Del Vecchio et al. 2011). The same research group used their TMA method to show  
229 female competitors spend longer engaged in low intensity standing movements, less time in high intensity standing  
230 movements and less groundwork time in total than males. The same study also reported the ratio of high to low  
231 intensity (H:L) to be 2:1, indicating that MMA participants utilise higher intensity movements more often than  
232 low intensity movements (Del Vecchio et al. 2015).

233 Further analyses demonstrated W:R differences between divisions, with the greatest differences in H:L being  
234 observed in the standing phases of each bout. For example, flyweight (FIW) reported values of 1:15 whereas  
235 heavyweights (HW) reported values of 1:7 (Miarka et al. 2015). In terms of total effort time, HW displayed shorter  
236 time than all other divisions in the first ( $212.4 \pm 101.5$  s) and third ( $246.3 \pm 89.1$  s) rounds, whilst bantamweight  
237 (BW) was shortest in the second round ( $132.8 \pm 90.9$  s). MMA competitors also increase their time spent in  
238 standing preparatory actions (non-striking or grappling) as the bout progresses, a potential effect of fatigue  
239 (Antonietto et al. 2019). In simulated competition, MMA participants had a W:R = 1:1.01, with most active time  
240 being spent in the clinch (aggressor =  $100.33 \pm 65.87$  s; defensive =  $95.5 \pm 58.48$  s) and striking low activity ( $95.5$   
241  $\pm 12.63$  s). Conversely, participants in this study only spent  $15.33 \pm 11.22$  s in striking high activity (Kirk et al.  
242 2015b). When comparing results of each of these papers, it becomes clear that actions in MMA are intermittent,  
243 with differing requirements for participants of different divisions and sex (Del Vecchio et al. 2015, Antonietto et  
244 al. 2019). Equally, relative intensities of competition and simulated competition can be very different, an  
245 observation that has also been made in other combat sports (Andreato et al. 2016, Slimani et al. 2017).

246 Importantly, there are no studies analysing the physical demands of amateur bouts, with each of the discussed  
247 studies using professional or semi-professional participants and most using data from high level international  
248 bouts. Because of this, it is unknown whether there are any differences in bout characteristics between this level  
249 and the developmental levels of the sport. Such knowledge would likely be important to the successful  
250 development of training frameworks. One key issue with much of the external load data is the subjective manner  
251 in which ‘rest’ and ‘work’ have been identified. For example, the load of isometric movements associated with  
252 grappling has not been quantified, so determining whether these are genuinely rest or work is largely a judgement-  
253 based decision. This needs to be considered when reviewing TMA data in MMA.

#### 254 **4.4.2– Internal Load of MMA competition**

255 Data regarding internal load is presented in Table 5. Such data demonstrate that MMA seems to induce  
256 physiological responses within the same ranges as other contact and non-contact sports (Klappinska et al. 2001,  
257 Lippi et al. 2004, Paccotti et al. 2005, Nunes et al. 2012). These responses appear to be similar regardless of  
258 whether it is official or simulated competition, with the exception of cortisol. The specific movements, techniques  
259 and tactical approaches that bring about these responses have not yet been identified, whilst the medium term (24  
260 – 48h post) metabolic and muscle damage responses have also not been fully determined. This would be a key  
261 area for identifying the true internal load of MMA. Also highlighted is the need for researchers to use a consistent  
262 protocol of sampling and measurement in terms of biomarkers and sampling points.

**Table 5 – Summary of internal load measures of MMA technical training, official competition and simulated competition**

Study	Mode	Variable measured	Pre	Post	24 hours post	48 hours post
(Amtmann et al. 2008)	Official competition	Lactate (mmol·L <sup>-1</sup> )		10.2 – 20.7		
	Simulated competition	RPE (AU)		13 – 19		
		Lactate (mmol·L <sup>-1</sup> )		13.3 – 18		
(Kirk et al. 2015b)	Simulated competition	RPE (AU)		15-18		
		Lactate (mmol·L <sup>-1</sup> )		13 – 19.7		
		RPE (au)		17 – 19		
(Kirk et al. 2015b)	Simulated competition	Lactate (mmol·L <sup>-1</sup> )	2.7 ± 1.46	9.25 ± 2.96 ↑		
(Coswig, Ramos, et al. 2016)	Official competition	Urea (mg·ml <sup>-1</sup> )	44.15 ± 8.93			36.31 ± 7.85 ↓
		Uric acid (mg·ml <sup>-1</sup> )	5.17 ± 0.91			4.62 ± 0.78 ↓
(Coswig, Fukuda, et al. 2016)	Official competition	Creatine kinase (U·L)	221 (median)	237 (median) ↑		
	Simulated competition	Lactate (mmol·L <sup>-1</sup> )	4 (median)	16.9 (median) ↑		
		Creatine kinase (U·L)	225 (median)	297 (median) ↑		
(Souza et al. 2017)	Official competition	Lactate (mmol·L <sup>-1</sup> )	3.8 (median)	16.8 (median) ↑		
		Cortisol (nmol·L <sup>-1</sup> )	594 ± 144.9	876 ± 107.84 ↑	436 ± 126.26 ↓↓	
		Glucose (mg·dl <sup>-1</sup> )	4.81 ± 0.92	10 ± 2.55 ↑	3.82 ± 0.28 ↓↓	
		Testosterone (nmol·L <sup>-1</sup> )	13.03 ± 3.9	11.92 ± 1.79 ↓	17.75 ± 2 ↑↑	
		Creatine kinase (U·L)	433.89 ± 215.95	491.81 ± 219.17 ↑	1,412.69 ± 758.63 ↑↑	
(Ghoul et al. 2017)	Simulated competition	Cortisol (ng·dl <sup>-1</sup> )	88.3 ± 19.6	131.2 ± 56.4 ↑	79.3 ± 23.2 ↓↓	
		Testosterone (ng·dl <sup>-1</sup> )	3.5 ± 0.3	4.6 ± 0.3 ↑	3.4 ± 0.3 =↓	
		Uric acid (μmol·L <sup>-1</sup> )	308 ± 53	342 ± 54 ↑	344 ± 48 ↑=	
		Lactate dehydrogenase (U·L)	169 ± 70	230 ± 116 ↑	226 ± 75 ↑=	

Nb. ↑/↓/= on Post and 48 hours post measurements indicate either increase/decrease/no practical change from Pre measurements; first ↑/↓/= on 24 hours post measurements indicate either increase/decrease/no practical change from Pre measurements; second ↑/↓/= on 24 hours post measurements indicate either increase/decrease/no practical change from Post measurements; RPE = rating of perceived exertion.

264

## 265 5 – ARMSS Stage 3: Predictors of Performance

### 266 5.1 – Physical and Physiological Characteristics

#### 267 5.1.1 - Physiology

268 Analysis of muscular force production in relation to success in MMA was found in one study (James, Beckman,  
 269 et al. 2017). High level semi-professional competitors (HL) were capable of superior lower body neuromuscular  
 270 force production than low level semi-professional and amateur competitors (LL). This was based on significant  
 271 differences in relative back squat one repetition maximum (1RM) (HL = 1.84 ± 0.23; LL = 1.56 ± 0.24 kg·kg<sup>-1</sup>),  
 272 squat jump (SJ) peak power (HL = 44.45 ± 7.54; LL = 38.47 ± 6.74 W·kg<sup>-1</sup>) and SJ peak velocity (HL = 3.06 ±  
 273 0.33; LL = 2.81 ± 0.33 m·s<sup>-1</sup>) between both groups (for other results see Tables 1 and 3). Professional and amateur  
 274 competitors may also be distinguished with at least a small or better effect size in 10 and 20 m sprint times, repeat

275 sprint ability and Yo-Yo Intermittent Recovery results (James et al. 2018). The authors of each of these studies  
276 suggest that training for this population should maximise lower body force capacity and impulse, as well as sprint  
277 and repeat sprint capability. Given the limited number of studies completed and the largely subjective distinction  
278 between high and low-level competitors, more work is required to unequivocally link these variables to success  
279 in MMA.

### 280 **5.1.2 – Anthropometry**

281 Anthropometry has an anecdotal relationship to success in combat sports, with a competitor's 'height and reach'  
282 being reported as a matter of course. This assumption has been tested in MMA, mostly by the lead author of this  
283 review. Stature and/or arm span differences have small to moderate correlations with technique use differences  
284 between winners and losers of professional bouts. These relationships were found in heavyweight (HW),  
285 welterweight (WW) and featherweight (FW) with less than half of the variance in technique use being explained  
286 by anthropometrical differences between opponents (Kirk 2018a). Anthropometry has also been studied in  
287 relation to divisional rank and attainment of world title bouts, with only a negligible effect being found in two  
288 divisions. These were in flyweight (FIW), where shorter competitors were ranked highest and in women's  
289 strawweight (WSW) where shorter competitors were ranked in the middle of the division. Arm span was found  
290 to have no influence at all(Kirk 2016a). Though taller participants were more likely to win individual bouts via  
291 strikes than submission or decision, this was found to be anecdotal using Bayesian inference ( $BF_{10}$  – see note  
292 marked \* in Table 6). In comparison, if the bout loser was taller than their opponent, the evidence for them losing  
293 due to strikes was classed as very strong. Equally, competitors with longer arm spans were more likely to lose  
294 due to strikes rather than decision, and at WW they were moderately more likely to lose due to submission(Kirk  
295 2016b).

296 Each of these papers also discussed the so called 'ape ratio', a measurement representing the scale of a person's  
297 arm span in comparison to their stature (S:W), showing that despite this variable influencing success in other  
298 sports, it has none in MMA(Kirk 2016a, 2016b, 2018a). This conclusion has been questioned as it was found that  
299 S:W can predict MMA participant's winning percentage. This result, however, showed that less than 1% of  
300 variance in winning percentage can be explained by this anthropometric measurement(Monson et al. 2018). The  
301 anecdotal link between success and anthropometry appears to be vastly over-estimated in MMA, potentially due  
302 to the use of grappling to negate any potential dis/advantages stature or arm span provides in the striking phases  
303 of competition.

### 304 **5.1.3 – Aging**

305 Competitive bout winners ( $30 \pm 4$  years) were likely to be younger than bouts losers ( $31 \pm 4$  years) when examined  
306 at as cross divisional cohort (Kirk 2016b). This age gap was different for each division with the greatest found at  
307 lightweight (LW) and bantamweight (BW) (~ 3 years difference in each). A potentially more important effect of  
308 age was found in the methods of winning or losing. If the winning participant was older they most likely won due  
309 to a decision and in cases where older participants lost, it was most likely due to strikes. This could provide  
310 evidence that participants become more susceptible to strikes as they age, possibly due to a reduction in speed and  
311 reaction times (Korhonen et al. 2009, Hunter et al. 2016). This is supported by the odds ratios of suffering a KO

312 or a TKO increasing with age(Hutchison et al. 2014). This could result in a cyclical pattern of performance and  
313 health deterioration, as physiological decrements have also been related to the effects of repeated concussive  
314 strikes in training and competition (De Beaumont et al. 2013, Richmond and Rogol 2014).

315 The effects of age related adaptations have been revealed over a 10 year period as reduced strike accuracy and  
316 rate, fewer successful takedowns and lower markers of intensity (Santos et al. 2018). Another potential  
317 consideration could be that as participants age they become more attuned to tactical requirements of the sport and  
318 are better able to achieve decision wins instead of taking increased risks to attain a stoppage victory. An analysis  
319 of the top 100 male competitors in each division indicated that a combination of these two factors could result in  
320 'peak performance windows'. Older participants were generally ranked higher, with the highest 20% of a cross-  
321 divisional cohort being older than the other 80%. Equally, this study also found that as divisional mass limit  
322 increases, so too does the average age of the top 100 competitors. This pattern starts at BW ( $29 \pm 4$  years) and  
323 progressively increases up to HW ( $33 \pm 5$  years) (Kirk 2018b). We conclude here that performances in lighter  
324 divisions could be more affected by physiological decrements of aging on speed, force production and reaction  
325 time than in heavier divisions, leading to lighter competitors dropping out of the top 100 earlier. However, without  
326 currently knowing the respective contributions of these variables to performance in MMA, this conclusion remains  
327 an early supposition.

## 328 **5.2 – Technical Actions of Competition**

329 Table 6 provides a brief summary of distinguishing technical factors of successful MMA performance. These  
330 results suggest that being competent in all phases of combat is a prerequisite of success, and that skills needed are  
331 highly dependent on situation and opponent. Though the studies listed elucidate technical requirements of MMA,  
332 they do not distinguish between divisions, which have understandably large differences in anthropometrical  
333 measurements (Kirk 2016a). It would be natural to assume that these differences would lead to distinguishable  
334 movement and skill characteristics. Such technical differences have been reported (Miarka, Brito, Dal Bello, et  
335 al. 2017), showing that each division displays differences in the time spent engaged in striking, clinch work,  
336 keeping distance and attacking the body, as well as diverse frequencies of technique use in each area.

337 In a similar vein, there is evidence of differences between each division in terms of which techniques contribute  
338 most to a winning performance (Kirk 2018a). The summary in Table 7 shows wide variations in technical  
339 characteristics separating winners and losers. WW and LW participants arguably require a more complete skill  
340 set than HW and light heavyweight (LHW), who appear to display little influence of grappling. These results  
341 strongly indicate that future studies should be conducted on a division-by-division basis, not treating MMA  
342 participants as a homogenous group. There could also be a need to perform replications of some of the studies  
343 named in this section for the same purpose, thus allowing a clearer picture with more applicable results to emerge.  
344 Equally, there may need to be more research of bouts that end due to stoppage(Miarka et al. 2018) and bouts that  
345 go beyond the standard three rounds(Miarka et al. 2019). Though a reasonable amount of work has been done to  
346 better understand technical and tactical performance in MMA, these criteria seem to be evolving as the sport  
347 matures (Santos et al. 2018), often in response to adaptations to rule changes in the sport (Fernandes et al. 2018).  
348 With this in mind, it is important that this work continues to monitor these changes to better inform coaching  
349 decisions.

**Table 6 – Summary of key determining factors of successful MMA performance**

<b>Study</b>	<b>Mode</b>	<b>Sample</b>	<b>Key determining factor</b>	<b>Other info</b>
(Kirk et al. 2015b)	Simulated competition	6 male participants	Successful takedowns achieved	
(Miarka, Brito, and Amtmann 2017)	Official competition	584 professional bouts	Head strikes landed at distance, guard passes, successful takedowns	
(Miarka, Del Vecchio, Camey, et al. 2016)	Official competition	215 professional bouts	Total strikes, submission attempts, ground positional changes	
(Miarka et al. 2018)	Official competition	678 professional bouts	Less time engaged in low intensity standing actions at any point in the bout, and more time in high intensity groundwork in the first 2 rounds	Bouts ending in judge's decision excluded
(dal Bello et al. 2019)	Official competition	304 professional rounds	Type and frequency of grappling techniques used (takedowns, sweeps and submission attempts) appear to have a direct influence on whether the bout ends in a unanimous/split decision, submission, KO or TKO	
(Miarka, Coswig, et al. 2016)	Official competition	174 professional rounds	Bouts that ended via stoppage (TKO, KO or submission) consist of more time spent striking and grappling; bouts that ended in a decision characterised by more time spent performing standing strikes only	Female bouts only
(Miarka et al. 2019)	Official competition	779 professional rounds from bouts ended in rounds 3 – 5	Activity profiles are different dependent on whether the bout ended in the 3 <sup>rd</sup> , 4 <sup>th</sup> or 5 <sup>th</sup> rounds	
(James, Robertson, et al. 2017)	Official competition	234 professional bouts	Strike and takedown accuracy concomitant with ground strikes landed	

*Nb. KO = knockout; TKO = technical knockout*

**Table 7 – Summary of variables found to distinguish between winners and losers of professional MMA bouts according to Bayes Factor (BF<sub>10</sub>)\* hypothesis test thresholds greater than moderate (Kirk 2018a)**

Distinguishing Performance Variables by BF <sub>10</sub> Likelihood Threshold						
Division	Decisive	Decisive to Very Strong	Very Strong	Very Strong to Strong	Strong	Strong to Moderate
Heavyweight	Significant strikes landed; significant strikes attempted					Significant ground strikes landed; knockdowns
Light Heavyweight	Strikes landed					Significant ground strikes landed
Middleweight	Significant strikes landed; strikes landed; strikes attempted; significant strikes attempted; significant ground strikes landed.		Significant distance strikes landed		Guard passes	
Welterweight	Significant ground strikes landed; strikes landed; significant strikes landed; guard passes; strikes attempted; knockdowns; successful takedowns; distance knockdowns	Takedowns attempted			Significant strikes attempted; significant distance strikes landed.	
Lightweight	Significant strikes landed; significant ground strikes landed; strikes landed; strikes attempted; significant strikes attempted; submissions attempted; knockdowns; distance knockdowns; guard passes.		Significant distance strikes landed	Successful takedowns; significant clinch strikes landed		Significant distance strikes attempted.
Featherweight	Knockdowns; Distance knockdowns		Submissions attempted			Significant strikes landed; significant ground strikes landed.
Bantamweight	Significant strikes landed; significant ground strikes landed; Significant strikes attempted.		Significant clinch strikes attempted.	Strikes attempted	Knockdowns	Strikes landed; significant distance strikes landed; submissions attempted
Flyweight	Significant strikes attempted	Successful takedowns	Submissions attempted	Strikes attempted	Passes	Significant ground strikes landed
Women's Bantamweight	No variables greater than moderate					
Women's Strawweight	Strikes landed	Strikes attempted	Significant strikes landed		Guard passes; significant strikes attempted	Significant distance strikes landed

*Nb. Variables for each division shown in order of BF<sub>10</sub> likelihood magnitude. \*For an explanation of the use and application of Bayesian inference and BF<sub>10</sub>, please see Wagenmakers et al. (2018)*

## 353 **6 – ARMSS Stage 4 Experimental Testing of Key Performance Predictors**

### 354 **6.1 – Physiological adaptations associated with MMA Training**

355 A small number of studies provide experimental data assessing effectiveness of training methods applied in MMA.  
356 A recent study suggested a 6-week strength and power training program likely caused increases in cross punch,  
357 roundhouse kick and lower limb force production amongst MMA and kickboxing participants (Vecchio et al.  
358 2019). However, the authors do not clarify how the MMA and kickboxing participants are distributed between  
359 the intervention and control groups. More importantly, there is no evidence that strike force in itself is a key  
360 determinant of success. The content of the 6-week training program, however, is provided in sufficient detail to  
361 allow replications. The physiological effects of a 4-week MMA specific strength and conditioning programme on  
362 a group of MMA competitors (specific training group (STG)) were compared to a control group of MMA  
363 competitors (regular training group (RTG)) completing their normal technical and non-MMA specific strength  
364 and conditioning sessions (Kostikiadis et al. 2018). Use of the specific strength and conditioning programme  
365 alongside technical training elicited greater increases in comparison to technical training alone: estimated  $\dot{V}O_2\text{max}$   
366 (STG =  $13.3 \pm 14.5\%$ , RTG =  $-0.1 \pm 6.9\%$ ); back squat 1RM (STG =  $19.5 \pm 10.4\%$ , RTG =  $2.05 \pm 8.45\%$ ); CMJ  
367 (STG =  $7.4 \pm 4.4\%$ , RTG =  $1.2 \pm 1\%$ ); 10m sprint (STG =  $-3.7 \pm 1.4\%$ , RTG =  $-0.4 \pm 1.1\%$ ).

368 Further evidence of the acute effect of strength and conditioning was observed between two groups of participants  
369 split by their preferred combat style (group A = striking based; group B = grappling based) (Chernozub et al.  
370 2018). Each group was prescribed a 3-month power training programme based on their combat style to be  
371 completed alongside their technical training. Upon completion of the programme both groups then completed the  
372 opposite style's training programme for the next 3 months. Bench press 1RM data was collected at months 0, 3  
373 and 6. Both groups significantly increased strength at each sampling point regardless of the style of training  
374 completed. Though between group analysis was absent, it does appear that both groups responded better to training  
375 designed for striking style than grappling style. Due to the programmes being completed in series, this cannot be  
376 confirmed. Equally, there is no comparison to a control group, and the measurement chosen does not necessarily  
377 reflect MMA performance.

378 Though each of these studies provide experimental data of training interventions, there is little indication that the  
379 variables being improved directly influence competitive performance, showing a clear deviation from ARMSS.  
380 These data do, however, provide a starting point for understanding which types of training may bring about  
381 positive adaptations.

### 382 **7 – ARMSS Stages 5-8**

383 As can be seen in Tables 1 and 2, there are currently no studies that can be placed in any of the four later stages  
384 of ARMSS. This is likely due to the conclusions of sections 5 and 6, that a causal link between predictors of  
385 performance and success has not yet been established. This is likely to be important if successful stage 5 (and  
386 beyond) research is to be completed. Conversely, while this may initially appear to be a failing of MMA research,  
387 it may also demonstrate that researchers are working to build the foundation of our collective understanding  
388 sufficiently before attempting to undertake potentially flawed latter stage research. From this viewpoint, MMA  
389 research could be said to be following the process of ARMSS, in that it appears to have been recognised that stage

390 5-8 work cannot currently be performed. It does also mean, however, that we are currently not in a position to  
391 confidently state what the determinants of successful performance are, or how we can optimally prepare an athlete  
392 to achieve them. Most of the topics discussed throughout this review have been categorised to sit within the  
393 foundational, descriptive stages of the ARMSS.

#### 394 **8 – Performance studies that are not applicable to the ARMSS model**

395 A number of retrieved articles do not fit into any stage of ARMSS. In response to the pressing needs of  
396 practitioners, several authors have presented suggestions for strength and conditioning models based on data from  
397 other combat sports and assumptions of the requirements of MMA (La Bounty et al. 2011, Tack 2013, Earnshaw  
398 2015). Other protocols are specific towards injury prevention (James 2014), female competitors (Schick et al.  
399 2012) or metabolic conditioning (Mikeska 2014, Harvey 2018). In summary, the recommendations are a  
400 combination of some or all of the following: strength and power training to improve torso stability and lower body  
401 force production; mobility and aerobic conditioning via the use of high intensity interval training (HIIT). Though  
402 each presents recommendations based on established strength and conditioning theory, none provide any data  
403 assessing the effectiveness of these models. Only one (Harvey 2018) provides details as to how their  
404 recommendations could be utilised alongside technical training to optimise preparation. The author in question  
405 presents an MMA specific HIIT protocol to integrate into technical training sessions, though admits that  
406 effectiveness of this protocol requires testing to demonstrate validity. This recommendation can be applied to each  
407 of the articles in this section: future research needs to address the absence of scientific evidence to support the  
408 suggested protocols for MMA.

#### 409 **9 – Conclusions and Future Research Directions**

410 When considering the body of research as presented in the current review, it is clear that most is descriptive in  
411 nature and does not strictly follow ARMSS. Not all factors studied in stage 3 (predictors) are linked to stage 2  
412 (descriptive) findings. Equally, none of the variables experimentally tested in stage 4 have been found to predict  
413 performance in stage 3. There has been little attempt at understanding the sport's current training practices.  
414 Research describing the movements and techniques used in professional competition is fairly well developed, but  
415 this does not seem to have transferred into experimental testing or training interventions. Training studies focus  
416 on physiological variables that have not been adequately linked to success. There is also a lack of distinction  
417 between sexes, with only two papers discussing females exclusively (Schick et al. 2012, Miarka, Coswig, et al.  
418 2016). Given the inherently different responses to training and performance between sexes (Pitchers and Elliot-  
419 Sale 2019), this would be an important area for research focus. These conclusions place the current review in stage  
420 1 of ARMSS, defining as it does the current gaps in MMA research.

421 Based on these findings, the authors have developed a series of questions within ARMSS to reflect the areas we  
422 feel are most in need of immediate attention (Table 8). These questions may begin the process of our collective  
423 foundational knowledge of MMA becoming better developed. This in turn would allow researchers and  
424 practitioners to access later stages of ARMSS. This would require active collaboration between researchers and  
425 practitioners with a coherent approach to issues raised, in turn reducing negative effects of disparate results caused  
426 by isolated questions. It should be noted that ARMSS is designed to improve the quality of information being  
427 delivered to practitioners only. Other models have been developed to improve the vital process of knowledge

428 management and transfer (Drust and Green 2013, Verhagen et al. 2014, Eisenmann 2017). Given the aims of the  
 429 current review, however, exploration of these models within MMA are beyond the scope of this article. Discussion  
 430 of this kind may also be premature, given the lack of foundational knowledge within MMA science demonstrated  
 431 here. Using ARMSS in MMA research at this point will allow researchers to fully develop early stage  
 432 understanding of the sport in a consistent and effective manner. Once this has been achieved then impactful  
 433 knowledge transfer can occur.

434 The current review provides an important resource for applied practitioners, coaches and researchers in outlining  
 435 our current understanding of the demands of MMA. We have collated all extant data (Table 2) for a clearer  
 436 appreciation of the differing requirements of success between divisions, the short term internal responses to  
 437 competition, the potential physiological responses to MMA training, and to highlight our lack of data regarding  
 438 the training practices used in the sport. This knowledge should be used by coaches as part of an ongoing evaluation  
 439 of training methods in relation to their athlete’s needs. Specifically, this could be avoiding reliance on unsupported  
 440 predictive variables such as anthropometry; incorporation of targeted strength and conditioning programs  
 441 alongside technical training; and devising tactical approaches to competition based on division specific  
 442 performance characteristics. Researchers should use this article as a reference text to guide future research plans  
 443 and questions to address the highlighted gaps in our knowledge. The application of ARMSS presented here also  
 444 demonstrates how MMA research can be structured to ensure knowledge generation is efficient and purposeful,  
 445 enabling future work to be conducted on a strong foundation. With MMA becoming more organised globally at  
 446 both professional and amateur levels, now is the time to influence real change and progress within the scientific  
 447 basis of the sport for everyone involved.

448

**Table 8 – Suggested questions in the first three stages of the ARMSS model to inform future MMA research**

Stage	
2	<p data-bbox="252 1330 443 1352"><b>Descriptive research</b></p> <ul style="list-style-type: none"> <li data-bbox="300 1357 1283 1379">• What are the physical, physiological and technical demands of professional and amateur MMA competition?</li> </ul> <p data-bbox="347 1406 1474 1456">Such research studies should use a range of methods to quantify external and internal loads in both male and female participants as well as be specific to each weight division.</p> <ul style="list-style-type: none"> <li data-bbox="300 1482 1401 1505">• What are the habitual training practices undertaken by professional and amateur MMA athletes preparing for competition?</li> </ul> <p data-bbox="347 1532 1458 1603">Such research should describe the range of training modalities, activities and specific combat techniques across a time-scale that athletes typically complete when preparing for competition. Similar to the above, such studies should use a range of methods to quantify external and internal loads in both male and female participants as well as be specific to each weight division.</p> <ul style="list-style-type: none"> <li data-bbox="300 1630 916 1653">• What are the physical and physiological effects of MMA training?</li> </ul> <p data-bbox="347 1680 1474 1749">Using a testing battery of reliable and valid performance tests, such research should examine the physiological adaptations associated with participating in an MMA training programme that is intended to prepare male and female professional and amateur athletes for competition.</p>
3	<p data-bbox="252 1776 494 1798"><b>Predictors of performance</b></p> <ul style="list-style-type: none"> <li data-bbox="300 1803 1449 1825">• Which physical, physiological and technical characteristics provide the strongest predictor of successful performance in MMA?</li> </ul> <p data-bbox="347 1852 1465 1921">On the basis of understanding the physical and physiological demands of training and competition, such studies should subsequently a range of factors that are associated with successful performance so as to provide a platform to then experimentally test each predictor.</p>

449

450 **10 - References**

- 451 ABC, 2018. Unified Rules of Mixed Martial Arts.  
452
- 453 Adam, M., Pujso, R., Kuzmicki, S., Szymanski, M., and Tabakov, S., 2015. MMA fighters' technical-tactical  
454 preparation-fight analysis: a case study. *Journal of Combat Sports and Martial Arts*, 6 (1), 35–41.  
455
- 456 Alm, P. and Yu, J., 2013. Physiological characters in mixed martial arts. *American Journal of Sports Science*, 1  
457 (2), 12–17.  
458
- 459 Amtmann, J., 2004. Self-reported training methods of mixed martial artists at a regional reality fighting event.  
460 *Journal of Strength and Conditioning Research*, 18 (1), 194–6.  
461
- 462 Amtmann, J., 2010. Training volume and methods of athletes competing at a mixed martial arts event.  
463 *Intermountain Journal of Sciences*, 16 (1-3), 51–54.  
464
- 465 Amtmann, J., 2011. Anticipating and training for the worst-case metabolic scenario: a comment on Del Vecchio,  
466 Hirata, and Franchini. *Perceptual and Motor Skills*, 114 (1), 123–124.  
467
- 468 Amtmann, J., Amtmann, K.A., and Spath, W.K., 2008. Lactate and rate of perceived exertion responses of  
469 athletes training for and competing in a mixed martial arts event. *The Journal of Strength & Conditioning*  
470 *Research*, 22 (2), 645–647.  
471
- 472 Andreato, L. and Branco, B.H.M., 2016. Different sports, but the same physical and physiological profiles?  
473 *Sports Medicine*, 46 (12), 1963–1965.  
474
- 475 Andreato, L.V., Follmer, B., Celidonio, C.L., and da Silva Honorato, A., 2016. Brazilian Jiu-Jitsu combat  
476 among different categories: time-motion and physiology. A systematic review. *Strength and Conditioning*  
477 *Journal*, 38 (6), 44–54.  
478
- 479 Antonietto, N.R., Bello, F., Carrenho, A.Q., de Carvalho Berbert, P., Brito, C.J., Amtmann, J., and Miarka, B.,  
480 2019. Suggestions for Professional Mixed Martial Arts Training With Pacing Strategy and Technical-  
481 Tactical Actions by Rounds. *Journal of Strength and Conditioning Research*.  
482
- 483 Antonio, J., Leaf, A., Carson, C., Ellerbroek, A., Silver, T., Peacock, C.A., Bommarito, P., Knafo, S., and  
484 Tartar, J., 2018. Bone Mineral Density in Competitive Athletes. *Journal of Exercise and Nutrition*, 1, 2.  
485
- 486 Argus, C.K., Gill, N.D., Keogh, J.W., McGuigan, M.R., and Hopkins, W.G., 2012. Effects of two contrast  
487 training programs on jump performance in rugby union players during a competition phase. *International*  
488 *Journal of Sports Physiology and Performance*, 7 (1), 68–75.  
489
- 490 De Azevedo, A.P., Guerra, M.A., Caldas, L.C., and Guimarães-Ferreira, L., 2019. Acute caffeine ingestion did  
491 not enhance punch performance in professional mixed-martial arts athletes. *Nutrients*, 11 (6), 1422.  
492
- 493 Baker, J. and Schorer, J., 2013. The southpaw advantage? Lateral preference in mixed martial arts. *PloS one*, 8  
494 (11), e79793.  
495
- 496 Barley, O.R., Chapman, D.W., and Abbiss, C.R., 2019. The current state of weight-cutting in combat sports.  
497 *Sports*, 7 (5), 123.  
498
- 499 De Beaumont, L., Tremblay, S., Henry, L.C., Poirier, J., Lassonde, M., and Théoret, H., 2013. Motor system  
500 alterations in retired former athletes: the role of aging and concussion history. *BMC Neurology*, 13 (1), 109.  
501
- 502 Dal Bello, F., Brito, C.J., Amtmann, J., and Miarka, B., 2019. Ending MMA combat, specific Grappling  
503 techniques according to the type of the outcome. *Journal of Human Kinetics*, 1 (1).  
504
- 505 Bishop, D., 2008. An applied research model for the sport sciences. *Sports Medicine*, 38 (3), 253–263.  
506
- 507 Bodden, J.G., Needham, R.A., and Chockalingam, N., 2015. The effect of an intervention program on functional  
508 movement screen test scores in mixed martial arts athletes. *Journal of Strength and Conditioning Research*,

509 29 (1), 219–225.  
510  
511 Bosquet, L., Duchene, A., Dupont, G., Leger, L., and Carter, H., 2007. VO<sub>2</sub> Kinetics during Supramaximal  
512 Exercise: Relationship with Oxygen Deficit and 800-m Running Performance. *International Journal of*  
513 *Sports Medicine*, 28 (06), 518–524.  
514  
515 La Bounty, P., Campbell, B.I., Galvan, E., Cooke, M., and Antonio, J., 2011. Strength and conditioning  
516 considerations for mixed martial arts. *Strength & Conditioning Journal*, 33 (1), 56–67.  
517  
518 Brito, C.J., Mozer, R.L., Bem, E.M.A. de, Carvalho, P.H.B. de, Queiroz, A.C.C., Bello, F. dal, Barreto, L.B.M.,  
519 and Miarka, B., 2018. Exploratory study on illegal pharmacologic agents in mixed martial arts performance.  
520 *Revista Brasileira de Cineantropometria & Desempenho Humano*, 20 (3), 269–279.  
521  
522 Bujak, Z., Gierczuk, D., and Litwiniuk, S., 2013. Professional activities of a coach of martial arts and combat  
523 sports. *Journal of Combat Sports and Martial Arts*, 4 (2), 191–195.  
524  
525 Buse, G.J., 2006. No holds barred sport fighting: a 10 year review of mixed martial arts competition. *British*  
526 *Journal of Sports Medicine*, 40 (2), 169–172.  
527  
528 Chernozub, A., Danylchenko, S., Imas, Y., Kochina, M., Natalia, I., Korobeynikov, G., Korobeynikova, L.,  
529 Potop, V., Cynarski, W.J., and Gorashchenko, A., 2019. Peculiarities of correcting load parameters in power  
530 training of mixed martial arts athletes. *Journal of Physical Education and Sport*, 19, 481–488.  
531  
532 Chernozub, A., Korobeynikov, G., Mytskan, B., Korobeinikova, L., and Cynarski, W.J., 2018. Modelling Mixed  
533 Martial Arts Power Training Needs Depending on the Predominance of the Strike or Wrestling Fighting  
534 Style. *Ido Movement for Culture. Journal of Martial Arts Anthropology*, 18 (3), 28–36.  
535  
536 Cormack, S.J., Smith, R.L., Mooney, M.M., Young, W.B., and O’Brien, B.J., 2014. Accelerometer load as a  
537 measure of activity profile in different standards of netball match play. *International Journal of Sports*  
538 *Physiology and Performance*, 9 (2), 283–91.  
539  
540 Coswig, V., Ramos, S. de P., and Del Vecchio, F.B., 2016. Time-motion and biological responses in simulated  
541 mixed martial arts sparring matches. *Journal of Strength & Conditioning Research*, 30 (8), 2156–2163.  
542  
543 Coswig, V.S., Fukuda, D.H., de Paula Ramos, S., and Del Vecchio, F.B., 2016. Biochemical differences  
544 between official and simulated mixed martial arts (MMA) matches. *Asian Journal of Sports Medicine*, 7 (2).  
545  
546 Domene, Á., 2013. Evaluation of movement and physiological demands of full-back and center-back soccer  
547 players using global positioning systems. *Journal of Human Sport and Exercise*, 8 (4).  
548  
549 Drust, B. and Green, M., 2013. Science and football: evaluating the influence of science on performance.  
550 *Journal of Sports Sciences*, 31 (13), 1377–1382.  
551  
552 Earnshaw, A., 2015. Planning a mesocycle for a mixed martial artist. *Journal of Australian Strength and*  
553 *Conditioning*, 23 (7), 34–43.  
554  
555 Eisenmann, J., 2017. Translational gap between laboratory and playing field: New era to solve old problems in  
556 sports science. *Translational Journal of the American College of Sports Medicine*, 2 (8), 37–43.  
557  
558 Fernandes, J., Bello, F., Duarte, M., de Carvalho, P., Queiroz, A., Brito, J., and Miarka, B., 2018. Effect of rule  
559 changes on technical-tactical actions correlated with injury incidence in Professional Mixed Martial Arts.  
560 *Journal of Physical Education and Sport*, 18 (3).  
561  
562 Fightmatrix, 2020. Current MMA Rankings [online]. Available from: <http://www.fightmatrix.com/mma-ranks/>  
563 [Accessed 12–9 2019].  
564  
565 Ghoul, N., Tabben, M., Miarka, B., Tourny, C., Chamari, K., and Coquart, J., 2017. Mixed martial arts induces  
566 significant fatigue and muscle damage up to 24 hours post-combat. *Journal of Strength and Conditioning*  
567 *Research*.  
568

569 Gracie, R. and Danaher, J., 2003. *Mastering jujitsu*. Human Kinetics.  
570

571 Guidetti, L., Musulin, A., and Baldari, C., 2002. Physiological factors in middleweight boxing performance.  
572 *Journal of Sports Medicine and Physical Fitness*, 42 (3), 309–314.  
573

574 Harvey, B., 2018. Energy system development in elite mixed martial arts. *Journal of Australian Strength and*  
575 *Conditioning*, 26 (4), 75–89.  
576

577 Hunter, S.K., Pereira, H.M., and Keenan, K.G., 2016. The aging neuromuscular system and motor performance.  
578 *Journal of Applied Physiology*, 121 (4), 982–995.  
579

580 Hurst, H.T., Atkins, S., and Kirk, C., 2014. Reliability of a portable accelerometer for measuring workload  
581 during mixed martial arts. *Journal of Athletic Enhancement* , 5 (2).  
582

583 Hutchison, M.G., Lawrence, D.W., Cusimano, M.D., and Schweizer, T.A., 2014. Head trauma in mixed martial  
584 arts. *The American Journal of Sports Medicine*, 42 (6), 1352–1358.  
585

586 IMMAF, 2019. About IMMAF [online]. Available from: <https://immaf.org/about/> [Accessed 16–9 2019].  
587

588 Institute, U.F.C.P., 2018. A cross-sectional performance analysis and projection of the UFC athlete: Volume  
589 one.  
590

591 James, L.P., 2014. Injury prevention strategies for mixed martial arts. *Strength & Conditioning Journal*, 36 (5),  
592 88–95.  
593

594 James, L.P., Beckman, E.M., Kelly, V.G., and Haff, G.G., 2017. The neuromuscular qualities of higher-and  
595 lower-level mixed-martial-arts competitors. *International Journal of Sports Physiology and Performance*, 12  
596 (5), 612–620.  
597

598 James, L.P., Haff, G.G., Kelly, V.G., and Beckman, E.M., 2016. Towards a determination of the physiological  
599 characteristics distinguishing successful mixed martial arts athletes: a systematic review of combat sport  
600 literature. *Sports Medicine*, 46 (10), 1525–1551.  
601

602 James, L.P., Haff, G.G., Kelly, V.G., and Beckman, E.M., 2018. Physiological determinants of mixed martial  
603 arts performance and method of competition outcome. *International Journal of Sports Science & Coaching*,  
604 1747954118780303.  
605

606 James, L.P., Kelly, V.G., and Beckman, E.M., 2013. Periodization for mixed martial arts. *Strength &*  
607 *Conditioning Journal*, 35 (6), 34–45.  
608

609 James, L.P., Robertson, S., Haff, G.G., Beckman, E.M., and Kelly, V.G., 2017. Identifying the performance  
610 characteristics of a winning outcome in elite mixed martial arts competition. *Journal of Science and*  
611 *Medicine in Sport*, 20 (3), 296–301.  
612

613 Jay, M., 2013. Physiological characteristics and training practices of mixed martial artists: considerations for the  
614 strength and conditioning coach. *Journal of Australian Strength and Conditioning*, 21 (3), 91–103.  
615

616 Jukic, I., Milanovic, L., Hopovac, A., Filipovic, M., Jukic, N., and Krakan, I., 2017. Physical conditioning in  
617 mixed martial arts: From evidence to practical experience. In: Drid, Patrik, ed. *Science and Medicine in*  
618 *Combat Sports*. New York: Nova Science, 101.  
619

620 Kasper, A.M., Crighton, B., Langan-Evans, C., Riley, P., Sharma, A., Close, G.L., and Morton, J.P., 2018. Case  
621 Study: Extreme Weight Making Causes Relative Energy Deficiency, Dehydration and Acute Kidney Injury  
622 in a Male Mixed Martial Arts Athlete. *International Journal of Sport Nutrition and Exercise Metabolism*, 1–  
623 20.  
624

625 Kirk, C., 2016a. Does stature or wingspan length have a positive effect on competitor rankings or attainment of  
626 world title bouts in international and elite mixed martial arts? *Sport Science Review*, 25 (5-6), 334–349.  
627

- 628 Kirk, C., 2016b. The influence of age and anthropometric variables on winning and losing in professional mixed  
629 martial arts. *Facta Universitatis, Series: Physical Education and Sport*, 14 (2), 227–236.  
630
- 631 Kirk, C., 2018a. Does anthropometry influence technical factors in competitive mixed martial arts? *Human*  
632 *Movement*, 19 (2), 46–59.  
633
- 634 Kirk, C., 2018b. The relationship between age and divisional rank in professional mixed martial arts. *Facta*  
635 *Universitatis Series: Physical Education and Sport*, 16 (1), 73–84.  
636
- 637 Kirk, C., Hurst, H.T., and Atkins, S., 2015a. Comparison of the training loads of mixed martial arts techniques  
638 in isolated training and open sparring. *Journal of Combat Sports & Martial Arts*, 6 (1).  
639
- 640 Kirk, C., Hurst, H.T., and Atkins, S., 2015b. Measuring the workload of mixed martial arts using accelerometry,  
641 time motion analysis and lactate. *International Journal of Performance Analysis in Sport*, 15 (1), 359–370.  
642
- 643 Klapcinska, B., Iskra, J., Poprezecki, S., and Grzesiok, K., 2001. The effects of sprint (300 m) running on  
644 plasma lactate uricacid, creatin kinase, and lactate dehydrogenas in competitive hardlers and untrained men.  
645 *Journal of Sports Medicine and Physical Fitness*, 41 (3), 306–311.  
646
- 647 Korhonen, M.T., Mero, A.A., Alén, M., Sipilä, S., Häkkinen, K., Liikavainio, T., Viitasalo, J.T., Haverinen,  
648 M.T., and Suominen, H., 2009. Biomechanical and skeletal muscle determinants of maximum running speed  
649 with aging. *Medicine and Science in Sports and Exercise*, 41 (4), 844–56.  
650
- 651 Kostikiadis, I.N., Methenitis, S., Tsoukos, A., Veligekas, P., Terzis, G., and Bogdanis, G.C., 2018. The effect of  
652 short-term sport-specific strength and conditioning training on physical fitness of well-trained mixed martial  
653 arts athletes. *Journal of Sports Science and Medicine*, 17 (3), 348–358.  
654
- 655 LaRocca, D., Barns, S., Hicks, S.D., Brindle, A., Williams, J., Uhlig, R., Johnson, P., Neville, C., and  
656 Middleton, F.A., 2019. Comparison of serum and saliva miRNAs for identification and characterization of  
657 mTBI in adult mixed martial arts fighters. *PloS one*, 14 (1).  
658
- 659 Lenetsky, S. and Harris, N., 2012. The mixed martial arts athlete: a physiological profile. *Strength &*  
660 *Conditioning Journal*, 34 (1), 32–47.  
661
- 662 Lippi, G., Brocco, G., Franchini, M., Schena, F., and Guidi, G., 2004. Comparison of serum creatinine, uric  
663 acid, albumin and glucose in male professional endurance athletes compared with healthy controls. *Clinical*  
664 *Chemistry and Laboratory Medicine*, 42 (6), 644–647.  
665
- 666 Lockie, R.G., Dawes, J.J., and Jones, M.T., 2018. Relationships between Linear Speed and Lower-Body Power  
667 with Change-of-Direction Speed in National Collegiate Athletic Association Divisions I and II Women  
668 Soccer Athletes. *Sports*, 6 (2), 30.  
669
- 670 Lockwood, J., Frape, L., Lin, S., and Ackery, A., 2018. Traumatic brain injuries in mixed martial arts: A  
671 systematic review. *Trauma*, 20 (4), 245 – 254.  
672
- 673 Lonergan, B., Tallent, L., and Lazarczuk, D., 2018. Needs analysis of mixed martial arts. *Journal of Australian*  
674 *Strength and Conditioning*, 26 (5), 40 – 46.  
675
- 676 Lovell, D., Bousson, M., and McLellan, C., 2013. The use of performance tests for the physiological monitoring  
677 of training in combat sports: A case study of a world ranked mixed martial arts fighter. *Journal of Athletic*  
678 *Enhancement*, 2 (1).  
679
- 680 Marinho, B.F., Follmer, B., Esteves, J.V.D.C., and Andreato, L.V., 2016. Body composition, somatotype, and  
681 physical fitness of mixed martial arts athletes. *Sport Sciences for Health*, 12 (2), 157–165.  
682
- 683 Marinho, B.F., Del Vecchio, F.B., and Franchini, E., 2011. Physical fitness and anthropometric profile of mixed  
684 martial arts athletes. *Revista de Artes Marciales Asiáticas*, 6 (2), 11–18.  
685
- 686 McGill, S.M., Chaimberg, J.D., Frost, D.M., and Fenwick, C.M.J., 2010. Evidence of a double peak in muscle  
687 activation to enhance strike speed and force: an example with elite mixed martial arts fighters. *Journal of*

688 *Strength and Conditioning Research*, 24 (2), 348–57.  
689  
690 Miarka, B., Brito, C.J., and Amtmann, J., 2017. Performance probabilities and outcome effects in technical-  
691 tactical factors with bout phase changes of mixed martial arts. *International Journal of Performance Analysis*  
692 *in Sport*, 17 (4), 510–520.  
693  
694 Miarka, B., Brito, C.J., Dal Bello, F., and Amtmann, J., 2017. Motor actions and spatiotemporal changes by  
695 weight divisions of mixed martial arts: applications for training. *Human Movement Science*, 55, 73–80.  
696  
697 Miarka, B., Brito, C.J., Moreira, D.G., and Amtmann, J., 2018. Differences by Ending Rounds and Other  
698 Rounds in Time-Motion Analysis of Mixed Martial Arts: Implications for Assessment and Training. *The*  
699 *Journal of Strength & Conditioning Research*, 32 (2), 534–544.  
700  
701 Miarka, B., Coswig, V., Brito, J.C., Slimani, M., Amtmann, J., and Del Vecchio, F.B., 2016. Comparison of  
702 combat outcomes: technical and tactical analysis of female MMA. *International Journal of Performance*  
703 *Analysis in Sport*, 16 (2), 539–552.  
704  
705 Miarka, B., Coswig, V.S., and Amtmann, J., 2019. Long MMA fights technical-tactical analysis of mixed  
706 martial arts: implications for assessment and training. *International Journal of Performance Analysis in*  
707 *Sport*, 1–14.  
708  
709 Miarka, B., Coswig, V.S., Del Vecchio, F.B., Brito, C.J., and Amtmann, J., 2015. Comparisons of time-motion  
710 analysis of mixed martial arts rounds by weight divisions. *International Journal of Performance Analysis in*  
711 *Sport*, 15 (3), 1189–1201.  
712  
713 Miarka, B., Del Vecchio, F.B., Brito, C.J., Tabben, M., Amtmann, J., and Chamari, K., 2016. Effects of home  
714 advantage in mixed martial arts performance with paired bouts of the same fighting opponents. *International*  
715 *Journal of Performance Analysis in Sport*, 16 (3), 948–960.  
716  
717 Miarka, B., Del Vecchio, F.B., Camey, S., and Amtmann, J.A., 2016. Comparisons: Technical-tactical and time-  
718 motion analysis of mixed martial arts by outcomes. *The Journal of Strength & Conditioning Research*, 30  
719 (7), 1975–1984.  
720  
721 Mikeska, J.D., 2014. A 12-week metabolic conditioning program for a mixed martial artist. *Strength &*  
722 *Conditioning Journal*, 36 (5), 61–67.  
723  
724 Monson, T.A., Brasil, M.F., and Hlusko, L.J., 2018. Allometric variation in modern humans and the relationship  
725 between body proportions and elite athletic success. *Journal of Anthropology of Sport and Physical*  
726 *Education*, 2 (3), 3–8.  
727  
728 Mooney, M., Cormack, S., O'Brien, B., and Coutts, A.J., 2013. Do physical capacity and interchange rest  
729 periods influence match exercise-intensity profile in Australian football? *International Journal of Sports*  
730 *Physiology and Performance*, 8 (2), 165–172.  
731  
732 Niiniluoto, I., 1993. The aim and structure of applied research. *Erkenntnis*, 38 (1), 1–21.  
733  
734 NJSAC, 2002. Mixed martial arts unified rules of conduct additional mixed martial arts rules [online]. Available  
735 from: <http://www.nj.gov/lps/sacb/docs/martial.html> [Accessed 19–6 2018].  
736  
737 Nunes, L.A.S., Lazarim, F.L., Brenzikofer, R., and Macedo, D.V., 2012. Applicability of the reference interval  
738 and reference change value of hematological and biochemical biomarkers to sport science. In: Zaslav,  
739 Kenneth, ed. *An International Perspective on Topics in Sports Medicine and Sports Injury*. IntechOpen, 77 –  
740 98.  
741  
742 De Oliveira, S.N., Follmer, B., de Moraes, M.A., dos Santos, J.O.L., de Souza Bezerra, E., Gonçalves, H.J.C.,  
743 and Rossato, M., 2015. Physiological profiles of north Brazilian mixed martial artists (MMA). *Journal of*  
744 *Exercise Physiology Online*, 18 (1), 56–61.  
745  
746 Paccotti, P., Minetto, M., Terzolo, M., Ventura, M., Ganzit, G., Borrione, P., Termine, A., and Angeli, A., 2005.  
747 Effects of high-intensity isokinetic exercise on salivary cortisol in athletes with different training schedules:

748 relationships to serum cortisol and lactate. *International Journal of Sports Medicine*, 26 (09), 747–755.  
749

750 Peacock, C., Mena, M., Sanders, G., Silver, T., Kalman, D., and Antonio, J., 2019. Sleep data, physical  
751 performance, and injuries in preparation for professional mixed martial arts. *Sports*, 7 (1), 1.  
752

753 Peacock, C.A., Sanders, G.J., Antonio, J., and Silver, T.A., 2018. The Reporting of a Multifaceted Mixed  
754 Martial Arts Strength and Conditioning Program. *Journal of Exercise Physiology Online*, 21 (1), 86–91.  
755

756 Pitchers, G. and Elliot-Sale, K., 2019. Considerations for coaches training female athletes. *Professional Strength  
757 and Conditioning*, 55, 19–30.  
758

759 Richmond, E. and Rogol, A.D., 2014. Traumatic brain injury: endocrine consequences in children and adults.  
760 *Endocrine*, 45 (1), 3–8.  
761

762 Rovell, D. and Okamoto, B., 2016. Dana White on \$4 billion UFC sale: “Sport is going to the next level”  
763 [online]. Available from: [https://www.espn.co.uk/mma/story/\\_/id/16970360/ufc-sold-unprecedented-4-](https://www.espn.co.uk/mma/story/_/id/16970360/ufc-sold-unprecedented-4-billion-dana-white-confirms)  
764 billion-dana-white-confirms [Accessed 16–10 2019].  
765

766 Ruiz, M.C., 2017. I Symposium of Elite Performance in Combat Sports. *Revista de Artes Marciales Asiáticas*,  
767 12 (1), 41–44.  
768

769 Santos, D.A. dos, Miarka, B., Bello, F. dal, Queiroz, A.C.C., de Carvalho, P.H.B., Brito, C.J., and Beneke, R.,  
770 2018. 10-Years on Time-Motion and Motor Actions of Paired Mixed Martial Arts Athletes. *International  
771 Journal of Sports Physiology and Performance*, 1–13.  
772

773 Schick, M., Brown, L., Coburn, J., Beam, W., Schick, E., and Dabbs, N., 2010. Physiological profile of mixed  
774 martial artists. *Medicina Sportiva*, 14 (4), 182–187.  
775

776 Schick, M., Brown, L., and Schick, E., 2012. Strength and conditioning considerations for female mixed martial  
777 artists. *Strength & Conditioning Journal*, 34 (1), 66–75.  
778

779 Seidenberg, P.H., 2011. Mixed martial arts: injury patterns and issues for the ringside physician. *Current Sports  
780 Medicine Reports*, 10 (3), 147–150.  
781

782 Slimani, M., Chaabene, H., Davis, P., Franchini, E., Cheour, F., and Chamari, K., 2017. Performance aspects  
783 and physiological responses in male amateur boxing competitions: A brief review. *Journal of Strength and  
784 Conditioning Research*, 31 (4), 1132–1141.  
785

786 Souza, R.P. de, Sousa, M. do S.C. de, Neves, E.B., Rosa, C., Cruz, I.R.D., Júnior, A.T., Macedo, J.O.R. de,  
787 Reis, V.M., and Vilaça-Alves, J., 2017. Acute effect of a fight of Mixed Martial Arts (MMA) on the serum  
788 concentrations of testosterone, cortisol, creatine kinase, lactate, and glucose. *Motricidade*, 13 (1), 30–37.  
789

790 Souza-Junior, T.P., Ide, B.N., Sasaki, J.E., Lima, R.F., Abad, C.C., Leite, R.D., Barros, M.P., and Utter, A.C.,  
791 2015. Mixed martial arts: History, physiology and training aspects. *The Open Sports Sciences Journal*, 8 (1).  
792

793 Spanias, C., Nikolaidis, P.T., Rosemann, T., and Knechtle, B., 2019. Anthropometric and physiological profile  
794 of mixed martial art athletes: A brief review. *Sports*, 7 (6), 146.  
795

796 Tack, C., 2013. Evidence-based guidelines for strength and conditioning in mixed martial arts. *Strength &  
797 Conditioning Journal*, 35 (5), 79–92.  
798

799 Tota, L., Drwal, T., Maciejczyk, M., Szygula, Z., Pilch, W., Palka, T., and Lech, G., 2014. Effects of original  
800 physical training programme on changes in body composition, upper limb peak power and aerobic  
801 performance of a mixed martial arts fighter. *Medicina Sportiva*, 18 (2).  
802

803 UFC, 2019. UFC Performance Institute [online]. Available from: <http://www.ufc.com/performance-institute/>  
804 [Accessed 16–9 2019].  
805

- 806 Del Vecchio, F.B. and Franchini, E., 2013. Specificity of high-intensity intermittent action remains important to  
807 MMA athletes' physical conditioning: response to Paillard (2011). *Perceptual & Motor Skills*, 116 (1).  
808
- 809 Del Vecchio, F.B., Hirata, S.M., and Franchini, E., 2011. A review of time-motion analysis and combat  
810 development in mixed martial arts matches at regional level tournaments. *Perceptual and Motor Skills*, 112  
811 (2), 639–648.  
812
- 813 Del Vecchio, F.B., da Silva, K.C., and Miarka, B., 2015. Time-motion analysis of mixed martial arts (MMA)  
814 combats: Comparison between genders. *Conexões*, 13 (3), 48–64.  
815
- 816 Vecchio, L., Stanton, R., Macgregor, C., Humphries, B., and Borges, N., 2019. Effects of a six-week strength  
817 and power program on punching and kicking impact power in amateur male combat athletes: A pilot study.  
818 *Journal of Athletic Enhancement*, 8 (1).  
819
- 820 Verhagen, E., Voogt, N., Bruinsma, A., and Finch, C.F., 2014. A knowledge transfer scheme to bridge the gap  
821 between science and practice: an integration of existing research frameworks into a tool for practice. *British*  
822 *Journal of Sports Medicine*, 48 (8), 698–701.  
823
- 824 Wagenmakers, E.-J., Love, J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., Selker, R., Gronau, Q.F.,  
825 Dropmann, D., Boutin, B., and others, 2018. Bayesian inference for psychology. Part II: Example  
826 applications with JASP. *Psychonomic Bulletin & Review*, 25 (1), 58–76.  
827  
828



<b>Table 2 – MMA performance related research placed within the ARMSS model</b>				
<b>Stage of ARMSS Model, followed by Research Category</b>	<b>Study</b>	<b>Cohort n and type</b>	<b>Key Data</b>	<b>Conclusions</b>
<b>1: Defining the problem</b>				
Athlete	(Lenetsky and Harris 2012)	N/A#	6 peer reviewed studies related to MMA performance.	Call for more MMA specific research to be conducted due to the paucity of extant research at the time. MMA training prescription based on physiological measurements of other combat sports.
	(Andreato and Branco 2016)	N/A#	Author communication.	Discussion of applicability of other combat sports physiologies to MMA research, and recommendation to base physiological profiling of MMA on work-to-rest (W:R) ratios and bout length.
	(James et al. 2016)	Systematic review of n = 23 combat sports related physiological studies	n of studies included and discussed: Brazilian jiu jitsu = 1 Boxing = 2 Judo = 8 Karate = 4 Wrestling = 8	The lack of robust physiological studies existing in MMA led authors to determine training requirements of MMA competitors based on findings from other related combat sports. Grappling based sports require a greater maximal strength in comparison to striking based sports, with greater movement velocity being characteristic of striking based sports.
	(Loneragan et al. 2018)	N/A#	Strength and conditioning (S&C) based needs analysis of MMA performance	S&C for MMA should be multifaceted and aimed towards the development of aerobic capacity and 'explosive' movements. <i>Nb. S&amp;C = strength and conditioning</i>
	(Spanias et al. 2019)	Systematic review of n = 19 MMA related physiological studies	Review of existing studies into anthropometry and physical performance on MMA competitors.	MMA competitors tend to have low body fat, high flexibility, muscle strength, muscular endurance and anaerobic power, with moderate cardiovascular capacity. Lack of consistent testing protocols between studies make comparisons difficult, but there does not appear to be differences between performance levels.
Training	(Amtmann 2004)	n = 28 regional MMA competitors, age range = 19 - 37	Freq of S&C sessions per week range = 1 - 7; freq of MMA sessions per week range = 3 – 12; 12 participants stated using neck strengthening exercises; 8 participants stated using Olympic lift variations; 5 participants admitted to anabolic steroid use.	Questionnaire conducted immediately prior to competition. Large variations in frequency and types of S&C training, poor education regarding training methods
	(Amtmann 2010)	n = 32 regional MMA competitors, age range = 19 – 41	Freq of S&C sessions per week range = 1 - 6; freq of MMA sessions per week range = 2 – 10; 16 participants stated using neck strengthening exercises; 8 participants stated using Olympic lift variations; 2 participants admitted to anabolic steroid use.	Questionnaire conducted immediately prior to competition. Large variations in frequency and types of S&C training, poor education regarding training methods. <i>Nb. S&amp;C = strength and conditioning</i>
	(Amtmann 2011)	N/A#	Author communication.	Discussion of MMA training based on other combat sports in lieu of detailed MMA research. Suggests basing training on work-to-rest ratio (W:R) found in MMA studies until more research is conducted.
	(Jay 2013)	N/A#	Narrative review of MMA research at time of publication in relation to <u>strength and conditioning research</u>	S&C recommendations based on general S&C research. Highlights the lack of MMA specific research as a barrier. <i>Nb. S&amp;C = strength and conditioning</i>
	(Souza-Junior et al. 2015)	N/A#	Non-systematic review of MMA literature and other related combat sports literature.	Too little MMA specific research has been conducted to form specific physiological profile or training framework. Research from other combat sports and sports physiology in general should be used in the interim
	(Del Vecchio and Franchini 2013)	N/A#	Author communication.	Correspondence discussing the need for more W:R research to investigate the intermittent nature of MMA. Suggests basing training on W:R and individual athlete's responses until more research is conducted. <i>Nb. W:R = work-to-rest ratio</i>
<b>2: Descriptive research</b>				

Training	(Hurst et al. 2014)	n = 8 male MMA competitors, age = 25.5 ± 4.5	Player load (PLd) intraclass correlation coefficient (ICC) = 0.700 – 0.970 and accumulated player load (PLd <sub>ACC</sub> ) ICC = 0.794 – 0.984. Coefficient of variation (CV) = 2.4 – 7.8%.	Demonstration of reliability of body worn accelerometry to measure external load of MMA participation. Measured across 8 Catapult Minimax units, with 5 repetitions of 10 standing striking techniques, 6 ground striking techniques and 2 takedown techniques.
	(Tota et al. 2014)	n = 1 male, professional MMA competitor preparing for one bout over an 11-week period	Mean power: Pre = 6.9, Post = 7.1 W·kg <sup>-1</sup> ; Peak power: Pre = 7.79, Post = 8.1 W·kg <sup>-1</sup> ; VO <sub>2</sub> max: Pre = 57.1, Post = 58.4 ml·kg <sup>-1</sup> ·min <sup>-1</sup> ; Lean body mass increase = 1.5 kg; Fat mass decrease = 1.4 kg.	11 weeks of a specific training program brought about positive, measurable changes in body composition and physiological capabilities. Though times, distance and prescribed intensities of the training program are reported, there is no detail about training methods or modes provided.
	(Kirk et al. 2015a)	n = 8 male MMA competitors, age = 25.5 ± 4.5	Significant differences in PLd (au) between sparring and isolation: jab = 2.04 ± .29 (sparring), 2.88 ± .37 (isolation); cross = 2.25 ± .26 (sparring), 3.37 ± .37 (isolation); left hook = 2.48 ± .31 (sparring), 3.18 ± .4 (isolation).	Accelerometry used to show that jabs, crosses and left hooks cause greater PLd in isolated training than in sparring, single leg takedowns cause greater PLd than double leg takedowns and takedowns cause greater PLd than strikes, with no differences in PLd between a successful and an unsuccessful takedown. Despite no significant differences found, right hook, left leg kick, right body kick, right high kick, successful double leg takedown (attacking and defending) all had either a moderate or large effect size (Cohen's d) between isolated training and sparring. <i>Nb. Pld = Playerload</i>
	(Jukic et al. 2017)	n = 1 male, professional MMA competitor preparing for a one night international tournament	Weekly sRPE (au): Week 1 = 2,210; Week 2 = 5,132; Week 3 = 6,035; Week 4 = 3,240; Week 5 = 5,443; Week 6 = 2,925; Week 7 = 3,442; Week 8 = 3,105; Week 9 = 1,345; Mean = 3,653; Max HR = 182 beats·min <sup>-1</sup> ; VO <sub>2</sub> max = 55.2 ml·kg <sup>-1</sup> ·min <sup>-1</sup>	Case study detailing the S&C training practices and weekly loads over a 9-week training period of an elite participant preparing for competition in 2006. Training included strength, power, strength endurance, aerobic conditioning and injury prevention strategies in keeping with accepted S&C and periodisation theory <i>Nb. sRPE = sessional rating of perceived exertion</i>
	(Chernozub et al. 2019)	n = 40 MMA participants (age 21 ± 0.8), split into two groups: group A: resistance training low intensity/high volume; group B resistance training high intensity/low volume	Fat free BM: A pre = 69.1 ± 3.3 kg, post 70.8 ± 3.4 kg; B pre = 68.7 ± 3.4, 71.5 ± 3.3 kg. Cortisol: A pre = 283.5 nmol·L <sup>-1</sup> , post 483.9 nmol·L <sup>-1</sup> ; B pre = 458.8 nmol·L <sup>-1</sup> , post = 641.3 nmol·L <sup>-1</sup> . LDH: A pre = 492 cu, post = 440 cu; B pre = 465 cu, post 366.5 cu.	MMA competitors respond as expected to high intensity/low volume and low intensity/high volume resistance training over 3 months. No indication of whether body composition or biomarker changes are related to MMA performance. <i>Nb. LDH = lactate dehydrogenase. cu = article authors do not define this unit</i>
Performance	(Amtmann et al. 2008)	n = 6 competitive males, age range = 21 - 41	Post MMA training (n = 4): Lac (mmol·L <sup>-1</sup> ) = 13 – 19.7; RPE (au) = 17 – 19; Post MMA sparring (n = 4) : Lac = 13.3 – 18; RPE = 15 – 18; Post competitive MMA bout (n = 6): Lac = 10.2 – 20.7, RPE = 13 -19	Suggests that full competitive bouts (3 x 5 mins) require equal exertion to sparring bouts. 4 of the 6 bouts ended in the first round producing lower values for lactate and/or RPE for these participants. <i>Nb. RPE = rating of perceived exertion</i>
	(Del Vecchio et al. 2011)	n = 52 male professional competitors, age = 24 ± 5, n of bouts = 26	W:R = 9:1 – 6:1 when including the break between rounds; 1:2 – 1:4 when excluding the break between rounds.	Only significant difference between rounds found in low intensity groundwork between round 2 (36 ± 26 s) and round 3 (21 ± 6 s). Over half of bouts end in high intensity ground positions, meaning training should be aimed at simulating these positions and intensities.
	(Miarka et al. 2015)	n = 2,097 male professional bouts	Effort time (s): 1 <sup>st</sup> round: Longest = 269.8 ± 64.1 (LW); Shortest = 212.4 ± 101.5 (HW); 2 <sup>nd</sup> round: Longest = 277.8 ± 61.6 (FW); Shortest = 132.8 ± 90.9 (BW);	W:R profile of MMA bouts is significantly different between weight divisions and between rounds, demonstrating intermittent nature of event and potential differences in pacing strategies between divisions. <i>Nb. W:R = work-to-rest ratio</i>

			3 <sup>rd</sup> round: Longest = 289.6 ± 42.3 (WW) Shortest = 246.3 ± 89.1 (HW)	
	(Del Vecchio et al. 2015)	n = 64 professional MMA bouts, 32 male bouts and 32 female bouts	High intensity to low intensity ratio = 2:1; Low intensity standing (min:s): Males = 04:19.7 ± 03:19.5; Females = 06:55.6 ± 04:47.7; High intensity ground (min:s): Males = 01:14.6 ± 01:16.4; Females = 0:40.8 ± 01:9.1; Freq of attacks: Males = 47 ± 22.83; Females = 21 ± 15.11; Attacks to head: Males = 31 ± 18.51; Females = 15 ± 6.28.	Female participants spend longer in competition engaged in low intensity standing movements than male participants, less time in low intensity standing and grounded movements than males.
	(Adam et al. 2015)	n = 1 professional, regional bout between two male MMA competitors (Comp A and Comp B)	Attack activity by round: Round 1 = 4.6 in favour of Comp A; Round 2 = 1.6 in favour of Comp B; Round 3 = 13.3 in favour of Comp A. Attack versatility index by round: Round 1: Comp A = 0.05, Comp B = 0.11; Round 2: Comp A = 0.12, Comp B = 0.1; Round 3: Comp A = 0.03, Comp B = 0.13	Single bout description of techniques used, showing one participant using stand up striking only, the other participant using a combination of stand up striking and ground grappling. <i>Nb. Attack activity = sum of attacks per minute minus sum of opponent's attacks per minute. Attack versatility = metric representing the range of different attacks from different technique groups</i>
	(Kirk et al. 2015b)	n = 6 male MMA competitors, age = 26.17 ± 5.04, one 3 x 5 minute sparring bout each	PL <sub>ACC</sub> per minute (PL <sub>ACC</sub> ·min <sup>-1</sup> ) (au) = 224 ± 26.59; Lactate (mmol·L <sup>-1</sup> ) = 9.25 ± 2.96; W:R = 1:1.01; bout winners takedowns = 3 ± 1 (attempted 4 ± 2), bout losers takedowns = 0 (attempted 2 ± 2)	Lactate measurements significantly different across each sampling point (rest, post warm up, end of each round and 5 minutes post bout). PL <sub>ACC</sub> and PL <sub>ACC</sub> ·min <sup>-1</sup> displayed a linear reduction in each round. Successful takedowns only significant differences between winners and losers of bouts. <i>Nb. PL<sub>ACC</sub> = accumulated Playerload; PL<sub>ACC</sub>·min<sup>-1</sup> = accumulated Playerload per minute</i>
	(Coswig, Fukuda, et al. 2016)	n = 25 male MMA competitors, age = 26.5 ± 5	Sparring bouts (13): Lac (mmol·L <sup>-1</sup> ) Pre = 3.8 [2.8 – 5.5], Post = 16.8 [12.3 – 19.2]; Creatine Kinase (CK) (U.L) Pre = 225 [136.5 – 330], Post = 297 [208.5 – 403.5] <i>Results reported as Median [interquartile range]</i>	No significant differences in pre or post lactate, blood glucose (Glu), alanine aminotransferase (ALT) or creatine kinase (CK) between competition and sparring bouts. Glu significantly increased pre to post in both competition and sparring bouts. Pre and post measurements of ALT significantly lower in sparring bouts than in competition bouts, possibly suggesting onset of overtraining during period between bouts.
	(Coswig, Ramos, et al. 2016)	n = 13 male MMA competitors, age = 25 ± 5	Sparring bouts: Urea (mg·ml <sup>-1</sup> ) Pre = 42.08 ± 8.77, Post = 44.15 ± 8.93, 48H post = 36.31 ± 7.85; Glu (mg·ml <sup>-1</sup> ) Pre = 80.38 ± 12.7, Post = 156.54 ± 19.09, 48H post = 87.69 ± 15.5	Urea significantly decreased from resting 48H post bout with a large ES; Glu significantly increased with a large ES immediately post sparring bout, and a significant reduction with a small ES 48H post bout. No significant changes in alanine aminotransferase (ALT), magnesium (Mg) or uric acid (UA) over same time periods. MMA sparring bouts present low muscle damage and require relatively short physiological recovery times during training periods.
	(Souza et al. 2017)	n = 20 male professional MMA competitors paired in sparring bouts, bout winners age = 26.2 ± 2.39; bout losers age = 24.3 ± 1.83	Testosterone (nmol·L <sup>-1</sup> ): Pre = 13.03 ± 3.9; Post = 9.53 ± 3.33; 24H post = 14.68 ± 4.02 (sig differences between bout winners and losers at all sample points). Cortisol (nmol·L <sup>-1</sup> ):	Significant increase in testosterone (T) and cortisol (C) immediately post bout with return to resting levels 24H post. Creatine kinase (CK) displayed a significant increase immediately post bout, with an almost three-fold increase over the next 24H. Suggests MMA involves significant physiological, metabolic and muscular strain including acute muscle damage. More sampling points required over a longer time frame to assess the extent of these affects.

			Pre = 549.02 ± 144.99; Post = 876.09 ± 107.84; 24H post = 436.96 ± 126.26 Creatine kinase (CK) (U·L): Pre = 433.89 ± 215.95, Post = 491.81 ± 219.17, 24H post = 1412.69 ± 758.63	
	(Ghoul et al. 2017)	n = 12 male MMA competitors paired in sparring bouts, age = 26 ± 5	C (ng·dL <sup>-1</sup> ): Pre = 88.3 ± 19.6, Post = 131.2 ± 56.4, 30min post = 117.1 ± 46.3, 24H post = 79.3 ± 23.2; T (ng·dL <sup>-1</sup> ): Pre = 3.5 ± 0.3, Post = 4.6 ± 0.3, 30min post = 3.7 ± 0.3, 24H post = 3.4 ± 0.3; Uric acid (UA)(μmol·L <sup>-1</sup> ): Pre = 308 ± 53, Post = 342 ± 54, 30min post = 414 ± 58, 24H post = 344 ± 48; Lactate dehydrogenase (LDH) (U·L): Pre = 169 ± 70, Post = 230 ± 116, 30min post = 228 ± 94, 24H post = 226 ± 75.	Cortisol (C) and testosterone (T) significantly raised at end of 3 round sparring bout, with uric acid and lactate remaining significantly elevated 24H after bout. MMA is a physiologically demanding event with fatigue indices remaining high 24 hours after a bout. Results could inform training and recovery protocols.
	(Fernandes et al. 2018)	n = 1,496 professional MMA bouts, analysed as pre 2012 rule changes versus post 2012 rule changes	Total strike attempts: Pre = 41.5 ± 25.9, Post = 43.6 ± 26.4; Participant exposure time (mins): Pre = 383.3, Post = 480.2; Bouts ending in final round: Pre: 90, Post = 150; Rounds with injuries: Pre = 68.2%, Post = 44.8%	The frequency of some striking techniques and bout exposure time increased after a 2012 change to the MMA bout scoring criteria, with a lower injury incidence and a greater number of bouts ending in the final round
	(Antonietto et al. 2019)	n = 678 professional MMA bouts	Standing preparatory time in bouts ending in: 1 <sup>st</sup> round = 95.6 ± 62.9 s; 2 <sup>nd</sup> round = 93.6 ± 67.9 s; 3 <sup>rd</sup> round = 144 ± 88.5 s.	MMA competitors become less active as the bout progresses, demonstrating pacing strategies potentially being employed in bouts, or the onset of fatigue as the bout progresses.
Athlete	(McGill et al. 2010)	n = 5 international/elite standard males, age = 29 ± 1.8	EMG study demonstrating double peak of muscle activation (rectus abdominus, latissimus dorsi, erector spinae, gluteus medius, gluteus maximus and rectus femoris) during strikes on heavy bag.	Double peak of muscle activation possible due to contraction-relaxation-contraction pulses of observed muscles to achieve strike speed and force. May be affected by different specific training methods.
	(Schick et al. 2010)	n = 11 male regional MMA competitors, age = 25.5 ± 5.7	Body fat = 11.7 ± 4%; VO <sub>2</sub> max = 55.5 ± 7.3 ml·kg <sup>-1</sup> ·min <sup>-1</sup> ; vertical jump = 57.6 ± 7.3 cm; relative bench press strength = 1.2 ± 0.1 kg·kg <sup>-1</sup> ; relative squat strength = 1.4 ± 0.1 kg·kg <sup>-1</sup> ; absolute grip strength = 45.8 ± 6.2 kg.	MMA participants displayed lower maximal oxygen consumption than kickboxers, lower power than wrestlers, similar bench press and squat strength to judokas, and lower grip strength than boxers.
	(Marinho et al. 2011)	n = 13 male MMA competitors, age = 30 ± 4	Body fat = 11.87 ± 5.11%; squat strength = 73 ± 15 kg; broad jump = 219 ± 0.25 cm	MMA participants have relatively low levels of strength and moderate levels of power.
	(Alm and Yu 2013)	n = 5 male MMA competitors, age = 29.6 ± 5.5	VO <sub>2</sub> max (ml·kg <sup>-1</sup> ·min <sup>-1</sup> ) Pre = 50 ± 6.5 , Post = 48.5 ± 5.7; Counter movement jump (CMJ) (cm) Pre = 43.1 ± 5.07, Post = 41.25 ± 2.13; Relative CMJ with arm swing Pre = 0.63 ± 0.4, Post = 0.59 ± 0.09;	Low strength and aerobic capacity before and after 12 months of MMA training, with some variables reducing (non significant). Only CMJ with arm swing displayed a statistically significant increase over the 12 month period. Suggests MMA training alone is not sufficient to bring about significant physiological performance adaptations.

			Relative hang clean (kg·kg <sup>-1</sup> ) Pre = 1.09 ± 0.07, Post = 1.06 ± 0.11;	
	(Lovell et al. 2013)	n = 1 case study, national standard MMA competitor, age = 25	Upper body $\dot{V}O_2$ max (ml·kg <sup>-1</sup> ·min <sup>-1</sup> ) Pre = 40.35, Post = 45.96; Lower body $\dot{V}O_2$ max Pre = 54.96, Post = 56.75; Upper body peak power (W·kg <sup>-1</sup> ) Pre = 798 W·kg, Post = 841; Lower body peak power Pre = 914, Post = 934; MHR (beats·min <sup>-1</sup> ) Pre = 183, Post = 191	Upper (14%) and lower (4%) body $\dot{V}O_2$ max, upper body peak power (5.4%), lower body peak power (2.2%) and MHR (4%) increased after 8 weeks of MMA training supplemented by a specific S&C program. Authors suggest use of Wingate anaerobic test (WAnT) to monitor adaptations to training for MMA competition. <i>Nb. S&amp;C = strength and conditioning; MHR = maximal heart rate</i>
	(de Oliveira et al. 2015)	n = 18 male regional MMA competitors, age = 27.9 ± 5.9	$\dot{V}O_2$ max = 44.22 ± 6.69 ml·kg <sup>-1</sup> ·min <sup>-1</sup> ; Right hand grip strength = 45.99 ± 8.99 kg; Left hand grip strength = 45 ± 8.5 kg	MMA participants have low $\dot{V}O_2$ max and low handgrip strength in comparison to other combat sports such as wrestling, boxing, karate and Brazilian jiu jitsu.
	(Marinho et al. 2016)	n = 8 male MMA competitors, age = 31 ± 5	Body mass index (BMI) = 26 ± 3.3 kg·m <sup>-2</sup> ; body fat = 11.8 ± 6.2 kg and 13.4 ± 5.6%; muscle mass = 69.6 ± 4.6%; flexed arm hang = 35 ± 10s; relative squat strength = 0.84 ± 0.1 kg·kg <sup>-1</sup> ; relative bench press = 1 ± 0.2 kg·kg <sup>-1</sup>	MMA participants display high lean body mass, excellent upper body muscular endurance and low levels of relative strength.
	(Antonio et al. 2018)	n = 15 male professional MMA competitors, age = 28.4 ± 4.4	Bone mineral content (BMC) (kg) = 3.9 ± 0.52; Bone mineral density (BMD) (g·cm <sup>3</sup> ) = 1.57 ± 0.1	BMC and BMD measured using DXA scanning. MMA participants display higher BMD than other sports (including track and distance athletics, swimming, American football and stand up paddling), with no significant differences to American football players. <i>Nb. BMC = bone mineral content; BMD = bone mineral density; DXA = dual energy x-ray absorptiometry</i>
<b>3: Predictors of performance</b>				
Performance	(Baker and Schorer 2013)	n = 1,468 professional MMA competitor records and profiles	Southpaw (left handed) winning % = 64 ± 20.4 Orthodox (right handed) winning % = 62.6 ± 21.3 No statistically significant difference	No indication of lateral preference influencing success in professional bouts
	(Miarka, Del Vecchio, Camey, et al. 2016)	n = 645 professional MMA rounds	Total strike attempts ES between winners and losers: 1 <sup>st</sup> round = -0.20, 2 <sup>nd</sup> round = -0.25, 3 <sup>rd</sup> round = -0.21; Advances to half guard, side control, mount and back mount all significantly different (≤0.05) in favour of bout winner (with exception of mount in 1 <sup>st</sup> round).	Winners and losers distinguished by the numbers of total strikes, submissions and positional improvements <i>Nb. ES = effect size</i>
	(Miarka, Del Vecchio, Brito, et al. 2016)	n = 202 professional MMA bouts (n = 101 participants classified as 'home', n = 101 classified as 'away')	Total strikes attempted: Home = 37.3 ± 23.9; Away = 46.4 ± 26.3; Head strikes attempted: Home = 21.5 ± 15.1; Away = 26.5 ± 16.2; Low intensity ground time (s): Home = 13.8 ± 6.2; away = 5 ± 0.1.	No effect of 'home vs away' in determining winner or loser, but some effect on numbers of strikes attempted and landed.

	(Miarka, Coswig, et al. 2016)	n = 174 professional female MMA rounds	Time (s) spent in low intensity standing combat by round for bout ending methods: Split decision = 160.4 ± 83.6; Unanimous decision = 158.4 ± 87.6; KO/TKO = 44.8 ± 38.8; Submission = 42.1 ± 44.1. Total combat time(s) by round for bout ending methods: Split decision = 300.7 ± 0.3; Unanimous decision = 300 ± 0.4; KO/TKO = 154.4 ± 95.2; Submission = 204.2 ± 96.6.	Winners and losers distinguished by striking and grappling actions in bouts that ended via strikes or submissions, distinguished by standing strikes only in bouts that ended via decision.
	(Kirk 2016a)	n = 474 professional MMA bouts	Shorter competitors ranked higher in FIW ( $\omega^2 = 0.14$ ); Shorter competitors ranked in the middle in WSW ( $\omega^2 = 0.2$ ); Shorter competitors more likely to have competed for/won a world title in FW ( $\omega^2 = -1.3$ ) and FIW ( $\omega^2 = -0.95$ )	Negligible effect of stature and wingspan on rankings, isolated to small number of divisions. <i>Nb. <math>\omega^2 = \text{omega squared effect size}</math></i>
	(Kirk 2016b)	n = 278 professional MMA bouts, bout winner's age = 29.79 ± 4.3; bout loser's age = 30.79 ± 4.3.	Bout winner stature (cm): Won by strikes = 181.4 ± 9; won by submission = 177.3 ± 8.3; won by decision = 177.1 ± 8.6. Bout loser stature (cm): Lost by strikes = 180.7 ± 8.2; lost by submission = 176.9 ± 8.5; lost by decision = 176.5 ± 9.1. Bout winner age: Won by strikes = 30.59 ± 4.8; won by submission = 28.44 ± 3.4; won by decision = 29.68 ± 4. Bout loser age: Lost by strikes = 31.8 ± 4.4; lost by submission = 29.19 ± 4.1; lost by decision = 30.6 ± 4.1.	Taller participants and participants with a longer armspan more likely to lose due to strikes, older participants more likely to lose in general, but when bout winner is older it is most likely due to decision. No other effects of anthropometry better than moderate found (n = 278 bouts).
	(James, Robertson, et al. 2017)	n = 234 professional MMA bouts.	% of competitors winning after achieving >4 significant ground strikes = 80.4%, increased to 84.9% when takedown accuracy >25%; 0.850 significant ground strikes per minute = 91.5% chance of winning, increased to 91.5% when strikes >4.19 per minute.	Decision tree used to determine grappling and technique accuracy determine success in professional competitive MMA bouts.
	(Miarka, Brito, and Amtmann 2017)	n = 584 professional MMA bouts	Probability of winning bout if achieved more than opponent: Head strikes landed = 14.8%; Successful takedowns = 76.79%; Offensive passes = 66.7%; Chokes attempted = 69.67%	Distance head strikes landed, guard passes and successful takedowns most contribute to winning a bout.
	(Miarka, Brito, Dal Bello, et al. 2017)	n = 2,814 professional MMA rounds	Time keeping distance (s) sig. different to than other divisions: FW = 131.4 ± 89.9; HW = 179.2 ± 93.4; LW = 163.4 ± 88.4. Clinch time without attack (s) sig. greater than other divisions:	Performance in MMA divisions distinguished by different frequencies of actions in both standing and grounded positions, with 'rest' time being a key difference between divisions.

			FIW = 11.4 ± 10.1; WW = 12.6 ± 13.	
	(Miarka et al. 2018)	n = 678 professional MMA bouts, 1,564 rounds	% of KO/TKO by round: 1 <sup>st</sup> = 63.5%, 2 <sup>nd</sup> = 58.1%, 3 <sup>rd</sup> = 64%; % of submissions by round: 1 <sup>st</sup> = 30.3%, 37.2%, 3 <sup>rd</sup> = 0.5%.	Lower time spent in low intensity striking with increased forcefulness of actions in grounded positions in first two rounds increase likelihood of winning. <i>Nb. KO = knockout, TKO = technical knockout</i>
	(Brito et al. 2018)	n = 54 MMA competitors tested positive for PEDs in post bout testing, 2001-2014	Bouts won by PED competitors = 34 (60%); Androgenic steroids = 55%, psychotropic = 27%, thermogenic/diuretic = 9%, opioids = 9%;	Competitors who tested positive for PEDs and bout winners displayed greater time spent in high intensity movements than bout losers. No differences between PED competitors and non-PED competitors in terms of technical variables or likelihood of winning. <i>Nb. PED = performance enhancing drug</i>
	(Kirk 2018a)	n = 461 professional MMA bouts	Divisions where bout winners/losers can be distinguished by: striking only (BF <sub>10</sub> = 399-10) = HW; by striking (BF <sub>10</sub> = 791,661 - 7) and moderately by grappling (BF <sub>10</sub> = 75-7) = LHW, MW, FW, BW, WBW, WSW; by striking (BF <sub>10</sub> = 3.533e <sup>+6</sup> - 221) and grappling (BF <sub>10</sub> = 17,100 - 50): WW, LW, FIW.	Technical differences between winners and losers distinct between divisions, with some differences having moderate to weak predictive relationships to anthropometry in HW, WW, LW and FW.
	(Kirk 2018b)	n= 100 professional MMA competitors in each division from HW - BW, each division split into 5 ranking groups (RG).	Divisional age: HW = 32.8 ± 5.3; LHW = 31.3 ± 4.4; MW = 31.4 ± 4.4; WW = 30.8 ± 4.1; LW = 30.1 ± 3.4; FW = 29.6 ± 3.9; BW = 29 ± 4.1 Cross divisional RG age: RG1 = 32.3 ± 4.4; RG2 = 30.5 ± 3.9; RG3 = 30.1 ± 4.1; RG5 = 29.9 ± 4.7	As divisional weight increases, so too does participant age (with a decisive BF <sub>10</sub> ) with general trend repeating across each RG. Competitors in RG1 are older than all other RG (decisive to strong BF <sub>10</sub> ) with general trend repeating across each division. MW displays a moderate correlation between age and rank.
	(Monson et al. 2018)	n = 1,284 professional MMA competitors	Stature (cm) = 178.3 ± 8.51; Armspan (cm) = 182.55 ± 10.22; Stature-to-armspan ratio (S:W) = 1.02	S:W found to predict success in MMA based on winning %. <1% (R <sup>2</sup> = 0.008) of winning % variance predicted by S:W.
	(Santos et al. 2018)	n = 45 professional male MMA competitors, assessed twice, 10 years apart (M1 and M2)	Differences between time points: Strikes landed: M1 = 22[12, 34], M2 = 18[18, 31.7]; strikes attempted: M1 = 41[24.5, 62], M2 = 35[21, 48]; takedowns attempted: M1 = 1[0.1,2], M2 = 1[0,2]	The variables most associated with winning in MMA changed from 2000-2014 with successful body and head strikes becoming more associated with winning. Head strikes attempted, total strikes attempted, and submissions attempted became less associated with winning. <i>Nb. Result shown as median[interquartile range]</i>
	(dal Bello et al. 2019)	n = 304 professional MMA rounds	Bouts ending in decision = greater freq. of takedowns attempted per round (ES = 0.13); Bouts ending in KO/TKO = lesser freq. of submissions attempted per round (ES = 0.15-0.30); Bouts ending in submission = greater freq. of sweeps per round (ES = 0.27)	The type and frequency of grappling techniques (takedowns, sweeps and submission attempts) used directly affects the outcome of the bout (unanimous/split decision, submission or KO) (n = 304 rounds). <i>Nb. ES = effect size, KO = knockout, TKO = technical knockout</i>
	(Miarka et al. 2019)	n = 779 rounds from professional MMA bouts which ended in rounds 3 - 5.	Low intensity combat: 3 <sup>rd</sup> round = 75%; 4 <sup>th</sup> round = 84%; 5 <sup>th</sup> round = 79%; Freq. of strikes attempted: 1 <sup>st</sup> round = 39.5 ± 17.6; 2 <sup>nd</sup> round = 46.1 ± 21.9; 3 <sup>rd</sup> round = 46.1 ± 27	Activity profiles of bouts ending in the 3 <sup>rd</sup> , 4 <sup>th</sup> or 5 <sup>th</sup> rounds are significantly different to bouts which end in rounds 1 or 2, indicating differing training requirements.

Athlete	(James, Beckman, et al. 2017)	n = 29 male semi-professional (HL n = 15, age = 29.5 ± 2.2) and amateur (LL n = 14, age = 26.6 ± 7.95) male competitors	Jump squat time to peak force at 50% BM load(s): HL = 0.718 ± 0.209; LL = 0.898 ± 0.169 (ES = -0.86); Jump squat mean RFD (W·kg <sup>-1</sup> ·s <sup>-1</sup> ) at 50% BM load: HL = 143.03 ± 51.28; LL = 104.82 ± 25.73 (ES = 0.93); Jump squat peak power (W·kg <sup>-1</sup> ): HL = 44.45 ± 7.54, LL = 38.47 ± 6.74 (ES = 0.78); Jump squat peak velocity (m·s <sup>-1</sup> ): HL = 3.06 ± 0.33, LL = 2.81 ± 0.33 (ES = 0.72)	Maximal lower body neuromuscular capabilities distinguish between high-level (based on number of wins being ≥50% of verified competition record) and low-level performers. No differences in reactive strength index, 1RM bench press strength or relative peak force measured using isometric mid-thigh pull (IMTP). Results used to suggest that success is related to lower body neuromuscular force capabilities and training should be aimed towards improving this area. <i>Nb. 1RM = one repetition maximum</i>
	(James et al. 2018)	n = 15 professional male MMA competitors, split into HL (n = 7, >50% professional wins, MMA training age = 6.14 ± 0.71) and LL (n = 8, amateur bouts only, MMA training age = 4.31 ± 1.53).	Differences between groups: 20M sprint: ES = -0.49[-1.39, 0.49]; 10M sprint: ES = -0.18[-1.11, 0.76]; Yo-Yo distance covered: ES = 0.57[-0.37, 1.5]; Repeat sprint ability: ES = small to moderate (-0.2 - -0.61) differences in times at each sprint interval	10 and 20M sprint times, repeat sprint ability, and Yo-Yo Intermittent Recovery test results distinguish between professional/semi-professional and amateur competitors to a small effect size or better in favour of the HL group. <i>Nb. ES = Cohen's d[90% CI]</i>
	(Peacock et al. 2019)	n = 7 professional, male MMA participants; age = 27.2 ± 3.4; mass = 77.3 ± 0.3 kg; stature = 180.7 ± 4.4 cm.	Sleep latency correlated to: Vertical jump (r = -0.787), Prowler push time (r = 0.776), VO <sub>2</sub> max (r = -0.860). Heart rate recovery correlated to: Sleep latency (r = -0.739), Sleep efficiency (r = -0.891), Sleep onset variance (r = 0.710). Number of missed sessions correlated to: Sleep latency (r = -0.789), Sleep onset variance (r = 0.788)	Sleep quality may directly affect physiological performance measures and training attendance amongst MMA participants.
	(de Azevedo et al. 2019)	n = 11 male MMA participants (age = 27.6 ± 4.3, mass = 83.5 ± 7.8 kg)	Repeat punch protocol onto force plate, intervention group consumed 5 ml·kg <sup>-1</sup> caffeine, control group consumed placebo.	No differences in punch force or frequency between groups.
<b>4: Experimental testing of key performance predictors</b>				
Training	(Bodden et al. 2015)	n = 25 male semi-professional MMA competitors, age = 24.31 ± 4.46, split into INT group and CON group	FMS scores pre: INT = 13.25 ± 0.87, CON = 13.23 ± 0.8; Week 4: INT = 15.17 ± 1.21, CON = ~13.4 (not specified) Week 8: INT = 15.33 ± 1.43, CON = ~13.34 (not specified)	Training intervention (corrective FMS exercises performed 4 times a week for 8 weeks for INT group) can improve FMS scores of MMA participants. <i>Nb. FMS = Functional Movement Screen</i>
	(Kostikiadis et al. 2018)	INT group n = 10 male professional MMA competitors, age = 28.9 ± 4.2; CON group n =	Post intervention changes ES in favour of INT group: Est. VO <sub>2</sub> max = 0.69; Bench press 1RM = 0.82;	Specific S&C programme alongside MMA training significantly improves estimated VO <sub>2</sub> max, 1RM strength, CMJ height and power, SJ height and 10M sprint in comparison to control group (regular self-selected training alongside MMA training). <i>Nb. ES = Hedge's g, CMJ = countermovement jump, 1RM = one repetition maximum</i>

		7 male professional MMA competitors, age = 25.7 ± 5	Back squat 1RM = 1.17; Dead lift 1RM = 1.63; CMJ height = 0.70; CMJ power = 0.70; 10m sprint time = 0.54	
	(Chernozub et al. 2018)	n = 30 male MMA competitors, age = 21 ± 1.2, split into two groups n= 15 based on using a striking based approach (STR) or a wrestling (WRE) based approach to bouts.	Bench Press 1RM (kg) pre: STR = 68.2 ± 3.7, WRE = 68.1 ± 2.3; 3 months post: STR = 99.2 ± 9.3, WRE = 83.8 ± 4.9; 6 months post: STR = 105.8 ± 7.4, WRE = 95 ± 6.4. Serum cortisol (C) (nmol·L <sup>-1</sup> ) pre: STR = 184.8[100,263], WRE = 197[109,269]; 3 months post: STR = 275[200, 311], WRE = 12[2,43]**; 6 months post: STR = 250[144, 276], WRE = 98[69, 166]**	Significant differences in serum C and bench press 1RM found between groups distinguished by combat style after a power training programme designed specifically for their combat style. No details of training programme provided.  ** Results as reported by authors in table appear to differ to results in figures displaying supposedly same data. May be some errors due to pre-publication translation in original paper.  Nb. C results reported as median[interquartile range]; 1RM = one repetition maximum
	(Peacock et al. 2018)	n = 12 male professional MMA competitors, age = 25.2 ± 2.3	BESS pre = 11.9 ± 31; post = 10.8 ± 2.8; Sit and reach (cm) pre = 17.4 ± 4.2, post = 18.8 ± 4.3	Balance and flexibility improved over the course of 6-week MMA training period incorporating Tai Chi movements, no control group for comparison Nb. BESS = Balance error scoring system
	(Vecchio et al. 2019)	n = 6 MMA participants and 10 kickboxing participants split into intervention group = 10 (INT age = 25.2 ± 1.8, mass = 76 ± 7.2 kg, stature = 178.1 ± 7.1 cm) and control group = 6 (CON age = 29 ± 2, mass = 79.8 ± 11.9 kg, stature = 177.7 ± 5.7 cm). Unclear how MMA participants are split between groups.	Post 6-week strength and power training program: Within INT group changes: cross punch force (W) = likely increase, roundhouse kick force (W) = likely increase, vertical jump (cm) = likely increase. No within CON group changes found. Between INT and CON differences: INT likely greater changes than CON for cross punch force and 5RM squat (kg).	6-week strength and power training program caused likely* increases in punch, kick and lower limb force production in comparison to CON group only completing regular sport specific training. *Nb. Inferred from magnitude-based inferences (MBI) and 90% confidence intervals (90%CI).
<b>5: Determinants of key performance predictors</b>	No studies found			
<b>6: Efficacy studies</b>	No studies found			
<b>7: Barriers to uptake</b>	No studies found			
<b>8: Implementation studies</b>	No studies found			
<b>No ARMSS Category</b>				
Training	(La Bounty et al. 2011)	N/A#	Narrative review of MMA research at time of publication in relation to strength and conditioning research	S&C considerations for MMA based on findings from other combat sport and general S&C research Nb. S&C = strength and conditioning
	(Schick et al. 2012)	N/A#	Narrative review of MMA research at time of publication in relation to strength and conditioning research for female athletes	S&C considerations for female MMA participants based on findings from other combat sports and general S&C research
	(Tack 2013)	N/A#	Narrative review of MMA research at time of publication in relation to strength and conditioning research	S&C considerations for MMA based on findings from other combat sport and general S&C research
	(James et al. 2013)	N/A#	Narrative review of MMA research at time of publication in relation to strength and conditioning research for basing conditioning on W:R and metabolic requirements of sport	Suggested training structure based on W:R studies in MMA and general S&C research Nb. W:R = work-to-rest ratio
	(Mikeska 2014)	N/A#	Narrative review of MMA research at time of publication in relation to	Metabolic conditioning program based on findings from other combat sports and general S&C research

			strength and conditioning research for field based metabolic conditioning	
	(James 2014)	N/A#	Narrative review of MMA research at time of publication in relation to strength and conditioning research for injury prevention	S&C injury prevention recommendations based on MMA injury studies and general S&C research.
	(Earnshaw 2015)	N/A#	Narrative review of MMA research at time of publication in relation to strength and conditioning research	Recommendations for the content of an MMA S&C mesocycle based on general S&C research
	(Harvey 2018)	N/A#	Narrative review of MMA research at time of publication in relation to HIIT training	Details the development of a potential MMA specific HIIT protocol (MMASIT) encouraging researchers and practitioners to test the validity of the protocol. <i>Nb. HIIT = high intensity interval training</i>
<i>Nb. #No experimental cohort used</i>				