

Daily Changes of Resting Metabolic Rate in Elite Rugby Union Players.

JAMES HUDSON¹, MATTHEW COLE², JAMES P. MORTON¹, CLAIRE STEWART¹ &
GRAEME L. CLOSE¹

*¹Research Institute for Sport and Exercise Sciences, Liverpool John Moores University,
Liverpool, UK*

²Sport and Exercise Science, Birmingham City University, Birmingham, UK

Address for correspondence:

Professor Graeme L. Close,
Research Institute for Sport and Exercise Sciences,
Liverpool John Moores University,
Tom Reilly Building,
Bryom St Campus,
Liverpool,
L3 3AF, UK
g.l.close@ljmu.ac.uk

21 **Abstract**

22 **INTRODUCTION:** Preparation for competitive contact sport has been extensively
23 researched. There are, however, limited data to guide players as to how the demands of their
24 sport affect the energy requirements of recovery. We aimed to provide novel data on changes
25 in resting metabolic rate (RMR) in contact sport athletes and relate these to the physical
26 demands of training and competition.

27 **METHODS:** 22 Elite professional Premiership Rugby Union players were recruited to the
28 study. Indirect calorimetry (Vyntus CPX canopy, CareFusion) was used to measure RMR
29 each morning of the competitive game week, in a fasted, rested state. External loads for
30 training and game play were monitored and recorded using global positioning systems
31 (Catapult Innovations, Australia), whilst internal loads were tracked using rate of perceived
32 exertion scales. Collisions were reviewed and recorded by expert video analysts for contacts
33 in general play (breakdown and tackle area) or the set piece (scrum or maul).

34 **RESULTS:** There were significant ($p=0.005$) mean increases in RMR of ≈ 231 kcal the
35 morning after (GD+1) and 3 days after the game (GD+3), compared with the day before the
36 game (GD-1). The players were exposed to internal and external loads during the training
37 week comparable to that of a match day, however, despite the equivocal loads between
38 training and game play, there were no significant increases in RMR following training.

39 **CONCLUSION:** The collisions experienced in rugby match play are likely to be responsible
40 for the significant increases in RMR at GD+1 and GD+3. Consequently, the measurement of
41 RMR via indirect calorimetry may provide a novel non-invasive measure of the effects of
42 collisions. This study provides a novel insight to the energy requirements of recovering from
43 contact sport.

44 **Keywords:** Damage, Team Sport, Contact, Injury, DOMS, Soreness

45

46 **Introduction.**

47 Rugby Union is a dynamic and combative team sport participated in globally (1). Two teams
48 of 15 players, broadly categorised as forwards (n=8) and backs (n=7) contest a match for 80
49 minutes (1). The sport is comprised of intermittent, high intensity activities incorporating high
50 speed running, sprinting, accelerations and decelerations (2-4). Rugby Union also involves
51 collision-based activities at the tackle area (tackle and breakdown contest) and the set piece
52 (scrum and maul). Time motion analysis and global positioning systems (GPS) studies report
53 that forwards experience ~60% more high level impacts during contact situations than backs
54 (4). However, there are significant limitations of using GPS technology to determine contact
55 occurrence and quantitative measurement of force, rendering it unreliable to determine the
56 physical strain placed on the players (5).

57 Whilst the technology to accurately quantify physical collisions in rugby is currently lacking,
58 the recognition of their impact made upon the athlete is not (6). The forces and mechanical
59 stress in rugby can cause exercise induced muscle damage (EIMD) and impact induced muscle
60 damage (IIMD) which may be distinct in their symptomology and recovery time course (7).
61 These physical collisions have been shown to increase indirect markers of muscle damage (8,
62 9), reduce neuromuscular function (10, 11), and increase perception of muscle soreness (11).
63 Sport scientists have examined a wide array of modalities to enhance recovery from the
64 damaging collisions of rugby match play, some of which may mildly alleviate symptoms (12).
65 However, despite multiple interventions being implemented, we have reported that elite rugby
66 players are in pain every day throughout a competitive rugby season (13). It is therefore crucial
67 that accurate and quantitative markers are developed to assess the extent of the IIMD to allow
68 more targeted interventions to be developed. One potential candidate is assessing the energy
69 expenditure of players given that the total energy expenditure (TEE) of young rugby league
70 players was 5% higher when training weeks involved collisions (14).

71 Resting metabolic rate (RMR) is the primary component of TEE and is the energy expended to
72 maintain homeostasis at rest. Indirect calorimetry (IC) requiring both oxygen ($\dot{V}O_2$) and carbon
73 dioxide ($\dot{V}CO_2$) to be measured is the most accurate method of assessing RMR (15). Large
74 variations in the estimation of RMR using prediction equations have been noted in a variety of
75 sports (16), especially athletes with a high fat free mass (17) such as rugby (18-20). It is
76 therefore imperative that RMR is accurately measured rather than predicted using equations.
77 Importantly, much of the existing understanding around effectively calculating an athlete's
78 energy requirements are based upon studies which primarily utilise recreational or youth
79 athletes and are thus limited by lower training ages and exposures to lower absolute intensities
80 of work. To our knowledge there are no data on the daily variations in RMR across an entire
81 competitive match week in any sport, including positional differences. It is therefore crucial
82 that potential changes in RMR are explored in highly trained professional athletes with indirect
83 calorimetry performed prior to and the days following a competitive fixture.

84 To facilitate recovery, it is essential that rugby players are provided with the correct nutrition
85 in terms of both the total energy intake and the provision of recovery promoting foods. The
86 majority of nutrition research in rugby has focussed upon preparation for match play, ensuring
87 muscle glycogen concentrations are optimal for performance (21). It appears elite players now
88 have a good understanding of this (22), however, the nutritional intakes in the days following
89 a match are much more variable (22) with many players decreasing total energy intake the day
90 after a game. If muscle damage arising from match play causes an increase in energy
91 requirements in recovery, current guidelines could be underestimating player's needs post
92 competition.

93 To this end, the objectives of the present study were twofold. 1) To assess, for the first-time,
94 changes in RMR in an elite group of professional rugby union players measured throughout a
95 competitive week, including the days before and after a professional game using indirect

96 calorimetry. 2) To explore the relationship between game day factors, (e.g. the number of
97 physical collisions), and changes in RMR. These data would provide more accurate
98 information into the energy requirements of players in the days after a game, which could help
99 recovery strategy, as well as providing a novel non-invasive assessment of the effects of the
100 physical collisions upon the players.

101 **Methods**

102 *Participants*

103 A convenience sample of twenty-two healthy elite rugby union players, all members of an
104 English Premiership squad, were recruited for this study. The participants included six
105 internationals, and many established Premiership or Super 15 players (mean \pm SD, age; 25.7
106 \pm 4.1 years, body mass; 104.6 \pm 12.6 kg). Five participants were excluded from the analysis
107 having sustained an injury during games which prevented them from completing all aspects of
108 the study. All playing positions were covered in the remaining 17 players who were eligible
109 for the full study analysis. All participants gave written informed consent prior to commencing
110 the study. Ethical approval (18/SPS/004) was granted by the university research ethics
111 committee at Liverpool John Moores University, UK.

112 *Research Design*

113 The study was designed to allow RMR to be measured within the training schedules of elite
114 rugby players during a complete microcycle. Timepoints throughout the study are described
115 relative to game day (GD) using +/- symbols for days before (-) and days after (+) GD. Due to
116 the timing of team selection defining when recruitment could occur, the first measurement was
117 taken at GD-2. Measurements were then repeated every day, apart from the game day itself, as
118 this was deemed too disruptive to the players' habitual routine. Table 1 details the training
119 schedule for the match week. Seven microcycles were used to attain the total data set, with all

games played on the Saturday afternoon (Game Day). This ensured that the training schedules throughout the microcycle were the same and there were no conflicting kick-off times, which would alter the time relative to match play of the subsequent measures. Internal and external loads for training and match play were recorded throughout the week. The weeks chosen were throughout the middle of the season (weeks 13-30) so the players were accustomed to the training load and rigours of match play.

Resting Metabolic Rate

The RMR of participants was assessed 6 times in total. All measures were completed at the same time between 7-9am and players arrived after an overnight fast, with their last meal at least 8 hours prior to measurement. Players awoke and came straight to the training ground as per reliable outpatient protocol (23). To ensure best practice, a private room was established at the training facility away from the main building where temperature was maintained at 21-23 °C, the room was dimly lit, and quiet (15). Players lay in a comfortable supine position and were reminded to stay awake. A twenty minute resting period was prescribed, as the minimum sufficient time to achieve rest (24). A ventilated hood was employed rather than mouth piece and nose clip to reduce day-to-day variance (25). The coefficient of variance for our protocol was measured at 1.13% for RMR and 1.62% for RER. The ventilated hood was placed over the head of the athlete and expired gas was analysed using the dilution canopy method (Vyntus CPX canopy, CareFusion, Hoechberg, Germany). The gas analyser was calibrated every day using the manufacturer's automated flow and digital volume transducer calibration (15.92% O₂ and 5.03% CO₂). The first 5 minutes of measurements were discarded following best practice guidelines (15). Measurements were subsequently recorded for 15 minutes continuously at 10 second intervals for $\dot{V}O_2$ and $\dot{V}CO_2$. Data were exported into Microsoft Excel (2018, Seattle, USA), and mean respiratory exchange ratio (RER) across the

measurement period generated, with the calorific value, carbohydrate and fat oxidation rates determined according to the table of Zuntz (26).

Measurement of lean body mass

Lean body mass was measured using a dual-energy-X-ray absorptiometry (DXA) fan beam scanner (Hologic Horizon W, Hologic, Bedford, MA), with scanning and analysis performed by the same trained individual using Apex software version 13.5.3.1(Hologic, Bedford, MA). Players were scanned twice during the period of data collection for this study and the scan corresponding closest to their week of participation used, which was no longer than 4 weeks. Measurements were taken first thing in the morning prior to eating, drinking, or exercise and protocols implemented to maximise reliability of positioning (27).

Training and Match loads

Internal loads for each training day and the game day were assessed by the session rating of perceived exertion (sRPE) using a modified Borg scale (28). This RPE of the training session was multiplied by the training duration to calculate a player load in arbitrary units (sRPE; AU) (28). External demands of all rugby training sessions and match play were recorded using micro-technological units worn by players containing GPS (10Hz) and accelerometer (100Hz) (Catapult Innovations, Melbourne, Australia). Data were downloaded and analysed using Catapult Sprint software (Catapult Innovations, Melbourne, Australia). The total distance covered, number of high-speed efforts (>60% positional average) and the number of very high-speed efforts (>80% individual average) were recorded (29, 30). The GPS sampling frequency of 10Hz is the most reliable in team sports measuring high speed running activities (31).

Contacts were analysed in match play by a professional rugby union analyst with over five years' experience working in English domestic and European rugby using NacSport (Analysis Pro, UK). The potential collisions were then further reviewed by an expert ex-professional player with 15 years and over 250 matches played in English domestic, European and

International Rugby Union. The games were reviewed to ensure contacts recorded involved an actual collision. For example, a scrum may be analysed as a single contact but there may have been more than one engagement process involving a full collision before the match restarted. Some players may also be analysed as having been involved at a breakdown to keep possession, but they may not necessarily have endured a collision as part of this. The nature of these were then also accounted for as either set piece based (scrum or maul), or general phase play (breakdown and tackle area).

Data analysis

All data are presented as mean (\pm SD). All statistical analyses were completed using SPSS (Version 24 for Windows, SPSS Inc., Chicago, IL). A one-way repeated measures ANOVA was used to compare all gas exchange measures and the work completed by players throughout training days and during the competitive game day. The tests of within subjects' effects provided values for Mauchly's test for sphericity. If this was violated, then a Greenhouse-Geisser correction was used. The difference between means were tested at a significance level of $p < 0.05$. The least significant difference (LSD) was used post hoc to compare specific time points when the ANOVA revealed a significant difference between measures over the week. This was examined in the whole group ($n=17$), sub-groups forwards ($n=11$) and backs ($n=6$). A Spearman's correlation was run to assess any associations between changes in RMR throughout the microcycle, with the metrics of physical load and collision data gathered from the competitive match play ($n=17$). A Spearman's rank-order correlation coefficient value (r_s) was generated and this was tested at $p < 0.05$ to test the significance of any relationships found (32).

****TABLE 1 ABOUT HERE****

Results

Training and match demands

The training schedule and structure of sessions can be seen in Table 1 with the internal and external demands of the week in Table 2. It should be noted that data are presented as n=14 for these analyses due to faults with GPS data collection, resulting in lost running metrics for some training sessions in three of the participants.

****TABLE 2 ABOUT HERE****

Player Load

There was no significant difference in player load on GD+3 compared with GD. This was also true for the sub-groups of forwards and backs. The player load on all other days of the training week were significantly lower than the game day in the whole group and when subdivided into forwards, and backs.

High Speed Running Distance

In the whole group, there was no significant difference in high speed running distance covered on GD+3 compared with GD. In the forwards sub-group, there was only significantly less HSR distance covered on GD-1($p=0.001$) and GD+2($p=0.013$) compared with GD. In the backs sub-group, there was significantly less HSR distance covered on GD-2 ($p=0.005$), GD-1 ($p<0.0005$), GD+2 ($p<0.0005$), and GD+3($p=0.019$) compared with GD.

Number of High Speed Running Efforts

In the whole group, there were significantly fewer HSR efforts on GD-2 ($p=0.002$), GD-1 ($p<0.0005$), GD+2 ($p<0.0005$), and GD+3 (0.031) compared with GD. In the forwards subgroup, significantly fewer HSR efforts were completed on GD-1 ($p=0.001$) and GD+2

($p=0.014$) compared with GD. In the backs sub-group, significantly fewer HSR efforts were completed on GD-2 ($p=0.003$), GD-1 ($p<0.0005$), GD+2 ($p=0.001$), and GD+3 ($p=0.001$) compared with GD.

Very High Speed Running Distance

In the whole group, VHRSR distance was only significantly lower on GD-1 ($p=0.002$) and GD+2 ($p=0.002$) compared with GD. Within the forwards sub-group, there was no significant difference in VHRSR distances covered on any day compared with GD. The backs covered significantly fewer VHRSR metres on GD-1 ($p=0.005$) and GD+2 ($p=0.006$).

Very High Speed Running Efforts

In the whole group, the number of VHRSR efforts completed was only significantly lower on GD-1 ($p=0.003$), and GD+2 ($p=0.013$) compared with GD. In the forwards sub-group, there was no significant difference in VHRSR efforts on all training days compared with GD. In the backs sub-group, there were only significantly less VHRSR efforts on GD-1 ($p=0.001$), and GD+2 ($p=0.013$) compared with GD.

****FIGURE 1. a-f ABOUT HERE****

****TABLE 3 ABOUT HERE****

Changes in resting metabolic rate

Changes in RMR adjusted for lean body mass across the microcycle can be seen in Figure 1a whilst the absolute ($\text{kcal}\cdot\text{day}^{-1}$) and relative ($\text{kcal}\cdot\text{kg}\cdot\text{day}^{-1}$) RMR measures are displayed in Table 3. Lean body mass (measured by DXA) was $74.8\pm 7.4\text{kg}$ for the whole group, $78.2\pm 5.6\text{kg}$ for the forwards, and $68.6\pm 6.0\text{kg}$ for the backs. In the whole group, there was a significant increase in RMR from GD-1 to GD+1 ($p=0.005$) and GD-1 to GD+3 ($p=0.04$). In the forwards sub-group, there was a significant increase in RMR between GD-1 to GD+1 ($p=0.017$) and

GD-1 to GD+3 ($p=0.045$). However, in the backs sub-group, there was no significant difference in RMR at any time point across the week.

Changes in respiratory exchange ratio

Changes in RER across the microcycle can be seen in Figure 1b. In the whole group, there were significant increases at GD+2 ($p=0.030$) and GD+3 ($p=0.006$) compared with GD-1. In the positional subgroups there were no significant differences across the microcycle $p=0.065$ and $p=0.177$ for forwards and backs respectively.

Changes in VO_2 and VCO_2

Figures 1c and 1d show the measures of VO_2 and VCO_2 . There were significant increases in VO_2 in the whole group at GD+1 ($p=0.008$) and GD+3 ($p=0.041$) compared with GD-1. These significant increases were also observed in the forwards at GD+1 ($p=0.025$) and GD+3 ($p=0.027$) compared with GD-1. There were no significant differences for VO_2 in the backs subgroup across the week. There were significant increases in VCO_2 in the whole group at GD+1 ($p=0.008$), GD+2 ($p=0.01$), and GD+3 ($p=0.001$) compared to GD-1. These significant increases were also observed in the forwards at GD+1 ($p=0.037$) and GD+3 ($p<0.001$) compared to GD-1. There were no significant differences across the week in measures of VCO_2 in the backs.

Changes in carbohydrate and fat oxidation

Measures of carbohydrate and fat oxidation are displayed in Figures 1e and 1f. Carbohydrate oxidation significantly increased at GD+2 ($p=0.044$) and GD+3 ($p=0.003$) compared with GD-1 in the whole group. In the forwards a significant increase was measured at GD+3 ($p=0.003$) compared with GD-1, whilst there were no significant differences across the microcycle in the backs for carbohydrate oxidation. Fat oxidation decreased significantly at GD+3 ($p=0.029$) in the whole group and at the same time point in the forwards ($p=0.028$) compared with GD-1.

262 There were no significant differences measured for fat oxidation across the microcycle in the
263 backs.

264 **TABLE 4 ABOUT HERE**

265 *Associations of match demands with changes in metabolic measurements*

266 Table 3 displays the Spearman's coefficient associations between the physical match demands,
267 and changes in RMR. In the whole group, there were no significant associations found between
268 phase contacts, total contacts, player load, HSR meters, HSR efforts, VHSR meters, VHSR
269 efforts and the change in RMR observed between GD-1 to GD+1. This was also true when the
270 positional sub-groups of forwards and backs were analysed.

Discussion

The aim of the present study was to assess changes in RMR in an elite group of professional RU players measured throughout a competitive week and explore the impact of game day factors on changes in RMR. To this end, we monitored RMR using indirect calorimetry alongside game day and training demands in 22 Premiership RU players throughout a game week. We report, for the first-time, that RMR increased significantly following elite rugby union match play, a change that was not observed following intense training with the same training loads. These data therefore illustrate that changes in RMR following match days exist, reflecting a yet unreported increased energy demand in the days after a game of elite rugby and allows the development of individualised nutritional strategies to help facilitate recovery. Furthermore, increased RMR may also represent the physical collisions of match play and indeed could suggest that RMR may be used as a non-invasive marker of muscle damage.

We have reported a mean increase in RMR following match play of ~231kcal per day at GD+1, a 10% increase from GD-1. We are confident this represents a truly significant increase given that it is greater than the suggested 6% required as meaningful change using the canopy method (25). The rigour in our protocol also resulted in a lower coefficient of variance than reported previously (25). Importantly, these increases in RMR were due to significant increases in VO₂ and VCO₂ and are not merely EPOC being measured as increased VO₂. The range of increased RMR was large, with individual responses between 240-1000kcal. The greatest increases in RMR were seen in the forwards, who underwent more physical collisions during a game at the scrum, maul and tackle area vs. backs (4). The whole group, and forwards positional group, also experienced increased RMR which remained elevated 3 days post-game. This sustained increase at GD+3 may be a result of the lower limb resistance training session on GD+2 given that resistance training, especially with an eccentric component, has been shown to increase RMR (33). It is possible that this sustained increase in RMR, as a result of the resistance

training session, negatively affected the recovery from match play, therefore extending the period during which RMR remained elevated, although this suggestion remains speculative and requires further investigation.

Along with changes in RMR in the days after the game we also report significant changes in RER. The increased RER at GD+2 and GD+3 corresponds with significant increases in resting carbohydrate oxidation coupled with a significant reduction in fat oxidation at GD+3. These significant changes in carbohydrate oxidation are occurring at a time where markers of muscle damage and soreness typically peak following match play (8, 9, 11). Muscle damage induced reductions in glucose transport may result in a decreased whole-body glucose tolerance which has been reported after a laboratory based muscle damage protocol (34). It should also be recognised that the inflammatory cytokine activity associated with muscle damaging exercise, together with the presence of various cell types such as neutrophils and macrophages (35), may alter substrate oxidation in the recovery period (36). Taken together, we have demonstrated increased RMR and altered carbohydrate oxidation, following match play, which suggests that post-exercise nutrition should be specifically tailored to the unique metabolic demands of this time period. Moreover, we have shown highly individual responses with some players increasing their RMR by 1000kcal. It is crucial to identify such players and tailor their dietary plans and recovery strategies accordingly.

Given that the participants in the present study were full-time professional players, in the middle of a competitive playing season, it was not possible to either control or record dietary intake. It is possible that some of the differences in RMR and RER seen between the forwards and backs could have been a result of differing diets of the 2 sub-groups. However, whilst there is evidence that the thermic effect of food and the total energy content of a meal may alter resting metabolic rate measures (15) we do not believe that the player to player variations in diet would have any meaningful effects on RMR or RER in the present study. Previous research

has reported that a large meal containing 1300kcal had negligible effects upon measuring RMR and RER when measured 7 hours later, and in lean male subjects both measures had returned to baseline at 8 hours following this meal (37). Given that both the forwards and backs in the present study had undergone a minimum of an 8 hour fast prior to having their RMR and RER assessed, it is unlikely that differences in diet would be a primary contributor to the observed changes. Moreover, we believe that this group of players consumed a more than adequate energy availability as indicated by no major changes in body mass over the testing periods. This group are unlikely to be in low energy availability, however, future studies should attempt to measure or control dietary intake to fully explore this hypothesis.

We propose that the muscle damage as a result of elite rugby union match play could be a key factor in accounting for the changes in metabolism we have witnessed. By carefully monitoring the internal and external demands of the competitive week we have shown that when contact sport athletes are exposed to comparable player load (including HSR and VHSR metrics) to that of a match day but without the physical collisions, there is no change in RMR in the following days. We therefore speculate that the collisions encountered on a game day could be responsible for the significant changes in RMR reported at GD+1. This may account for the increases in TEE previously observed in youth players when a training session contained collisions similar to that of match play (14).

When we investigated the positional groups of forwards and backs there were differences in how they reacted to match play. The backs sub-group did not show any significant changes in RMR or RER post-match, albeit they did show a similar pattern across the week as seen in the forwards sub-group. The backs did not experience as many contact incidents as the forwards as has previously been shown (4), and they were not involved in the static exertions of the scrum and maul which are potentially damaging. These positional differences may further

substantiate our hypothesis that the contact-based activities are responsible for the metabolic changes reported here.

The total number of contacts were rigorously evaluated; however, the Spearman's correlations did not show any significant correlations of changes in RMR with the match demands or collisions experienced. There was one back who exhibited a large increase of ≈ 796 kcal in RMR. Although the actual number of contacts performed by this player were not significantly different to the mean of the backs group, subjective analysis of these collisions (by experienced rugby staff) classified the magnitude and intensity of these as being much greater than typical. Examples like this, coupled with the current inability to accurately quantify collision activities, emphasises the need for a practical measure of the impact contact sports have upon these athletes to be developed.

Practical implications.

From an applied perspective the periodisation of nutrition throughout microcycles to optimise adaptation and ultimately performance is well established under the 'Fuel for the work required' paradigm (38). The novel data presented here could enhance the application of this in team sports, especially those involving muscle damage due to collision-based activities. Even using a modest physical activity level (PAL) of 1.3-1.4 for a GD+1 rest day, would translate these findings into a required increase in energy intake of >300 kcal. This, on a day where the continued restoration of muscle glycogen is a primary concern, in a population who habitually appear to consume lower than the recommended carbohydrate intakes, may require a conscious intervention (22). Carbohydrate intake as part of an in-season week in elite rugby union players appears to be 3 g/kg on GD+1 (22), therefore an extra ~ 70 g carbohydrate could be an increase of $\sim 20\%$ required on that day. We speculate the timing of carbohydrate feeding may also require further investigation though, if indeed substrate oxidation is altered until the muscle damage due to match play is resolved (34, 39).

Given that the true definition of resting metabolic rate involves ‘strict and steady resting conditions’ it could be argued that the present study did not actually measure RMR at any time point where in fact Morning Metabolic Rate (MMR) was actually measured. Indeed, it could be argued that rugby players (and indeed many athletes) during a competitive season are never truly at ‘rest’ bringing about methodological questions over when during a training period RMR should be measured to accurately predict energy requirements. A protocol according to best practice and adhering strictly to a minimum rest time, fasted measurement and proper outpatient protocols *as per* resting metabolic rate are crucial for reliability but this measure may need to be categorised differently (15, 23). In the applied world, the term ‘Morning Metabolic Rate’ may be a more accurate description of what is actually being measured and future studies may choose to adopt this terminology.

Conclusions.

In conclusion, the present study has for the first time assessed the resting metabolic rate of elite rugby union players across a competitive match week using indirect calorimetry. We report a significant increase in the RMR of these contact sport athletes in the days after match play. There were also significant shifts in RER at two and three days after competition. We propose these changes could be attributed to the collisions experienced in match play rather than the internal and external loads the athletes are exposed to throughout the microcycle. The metabolites and markers of these muscle damaging actions need to be researched further to help guide athletes as how best to feed their recovery after competition. This research is the first step in working towards a novel non-invasive marker of muscle damage. Further studies need to control factors of energy availability and exercise modalities responsible for the forms of muscle damage. Protocols more readily useable in the ‘real world’ of applied performance science then need to be designed to truly shift the paradigm of athlete monitoring and optimise recovery from contact sports.

395 **Acknowledgements**

396 The research was funded by the rugby club as part of the post graduate work of JH. Thank you
397 to MC for providing equipment support for the indirect calorimetry. All authors contributed to
398 the study design and preparation of the manuscript. The authors would like to thank all the
399 players for the time taken to perform the RMR analysis and the rugby club for their support.

400 **Conflict of interest.**

401 The authors reported no potential conflict of interest. The results of the current study do not
402 constitute endorsement by ACSM. All results presented here are done so clearly, honestly, and
403 without fabrication, falsification, or inappropriate data manipulation.

404 **Word Count 4444.**

405

406

407

408

409

410

411

412

413

414

415

416 **References.**

- 417 1. Duthie G, Pyne D, Hooper S. Applied physiology and game analysis of rugby union.
418 Sports Med. 2003;33(13):973-91. Epub 2003/11/11. doi: 10.2165/00007256-200333130-
419 00003. PubMed PMID: 14606925.
- 420 2. Roberts SP, Trewartha G, Higgitt RJ, El-Abd J, Stokes KA. The physical demands of
421 elite English rugby union. J Sports Sci. 2008;26(8):825-33. Epub 2008/06/24. doi:
422 10.1080/02640410801942122. PubMed PMID: 18569548.
- 423 3. Austin D, Gabbett T, Jenkins D. The physical demands of Super 14 rugby union. J Sci
424 Med Sport. 2011;14(3):259-63. Epub 2011/02/18. doi: 10.1016/j.jsams.2011.01.003. PubMed
425 PMID: 21324741.
- 426 4. Cunniffe B, Proctor W, Baker JS, Davies B. An evaluation of the physiological
427 demands of elite rugby union using Global Positioning System tracking software. J Strength
428 Cond Res. 2009;23(4):1195-203. Epub 2009/06/17. doi: 10.1519/JSC.0b013e3181a3928b.
429 PubMed PMID: 19528840.
- 430 5. Reardon C, Tobin DP, Tierney P, Delahunt E. Collision count in rugby union: A
431 comparison of micro-technology and video analysis methods. J Sports Sci.
432 2017;35(20):2028-34. Epub 2016/11/22. doi: 10.1080/02640414.2016.1252051. PubMed
433 PMID: 27868475.
- 434 6. Tavares F, Smith TB, Driller M. Fatigue and Recovery in Rugby: A Review. Sports
435 Med. 2017;47(8):1515-30. Epub 2017/02/06. doi: 10.1007/s40279-017-0679-1. PubMed
436 PMID: 28150163.
- 437 7. Naughton M, Miller J, Slater GJ. Impact-Induced Muscle Damage and Contact
438 Sports: Etiology, Effects on Neuromuscular Function and Recovery, and the Modulating
439 Effects of Adaptation and Recovery Strategies. Int J Sports Physiol Perform. 2018;13(8):962-
440 9. Epub 2017/11/29. doi: 10.1123/ijsp.2017-0268. PubMed PMID: 29182431.

- 441 8. McLellan CP, Lovell DI, Gass GC. Biochemical and endocrine responses to impact
442 and collision during elite Rugby League match play. *J Strength Cond Res.* 2011;25(6):1553-
443 62. Epub 2011/05/24. doi: 10.1519/JSC.0b013e3181db9bdd. PubMed PMID: 21602645.
- 444 9. Takarada Y. Evaluation of muscle damage after a rugby match with special reference
445 to tackle plays. *Br J Sports Med.* 2003;37(5):416-9. Epub 2003/09/30. PubMed PMID:
446 14514532; PubMed Central PMCID: PMCPMC1751364.
- 447 10. McLellan CP, Lovell DI. Neuromuscular responses to impact and collision during
448 elite rugby league match play. *J Strength Cond Res.* 2012;26(5):1431-40. Epub 2012/04/21.
449 doi: 10.1519/JSC.0b013e318231a627. PubMed PMID: 22516913.
- 450 11. McLean BD, Coutts AJ, Kelly V, McGuigan MR, Cormack SJ. Neuromuscular,
451 endocrine, and perceptual fatigue responses during different length between-match
452 microcycles in professional rugby league players. *Int J Sports Physiol Perform.*
453 2010;5(3):367-83. Epub 2010/09/24. PubMed PMID: 20861526.
- 454 12. Calleja-Gonzalez J, Mielgo-Ayuso J, Ostojic SM, Jones MT, Marques-Jimenez D,
455 Caparros T, et al. Evidence-based post-exercise recovery strategies in rugby: a narrative
456 review. *Phys Sportsmed.* 2018:1-11. Epub 2018/10/30. doi:
457 10.1080/00913847.2018.1541701. PubMed PMID: 30369286.
- 458 13. Fletcher BD, Twist C, Haigh JD, Brewer C, Morton JP, Close GL. Season-long
459 increases in perceived muscle soreness in professional rugby league players: role of player
460 position, match characteristics and playing surface. *J Sports Sci.* 2016;34(11):1067-72. Epub
461 2015/09/15. doi: 10.1080/02640414.2015.1088166. PubMed PMID: 26368285.
- 462 14. Costello N, Deighton K, Preston T, Matu J, Rowe J, Sawczuk T, et al. Collision
463 activity during training increases total energy expenditure measured via doubly labelled
464 water. *Eur J Appl Physiol.* 2018;118(6):1169-77. Epub 2018/03/24. doi: 10.1007/s00421-
465 018-3846-7. PubMed PMID: 29569055; PubMed Central PMCID: PMCPMC5966477.

- 466 15. Fullmer S, Benson-Davies S, Earthman CP, Frankenfield DC, Gradwell E, Lee PS, et
467 al. Evidence analysis library review of best practices for performing indirect calorimetry in
468 healthy and non-critically ill individuals. *J Acad Nutr Diet*. 2015;115(9):1417-46 e2. Epub
469 2015/06/04. doi: 10.1016/j.jand.2015.04.003. PubMed PMID: 26038298.
- 470 16. Jagim AR, Camic CL, Kisiolek J, Luedke J, Erickson J, Jones MT, et al. Accuracy of
471 Resting Metabolic Rate Prediction Equations in Athletes. *J Strength Cond Res*.
472 2018;32(7):1875-81. Epub 2017/07/07. doi: 10.1519/JSC.0000000000002111. PubMed
473 PMID: 28682934.
- 474 17. Carlsohn A, Scharhag-Rosenberger F, Cassel M, Mayer F. Resting metabolic rate in
475 elite rowers and canoeists: difference between indirect calorimetry and prediction. *Ann Nutr*
476 *Metab*. 2011;58(3):239-44. Epub 2011/08/04. doi: 10.1159/000330119. PubMed PMID:
477 21811063.
- 478 18. Smith DR, King R, Duckworth LC, Sutton L, Preston T, O'Hara JP, et al. Energy
479 expenditure of rugby players during a 14-day in-season period, measured using doubly
480 labelled water. *Eur J Appl Physiol*. 2018;118(3):647-56. Epub 2018/01/22. doi:
481 10.1007/s00421-018-3804-4. PubMed PMID: 29353321; PubMed Central PMCID:
482 PMC5805815.
- 483 19. Morehen JC, Bradley WJ, Clarke J, Twist C, Hambly C, Speakman JR, et al. The
484 Assessment of Total Energy Expenditure During a 14-Day In-Season Period of Professional
485 Rugby League Players Using the Doubly Labelled Water Method. *Int J Sport Nutr Exerc*
486 *Metab*. 2016;26(5):464-72. Epub 2016/10/30. doi: 10.1123/ijsnem.2015-0335. PubMed
487 PMID: 27096279.
- 488 20. MacKenzie-Shalders KL, Byrne NM, King NA, Slater GJ. Are increases in skeletal
489 muscle mass accompanied by changes to resting metabolic rate in rugby athletes over a pre-

490 season training period? Eur J Sport Sci. 2019;1-8. Epub 2019/01/08. doi:
 491 10.1080/17461391.2018.1561951. PubMed PMID: 30614386.

492 21. Bradley WJ, Morehen JC, Haigh J, Clarke J, Donovan TF, Twist C, et al. Muscle
 493 glycogen utilisation during Rugby match play: Effects of pre-game carbohydrate. J Sci Med
 494 Sport. 2016;19(12):1033-8. Epub 2016/05/03. doi: 10.1016/j.jsams.2016.03.008. PubMed
 495 PMID: 27134132.

496 22. Bradley WJ, Cavanagh B, Douglas W, Donovan TF, Twist C, Morton JP, et al.
 497 Energy intake and expenditure assessed 'in-season' in an elite European rugby union squad.
 498 Eur J Sport Sci. 2015;15(6):469-79. Epub 2015/06/10. doi: 10.1080/17461391.2015.1042528.
 499 PubMed PMID: 26055695.

500 23. Bone JL, Burke LM. No Difference in Young Adult Athletes' Resting Energy
 501 Expenditure When Measured Under Inpatient or Outpatient Conditions. Int J Sport Nutr
 502 Exerc Metab. 2018;28(5):464-7. Epub 2017/11/16. doi: 10.1123/ijsnem.2016-0315. PubMed
 503 PMID: 29140154.

504 24. Schols AM, Schoffelen PF, Ceulemans H, Wouters EF, Saris WH. Measurement of
 505 resting energy expenditure in patients with chronic obstructive pulmonary disease in a
 506 clinical setting. JPEN J Parenter Enteral Nutr. 1992;16(4):364-8. Epub 1992/07/01. doi:
 507 10.1177/0148607192016004364. PubMed PMID: 1640635.

508 25. Roffey DM, Byrne NM, Hills AP. Day-to-day variance in measurement of resting
 509 metabolic rate using ventilated-hood and mouthpiece & nose-clip indirect calorimetry
 510 systems. JPEN J Parenter Enteral Nutr. 2006;30(5):426-32. Epub 2006/08/26. PubMed
 511 PMID: 16931612.

512 26. Zuntz N. Ueber die Bedeutung der verschiedenen Nahrstoffe. (About the importance
 513 of different nutrients). Archive European Journal of Physiology. 1901(83):557-71. doi:
 514 10.1007/BF01746509.

515 27. Nana A, Slater GJ, Stewart AD, Burke LM. Methodology review: using dual-energy
516 X-ray absorptiometry (DXA) for the assessment of body composition in athletes and active
517 people. *Int J Sport Nutr Exerc Metab.* 2015;25(2):198-215. Epub 2014/07/17. doi:
518 10.1123/ijsnem.2013-0228. PubMed PMID: 25029265.

519 28. Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, et al. A new
520 approach to monitoring exercise training. *J Strength Cond Res.* 2001;15(1):109-15. Epub
521 2001/11/16. PubMed PMID: 11708692.

522 29. Reardon C, Tobin DP, Delahunt E. Application of Individualized Speed Thresholds to
523 Interpret Position Specific Running Demands in Elite Professional Rugby Union: A GPS
524 Study. *PLoS One.* 2015;10(7):e0133410. Epub 2015/07/25. doi:
525 10.1371/journal.pone.0133410. PubMed PMID: 26208315; PubMed Central PMCID:
526 PMC4514747.

527 30. Tierney P, Tobin DP, Blake C, Delahunt E. Attacking 22 entries in rugby union:
528 running demands and differences between successful and unsuccessful entries. *Scand J Med*
529 *Sci Sports.* 2017;27(12):1934-41. Epub 2016/12/29. doi: 10.1111/sms.12816. PubMed
530 PMID: 28028894.

531 31. Rampinini E, Alberti G, Fiorenza M, Riggio M, Sassi R, Borges TO, et al. Accuracy
532 of GPS devices for measuring high-intensity running in field-based team sports. *Int J Sports*
533 *Med.* 2015;36(1):49-53. Epub 2014/09/26. doi: 10.1055/s-0034-1385866. PubMed PMID:
534 25254901.

535 32. Spearman C. The proof and measurement of association between two things. By C.
536 Spearman, 1904. *Am J Psychol.* 1987;100(3-4):441-71. Epub 1987/01/01. PubMed PMID:
537 3322052.

- 538 33. Dolezal BA, Pottleiger JA, Jacobsen DJ, Benedict SH. Muscle damage and resting
539 metabolic rate after acute resistance exercise with an eccentric overload. *Med Sci Sports*
540 *Exerc.* 2000;32(7):1202-7. Epub 2000/07/27. PubMed PMID: 10912882.
- 541 34. Gonzalez JT, Barwood MJ, Goodall S, Thomas K, Howatson G. Alterations in
542 Whole-Body Insulin Sensitivity Resulting From Repeated Eccentric Exercise of a Single
543 Muscle Group: A Pilot Investigation. *Int J Sport Nutr Exerc Metab.* 2015;25(4):405-10. Epub
544 2015/02/13. doi: 10.1123/ijsnem.2014-0211. PubMed PMID: 25675160.
- 545 35. Peake JM, Neubauer O, Della Gatta PA, Nosaka K. Muscle damage and inflammation
546 during recovery from exercise. *J Appl Physiol (1985).* 2017;122(3):559-70. Epub 2016/12/31.
547 doi: 10.1152/jappphysiol.00971.2016. PubMed PMID: 28035017.
- 548 36. Pearce EL, Pearce EJ. Metabolic pathways in immune cell activation and quiescence.
549 *Immunity.* 2013;38(4):633-43. Epub 2013/04/23. doi: 10.1016/j.immuni.2013.04.005.
550 PubMed PMID: 23601682; PubMed Central PMCID: PMC3654249.
- 551 37. D'Alessio DA, Kavle EC, Mozzoli MA, Smalley KJ, Polansky M, Kendrick ZV, et al.
552 Thermic effect of food in lean and obese men. *J Clin Invest.* 1988;81(6):1781-9. Epub
553 1988/06/01. doi: 10.1172/JCI113520. PubMed PMID: 3384951; PubMed Central PMCID:
554 PMCPMC442625.
- 555 38. Impey SG, Hearris MA, Hammond KM, Bartlett JD, Louis J, Close GL, et al. Fuel for
556 the Work Required: A Theoretical Framework for Carbohydrate Periodization and the
557 Glycogen Threshold Hypothesis. *Sports Med.* 2018;48(5):1031-48. Epub 2018/02/18. doi:
558 10.1007/s40279-018-0867-7. PubMed PMID: 29453741; PubMed Central PMCID:
559 PMCPMC5889771.
- 560 39. Costill DL, Pascoe DD, Fink WJ, Robergs RA, Barr SI, Pearson D. Impaired muscle
561 glycogen resynthesis after eccentric exercise. *J Appl Physiol (1985).* 1990;69(1):46-50. Epub
562 1990/07/01. doi: 10.1152/jappl.1990.69.1.46. PubMed PMID: 2394662.

Table 1. The training sessions throughout the competitive micro cycle. Game Day – GD.

Table 2. Comparison of metrics recorded for training and match play throughout the competitive micro cycle.

*Denotes values significantly different ($p < 0.05$) when compared with **game day (GD)** shown in bold.

Figure 1. Gas exchange measurements across the microcycle. a. RMR ($\text{kcal} \cdot \text{kg} \cdot \text{day}^{-1}$). b. RER. c. VO_2 (L/min). d. VCO_2 (L/min). e. Carbohydrate (CHO) oxidation (g/min). f. Fat oxidation (g/min).

Measurements displayed as mean \pm S.D. with individual data points for all participants.

Forwards – ▲ (filled black triangle), Backs- ○ (empty circle).

*Denotes significant difference ($p < 0.05$) for the whole group when compared to GD-1.

+Denotes significant difference ($p < 0.05$) for the forwards group when compared to GD-1.

Table 3. Absolute and adjusted measurements of RMR across the competitive microcycle for all players ($n=17$).

Table 4. Spearman's coefficient (r_s) associations derived from changes in RMR between GD-1 and GD+1.

*denotes significant $p < 0.05$ association

Time Point	GD-3	GD-2	GD-1	GD	GD+1	GD+2	GD+3	GD+4
Purpose	Rest & Recovery	Intensity	Team Run	Match Play	Rest & Recovery	Installation	Volume	Rest & Recovery
Resistance Training Content	None	Upper Limb Strength (30 min)	None	None	None	Lower Limb Strength (45 mins)	Upper Limb Strength (45min)	None
Rugby Content	None	Specific Game Prep (35 mins) Unit Split (15-25mins)	Agility warm-up, Execution of specific game prep at a low-moderate intensity (35min)	Individual & Team Warm Ups. Rugby Match Play (80 mins).	None	Low-moderate intensity attack shapes and defensive systems. Running top-ups for some players. (60 mins)	High Intensity throughout rugby specific drills. Units Split- Forwards - Scrum/Maul Backs - Strike plays. (75 min)	None
Targets	Recovery	Execution of tactical game specifics at a high intensity. Rehearsal of set pieces in a unit split. Forwards – Lineouts Backs – Strike and skill execution.	Execution of specifics at a lower intensity, low intensity unit rehearsal of set pieces. Forwards – Lineouts Backs – Strike plays.	Full competitive rugby match play. Target physical performance and win.	Recovery	Learning of specifics for the following fixture and recovery.	High running volume, aiming to overload running volume relative to time.	Recovery

Table 1. The training sessions throughout the competitive micro cycle. Game Day – GD.

Time Point		GD -2	GD -1	GD	GD +1	GD +2	GD +3	GD +4
Player Load (sRPE x Time)	Whole group	404.07± 103.88*	27.00± 34.52*	622.36± 98.70	Rest & Recovery	238.14± 186.90*	631.07± 110.67	Rest & Recovery
	Forwards	391.25± 83.93*	31.25± 35.65*	595.13± 106.45		243.13± 212.10*	654.37± 100.19	
	Backs	421.17± 132.60*	21.33± 35.39*	658.67± 81.85		231.50± 166.58*	600.00± 125.57	
High Speed Running Distance (m)	Whole group	168.00± 60.12*	6.21± 10.15*	285.43± 113.09		63.29± 95.43*	254.64± 214.15	
	Forwards	132.88± 45.89	9.25± 12.10*	215.88± 96.82		72.00± 102.86*	279.63± 268.19	
	Backs	214.83± 42.94*	2.17± 5.31*	378.17± 45.46		51.67± 92.63*	221.33± 127.33*	
High Speed Running Efforts (n)	Whole group	11.43± 3.34*	0.93± 1.69*	20.29± 7.23		4.93± 6.86*	14.79± 8.91*	
	Forwards	10.50± 3.55	1.38± 2.07*	16.63± 6.99		5.63± 7.05*	15.50± 10.90	
	Backs	12.67± 2.88*	0.33± 0.82*	25.17± 4.17		4.00± 7.13*	13.83± 6.18*	
Very High-Speed Running Distance (m)	Whole group	17.86± 16.28	0.00*	16.50± 15.89		0.36± 0.93*	23.29± 31.10	
	Forwards	9.88± 10.64	0.00	10.75± 16.46		0.38± 1.06	23.75± 38.99	
	Backs	28.50± 17.10	0.00*	24.17± 12.42		0.33± 0.82*	22.67± 19.66	
Very High-Speed Running Efforts (n)	Whole group	1.29± 0.91	0.00*	1.14± 1.17		0.14± 0.36*	1.29± 1.14	
	Forwards	1.13± 0.99	0.00	1.00± 1.51		0.13± 0.35	1.25± 1.28	
	Backs	1.50± 0.84	0.00*	1.33± 0.52		0.17± 0.41*	1.33± 1.03	

Table 2. Comparison of metrics recorded for training and match play throughout the competitive micro cycle.

*Denotes values significantly different (p<0.05) when compared with **game day (GD)** shown in bold.

Time Point	GD-2	GD-1	GD+1	GD+2	GD+3	GD+4
Absolute RMR (kcal)	2318± 182.1	2313± 283.0	2544± 396.9	2391± 274.2	2424± 312.0	2327± 305.3
Adjusted RMR (kcal·kg·day⁻¹)	31.2± 3.0	31.1± 4.7	34.1± 5.3	32.1± 4.1	32.5± 4.2	31.3± 4.7

Table 3. Absolute and adjusted measurements of RMR across the competitive microcycle for all players (n=17).

Timepoints for comparison	Group	Phase Contacts		Total Contacts (Phase+set piece)		Player Load (sRPE×Time)		HSR (m)		HSR (efforts)		VHSR (m)		VHSR (efforts)	
		r_s	p	r_s	p	r_s	p	r_s	p	r_s	p	r_s	p	r_s	p
Change in RMR GD-1 to GD+1	Whole Group	0.05	0.84	0.23	0.38	-0.17	0.95	-0.13	0.62	-0.26	0.31	-0.11	0.97	0.19	0.48
	Forwards	-0.10	0.77	0.16	0.63	-0.19	0.58	-0.28	0.40	-0.24	0.47	0.19	0.57	0.32	0.34
	Backs	0.09	0.87	0.09	0.87	0.34	0.51	0.37	0.47	-0.44	0.39	-0.09	0.87	0.00	1.00

Table 4. Spearman's coefficient (r_s) associations derived from changes in RMR between GD-1 and GD+1.

*denotes significant $p < 0.05$ association.




Figure 1. Gas exchange measurements across the microcycle. a. RMR ($\text{kcal} \cdot \text{kg} \cdot \text{day}^{-1}$). b. RER. c. VO_2 (L/min). d. VCO_2 (L/min). e. Carbohydrate (CHO) oxidation (g/min). f. Fat oxidation (g/min).

Measurements displayed as mean \pm S.D. with individual data points for all participants.

Forwards – ▲ (filled black triangle), Backs- ○ (empty circle).

*Denotes significant difference ($p < 0.05$) for the whole group when compared to GD-1.

+Denotes significant difference ($p < 0.05$) for the forwards group when compared to GD-1.

