



LJMU Research Online

Sirichaiwetchakoon, K, Lowe, GM and Eumkeb, G

The Free Radical Scavenging and Anti-Isolated Human LDL Oxidation Activities of *Pluchea indica* (L.) Less. Tea Compared to Green Tea (*Camellia sinensis*).

<http://researchonline.ljmu.ac.uk/id/eprint/14077/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Sirichaiwetchakoon, K, Lowe, GM and Eumkeb, G (2020) The Free Radical Scavenging and Anti-Isolated Human LDL Oxidation Activities of *Pluchea indica* (L.) Less. Tea Compared to Green Tea (*Camellia sinensis*). *BioMed Research International*. 2020. ISSN 2314-6133

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Research Article

The Free Radical Scavenging and Anti-Isolated Human LDL Oxidation Activities of *Pluchea indica* (L.) Less. Tea Compared to Green Tea (*Camellia sinensis*)

Kittipot Sirichaiwetchakoon ¹, Gordon Matthew Lowe ², and Griangsak Eumkeb ¹

¹School of Preclinic, Institute of Science, Suranaree University of Technology, 111 University Avenue, Suranaree Subdistrict, Muang District, Nakhonratchasima 30000, Thailand

²School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool, UK

Correspondence should be addressed to Gordon Matthew Lowe; g.m.lowe@ljmu.ac.uk and Griangsak Eumkeb; griang@sut.ac.th

Received 21 July 2020; Revised 26 August 2020; Accepted 15 September 2020; Published 25 September 2020

Academic Editor: Roland E. Akhigbe

Copyright © 2020 Kittipot Sirichaiwetchakoon et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tea is one of the most popular beverages in the world. *Camellia sinensis* tea (CST) or green tea is widely regarded as a potent antioxidant. In Thailand, *Pluchea indica* (L.) Less. tea (PIT) has been commercially available as a health-promoting drink. This study focused on free radical scavenging activities of PIT, and its ability to protect isolated human low-density lipoproteins (LDL) from oxidation by chemical agents. A preliminary study to investigate the antioxidant nature of PIT was undertaken. These included common antioxidant assays involving 2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,2-azinobis-(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS), hypochlorous acid (HOCl), and its potential to scavenge peroxynitrite. In separated experiments, isolated human LDL was challenged with either 2,2'-azobis(2-amidinopropane) dihydrochloride (AAPH), copper (Cu^{2+}), or 3-Morpholinopyridone hydrochloride (SIN-1) to induce LDL oxidation. PIT exhibited antioxidant activity in all test systems and performed significantly better than CST in both DPPH ($P < 0.05$; $\text{IC}_{50}\text{PIT} = 245.85 \pm 15.83$ and $\text{CST} = 315.41 \pm 24.18 \mu\text{g/ml}$) and peroxynitrite scavenging assays. PIT at $75 \mu\text{g/ml}$ almost fully prevented the peroxynitrite over a 5 h period. Moreover, it displayed similar properties to CST during the antioxidantation of isolated human LDL using AAPH, Cu^{2+} , SIN-1, and hypochlorous acid scavenging assays. However, it revealed a significantly lower ABTS scavenging activity than CST ($P < 0.05$; $\text{IC}_{50}\text{PIT} = 30.47 \pm 2.20$ and $\text{CST} = 21.59 \pm 0.67 \mu\text{g/ml}$). The main constituents of the PIT were identified using LC-MS/MS. It contained 4-O-caffeoylquinic acid (4-CQ), 5-O-caffeoylquinic acid (5-CQ), 3,4-O-dicaffeoylquinic acid (3,4-CQ), 3,5-O-dicaffeoylquinic acid (3,5-CQ), and 4,5-O-dicaffeoylquinic acid (4,5-CQ). In conclusion, caffeoyl derivatives in PIT could play an important role in potent antioxidant properties. So, it may be further developed to be antioxidant beverages for preventing atherosclerosis and cardiovascular diseases associated with oxidative stress.

1. Introduction

Oxidative stress is defined as the imbalance between the production of free radicals and defense mechanisms, which are natural physiological processes in biological systems [1]. The excesses of intracellular reactive oxygen species (ROS) and reactive nitrogen species (RNS) is the major causes of oxidative stress that is associated with the development of

chronic and degenerative diseases such as cancer, arthritis, aging, autoimmune disorders, cardiovascular, and neurodegenerative diseases [2–5]. Moreover, oxidative stress may also modify the structure and function of certain biomolecules, including proteins, lipids, and DNA [6].

Oxidative stress may also result in the oxidation of human LDL. LDL oxidation may result in lipid peroxidation or the direct oxidation of apolipoprotein. Oxidized LDL is

thought to have a vital role in the etiology of atherosclerosis, which ultimately has a profound effect on cardiovascular function [7–9].

Herbal supplements derived from fruit and vegetables tend to be rich in both water and lipid-soluble antioxidants [10]. One of the most popular herbal supplements is a beverage, such as a tea infusion. Many herbal tea infusions have a strong antioxidant capacity [11–14], and it has the potential to prevent diseases associated with oxidative stress such as atherosclerosis [15, 16].

Tea is the most widely consumed beverage in the world, second only to water [17], and one of the most popular beverages in Southeast Asia. Green tea is derived from the tea plant *Camellia sinensis*. It has been demonstrated that some of the components of green tea have potent antioxidative properties and have free radical scavenging properties towards the DPPH, ABTS, and Fluorescence Recovery After Photobleaching (FRAP) assays [18]. Also, flavonols from green tea have potent antioxidant capacities and reduced oxidative stress [19]. Moreover, green tea can prevent lipid oxidation induced by copper ions [20]. The main active ingredients of green tea are polyphenol catechins. The major catechins present in green tea are Epicatechin (EC), Epigallocatechin gallate (EGCG), Epicatechin gallate (ECG), Epigallocatechin (EGC), and Gallic catechin gallate (GCG) [21]. EGCG, which acts as an antioxidant, might exert a preventive effect against cardiovascular disease [22].

The plant *Pluchea indica* (L.) Less. (*P. indica*) is a large evergreen shrub found abundantly in salt marshes. It is widely distributed in India, Southern China, and Southeast Asia. In Thailand, The PIT has been commercially available for approximately ten years as a health-promoting drink [23]. The various biological activities of *P. indica* leaves have been widely reported. Several studies revealed that its methanolic extract had diuretic [24], hypoglycemic, and antihyperglycemic effects [25]. Moreover, the ethyl acetate fraction from an ethanolic extract of *P. indica* in lipopolysaccharide (LPS-) stimulated RAW 264.7 macrophages displayed anti-inflammatory activities [26]. Besides, the volatile oil of *P. indica* exhibited antioxidant activity [27]. Although some biological activities of *P. indica* have been reported, there is little information published on the antioxidative properties of the aqueous tea.

The aim of this study was to examine the free radical scavenging potential of PIT with regard to established antioxidant assays and its potential to inhibit the oxidation of isolated human LDL by Cu^{2+} , AAPH, or SIN-1. The antioxidant properties of PIT were compared to a commercially available green tea.

2. Materials and Methods

2.1. Tea Materials. PIT was supplied by the Crystal Biotechnology Company, Thailand. Commercial green tea was purchased from a supermarket in the United Kingdom. The tea samples were kept in the dark, dry place until required. Beverages were prepared by brewing ground tea leaves in 80°C 1x phosphate buffer saline (PBS) for 5 min and filtered by What-

man No. 1 filter paper. The concentration of PIT was calculated from the ground tea leaves dry weight in PBS volume ($\mu\text{g}/\text{ml}$). Tea samples were kept at -20°C until used.

2.2. Chemicals and Reagents. 2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,2-azinobis-(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS), potassium persulfate, sodium hypochlorite (NaOCl), catalase, sulfanilamide, naphthylethylenediamine dihydrochloride (NED), 3-Morpholinolonylamine hydrochloride (SIN-1), Diethylene-triamine-pentaacetic acid (DTPA), Evans blue, iodixanol (Optiprep™), 2,2'-azobis (2-amidinopropane) dihydrochloride (AAPH), copper sulfate (Cu_2SO_4), trichloroacetic acid (TCA), thiobarbituric acid were purchased from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). 4-O-caffeoylquinic acid (4-CQ), 5-O-caffeoylquinic acid (5-CQ), 3,4-O-dicaffeoylquinic acid (3,4-CQ), 3,5-O-dicaffeoylquinic acid (3,5-CQ), and 4,5-O-dicaffeoylquinic acid (4,5-CQ) were purchased from Chengdu Biopurify Phytochemicals Ltd., China. Other reagents used were all analytical grade.

2.3. DPPH Scavenging Assay. The %DPPH scavenging activity of PIT and CST were evaluated by the method of Brand-Williams et al. [28] with slight modifications. In brief, 0–300 $\mu\text{g}/\text{ml}$ of the tea extracts was prepared using 0.002% DPPH reagent in a 96-well plate. The samples were suitably mixed prior to incubation in the dark at room temperature for 30 min. PBS was used as a negative control. The absorbance of all samples was measured at 515 nm. Radical scavenging activity was repeated for six times and expressed as a DPPH scavenging percentage as,

$$\% \text{DPPH scavenging} = \left(1 - \frac{\text{OD}_{\text{sample}} - \text{OD}_{\text{sample blank}}}{\text{OD}_{\text{control}} - \text{OD}_{\text{sample blank}}} \right) \times 100. \quad (1)$$

2.4. ABTS Scavenging Assay. The ABTS scavenging activity of PIT was measured and compared with CST. The ABTS^+ radical cation was prepared by mixing 7 mM ABTS stock solution with 2.45 mM potassium persulfate (final concentration) in methanol and kept them in the dark at room temperature for 16 h. The solution was adjusted to an absorbance of 0.70 (± 0.02) with ethanol. The reaction was performed in a 96-well plate by mixing and shaking 90 μl ABTS^+ radical cation solution with 10 μl of the test sample at various concentrations (0–40 $\mu\text{g}/\text{ml}$ final concentration) for 45 seconds. The absorbance was measured at 734 nm. The assay was repeated six times, and the percentage inhibition of absorbance was reported as an ABTS scavenging percentage as

$$\% \text{ABTS scavenging} = \left(1 - \frac{\text{OD}_{\text{sample}} - \text{OD}_{\text{sample blank}}}{\text{OD}_{\text{control}} - \text{OD}_{\text{sample blank}}} \right) \times 100. \quad (2)$$

2.5. Hypochlorous Acid (HOCl) Scavenging Assay. This assay was performed as previously described by Aruoma and Halliwell [29] with minor modifications. HOCl was generated by

adjusting the pH of a 10% (*v/v*) solution of NaOCl to 6.2% with 0.6 M H₂SO₄. The HOCl was mixed with 50 mM phosphate buffer (pH 6.8), catalase (7.2 μM), and tea samples at various concentrations (0–300 μg/ml) in 96 well-plate. The mixture was incubated at 25°C for 20 min. The scavenging activity was measured by the decrease in absorbance of catalase at 404 nm. The assay was performed in triplicate. %HOCl scavenging was calculated as

$$\% \text{HOCl scavenging} = \left(1 - \frac{\text{OD}_{\text{sample blank}} - \text{OD}_{\text{sample}}}{\text{OD}_{\text{sample blank}} - \text{OD}_{\text{control}}} \right) \times 100. \quad (3)$$

2.6. Nitric Oxide (NO) Scavenging Assay. SIN-1 was used for generating RNS in this experiment. One of the end products of RNS chemistry is nitrite. Nitrite concentration was determined using the Griess assay. The test solution was prepared in a 96-well plate by mixing 0.25 mM SIN-1 with PBS (pH 7.4) and various doses of tea samples (0–100 μg/ml final concentration). The test solutions were incubated at 25°C for 30 min. Then, 80 μl of 0.33% sulfanilamide in 20% glacial acetic acid was added to the produced solution and was suitably shaken for 5 min before adding 80 μl of 0.1% NED and incubated at 25°C for 15 min. The nitric oxide scavenging was measured spectrophotometrically at 540 nm against a blank sample. All tests were repeated six times, and %Nitric oxide radical scavenging was expressed as

$$\% \text{Nitric oxide radical scavenging} = \left(1 - \frac{\text{OD}_{\text{sample}} - \text{OD}_{\text{sample blank}}}{\text{OD}_{\text{control}} - \text{OD}_{\text{sample blank}}} \right) \times 100. \quad (4)$$

2.7. Peroxynitrite Scavenging Assay. SIN-1 was used as a peroxynitrite donor, and peroxynitrite scavenging activity was measured by an Evans blue bleaching assay. Briefly, the reaction mixture comprised of 50 mM phosphate buffer (pH 7.4), 0.1 mM DTPA, 90 mM NaCl, 5 mM KCl, 12.5 μM Evans blue, 1 mM SIN-1, 37.5, and 75 μg/ml of

either PIT or CST in a 96-well plate. The mixture was incubated at 37°C for 300 min, and the absorbance was measured at 611 nm every 30 min. The assay was performed six times. The percentage scavenging of peroxynitrite at various times was calculated by the %Optical density of Evans blue as;

$$\% \text{Optical density of Evans blue} = \left(1 - \frac{\text{OD}_{\text{sample blank}} - \text{OD}_{\text{sample}}}{\text{OD}_{\text{sample blank}} - \text{OD}_{\text{control}}} \right) \times 100. \quad (5)$$

2.8. LDL Isolation by Ultracentrifugation Technique. The Liverpool John Moores University (LJMU) ethics committee approved the use of human blood and the preparation of LDL. Whole blood was obtained from the vein of healthy volunteers aged between 24 and 70 years who were normolipidemic, nonsmoking, had not taken any medications or supplements within the last two weeks. Whole blood was added to 3.8% (*w/v*) sodium citrate at a ratio of blood against anticoagulant as 9:1. The whole blood was centrifuged at 1500 x g for 20 min, and the platelet-poor plasma was transferred to a separate plastic tube. LDL was isolated by density gradient ultracentrifugation using a method developed by Graham et al. (1996) with minor modifications [30]. In brief, 0.5 ml of 60% (*v/v*) iodixanol (Optiprep™) was mixed with 4.5 ml of plasma and transferred to an 11.2 ml capacity Optiseal tube. Then, it was overlaid with 5 ml of 12% *v/v* iodixanol prepared with human plasma, and PBS was used to fill the rest of the tube. The tubes were centrifuged in a Beckman L8-80 ultracentrifuge using a vertical rotor V65.1 at 350,000 x g for 3 h. An Auto Densi-Flow gradient fractionator (Labconco, UK) was used for fractionating

the gradient by unloading the gradient from the top to the bottom of the tube. Each tube of 11.2 ml was fractionated into 0.5 ml aliquots per tube. All fractions were measured for triglycerides, cholesterol, LDL, and apoB100 components by using reagents and standards from Randox (Eire). Pooled LDL fraction was measured protein concentration by Bradford assay and stored at -20°C until required.

2.9. AAPH Induce LDL Oxidation Assay. AAPH is a reactive oxygen species generator. The sample solution was prepared by adding 20 mM AAPH to 50 μg/ml of LDL protein. Either 50 μg/ml or 75 μg/ml of PIT or CST was added in the test samples. The control sample was 50 μg/ml of LDL protein added with PBS. The LDL protein was added with AAPH 20 mM. All samples were incubated at 37°C. At this temperature, AAPH produces a range of ROS. At a specific time point (0, 30, 60, 90, 120, 150, and 180 min), sample aliquots were removed and placed at -20°C prior to thiobarbituric acid reactive substances (TBARS) measurements. The first stage of the TBARS assay was to precipitate protein by the addition of 20% *v/v* TCA. The samples were then spun at

10,000 rpm for 10 min at 4°C. The supernatant was collected and treated with 1% *w/v* thiobarbituric acid. The samples were heated at 95°C for 20 min. TBARS concentration was determined by UV absorption at 532 nm. The malondialdehyde (MDA) concentration was determined by a calibration curve. The assay was performed in triplicate.

2.10. Copper Induce LDL Oxidation Assay. LDL protein was challenged with Cu^{2+} by adding Cu_2SO_4 at a final concentration of 40 μM to 50 $\mu\text{g/ml}$ of LDL protein. The negative control sample was 50 $\mu\text{g/ml}$ of LDL protein with PBS. The positive control was 50 $\mu\text{g/ml}$ of LDL protein incubated with 40 μM Cu_2SO_4 . The 15 $\mu\text{g/ml}$ of PIT or CST was added in the test samples that contained 50 $\mu\text{g/ml}$ LDL protein plus 40 μM Cu_2SO_4 . All samples were incubated at 37°C for 3 h. At various time points, the samples were assessed for MDA equivalence as TBARS products. The assay was performed in triplicate.

2.11. SIN-1 Induce LDL Oxidation Assay. SIN-1, which is a peroxynitrite donor, has been used to investigate RNS-mediated LDL oxidation. In this assay, 50 $\mu\text{g/ml}$ of LDL protein was challenged with 1 mM of SIN-1. In test samples, 15 $\mu\text{g/ml}$ of PIT or CST was mixed with 50 $\mu\text{g/ml}$ of LDL protein and 1 mM of SIN-1. The samples were incubated at 37°C for 18 h. Finally, the samples were stored at -20°C before further analysis by TBARS assay. The assay was repeated three times.

2.12. LC-MS/MS Instrument and Conditions. LC-MS/MS technique has been used for identifying the main chemical constituents of PIT as previously described by Kongkiatpaiboon et al. [23]. The combination of chromatographic separation of LC-MS/MS system was combined by Agilent HPLC 1290 Infinity and mass analyzer 6490 Triple Quad LC/MS Agilent Technologies, which equipped with an electrospray ionization (ESI) source system. Agilent ZORBAX Rapid Resolution High Definition (RRHD) SB-C18, 2.1 mm id \times 150 mm (1.8 μm) was used for chromatographic separation. The mobile phase system used 1% formic acid in water as solvent A and 1% formic acid in acetonitrile as solvent B. The gradient of the mobile phase was set at a ratio of solvent A : solvent B, 100 : 0, with gradient elution, from 30% solvent B at 10 min and 100% solvent B at 30 min at a flow-rate of 0.2 ml/min. The sample injection volume was 5 μL , and the column was set at 25°C. 4-CQ, 5-CQ, 3,4-CQ, 3,5-CQ, and 4,5-CQ were used as standard.

2.13. Statistical Analysis. All the data were presented as a mean \pm standard deviation (S.D.) or standard error of the mean (S.E.M.). Statistical analysis was performed using SPSS version 18.0. The significant statistical differences between groups of DPPH, ABTS, HOCl, and NO scavenging assay were analyzed by an independent *t*-test, whereas peroxynitrite, Cu^{2+} , AAPH, and SIN-1 scavenging assay were compared by one-way analysis of variance (ANOVA) with a Tukey's HSD post hoc test. Values were considered statistically significant when $P < 0.05$, and data were the representative of at least three independent experiments.

3. Results

3.1. DPPH Scavenging Assay. The activities of both teas were investigated using the DPPH radical scavenging assay. The antioxidant activity was calculated spectrophotometrically at 515 nm. The result indicated that PIT generated significantly stronger antioxidant capacity than CST at a concentration of 75 $\mu\text{g/ml}$ to 300 $\mu\text{g/ml}$ CST ($P < 0.05$) (Figure 1). At a concentration of 300 $\mu\text{g/ml}$, the %DPPH scavenging value of PIT and CST was 51.19 ± 4.02 and 41.46 ± 3.83 , respectively. The IC_{50} value of PIT was determined at 245.85 ± 15.83 $\mu\text{g/ml}$, which was lower than CST at 315.41 ± 24.18 $\mu\text{g/ml}$.

3.2. ABTS Scavenging Assay. This assay shows the abilities of the extract to quench the ABTS^+ radical. The extracts interacted with ABTS^+ , which decreased the absorbance of the solution. The absorbance was measured spectrophotometrically at 734 nm. Figure 2 displays %ABTS radical scavenging of PIT and CST. CST significantly inhibited ABTS^+ radical stronger than PIT at 5-40 $\mu\text{g/ml}$ ($P < 0.05$). The IC_{50} of PIT and CST was 30.47 ± 2.20 and 21.59 ± 0.67 $\mu\text{g/ml}$, respectively.

3.3. Hypochlorous Acid Scavenging Assay. The hypochlorous acid scavenging activity of the PIT and CST are shown in Figure 3. The results showed that PIT significantly scavenged more hypochlorous acid than CST at a concentration of 18.75 $\mu\text{g/ml}$ ($P < 0.05$). Nevertheless, the other strengths of PIT inhibited more hypochlorous acid than CST but not significantly.

3.4. Nitric Oxide Scavenging Assay. Nitric oxide scavenging could be detected by determining the nitrite concentration using the decolorization by the Griess reaction method. Figure 4 shows % inhibition of nitric oxide of PIT and CST. The result indicated that the %nitric oxide scavenging activity of PIT was significantly higher than CST ($P < 0.01$). The IC_{50} of PIT was 116.48 ± 5.08 $\mu\text{g/ml}$, while the IC_{50} of CST was 178.42 ± 15.52 $\mu\text{g/ml}$.

3.5. Peroxynitrite Scavenging Assay. Evans blue assay is used to measure peroxynitrite scavenging, which is generated by SIN-1. The peroxynitrite is thought to bleach the Evans blue dye. The results are shown in Figure 5. At higher concentrations, PIT (75 $\mu\text{g/ml}$) almost fully prevented the peroxynitrite bleaching the dye over a 5 h period. Whereas at 37.5 $\mu\text{g/ml}$, the %peroxynitrite scavenging was $64.50 \pm 8.07\%$ at 2 h. In contrast, a higher concentration of CST (75 $\mu\text{g/ml}$) exhibited significantly lower peroxynitrite scavenging activity compared to PIT.

3.6. AAPH Induce LDL Oxidation Assay. The ROS induce LDL oxidation scavenging effect has been investigated by challenging isolated human LDL with AAPH, which is ROS. PIT and CST at 50 and 75 $\mu\text{g/ml}$ were used for the experiment. The samples were incubated for 3 h, and aliquots were collected at various times for analysis using the TBARS assay. The result indicated that after 60 min of AAPH activity, both teas could significantly decrease the TBARS formation (Figure 6). Furthermore, a stronger effect was observed




FIGURE 1: The DPPH radical scavenging activity. PIT: *P. indica* (L.) Less. tea; CST: *C. sinensis* tea at concentrations ranging from 0-300 µg/ml. The data represent the percentage of DPPH inhibition. Each value represents mean ± S.D. (n = 6). ** indicates a significant difference between groups at P < 0.01.




FIGURE 2: The ABTS radical scavenging activity. PIT: *P. indica* (L.) Less. tea; CST: *C. sinensis* tea at concentrations ranging from 0-40 µg/ml. The data represent the percentage of ABTS inhibition. Each value represents mean ± S.D. (n = 6). ** indicates a significant difference between groups at P < 0.01.

at higher concentrations of both teas. The lag time of LDL oxidation was increased from 40 to 70 min at all strengths of testing teas. These results suggest that both teas have approximately similar LDL oxidation scavenging properties.

3.7. *Copper Induce LDL Oxidation Assay.* Isolated human LDL was incubated with either PIT or CST at 15 µg/ml and

challenged with Cu²⁺. The result was presented as MDA equivalence (nmol/mgprotein), which was calculated from the MDA calibration standard curve. Figure 7 showed that in the Cu²⁺-treated group, the MDA equivalence was continually increased by 210 min at 222.73 ± 2.22 nmol/mgprotein, while the PIT and CST treated group could almost entirely prevent Cu²⁺ oxidation of human LDL.




FIGURE 3: The hypochlorous acid radical scavenging activity. PIT: *P. indica* (L.) Less. tea; CST: *C. sinensis* tea at concentrations ranging from 0-300 µg/ml. The data represent the percentage of HOCl inhibition. Each value represents mean \pm S.E.M ($n = 3$). * indicates a significant difference between groups at $P < 0.05$.




FIGURE 4: The nitric oxide radical scavenging activity. PIT: *P. indica* (L.) Less. tea; CST: *C. sinensis* tea at concentrations ranging from 0-150 µg/ml in various concentrations. The data represent the percentage of nitric oxide inhibition. Each value represents mean \pm S.D. ($n = 6$). ** indicates a significant difference between groups at $P < 0.01$.

3.8. *SIN-1 Induce LDL Oxidation Assay.* SIN-1, which is an RNS generator, is used to induce LDL oxidation. Isolated human LDL was incubated with SIN-1 and tea extracts. The result expressed the RNS scavenging effect of both

PIT and CST at a dose of 15 µg/ml (Figure 8). PIT and CST displayed nearly similar properties and could decrease MDA concentration from 86.73 ± 2.55 to 20.41 ± 2.55 and 23.81 ± 3.90 nmol/mgprotein, respectively.




FIGURE 5: The peroxynitrite radical scavenging activity. PIT(75): *P. indica* (L.) Less. tea at 75 $\mu\text{g/ml}$; CST(75): *C. sinensis* tea at 75 $\mu\text{g/ml}$. The data represent the percentage of the optical density of Evans blue at 611 nm at various times over 5 h. Means \pm S.E.M is illustrated for six replicates. Means with the same superscript are not significantly different from each other (Tukey's HSD test, $P < 0.05$).




FIGURE 6: The AAPH induces LDL oxidation scavenging activity. LDL(AAPH): LDL-treated with 20 mM of AAPH; PIT(75): *P. indica* (L.) Less. tea at 75 $\mu\text{g/ml}$; CST(75): *C. sinensis* tea at 75 $\mu\text{g/ml}$. The data represent the MDA equivalence at various times over 3 h. Means \pm S.D. is illustrated for three replicates. Means with the same superscript are not significantly different from each other (Tukey's HSD test, $P < 0.05$).

3.9. LC-MS/MS Analysis of PIT. In this experiment, ESI-MS analysis used negative ion mode and identified the chemical constituents by comparing the profiles with authentic stan-

dards using the Multiple Reaction Monitoring (MRM) modes. Two pairs of MRM transitions were selected at m/z 353.1 \rightarrow 191.0 and 515 \rightarrow 353. Table 1 displayed that 4-CQ,




FIGURE 7: The copper induces LDL oxidation scavenging activity. LDL(Cu²⁺): LDL-treated with 40 μ M of copper sulfate; PIT(15): *P. indica* (L.) Less. tea at 15 μ g/ml; CST(15): *C. sinensis* tea at 15 μ g/ml. The data represent the MDA equivalence at various times over 5 h. Means \pm S.D. is illustrated for three replicates. Means with the same superscript are not significantly different from each other (Tukey's HSD test, $P < 0.05$).




FIGURE 8: The SIN-1 induces LDL oxidation scavenging activity. LDL(SIN-1): LDL-treated with 1 mM of SIN-1; PIT: *P. indica* (L.) Less. tea at 15 μ g/ml; CST: *C. sinensis* tea at 15 μ g/ml. The data represent the MDA equivalence at 18 h. Means \pm S.D. is illustrated for three replicates. Means with the same superscript are not significantly different from each other (Tukey's HSD test, $P < 0.05$).

5-CQ, 3,4-CQ, 3,5-CQ, and 4,5-CQ were detected in the PIT (Table 1). Moreover, the results showed a concentration of 3,5-CQ, which was the highest peak of the chromatogram at 169.93 μ g/ml of 1,500 μ g/ml PIT (data not showed).

4. Discussion

Free radicals encompassing the ROS and RNS are derived from both endogenous sources (mitochondria, peroxisomes, endoplasmic reticulum, phagocytic cells, etc.) and exogenous sources (pollution, alcohol, tobacco smoke, heavy metals,

transition metals, industrial solvents, pesticides, certain drugs like halothane, paracetamol, and radiation) [31]. The imbalance between free radical and antioxidant systems can cause extensive damage to tissues and biomolecules [32], leading to various diseases especially degenerative diseases of aging such as cancer, immune-system decline, brain dysfunction, and cardiovascular [33]. Antioxidants derived from the diet assist physiological protective mechanisms in preventing damage from ROS or RNS. Dietary antioxidants can be derived from either supplements, fruit, vegetables, or herbal beverages, including teas. These are popular with

TABLE 1: The main chemical constituent of *P. indica* tea. was analyzed by Liquid Chromatography-Mass Spectrometer/Mass Spectrometer (LC-MS/MS).

Main chemical constituent	Detection
4-O-caffeoylquinic acid (4-CQ)	+
5-O-caffeoylquinic acid (5-CQ)	+
3,4-O-dicaffeoylquinic acid (3,4-CQ)	+
3,5-O-dicaffeoylquinic acid (3,5-CQ)	+
4,5-O-dicaffeoylquinic acid (4,5-CQ)	+

consumers and are widely used to prevent the diseases generated by free radicals [34, 35].

PIT has been used for health-promoting tea, but its antioxidative properties have not been fully explored. In this study, the antioxidative properties of PIT were compared with a well-known and commercially available green tea. The properties were investigated using well-established techniques, including DPPH and ABTS. The properties of the tea to scavenge peroxynitrite, RNS, and hypochlorous acid and prevent the oxidation of isolated human LDL were also undertaken.

DPPH radical scavenging assay has been widely used in the determination of the antioxidant activity of natural antioxidants from plant sources [28, 36]. This assay determines the reduction of DPPH radical by measuring the color changing from the violet color of DPPH radical to yellow of the nonradical DPPH derivative at 515 nm. Several studies indicated that CST scavenged the DPPH radical [37, 38]. Interestingly, PIT showed significantly stronger antioxidant activity in this assay than CST at all concentrations. These results are in substantial agreement with Srisook et al. [39] that hot water extract of *P. indica* leaves shows the DPPH radical scavenging activity (EC_{50} value = $23.8 \pm 1.0 \mu\text{g/ml}$).

ABTS⁺ radical cation decolorization assay was used to measure the antioxidant capacity of PIT compared to CST. These results provide evidence that CST could significantly reduce ABTS⁺ radical better than PIT. This higher action of CST may come from various classes of polyphenols in CST, which act as a potent antioxidant for the ABTS⁺ radical [40, 41].

Hypochlorous acid is a weak acid that could inactivate the antioxidant enzyme catalase by breaking down the heme prosthetic group [42]. Our results demonstrated that PIT was a higher potent hypochlorous acid scavenger than CST, which has been previously reported its hypochlorous acid scavenging capacity [43, 44].

Nitric oxide has an important role in several physiological processes like neural signal transmission, immune response, control vasodilation, and control of blood pressure [45, 46]. Nevertheless, the elevation of the nitric oxide causes inflammation and sustained levels of nitric oxide results in tissue toxicity and several pathological, including in vascular disease [47]. The present study examined the nitric oxide scavenging effect of PIT. Tsai et al. reported that CST had IC_{50} values of nitric oxide scavenging less than $500 \mu\text{g/ml}$ and was proven to be a good nitric oxide suppressor [48].

Interestingly, our findings provide evidence that PIT displays significantly higher nitric oxide scavenging activity than CST.

Peroxynitrite, which is one of the nitrogen-containing species, is indicated as RNS. Excess peroxynitrite represents a crucial pathogenic mechanism in conditions, such as stroke, myocardial infarction, chronic heart failure, diabetes, circulatory shock, chronic inflammatory diseases, cancer, and neurodegenerative disorders [49]. PIT at a concentration of $75 \mu\text{g/ml}$ had the capacity to fully inhibit the ability of peroxynitrite to bleach the color of Evan blue dye. Noticeably, PIT demonstrated significantly better peroxynitrite scavenging activity than the green tea preparation at the same concentration of $75 \mu\text{g/ml}$. Chung et al. reported that catechins, a galloyl group containing in green tea, inhibited peroxynitrite formation by both SIN-1 and scavenged peroxynitrite itself [50]. These findings lead us to believe that the effect of PIT on DPPH, ABTS, hypochlorous acid, nitric oxide, and peroxynitrite scavenging is better than CST.

LDL lipid oxidation is considered to be essential in the pathogenesis of atherosclerotic vascular diseases [51]. Several lines of evidence suggest that the important mechanisms of LDL lipid oxidation occur by ROS, RNS, and Cu^{2+} [52]. The natural compounds with anti-LDL oxidation activity could have some beneficial effects in the prevention of the disease [53, 54]. The investigation of anti-LDL oxidation activity *in vivo* can be measured *in vitro* by using whole plasma/serum [55]. In this study, we measured the LDL oxidation in AAPH, Cu^{2+} , and SIN-1 challenged isolated human LDL using the TBARS assay and presented the results by MDA equivalence. AAPH is a ROS generator that can initiate lipid peroxidation and protein oxidation in isolated LDL particles. One measure of antioxidative protection in LDL is the lag-time. The inclusion of both PIT and CST extended the lag time compared to the control. From the results, the antioxidative properties of CST observed toward AAPH activity are consistent with those of Liu et al. reported that the polyphenolic components derived from green tea leaves are effective antioxidants against AAPH-initiated photosensitized LDL oxidation [56]. Noticeably, our studies found that PIT demonstrated better AAPH scavenging activity than green tea from 150 min onward. Besides, the Cu^{2+} scavenging activity effect of both PIT and CST was also investigated. The results expressed that PIT could favorably inhibit Cu^{2+} induced LDL oxidation compared to CST. It was reported by Yokozawa et al. that green tea extract markedly delayed Cu^{2+} induced LDL oxidation with a dose-dependent pattern [57]. Green tea contains a rich array of polyphenols, and these may chelate the copper ions from solution, the same may be true for PIT. Moreover, SIN-1, which is an RNS generator, was also used to investigate LDL oxidation. The RNS, which is generated by SIN-1, was scavenged by both teas. This result is consistent with the Evan blue dye study presented in this study. These results suggest that PIT may have a role in preventing the initial stages of atherogenic events by inhibiting ROS, RNS, and Cu^{2+} induce lipid peroxidation.

The PIT has been investigated as the main chemical constituent by using LC-MS/MS technique. The result expressed that caffeoylquinic acid derivatives were the main chemical compositions of PIT. This result is in correspondence with

the study from Kongkiatpaiboon et al. [23], which reported that PIT contained six caffeoylquinic acid derivatives. Interestingly, caffeoylquinic acid derivatives were reported that they had potent antioxidant properties [58]. Caffeoylquinic acid derivatives from *Dipsacus asper* Wall (Dipsacaceae) showed antioxidant activity against free radical and Cu²⁺-mediated LDL oxidation. They may have an essential role in preventing the development and progression of atherosclerotic disease [58]. Furthermore, these derivatives also showed high DPPH-radical and peroxynitrite scavenging activity [59, 60]. These findings provide evidence that the antioxidative nature, which is free radical scavenging activities and anti-LDL oxidation, of PIT, is better than or approximately equal CST. These effects may be acted by caffeoylquinic acid derivatives rather than polyphenolic catechins found in CST [12].

5. Conclusion

The results of the present work show free radical scavenging activities and anti-LDL oxidation effects of PIT. The PIT has the potential to be developed as a health supplement product to provide antioxidants for atherosclerosis or other diseases associated with oxidative stress prevention. Further studies should focus on *in vivo* investigation, including pharmacokinetics, pharmacodynamic, efficacious, and safe dose in humans.

Data Availability

The data used and analyzed in this study are available from the corresponding author on reasonable request.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

Authors' Contributions

KS performed the experiments and wrote the report. GE and GL designed the project, supervised the experiments, analyzed data, gave comments, and wrote the full manuscript. All authors have read and approved the final manuscript.

Acknowledgments

The authors are indebted and grateful to the Thailand Research Fund, Newton Fund (PhD Placement grant for Scholars), and the Crystal Biotechnology Co., Ltd for the assistance in research fund support through The Research and Researchers for Industries Ph.D. scholarships (Grant No. PHD58I0015 Code 5712035). Commenting on the report's findings is responsible for a grant recipient; the Thailand Research Fund, Newton Fund, and the Crystal Biotechnology Co., Ltd. are not necessarily always agreeing.

References

- [1] V. Dhawan, "Reactive oxygen and nitrogen species: general considerations," in *Studies on Respiratory Disorders*, pp. 27–47, Humana press, New York, 2014.
- [2] L. A. Pham-Huy, H. He, and C. Pham-Huy, "Free radicals, antioxidants in disease and health," *International Journal of Biomedical Science*, vol. 4, no. 2, p. 89, 2008.
- [3] H. Bartsch and J. Nair, "Chronic inflammation and oxidative stress in the genesis and perpetuation of cancer: role of lipid peroxidation, DNA damage, and repair," *Langenbeck's Archives of Surgery*, vol. 391, no. 5, pp. 499–510, 2006.
- [4] P. P. Tak, N. J. Zvaifler, D. R. Green, and G. S. Firestein, "Rheumatoid arthritis and p53: how oxidative stress might alter the course of inflammatory diseases," *Immunology Today*, vol. 21, no. 2, pp. 78–82, 2000.
- [5] E. R. Stadtman, "Role of oxidant species in aging," *Current Medicinal Chemistry*, vol. 11, no. 9, pp. 1105–1112, 2004.
- [6] S. Toyokuni, "Reactive oxygen species-induced molecular damage and its application in pathology," *Pathology International*, vol. 49, no. 2, pp. 91–102, 2002.
- [7] T. Bahorun, M. Soobrattee, V. Luximon-Ramma, and O. Aruoma, "Free radicals and antioxidants in cardiovascular health and disease," *Internet Journal of Medical Update*, vol. 1, no. 2, pp. 25–41, 2006.
- [8] P. Marchio, S. Guerra-Ojeda, J. M. Vila, M. Aldasoro, V. M. Victor, and M. D. Mauricio, "Targeting early atherosclerosis: a focus on oxidative stress and inflammation," *Oxidative Medicine and Cellular Longevity*, vol. 2019, 32 pages, 2019.
- [9] T. Yuan, T. Yang, H. Chen et al., "New insights into oxidative stress and inflammation during diabetes mellitus-accelerated atherosclerosis," *Redox Biology*, vol. 20, pp. 247–260, 2019.
- [10] J. L. Slavin and B. Lloyd, "Health benefits of fruits and vegetables," *Advances in Nutrition*, vol. 3, no. 4, pp. 506–516, 2012.
- [11] S. Li, S. K. Li, H. B. Li, X. R. Xu, G. F. Deng, and D. P. Xu, "Antioxidant capacities of herbal infusions," in *Processing and Impact on Antioxidants in Beverages*, pp. 41–50, 2014.
- [12] A. K. Atoui, A. Mansouri, G. Boskou, and P. Kefalas, "Tea and herbal infusions: their antioxidant activity and phenolic profile," *Food Chemistry*, vol. 89, no. 1, pp. 27–36, 2005.
- [13] B. Dimitrios, "Sources of natural phenolic antioxidants," *Trends in Food Science & Technology*, vol. 17, no. 9, pp. 505–512, 2006.
- [14] E. W. C. Chan, Y. Y. Lim, K. L. Chong, J. B. L. Tan, and S. K. Wong, "Antioxidant properties of tropical and temperate herbal teas," *Journal of Food Composition and Analysis*, vol. 23, no. 2, pp. 185–189, 2010.
- [15] A. Babu, V. Pon, and D. Liu, "Green tea catechins and cardiovascular health: an update," *Current Medicinal Chemistry*, vol. 15, no. 18, pp. 1840–1850, 2008.
- [16] Y. Miura, T. Chiba, I. Tomita et al., "Tea catechins prevent the development of atherosclerosis in apoprotein E-deficient mice," *The Journal of Nutrition*, vol. 131, no. 1, pp. 27–32, 2001.
- [17] S. I. Koo and S. K. Noh, "Green tea as inhibitor of the intestinal absorption of lipids: potential mechanism for its lipid-lowering effect," *The Journal of Nutritional Biochemistry*, vol. 18, no. 3, pp. 179–183, 2007.
- [18] S. Yan, H. Shao, Z. Zhou, Q. Wang, L. Zhao, and X. Yang, "Non-extractable polyphenols of green tea and their antioxidant, anti- α -glucosidase capacity, and release during *in vitro*

- digestion," *Journal of Functional Foods*, vol. 42, pp. 129–136, 2018.
- [19] C.-S. Rha, H. W. Jeong, S. Park, S. Lee, Y. S. Jung, and D.-O. Kim, "Antioxidative, anti-inflammatory, and anticancer effects of purified flavonol glycosides and aglycones in green tea," *Antioxidants*, vol. 8, no. 8, p. 278, 2019.
- [20] L. K. Leung, Y. Su, R. Chen, Z. Zhang, Y. Huang, and Z. Y. Chen, "Theaflavins in black tea and catechins in green tea are equally effective antioxidants," *The Journal of Nutrition*, vol. 131, no. 9, pp. 2248–2251, 2001.
- [21] C. Folch-Cano, C. Jullian, H. Speisky, and C. Olea-Azar, "Antioxidant activity of inclusion complexes of tea catechins with β -cyclodextrins by ORAC assays," *Food Research International*, vol. 43, no. 8, pp. 2039–2044, 2010.
- [22] K. Yamagata, "Protective effect of epigallocatechin gallate on endothelial disorders in atherosclerosis," *Journal of Cardiovascular Pharmacology*, vol. 75, no. 4, pp. 292–298, 2020.
- [23] S. Kongkiatpaiboon, S. Chewchinda, and B. Vongsak, "Optimization of extraction method and HPLC analysis of six caffeoylquinic acids in *Pluchea indica* leaves from different provenances in Thailand," *Revista Brasileira de Farmacognosia*, vol. 28, no. 2, pp. 145–150, 2018.
- [24] K. C. Pramanik, R. Biswas, A. Mitra, D. Bandyopadhyay, M. Mishra, and T. K. Chatterjee, "Tissue culture of the plant *Pluchea indica* (L.) Less. and evaluation of diuretic potential of its leaves," *Oriental Pharmacy and Experimental Medicine*, vol. 7, no. 2, pp. 197–204, 2007.
- [25] K. C. Pramanik, P. Bhattacharya, R. Biswas, D. Bandyopadhyay, M. Mishra, and T. Chatterjee, "Hypoglycemic and antihyperglycemic activity of leaf extract of *Pluchea indica* Less.," *Oriental Pharmacy and Experimental Medicine*, vol. 6, pp. 232–236, 2006.
- [26] D. Buapool, N. Mongkol, J. Chantimal, S. Roytrakul, E. Srisook, and K. Srisook, "Molecular mechanism of anti-inflammatory activity of *Pluchea indica* leaves in macrophages RAW 264.7 and its action in animal models of inflammation," *Journal of Ethnopharmacology*, vol. 146, no. 2, pp. 495–504, 2013.
- [27] P. S. Widyawati, C. H. Wijaya, P. S. Hardjosworo, and D. Sajuthi, "Volatile compounds of *Pluchea indica* Less and *Ocimum basilicum* Linn essential oil and potency as antioxidant," *Journal of Biosciences*, vol. 20, no. 3, pp. 117–126, 2013.
- [28] W. Brand-Williams, M.-E. Cuvelier, and C. Berset, "Use of a free radical method to evaluate antioxidant activity," *LWT-Food Science and Technology*, vol. 28, no. 1, pp. 25–30, 1995.
- [29] O. I. Aruoma and B. Halliwell, "Action of hypochlorous acid on the antioxidant protective enzymes superoxide dismutase, catalase and glutathione peroxidase," *Biochemical Journal*, vol. 248, no. 3, pp. 973–976, 1987.
- [30] J. M. Graham, J. A. Higgins, T. Gillott et al., "A novel method for the rapid separation of plasma lipoproteins using self-generating gradients of iodixanol," *Atherosclerosis*, vol. 124, no. 1, pp. 125–135, 1996.
- [31] A. Phaniendra, D. B. Jestadi, and L. Periyasamy, "Free radicals: properties, sources, targets, and their implication in various diseases," *Indian Journal of Clinical Biochemistry*, vol. 30, no. 1, pp. 11–26, 2015.
- [32] B. Halliwell and J. M. Gutteridge, *Free Radicals in Biology and Medicine*, Oxford university press, USA, 2015.
- [33] B. N. Ames, M. K. Shigenaga, and T. M. Hagen, "Oxidants, antioxidants, and the degenerative diseases of aging," *Proceedings of the National Academy of Sciences*, vol. 90, no. 17, pp. 7915–7922, 1993.
- [34] M. Wojcik, I. Burzynska-Pedziwiatr, and L. Wozniak, "A review of natural and synthetic antioxidants important for health and longevity," *Current Medicinal Chemistry*, vol. 17, no. 28, pp. 3262–3288, 2010.
- [35] Y. Fukushima, T. Ohie, Y. Yonekawa et al., "Coffee and green tea as a large source of antioxidant polyphenols in the Japanese population," *Journal of Agricultural and Food Chemistry*, vol. 57, no. 4, pp. 1253–1259, 2009.
- [36] G. C. Yen and P. D. Duh, "Scavenging effect of methanolic extracts of peanut hulls on free-radical and active-oxygen species," *Journal of Agricultural and Food Chemistry*, vol. 42, no. 3, pp. 629–632, 1994.
- [37] D. Bastos, L. Saldanha, R. Catharino et al., "Phenolic antioxidants identified by ESI-MS from yerba maté (*Ilex paraguariensis*) and green tea (*Camellia sinensis*) extracts," *Molecules*, vol. 12, no. 3, pp. 423–432, 2007.
- [38] R. Manian, N. Anusuya, P. Siddhuraju, and S. Manian, "The antioxidant activity and free radical scavenging potential of two different solvent extracts of *Camellia sinensis* (L.) O. Kuntz, *Ficus bengalensis* L. and *Ficus racemosa* L.," *Food Chemistry*, vol. 107, no. 3, pp. 1000–1007, 2008.
- [39] K. Srisook, D. Buapool, R. Boonbai, P. Simmasut, Y. Charoensuk, and E. Srisook, "Antioxidant and anti-inflammatory activities of hot water extract from *Pluchea indica* Less. herbal tea," *Journal of Medicinal Plants Research*, vol. 6, no. 23, pp. 4077–4408, 2012.
- [40] R. Tsao, "Chemistry and biochemistry of dietary polyphenols," *Nutrients*, vol. 2, no. 12, pp. 1231–1246, 2010.
- [41] L. S. Lee, S. H. Kim, Y. B. Kim, and Y. C. Kim, "Quantitative analysis of major constituents in green tea with different plucking periods and their antioxidant activity," *Molecules*, vol. 19, no. 7, pp. 9173–9186, 2014.
- [42] R. Sarkar, B. Hazra, S. Mandal, S. Biswas, and N. Mandal, "Assessment of *in vitro* antioxidant and free radical scavenging activity of *Cajanus cajan*," *Journal of Complementary and Integrative Medicine*, vol. 6, no. 1, p. 1248, 2009.
- [43] P. Valentão, E. Fernandes, F. Carvalho, P. Andrade, R. Seabra, and M. Bastos, "Hydroxyl radical and hypochlorous acid scavenging activity of small centaury (*Centaureum erythraea*) infusion. A comparative study with green tea (*Camellia sinensis*)," *Phytomedicine*, vol. 10, no. 6-7, pp. 517–522, 2003.
- [44] C. Cabrera, R. Artacho, and R. Giménez, "Beneficial effects of green tea—a review," *Journal of the American College of Nutrition*, vol. 25, no. 2, pp. 79–99, 2006.
- [45] D. Rees, R. Palmer, and S. Moncada, "Role of endothelium-derived nitric oxide in the regulation of blood pressure," *Proceedings of the National Academy of Sciences*, vol. 86, no. 9, pp. 3375–3378, 1989.
- [46] R. M. Palmer, A. Ferrige, and S. Moncada, "Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor," *Nature*, vol. 327, no. 6122, pp. 524–526, 1987.
- [47] H. Li and U. Förstermann, "Nitric oxide in the pathogenesis of vascular disease," *The Journal of Pathology*, vol. 190, no. 3, pp. 244–254, 2000.
- [48] P. J. Tsai, T. H. Tsai, C. H. Yu, and S. C. Ho, "Comparison of NO-scavenging and NO-suppressing activities of different herbal teas with those of green tea," *Food Chemistry*, vol. 103, no. 1, pp. 181–187, 2007.

- [49] P. Pacher, J. S. Beckman, and L. Liaudet, "Nitric oxide and peroxynitrite in health and disease," *Physiological Reviews*, vol. 87, no. 1, pp. 315–424, 2007.
- [50] H. Y. Chung, T. Yokozawa, D. Y. Soung, I. S. Kye, J. K. No, and B. S. Baek, "Peroxynitrite-scavenging activity of green tea tannin," *Journal of Agricultural and Food Chemistry*, vol. 46, no. 11, pp. 4484–4486, 1998.
- [51] H. Yoshida and R. Kisugi, "Mechanisms of LDL oxidation," *Clinica Chimica Acta*, vol. 411, no. 23-24, pp. 1875–1882, 2010.
- [52] R. Govindarajan, M. Vijayakumar, and P. Pushpangadan, "Antioxidant approach to disease management and the role of 'Rasayana' herbs of Ayurveda," *Journal of Ethnopharmacology*, vol. 99, no. 2, pp. 165–178, 2005.
- [53] B. Fuhrman and M. Aviram, "Flavonoids protect LDL from oxidation and attenuate atherosclerosis," *Current Opinion in Lipidology*, vol. 12, no. 1, pp. 41–48, 2001.
- [54] W. J. Craig, "Health-promoting properties of common herbs," *The American Journal of Clinical Nutrition*, vol. 70, no. 3, pp. 491s–499s, 1999.
- [55] T. Spranger, B. Finckh, R. Fingerhut, A. Kohlschütter, U. Beisiegel, and A. Kontush, "How different constituents of human plasma and low density lipoprotein determine plasma oxidizability by copper," *Chemistry and Physics of Lipids*, vol. 91, no. 1, pp. 39–52, 1998.
- [56] Z. Q. Liu, L. P. Ma, B. Zhou, L. Yang, and Z. L. Liu, "Antioxidative effects of green tea polyphenols on free radical initiated and photosensitized peroxidation of human low density lipoprotein," *Chemistry and Physics of Lipids*, vol. 106, no. 1, pp. 53–63, 2000.
- [57] T. Yokozawa and E. Dong, "Influence of green tea and its three major components upon low-density lipoprotein oxidation," *Experimental and Toxicologic Pathology*, vol. 49, no. 5, pp. 329–335, 1997.
- [58] T. M. Hung, M. Na, P. T. Thuong et al., "Antioxidant activity of caffeoyl quinic acid derivatives from the roots of *Dipsacus asper* Wall," *Journal of Ethnopharmacology*, vol. 108, no. 2, pp. 188–192, 2006.
- [59] M. S. Islam, M. Yoshimoto, and O. Yamakawa, "Distribution and physiological functions of caffeoylquinic acid derivatives in leaves of sweetpotato genotypes," *Journal of Food Science*, vol. 68, no. 1, pp. 111–116, 2003.
- [60] H. J. Park, A. Nugroho, J. H. Lee et al., "HPLC analysis of caffeoylquinic acids in the extract of *Cacalia firma* and peroxynitrite scavenging effect," *Korean Journal of Pharmacognosy*, vol. 40, no. 4, pp. 365–369, 2009.