



LJMU Research Online

Malhotra, A, Oxborough, D, Rao, P, Finocchiaro, G, Dhutia, H, Prasad, V, Miller, C, Keavney, B, Papadakis, M and Sharma, S

Defining the Normal Spectrum of Electrocardiographic and Left Ventricular Adaptations in Mixed-Race Male Adolescent Soccer Players.

<http://researchonline.ljmu.ac.uk/id/eprint/14225/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Malhotra, A, Oxborough, D, Rao, P, Finocchiaro, G, Dhutia, H, Prasad, V, Miller, C, Keavney, B, Papadakis, M and Sharma, S (2020) Defining the Normal Spectrum of Electrocardiographic and Left Ventricular Adaptations in Mixed-Race Male Adolescent Soccer Players. *Circulation*. 143 (1). pp. 94-

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

RESEARCH LETTER

Defining the Normal Spectrum of Electrocardiographic and Left Ventricular Adaptations in Mixed-Race Male Adolescent Soccer Players

Contemporary data on exercise-related sudden cardiac death reveal that young adolescent athletes (14 to 18 years old) are the most vulnerable group.¹ Several studies have described the adolescent athlete's heart but have focused only on White or Black athletes.^{2,3} The demographic landscape for adolescent athletes has changed over the past few decades and consists of an increasing population of mixed-race individuals in whom one parent is Black (African or Afro-Caribbean) and the other parent is White. We examined the electric and structural adaptations in the heart in healthy mixed-race male soccer players and compared them with those of White and Black male soccer players.

Ethics approval was granted in accordance with the Research Governance Framework for Health and Social Care (2005) and the English Football Association. Written consent was obtained from all athletes in accordance with the Data Protection Act (approved in 1998). The data that support the findings of this study are available from the corresponding author on reasonable request.

Between 2015 and 2018, 3000 consecutive cases of healthy mixed-race, Black, and White athletes were assessed with ECG and echocardiography. A mixed-race athlete was classified as an individual with 1 White parent of European origin and 1 Black parent of African/Afro-Caribbean origin.

The 12-lead ECG was interpreted in accordance with international recommendations.⁴ Transthoracic echocardiography was performed in accordance with standard American and European protocols. The χ^2 test was used to compare the prevalence of ECG and echocardiogram features among Black, White, and mixed-race groups.

The mean age was 16.4 ± 1.3 years. Among mixed-race athletes, 1 parent was White British in all cases; the other parent was either of African (60%) or Caribbean (40%) origin. Black athletes were of African (61%) or Caribbean (39%) origin.

ELECTRIC CHANGES

Sinus bradycardia and Sokolow-Lyon criterion for left ventricular (LV) hypertrophy were most prevalent among mixed-race athletes. Mixed-race and Black athletes demonstrated a similar prevalence of voltage criteria for left and right atrial enlargement, which was higher compared with White athletes.

T-wave inversion (TWI) was most common in Black athletes, followed by mixed-race and White athletes (12.6%, 8.6%, and 2.3%, respectively). The prevalence of anterior TWI was higher in Black athletes (9.5%) than in mixed-race and White athletes (1.5%).

Mixed-race and Black athletes had a similar prevalence of inferior TWI (2% versus 1.5%; $P=0.49$). This was higher in mixed-race than in White athletes (2% versus 0.5%; $P=0.004$) and in Black versus White athletes (1.5% versus

Aneil Malhotra , MB, BChir, MA, MRCP(UK), MSc, PhD
David Oxborough, MSc, PhD
Prashant Rao, MRCP(UK), MD
Gherardo Finocchiaro, MD, MRCP(UK)
Harshil Dhutia, BSc, MBBS, MRCP(UK), MD
Vivek Prasad, BSc
Chris Miller, BSc, MBBS, MRCP(UK), PhD
Bernard Keavney , BSc, BM BCh, DM
Michael Papadakis , BSc, MRCP(UK), MD
Sanjay Sharma , BSc, MRCP(UK), MD

Key Words: athletes
■ electrocardiography ■ physiological adaptation

© 2020 American Heart Association, Inc.

<https://www.ahajournals.org/journal/circ>

0.5%; $P=0.04$). Mixed-raced and Black athletes had a similar prevalence of lateral/apical TWI (0.6% versus 1.0%; $P=0.33$), which was also higher among Black athletes than among White athletes (1.0% versus 0.2%; $P=0.02$; Table). No arrhythmias were identified in any group.

STRUCTURAL CHANGES

Mixed-race athletes revealed a greater LV wall thickness than White athletes even after indexing for body surface area. Black athletes revealed a greater LV wall thickness compared with both White and mixed-race athletes. In absolute terms, LV hypertrophy >12 mm was present in 7.1% of Black athletes, 5.9% of mixed-race athletes, and 1.3% of White athletes. The maximal LV wall thickness did not exceed 15 mm in any athlete.

White athletes demonstrated the largest LV cavity size and mixed-race athletes had larger LV cavities than Black athletes. These differences became more pronounced when indexed for body surface area. A total of 5.9% of White athletes demonstrated an LV end diastolic dimension >58 mm, as did 4.4% of mixed-race athletes and 3.2% of Black athletes. None of the athletes showed an LV end diastolic dimension >62 mm.

Among athletes with an abnormal ECG (mixed-race interpretation as in White athletes), a LV wall thickness >12 mm, LV cavity >60 mm, or LV ejection fraction $<50\%$, we had excluded overt cardiomyopathy on the basis of a normal exercise stress test/echocardiogram, 24-hour ECG monitor, and absence of scar on cardiovascular MRI before inclusion in the study.

Mixed-race athletes revealed ECG characteristics that were more similar to Black athletes than White athletes,

Table. Electrocardiographic and Echocardiographic Features of 3000 Male Adolescent Athletes, by Ethnicity

ECG/echocardiogram feature	White (n=1000), %	Black (n= 1000), %	Mixed (n=1000), %	White vs Black, P value	White vs mixed race, P value	Black vs mixed race, P value
Sinus bradycardia <60 bpm	45.4	46.5	67	0.65	$<0.0001^*$	$<0.0001^*$
Sinus arrhythmia	37.3	36.8	35.5	0.85	0.43	0.58
LV hypertrophy	25.6	17.6	30	$<0.0001^*$	0.03*	$<0.0001^*$
RV hypertrophy	2.3	4.2	3.0	0.02*	0.4	0.19
Left atrial enlargement	1	5.9	4.7	$<0.0001^*$	$<0.0001^*$	0.27
Right atrial enlargement	0.7	3.9	3.0	$<0.0001^*$	0.0002*	0.33
TWI, all	2.3	13.0	8.6	$<0.0001^*$	$<0.0001^*$	0.0019*
TWI, anterior V1, V2	0.9	5.8	3.7	$<0.0001^*$	$<0.0001^*$	0.04*
TWI, anterior beyond V2	0.6	3.7	2.0	$<0.0001^*$	0.0089*	0.03*
TWI, inferior	0.5	1.5	2.0	0.04*	0.0039*	0.49
TWI, lateral/apical	0.2	1	0.6	0.02*	0.29	0.33
TWI, inferolateral/apical	0.1	0.9	0.2	0.02*	1	0.06
ST segment elevation	48.4	63.6	54.7	$<0.0001^*$	0.0055*	$<0.0001^*$
Ascending convex	28.8	38.0	32.7	$<0.0001^*$	0.04*	0.0150*
Ascending concave	16.2	21.8	19.8	0.0017*	0.04*	0.3
Isoelectric	3.2	3.3	2.0	1	0.12	0.09
Depressed	0.2	0.5	0.2	0.45	1	0.45
Left atrial diameter, mm	32.9 \pm 5.2	32.8 \pm 4.7	32.6 \pm 4.4	0.65	0.16	0.33
LVWT, mm	9.15 \pm 1.3	10.1 \pm 1.4	9.8 \pm 1.41	$<0.0001^*$	$<0.0001^*$	$<0.0001^*$
Relative wall thickness	0.36 \pm 0.05	0.38 \pm 0.04	0.37 \pm 0.05	$<0.0001^*$	$<0.0001^*$	$<0.0001^*$
LVEDd, mm	52.4 \pm 3.9	50.6 \pm 4	51.1 \pm 4.03	$<0.0001^*$	$<0.0001^*$	0.0054*
LVEDdi, mm/m ²	27.9 \pm 2.12	26.6 \pm 2.19	27.2 \pm 2.05	$<0.0001^*$	$<0.0001^*$	$<0.0001^*$
LVM, g	168 \pm 33	171 \pm 35	173 \pm 31	0.048*	$<0.0001^*$	0.18
LVMi, b/m ²	88.1 \pm 16.5	89.6 \pm 17.5	91.2 \pm 16.5	0.04*	$<0.0001^*$	0.03*
LV E/A	2.10 \pm 0.6	2.02 \pm 0.5	2.05 \pm 0.5	0.0012*	0.04*	0.18
RVD1, mm	36.58 \pm 4.76	36.35 \pm 4.80	36.45 \pm 4.11	0.28	0.51	0.62
Aortic root, mm	27.56 \pm 3.22	27.93 \pm 3.16	27.90 \pm 3.81	0.01*	0.03*	0.85

E/A indicates early filling velocity/atrial filling velocity; LV, left ventricular; LVEDd, left ventricular end diastolic diameter; LVEDdi, left ventricular end diastolic dimension index; LVM, left ventricular mass; LVMi, left ventricular mass indexed to body surface area; LVWT, left ventricular wall thickness; RV, right ventricular; RVD1, right ventricular diameter at level of tricuspid valve; and TWI, T-wave inversion.

*Statistically significant ($P<0.05$).

with specific reference to left and right atrial enlargement and TWI. Inferior and lateral/apical TWI was rare in all groups, indicating the need for further investigations and clinical surveillance.

Mixed-race athletes demonstrated a mean LV wall thickness and LV cavity size that fell between values observed in the Black and White athlete groups. The electric and structural parameters for the mixed-race athlete seem to be more similar to those of the Black athlete. One in 14 healthy mixed-race athletes demonstrate TWI or have an LV wall thickness >13 mm. On the basis of our observations, anterior TWI and a marginally increased LV wall thickness may be normal adaptation to athletic training in mixed-race athletes.

The study included only elite male soccer players, but soccer involves a combination of strength and endurance exercises and is the most popular sport worldwide. The cross-sectional nature of the study precludes comment on the longitudinal implications of repolarization changes, mild LV hypertrophy, or LV dilation with reference to development of cardiomyopathy in the future.

ARTICLE INFORMATION

Correspondence

Aneil Malhotra, St George's, University of London, Cranmer Terrace, London, United Kingdom SW17 0RE. Email amalhotr@sgul.ac.uk

Affiliations

Cardiology Clinical Academic Group, St George's, University of London, United Kingdom (A.M., M.P., S.S.). Division of Cardiovascular Sciences, School of Medical

Sciences, Faculty of Biology, Medicine and Health, University of Manchester, United Kingdom (A.M., C.M., B.K.). Manchester University National Health Service Foundation Trust, Manchester Academic Health Science Centre, United Kingdom (A.M., V.P., C.M., B.K.). Research Institute for Sports and Exercise Science, Liverpool John Moore's University, United Kingdom (D.O.). Beth Israel Deaconess Medical Centre, Harvard Medical School, Boston, MA (P.R.). Cardiothoracic Centre, Guy's and St Thomas' Hospital, London, United Kingdom (G.F.). Department of Cardiology, Glenfield Hospital, Leicester, United Kingdom (H.D.).

Acknowledgments

The authors thank The Football Association Cardiology Consensus panel and Cardiac Risk in the Young charity for support. Dr Miller is supported by the National Institute of Health and Research. Professor Bernard Keavney is supported by the British Heart Foundation.

Disclosures

None.

REFERENCES

1. Malhotra A, Dhutia H, Finocchiaro G, Gati S, Beasley I, Clift P, Cowie C, Kenny A, Mayet J, Oxborough D, et al. Outcomes of cardiac screening in adolescent soccer players. *N Engl J Med*. 2018;379:524–534. doi: 10.1056/NEJMoa1714719
2. Papadakis M, Carre F, Kervio G, Rawlins J, Panoulas VF, Chandra N, Basavarajiah S, Carby L, Fonseca T, Sharma S. The prevalence, distribution, and clinical outcomes of electrocardiographic repolarization patterns in male athletes of African/Afro-Caribbean origin. *Eur Heart J*. 2011;32:2304–2313. doi: 10.1093/eurheartj/ehr140
3. Sheikh N, Papadakis M, Carre F, Kervio G, Panoulas VF, Ghani S, Zaidi A, Gati S, Rawlins J, Wilson MG, et al. Cardiac adaptation to exercise in adolescent athletes of African ethnicity: an emergent elite athletic population. *Br J Sports Med*. 2013;47:585–592. doi: 10.1136/bjsports-2012-091874
4. Sharma S, Drezner JA, Baggish A, Papadakis M, Wilson MG, Prutkin JM, La Gerche A, Ackerman MJ, Borjesson M, Salerno JC, et al. International recommendations for electrocardiographic interpretation in athletes. *Eur Heart J*. 2018;39:1466–1480. doi: 10.1093/eurheartj/ehw631