



LJMU Research Online

Blackman, RC, Ling, KKS, Harper, LR, Shum, P, Hanfling, B and Lawson-Handley, L

Targeted and passive environmental DNA approaches outperform established methods for detection of quagga mussels, *Dreissena rostriformis bugensis* in flowing water

<http://researchonline.ljmu.ac.uk/id/eprint/14458/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Blackman, RC, Ling, KKS, Harper, LR, Shum, P, Hanfling, B and Lawson-Handley, L (2020) Targeted and passive environmental DNA approaches outperform established methods for detection of quagga mussels, *Dreissena rostriformis bugensis* in flowing water. *Ecology and Evolution*. 10

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Targeted and passive environmental DNA approaches outperform established methods for detection of quagga mussels, *Dreissena rostriformis bugensis* in flowing water

Rosetta C. Blackman^{1,2,3}  | Kar Keun Sean Ling³ | Lynsey R. Harper^{3,4}  | Peter Shum^{3,4}  | Bernd Hänfling³  | Lori Lawson-Handley³ 

¹Department of Aquatic Ecology, Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

²Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zürich, Switzerland

³Evolutionary and Environmental Genomics Group (EvoHull), Department of Biological and Marine Sciences, University of Hull, Hull, UK

⁴School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, UK

Correspondence

Rosetta Blackman, Department of Aquatic Ecology, Eawag, Swiss Federal Institute of Aquatic Science and Technology, Überlandstrasse 133, CH-8600 Dübendorf, Switzerland.
Email: Rosetta.Blackman@eawag.com

Funding information

Gilson Le Cren Award

Abstract

1. The early detection of invasive non-native species (INNS) is important for informing management actions. Established monitoring methods require the collection or observation of specimens, which is unlikely at the beginning of an invasion when densities are likely to be low. Environmental DNA (eDNA) analysis is a highly promising technique for the detection of INNS—particularly during the early stages of an invasion.
2. Here, we compared the use of traditional kick-net sampling with two eDNA approaches (targeted detection using both conventional and quantitative PCR and passive detection via metabarcoding with conserved primers) for detection of quagga mussel, *Dreissena rostriformis bugensis*, a high priority INNS, along a density gradient on the River Wraysbury, UK.
3. All three molecular tools outperformed traditional sampling in terms of detection. Conventional PCR and qPCR both had 100% detection rate in all samples and outperformed metabarcoding when the target species was at low densities. Additionally, quagga mussel DNA copy number (qPCR) and relative read count (metabarcoding) were significantly influenced by both mussel density and distance from source population, with distance being the most significant predictor.
4. *Synthesis and application.* All three molecular approaches were more sensitive than traditional kick-net sampling for the detection of the quagga mussel in flowing water, and both qPCR and metabarcoding enabled estimates of relative abundance. Targeted approaches were more sensitive than metabarcoding, but metabarcoding has the advantage of providing information on the wider community and consequently the impacts of INNS.

KEYWORDS

early detection, eDNA, invasive non-native species, metabarcoding, PCR, qPCR, targeted assay

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Invasive non-native species (INNS) have wide ranging effects on ecosystems, from biodiversity loss (Doherty et al., 2016; Gallardo et al., 2016) to economic impacts, including damage to infrastructure (Connelly et al., 2007; Pimentel et al., 2005). Eradication or containment methods are most effective during the early stages of an invasion, both in terms of success and lower financial cost (Hulme, 2006; Mehta et al., 2007). Early detection is particularly challenging for species that are small, elusive, or cryptic, and these species often go unnoticed until they become established (Simmons et al., 2015; Blackman et al., 2017). Monitoring organisms such as freshwater invertebrates generally relies on specimen collection and morphological identification, and newly invading species can be easily overlooked. Alternative methods that offer rapid and cost-effective detection of new INNS are crucial at a time when rapid increase in trade and transport has caused a surge in the number of successful invasions to new environments (Hulme, 2006).

The use of molecular methods to identify DNA taken from environmental samples (i.e., environmental DNA or “eDNA”) has been a major research focus in biodiversity monitoring over the last 10 years, including the detection of INNS (Ficetola et al., 2008; Thomsen et al., 2012; Blackman et al., 2018). Two broad approaches can be applied to detect a species of interest. The “targeted” approach utilizes species-specific primers with conventional PCR (cPCR), quantitative PCR (qPCR), or droplet digital PCR (ddPCR) to detect a single species. By contrast, metabarcoding can be considered a “passive approach” as it does not focus on single species (Lawson Handley, 2015; Creer et al., 2016; Deiner et al., 2017). Metabarcoding utilizes conserved primers to amplify DNA from a group of taxa, for example, fish and other vertebrates (e.g., Kelly et al., 2014; Miya et al., 2015; Simmons et al., 2015), zooplankton (e.g., Brown et al., 2016), or insects (e.g., Yu et al., 2012; Elbrecht and Leese, 2017). Amplicons are then sequenced using high-throughput sequencing (HTS) platforms and bioinformatically assigned to taxa. Metabarcoding provides an opportunity to detect multiple species simultaneously, including those, which have not been listed as priority INNS (Simmons et al., 2015; Blackman et al., 2017). Comparisons between traditional methods for biodiversity monitoring and targeted (Dejean et al., 2012; Tréguier et al., 2014) or passive (Smart et al., 2015; Hänfling et al., 2016; Wilcox et al., 2016) molecular approaches have demonstrated an increased probability of detection using eDNA methods.

The targeted approach was first used by Ficetola et al. (2008) to detect invasive American bullfrog *Lithobates catesbeianus*, in lentic waterbodies, and has since been applied to over 100 target INNS (Blackman et al., 2018). Conventional PCR is a well-established and widely practiced method for detecting presence/absence of target DNA by visualization of PCR products on agarose gels. However, quantitative PCR (qPCR) has on many occasions been shown to be more sensitive than PCR (e.g., Nathan et al., 2014; Simmons et al., 2015; De Ventura et al., 2017) and therefore more suitable for detecting INNS at low density, for example, the early stages of

invasion. Quantitative PCR has the added advantage of providing the number of DNA copies in a sample, which has been successfully linked to biomass or abundance of some taxa (e.g., Amphibians, Thomsen et al., 2012; Fish, Takahara et al., 2012; Gastropods, Goldberg et al., 2013). However, qPCR requires the construction of calibration curves from known standard concentrations and is substantially more expensive and time-consuming than cPCR (Nathan et al., 2014) and maybe less accessible to end-users than cPCR (Blackman et al., 2020).

The potential of eDNA metabarcoding for INNS monitoring has already been shown as an effective tool (Simmons et al., 2015; Brown et al., 2016; Blackman et al., 2017; Borrell et al., 2017; Holman et al., 2019). These studies include the detection of new INNS, which had been previously overlooked by traditional methods. For example, Blackman et al. (2017) used a metabarcoding approach to investigate macroinvertebrate diversity and discovered a non-native Gammaridae species, *Gammarus fossarum*, which traditional methods had failed to identify for over 50 years in the UK. Similarly, Simmons et al. (2015) detected the Northern Snakehead, *Channa argus*, in the Muskingum River Watershed, Ohio, which had not been included on INNS priority lists. Metabarcoding also has the potential to act as a surveillance tool and has been applied to monitor high-risk pathways, such as ballast water (Ardura et al., 2015; Zaiko et al., 2015), the live bait trade (Mahon et al., 2014), and ports (Borrell et al., 2017; Brown et al., 2016). However, comparisons of targeted and passive approaches for INNS or other focal species have been exemplified in the bighead carp, *Hypophthalmichthys nobilis* (Simmons et al., 2015) and great crested newts, *Triturus cristatus* (Harper et al., 2018). These studies both showed that a targeted approach had a higher sensitivity than the passive approach. Possible causes of this include primer choice and PCR bias, that is, species successfully amplifying and masking others in the PCR reaction during metabarcoding (Harper et al., 2018). Further comparisons of these methods covering different target taxa are required to better inform management strategies.

In this study, we focus on the quagga mussel, *Dreissena rostriformis bugensis*, a highly invasive bivalve from the Ponto-Caspian region, which has a significant impact on all trophic levels within the environments it invades (Karatayev et al., 2007; Roy et al., 2014). In a horizon scanning exercise from 2014, the quagga mussel scored the highest out of 591 species evaluated for its potential to invade, establish, and have negative impact in the UK (Roy et al., 2014). Consistent with predictions, quagga mussels were subsequently detected later the same year (Aldridge et al., 2014). Due to its wider tolerance to environmental conditions, it has the potential to spread rapidly and occupies a greater range of habitat than the closely related zebra mussel, *Dreissena polymorpha*, which has already spread throughout much of Europe and North America (Nalepa et al., 2010; Gallardo & Aldridge, 2013; Quinn et al., 2014). Motivated by a need for cost-effective tools for quagga mussel detection, we previously designed a cPCR assay, which provided 100% detection rate in field trials with quagga mussel populations (Blackman et al., 2020). Here, we further developed and validated this assay for abundance estimation using a dye-based qPCR approach

in mesocosm experiments and field trials. Second, we directly compared the three molecular methods (cPCR, qPCR, and metabarcoding) to traditional kick-net sampling for detection of quagga mussels in the field. Finally, we investigated whether qPCR and metabarcoding are suitable for estimating relative abundance of quagga mussels by sampling a density gradient away from the main population. We hypothesized that: (1) The probability of detecting quagga mussels will be higher for eDNA methods than traditional methods, (2) targeted methods will have a higher probability of detecting quagga mussel than the passive method, (3) qPCR will be more sensitive than cPCR for quagga mussel detection, and (4) both read count and DNA copy number will decline with distance downstream of the main population.

2 | METHODS

2.1 | Field sample collection, DNA capture, and extraction

Quagga mussels were first detected in the UK in Wraysbury Reservoir, Berkshire, in October 2014. This site is considered the founding population of quagga mussels in the UK. The mussels spread through the outfall and along the adjacent River Wraysbury, which is a tributary of the River Colne. The species currently has a restricted distribution in the South East of England but has also been recorded in reservoirs adjacent to the River Lee, north of London, and in the River Thames. Samples were collected from Wraysbury Reservoir and River in May 2016, working in an upstream direction to ensure DNA availability and inhibitor levels were not altered by sampling activity. In order to compare established and molecular methods directly, we collected eDNA and kick-net samples at six sites across the river at each of three main sampling locations downstream of Wraysbury Reservoir: Wraysbury Weir (WW, 0.61 km downstream of Wraysbury Reservoir), Wraysbury Bridge (WB, 1.10 km), and Wraysbury Gardens (WG, 2.7 km) (orange diamonds, Figure 1 and Supplementary Information I Table S1). Water samples from these three locations consisted of three 500 ml replicates, collected at each of six sites across the width of the river ($n = 54$). Environmental DNA samples were collected from the river surface, using sterile gloves and Gosselin™ HDPE plastic bottles (Fisher Scientific UK Ltd) prior to kick-net sampling. Three-minute kick-net samples were collected from the same six sites at each of these three locations, in accordance with established methods (Murray-Bligh, 1999). Environmental DNA sampling of three 500 ml replicates was also carried out at four additional locations along Wraysbury River (Reservoir Outfall, RO, 0.2 km; Moor Lane, ML, 1.7 km; Hale Street, HS, 2.72 km; and Upper Thames Confluence, UT, 2.75 km, blue diamonds, Figure 1, Supplementary Information I Table S1). These additional eDNA samples were collected to provide additional fine-scale resolution in the eDNA analysis and were analyzed with cPCR and qPCR only, whereas samples from the three main locations were analyzed with cPCR, qPCR, and metabarcoding. Sampling locations within the river were chosen based primarily on accessibility, in accordance with standard macroinvertebrate monitoring procedures carried out by UK




FIGURE 1 Sampling locations on the Wraysbury River. Diamond symbols represent the sampling locations: Reservoir Outfall (RO, 0.2 km downstream of the reservoir), Wraysbury Weir (WW, 0.61 km), Wraysbury Bridge (WB, 1.10 km), Moor Lane (ML, 1.7 km), Wraysbury Gardens (WG, 2.7 km), Hale Street (HS, 2.72 km), and Upstream Thames confluence (UT, 2.75 km). Environmental DNA and kick-net samples were collected from sites with an orange outline, and sites sampled for eDNA only have a blue outlined symbol (Full grid references are provided in Supplementary Information I Table S1)

agencies. We did not target specific locations with previous records of quagga mussels or their preferred habitat. At the time of sampling, quagga mussels had not been reported upstream of the reservoir. We therefore collected an eDNA sample upstream of the reservoir with the expectation that this would be negative for quagga mussel (Waterside Car Park, $n = 3$). Sample bottles filled with ddH₂O were also taken into the field as filtering blanks ($n = 3$). Hence, the total number of eDNA samples collected in the field was $n = 72$. All eDNA samples and blanks were stored in a cool box with ice packs and transported to the laboratory for filtration within 24 hr. Water samples were filtered through 0.45- μ m cellulose nitrate membrane filters attached to a vacuum pump and DNA extractions performed following Brolaski et al. (2008) with minor modifications (see Supplementary Information I and II for full details). Kick-net samples were placed in individual 42 fluid oz. sterile Whirl-pak® bags (Cole-Palmer, Hanwell, London) and stored in a cool box before being frozen at -20°C in the laboratory. Each sample was thawed at room temperature and rinsed with ddH₂O separately prior to analysis. Samples were then sorted for the presence of *D. r. bugensis* only, and all specimens were counted and subsequently stored at -20°C .

2.2 | Performance of targeted molecular detection

The performance of targeted DNA assays for the detection of invasive quagga mussel DNA was first examined through

laboratory-based mesocosm experiments. The objectives of these experiments were (a) to validate species-specific DNA assays, using both conventional and quantitative PCR, to target eDNA detection of quagga mussels; (b) to assess the duration and persistence of eDNA signal over a 42-day period; and (c) to examine a correlation between eDNA concentration and known densities of quagga mussel. The mesocosm experiment was set up to monitor DNA production of mussels at three densities (1, 5, and 20 individuals, with total and individual biomass weights recorded) over 21 days, with three replicates of each density and a control tank with no mussels present. Fifteen liter plastic tanks with individual aeration stones were set up in a climate controlled facility where temperature averaged 16°C (range 14–18°C), with light:dark cycles of 16 hr:8 hr. Sampling events took place over 42 days at: 0 hrs (prior to specimens being added), 4 hr, 8 hr, 24 hr, 48 hr, 72 hr, 7 days, 15 days, and 21 days. On day 21, the specimens were removed, and sampling continued at: 22, 28, 35, and 42 days ($n = 130$). Two hundred milliliter water samples were collected at each time point, and each sample was filtered through a single filter membrane following the same protocol as the field samples (Blackman et al., 2020).

2.2.1 | Conventional PCR (cPCR) assay

We used a cPCR assay to detect the quagga mussel, *D. r. bugensis*, using a species-specific primer pair targeting a 188 bp fragment of the cytochrome oxidase I (COI) gene (DRB1_F - 5'-GGAACTGGTTGGTCCCGAT3'- and DRB1_R 5'-GGCCCTGAATGCCCCATAAT-3', see Supplementary Material I for full cPCR protocol). The DRB1 cPCR assay has shown to be highly sensitive (positive detection with 0.001 ng of target tissue DNA per reaction) and does not cross amplify closely related Dreissenid or native mussel species (Blackman et al., 2020). All mesocosm ($n = 130$), field samples ($n = 66$), and control samples (field (3), filter (3) and cPCR (6) blanks) were tested with the DRB1 assay. A tissue DNA sample from closely related zebra mussel was included as an additional negative control.

2.2.2 | Quantitative PCR (qPCR) assay

The DRB1 assay was further developed using fluorescent dye-based qPCR (PowerUp SYBR green, Thermo Fisher, see Supplementary Material I for full details). Briefly, we performed serial dilutions of DNA from tissue samples (from 1 to 1×10^{-7} ng/ μ l) to estimate the LoD and performed the DRB1 qPCR assay on samples collected from the mesocosm experiment, field and control samples. Zebra mussel tissue DNA was used as a negative control in the qPCR assays. Amplification curves, C_q values, and melt curves were analyzed using StepOne-Plus™ software 2.0. Quantification calibration curves using a synthesized gBlock of the target fragment were used to estimate the DNA copy number for each sample (See Supplementary Information I and IV for full details).

2.3 | Passive detection: eDNA metabarcoding

Degenerate metazoan primers that target the conserved COI region were used for metabarcoding (Geller et al., 2013; Leray et al., 2013): jgHCO2198: TAIACYTCIGGRTGICC RAARAAAYCA and mICOLintF: GGWACWGGWTGAACWGTWTAYCCYCC. These primers were chosen as they (a) target the same mitochondrial gene as our targeted assays, (b) are the most commonly used COI metabarcoding primers and are good candidates for studying the entire river community, and (c) we confirmed in previous work that they amplify Dreissenid mussels. We used universal primers rather than mollusk-specific metabarcoding primers (which are also available, Klymus et al., 2017), since universal primers are more likely to be useful in routine community monitoring of INNS. We confirmed amplification of the target taxa from tissue extractions prior to library preparation. Full details of the library preparation are provided in Supplementary Information I. In brief, the 54 field samples from WW, WB, and WG (Figure 1) were sequenced, with the addition of 3 field blanks, 3 filter blanks, 6 PCR blanks, and 3 positive non-target tissue samples (2 of *Triops cancriformis* and 1 of *Osmia bicornis* total $n = 69$). A two-step PCR protocol was used for library preparation: First, the target region was amplified for each sample and a second amplification was performed to add the Illumina adaptor sequences onto the initial amplicons. Samples were normalized following the second PCR using SequalPrep Normalization plates (Invitrogen), and samples were then pooled in equimolar concentrations. The final pooled library was concentrated, cleaned with a QIAquick Gel Extraction Kit (Qiagen), quantified, and sequenced at 13 pM concentration with 20% PhiX, on an Illumina MiSeq platform using 2x 250bp v2 chemistry at the University of Hull.

2.4 | Bioinformatics and data analysis

Full details of the bioinformatics pipeline are provided in Supplementary Material I. In brief, processing of Illumina data and taxonomic assignment were performed using a custom pipeline, MetaBEAT (metaBarcoding and eDNA Analysis Tool) v 0.8 (<https://github.com/HullUnibioinformatics/metaBEAT>) but with minor modifications including filtering for the presence of the target taxa, *D. r. bugensis*, by comparing against curated reference sequences from GenBank, listed in Supplementary Material I.

We normalized the metabarcoding data to account for uneven sequencing depth between samples, by dividing the number of reads for each OTU within a sample by the total number of reads in that sample and multiplying by 100 (hereafter referred to as relative read count). We used generalized linear models (GLMs) to investigate (1) the influence of mussel density, biomass, and time on the DNA copy number (qPCR) since the start of the mesocosm experiment (before the mussels were removed from the tanks only) and (2) the influence of mussel density and distance from source population on the DNA copy number (qPCR) from the field samples and relative read count (metabarcoding). Initial model tests showed all data sets were

overdispersed; therefore, a negative binomial GLM with a log link function was used to resolve this issue. To check for model suitability, we checked the residual deviance of each model fitted a chi-square distribution (see Supplementary Information III Tables S2, S3 and S4 for GLM results). Finally, we assessed the relationship between DNA copy number and relative read count with the distance along the river using all seven sites included in the qPCR experiment, using a Spearman rank correlation since the data were not normally distributed. All calculations and visualizations of the data were carried out in R version 3.6.3 (R Core Team, 2019), with GLMs performed using the MASS package, version 7.3-51.6 (Venables & Ripley, 2002).

3 | RESULTS

We successfully adapted the DRB1 primer pair for detection of quagga mussel with the addition of SYBR green. Across both traditional and eDNA field experiments, we were able to detect quagga mussel with kick-net sampling, cPCR, qPCR, and metabarcoding. All negative controls (field blanks, filtration blanks, PCR blanks, and non-target tissue blanks) were negative for cPCR and qPCR. Minor contamination was detected in 3 field blanks with metabarcoding, with low reads of *Homo sapiens* (5 and 14 reads), *Triops cancriformis* (the positive tissue extract) (10 reads), and unassigned reads (42–189 reads). We therefore applied a 0.2% read threshold to remove possible contamination from the metabarcoding data set (i.e., sequences that had a frequency of 0.2% or less in each sample were removed).

3.1 | qPCR assay development and validation in mesocosms

The limit of detection (LoD) determined from the serial dilutions of tissue DNA was 1×10^{-4} ng/ μ l per reaction with both cPCR and qPCR, which equates to approximately 5 copies per reaction (See Supplementary Information III Figures S1 and S2 for amplification plots). Rapid DNA accumulation in the first two days of the mesocosm experiments was followed by a depletion in DNA concentration until removal of mussels on day 21 of the experiment (Figure 2).




FIGURE 2 *Dreissena rostriformis bugensis* qPCR mesocosm assay validation. The graph shows the mean and standard deviation of DNA copy number recorded for each density treatment during the mesocosm experiment: red line—1 specimen, green line—5 specimens, blue line—20 specimens, and the black line is the mean copy number for all three densities. The vertical dashed line indicates when the mussels were removed from the mesocosm experiment

This pattern of DNA accumulation and depletion is highly consistent across all three density treatments. Time since the beginning of the experiment (until removal of the mussels) had a significant effect on the DNA copy number in each mesocosm experiment when all density treatments were analyzed together (GLM, $Z = -7.224$, $p < .0001$, $AIC = 912.95$). There is still some overdispersion with this model; however, the overdispersion parameter indicates this is not substantial (overdispersion statistic = 1.3, Residual Deviance 91.142 on 70 df , $\chi^2 p = .0457$). There is substantial overlap between the DNA copy number estimates for the three density treatments (Figure 2), and neither density nor total biomass significantly explained the number of DNA copies amplified by qPCR (Density $Z = -1.019$, $p = .308$; Total biomass $Z = -0.776$, $p = .438$). DNA was still detectable 24 hr after removal of the mussels, but not at 7 days after removal (Day 28) (Figure 2). All blanks and mesocosm control tank samples were negative of target DNA throughout the experiment.

3.2 | Detection of *D. r. bugensis* by traditional and molecular approaches in the field

3.2.1 | Kick-net sampling

The number of *D. r. bugensis* specimens varied from 14 at Wraysbury Weir, the location closest to the reservoir outfall (source population), to a single specimen at Wraysbury Gardens, the location furthest from the outfall (Table 1). Detection rate was also highest at Wraysbury Weir (mussels detected in 4/6 sampling events). Mussels were detected in 1/6 samples at each of the other two sampling locations (Table 1).

3.3 | Targeted approaches

Detection rate was 100% in all three field locations, and all replicates tested for both targeted approaches (cPCR and qPCR) (Table 1 and Figure S3). Figure 3a shows the DNA copy number (\log_{10}) of the qPCR assay for *D. r. bugensis* found at each of the sampling locations. Both distance from source and number of mussels found at

the three locations, significantly influenced the DNA copy number, but distance was the strongest predictor (density $Z = -3.367$, $p = .000761$; distance $Z = -17.223$, $p < .0001$; AIC: 1392.1, residual deviance 57.621 on 51 df , $\chi^2 p = .244$). A Spearman Rank correlation between the DNA copy number and distance from the source

TABLE 1 (a–c) Comparison of methods for the detection of *D. r. bugensis* at three locations on the Wraybury River

Sample no.	Kick sampling		Environmental DNA	
	No. of individuals	PCR	qPCR	Metabarcoding
A. Wraybury Weir (WW)				
1	–	+++	+++	+++
2	+ (4)	+++	+++	+++
3	+ (4)	+++	+++	+++
4	–	+++	+++	+++
5	+ (3)	+++	+++	+++
6	+ (3)	+++	+++	+++
B. Wraybury Bridge (WB)				
1	–	+++	+++	+++
2	–	+++	+++	+++
3	–	+++	+++	+++
4	–	+++	+++	+++
5	+ (4)	+++	+++	+++
6	–	+++	+++	+++
C. Wraybury Gardens (WG)				
1	–	+++	+++	– + –
2	–	+++	+++	– – +
3	+ (1)	+++	+++	– – +
4	–	+++	+++	– – –
5	–	+++	+++	– – –
6	–	+++	+++	– – –

Note: “+” indicates a positive detection and “–” indicates no detection for each technical replicate (i.e., three per sample for the eDNA methods and one for the kick sample). The numbers in parentheses under Kick Sampling correspond to the number of individuals found in each sample.

population at each of the seven locations showed a weak correlation ($\rho = -0.25$, $p = .5948$). However, the low copy numbers found immediately downstream of the reservoir outfall (Figure 3a) affected this. If this site is treated as an outlier and removed from the analysis, the correlation is significant ($\rho = -1$, $p = .002778$).

3.4 | Passive approach

The total number of reads produced following quality control and chimera removal was 4,928,265, of which 576,365 reads were assigned to the samples in this study. A total of 5.19% reads (29,936)

were assigned to *D. r. bugensis*. After applying a 0.2% threshold to remove minor contamination, the average read depth per sample was 2,921 over all taxa and 548 for *D. r. bugensis* (excluding controls). The clustering procedure identified 81 OTUs revealing a range of macroinvertebrate and algae/diatom species (Supplementary information III, Figure S4). The proportion of reads assigned to *D. r. bugensis* from each sample within each location varied greatly as seen in Figure 3b ($n = 54$). The target species was detected in all samples from both Wraybury Weir and Wraybury Bridge locations, while detection at Wraybury Gardens was much lower with only 3 of the 18 replicates present for *D. r. bugensis* (Table 1). Detection rate was therefore 86% and is lower than the targeted methods. The average raw read count of *D. r. bugensis* was 843 and 801 at Wraybury Bridge and Wraybury Weir respectively, and only 1 read on average at Wraybury Gardens. Distance was the strongest predictor of the relative read count ($Z = 13.47$, $p < .0001$; AIC = 301.86, residual deviance 54.989 on 52 df , $\chi^2 p = .36$) in agreement with the qPCR results and when compared to the distance and density model (distance- $Z = -4.707$, $p < .0001$; Density $Z = -3.773$, $p = .0001$; AIC = 258.64, residual deviance 49.696 on 51 df , $\chi^2 p = .820$), and density on its own (which was overdispersed).

4 | DISCUSSION

This study investigates the detection, persistence, and quantification of the invasive non-native quagga mussel (*D. r. bugensis*) using kick-net sampling and eDNA. Through experimental mesocosms, we validated a species-specific DNA assay (cPCR) to detect quagga mussel and further developed a dye-based quantitative PCR (qPCR) assay to evaluate the performance of both targeted approaches. We carried out an eDNA survey of water samples and compared targeted (cPCR and qPCR) and passive (metabarcoding) approaches to traditional kick-net sampling for qualitative and semiquantitative detection in the River Wraybury, UK. In line with our first hypothesis, detection rate was higher for all three eDNA methods than with the traditional method. Second, targeted approaches were more sensitive than the passive approach (100% detection rates vs. 86%). Contrary to our third hypothesis, we observed no increased sensitivity using a dye-based qPCR over cPCR. Lastly, both density and distance from the source population were significant predictors of relative read count (metabarcoding) and copy number (qPCR) indicating that both approaches could provide estimates of quagga mussel abundance in the field.

4.1 | qPCR validation

Mesocosm validation provides an opportunity to test the LoD of eDNA assays and to investigate the relationship between density, time, and DNA copy number (Dejean et al., 2011; Blackman et al., 2020; Thomsen et al., 2012). DNA was detected with qPCR at all three densities (1, 5, and 20 individuals) within 4 hr of being




FIGURE 3 *Dreissena rostriformis bugensis* DNA signal recorded at field sites. (a) qPCR DNA copy number per μl (\log^{10}) at all 7 sample sites. (b) Normalized relative read count of taxa assigned to quagga mussel at the three locations sequenced

present, in agreement with previous cPCR results (Blackman et al., 2020). The LoD from serial dilutions of tissue DNA was also identical for cPCR and qPCR (1×10^{-4} ng/ μl). Surprisingly, the mesocosm experiments highlighted unexpected dynamics of DNA in closed systems. First, time since the start of the experiment was a significant predictor of DNA copy number, but rather than increasing over time or reaching a plateau, DNA accumulated quickly in the first 48 hr then rapidly depleted, and this trend was seen in all three density treatments. A possible explanation for this is stress induced on the mussel caused by a change in environment, as reported in other eDNA studies (Li et al., 2019). Dreissenid mussels, however, are known to “shut-down” in times of stress, such as chemical dosing, and can remain with their shells closed for up for three weeks (Aldridge et al., 2006). A further explanation for the rapid accumulation in DNA may be due to the production of protein-based byssal threads which allow the mussels to attach to the substrate (Ricciardi et al., 1998; Peyer et al., 2009; Timar &

Phaneuf, 2009; Aldridge et al., 2004). During the first 48 hr of the experiment, the mussels are likely to secure themselves to the new substrate. All mussels were found to have done this when they were removed at the end of the experiment. The mussels are also filter feeders and were actively feeding (on microorganisms found in the river water) throughout the experiment producing sloughed cells, feces, and pseudofeces (Sansom & Sassoubre, 2017). This feeding behavior is likely to be a major contributor to successful detection in the field and, in a closed system, effect the DNA copy number detected with the qPCR assay.

The second surprising result from the mesocosm experiments was that we found no effect of density or biomass on DNA copy number. This contradicts several previous studies that have demonstrated a relationship between biomass or density and DNA copy number (Thomsen et al., 2012; Takahara et al., 2012; Doi et al., 2015). This could in part be due to the physiology and feeding behavior of mussels, since eDNA is being consumed as well as

produced (Friebertshauer et al., 2019). The complex pattern of eDNA accumulation and depletion together with the lack of discrimination between different density treatments presents challenges for estimation of quagga mussel abundance. However, eDNA is unlikely to accumulate in the same way in natural (particularly lotic) environments, and this case highlights that mesocosm experiments may poorly reflect the true relationship between DNA copy number and mussel density in the field. It would be worthwhile to examine whether this pattern is also observed with other filter-feeding species. In keeping with previous studies (e.g., Thomsen et al., 2012), rapid degradation of eDNA was observed in mesocosm experiments, with a complete loss of signal by one week after mussel removal. Unfortunately, our experiments did not allow us to fully investigate the degradation rate of eDNA in mesocosms since our sampling was too limited. We recommend future studies carry out more intensive sampling once specimens have been removed.

4.2 | Comparison of eDNA and kick-net sampling for quagga mussel detection

The lower sensitivity of kick-net sampling is highlighted by the failure to collect a *D. r. bugensis* specimen in any of the first kick-net sampling attempts at each site. As demonstrated in a previous study, species in low abundance are likely to be missed, with only 62% of macroinvertebrate families detected in a three-minute kick-net sample (Furse et al., 1981). This highlights the need for multiple replicates for detection of low abundance species if using kick-net sampling. It should be noted that surveillance methods specifically for *D. r. bugensis* would be better focussed on the preferred habitat of mussels, that is, hard substrates such as man-made structures (Aldridge et al., 2014), which is more likely to detect quagga mussels than the randomized method used here. However, randomized kick-net sampling along a range of substrate types is the most widely used sampling method for freshwater macroinvertebrate monitoring, and therefore, our experiment reflects the most likely methods to detect new INNS through routine monitoring schemes.

Of the three molecular approaches compared in this study, we show that both targeted approaches were more sensitive for detecting *D. r. bugensis* than metabarcoding, but that metabarcoding still performed encouragingly well. Given that the metabarcoding assay targeted metazoa in general rather than the focal species, it is likely that the targeted approach would perform better. Few studies have compared targeted approaches with metabarcoding; however, our findings are similar to those of Simmons et al. (2015) and Harper et al. (2018), who showed that metabarcoding using conserved primers was less sensitive than targeted approaches. In this study, there was no increase in the detection of *D. r. bugensis* when the 0.2% contamination threshold was removed. However, it would be pertinent to examine the raw output of metabarcoding runs before thresholds are applied in order to identify any potential INNS with low read counts. As this form of INNS monitoring is still in its infancy, quality control measures to ensure these detections are correct are yet to

be put in place (Darling et al., 2020). We would therefore recommend any new INNS detections with metabarcoding are followed up with species-specific primers or collection of specimens prior to management actions to verify the detection.

A significant advantage of metabarcoding with conserved primers is the added information from non-target species detection: In our study, a further 81 OTUs were identified, and although no additional INNS were detected, previous studies have demonstrated the power of this approach for detecting new and unexpected INNS (Blackman et al., 2017). Metabarcoding also provides the opportunity to monitor changes in community composition as the result of INNS or other ecological stressors (Simmons et al., 2015; Hänfling et al., 2016), which is extremely promising to better understand INNS impacts and for more effective prioritization and management. This information on the wider community is a justification for the use of universal primers rather than more taxon-specific metabarcoding assays, but if more targeted information is needed, more specific metabarcoding primers can be developed. Indeed, specific mollusk metabarcoding primers have recently been developed and are very promising for more targeted investigation of mollusk communities (Klymus et al., 2017). We hypothesize that more focussed metabarcoding assays will yield even higher detection rates than the one reported here, reducing the difference in sensitivity between targeted and passive approaches. It could also be possible to improve the power of metabarcoding by increasing spatial sampling. Future studies should investigate how many samples are needed in order to maximize the probability of detecting rare species with metabarcoding and also the required read depth needed per sample in order to maximize species detection at a given sampling site.

Conventional PCR and dye-based qPCR had identical detection rates for *D. r. bugensis* in our study. This is somewhat surprising as qPCR is generally considered to be more sensitive than cPCR (e.g., Thomsen et al., 2012; Nathan et al., 2014). Conventional PCR has the advantage of being quick and cost-effective, requiring limited lab equipment and up to half the time for sample processing than qPCR (Davison et al., 2016). Drawbacks to cPCR are typically lower sensitivity, and gel-quality issues may make interpretation of results more subjective than with qPCR. This is dependent on the primer specificity and optimization of the PCR. Our conventional *D. r. bugensis* assay is highly specific and fully optimized, yielding clear bands on agarose gels even in samples with low concentration of target species DNA (as inferred from qPCR). However, we acknowledge that our experience with this assay may be unusual. An obvious advantage to qPCR is the quantitative information it provides, in addition to increased sensitivity. The added information on DNA concentration from qPCR in this study yielded important insights into the mussel density and distance from the source population (see below), which was not possible with cPCR. An important consideration is whether the assay could be improved by using probe-based rather than dye-based qPCR. Addition of a probe is generally considered beneficial to prevent the possibility of non-target amplification. Given the high specificity of the DRB1

primer pair described here, we believe the addition of a probe is not necessary in this case, but we stress that adding a probe should be considered if an extra layer of specificity is needed and/or if non-target amplification is possible.

4.3 | Quagga mussel eDNA declines with distance from the source and decreasing mussel density

Results from field trials for both dye-based qPCR and metabarcoding demonstrate a significant decrease in DNA copy number and relative read count of quagga mussel, respectively, with increasing distance from the main source population (Wraysbury reservoir) and with decreasing mussel density. Kick-net sample data also indicate that mussel density decreases along the course of the river, with only one mussel found at Wraysbury Gardens, 2.7 km downstream of the reservoir. The only site that did not fit this trend was the first site, 20 m downstream of the outfall from Wraysbury reservoir, which had a very low DNA copy number. The decrease in concentration of quagga mussel eDNA detected with distance from Wraysbury Reservoir, confirmed with both qPCR and metabarcoding, reflects both the decreasing density of mussels and a gradient in the transport of DNA downstream. The fact that eDNA concentration is not homogeneously distributed in the river suggests that it may be possible, at least in some cases, to infer the source population in cases where it is unknown. This is somewhat surprising given that our sampling took place after a period of heavy rain and eDNA distribution in rivers is often considered to be uniform and heavily confounded by flow rate and mixing (e.g., Jane et al., 2015). Characteristics including discharge (Jane et al., 2015), physical structure and substrate type (Wilcox et al., 2016; Shogren et al., 2017), and water chemistry (Seymour et al., 2018) all influence the transport and retention of eDNA in lotic systems. Further empirical studies in rivers of all sizes and characteristics, ideally in combination with hydrological modeling (Carraro et al., 2018), are needed to fully understand the influence of flow dynamics and substrate type on retention and transport of eDNA in rivers.

Estimating abundance with eDNA is challenging because of the combined physical, chemical, and biological factors that influence production, transport, and retention of DNA in a water body (Barnes et al., 2014; Barnes & Turner, 2020). Abundance estimation is expected to be particularly challenging in rivers due to their complex hydrodynamics, and the transport of eDNA from upstream (e.g., Jerde et al., 2016). Abundance estimation with metabarcoding is also controversial as read counts are influenced by bias during sampling, laboratory, and bioinformatics steps (Valentini et al., 2016). Despite these challenges, our data add to the growing body of evidence that both qPCR and metabarcoding can yield insight into relative biomass, which has rarely been demonstrated in lotic environments (but see e.g., Doi et al., 2015; Akamatsu et al., 2020). Temporal sampling and controlling for environmental variables, combined with improved statistical approaches such as site occupancy modeling, which is becoming increasingly popular in eDNA studies (e.g., Pilliod et al., 2013;

Schmidt et al., 2013; Chambert et al., 2018; Sales et al., 2020) or spatially explicit approaches that take hydrological conditions into account (e.g., Shogren et al., 2017; Carraro et al., 2018) will provide better quantitative information and allow changes in biomass to be monitored.

4.4 | Conclusions and practical considerations for future INNS biomonitoring with eDNA

We have demonstrated that eDNA from quagga mussels can be successfully detected using cPCR, qPCR, and metabarcoding in a UK river. Both qPCR and metabarcoding provide quantitative information, with DNA copy number and relative read count significantly decreasing with a combination of distance from the source population and mussel density. In our previous work, we demonstrated that metabarcoding is also an effective tool for passive detection of a previously unrecorded cryptic non-native species, *Gammarus fossarum* (Blackman et al., 2017). The present study provides further demonstration that metabarcoding with conserved metazoan primers is suitable as an early warning tool for non-targeted monitoring or routine surveillance even at a very low population density. The optimal approach for detection of INNS using eDNA will depend on the objectives of the research or monitoring program. For early detection of previously unrecorded INNS, we recommend a comprehensive, randomized sampling design, combined with metabarcoding using universal primers, which can be supplemented with targeted approaches if additional verification is needed. Targeted qPCR assays provide optimal sensitivity and specificity and are particularly suited to surveys of INNS that have been previously recorded. In terms of future monitoring of quagga mussels, we believe that the sampling and laboratory methods adopted in the present study are widely applicable, but a temporal study would be beneficial to determine the optimum timing for sampling, and greater spatial coverage is needed to investigate the influence of environmental variables on detection rates.

ACKNOWLEDGMENTS

This project was funded by the Freshwater Biological Association Gilson Le Cren Award 2016 and the UK Environment Agency. We would like to thank staff from the UK Environment Agency—Drs Alice Hiley and Kerry Walsh for their advice and support and Dr Patrycja Meadows and Colette Sales for their help to identify field sites, arrange access and collect specimens. We also thank two anonymous reviewers for helpful and constructive suggestions on a previous version of our manuscript.

CONFLICT OF INTEREST

The authors declare no competing interests.

AUTHOR CONTRIBUTION

Rosetta C. Blackman: Conceptualization (equal); Data curation (equal); Formal analysis (lead); Investigation (lead); Methodology

(lead); Project administration (equal); Validation (lead); Writing-original draft (lead); Writing-review & editing (lead). **Kar Keun Sean Ling:** Methodology (supporting); Writing-review & editing (supporting). **Lynsey R. Harper:** Methodology (supporting); Writing-review & editing (supporting). **Peter Shum:** Methodology (supporting); Writing-review & editing (supporting). **Bernd Hänfling:** Methodology (supporting); Supervision (supporting); Writing-review & editing (supporting). **Lori Lawson Handley:** Conceptualization (equal); Data curation (supporting); Formal analysis (supporting); Investigation (equal); Methodology (supporting); Project administration (equal); Supervision (lead); Writing-original draft (supporting); Writing-review & editing (supporting).

OPEN RESEARCH BADGES



This article has earned an Open Data Badge for making publicly available the digitally-shareable data necessary to reproduce the reported results. The data is available at <https://doi.org/10.5061/dryad.mkkwh70xm>.

DATA AVAILABILITY STATEMENT

Data generated in this study are available from the dryad repository: <https://doi.org/10.5061/dryad.mkkwh70xm>.

ORCID

Rosetta C. Blackman  <https://orcid.org/0000-0002-6182-8691>

Lynsey R. Harper  <https://orcid.org/0000-0003-0923-1801>

Peter Shum  <https://orcid.org/0000-0001-8154-9828>

Bernd Hänfling  <https://orcid.org/0000-0001-7630-9360>

Lori Lawson-Handley  <https://orcid.org/0000-0002-8153-5511>

REFERENCES

- Akamatsu, Y., Kume, G., Gotou, M., Kono, T., Fujii, T., Inui, R., & Kurita, Y. (2020). Using environmental DNA analyses to assess the occurrence and abundance of the endangered amphidromous fish *Plecoglossus altivelis ryukyensis*. *Biodiversity Data Journal*, 8, e39679. <https://doi.org/10.3897/BDJ.8.e39679>
- Aldridge, D. C., Elliott, P., & Moggridge, G. D. (2004). The recent and rapid spread of the Zebra Mussel (*Dreissena Polymorpha*) in Great Britain. *Biological Conservation*, 119(2), 253–261. <https://doi.org/10.1016/j.biocon.2003.11.008>
- Aldridge, D. C., Elliott, P., & Moggridge, G. D. (2006). Microencapsulated BioBullets for the Control of Biofouling Zebra Mussels. *Environmental Science & Technology*, 40(3), 975–979. <https://doi.org/10.1021/es050614%2B>
- Aldridge, D. C., Ho, S., & Froufe, E. (2014). The Ponto-Caspian Quagga Mussel, *Dreissena Rostriformis Bugensis* (Andrusov, 1897), Invades Great Britain. *Aquatic Invasions*, 9(4), 529–535. <https://doi.org/10.3391/ai.2014.9.4.11>
- Ardura, A., Zaiko, A., Martinez, J. L., Samuiloviene, A., Borrell, Y., & Garcia-Vazquez, E. (2015). Environmental DNA evidence of transfer of North Sea Molluscs across Tropical Waters through Ballast Water. *The Journal of Molluscan Studies*, 81(4), 495–501. <https://doi.org/10.1093/mollus/eyv022>
- Barnes, M. A., Turner, C. R., Jerde, C. L., Renshaw, M. A., Chadderton, W. L., & Lodge, D. M. (2014). Environmental Conditions Influence eDNA Persistence in Aquatic Systems. *Environmental Science & Technology*, 48(3), 1819–27. <https://doi.org/10.1021/es404734p>
- Barnes, M. A., & Turner, C. R. (2016). The Ecology of Environmental DNA and Implications for Conservation Genetics. *Conservation Genetics*, 17(1), 1–17. <https://doi.org/10.1007/s10592-015-0775-4>
- Blackman, R. C., Benucci, M., Donnelly, R. C., Hänfling, B., Harper, L. R., Sellers, G. S., & Lawson Handley, L. (2020). Simple, sensitive and species-specific assays for detecting quagga and zebra mussels (*Dreissena rostriformis bugensis* and *D. polymorpha*) using environmental DNA. *Management of Biological Invasions*, 11(2), 218–236. <https://doi.org/10.3391/mbi.2020.11.2.04>
- Blackman, R. C., Constable, D., Hahn, C., Sheard, A., Durkota, J., Hänfling, B., & Lawson Handley, L. (2017). Detection of a new non-native freshwater species by DNA metabarcoding of environmental samples – First Record of *Gammarus Fossarum* in the UK. *Aquatic Invasions*, 12(2), 177–189. <https://doi.org/10.3391/ai.2017.12.2.06>
- Blackman, R. C., Hänfling, B., & Lawson Handley, L. (2018). The use of environmental DNA as an early warning tool in the detection of new freshwater invasive non-native species. *CAB Reviews*, 12(10), 1–15. <https://doi.org/10.1079/PAVSNR201813010>
- Borrell, Y. J., Miralles, L., Do Huu, H., Mohammed-Geba, K., & Garcia-Vazquez, E. (2017). DNA in a bottle—Rapid metabarcoding survey for early alerts of invasive species in Ports. *PLoS One*, 12(9), e0183347. <https://doi.org/10.1371/journal.pone.0183347>
- Brolaski MN, Venugopal RJ, Stolow D (2008) Kits and Processes for removing contaminants from nucleic acids in environmental and biological samples. *US Patent*.
- Brown, E. A., Chain, F. C. C., Zhan, A., Maclsaac, H. J., & Cristescu, M. C. (2016). Early detection of aquatic invaders using metabarcoding reveals a high number of non-indigenous species in Canadian Ports. *Diversity & Distributions*, 22(10), 1045–1059. <https://doi.org/10.1111/ddi.12465>
- Carraro, L., Hartikaninen, H., Jokela, J., Bertuzzo, E., & Rinaldo, A. (2018). Estimating species distribution and abundance in river networks using environmental DNA. *Proceedings of the National Academy of Sciences of the United States of America*, 115(46), 11724–11729. <https://doi.org/10.1073/pnas.1813843115>
- Chambert, T., Pilliod, D. S., Goldbrg, C. S., Doi, H., & Takahara, T. (2018). An analytical framework for estimating aquatic species density from environmental DNA. *Ecology and Evolution*, 8(6), 3468–3477. <https://doi.org/10.1002/ece3.3764>
- Connelly, N. A., O'Neill, C. R., Knuth, B. A., & Brown, T. L. (2007). Economic impacts of zebra mussels on drinking water treatment and electric power generation facilities. *Environmental Management*, 40(1), 105–112. <https://doi.org/10.1007/s00267-006-0296-5>
- Creer, S., Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, W. K., Potter, C., & Bik, H. M. (2016). The Ecologist's Field Guide to Sequence-Based Identification of Biodiversity. *Methods in Ecology and Evolution / British Ecological Society*, 7(9), 1008–1018. <https://doi.org/10.1111/2041-210X.12574>
- Darling, J. A., Pochon, X., Abbott, C. L., Inglis, G. J., & Zaiko, A. (2020). The risks of using molecular biodiversity data for incidental detection of species of concern. *Diversity and Distributions*, 26, 1116–1121. <https://doi.org/10.1111/ddi.13108>
- Davison, P. I., Créach, V., Liang, W.-J., Andreou, D., Britton, J. R., & Copp, G. H. (2016). Laboratory and field validation of a simple method for detecting four species of non-native freshwater fish using eDNA. *Journal of Fish Biology*, 89(3), 1782–1793. <https://doi.org/10.1111/jfb.13086>
- De Ventura, L., Kopp, K., Seppälä, K., & Jokela, J. (2017). Tracing the Quagga Mussel Invasion along the Rhine River System Using eDNA Markers: Early detection and surveillance of invasive zebra and quagga mussels. *Management of Biological Invasions*, 8(1), 101–112. <https://doi.org/10.3391/mbi.2017.8.1.10>

- Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., & Creer, S. (2017). Environmental DNA Metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*, 26(21), 5872–5895. <https://doi.org/10.1111/mec.14350>
- Dejean, T., Valentini, A., Duparc, A., Pellier-Cuit, S., Pompanon, F., Taberlet, P., & Miaud, C. (2011). Persistence of environmental DNA in freshwater ecosystems. *PLoS One*, 6(8), e23398. <https://doi.org/10.1371/journal.pone.0023398>
- Dejean, T., Valentini, A., Miquel, C., Taberlet, P., Bellemain, E., & Miaud, C. (2012). Improved detection of an alien invasive species through environmental DNA barcoding: The example of the American Bullfrog *Lithobates Catesbeianus*. *The Journal of Applied Ecology*, 49(4), 953–959. <https://doi.org/10.1111/j.1365-2664.2012.02171.x>
- Doherty, T. S., Glen, A. S., Nimmo, D. G., Ritchie, E. G., & Dickman, C. R. (2016). Invasive predators and global biodiversity loss. *Proceedings of the National Academy of Sciences of the United States of America*, 113(40), 11261–11265. <https://doi.org/10.1073/pnas.1602480113>
- Doi, H., Uchii, K., Takahara, T., Matsushashi, S., Yamanaka, H., & Minamoto, T. (2015). Use of Droplet Digital PCR for Estimation of Fish Abundance and Biomass in Environmental DNA Surveys. *PLoS ONE*, 10(3), e0122763. <https://doi.org/10.1371/journal.pone.0122763>
- Elbrecht, V., & Leese, F. (2017). Validation and development of freshwater invertebrate metabarcoding COI primers for environmental impact assessment. *Frontiers in Freshwater Science*, 5(11), 1–11. <https://doi.org/10.3389/fenvs.2017.00011>
- Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using environmental DNA from water samples. *Biology Letters*, 4(4), 423–425. <https://doi.org/10.1098/rsbl.2008.0118>
- Friebertshausen, R., Shollenberger, K., Janosik, A., Garner, J. T., & Johnston, C. (2019). The effect of bivalve filtration on eDNA-based detection of aquatic organisms. *PLoS One*, 14(11), e0222830. <https://doi.org/10.1371/journal.pone.0222830>
- Furse, M. T., Wright, J. F., Armitage, P. D., & Moss, D. (1981). An appraisal of pond-net samples for biological monitoring of lotic macro-invertebrates. *Water Research*, 15(6), 679–689. [https://doi.org/10.1016/0043-1354\(81\)90160-3](https://doi.org/10.1016/0043-1354(81)90160-3)
- Gallardo, B., & Aldridge, D. C. (2013). The 'dirty Dozen': Socio-economic factors amplify the invasion potential of 12 high-risk aquatic invasive species in Great Britain and Ireland. *The Journal of Applied Ecology*, 50(3), 757–766. <https://doi.org/10.1111/1365-2664.12079>
- Gallardo, B., Clavero, M., Sánchez, M. I., & Vilà, M. (2016). Global ecological impacts of invasive species in aquatic ecosystems. *Global Change Biology*, 22(1), 151–163. <https://doi.org/10.1111/gcb.13004>
- Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Molecular Ecology Resources*, 13, 851–861. <https://doi.org/10.1111/1755-0998.12138>
- Goldberg, C. S., Sepulveda, A., Ray, A., Baumgardt, J., & Waits, L. P. (2013). Environmental DNA as a new method for Early detection of New Zealand Mudsnailed (Potamopyrgus Antipodarum). *Freshwater Science*, 32(3), 792–800. <https://doi.org/10.1899/13-046.1>
- Hänfling, B., Lawson Handley, L., Read, D. S., Hahn, C., Li, J., Nichols, P., Blackman, R. C., Oliver, A., & Winfield, I. J. (2016). Environmental DNA Metabarcoding of Lake Fish Communities reflects long-term data from established survey methods. *Molecular Ecology*, 25(13), 3101–3119. <https://doi.org/10.1111/mec.13660>
- Harper, L. R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H. C., Gough, K. C., & Hänfling, B. (2018). Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of great crested newt (*Triturus cristatus*). *Ecology and Evolution*, 8(12), 6330–6341. <https://doi.org/10.1002/ece3.4013>
- Holman, L. E., de Bruyn, M., Creer, S., Carvalho, G., Robidart, J., & Rius, M. (2019). Detection of introduced and resident marine species using environmental DNA metabarcoding of sediment and water. *Scientific Reports*, 9(1), 11559. <https://doi.org/10.1038/s41598-019-47899-7>
- Hulme, P. E. (2006). Beyond Control: Wider implications for the management of biological invasions. *The Journal of Applied Ecology*, 43(5), 835–847. <https://doi.org/10.1111/j.1365-2664.2006.01227.x>
- Jane, S. F., Wilcox, T. M., McKelvey, K. S., Young, M. K., Schwartz, M. K., Lowe, W. H., Letcher, B. G., & Whiteley, A. R. (2015). Distance, flow and PCR inhibition: eDNA dynamics in two headwater streams. *Molecular Ecology Resources*, 15(1), 216–227. <https://doi.org/10.1111/1755-0998.12285>
- Jerde, C. L., Olds, B. P., Shogren, A. J., Andruszkiewicz, E. A., Mahon, A. R., Bolster, D., & Tank, J. L. (2016). Influence of stream bottom substrate on retention and transport of vertebrate environmental DNA. *Environmental Science & Technology*, 50(16), 8770–8779. <https://doi.org/10.1021/acs.est.6b01761>
- Karatayev, A. Y., Padilla, D. K., Minchin, D., Boltovskoy, D., & Burlakova, L. E. (2007). Changes in global economies and trade: The potential spread of exotic freshwater bivalves. *Biological Invasions*, 9(2), 161–180. <https://doi.org/10.1007/s10530-006-9013-9>
- Kelly, R. P., Port, J. A., Yamahara, K. M., & Crowder, L. B. (2014). Using environmental DNA to census marine fishes in a large mesocosm. *PLoS One*, 9(1), e86175. <https://doi.org/10.1371/journal.pone.0086175>
- Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA) metabarcoding assays to detect invasive invertebrate species in the Great Lakes. *PLoS One*, 12(5), e0177643. <https://doi.org/10.1371/journal.pone.0177643>
- Lawson Handley, L. (2015). How will the 'molecular Revolution' contribute to biological recording? *Biological Journal of the Linnean Society*, 115(3), 750–766. <https://doi.org/10.1111/bij.12516>
- Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., & Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI Region for Metabarcoding Metazoan Diversity: Application for characterizing coral reef fish gut contents. *Frontiers in Zoology*, 10, 34. <https://doi.org/10.1186/1742-9994-10-34>
- Li, J., Lawson Handley, L. J., Harper, L. R., Brys, R., Watson, H. V., Di Muri, C., Zhang, X., & Hänfling, B. (2019). Limited dispersion and quick degradation of environmental DNA in fish ponds inferred by metabarcoding. *Environmental DNA*, 1(3), 238–250. <https://doi.org/10.1002/edn3.24>
- Mahon, A. R., Nathan, L. H., & Jerde, C. L. (2014). Meta-genomic surveillance of invasive species in the Bait Trade. *Conservation Genetics Resources*, 6, 563–567. <https://doi.org/10.1007/s12686-014-0213-9>
- Mehta, S. V., Haight, R. G., Homans, F. R., Polasky, S., & Venette, R. C. (2007). Optimal detection and control strategies for invasive species management. *Ecological Economics*, 61(2), 237–245. <https://doi.org/10.1016/j.ecolecon.2006.10.024>
- Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., Minamoto, T., Yamamoto, S., Yamanaka, H., Araki, H., Kondoh, M., & Iwasaki, W. (2015). MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: Detection of more than 230 subtropical marine species. *Royal Society Open Science*, 2(7), 150088. <https://doi.org/10.1098/rsos.150088>
- Murray-Bligh, J. A. D. (1999). *Procedure for collecting and analysing macro-invertebrate samples*. Environment Agency.
- Nalepa, T. F., Fanslow, D. L., & Pothoven, S. A. (2010). Recent changes in density, biomass, recruitment, size structure, and nutritional state of Dreissena populations in Southern Lake Michigan. *Journal of Great Lakes Research*, 36, 5–19. <https://doi.org/10.1016/j.jglr.2010.03.013>
- Nathan, L. M., Simmons, M., Wegleitner, B. J., Jerde, C. L., & Mahon, A. R. (2014). Quantifying environmental DNA signals for aquatic invasive species across multiple detection platforms. *Environmental Science & Technology*, 48(21), 12800–12806. <https://doi.org/10.1021/es5034052>

- Peyer, S. M., McCarthy, A. J., & Lee, C. E. (2009). Zebra mussels anchor byssal threads faster and tighter than Quagga Mussels in flow. *The Journal of Experimental Biology*, 212(Pt 13), 2027–2036. <https://doi.org/10.1242/jeb.028688>
- Pilliod, D. S., Goldberg, C. S., Arkle, R. S., & Waits, L. P. (2013). Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples. *Canadian Journal of Fisheries and Aquatic Sciences*, 70(8), 1123–1130. <https://doi.org/10.1139/cjfas-2013-0047>
- Pimentel, D., Zuniga, R., & Morrison, D. (2005). Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics*, 52(3), 273–288. <https://doi.org/10.1016/j.ecolecon.2004.10.002>
- Quinn, A., Gallardo, B., & Aldridge, D. C. (2014). Quantifying the ecological Niche Overlap between two interacting invasive species: The Zebra Mussel (*Dreissena Polymorpha*) and the Quagga Mussel (*Dreissena Rostriformis Bugensis*). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(3), 324–337. <https://doi.org/10.1002/aqc.2414>
- R Core Team (2019). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- Ricciardi, A., Neves, R. J., & Rasmussen, J. B. (1998). Impending Extinctions of North American Freshwater Mussels (Unionoida) Following the Zebra Mussel (*Dreissena Polymorpha*) Invasion. *The Journal of Animal Ecology*, 67(4), 613–619. <https://doi.org/10.1046/j.1365-2656.1998.00220.x>
- Roy, H. E., Peyton, J., Aldridge, D. C., Bantock, T., Blackburn, T. M., Britton, R., Clark, P., Cook, E., Dehnen-Schmutz, K., Dines, T., Dobson, M., Edwards, F., Harrower, C., Harvey, M. C., Minchin, D., Noble, D. G., Parrott, D., Pocock, M. J. O., Preston, C. D., ... Walker, K. J. (2014). Horizon scanning for invasive alien species with the potential to threaten biodiversity in Great Britain. *Global Change Biology*, 20(12), 3859–3871. <https://doi.org/10.1111/gcb.12603>
- Sales, N. G., McKenzie, M. B., Drake, J., Harper, L. R., Browett, S. S., Coscia, I., Wangenstein, O. S., Baillie, C., Bryce, E., Dawson, D. A., Ochu, E., Hänfling, B., Lawson Handley, L., Mariani, S., Lambin, X., Sutherland, C., & McDevitt, A. D. (2020). Fishing for mammals: Landscape-level monitoring of terrestrial and semi-aquatic communities using eDNA from riverine systems. *The Journal of Applied Ecology*, 57(4), 707–716. <https://doi.org/10.1111/1365-2664.13592>
- Sansou, B. J., & Sassoubre, L. M. (2017). Environmental DNA (eDNA) shedding and decay rates to model freshwater mussel eDNA transport in a river. *Environmental Science & Technology*, 51(24), 14244–14253. <https://doi.org/10.1021/acs.est.7b05199>
- Schmidt, B. R., Kéry, M., Ursenbacher, S., Hyman, O. J., & Collins, J. P. (2013). Site occupancy models in the analysis of environmental DNA presence/absence surveys: A case study of an emerging amphibian pathogen. *Methods in Ecology and Evolution*, 4(7), 646–653. <https://doi.org/10.1111/2041-210X.12052>
- Seymour, M., Durance, I., Cosby, B. J., Ransom-Jones, E., Deiner, K., Ormerod, S. J., Colbourne, J. K., Wilgar, G., Carvalho, G. R., de Bruyn, M., Edwards, F., Emmett, B. A., Bik, H. M., & Creer, S. (2018). Acidity promotes degradation of multi-species environmental DNA in lotic mesocosms. *Communications Biology*, 1(1), 4. <https://doi.org/10.1038/s42003-017-0005-3>
- Shogren, A. J., Tank, J. L., Andruszkiewicz, E., Olds, B., Mahon, A. R., Jerde, C. L., & Bolster, D. (2017). Controls on eDNA movement in streams: Transport, retention, and resuspension. *Scientific Reports*, 7(1), 5065. <https://doi.org/10.1038/s41598-017-05223-1>
- Simmons, M., Tucker, A., Chadderton, L. W., Jerde, C. L., & Mahon, A. R. (2015). Active and passive environmental DNA surveillance of aquatic invasive species. *Canadian Journal of Fisheries and Aquatic Sciences*, 73(1), 76–83. <https://doi.org/10.1139/cjfas-2015-0262>
- Smart, A. S., Tingley, R., Weeks, A. R., van Rooyen, A. R., & McCarthy, M. A. (2015). Environmental DNA sampling is more sensitive than a traditional survey technique for detecting an aquatic invader. *Ecological Applications*, 25(7), 1944–1952. <https://doi.org/10.1890/14-1751.1>
- Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata, Z. (2012). Estimation of fish biomass using environmental DNA. *PLoS One*, 7(4), e35868. <https://doi.org/10.1371/journal.pone.0035868>
- Thomsen, P. F., Kielgast, J., Iversen, L. L., Wiuf, C., Rasmussen, M., Gilbert, M. T. P., Orlando, L., & Willerslev, E. (2012). Monitoring endangered freshwater biodiversity using environmental DNA. *Molecular Ecology*, 21(11), 2565–2573. <https://doi.org/10.1111/j.1365-294X.2011.05418.x>
- Timar, L., & Phaneuf, D. J. (2009). Modeling the human-induced spread of an aquatic invasive: The case of the Zebra Mussel. *Ecological Economics*, 68(12), 3060–3071. <https://doi.org/10.1016/j.ecolecon.2009.07.011>
- Tréguier, A., Paillisson, J., Dejean, T., Valentini, A., Schlaepfer, M. A., & Roussel, J. (2014). Environmental DNA surveillance for invertebrate species: Advantages and technical limitations to detect invasive Crayfish *Procambarus Clarkii* in freshwater ponds. *The Journal of Applied Ecology*, 51(4), 871–879. <https://doi.org/10.1111/1365-2664.12262>
- Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., Dejean, T. (2016). Next-Generation Monitoring of Aquatic Biodiversity Using Environmental DNA Metabarcoding. *Molecular Ecology*, 25, 929–942. <https://doi.org/10.1111/mec.13428>
- Venables, W. N., & Ripley, B. D. (2002). *Modern applied statistics with S*. 4th ed. Springer.
- Wilcox, T. M., McKelvey, K. S., Young, M. K., Sepulveda, A. J., Shepard, B. B., Jane, S. F., Whiteley, A. R., Lowe, W. H., & Schwartz, M. K. (2016). Understanding environmental DNA detection probabilities: A Case Study Using A Stream-Dwelling Char *Salvelinus Fontinalis*. *Biological Conservation*, 194, 209–216. <https://doi.org/10.1016/j.biocon.2015.12.023>
- Yu, D. W., Ji, Y., Emerson, B. C., Wang, X., Ye, C., Yang, C., & Ding, Z. (2012). Biodiversity Soup: Metabarcoding of arthropods for rapid biodiversity assessment and biomonitoring. *Methods in Ecology and Evolution*, 3(4), 613–623. <https://doi.org/10.1111/j.2041-210X.2012.00198.x>
- Zaiko, A., Martinez, J. L., Schmidt-Petersen, J., Ribicic, D., Samuiloviene, A., & Garcia-Vazquez, E. (2015). Metabarcoding Approach for the Ballast Water Surveillance—An advantageous solution or an Awkward Challenge? *Marine Pollution Bulletin*, 92(1–2), 25–34. <https://doi.org/10.1016/j.marpolbul.2015.01.008>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Blackman RC, Ling KKS, Harper LR, Shum P, Hänfling B, Lawson-Hanley L. Targeted and passive environmental DNA approaches outperform established methods for detection of quagga mussels, *Dreissena rostriformis bugensis* in flowing water. *Ecol Evol*. 2020;10:13248–13259. <https://doi.org/10.1002/ece3.6921>