

**Physical activity and the menstrual cycle: A mixed-methods study of women's experiences.**

Petra V. Kolić<sup>a\*</sup>, David T. Sims<sup>a</sup>, Kirsty Hicks<sup>b</sup>, Laura Thomas<sup>c</sup> and Christopher I. Morse<sup>a</sup>

*<sup>a</sup>Department of Sport and Exercise Sciences, Manchester Metropolitan University, Manchester, United Kingdom (UK); <sup>b</sup>Department of Sport, Exercise & Rehabilitation, Northumbria University, Newcastle, UK; <sup>c</sup>School of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK*

\*Corresponding author. Department of Sport and Exercise Sciences, Manchester

Metropolitan University, Oxford Road, Manchester M15 6BH, UK. Email:

[P.Kolic@mmu.ac.uk](mailto:P.Kolic@mmu.ac.uk). ORCID ID: <https://orcid.org/0000-0003-3402-1857>

**Physical activity and the menstrual cycle: A mixed-methods study of women's experiences.**

The menstrual cycle is an important biological process in women that is associated with a range of physical symptoms, which can shape how women think, feel, and participate in activities of daily life. This study employed a mixed-methods design to investigate adult women's physical activity throughout the menstrual cycle. One hundred and twenty-eight participants completed online questionnaires that explored events of the menstrual cycle (e.g., bleeding, pain, fatigue) and physical activity. Semi-structured interviews with 21 questionnaire respondents unpacked individual experiences of physical activity throughout the menstrual cycle. From the questionnaire data, 44 participants were categorised as avoiders and 84 as non-avoiders of physical activity due to menstrual events. Avoiders of physical activity reported longer periods, heavier menstrual flow, higher levels of fatigue and pain compared to non-avoiders. Interviews revealed that avoidance of physical activity ranged from complete avoidance to adaptation (e.g., types of exercise). Reasons for avoidance and adaptation of physical activity included menstrual symptoms, personal thoughts, and concerns about other people's views of the period. The present study findings emphasise the importance of recognising women's individual perspectives and established societal norms in addition to physical symptoms in order to better understand and normalise physical activity throughout the menstrual cycle.

Keywords: menstruation; period; questionnaire; interview; symptoms; avoidance; self

35 **Introduction**

36 The menstrual cycle is a natural process in women's reproductive years that represents a  
37 highly individual and personal experience (Brantelid et al., 2014). The start of every  
38 menstrual cycle is marked by the period, also known as menstruation, which can lead to  
39 symptoms, such as bleeding (Santer et al., 2007, 2008), pain (Chen & Hu, 2019), mood  
40 changes (Samy et al., 2019), lethargy and fatigue (Bruinvels et al., 2016). The  
41 symptoms can extend beyond the period and throughout the entire menstrual cycle and  
42 affect how women feel and think about their bodies (Chrisler et al., 2015; Spadaro et al.,  
43 2018). How women feel, think, and act is framed further by normative expectations,  
44 stereotypes, and myths about the menstrual cycle (Kowalski & Chapple, 2000; Marván  
45 et al., 2006). A combination of menstrual symptoms, women's thoughts and feelings,  
46 and social norms could contribute to changes in activities of daily life (Brantelid et al.,  
47 2014; Chen et al., 2016; Houston et al., 2006).

48       The most commonly reported menstrual symptoms represent painful cramps  
49 (dysmenorrhea) and tiredness (Schoep et al., 2019), as well as heavy bleeding  
50 (Bruinvels et al., 2016). On days of the period, the burden of these symptoms has been  
51 shown to prevent women from participating in daily activities (Schoep et al., 2019). For  
52 instance, 43% of adolescent women avoided aspects of daily life due to menstrual  
53 events (e.g., bleeding and pain) of which 21% missed at least one in 30 school days  
54 (Houston et al., 2006). Similarly, 64% of adult women missed an average of 2.6  
55 workdays per month due to severe menstrual bleeding and pain (Fourquet et al., 2010).  
56 Aside from the severity of menstrual symptoms, women's behaviour changes could be  
57 shaped by upbringing (Marván & Molina-Abolnik, 2012), education (Stubbs, 2008), and  
58 media coverage (e.g., advertisements) (Spadaro et al., 2018). Presentations of the period  
59 as a matter of secrecy led women to be vigilant, self-conscious, and selective of daily

activities they undertook and avoided (Johnston-Robledo & Chrisler, 2013). Beyond daily activities, however, limited studies have investigated how women experienced and perceived the menstrual cycle and its impact on sporting performance (Findlay et al., 2020; Moreno-Black & Vallianatos, 2005). A study conducted by Moreno-Black and Vallianatos (2005) with first-year students, who had a history of sport participation, found that women associated menstruation with shame and therefore disguised any associated signs (e.g., sanitary product). The participants were concerned and anxious about the visibility of menstruation (e.g., through leakage) and subsequent, negative public attention. Such negative connotations were perpetuated by comments from parents, teammates, and coaches of young women (Moreno-Black & Vallianatos, 2005). Similarly, Findlay et al.'s (2020) study with international rugby players demonstrated that athletes were highly aware and, at times, worried about the impact of their menstrual symptoms on performance. Although severe symptoms, such as dysmenorrhea, limited the participants during strenuous exercise (e.g., high intensity training), the athletes had established coping strategies (e.g., by accepting their menstrual experience or adapting to it), they had sought advice from medical professionals to manage menstrual symptoms, and spoke to trusted others to process their experiences of the menstrual cycle (Findlay et al., 2020). There remains no comparison, however, how physical activity in a non-athlete population might be affected by not only menstrual symptoms, but also women's individual experiences and social understandings of the menstrual cycle.

Identifying that there is no extent data on the menstrual impact of physical activity, broader comparisons could be made with examples from the effect of pregnancy, adult life transitions, and body awareness on physical activity. In physically active women, who adapted routines throughout pregnancy, decisions whether to

participate in physical activity and, if so, to which extent, depended on the nature and severity of pregnancy-related symptoms (e.g., nausea, fatigue, backache), as well as women's perceptions of their own pregnancy and social expectations regarding physical activity throughout pregnancy (Cioffi et al., 2010). Women self-managed the type, intensity, and amount of physical activity they undertook, thoroughly evaluating societal expectations, information received from health professionals, and myths that relatives and friends shared with them (Cioffi et al., 2010). Similarly, a survey with 14,779 women aged 18–23 years showed that marriage and motherhood, both significant transitions in adult life, put pressure on the time available for physical activity and led to an increase in inactivity (Brown & Trost, 2003). Comparably, body consciousness and the awareness of social norms regarding a desirable body image led women to avoid physical activity (Markland, 2009). The perceived pressure to look a certain way and comparisons to other women in physical activity settings reduced already deflated levels of body satisfaction and affected drop out from physical activity (Pridgeon & Grogan, 2012). This drop out is concerning as it has been shown that physical activity could have a positive effect on dysmenorrhea (Dehnavi et al., 2018) and self-esteem (Zamani Sani et al., 2016).

Within the United Kingdom (UK), 15% fewer women than men aged 16–34 meet aerobic exercise guidelines and 47% of young adult women undertake no structured exercise compared to 32% of men (NHS Digital, 2017). Barriers to physical activity are multifactorial and the lower level of daily physical activity in women could be attributed to numerous factors. It is however impossible to discount that the menstrual cycle, known to influence other aspects of daily life (e.g., school [Houston et al., 2006], work attendance [Fourquet et al., 2010]), may account in part for lower physical activity in women compared to men. One approach to determining whether

physical activity is impacted by the menstrual cycle is to combine quantitative and qualitative methodology and facilitate a holistic understanding of the quantifiable (e.g., pain and flow) and the personal (e.g., experiences and perceptions) (Santer et al., 2007; 2008). Within the context of limited qualitative studies in athletic populations (Findlay et al., 2020; Moreno-Black & Vallianatos, 2005), there remains no broader analysis of physical activity and the menstrual cycle in the general population. The aim of this study was therefore to, (1) quantify events of the menstrual cycle and self-reported physical activity avoidance in women and (2) understand women's lived experiences of physical activity throughout the menstrual cycle. The use of mixed methodology was envisaged to aid an in-depth understanding of the multi-faceted factors affecting physical activity throughout the menstrual cycle, which could usefully inform physical activity recommendations and agendas. We hypothesise based on the avoidance of school (Houston et al., 2006), work (Fourquet et al., 2010) and elite athlete training commitments (Findlay et al., 2020), that women in the present study will show avoidance of physical activity as a result of menstrual events.

## **Methodology**

### ***Design***

The authors utilised a mixed-methods design for the purpose of which they adopted a pragmatic research positioning in line with Morgan (2007). Within this view, emphasis is on (1) drawing on the strengths of quantitative and qualitative methods, (2) following an abductive research process, and (3) acknowledging intersubjectivity. In the present study, quantitative research methods were utilised to quantify the extent to which women experienced menstrual symptoms and participated in physical activity while the

use of qualitative research methods gave insight into the different ways in which individual experiences of the menstrual cycle affected women's physical activity. By working back and forth between quantitative and qualitative components of this study, the authors followed an abductive research process in line with Morgan (2007). Finally, equal value was placed on the worldviews of the research team members, who recognised and accepted that their individual interpretations about "one world" might differ (Morgan, 2007). In doing so, the authors sought to aid the development of the research (Sandelowski, 2000) and the comprehensiveness of the study findings (O'Cathain, 2010).

A questionnaire was developed to examine the frequency and extent of menstrual events as well as self-reported physical activity throughout the menstrual cycle. Semi-structured interviews explored in-depth women's individual experiences and meanings of physical activity and the menstrual cycle (Creswell et al., 2010). Figure 1 visualises how the quantitative and qualitative components were prepared, utilised, and integrated.

To ensure research quality, the researchers employed traditional and alternative criteria (Bryman et al., 2008) and aligned their thinking with the integrative framework by Tashakkori and Teddlie (2008). For design quality, focus was on the suitability and adequacy of research design and methods, on whether the study components flowed in a sound manner, and whether the strategies for data analysis were appropriate to meet the research aims. To achieve interpretive rigor, it was of importance to engage in critical debate and to seek interpretive agreement in the research team, to ensure distinctiveness of interpretations, and adequate integration of quantitative and qualitative results. Finally, the transferability of quantitative and qualitative results into other contexts and

participant groups was prioritised over rigid emphasis on generalisability or context-dependence of the study results (see discussion).

Insert figure 1 here.

### ***Participants***

Following ethical approval from the Institution Review Board, participant recruitment took place via social media (Twitter and Facebook) and leaflets, which were distributed at fitness centres affiliated to the authors' universities by the research team. The authors sought to gain a broad understanding of physical activity and the menstrual cycle from women in the generic population, however, equally adopted a purposeful recruitment strategy, as only women of menstruating age were eligible for participation (Collins, 2010). This approach proved successful as patterns were later observed in the data gathered from the study participants. The recruitment process led to a sample of 128 women of multiple races and ethnicities (27.9 (7.5) years, 1.65 (0.06) m, 63.2 (12.2) Kg, Table 1), who completed the online questionnaire. Through the questionnaire, 82 of 128 participants provided consent to be contacted for further research participation and of these, 38 shared their contact information (email address).

The 38 questionnaire respondents, who had provided contact details, were then invited to a follow-up interview and 21 women took part in interviews. All of whom completed the online questionnaire first and then participated in an interview. Interview invitations and interviews themselves were led by the principal author, one of the female researchers on the team, to facilitate the establishment of rapport and trust



(Miller, 2017), which was considered important to intimate discussions about the menstrual cycle (Dempsey et al., 2016). Interviews were conducted at times and locations that were not only convenient, but also facilitated comfort between her and the participants. Everyone, who stepped forward for an interview, was interviewed and as the final interviews were arranged, saturation had been reached and the research team were confident to halt data collection (Beitin, 2012).

## ***Data collection***

### *Questionnaire*

For the purpose of this research, participants completed a questionnaire about their menstrual cycle duration, menstrual flow, pain or discomfort, and lethargy, combined with questions related to exercise avoidance and self-reported physical activity status. All aspects of the questionnaire were completed online (Jisc, UK). Informed consent was obtained via a compulsory drop down selection at the start of the questionnaire, which was presented following a participant information page. Those who responded to “I do not provide consent for my answers to be used in research” were directed to an exit page of the questionnaire. All questions required a compulsory answer, however two participants selected “not applicable” for their self-reported body mass, which is therefore presented from N = 126. The questionnaire was developed adopting aspects of previous studies that had reported menstrual symptoms, including items from the Menstrual Bleeding Questionnaire (3 items, Matteson et al., 2015), assessment of menstrual pain (1 item, Larroy, 2002), and menstrual flow heaviness (1 item, Fraser et al., 2015).

204

205     Participant demographics

206     Participants completed drop-down selections for age (yrs), height and body mass.

207     Options were given for the height in metres and feet and inches, and for body mass in

208     kg and lbs.

209

210     Hormonal contraceptive

211     Participants were given the option of seven choices related to their use of hormonal

212     contraceptive, from “none” to different forms of hormonal contraceptives, such as oral

213     contraceptive pill (including type and exogenous hormone dosage), patch, injection, or

214     intrauterine devices. For the purpose of data analysis, they are classified as: “None” if

215     they use no form of hormonal contraceptive, “Pill” if they use any form of oral

216     contraceptive, and “non-oral contraceptive” if they use any form of indwelling, injected

217     or cutaneous hormonal contraceptive.

218

219     Menstrual events

220     Participants were asked “Over the last three months, roughly how many days on

221     average has your period lasted?”; these data are reported in the present results as “length

222     of period”. In addition, participants completed a calendar style grid for the symptoms

223     experienced throughout the month for: “bleeding”, “spotting”, “discomfort, cramps or

224     pain” and “lethargy and fatigue”. Within the questionnaire, each of these menstrual

symptoms contained a more detailed description. Participants were instructed to select the days of the month when these symptoms normally occur, with specific reference to the previous month. The frequency of each symptom was summed throughout the month and is presented below as the number of days each symptom occurs during a menstrual cycle (days/MC). Although reliability data is available on retrospective recall for some elements of the menstrual cycle, for instance 80% of women are able to recall their last period date with two days of accuracy (Wegienaka & Baird, 2005) and menstrual cycle length is underestimated by one day compared to using prospective logging (29.4 and 30.9 days, respectively [Small et al., 2007]), the reliability of recalling period length as described above is not presently reported. There is however, “excellent agreement” between daily compared to 1-month recall for menstrual symptoms of bleeding heaviness and pain (Matteson et al., 2015), with the risk of recall bias on these specific elements of the Menstrual Bleeding Questionnaire being described by those authors as “not a problem”. It should also be noted that 52% of our participants self-reported that they tracked their menstrual and pre-menstrual events through an App or diary, and 88% reported that they were somewhat or very confident that they “can accurately recall [their] menstrual cycle dates and pre-menstrual symptoms”.

#### Heavy or normal menstrual bleeding

Participants were classified as “heavy” menstrual bleeders based on the selection of two or more of the following symptoms, consistent with Fraser et al. (2015): 1) a need for double sanitary products (e.g., tampons and towels) at the same time, 2) a need for frequent changes of sanitary towels or tampons (every two hours or less, or 12 sanitary

items per day), 3) bleeding through sanitary products onto clothes or bedding, and 4) the presence of large clots within period blood. A final option of “none of the above” was also included. Participants reporting one or none of the above symptoms were classified as “normal” menstrual bleeders (Fraser et al., 2015). These terminologies are consistent with the International Federation of Gynecology and Obstetrics (FIGO) systems for nomenclature of symptoms of normal and abnormal uterine bleeding (Fraser et al., 2011). Despite the accepted terminology and classification of heavy menstrual bleeding, there is presently no reliability data available for the classification of “heavy menstrual bleeding” as established by Fraser et al. (2015). Concurrent validity has however been previously reported based on menstrual symptom severity being higher in women who are classified as experiencing heavy compared to normal menstrual bleeding (Matteson et al., 2015), with daily and monthly flow scores showing excellent agreement ( $\rho = 0.82$ , Matteson et al., 2015).

## Pain

Participants were provided with a numerical scale from 0-10 (Larroy, 2002), where 0 was labelled “no pain”, 5 was labelled “moderate pain” and 10 was labelled “worst possible pain”. Within the questionnaire, a visual analogue scale was also provided under “more info”. Participants were classified into “mild pain” if pain was between 1 and 3, “moderate pain” between 4 and 7 points, and “severe pain” between 8 and 10 points (Kural et al., 2015). Due to there being insufficient “avoider” participants (see below) falling within the “no pain” category, the “no-pain” and “mild pain” were compressed into a “no-to-mild pain” category representing all participants reporting pain of  $\leq 3$ . Analysis of menstrual pain was conducted based on both continuous data

from the numerical scale (termed pain severity, Table 1), and pain classification.

#### Avoidance

Participants were asked “In the last 3 months, which, if any, of the following activities have you avoided or postponed due to menstrual events or discomfort?” with the options provided of “social activities”, “work or university”, “playing sport, going to the gym, or other physical activities” and “none”. An open text option of “other” was also provided. Where these included recreational activities such as “swimming”, “walking” or “hiking” they were categorised in the “physical activity” avoidance response if this was not already marked in the affirmative. Due to the focus of the present study, participants were classified as “avoiders” and “non-avoiders” based on whether they had reported to have previously avoided physical activity due to the events of the menstrual cycle. The impact of menstrual symptoms on physical activity avoidance was based on similar single question components of absenteeism from work and school (Fourquet et al., 2010; Houston et al., 2006).

#### Physical activity

Self-reported physical activity status was assessed based on the following criteria: sedentary (walking less than 20 mins a day); slightly active (walk over 20 mins per day); moderately active (undertake at least 20 mins of moderate physical activity per day); very active (undertake 40 mins of moderate intensity physical activity per day); athlete (high intensity exercise 5+ days a week). For the purpose of subsequent analysis,

as there were fewer than five participants within the sedentary and athlete categories, the groups were condensed with their closest categories. The physical activity classifications were therefore: sedentary-to-low activity, moderately active, high activity-to-athlete. Despite retrospective recall having known limitations for accurately quantifying daily physical activity (Lee et al., 2011), a single item physical activity question is a valid approach for the purposes of participant classification as adopted in the present study (Milton et al., 2013; Schechtman et al., 1991), with high test-retest reliability ( $r = 0.82$ , Milton et al., 2011).

#### *Interviews*

Semi-structured interviews were conducted to (1) expand on the data obtained from questionnaires (O’Cathain, 2010), (2) to facilitate the generation of new ideas that might not have emerged from the sole use of quantitative methodology, and (3) to uncover the subtleties underpinning participant perceptions ‘in diverse and enriched ways’ (Smith & Sparkes, 2016, p. 3). They were useful to cover aspects important to the inquiry, while rendering the flexibility to pose impromptu questions that encouraged participants to express thoughts freely (Brinkmann & Kvale, 2018).

In this study, 21 interviews were conducted, of which seven took place face-to-face and 14 via video call (FaceTime or WhatsApp). Prior to interview participation, the women gave written consent to voluntary interview participation, to audio recording of interviews, to transcription of these audio recordings, and to the use of anonymised, direct quotes in published work (e.g., conference presentations, journal articles). Interviews were informed by an interview guide, which outlined topics of interest and example questions. Following a summary by the principal author of the purpose of

interviews, conversations then focused on understanding women's physical activity throughout the menstrual cycle (Warren, 2012). Interview questions explored how the participants felt on days before, during, and after the period, what the participants' physical activity looked like throughout the menstrual cycle, how participants felt about themselves throughout the menstrual cycle, how menstrual symptoms, self-perceptions, and social situations shaped physical activity on days of the period, and why the participants thought, felt, and acted in certain ways. Participants were prompted to expand on initial thoughts with questions, such as "How did this make you feel?", "Why do you think this way?" or "Can you give me an example?" Interviews were audio-recorded and transcribed by the principal author for further interpretation (Davidson, 2009).

### ***Data analysis***

#### ***Questionnaire***

All quantitative analyses were performed using IBM SPSS Statistics 24 software. Where appropriate (i.e. ratio data), parametric assumptions of normal distribution were confirmed using Shapiro-Wilk's test ( $p>0.05$ ) in all dependent variables, except for height in both avoiders and non-avoiders ( $p<0.05$ ). Group comparisons were made using independent t-tests with equal variance (Levene's,  $p>0.05$ ) in all dependent variables, other than pain (days/MC), fatigue (days/MC), and spotting (days/MC) (Levene's  $p<0.05$ ). For comparison of group differences in height, the Mann-Whitney U test was performed. All data are presented as mean (SD), with group differences reported as:  $p$ , 95% confidence interval (CI) and effect size ( $d$ ).

Nominal data was assessed using Chi-square associations for classifications of pain, flow, contraception, and physical activity. Participants were grouped as avoiders or non-avoiders, with subsequent post-hoc analysis performed if the Chi-square reached significance ( $p < 0.05$ , for pain and flow). As only two classification were made for flow, no post hoc was necessary. Whereas for pain, there were three classifications (no-mild, moderate, and severe), in this instance cell-wise residual analysis was performed (Garcia-Perez & Nunez-Anton, 2003), with the level of significance adjusted for the three sub-classifications. For the nominal variables, a significant outcome of Chi square is described as a significant association rather than a group difference based on previous recommendations (Field, 2013; McHugh, 2013; Scott et al., 2013).

### *Interviews*

Thematic analysis was utilised to identify patterns in the interview data and make sense of their meaning (Braun et al., 2016). The principal author read anonymised interview transcripts recurrently to define codes, which constituted of links between data and ideas, and themes that helped identify commonalities and distinctions (Nowell et al., 2017). She recognised this phase as an active process of meaning making and adopted an iterative approach by working “back and forth” between interview data, preliminary themes, and interpretations (Braun et al., 2016). Analytical questions included “What do the participants’ physical activity routines look like?”, “How do menstrual events affect the participants’ physical activity? Why is this so?”, “How do the participants think and feel about factors affecting physical activity? Why do they think and feel in certain ways?” To strengthen the credibility of interpretations, the principal author liaised with members of the research team, who acted as critical friends and discussed



interpretations, prompted reflection, and explored interpretive avenues (Smith & McGannon, 2018). In the latter stages of analysis and write-up, the dialogues also explored opportunities to integrate the insights gained from questionnaires and interviews to develop rich discussions of physical activity and the menstrual cycle.

## Results

### *Questionnaire results*

Of the 128 responders, 44 participants (34%) were classified as “avoider” and 84 participants (66%) were classified as “non-avoider” based on whether they had avoided playing sport, going to the gym, or other physical activities due to their menstrual events or discomfort.

### *Participant demographics*

There was no significant difference in age, height or body mass between avoiders and non-avoiders (Table 1).

### *Menstrual characteristics*

Compared to non-avoiders, avoiders had periods that lasted 0.65 days longer ( $t(126) = -2.34, p < 0.05, CI [0.10, 1.20], d = 0.40$ , Table 1). In terms of specific menstrual symptoms, compared to non-avoiders over the course of a menstrual cycle, avoiders had 0.94 more days of bleeding ( $t(126) = -2.464, p < 0.05, CI [0.18, 1.71], d = 0.46$ ), 0.98

more days of fatigue ( $t(126) = -4.769, p < 0.01, CI [1.30, 3.61], d = 0.33$ ) and 2.44 more days of pain ( $t(126) = -4.191, p < 0.01, CI [1.28, 3.61], d = 0.84$ , Table 1). There was no difference in days of spotting between avoiders and non-avoiders (Table 1).

Pain severity (numerical scale) was 62% higher in avoiders than non-avoiders ( $t(126) = -4.116, p < 0.01, CI [1.09, 2.96], d = 0.78$ , Table 1). There was no significant association between contraception use and physical activity avoidance ( $\chi^2 (2, N = 128) = 2.06, p = 0.35$ ); with 61% and 57% using no form of hormonal contraceptive, 18.2% and 11.9% using non-oral contraceptive, and 20.5% and 31% using some form of oral contraceptive pill, in avoiders and non-avoiders, respectively.

There was no significant association between physical activity classification and avoidance ( $\chi^2 (2, N = 128) = 2.73, p = 0.26$ ), with 20.5% avoiders and 11.9% non-avoiders being sedentary or low physical activity, 54.5% avoiders and 51.2% non-avoiders being active, and 25% avoiders and 36.9% non-avoiders being very active or athlete.

There was a significant association between pain classification and avoidance ( $\chi^2 (2, N = 128) = 14.5, p < 0.01$ ). Post hoc revealed that significantly less avoiders were in the “no-to-mild” pain classification (20.5%) than non-avoiders (52.4%,  $p < 0.01$ ). Similar participant numbers were found for avoiders and non-avoiders in the moderate pain classification (59.1% avoiders, and 41.7% of non-avoiders), and in the severe pain classification (20.5% of avoiders and 6% of non-avoiders,  $p < 0.04$ , no significant difference at the adjusted  $\alpha = p < 0.017$ ).

There was a significant association between menstrual flow and avoidance ( $\chi^2 (1, N = 128) = 22.3, p < 0.01$ ), 63.6% of avoiders were classified as heavy flow, compared to 21.4% of non-avoiders. Similarly, 36.4% of avoiders were classified as normal flow compared to 78.6% of non-avoiders.

412

413 Insert table 1 here.

414

415 ***Interview results***

416 Following on from the questions posed throughout the data analysis, the results explore  
417 women's experiences and management of physical activity as well as factors affecting  
418 physical activity throughout the menstrual cycle, including menstrual symptoms,  
419 personal perspectives, and social expectations. Throughout this section, pseudonyms  
420 were used to protect the identity of participants.

421

422 *Periods and physical activity: Realities and experiences*

423 The women, who took part in interviews, spoke openly about the menstrual cycle and  
424 shared personal experiences. Although participants referred to the period as the  
425 menstrual cycle phase that had the greatest impact on physical activity, they equally  
426 recognised the value of physical activity when coping with menstrual symptoms. It  
427 became apparent that their perspectives were informed by an understanding of the  
428 effects that physical activity had on physical and mental health. For instance, Val, Lisa,  
429 and Elaine explained:

430 I think exercise has been a big help with my mental health. Even going for a walk.  
431 I'll consciously do it because it makes me feel better. I feel lazy and more bloated  
432 if I don't move much. (Val, Non-avoider)

## Physical activity and the menstrual cycle

433 I think it's good for your longer term health. I know that people say exercise helps  
434 with your periods. (Lisa, Avoider)

435 I think to begin with I was quite emotional because I wasn't active. With being  
436 more consistently physically active, my mood swings are not bad at all. (Elaine,  
437 Non-avoider)

438 Not only did the participants recognise the benefits of physical activity, but they also  
439 sought to undertake it in everyday life. These women reported continuous physical  
440 activity throughout the menstrual cycle and, in their recollections, demonstrated an  
441 instilled sense of commitment. As an example, Olivia (Non-avoider) described why she  
442 pushed herself to exercise:

443 I force myself to do things. I'm very active, so I tend to stay physically active. I  
444 thank myself afterwards. It makes things better, especially the cramping and  
445 sluggishness. All I want to do is tuck up in a ball on my bed. So forcing myself to  
446 go to the gym is worth it because I do feel so much better after.

447 Sharing a perception of the period as something that should not limit activities of  
448 everyday life, some of the participants felt as though they had no reason to skip the  
449 gym. As a former professional dancer, Ava expected herself to commit to physical  
450 activity even on days of the period. Throughout her career as a professional dancer, she  
451 had developed a strong sense of commitment to routine training, which she had  
452 transferred into her approaches to physical activity (Stephan, 2003). Holding high  
453 expectations of herself, Ava (Non-avoider) differentiated the period from health  
454 problems or injuries and explained:

455 There's no injury or pain that's preventing me from going. I just feel unwell. I'd  
456 rather not go when I'm too tired or something is wrong with me. I think, "Okay  
457 well at least you went and maybe don't go another time." There's no actual  
458 physical reason for me not to go. It's a subjective feeling.

459 Likewise, Nina (Non-avoider) described that she sought continuity in her routines:

460 I usually have a day at the start of my period, when I'm a bit off. If I planned a  
461 session, I would do it and leave it behind me. I'm keen to follow my routines  
462 because I don't think that my periods are strong enough. There is no need for me to  
463 change things. I'm always challenging my brain whether I need to change  
464 something or whether I can try it.

465 It is noteworthy that the above-presented data extracts stem from interviews with  
466 women, who described their menstrual symptoms as manageable and therefore did not  
467 feel the need to avoid physical activity. Many participants, however, did describe  
468 avoidance of behaviours; either of physical activity environments (e.g., the gym) or of  
469 exercises (e.g., abdominal work) and types of exercise (e.g., cardiovascular training).  
470 The majority of conversations in this context therefore focused on the adaptations made  
471 to maintain some level of physical activity. None of the participants scheduled  
472 avoidance of physical activity due to menstrual events. Rather, symptoms had to be  
473 managed *in situ* and often left the participants feel unable to follow their "normal"  
474 physical activity routines. As an example, Willow (Non-avoider) changed her workouts  
475 to avoid the frustration she associated with not being able to perform the way she would  
476 on days when she did not have the period. She explained:

477 On my period, I don't feel like my workouts are as productive. I will swap what I  
478 do. I tend to focus more on upper body to avoid the annoyance I have if I have a  
479 bad session and I'd probably stay away from cardio.

480 For Hannah (Avoider), the first day of the period usually meant avoiding physical  
481 activity completely:

482 Sometimes I get home from work on the first day and think, "Forget it, not this  
483 week", and just stay at home. Then on later days, when I do decide to go for a run

## Physical activity and the menstrual cycle

484 or do exercise, I wouldn't put as much effort in. I'd choose an easier class rather  
485 than a more intense one.

486 For Scarlett and Harper, the discomfort of using sanitary products led them to alter  
487 routines:

488 I do a lot of running and I don't really do that when I'm on my period. The  
489 sensation feels weird, so I stick to weight bearing exercises. The feeling's not there  
490 as much when I do that. I don't wear tampons because I find them uncomfortable,  
491 so I wear sanitary towels and I always wonder, "Oh have they moved?" ... it's  
492 about where the pad is and it can get sore. It doesn't stop me but instead of doing  
493 half an hour, I might only do ten minutes of cardio. Or I'd sit on a bike instead of  
494 running because it's not moving as much. (Scarlett, Non-avoider)

495 I'd go to the gym, but perhaps not do cardio. Because of my injured knee, I can't  
496 run and sitting on a bike with a pad or tampon is really uncomfortable. That would  
497 ruin my gym session. So, sessions don't last as long. I probably do more upper  
498 body than lower body. I don't think I do abs when I'm on my period. Especially  
499 crunches or sit-ups when you've got a pad on. It's not nice at all. (Harper, Avoider)

500

### 501 *Symptoms, self, and social – Factors shaping physical activity*

502 Interview participants recalled a variety of factors affecting the ways in which they  
503 managed physical activity throughout the menstrual cycle. Their suggestions broadly  
504 related to the symptoms associated with the period, to perceptions of self, and to social  
505 expectations.

506 The most prominent influence on women's physical activity routines was the  
507 perceived severity of menstrual symptoms. Although the types of symptoms, their  
508 duration, and occurrence varied, they commonly included abdominal cramps, headaches

509 and backaches, leg pains, lethargy and fatigue. The following data extracts are examples  
510 of the symptoms described by interview participants:

511 I suffer most with sciatica-type leg pain. I'm extremely lethargic to the point where  
512 it's almost debilitating. Migraines as well, feeling dizzy and muggy. (Piper,  
513 Avider)

514 Normally I'm an energetic person, but when it's that time I'm tired. I can be  
515 moody, but it's mainly tiredness. I don't feel I have my normal amount of energy.  
516 (Hannah, Avider)

517 My body changes completely. When I put on a pair of trousers that are normally  
518 loose, they're tighter. My stomach feels heavy. And the fatigue! (Mary, Avider)

519 There were a couple of times, where I woke up in the morning and planned to go  
520 for a run, but then the cramps were so bad that I couldn't get out of bed. (Olivia,  
521 Non-avider)

522 In addition to feeling physically unwell, many participants also spoke of a heightened  
523 sense of self-consciousness. The period affected how women thought about themselves  
524 and how they felt in their own skin. Interestingly, these perceptions were not only  
525 limited to those who avoided (certain aspects of) physical activity, but were also shared  
526 by those who made efforts to maintain activity levels throughout the menstrual cycle.  
527 As an example, Harper (Avider), described:

528 You're not your usual self. You just feel you're on the period. For me it's at the  
529 start, but at that point, it's always there that thought. Just like hyper aware of it.

530 For Val and Caroline, feelings of consciousness led them to carefully select clothes  
531 worn in physical activity environments. The participants recalled, respectively:

## Physical activity and the menstrual cycle

532 I'm really conscious if I wear shorts or if my leggings are see through or thinking  
533 whether I've leaked. I'm more aware of what I do and how that might look. (Val,  
534 Non-avoider)  
535 I suppose with netball, I am conscious of what to wear because of the small shorts  
536 and dresses that we have to wear. I always wear a tampon over a towel and  
537 underwear that keeps everything in check. I actually saw a girl at netball a couple  
538 of years ago. You could see her pad through the shorts and I felt bad for her but  
539 nobody said anything. (Caroline, Non-avoider)

540 Beyond the sense of self-awareness, the perceived expectations others might have about  
541 menstrual events, shaped women's physical activity routines. Interview participants  
542 commonly referred to worries of being "found out" by others, recalled an awkwardness  
543 surrounding periods, and described subsequent discomfort in certain physical activity  
544 environments. For example, Lisa and Scarlett preferred to avoid the gym on days of  
545 heavier flow due to concerns over how others might react if they discovered the reasons  
546 for which they did not adhere to usual routines:

547 The gym is a social place for me. I know a lot of people. They would know I  
548 wasn't on top form. My PT would say, "Why are you lifting this today?" I  
549 wouldn't wanna turn around, "I feel shit because of my period." I don't know how  
550 he would handle that, what his feelings would be. That's where I suppose that  
551 taboo comes in because I wouldn't go, "I'm taking it easy today because I'm on  
552 my period." Even though I say I talk about periods, I guess I don't because I'd  
553 rather avoid telling them that I was going easy because of my period. (Lisa,  
554 Avoider)

555 In the first days when I'm heaviest, I feel embarrassed. I don't want other people to  
556 know. I'm scared that if I get out of the pool, blood is gonna run down my leg. I'm  
557 worried what other people would say. Especially with men in the gym. Someone  
558 might put me down and I wouldn't wanna go to the gym anymore. (Scarlett, Non-  
559 avoider)

560 Other participants perceived pressure to maintain their attendance at training sessions



and believed there were expectations for women to carry on “as normal” regardless of discomfort due to menstrual events. For Hannah and Harper, playing netball was not only important at a personal level, but also in a social context (Weiss, 2015). It was therefore to be continued throughout the menstrual cycle. The participants’ perceptions of expectations and understandings that teammates and coaches might hold shaped their dedication to the sport (Weiss et al., 2010). They explained:

With netball, you’ve got to go. There’s no choice. In my netball team, if you don’t train, you don’t get played at the match. You’ve got that pressure. Also we pay for training and matches, so you don’t want to waste money by not going. (Hannah, Avoider)

When I played netball, I couldn’t turn around to the team and say, “I can’t play. I’m on my period.” Someone might say, “I’m on my period too and I’m playing.” Especially in a team of girls. You can’t let them down when they all know how it feels. (Harper, Avoider)

Similarly, Mila and Nina believed that it was expected of women to disguise any signs of menstrual events. In Mila’s (Non-avoider) words:

When I’m on the period I don’t want anyone to find out I’m not feeling hundred percent. Maybe I complain to a female co-worker or friend but I would never, even, if I called in sick, I wouldn’t admit it’s period-related. I would say I am not well. I think the expectation to carry on as usual is always there.

Nina (Non-avoider) explained:

Nobody ever told me that I need to hide my tampon when I go to the loo at the gym, but I do it. Because I think, other people aren’t as relaxed as me about periods. We don’t talk about them. If we do, it’s awkward.

## Discussion

This mixed-methods study utilised an online questionnaire to quantify events relating to the menstrual cycle and physical activity as well as semi-structured interviews to explore women's lived experiences of physical activity throughout the menstrual cycle. The discussion draws together quantitative and qualitative results to provide rich interpretations of the questionnaire and interview data (O'Cathain, 2010).

In the present study, the majority of women who completed the questionnaire (84 of 128 women – Table 1) were categorised as non-avoiders of physical activity due to events of the menstrual cycle. Interview data revealed that the participants recognised the importance of physical activity and the benefits it brought to their lives. Through experiences in their upbringing, education, and employment, they had learned about the advantages of physical activity, which in turn affected their experiences of and commitment to physical activity throughout the menstrual cycle. Consistent with the knowledge that the present participants possessed, previous research has shown that connections existed between people's awareness of physical activity recommendations and their actual levels of physical activity (Plotnikoff et al., 2011). Those who knew about health recommendations tended to be more physically active in everyday life compared to those, who did not possess this level of understanding (Fredriksson et al., 2018). This was so especially when individuals knew not only about guidelines, but also about the health benefits associated with moderate physical activity (Heinrich et al., 2011). Interview data in the present study revealed that the participants had actually experienced positive post-workout effects on days of the period, for instance, uplifted emotional state, reduced pain, and increased feelings of accomplishment. This finding is in line with studies illustrating that exercise could alleviate menstrual symptoms (e.g., backaches – Chen & Hu, 2019) and enhance mental wellbeing throughout the menstrual

cycle (Kanojia et al., 2013). Physical activity has also been considered as a complementary treatment for dysmenorrhea (Samy et al., 2019) as it reduced the duration and severity of menstrual cramps (Dehnavi et al., 2018; Kannan et al., 2015).

In partial agreement with our hypothesis, one third of questionnaire respondents (44 of 128 participants) were categorised as avoiders of physical activity due to menstrual events (Table 1). This could be attributed to these avoiders having more severe menstrual symptoms, including longer periods, higher levels of pain and fatigue, and heavier flow as reflected in the questionnaire data. Despite the potential bias of menstrual recall (as discussed in detail in the methods) the qualitative data confirmed that women could not overlook menstrual events (e.g., bleeding and pain) and when considering whether to avoid physical activity, interviews revealed that the participants evaluated the severity of menstrual symptoms. For instance, symptoms that the women perceived as severe (e.g., migraines and abdominal pain) led the participants to avoid physical activity completely. This complete avoidance is comparable to recurrent school absence in adolescent women who suffered from dysmenorrhea (Houston et al., 2006). In adult women, missing university or workdays represented a common self-management strategy for painful periods (Chen et al., 2016). The present study participants confirmed that the period had to be managed (e.g., sanitary products) and taken into account in physical activity routines (e.g., types of exercises). The women in this research described the days of the period as limiting their ability to function as what they perceived “normal” for themselves. In particular, a combination of physical symptoms, lacking “energy” and motivation made it more challenging for the participants to perform physical activity as normal. These findings resonate with studies on the impact of the menstrual cycle on physical performance, such as through anaemia from heavy menstrual bleeding (Bruinvels et al., 2016), higher body temperature during

exercise in the luteal phase (De Jonge & Janse, 2003), and lower fatigue resistance in the pre-menstrual phase (Ansdell et al., 2019).

The present study participants, however, did not necessarily avoid all forms of physical activity on days of the period (a nuance not captured within the questionnaire). Although the questionnaire data showed no significant relationship between physical activity classification and avoidance, interviews were particularly useful to make sense of this observation and to unpack the subtleties underpinning women's reports of physical activity avoidance due to menstrual events without necessarily being less physically active overall. Indeed, interviews uncovered aspects of physical activity adaptation that have been previously unreported. Some of the participants, who reported to avoid physical activity, did not avoid all forms of activity and instead, changed exercise routines, used alternative physical activity or made up for missed physical activity on days when they felt less affected by menstrual symptoms. Often, it was about adapting routines depending on the perceived severity of menstrual symptoms and levels of discomfort, consistent with previous work showing adaptations to activities of daily life due to menstrual symptoms (Kennett et al., 2016). The selective avoidance and modification of physical activity was routine to some participants and could reflect women's knowledge about the benefits of physical activity to their health, physical wellbeing, and mood (Fredriksson et al., 2018). Similar to women who selectively avoided tasks in the workplace (e.g., meetings) due to reduced concentration, patience, and efficiency (Brantelid et al., 2014), some of the present interview participants only avoided physical activity that caused discomfort (e.g., lower body workouts), while continuing more manageable activities (e.g., walking and upper body workouts), when they believed that they could maintain some level of activity. This selective avoidance of physical activity is comparable to reports of adaptation as a way to maintain physical

activity levels in other populations, such as pregnant women suffering from backaches (Cioffi et al., 2010) as well as individuals suffering from chronic fatigue syndrome (Larun & Malterud, 2011) and migraines (Buse et al., 2012).

Aside from the menstrual symptoms, our qualitative findings demonstrated how self-perceptions and perceptions of social expectations influenced physical activity.

Interviews revealed that the participants had concerns about leakage of blood, product use, and pain management in the presence of other people in physical activity

environments. Similar to women, whose commitment to exercise suffered from heightened body consciousness in physical activity environments (Pridgeon & Grogan,

2012), several women in the present study described a heightened sense of self-

awareness on days of the period that led to careful selection of clothes and exercises.

Comparably, heightened consciousness in public spaces was previously described in

women, who felt their bodies did not conform to predefined norms of femininity, such

as female bodybuilders in the presence of casual gym visitors, in public (Shilling &

Bunsell, 2009), and when sharing photos of their physique on social media (Marshall et

al., 2019). In the present study, the perceptions of self in physical activity environments

were closely connected to concerns over what others might say if they found out about

women's menstrual events. Sensations, such as bleeding, pain, and fatigue, associated

with the period in particular, were to be "dealt with" in private. This persisted even

though the interview participants had reframed their own views of menstrual events and

felt comfortable to talk about them (Lee, 2002). The participants believed, however, that

upbringing and societal norms of the period in particular as something shameful and

embarrassing led women to silence and conceal (signs of) the period (e.g., blood and

sanitary product) (Chrisler et al., 2015; Spadaro et al., 2018). The concerns of women in

the present study about how others might react to signs of the period point towards

social stigma surrounding menstrual events that led women to avoid and adapt physical activity (Johnston-Robledo & Chrisler, 2013; Kowalski & Chapple, 2000). Such stigma surrounding exercise during the period has been similarly reported in overweight individuals (Vartanian & Novak, 2011), people with mobility impairments (Mulligan et al., 2017), and the LGBTQ+ community (Herrick & Duncan, 2018) undertaking physical activity.

Due to the paucity of data on the impact of the menstrual cycle on physical activity, in order to test our proposed hypothesis, it was necessary to construct the components of the questionnaire from a number of different sources. These sources have been acknowledged throughout the methods section including recall of menstrual symptoms (Matteson et al., 2015), menstrual pain (Larroy, 2002), menstrual flow heaviness (Fraser et al., 2015), avoidance (Fourquet et al., 2010), and physical activity (Milton et al., 2011). Through adopting these previously validated components of the questionnaire, we can state our main finding that physical activity avoiders have heavier, longer, and more painful periods, despite the limitation that no assessment of reliability or validity has been performed on the entirety of the questionnaire. It is important to note however, that based on the nuanced responses from the interviews, emphasising adaptation, over avoidance, future questionnaires on physical activity within the menstrual cycle should be validated for both avoidance and adaptation.

In developing our questionnaire, our aim was to identify whether physical activity avoidance could be attributed to menstrual symptoms. There are a wide range of demographic co-variables that are unreported in the present study including education, employment status, religious background, socio-economic level and ethnicity (e.g., Mondragon & Txertudi, 2019). As the area of physical activity avoidance due to menstrual symptoms is underreported in general, our aim was to provide the first

overview of the impact of the menstrual cycle on physical activity. In future studies, the prevalence of physical activity avoidance may therefore be different to the present study, if presented based on some of these demographic groupings.

Our study demonstrated that binary measures of physical activity, such as those utilised in the present questionnaire, miss the nuances that underpin avoidance behaviours. The use of quantitative and qualitative methodology was therefore particularly useful to developing nuanced views of factors affecting physical activity, which included, but were not exclusive to, menstrual symptoms. Other important considerations that shaped women's physical activity related to perceptions of "self" and social expectations about menstrual events; two areas that need to be confronted in greater depth. Future research might therefore seek to employ qualitative methodology to understand women's self-presentations (e.g., Goffman, 1956) and the role of social stigma (e.g., Goffman, 1963) in physical activity environments. The participants in the present study understood the importance of physical activity to maintain a healthy lifestyle and they enjoyed being physically active. We acknowledge, however, that the limited sample size and volunteer nature of interview participation could be considered limitations of this study. Although comparable sample sizes were suggested in methodological texts (Onwuegbuzie & Collins, 2007) and mixed-methods research investigating the menstrual cycle (Moreno-Black & Vallianatos, 2005) and voluntary interviews were conducted to ensure ethical research practice (e.g., Hammersley & Traianou, 2012), we recognise that our decisions might have resulted in unreported differences between the present participants and those, who did not volunteer for interview participation. As with the limitations of the questionnaire outlined above, a worthwhile line of inquiry would therefore be to investigate how other demographic groups approach physical activity throughout the menstrual cycle. In particular, the

impact that the combination of menstrual symptoms, self-perceptions, and social norms might have on physical activity warrants further research attention. In this context, it would be of interest to explore women's familiarity with their own menstrual cycle and to unpack their typical involvement in physical activity in order to understand how differences among women in these areas might affect their management of physical activity throughout the menstrual cycle.

In conclusion, the present study suggests that menstrual events are complex, individual, and personal to the women experiencing them (e.g., Brantelid et al., 2014). The results demonstrated that women did not simply avoid all physical activity. Rather, they adapted physical activity depending on their distinct experiences of the menstrual cycle. This insight into the multi-faceted effects that the menstrual cycle could have on women's physical activity is pertinent to practitioners in the field. Creating safe spaces for women who wish to speak about how their physical comfort, personal thoughts, and perceptions of social norms might affect physical activity is important. Efforts in academic and practitioner fields could contribute to normalising conversations about the menstrual cycle (Johnston-Robledo & Chrisler, 2013; Kowalski & Chapple, 2000) and support physical activity throughout the menstrual cycle in women of all ages and abilities.

## References

Ansdell, P., Brownstein, C. G., Škarabot, J., Hicks, K. M., Simoes, D. C. M., Thomas, K., Howatson, G., Hunter, S. K. & Goodall, S. (2019). Menstrual cycle associated modulations in neuromuscular function and fatigability of the knee extensors in eumenorrheic females. *Journal of Applied Physiology*, 126, 1701-1712. <https://doi.org/10.1152/jappphysiol.01041.2018>



- Beitin, B. K. (2012). Interview and sampling: How many and whom. In J. F. Gubrium, J. A. Holstein, A. B. Marvasti, & K. D. McKinney (Eds.), *The SAGE handbook of interview research: The complexity of the craft* (2nd ed., pp. 243-254). SAGE Publications. <https://doi.org/http://dx.doi.org.mmu.idm.oclc.org/10.4135/9781452218403.n17>
- Brantelid, I. E., Nilvér, H. & Alehagen, S. (2014). Menstruation during a lifespan: A qualitative study of women's experiences. *Health Care for Women International*, 35(6), 600-616. <https://doi.org/10.1080/07399332.2013.868465>
- Braun, V., Clarke, V. & Weate, P. (2016). Using thematic analysis in sport and exercise research." In B. Smith & A. C. Sparkes (Eds.), *Routledge handbook of qualitative research in sport and exercise* (pp. 191-205). Routledge.
- Brinkmann, S., & Kvale, S. (2018). *Doing interviews* (2nd ed.). SAGE Publications. <https://dx.doi.org/10.4135/9781529716665>
- Brown, W. J., & Trost, S. G. (2003). Life transitions and changing physical activity patterns in young women. *American Journal of Preventive Medicine*, 25(2), 140-143. [https://doi.org/10.1016/S0749-3797\(03\)00119-3](https://doi.org/10.1016/S0749-3797(03)00119-3)
- Bruinvels, G., Burden, R., Brown, N., Richards, T., & Pedlar, C. (2016). The prevalence and impact of heavy menstrual bleeding (menorrhagia) in elite and non-elite athletes. *PLoS One*, 11(2), e0149881. <https://doi.org/10.1371/journal.pone.0149881>
- Bryman, A., Becker, S., & Sempik, J. (2008). Quality criteria for quantitative, qualitative and mixed methods research: A view from social policy. *International Journal of Social Research Methodology*, 11(4), 261-276. <https://doi.org/10.1080/13645570701401644>
- Buse, D., Manack, A., Serrano, D., Reed, M., Varon, S., Turkel, C., & Lipton, R. (2012). Headache impact of chronic and episodic migraine: Results from the American migraine prevalence and prevention study. *Headache: The Journal of Head and Face Pain*, 52(1), 3-17. <https://doi.org/10.1111/j.1526-4610.2011.02046.x>
- Chen, C. X., Kwekkeboom, K. L., & Ward, S. E. (2016). Beliefs about dysmenorrhea and their relationship to self-management. *Research in Nursing & Health*, 39(4), 263-276. <https://doi.org/10.1002/nur.21726>

- 793 Chen, H.-M., & Hu, H.-M. (2019). Randomized trial of modified stretching exercise  
794 program for menstrual low back pain. *Western Journal of Nursing Research*,  
795 41(2), 238-257. <https://doi.org/10.1177/0193945918763817>
- 796 Chrisler, J. C., Marván, M. L., Gorman, J. A., & Rossini, M. (2015). Body appreciation  
797 and attitudes toward menstruation. *Body Image*, 12, 78-81.  
798 <https://doi.org/10.1016/j.bodyim.2014.10.003>
- 799 Cioffi, J., Schmied, V., Dahlen, H., Mills, A., Thornton, C., Duff, M., Cummings, J., &  
800 Kolt, G. S. (2010). Physical activity in pregnancy: Women's perceptions,  
801 practices, and influencing Factors. *The Journal of Midwifery & Women's*  
802 *Health*, 55(5), 455-461. <https://doi.org/10.1016/j.jmwh.2009.12.003>
- 803 Collins, K. M. T. (2010). Advanced sampling designs in mixed research: Current  
804 practices and emerging trends in the social and behavioral sciences. In A.  
805 Tashakkori & C. Teddlie (Eds.), *SAGE handbook of mixed methods in social &*  
806 *behavioral research* (2nd ed., pp. 353-378). SAGE  
807 Publications. [https://doi.org/https://dx-doi-](https://doi.org/https://dx-doi-org.mmu.idm.oclc.org/10.4135/9781506335193.n15)  
808 [org.mmu.idm.oclc.org/10.4135/9781506335193.n15](https://doi.org/https://dx-doi-org.mmu.idm.oclc.org/10.4135/9781506335193.n15)
- 809 Creswell, J. W., Klassen, A. C., Plano Clark, V. L., & Clegg Smith, K. (2010). *Best*  
810 *practices for mixed methods research in the health sciences*. Office for  
811 behavioral and social sciences research (OBSSR).
- 812 Davidson, C. (2009). Transcription: Imperatives for qualitative research. *International*  
813 *Journal of Qualitative Methods*, 8(2), 35-52.  
814 <https://doi.org/10.1177/160940690900800206>
- 815 De Jonge, X., & Janse, A. (2003). Effects of the menstrual cycle on exercise  
816 performance. *Sports medicine*, 33(11), 833-851.  
817 <https://doi.org/10.2165/00007256-200333110-00004>
- 818 Dehnavi, Z. M., Farzaneh, J., & Kamali, Z. (2018). The effect of aerobic exercise on  
819 primary dysmenorrhea: A clinical trial study. *Journal of Education and Health*  
820 *Promotion*, 7(3), 1-8. [https://doi.org/10.4103/jehp.jehp\\_79\\_17](https://doi.org/10.4103/jehp.jehp_79_17)
- 821 Dempsey, L., Dowling, M., Larkin, P., & Murphy, K. (2016). Sensitive interviewing in  
822 qualitative research. *Research in Nursing & Health*, 39(6), 480-490.  
823 <https://doi.org/10.1002/nur.21743>
- 824 Field, A. (2013). *Discovering statistics using IBM SPSS statistics*. SAGE Publications.

- Findlay, R. J., Macrae, E. H. R., Whyte, I. Y., Easton, C., & Forrest, L. J. (2020). How the menstrual cycle and menstruation affect sporting performance: experiences and perceptions of elite female rugby players. *British Journal of Sports Medicine*, 0, 1-7. <https://doi.org/10.1136/bjsports-2019-101486>
- Fourquet, J., Gao, X., Zavala, D., Orengo, J. C., Abac, S., Ruiz, A., Laboy, J., & Flores, I. (2010). Patients' report on how endometriosis affects health, work, and daily life. *Fertility and Sterility*, 93(7), 2424-2428. <https://doi.org/10.1016/j.fertnstert.2009.09.017>
- Fraser, I. S., Critchley, H. O., Broder, M., & Munro, M. G. (2011). The FIGO recommendations on terminologies and definitions for normal and abnormal uterine bleeding. *Seminars in Reproductive Medicine*, 29(5), 383-390. <https://doi.org/10.1055/s-0031-1287662>
- Fraser, I. S., Mansour, D., Breymann, C., Hoffman, C., Mezzacasa, A., & Petraglia, F. (2015). Prevalence of heavy menstrual bleeding and experiences of affected women in a European patient survey. *International Journal of Gynecology & Obstetrics*, 128(3), 196-200. <https://doi.org/10.1016/j.ijgo.2014.09.027>
- Fredriksson, S. V., Alley, S. J., Rebar, A. L., Hayman, M., Vandelanotte, C., & Schoeppe, S. (2018). How are different levels of knowledge about physical activity associated with physical activity behaviour in Australian adults? *PloS One*, 13(11), e0207003. <https://doi.org/10.1371/journal.pone.0207003>
- Garcia-Perez, M. A., & Nunez-Anton, V. (2003). Cellwise residual analysis in two-way contingency tables. *Educational and Psychological Measurement*, 63(5), 825-839. <https://doi.org/10.1177/0013164403251280>
- Goffman, E. (1956). *The presentation of self in everyday life*. Penguin.
- Goffman, E. (1963). *Stigma: Notes on a spoiled identity*. Simon and Schuster, Inc.
- Hammersley, M., & Traianou, A. (2012). *Ethics in qualitative research: Controversies and contexts*. SAGE Publications. <https://doi.org/10.4135/9781473957619>
- Heinrich, K. M., Maddock, J., & Bauman, A. (2011). Exploring the relationship between physical activity knowledge, health outcomes expectancies, and behavior. *Journal of Physical Activity and Health*, 8(3), 404-409. <https://doi.org/10.1123/jpah.8.3.404>

- 856 Herrick, S. S., & Duncan, L. R. (2018). A qualitative exploration of LGBTQ+ and  
857 intersecting identities within physical activity contexts. *Journal of Sport and*  
858 *Exercise Psychology*, 40(6), 325-335. <https://doi.org/10.1123/jsep.2018-0090>
- 859 Houston, A. M., Abraham, A., Huang, Z., & D'Angelo, L. J. (2006). Knowledge,  
860 attitudes, and consequences of menstrual health in urban adolescent  
861 females. *Journal of Pediatric and Adolescent Gynecology*, 19(4), 271-275.  
862 <https://doi.org/10.1016/j.jpag.2006.05.002>
- 863 Ivankova, N. V. (2014). Implementing quality criteria in designing and conducting a  
864 sequential QUAN → QUAL mixed methods study of student engagement with  
865 learning applied research methods online. *Journal of Mixed Methods*  
866 *Research*, 8(1), 25-51. <https://doi.org/10.1177/1558689813487945>
- 867 Johnston-Robledo, I., & Chrisler, J. C. (2013). The menstrual mark: Menstruation as  
868 social stigma. *Sex Roles*, 68(1), 9-18. <https://doi.org/10.1007/s11199-011-0052-z>
- 869 Kannan, P., Chapple, C. M., Miller, D., Claydon, L. S., & Baxter, G. D. (2015).  
870 Menstrual pain and quality of life in women with primary dysmenorrhea:  
871 Rationale, design, and interventions of a randomized controlled trial of effects of  
872 a treadmill-based exercise intervention. *Contemporary Clinical Trials*, 42, 81-  
873 89. <https://doi.org/10.1016/j.cct.2015.03.010>
- 874 Kanojia, S., Sharma, V. K., Gandhi, A., Kapoor, R., Kukreja, A., & Subramanian, S. K.  
875 (2013). Effect of yoga on autonomic functions and psychological status during  
876 both phases of menstrual cycle in young healthy females. *Journal of Clinical*  
877 *and Diagnostic Research: JCDR*, 7(10), 2133-2139.  
878 <https://doi.org/10.7860/JCDR/2013/6912.3451>
- 879 Kennett, D. J., O'Hagan, F. T., & Meyerhoff, T. J. (2016). Managing menstruation:  
880 Moderating role of symptom severity on active coping and acceptance. *Western*  
881 *Journal of Nursing Research*, 38(5), 553-571.  
882 <https://doi.org/10.1177/0193945915620055>
- 883 Kowalski, R. M., & Chapple, T. (2000). The social stigma of menstruation: Fact or  
884 fiction? *Psychology of Women Quarterly*, 24(1), 74-80.  
885 <https://doi.org/10.1111/j.1471-6402.2000.tb01023.x>
- 886 Kural, M., Noor, N. N., Pandit, D., Joshi, T., & Patil, A. (2015). Menstrual  
887 characteristics and prevalence of dysmenorrhea in college going girls. *Journal of*

- 888 *Family Medicine and Primary Care*, 4(3), 426-431. [https://doi.org/](https://doi.org/10.4103/2249-4863.161345)  
 889 [10.4103/2249-4863.161345](https://doi.org/10.4103/2249-4863.161345)
- 890 Larroy, C. (2002). Comparing visual-analog and numeric scales for assessing menstrual  
 891 pain. *Behavioral Medicine*, 27(4), 179-181.  
 892 <https://doi.org/10.1080/08964280209596043>
- 893 Larun, L., & Malterud, K. (2011). Finding the right balance of physical activity: a focus  
 894 group study about experiences among patients with chronic fatigue  
 895 syndrome. *Patient Education and Counseling*, 83(2), 222-226.  
 896 <https://doi.org/10.1016/j.pec.2010.05.027>
- 897 Lee, S. (2002). Health and sickness: The meaning of menstruation and premenstrual  
 898 syndrome in women's lives. *Sex Roles*, 46(1), 25-35.  
 899 <https://doi.org/10.1023/A:1016033517659>
- 900 Lee, P. H., Macfarlane, D. J., Lam, T. H., & Stewart, S. M. (2011). Validity of the  
 901 international physical activity questionnaire short form (IPAQ-SF): A systematic  
 902 review. *International Journal of Behavioral Nutrition and Physical*  
 903 *Activity*, 8(1), 1-11. [https://doi.org/https://doi.org/10.1186/1479-5868-8-115](https://doi.org/10.1186/1479-5868-8-115)
- 904 Markland, D. (2009). The mediating role of behavioural regulations in the relationship  
 905 between perceived body size discrepancies and physical activity among adult  
 906 women. *Hellenic Journal of Psychology*, 6(2), 169-182.
- 907 Marshall, K., Chamberlain, K., & Hodgetts, D. (2019). Female bodybuilders on  
 908 Instagram: Negotiating an empowered femininity. *Feminism & Psychology*,  
 909 29(1), 96-119. <https://doi.org/10.1177/0959353518808319>
- 910 Marván, L., & Molina-Abolnik, M. (2012). Mexican adolescents' experience of  
 911 menarche and attitudes toward menstruation: Role of communication between  
 912 mothers and daughters. *Journal of Pediatric and Adolescent Gynecology*, 25(6),  
 913 358-363. <https://doi.org/10.1016/j.jpag.2012.05.003>
- 914 Marván, L., Ramírez-Esparza, D., Cortés-Iniestra, S., & Chrisler, J. C. (2006).  
 915 Development of a new scale to measure beliefs about and attitudes toward  
 916 menstruation (BATM): Data from Mexico and the United States. *Health Care*  
 917 *for Women International*, 27(5), 453-473.  
 918 <https://doi.org/10.1080/07399330600629658>
- 919 Matteson, K. A., Scott, D. M., Raker, C. A., & Clark, M. A. (2015). The menstrual  
 920 bleeding questionnaire: Development and validation of a comprehensive patient

- 921 - reported outcome instrument for heavy menstrual bleeding. *BJOG: An*  
 922 *International Journal of Obstetrics & Gynaecology*, 122(5), 681-  
 923 689. <https://doi.org/10.1111/1471-0528.13273>
- 924 McHugh, M. L. (2013). The chi-square test of independence. *Biochemia Medica:*  
 925 *Biochemia Medica*, 23(2), 143-149. <https://doi.org/10.11613/BM.2013.018>
- 926 Miller, T. (2017). Telling the difficult things: Creating spaces for disclosure, rapport  
 927 and 'collusion' in qualitative interviews. *Women's Studies International Forum*,  
 928 61, 81-86. <https://doi.org/10.1016/j.wsif.2016.07.005>
- 929 Milton, K., Bull, F.C., & Bauman, A., (2011). Reliability and validity testing of a  
 930 single-item physical activity measure. *British Journal of Sports Medicine*, 45(3),  
 931 203-208.
- 932 Milton, K., Clemes, S., & Bull, F. (2013). Can a single question provide an accurate  
 933 measure of physical activity? *British Journal of Sports Medicine*, 47(1), 44-48.
- 934 Mondragon, N. I., & Txertudi, M. B., 2019. Understanding menstruation: Influence of  
 935 gender and ideological factors. A study of young people's social  
 936 representations. *Feminism & Psychology*, 29(3), 357-373.  
 937 <https://doi.org/10.1177/0959353519836445>
- 938 Moreno-Black, G., & Vallianatos, H. (2005). Young women's experiences of  
 939 menstruation and athletics. *Women's Studies Quarterly*, 33(1/2), 50-67.
- 940 Morgan, D. L. (2007). Paradigms lost and pragmatism regained: Methodological  
 941 implications of combining qualitative and quantitative methods. *Journal of*  
 942 *Mixed Methods Research*, 1(1), 48-76. [http://dx.doi.org/10.1136/bjsports-2011-](http://dx.doi.org/10.1136/bjsports-2011-090899)  
 943 [090899](http://dx.doi.org/10.1136/bjsports-2011-090899)
- 944 Mulligan, H., Miyahara, M., & Nichols-Dunsmuir, A. (2017). Multiple perspectives on  
 945 accessibility to physical activity for people with long-term mobility  
 946 impairment. *Scandinavian Journal of Disability Research*, 19(4), 295-306.  
 947 <https://doi.org/10.1080/15017419.2016.1167772>
- 948 NHS Digital. (2017). Health survey for England 2016 – Physical activity in adults.  
 949 <https://files.digital.nhs.uk/publication/m/3/hse16-adult-phy-act.pdf>
- 950 Nowell, L. S., Norris, J. M., White, D. E., & Moules, N. J. (2017). Thematic analysis:  
 951 Striving to meet the trustworthiness criteria. *International Journal of Qualitative*  
 952 *Methods*, 16(1), 1-13. <https://doi.org/10.1177/1609406917733847>



- 953 O'Cathain, A. (2010). Mixed methods involving qualitative research. In I. Bourgeault,  
954 R. Dingwall, & R. De Vries (Eds.), *The SAGE handbook of qualitative methods*  
955 *in health research* (pp. 575-588). SAGE Publications.  
956 <https://dx.doi.org/10.4135/9781446268247>
- 957 Onwuegbuzie, A. J., & Collins, K. M. T. (2007). A typology of mixed methods  
958 sampling designs in social science research. *The Qualitative Report*, 12(2), 281-  
959 316.
- 960 Plotnikoff, R. C., Lippke, S., Johnson, S. T., Hugo, K., Rodgers, W., & Spence, J. C.  
961 (2011). Awareness of Canada's physical activity guide to healthy active living in  
962 a large community sample. *American Journal of Health Promotion*, 25(5), 294-  
963 297. <https://doi.org/10.4278/ajhp.090211-ARB-60>
- 964 Pridgeon, L., & Grogan, S. (2012). Understanding exercise adherence and dropout: An  
965 interpretative phenomenological analysis of men and women's accounts of gym  
966 attendance and non-attendance. *Qualitative Research in Sport, Exercise and*  
967 *Health*, 4(3), 382-399. <https://doi.org/10.1080/2159676X.2012.712984>
- 968 Samy, A., Zaki, S. S., Metwally, A. A., Mahmoud, D. S. E., Elzahaby, I. M., Amin, A.  
969 H., Eissa, A. I., Abbas, A. M., Hussein, A. H., & Talaat, B. (2019). The effect of  
970 zumba exercise on reducing menstrual pain in young women with primary  
971 dysmenorrhea: A randomized controlled trial. *Journal of Pediatric and*  
972 *Adolescent Gynecology*, 32(5), 541-545.  
973 <https://doi.org/10.1016/j.jpap.2019.06.001>
- 974 Sandelowski, M. (2000). Combining quantitative and qualitative sampling, data  
975 collection, and analysis techniques in mixed-method studies. *Research in*  
976 *Nursing and Health*, 23(3), 246-255.
- 977 Santer, M., Wyke, S., & Warner, P. (2007). What aspects of periods are most  
978 bothersome for women reporting heavy menstrual bleeding? Community survey  
979 and qualitative study. *BMC Women's Health*, 7(1), 8-13.  
980 <https://doi.org/10.1186/1472-6874-7-8>
- 981 Santer, M., Wyke, S., & Warner, P. (2008). Women's management of menstrual  
982 symptoms: Findings from a postal survey and qualitative interviews. *Social*  
983 *Science & Medicine*, 66(2), 276-288.  
984 <https://doi.org/10.1016/j.socscimed.2007.08.018>

- 985 Schechtman, K. B., Barzilai, B., Rost, K., & Fisher, E. B. (1991). Measuring physical  
986 activity within a single question. *American Journal of Public Health*, 81(6), 771-  
987 773. <https://doi.org/10.2105/AJPH.81.6.771>
- 988 Schoep, M. E., Nieboer, T. E., Van der Zanden, M., Braat, D. D. M., & Nap, A. W.  
989 (2019). The impact of menstrual symptoms on everyday life: A survey among  
990 42,879 women. *Gynecology*, 220(6), 561-567.  
991 <https://doi.org/10.1016/j.ajog.2019.02.048>
- 992 Scott, M., Flaherty, D., & Currall, J. (2013). Statistics: Dealing with categorical  
993 data. *Journal of Small Animal Practice*, 54(1), 3-  
994 8. <https://doi.org/10.1111/j.1748-5827.2012.01298.x>
- 995 Shilling, C., & Bunsell, T. (2009). The female bodybuilder as a gender  
996 outlaw. *Qualitative Research in Sport and Exercise*, 1(2), 141-159.  
997 <https://doi.org/10.1080/19398440902909009>
- 998 Small, C. M., Manatunga, A. K., & Marcus, M. (2007). Validity of self-reported  
999 menstrual cycle length. *Annals of Epidemiology*, 17(3), 163-  
1000 170. <https://doi.org/10.1016/j.annepidem.2006.05.005>
- 1001 Smith, B., & McGannon, K. R. (2018). Developing rigor in qualitative research:  
1002 Problems and opportunities within sport and exercise psychology. *International*  
1003 *Review of Sport and Exercise Psychology*, 11(1), 101-121.  
1004 <https://doi.org/10.1080/1750984X.2017.1317357>
- 1005 Smith, B., & Sparkes, A. (2016). Introduction - An invitation to qualitative research. In  
1006 B. Smith & A. Sparkes (Eds.), *The Routledge handbook of qualitative research*  
1007 *in sport, exercise and health* (pp. 1-8). Routledge.
- 1008 Spadaro, G., d'Elia, S. R. G., & Mosso, C. O. (2018). Menstrual knowledge and taboo  
1009 TV commercials: Effects on self-objectification among Italian and Swedish  
1010 women. *Sex Roles*, 78(9-10), 685-696. [https://doi.org/10.1007/s11199-017-](https://doi.org/10.1007/s11199-017-0825-0)  
1011 [0825-0](https://doi.org/10.1007/s11199-017-0825-0)
- 1012 Stephan, Y. (2003). Repercussions of transition out of elite sport on subjective well-  
1013 being: A one-year study. *Journal of Applied Sport Psychology*, 15(4), 354-  
1014 371. <https://doi.org/10.1080/714044202>
- 1015 Stubbs, M. L. (2008). Cultural perceptions and practices around menarche and  
1016 adolescent menstruation in the United States. *Annals of the New York Academy*  
1017 *of Sciences*, 1135(1), 58-66. <https://doi.org/10.1196/annals.1429.008>



- 1018 Tashakkori, A., & Teddlie, C. (2008). Quality of inferences in mixed methods research:  
1019 Calling for an integrative framework. In M. M. Bergman (Ed.), *Advances in*  
1020 *mixed methods research: Theories and application* (pp. 101-119). SAGE  
1021 Publications. <https://dx.doi.org/10.4135/9780857024329>
- 1022 Vartanian, L. R., & Novak, S. A. (2011). Internalized societal attitudes moderate the  
1023 impact of weight stigma on avoidance of exercise. *Obesity*, 19(4), 757-762.  
1024 <https://doi.org/10.1038/oby.2010.234>
- 1025 Warren, C. A. B. (2012). Interviewing as social interaction. In J. F. Gubrium, J. A.  
1026 Holstein, A. B. Marvasti, & K. D. McKinney (Eds.), *The SAGE handbook of*  
1027 *interview research: The complexity of the craft* (2nd ed., pp. 129-142). SAGE  
1028 Publications. <https://dx.doi.org/10.4135/9781452218403>
- 1029 Wegienka, G., & Baird, D. D. (2005). A comparison of recalled date of last menstrual  
1030 period with prospectively recorded dates. *Journal of Women's Health*, 14(3),  
1031 248-252. <https://doi.org/https://doi.org/10.1089/jwh.2005.14.248>
- 1032 Weiss, W. M. (2015). Competitive-level differences on sport commitment among high  
1033 school and collegiate-level athletes. *International Journal of Sport and Exercise*  
1034 *Psychology*, 13(3), 286-303. <https://doi.org/10.1080/1612197X.2014.958517>
- 1035 Weiss, W. M., Weiss, M. R., & Amorose, A. J. (2010). Sport commitment among  
1036 competitive female athletes: Test of an expanded model. *Journal of Sports*  
1037 *Sciences*, 28(4), 423-434. <https://doi.org/10.1080/02640410903536442>
- 1038 Zamani Sani, S. H., Fathirezaie, Z., Brand, S., Pühse, U., Holsboer-Trachsler, E.,  
1039 Gerber, M., & Talepasand, S. (2016). Physical activity and self-esteem: Testing  
1040 direct and indirect relationships associated with psychological and physical  
1041 mechanisms. *Neuropsychiatric Disease and Treatment*, 12, 2617-2625.  
1042 <https://doi.org/10.2147/NDT.S116811>
- 1043

1044 **Tables**

1045 Table 1. Mean (SD) participant characteristics and menstrual cycle symptoms taken  
 1046 from the questionnaire.

	All	Avoider	Non-avoider
N	128	44	84
Age (yrs)	27.9 (7.5)	28.2 (8.0)	27.2 (7.3)
Height (m)	1.65 (0.06)	1.65 (0.06)	1.65 (0.06)
Body mass (kg)	63.2 (12.2)	61.9 (12.8)	63.8 (11.8)
Length of period (days)	4.64 (1.80)	5.07 (1.11)*	4.42 (2.04)
Bleeding (days/MC)	3.81 (2.10)	4.43 (2.07)*	3.49 (2.05)
Spotting (days/MC)	1.51 (1.80)	2.02 (2.37)	1.24 (1.35)
Fatigue (days/MC)	2.82 (2.99)	4.43 (3.45)†	3.45 (2.34)
Pain (days/MC)	2.60 (2.87)	4.20 (3.60)†	1.76 (1.95)
Pain severity (numerical scale)	3.95 (2.80)	5.27 (2.41)†	3.25 (2.75)


1047 MC, menstrual cycle. \*denotes significant difference from non-avoider ( $p<0.05$ ),

1048 †denotes significant difference from non-avoider ( $p<0.01$ ).

1049

1050 **Figures**

1051 Figure 1. Overview of study design (adapted from Ivankova, 2014)



1052