

Potential Health Benefits of Anthocyanins in Oxidative Stress Related Disorders

Md. Ashraful Alam¹, Priota Islam¹, Nusrat Subhan^{1*}, Md. Mizanur Rahman¹, Ferdous Khan¹, Geoffrey E. Burrows²,
Lutfun Nahar^{3,4} and Satyajit Dey Sarker³

¹Department of Pharmaceutical Sciences, North South University, Dhaka, Bangladesh

²School of Agricultural and Wine Sciences, Charles Sturt University, Wagga Wagga, Locked Bag 588, NSW 2678, Australia.

³Centre for Natural Products Discovery (CNPD), School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3 3AF, United Kingdom

^{3,4}Laboratory of Growth Regulators, Institute of Experimental Botany ASCR & Palacký University, Šlechtitelů 27, 78371 Olomouc, Czech Republic

Corresponding Author

*Dr Nusrat Subhan, Department of Pharmaceutical Sciences, North South University, Dhaka, Bangladesh, email-
rimmi04@yahoo.com

Abstract

Anthocyanins are naturally occurring water-soluble plant pigments belonging to the flavonoids chemical class. The red, blue and purple colours of leaves, flowers and fruits of plants confirm that they are rich sources of anthocyanins. Many *in vivo* and *in vitro* studies reveal that anthocyanins have different health beneficial effects such as antioxidant, antidiabetic, anti-inflammatory, anti-obesity, antihypertensive and anticancer properties. Major benefits of anthocyanin administration are owing to their potent anti-inflammatory and antioxidant activities. Recent investigations have revealed that anti-inflammatory activities of anthocyanins follow the inhibitory pathways of NF- κ B-mediated decline of inflammatory cytokines production. Inhibition of the anti-inflammatory pathways also influences the modulation of arteriolar disorders and cardiovascular complications due to anthocyanin administration. Moreover, anthocyanins improve diabetes, obesity and cancer pathology by inhibiting NF- κ B-mediated inflammatory pathways. However, considerable variations in activities do exist among structurally diverse anthocyanins. This review appraises the recent literature regarding the health benefits of anthocyanins and their molecular mechanisms in various oxidative stress related pathophysiological conditions.

Key words: Anthocyanins, cancer, diabetes, inflammation, obesity and oxidative stress.

Introduction

The healthy human body always maintains a good balance between free radical production and antioxidant systems. Free-radicals are species that exist independently, having one or more unpaired free electrons and may react with other molecules in cells by taking or supplying electrons. The imbalanced production of reactive oxygen species (ROS) or reactive nitrogen species (RNS) and their limited degradation through compromised enzymatic defence may turn into oxidative stress (García-Sánchez et al., 2020), playing an important role in the initiation, development and progression of several non-communicable diseases such as obesity, diabetes (Pang et al., 2020; Yaribeygi et al., 2020), cardiovascular disease, neurodegenerative diseases (Bhatt et al., 2020), as well as the modulation of gut bacterial environment (Hu et al., 2020), and bone metabolism (Bernatoniene and Kopustinskiene, 2018). Researchers also suggest that oxidative stress and free-radical generated reactions are prime contributors to degenerative processes like aging (Speer et al., 2020; Lobo et al. 2010; Kattoor et al. 2017). Major ROS/RNS species include superoxide anion radical ($O_2^{\bullet-}$), hydroxyl radical (HO^{\bullet}), nitric oxide radical (NO^{\bullet}) and other molecular species such as hydrogen peroxide (H_2O_2), hypochlorous acid ($HOCl$), peroxynitrite ($ONOO^-$) etc. are generated endogenously in cells and tissues (Valko et al., 2007). Several other oxygen species such as singlet oxygen (1O_2), lipid radicals (R^{\bullet} , RO^{\bullet} , ROO^{\bullet}), and hydroperoxyl radical (HOO^{\bullet}) are also contributing to the oxidative stress in biological system which may produce on later part and are linked to ROS/RNS mediated reactions (Phaniendra et al., 2015). The species $O_2^{\bullet-}$, H_2O_2 , and HO^{\bullet} are generated from molecular oxygen by one-, two-, and three-electron reduction of molecular oxygen primarily from the mitochondrial electron transport chain (Phaniendra et al., 2015; Pospíšil et al., 2019). Other sources of ROS/RNS species are the nicotinamide adenine dinucleotide phosphate oxidase (NADPH oxidase) in phagocytes, myeloperoxidase in neutrophil, xanthin oxidase system, nitric oxide synthase system and cytochrome p450 phase II metabolizing enzyme system etc. (Di Meo et al., 2016).

Endogenous enzymatic antioxidants such as superoxide dismutase (SOD) (removes $O_2^{\bullet-}$), catalase (decomposes H_2O_2) and glutathione peroxidase (GPX) may serve as the primary defence against oxidative stress and related disorders (Ighodaro and Akinloye, 2018; Kurutas, 2016). Several other non-enzymatic antioxidants such as glutathione, lipoic acid, albumin, uric acid, and bilirubin, are also some contributing agents against the free radical mediated oxidative stress (Kuciel-Lewandowska et al., 2020). Exogenous antioxidants are also available from dietary sources such as ascorbic acid, retinol, α -tocopherol, carotenoids, polyphenolic compounds (flavonoids and phenolic acids), and trace elements such as zinc, manganese, selenium, and chromium (Bouayed and Bohn, 2010; Singh et al., 2010). Deficiency of these antioxidants in diet may also lead to degenerative diseases in human (Bouayed and Bohn, 2010).

Anthocyanins, natural dietary antioxidants that provide protective effects against harmful effect of oxidative stress (Ullah et al. 2019), have become one of the key topics for research linking to diabetes, obesity, inflammation, cancer and degenerative neurological disorders (Fallah et al., 2020; Pandey and Rizvi 2009). These polyphenolic compounds are capable of trapping free-radicals in the human body. Plants are rich sources of natural antioxidants such as phenolic acids, flavonoids and anthocyanins. Anthocyanins are an important group of pigments that are water-soluble and belong to the flavonoids family. They give the distinctive red, purple and blue pigments in most fruits and vegetables

(Fallah et al., 2020; Passeri et al. 2016). Anthocyanins display numerous potential health benefits and they have been investigated recently for their use as possible clinical treatments for many human disorders (Bakuradze et al. 2019). They have both anti-inflammatory and antioxidant properties, which have proven effectiveness in *in vivo* and *in vitro* models of various chronic disease conditions such as cardiovascular disease, ophthalmic disorders, obesity, type II diabetes and atherosclerosis (Toufektsian et al. 2008; Seymour et al. 2009; Basu et al. 2010; Kalt et al. 2010; Mauray et al. 2012; Blesso, 2019).

This review appraises recent studies on the health benefits of anthocyanins that include their roles in cardiovascular diseases, neurodegenerative diseases, visual acuity, cancer, diabetes and several other health-related issues.

Methodology:

In order to assess the current data on anthocyanin, a comprehensive search of the scientific literature was conducted using Google Scholar, PubMed, Web of Knowledge and Science Directory with the key words or phrases ‘anthocyanin’, ‘anthocyanin in diabetes’, ‘anthocyanin in inflammation’, ‘anthocyanin in cardiovascular diseases’, ‘anthocyanin in cancer’ and ‘anthocyanin in neurological disorder’. Furthermore, the reference lists of the selected literatures have also carefully analysed to clarify information. Our exploration indicated that the most of the research done in the last decade was focused on the effect of dietary anthocyanins in minimizing health risks linked to various diseases. This information was then tabulated and discussed critically to evaluate the mechanism of anthocyanins in various cellular processes. A further prospective research goal related to anthocyanin molecules was also warranted.

Chemistry of Anthocyanins

Anthocyanin structurally belongs to the flavonoid class of natural products (Nahar and Sarker, 2019). Free anthocyanins are rare in fruits and vegetables and generally, they form glycosides, with several carbohydrates such as arabinose, galactose, glucose, rhamnose or xylose, which are attached to an aglycone anthocyanin nucleus (Harborne and Grayer 1988; Mazza and Francis 1995; Khoo *et al.* 2017; Salehi et al., 2020). Anthocyanins have a unique ability to form flavylium cations that distinguish them from other flavonoids (Ullah *et al.* 2019). Unlike other flavonoids, the positive charge makes anthocyanins more stable in acidic environments, such as in the stomach (Mazza and Francis 1995). The anthocyanidins are the aglycone or de-glycosylated forms of anthocyanins, including cyanidin, delphinidin, malvidin, pelargonidin, petunidin and peonidin (**Figure-1, Table 1**). The glycosides of the three non-methylated anthocyanidins (cyanidin, delphinidin and pelargonidin) (**Figure-1, Table 1**) are widespread in nature, being 80, 69 and 50% abundant in pigmented fruits and flowers, respectively. The most plentiful anthocyanins in edible parts of plants are cyanidin, followed by pelargonidin, peonidin, delphinidin, petunidin and malvidin (Kang *et al.* 2003; Su et al., 2019). The sugar parts of anthocyanins are usually attached to the anthocyanidin skeleton through the C₃ hydroxyl group. Numerous anthocyanins vary in the basic anthocyanidin skeleton, and the extent and position by which the glycosides attach to the skeleton (**Table 2**) (Harborne and Grayer 1988). Structural diversities among

anthocyanins render significant differences in their physicochemical characters implicating their various pathways of absorption (Prior and Wu 2006; Bonesi et al., 2020).

The intensity of colour of anthocyanins depend on pH and the presence of chelating metal ions, mainly occurring as blue, purple or red colour (Ibrahim *et al.* 2011). Flavylium cation is red at pH 1-3, at pH 5 the resultant carbinol *pseudo* base is colourless, while at pH 7-8 (**Figure 2**), the blue to purple quinoidal is formed (Harborne and Williams 2001). The cationic and polyphenolic nature of anthocyanins and their metabolites initiate various cellular responses among which the polyphenolic nature is mainly responsible for their strong free-radical scavenging and antioxidant activities, and thus produce their pharmacological effect.




Figure 1: Basic structure of anthocyanidin. It has two benzoyl rings (A and B) which are connected by a heterocyclic (C) ring. Glycosides are attached through the R₁ hydroxyl group.

Table 1. Various groups present in anthocyanidin ring structure.

	R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	Colour
Common Anthocyanidins*								
Cyanidin	-OH	-OH	-H	-OH	-OH	-OH	-H	Orange-red
Delphinidin	-OH	-OH	-H	-OH	-OH	-OH	-OH	Bluish-red
Pelargonidin	-OH	-OH	-H	-OH	-H	-OH	-H	Orange
Peonidin	-OH	-OH	-H	-OH	-OCH ₃	-OH	-H	Orange-red
Petunidin	-OH	-OH	-H	-OH	-OH	-OH	-OCH ₃	Bluish-red
Malvidin	-OH	-OH	-H	-OH	-OCH ₃	-OH	-OCH ₃	Bluish-red
6-Hydroxylated Anthocyanidins								
6-Hydroxycyanidin	-OH	-OH	-OH	-OH	-OH	-OH	-H	Red
6-Hydroxydelphinidin	-OH	-OH	-OH	-OH	-OH	-OH	-OH	Bluish-red
6-Hydroxypelargonidin (aurantinidin)	-OH	-OH	-OH	-OH	-H	-OH	-H	Orange
Methylated Anthocyanidins								
5-Methylcyanidin	-OH	-OCH ₃	-H	-OH	-OH	-OH	-H	Orange-red
5-Methyldelphinidin (pulchellidin)	-OH	-OCH ₃	-H	-OH	-OH	-OH	-OH	Bluish-red
7-Methylmalvidin (hirsutidin)	-OH	-OH	-H	-OCH ₃	-OCH ₃	-OH	-OCH ₃	Bluish-red
5-Methylmalvidin (capensinidin)	-OH	-OCH ₃	-H	-OH	-OCH ₃	-OH	-OCH ₃	Bluish-red
7-Methylpeonidin (rosinidin)	-OH	-OH	-H	-OCH ₃	-OCH ₃	-OH	-H	Red
5-Methylpetunidin (europinidin)	-OH	-OCH ₃	-H	-OH	-OCH ₃	-OH	-OH	Bluish-red
Others								
Apigeninidin	-H	-OH	-H	-OH	-H	-OH	-H	Orange
Luteolinidin	-H	-OH	-H	-OH	-OH	-OH	-H	Orange
Tricetinidin	-H	-OH	-H	-OH	-OH	-OH	-OH	Red

*Common anthocyanidins present in naturally occurring important foods

Table 2. Anthocyanins and their derivatives in natural food sources.

Source	Anthocyanins and their derivatives		Acyl	Origin	References
	Aglycones	Glycoside types			
Acai berry (<i>Euterpe oleracea</i>)	Cyn, Mav, Plg, Pen Cyn, Mav, Plg, Pen	3- <i>O</i> -glucoside.		Fruit pulp	(Poulose <i>et al.</i> 2012)
Bilberry (<i>Vaccinium myrtillus</i>)	Cyn	3- <i>O</i> -arabinoside, 3- <i>O</i> -galactoside, 3- <i>O</i> -glucoside		Fruit	(Madhavi <i>et al.</i> 1998; Cooke <i>et al.</i> 2006)
Blackberry (<i>Rubus fruticosus</i>)	Cyn Cyn, Plg, Pen	3- <i>O</i> -arabinoside, 3- <i>O</i> -rutinoside 3- <i>O</i> -glucoside.		Fruit	(Skrovankova <i>et al.</i> 2015)
Black carrot (<i>Daucus carota</i>)	Cyn, Pen Cyn, Pen, Plg. Cyn, Pen.	3- <i>O</i> -xylosylgalactoside, 3- <i>O</i> -xylosyl (feruloylglucosyl) galactoside 3- <i>O</i> -xylosyl (sinapoylglucosyl) galactoside		Root	(Montilla <i>et al.</i> 2011)
Black currant (<i>Ribes nigrum</i>)	Cyn, Del	3- <i>O</i> -rutinoside		Fruit	(Abo El-Ella and Bishayee 2019)
Black raspberry (<i>Rubus occidentalis</i>)	Cyn Cyn, Plg	3- <i>O</i> -sambubioside, 3- <i>O</i> -glucoside, 3- <i>O</i> -(xylosyl) rutinoside, 3- <i>O</i> -rutinoside		Fruit	(Overall <i>et al.</i> 2017)
Blueberry (<i>Vaccinium corymbosum</i>)	Cyn, Del, Mal. Pen, Pet.	3- <i>O</i> -arabinoside, 3- <i>O</i> -galactoside, 3- <i>O</i> -glucoside, 3- <i>O</i> -arabinoside, 3- <i>O</i> -galactoside	3- <i>O</i> -acetylglucoside	Fruit	(Nile and Park 2014; Skrovankova <i>et al.</i> 2015)
Cabbage (<i>Brassica oleracea</i>)	Del, Plg, Pen, Pet Cyn	3- <i>O</i> -diglucoside-5- <i>O</i> -glucoside		Flower	(Ahmadiani <i>et al.</i> 2019; Burton-Freeman <i>et al.</i> 2019)

Source	Anthocyanins and their derivatives		Acyl	Origin	References
	Aglycones	Glycoside types			
Cherry sweet (<i>Prunus avium</i>) and sour (<i>Prunus cerasus</i>)	Cyn, Del, Plg, Pen, Pet. Cyn, Plg, pen. Cyn	3- <i>O</i> -glucoside, 3- <i>O</i> -rutinoside, 3- <i>O</i> -glucosylrutinoside, 3- <i>O</i> -sophoroside		Fruit	(Mulabagal <i>et al.</i> 2009; Burton-Freeman <i>et al.</i> 2019)
Chokeberry (<i>Aronia melanocarpa</i>)	Cyn, Plg Cyn	3- <i>O</i> -arabinoside, 3- <i>O</i> -glucoside, 3- <i>O</i> -galactoside, 3- <i>O</i> -xyloside		Fruit	(Kulling and Rawel 2008)
Concord grape (<i>Vitis labrusca</i>)	Cyn, Del, Pet, Pen Cyn, Del	3- <i>O</i> -glucoside, 3- <i>O</i> -(coumaroyl) glucoside, 3,5- <i>O</i> -(coumaroyl) diglucoside,	3- <i>O</i> -acetyl and coumaroyl	Fruit	(Overall <i>et al.</i> 2017)
Cranberry (<i>Vaccinium macrocarpon</i>)	Cyn, Pen, Plg, Mal, Del Cyn, Pen, Pet	3- <i>O</i> -arabinoside, 3- <i>O</i> -galactoside, 3- <i>O</i> -glucoside 3- <i>O</i> -galactoside		Fruit	(Skrovankova <i>et al.</i> 2015; Abo El-Ella and Bishayee 2019;)
Eggplant (<i>Solanum melongena</i>)	Del	3- <i>O</i> -glucoside, 3- <i>O</i> -rutinoside, 3- <i>O</i> -rutinoside 5- <i>O</i> - glucoside		Fruit	(Sadilova <i>et al.</i> 2006)
Elderberry (<i>Sambucus nigra</i>)	Cyn	3- <i>O</i> -glucoside, 3- <i>O</i> -sambubioside, 3- <i>O</i> -sambubioside-5- <i>O</i> -glucoside		Fruit	(da Silva <i>et al.</i> 2019)
Grapes (<i>Vitis vinifera</i>)	Cyn, Del, Mal, Pen. Pet.	3- <i>O</i> -glucoside, 3,5- <i>O</i> -diglucoside		Fruit	(Mazza and Francis 1995; Grimes <i>et al.</i> 2018)
Jamun berry (<i>Syzygium cumini</i>)	Cyn, Del, Mal, Pen, Pet.	-		Fruit	(Aqil <i>et al.</i> 2012)

Source	Anthocyanins and their derivatives		Acyl	Origin	References
	Aglycones	Glycoside types			
Lingonberry (<i>Vaccinium vitis-idaea</i>)	Cyn	3- <i>O</i> -arabinoside, 3- <i>O</i> -galactoside, 3- <i>O</i> -glucoside		Fruit	(Isaak <i>et al.</i> 2017)
Pigmented rice (black and brown) (<i>Oryza sativa</i>)	Cyn, Pen	3- <i>O</i> -glucoside		Seed	(Yawadio <i>et al.</i> 2007)
Pigmented wheat (blue and purple) (<i>Triticum aestivum</i> , <i>Triticum durum</i>)	Cyn, Del, Pen, Mal	3- <i>O</i> -glucoside		Seed	(Zhu 2018)
Pomegranate (<i>Punica granatum</i>)	Cyn, Del, Pen Cyn, Del	3- <i>O</i> -glucoside 3,5- <i>O</i> -diglucoside		Fruit	(Sharma <i>et al.</i> 2017)
Purple cauliflower (<i>Brassica oleracea</i> var. botrytis)	Cyn	3- <i>O</i> -sophoroside-5- <i>O</i> -glucoside, 3- <i>O</i> -sophoroside-5-(malonyl) glucoside, glucoside-5-malonyl) glucoside, 3- <i>O</i> -(coumaryl)sophoroside-5- <i>O</i> -glucoside	3- <i>O</i> -(coumaryl-caffeyl)	Flower	(Chiu <i>et al.</i> 2010)
Purple corn (<i>Zea mays</i>)	Cyn, Plg, Pen	3- <i>O</i> -glucoside, 3-6- <i>O</i> -malonyl-glucoside		Seed	(Aoki <i>et al.</i> 2002)
Purple-fleshed sweet potato (<i>Ipomoea batatas</i>)	Cyn, Pen, Cyn	3- <i>O</i> -sophoroside-5- <i>O</i> -glucoside, 3- <i>O</i> -(6,6-dicaffeoyl sophoroside)-5- <i>O</i> -glucoside.		Root	(Su <i>et al.</i> 2019)

Source	Anthocyanins and their derivatives		Acyl	Origin	References
	Aglycones	Glycoside types			
		<i>p</i> -hydroxybenzoylated-3-sophoroside-5-glucoside), Caffeoylated-3- <i>O</i> -sophoroside-5- <i>O</i> -glucoside			
Radish (<i>Raphanus sativus</i>)	Cyn, Cyn, Pet	3- <i>O</i> -glucoside, 3- <i>O</i> -rutinoside-5-glucoside, 3,5- <i>O</i> -diglucoside		Root	(Zhang <i>et al.</i> 2019b)
Raspberry (<i>Rubus idaeus</i>)	Cyn	3- <i>O</i> -arabinose, 3- <i>O</i> -rutinoside, 3- <i>O</i> -sophoroside		Fruit	(Probst 2015; Skrovankova <i>et al.</i> 2015)
Red onion (<i>Allium cepa</i>)	Cyn,	3- <i>O</i> -glucoside, 3- <i>O</i> -laminaribioside, 3- <i>O</i> -(6-malonyl-glucoside), 3- <i>O</i> -(6-malonyl- laminaribioside)		Root	(Fossen and Andersen 2003; Frond <i>et al.</i> 2019)
Sweet potato (<i>Ipomoea batatas</i>)	Pen Cyn, Pen, Plg Cyn, Pen	3- <i>O</i> -glucoside. 3- <i>O</i> -sophoroside-5- <i>O</i> -glucoside. 3- <i>p</i> -hydroxybenzoylsophoroside-5- <i>O</i> -glucoside		Root	(Lim <i>et al.</i> 2013)
Strawberry (<i>Fragaria</i> × <i>ananassa</i>)	Plg Cyn, Plg, Pen	3- <i>O</i> -arabinoside, 3- <i>O</i> -malonylglucoside, malyglucoside, dissacharide (hexose + pentose) acylated with acetic acid. 3- <i>O</i> -glucoside, 3- <i>O</i> -galactoside, 3- <i>O</i> -rutinoside 3- <i>O</i> -glucoside		Fruit	(Giampieri <i>et al.</i> 2012; Skrovankova <i>et al.</i> 2015)

Source	Anthocyanins and their derivatives		Acyl	Origin	References
	Aglycones	Glycoside types			
Violet pepper or capsicum (<i>Capsicum annuum</i>)	Del	3- <i>O</i> -glucoside, 5- <i>O</i> -glucoside, 3- <i>O</i> -rhamnoside, 3- <i>O</i> -rutinoside, 3- <i>O</i> -caffeoyl rutinoside	3- <i>O</i> -coumaroyl-hexoside	Fruit	(Sadilova <i>et al.</i> 2006)

Cyn=Cyanidin, Plg= Pelargonidin, Mal= Malvidin, Pen=Peonidin, Pet=Petunidin, Del=Delphinidin, and ‘_’= No glycoside




Figure 2. Different forms of anthocyanins vs. pH values. Adopted and modified from previously reported articles (Borkowski *et al.* 2005; Marco *et al.* 2011).

Pharmacokinetics and Bioavailability

Recent studies on the bioavailability of anthocyanins suggest that they are rapidly absorbed from the small intestine and the stomach (McGhie and Walton 2019). Anthocyanins are quickly metabolized resulting in low concentrations of the parent compounds being present in plasma within a few hours of ingestion (Woodward *et al.* 2009; Azzini *et al.* 2010). In general, anthocyanins are absorbed intact as glycosides with their absorption and elimination being relatively quick. The absorption and subsequent metabolism of black raspberry (*R. occidentalis*) anthocyanins were investigated in humans, where high doses of anthocyanins were taken (2.69 g/day) (Stoner *et al.* 2005). Four anthocyanins from the raspberries were observed in plasma within 2 h of oral berry administration and their elimination was found to follow first-order kinetics. They were excreted, both as intact anthocyanins and as methylated derivatives. Similar results were recorded in bilberry (*V. myrtillus*) studies conducted in rodents (He *et al.* 2006). Maximum concentrations of anthocyanins were found in plasma within 1 to 2 h, and maximum quantities in urine appeared in 4 h. Overall, less than 1% of these compounds were absorbed and excreted in urine.

The biological activities of anthocyanins are closely correlated to their absorption and metabolism (Tian *et al.* 2019). Acylation and glycosylation patterns decrease the anthocyanin bioavailability; however, glycosidases present in the gastrointestinal tract may hydrolyze anthocyanins into anthocyanidins, thereby enhances their biological potential but

reduces their stability. The presence of a glucose substituent compared to an arabinose or galactose on the cyanidin and peonidin anthocyanidins present in cranberry (*V. macrocarpon*) juice appeared to increase their bioavailability as a percentage of the administered dose (Milbury *et al.* 2010).

Anthocyanins exist in the blood circulation and in urine in various forms such as intact, methylated, glucuronide derivatives and/or sulphate conjugated forms (Mazza *et al.* 2002; Felgines *et al.* 2003; Kay *et al.* 2005; Kay *et al.* 2007), reaching peak plasma concentrations 1 to 3 h after consumption. This depends on the individual compound along with the food matrix. The metabolites may remain in urine for up to 24 h and can retain their basic anthocyanin structure (Kay *et al.* 2005; Kay *et al.* 2007). Pharmacokinetic evidence indicated that parent glycosides and glucuronide derivatives were conspicuous in the bloodstream for 0 to 5 h after ingestion but became increasingly methylated after 6 to 24 h, which suggested that the bioactivity of anthocyanins was likely altered from metabolic transformation (Kay and Mazza, 2008). Various *in vivo* studies have suggested that the food matrix had a significant effect on the absorption and metabolism of anthocyanins. The bioavailability of an individual anthocyanin may vary in xenobiotic metabolism in the liver, GIT and other tissues. Human polymorphisms have been reported in the genes for catechol-*O*-methyltransferase, glucuronosyl transferase and glutathione *S*-transferases (Lampe and Chang 2007). The variation of human gut microflora may also have an important role in anthocyanin bioavailability (Zhu *et al.* 2018; Tian *et al.* 2019). Anthocyanins may be metabolized by microbiota occurring in the GIT and produce smaller, and more bioavailable end-products (Zhu *et al.* 2018).

Gut microflora are able to metabolize anthocyanins Chen *et al.* 2018; Zhu *et al.* 2018; (Li *et al.* 2019; Tian *et al.* 2019). By the use of a bacterial preparation imitating the normal human microbiota population, Williamson *et al.* (2009) were able to demonstrate the conversion of larger polyphenols to phenolic acids, which had similar anti-inflammatory effects as the parent compounds. In addition, smaller phenolic acids and other anthocyanin metabolites possessed greater chemical and microbial stability, suggesting that they may have an important role in the antioxidant activity and physiological effects observed in many studies (Keppler and Humpf 2005).

Health Benefits of Anthocyanins

The chemical nature of anthocyanins reveals that they have beneficial roles in the control of many diseases. Their beneficial roles have been found significant in various pathological conditions such as heart disease, neurodegenerative diseases, in improving visual acuity, cancer, diabetes and obesity etc. Some of the health benefits of anthocyanins are discussed below.

Antioxidant properties of anthocyanins

The beneficial effect of anthocyanins primarily relies on their antioxidants and free-radical scavenging properties. Antioxidant activity of anthocyanins were tested in various assay systems such as ferric reducing antioxidant potential (FRAP), trolox equivalent antioxidant capacity (TEAC), oxygen radical absorbance capacity (ORAC), peroxynitrite ('ONOO⁻') scavenging activity, lipid peroxidation inhibition, ability to bind heavy metals such as copper, iron and zinc

and the free-radical scavenging activity of 2,2-diphenyl-1-picrylhydrazyl (DPPH) (Kahkonen and Heinonen 2003; Yang and Zhai 2010; Ge and Ma 2013). However, antioxidant activities of anthocyanin derivatives are greatly dependent on anthocyanin chemical structure. Thus, various anthocyanin derivatives possess different capacities for scavenging diverse radioactive oxygen species (ROS) and radioactive nitrogen species (RNS). Moreover, induction of antioxidant enzymes such as glutathione-*S*-transferase (GST), glutathione reductase (GR), glutathione peroxidases (GPx) and superoxide dismutases were also observed due to anthocyanins (Turner 2009, Huang *et al.* 2016b).

Cyanidin 3-*O*-glycosides (arabinoside, galactoside, glucoside and rutinoside) and delphinidin 3-*O*-rutinoside were purified from various berries and *in vitro* investigation was performed to evaluate their effect on lipid peroxidation induced either by UV irradiation, Fe(II) ions or scavenging of 2,2'-azobis (2-amidinopropane) dihydrochloride peroxy radicals at the concentrations of 15-20 μ M (Gabrielska and Oszmiański 2005). Delphinidin-3-rutinoside produced a higher antioxidant activity against Fe(II)-induced liposome oxidation than cyanidin-3-rutinoside (Gabrielska and Oszmiański 2005). However, in terms of Fe(II)-induced liposome oxidation the antioxidant activity of the anthocyanins was higher than that of trolox (Gabrielska and Oszmiański 2005). Huang *et al.* (2016b) investigated malvidin and its two glycosides on cell lines to evaluate the effects on the ROS, heme oxygenase-1 (HO-1), superoxide dismutase (SOD) and xanthine oxidase-1 (XO-1). Malvidin glycosides displayed a greater inhibitory effect than malvidin in inhibiting xanthine oxidase activity; however, malvidin glycosides showed synergistic effects in HO-1 production in the cells (Huang *et al.* 2016b). Gabrielska and Oszmiański (2005) and Huang *et al.* (2016b) found the antioxidant activity in anthocyanin glycosides though they performed different antioxidant methods and at different concentrations. The peonidin-based anthocyanin components in purple sweet potato (*Ipomoea batatas*) were investigated in 2,2-diphenyl-1-picrylhydrazyl (DPPH) radicals and superoxide anions scavenging assay system (Sun *et al.* 2018). Significant increase of DPPH radicals and superoxide anions scavenging activity were observed with an increase in anthocyanin concentration (Sun *et al.* 2018). Moreover, these anthocyanin components also showed good potential in reducing Fe²⁺ chelating ability and total power activity (Sun *et al.* 2018).

Anti-Inflammatory Properties of Anthocyanins

Anthocyanins also possess anti-inflammatory properties. Inflammation is an integral part of tissue regeneration and wound healing due to infection. However, several pathways are considered responsible for inflammation such as nuclear factor-kappa B (NF- κ B)-mediated cytokines production, cyclooxygenase mediated pathway, mitogen activated protein kinase activity, inducible nitric oxide synthase (iNOS) expressed signalling, LPS-induced macrophage activation, endothelial adhesion molecules expression (Chen *et al.* 2017; Liu *et al.* 2017). Various reports suggest that anthocyanins have shown promising results in inhibiting most of the inflammatory signalling cascade and limiting inflammation mediated tissue damage (Pereira *et al.* 2017; Huang *et al.* 2018; Valenza *et al.* 2018). Anthocyanin, or anthocyanin-extract, inhibited NF- κ B through down-regulation of mitogen activated protein kinase (MAPK) pathways and reduced the expression of some pro-inflammatory cytokines *in vitro* (Pergola *et al.* 2006; Vendrame and Klimis-Zacas 2015). Malvidin and its glycosides inhibited high glucose-induced expression of intercellular adhesion molecule-1 (ICAM-1) and NF- κ B in human retinal capillary endothelial cells and significantly

increased cell viability (Huang *et al.* 2018). The anthocyanin-rich fraction of Portuguese blueberries (*V. corymbosum*) showed down regulation of cyclooxygenase-2 (COX-2) and iNOS in colon tissue homogenates (Pereira *et al.* 2017). The strong inhibition of COX-2 expression in colon tissue appears to be a key anti-inflammatory mechanism (Pereira *et al.* 2017).

Several anthocyanins such as delphinidin-3-*O*-glucoside, cyaniding-3-*O*-glucoside and petunidin-3-*O*-glucoside showed decreased NF- κ B activities via mitogen activated MAPK pathways (Afaq *et al.* 2005b; Jeong *et al.* 2013). Mulberry (*Morus australis*) anthocyanins such as cyanidin and pelargonidin showed decreased tumour necrosis factor alpha (TNF- α) and Interleukin 6 (IL-6) levels in animals fed a high fat diet (Wu *et al.* 2013). Joo *et al.* (2018) found that cyanidin collected from red Chinese cabbage (*Brassica rapa*) also inhibited TNF- α -induced NF- κ B. Black soybean (*Glycine max*) extract, rich in delphinine and petunidin, decreased the TNF- α and IL-6 level in high fat diet fed animals (Kim *et al.* 2015).

Anthocyanins in Cardiovascular Diseases

The development of cardiac and vascular dysfunction are attributed to the generation of free radicals from various sources such as mitochondrial electron transport chain, angiotensin II (ANG-II) mediated NADPH oxidase system, xanthin oxidase and inducible nitric oxide synthase (Münzel *et al.*, 2017). Free radicals and reactive oxygen species may contribute to the cardiomyocyte loss in the heart and destroy the endothelial cells in the blood vessel as well as leading to cellular signalling for hypertrophic responses. Previous reports suggest that antioxidant treatment may ameliorate oxidative stress and cardiac remodelling in oxidative stress (Alam, 2019; Ulla *et al.*, 2017). A strong relationship has been found between oxidative stress protection and the role of anthocyanins in cardiovascular diseases. Key findings of anthocyanin and its derivatives for cardiovascular benefits are summarized in **Table 3**. In one study, four anthocyanins were isolated from elderberries and incorporated into the cytosol and plasmalemma of endothelial cells to examine any protective roles, as endothelial dysfunction is part of the initiation and development of vascular disease (Youdim *et al.* 2000). The test result revealed that anthocyanin could be incorporated into endothelial cells and that significant protection was evident against oxidative stress. Endothelium-dependent vasorelaxation was provided by delphinidin in the rat aorta, giving a pharmacological benefit that can be compared with the polyphenolics of red wine from Cabernet-Sauvignon grape variety (Andriambeloson *et al.* 1998). Feeding of purified anthocyanins or anthocyanin rich extracts from black currant (*R. nigrum*) or elderberry (*S. nigra*) showed little influence on the fatty acid patterns or cholesterol levels in the liver of a rat model, but the pigments were able to spare vitamin E (Frank *et al.* 2002). Capillary permeability has been found to be reduced by administration of crude bilberry (*V. myrtillus*) anthocyanin extracts, both orally and via intravenous injection (Kong *et al.* 2003). Prevention of heart attacks through administration of red grape juice or wine have been found to be strongly linked to the roles of these anthocyanin rich preparation in enhancing the capillary strength and permeability, enhancing the nitric oxide (NO) release, reducing inflammation and inhibiting the platelet aggregate formation (Folts 1998). In addition, administration of black currant (*R. nigrum*) extract containing high concentration of anthocyanins, resulted in endothelial-dependent vasorelaxation in rings of rat aorta *in vitro* (Nakamura *et al.* 2002). Also, when rats were pre-

treated to be more susceptible to oxidative damage and fed with anthocyanin-rich extracts, a significant reduction in lipid peroxidation indices and decreased DNA damage were observed (Ramirez-Tortosa *et al.* 2001). Anthocyanins of blueberries (*V. corymbosum*) such as malvidin and its glycosides (malvidin-3-galactoside and malvidin 3-*O*-glucoside) increased the levels of HO-1 and SOD in endothelial cells followed by a decrease in ROS and XO-1 (Huang *et al.* 2016b). The effects of cyanidin 3-*O*-glucoside, delphinidin 3-*O*-glucoside, and pelargonidin 3-*O*-glucoside were investigated on mitochondrial respiratory chain complex I activity in rat hearts, subjected to ischemia for 45 min, which provides evidence that anthocyanins may regulate energy metabolism in ischemia-induced inhibition of ATP production after ischemia (Skemiene *et al.* 2015).

Postmenopausal women who participated in an Iowa Women's Health study showed significant reduction in mortality from the cardiovascular diseases (CVD) after being treated with strawberry (*Fragaria x ananassa*) extracts for a 16 year follow up period (Mink *et al.* 2007). Blueberries-enriched diet also produced a significant decrease in coronary heart diseases related mortality in a study model adjusted for age and energy (Ahmet *et al.* 2009). Red wine intake has been shown to reduce CVD mortality in several studies (Rimm *et al.* 1991; Klatsky 2001). A consistent dose-responsive cardio-preventive effect has been suggested in an analysis of wine consumption relative to CVD risk (Di Castelnuovo *et al.* 2002). Red wine has proven to have greater beneficial effects on lipid metabolism than white wine, probably due to its increased phytochemical content (van Velden *et al.* 2002). There have been significant reductions in blood pressure, inflammatory status, ischemia, and lipid levels in patients with clinically diagnosed vascular diseases when given relatively low-dose anthocyanin therapy (Aviram *et al.* 2004; Sumner *et al.* 2005; Gorinstein *et al.* 2006; Naruszewicz *et al.* 2007). Commercially available grape juice (10 mL/kg) can markedly inhibit platelet activity and experimental coronary thrombosis *in vivo* (Demrow *et al.* 1995). Corn-derived anthocyanins resulted in the myocardium being less vulnerable to ischemia reperfusion injury, in both *in vivo* and *ex vivo* studies, as compared with the anthocyanin-free control (Toufektsian *et al.* 2008). A previous report suggests that anthocyanins containing purple barley extracts (pigmented genotypes of *Hordeum vulgare*) inhibited angiotensin converting enzyme (ACE) significantly, which is a crucial enzyme for hypertension and oxidative stress development in the heart (Lee *et al.* 2013). Delphinidin and cyanin showed interruption of the renin-angiotensin system mediated signalling pathway by inhibiting the ACE activity and decreasing production of mRNA (Parichatikanond *et al.* 2012). Anthocyanins from *Hibiscus sabdariffa* (delphinidin- and cyanidin 3-*O*-sambubiosides) inhibited ACE enzyme activity by competing with the substrate for the active site (Ojeda *et al.* 2010).

Anthocyanins may be effective in improving endothelial function through the adjustment of NO levels. Bilberry (*V. myrtillus*) and chokeberry (*A. melanocarpa*) and other anthocyanin-rich extracts can prevent loss of endothelium-dependent and NO mediated relaxation in porcine arteries *in vitro* (Bell and Gochenaur 2006). Another report suggests that delphinidin enhances NO release and endothelial nitric oxide synthase (eNOS) phosphorylation (Martin *et al.* 2003). A recent investigation showed that cyanidin 3-*O*-glucoside prevented a rise of blood pressure in spontaneously hypertensive rats (Aloud *et al.* 2018). Cyanidin 3-*O*-glucoside affects the interaction between soluble guanylyl cyclase and eNOS, thus increasing production of cyclic guanosine monophosphate (cGMP) by regulating phosphorylation of

eNOS and protein kinase B (Akt) (Xu *et al.* 2004). Moreover, anthocyanin rich extract of purple barley grain showed potent *in vitro* inhibitory activity of angiotensin converting enzymes (Lee *et al.* 2013). A cross-sectional study reported that a higher intake of anthocyanins is associated with lower arterial stiffness (Jennings *et al.* 2012). *A. melanocarpa* extract rich in anthocyanins showed significant decreases in blood pressure in patients suffering from metabolic syndrome (Broncel *et al.* 2010).

Grape products associated with other foods containing polyphenols protect the heart from oxidative stress and inflammation, and activating novel proteins, e.g. Sirtuin 1 that prevent cellular senescence (Dohadwala and Vita 2009). Malvidin 3-*O*-glucoside inhibits NF- κ B in bovine arterial endothelial cells which was also involved in suppression of pro-inflammatory mediators (Paixao *et al.* 2012). Malvidin also inhibited the ICAM-1, monocyte chemo-attractant protein-1(MCP-1) and vascular cell adhesion molecule-1 (VCAM-1) expression that was induced in endothelial cells by TNF- α (Huang *et al.* 2014). Human umbilical vein endothelial cells, when treated with anthocyanins, resulted in the regulation of cholesterol distribution by interfering with recruitment of TNF receptor-associated factors-2 in lipid rafts, thus inhibiting glycoprotein CD40-induced pro-inflammatory signalling (Atalay *et al.* 2003). Delphinidin can reduce the degree of necrotic and apoptotic cell death in cultured cardiomyocytes and also the infarct size after ischemia in rats. The process was mediated by inhibition of activators and signal transducers of transcription-1 (Cines *et al.* 1998). In this study, purple rice (*O. sativa*) extract prevented the rise of toll-like receptor-4, NF- κ B and transforming growth factor- β (TGF- β) expression in heart as well as reduced activation of phospho extracellular signal regulated kinases-1/2, basic fibroblast growth factor and urokinase plasminogen activator in the heart of a diabetic group (Chen *et al.* 2016).

Anthocyanins may also protect against the production of adhesion molecules induced by activated platelets. An investigation involving optimal platelet function revealed that anthocyanins and their colonic metabolites inhibited thrombin peptide-induced, receptor-activating platelet aggregation but had no influence on the reactivity of platelets when strong agonists such as ADP and collagen were present (Liang *et al.* 2006). The beneficial effect of polyphenols on the suppression of platelet-mediated thrombosis was reported previously. Fractions from purple grapes (*V. vinifera*) containing delphinidin inhibited whole-blood aggregation, indicating a possible mechanism for the improvement of CVD (Freedman *et al.* 2001).




Figure 3: Schematic representation of anthocyanin mediated prevention of inflammation, hypertrophy and fibrosis signal in the heart. Angiotensin-II is the regulating factor for the development of hypertrophic signal in the heart through activation of Janus kinase/signal transducers and activators of transcription (JAK-STAT) pathway which may stimulate the transforming growth factor beta (TGF- β) and fibrosis in the heart. Moreover, anthocyanins such as delphinidin, malvidin 3-*O*-glucoside, cyanidine may inhibit the angiotensin-II mediated pathway and nuclear factor kappa B (NF- κ B) to reduce the pro-inflammatory and inflammatory cytokines.

Increased levels of C-reactive protein (CRP) due to low-grade chronic inflammation can be considered an independent risk factor for CVD (Ellulu *et al.* 2017). Among adults in the United States, a significant inverse association between anthocyanin intakes and serum CRP was found upon analysis of the national health and nutrition examination survey data (Chun *et al.* 2008). Data from the United States department of agriculture flavonoid databases also indicated that anthocyanidin intakes were inversely linked with serum CRP concentration (Chun *et al.* 2008). By using sweet cherries (*Prunus auiun*) that were anthocyanin-rich, a recent clinical study showed a decrease in serum CRP level after four weeks of intervention (Carluccio Maria *et al.* 2003). Based on this discussion a possible mechanism for anthocyanin mediated prevention of inflammation, hypertrophy and fibrosis signal in the heart has been proposed in **Figure 3 and Figure 4**.

Table 3. Function of anthocyanin rich natural extracts and anthocyanins in the prevention of cardiovascular complications and diseases

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Anthocyanins	Human aortic culture cardiomyocytes	1 mg/mL used for cell cultured experiments	-Endothelium-dependent vasorelaxant activity. -Reduces the degree of apoptotic and necrotic cell death and also the infarct size after ischemia mediated by the inhibition of signal transducers and activators of transcriptional factors.	(Youdim <i>et al.</i> 2000)
Anthocyanin rich extract of purple barley grain (pigmented genotypes of <i>Hordeum vulgare</i>)	<i>In vitro</i> ACE inhibition	IC ₅₀ 8.77 mg/mL	-Have antioxidant activity and potent angiotensin I-converting enzyme inhibitory capacity.	(Lee <i>et al.</i> 2013)
Anthocyanins and colonic metabolites	Endothelioma cell line	50 to 250 µg/mL	-Inhibits CD40-induced pro-inflammatory signaling, -Protect against production of adhesion molecule induced by activated platelets.	(Atalay <i>et al.</i> 2003)
Extracts of elderberry (<i>Sambucus nigra</i>), bilberry (<i>Vaccinium myrtillus</i>), and chokeberry (<i>Aronia melanocarpa</i>)	64 porcine coronary arteries were isolated and performed <i>in vitro</i> study	0.005–5 mg total anthocyanins/L	-Prevent loss of endothelium-dependent and NO mediated relaxation.	(Bell and Gochenaur 2006)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Blackcurrant extract	Human endothelial cells and 10 female Sprague-Dawley ovariectomized rats	Orally intake 3% (with or without) blackcurrant extract for 3 months	-Up-regulated eNOS mRNA levels and NO synthesis both <i>in vitro</i> human endothelial cells and <i>in vivo</i> ovariectomized rats.	(Horie <i>et al.</i> 2019)
Blueberries	Pulmonary arterial hypertension in rats. Forty-eight male Wistar rats in 7 groups	50, 100, and 200 mg/kg via oral gavage for 5 weeks	-Increased the Early/Late ratio of blood flow across the tricuspid valve and tricuspid annular phase systolic excursion. -Decreased total reactive species concentration and lipid oxidation, reduced activity of nicotinamide adenine dinucleotide phosphate oxidase and expression of xanthine oxidase. -Increased the activity of superoxide dismutase and restored sulfhydryl content	(Turck <i>et al.</i> 2020)
Blueberry anthocyanin-enriched extract	Transverse aortic constriction (TAC)-induced myocardial dysfunction in 30 male mice	0.5 g/kg of blueberry anthocyanin enriched extract was administered daily by oral gavage for 6 consecutive weeks	-Treatment markedly reduced asymmetric dimethylarginine (ADMA) concentration. -Significantly ameliorated heart weight, left ventricular weight, myocardial dysfunction, left ventricular hypertrophy and fibrosis.	(Hu <i>et al.</i> 2020)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Blueberry (Rabbiteye blueberries <i>Vaccinium ashei</i>) supplement	80 diet induced obese C57BL/6 mice were divided into eight groups (n = 10)	Orally intake 6.4 g/kg body weight/ day of blueberries for 8 weeks	-Reduced diastolic and systolic blood pressure in diet-induced obese mice. -Prevention of heart attacks by reducing inflammation, enhancing capillary permeability and strength, inhibiting formation of platelets and enhancing release of NO.	(Shi <i>et al.</i> 2019)
Black currant (<i>Ribes nigrum</i>) extract	Male Sprague-Dawley rats	The thoracic aorta was removed from the rats and induced 10-30 µg/mL of black currant extract	-Endothelial-dependent vasorelaxation.	(Nakamura <i>et al.</i> 2002)
Black mulberry (<i>Morus nigra</i>) fruit extract	50 male Sprague-Dawley rats divided into 5 groups (n=10)	25 and 50 mg/kg/day via intraperitoneal route for 6 weeks	-Significantly reduced total cholesterol, low-density lipoprotein-cholesterol and triglyceride levels and reduced atherosclerotic lesions.	(Jiang <i>et al.</i> 2017)
Cyanidin 3- <i>O</i> -glucoside	Spontaneously hypertensive rats (SHRs)	10 mg/ kg/day gavage for 15 weeks.	-Reduced blood pressure and cardiac hypertrophy.	(Aloud <i>et al.</i> 2018)
Chokeberry (<i>A. melanocarpa</i>) and purple maize (<i>Z. mays</i>)	72 male Wistar rats divided 6 groups each contained 12 rats	Orally approx. 0.8 mg /kg/day for 16 weeks	-Improved cardiovascular structure and function reduced systolic blood pressure, decreased plasma triglycerides and total cholesterol compared to high-fat diet rats.	(Bhaswant <i>et al.</i> 2017)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Commercial grape juice	47 Adult mongrel dogs of either sex	Ranges from 2 to 10 mL/kg intravenous infusion.	-Inhibit platelet; activity and experimental coronary thrombosis <i>in vivo</i> .	(Demrow <i>et al.</i> 1995)
<i>Hibiscus sabdariffa</i> aqueous extract	Two-kidney-one-clip (2K1C) model of hypertension in 42 rats seven groups (n=6/group)	15, 30, or 60 mg/200 g body weight; orally for 2 weeks	-Serum ACE activity and plasma angiotensin II level were significantly reduced. - Reduced blood pressure.	(Nurfaradilla <i>et al.</i> 2019)
Maize (<i>Z. mays</i>)	62 male Wistar rats divided into 2 groups	Special diet containing 20% anthocyanins rich or anthocyanins free diet taken orally for 8 weeks	-Reduced vulnerability of myocardium to ischemia reperfusion injury in both <i>ex vivo</i> and <i>in vivo</i> studies.	(Toufektsian <i>et al.</i> 2008)
Pure delphinidin	64 male Sprague-Dawley rat's left coronary artery isolated	Ventricular myocytes isolated from the rats and applied 10 μ M pure sample to cultured myocytes 2 h prior to the hypoxic injury and during reoxygenation	-Protection from heart treatment was linked with reduced NF-kB expression.	(Scarabelli <i>et al.</i> 2008)
Red Chinese cabbage (<i>B. rapa</i>)	40 male ApoE ^{-/-} mice subdivided into	150 or 300 mg/kg/day (27 or 54 mg cyanidin	-Reduced: (i) plaque formation, (ii) infiltration of leukocytes, (iii) concentrations of blood	(Joo <i>et al.</i> 2018)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
	5 groups each consisted of 8 mice	/kg/day) were fed by gavage for 12 weeks	inflammatory cytokines which lowered the risk of vascular inflammatory diseases.	
Anthocyanidins (blueberries, strawberries, red wine)	34489 Postmenopausal women	1%, 4% and <1% of total consumption of food one time/week for 16 years follow up	-Reduction in CVD mortality.	(Mink <i>et al.</i> 2007)
Anthocyanin-rich blueberry	A double-blind, parallel RCT, 138 eligible aged adults	2 dietary achievable blueberry oral intakes equivalent to 1/2 and 1 cup/day (75/150 g) for 6 months	-Improved endothelial function, systemic arterial stiffness, and attenuated cyclic guanosine monophosphate concentrations.	(Curtis <i>et al.</i> 2019)
Blueberries anthocyanins	63 healthy male volunteers randomized, double-blind, parallel controlled trial	300 mg anthocyanins, equivalent to 200 g of fresh blueberries for 1 month	-Dose-dependent improvement of endothelial function in healthy humans measured by flow-mediated dilation.	(Rodriguez-Mateos <i>et al.</i> 2019)
Blue honeysuckle, cyanidin-3- <i>O</i> -glucoside	A double-blind, counterbalanced, crossover intervention study carried in 20 older adults, aged 62-81 years	Oral drink containing 100 mg, 200 mg, and 400 mg anthocyanins for 1 week	-The 400 mg dose elicited significantly lower diastolic blood pressure and heart rate.	(Bell and Williams 2019)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Commercial red wine	Human	25 g (approximately 2 standard drinks)/ day	-Reduction in CVD mortality.	(Klatsky 2001)
Concord grape juice supplementation (<i>Vitis vinifera</i>) juice or wine	40 men with mild hypertension	Orally intake 5.5 ml/kg body weight/day split over two servings per day, for 8 weeks	-Decreased systolic blood pressure both in SHR and in hypertensive humans and also improved aortic elasticity in stroke-prone SHR.	(Dohadwala and Vita 2009)
Freeze-dried bilberry (<i>V. myrtillus</i>) as a dietary supplement	50 ST-segment elevation myocardial infarction patients	Orally 40 g/day, equivalent to 480 g fresh bilberries for 8 weeks	-A significant improvement was found in walk test and blood lipid profiles were altered within the bilberry group that could potentially translate the reduction of CVD.	(Arevstrom <i>et al.</i> 2019)
New Zealand blackcurrant extract	Ten adult healthy male	Orally 1.87 mg total anthocyanins/kg bodyweight for 1 week	-Acute ingestion of a single dose of blackcurrant extract maintained forearm blood flow and forearm vascular resistance during an extended period of sitting.	(Barnes <i>et al.</i> 2020)
New Zealand blackcurrant extract	14 older adults, randomized, double-blind, placebo-controlled, cross-over design	600 mg/day for 1 week	-Decrease in systolic and diastolic blood pressure.	(Cook <i>et al.</i> 2020)
<i>V. arctostaphylos</i> , berry hydro-alcoholic extract	Randomized placebo-controlled trial in	3-month intake of 400 mg extract capsule three times daily	-Systolic blood pressure and diastolic blood pressure decreased significantly.	(Kianbakht and Hashem-Dabaghian 2019)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
	overweight/obese 50 hypertensive patients			




Figure 4: Schematic representation of anthocyanin mediated prevention of oxidative stress in endothelial cells and relaxation of smooth muscle in blood vessels. Endothelial nitric oxide (eNOS) regulates the production of nitric oxide (NO) in the endothelium which is a signalling molecule responsible for vasodilation and vascular tone. Superoxide can directly interact with the NO and produces peroxynitrite radicles ($^{\cdot}\text{ONOO}^{\cdot}$). These peroxynitrite radicles may trigger endothelial cell destruction and causes vascular dysfunction. Anthocyanins such as cyanidin 3-*O*-glucoside may directly interact with superoxide to scavenge them or may increase the cyclic guanosine mono phosphate (cGMP) level to relax the smooth muscle of blood vessels.

Anthocyanins in diabetes and obesity

Many recent studies suggest that eating fruits and vegetables, especially those rich in polyphenols, reduce the occurrence of type-2 diabetes, a condition associated with insulin resistance (Anderson and Polansky 2002; Landrault *et al.* 2003; Anderson *et al.* 2004; Lachin and Reza 2012; Putta *et al.*, 2018). Our previous report also showed that antioxidant compounds prevented the oxidative stress and associated complications in insulin resistance and obesity (Alam *et al.*, 2016). In this part, we discussed that anthocyanins and natural fruits and vegetables rich in anthocyanins may influence and modulate the diabetes condition by improving insulin resistance, improved glucose absorption in cellular level, prevented glucose absorption by inhibiting key enzymes such as alpha amylase and beta glucosidase, decreased cholesterol level by increasing lipid metabolism and lowered inflammatory states in the adipose tissues. **Table 5** summarizes the beneficial effects of anthocyanins in diabetes and obesity. Insulin resistance is a disorder, where there is inadequate stimulation of glucose transport in skeletal fat and muscle by insulin, and it also enhances

hepatic glucose production. Anthocyanins and anthocyanidins were found to protect pancreatic β -cells from glucose-induced oxidative stress in several studies (Al-Awwadi *et al.* 2005; Jayaprakasam *et al.* 2005). The glycoside and dimethoxy ether of leucopelargonidin, isolated from the bark of the Indian banyan tree (*Ficus bengalensis*), have shown significant hypolipidemic, hypoglycaemic and serum insulin-raising effects in moderately diabetic rats, being comparable with the effects of glibenclamide (Cherian *et al.* 1992; Augusti *et al.* 1994; Daniel *et al.* 2003). In addition, *Cornus* fruits which are a rich source of anthocyanins have been reported to possess anti-diabetic activity (Yamahara *et al.* 1981; Seeram *et al.* 2002). White skin sweet potato resulted in improved diabetes, glucose tolerance, hyperinsulinemia and hyperlipidaemia, as well as lowered free fatty acids in Zucker fatty rats (Kusano *et al.* 2001). Recent investigation has shown that purple sweet potatoes (*I. batatas*) have also improved diabetes (Qin *et al.* 2019). Anthocyanins improved glucose stimulated glucose absorption and insulin secretion in INS-1E and human hepatoma cell (HepG2) cells (Luna-Vital and Gonzalez de Mejia 2018). It was also suggested that anthocyanins from purple corn significantly reduced gluconeogenesis by suppressing phosphoenolpyruvate carboxykinase expression and AMPK phosphorylation in HepG2 cells (Luna-Vital and Gonzalez de Mejia 2018). Mulberry (*Morus alba*) extract containing anthocyanins also improved glucose uptake, utilization, increased glycogen formation and diminished the insulin resistance in HepG2 cells (Yan *et al.* 2016).

Anthocyanins (Lachin and Reza 2012) especially pelargonidin-3-*O*-galactoside and its aglycone, pelargonidin was able to increase insulin secretion by selectively inhibiting COX-2 enzyme (Zhang *et al.* 2004). The study suggested that cherries (*Prunus* genus), grapes (*V. vinifera*) and other anthocyanin containing berries might have a role in the prevention of type-2 diabetes. Anthocyanin extracts showed potent α -glucosidase inhibitory activity, suppressing an increased postprandial glucose level in some *in vitro* and animal studies (Matsui *et al.* 2001; Matsui *et al.* 2004). Cyanidin and its glycosides showed inhibitory activities of intestinal α -glucosidase and pancreatic α -amylase *in vitro* and also showed synergistic effect with acarbose (Akkarachiyasit *et al.* 2010). α -Glucosidase and pancreatic lipase enzyme inhibition was associated with anthocyanins isolated from whole berries and skin of Muscadine grapes (*V. vinifera*) (You *et al.* 2011). Extracts from the pulp from several sweet cherry (*P. avium*) cultivars showed α -glucosidase enzyme inhibition (Goncalves *et al.* 2017). Black chokeberry (*A. melanocarpa*) extracts also showed α -amylase and lipase/enzyme inhibition (Worsztynowicz *et al.* 2014).

Diabetes results in various microcirculatory disorders. Many of them may occur before microangiopathic lesions begin to form thickening of capillaries in many areas such as the eye and are assumed critical in the pathogenesis of microcirculatory complications involved with diabetes. The number of leucocytes sticking to the venular endothelium and microvascular permeability are increased in the diabetic microangiopathic condition (Valensi *et al.* 1997; Valensi *et al.* 1998;). Delphinidin chloride showed a reduction of leucocytes adhering to the venular vessels and increased microvascular permeability in diabetic hamsters (Bertuglia *et al.* 1995). Several flavonoids, including anthocyanosides, have been effective against experimentally induced capillary filtration (Gabor 1972; Parmar and Ghosh 1977). In one animal study, it was shown that anthocyanosides can improve and even normalize capillary filtration of albumin (Cohen-Boulakia *et al.* 2000). Endothelium-dependent vasorelaxation by different vasodilator

agonists is reduced in various conditions including diabetes (Griffiths and Smith 1972). One of the mechanisms that resulted in dysfunction of the endothelium was a decrease in the release of nitric oxide (NO) (Barton *et al.* 1997). Extracts from red wines, other grape products and various plant polyphenols (mainly anthocyanins) were found to produce endothelium-dependent vasorelaxation, probably through NO release or due to enhanced biological activity of NO (Fitzpatrick *et al.* 1993; Fitzpatrick *et al.* 1995; Andriambeloson *et al.* 1998). A combination of anthocyanins of bilberry (*V. myrtillus*) was reported to have pharmacological and biological properties, including vasorelaxation and prevention of hypertension (Mykkänen *et al.* 2014).

Increased levels of triglyceride (hypertriglyceridemia) were strongly associated with the insulin resistance syndrome, with obesity being strongly associated with insulin resistance. Thus, a reduction in insulin resistance is important in preventing the development of type-2 diabetes. It was demonstrated that cyanidin 3-*O*-glucoside-rich purple corn may improve high fat diet-induced insulin resistance in mice (Tsuda *et al.* 2003). Consumption by diabetic patients of pomegranate (*P. granatum*) juice resulted in antioxidative effects in their serum and reduced the oxidative stress in their monocytes/macrophages levels. These changes were attributed specifically to anthocyanins (Gil *et al.* 2000; Rosenblat *et al.* 2006). Extracts of anthocyanin and procyanidins increased high-density lipoprotein (HDL) cholesterol levels while decreasing triglycerides in rats (Al-Awwadi *et al.* 2005). A recent investigation showed that delphinidin effectively modulated lipid metabolic gene expression in human HepG2 Cells and reduced triglyceride accumulation *in vitro* (Parra-Vargas *et al.* 2018). However, delphinidin failed to change body weight gain, energy intake, histological abnormalities, hyperglycemia or insulin resistance elicited by the high fat high carbohydrate diet (Parra-Vargas *et al.* 2018). Another report suggested that cyanidin and delphinidin consumption mitigated high fat diet-induced obesity, dyslipidaemia, insulin resistance and oxidative stress followed by the inhibition of NF- κ B and Jun *N*-terminal kinases (JNK) activation as well as protein tyrosine phosphatase-1B overexpression (Daveri *et al.* 2018). Blackberries containing 57% malvidin and 33% petunidin or peonidin increased mitochondrial respiration thus reducing metabolic damage related to a high-fat diet (Skates *et al.* 2018). Blueberry (*V. corymbosum* and *V. ashei*) containing anthocyanins were responsible for the reduction of glucose levels, triglyceride, cholesterol and leptin in high fat diet fed C57BL/6 mice (DeFuria *et al.* 2009; Wu *et al.* 2016a). Red cabbage (*Brassica oleracea*) and red cabbage microgreen (were harvested without roots, shipped overnight with specialized clamshell containers) supplementation reduced low-density lipoprotein and cholesterol in high fat diet fed C57BL/6 mice (Huang *et al.* 2016a). Mulberry (*M. australis*) and sweet cherry (*P. avium*) extract rich in cyaniding prevented the inflammatory cytokines (e.g. TNF- α , IL-6, iNOS and NF- κ B) and improved insulin resistance in high fat diet fed mice (Wu *et al.* 2016b). Black soybean (*G. max*) extract also showed decreased triglyceride and cholesterol as well as reduced cytokine production in high fat diet fed rats (Kim *et al.* 2015). Black soybean (*G. max*) also showed reduced triglyceride and cholesterol level in overweight and obese Korean adults (Lee *et al.* 2016). A recent investigation suggested that raspberry (*R. idaeus*) anthocyanin consumption elevated GPx and serum SOD activities as well as fecal butyric acid levels which can ameliorates diet induced obesity by alleviating oxidative stress (Wu *et al.* 2018). Raspberry (*R. idaeus*) anthocyanin consumption also reduced hepatic lipid and serum profiles, while markedly down-regulating the expression of TNF α , IL-6 and NF- κ B

genes (Wu *et al.* 2018). Blue berry anthocyanins showed altered mitogen-activated protein kinase and NF- κ B-mediated stress signalling pathways and gene expression in high fat diet fed male C57Bl/6j mice (DeFuria *et al.* 2009). The primary site of energy storage is the adipocytes that are known to accumulate triacylglycerol during nutritional excess. Recently, it has been established that adipocyte dysfunction has an important role in the development of obesity and insulin resistance. Adipocytes synthesize and secrete biologically active molecules called adipocytokines among which adiponectin is important (Shimomura *et al.* 1996). In obese and insulin resistant state, plasma adiponectin concentration and mRNA expression level are decreased (Arita *et al.* 2012). Purple sweet potato (*I. batatas*) containing 3-*O*-caffeoylferuloylsophoroside-5-*O*-glucoside and 3-*O*-caffeoylferuloylsophoroside-5-*O*-glucoside decreased the leptin (adipogenic marker) as well as decreased the production of COX-2, MCP-1 and IL-6 in 3T3-L1 cells (Ju *et al.* 2011). Anthocyanins can regulate obesity and insulin sensitivity associated with adipocytokine secretion in adipocytes. This provides a biochemical basis for the use of anthocyanins which will have significant implications for the prevention of diabetes and obesity (Tsuda *et al.* 2003). Anthocyanin-rich mixed grape-bilberry juice supplementation in Fischer rats reduced serum leptin and resistin, but did not influence serum adiponectin and secretion of adipokines from mesenteric adipose tissue (Graf *et al.* 2013). Blueberry (*V. corymbosum*) powder supplementation reduced triglycerides, fasting insulin, insulin resistance, and plasma glucose level in Zucker fatty rats fed with a high fat diet (Seymour *et al.* 2011). Blueberry (*V. corymbosum*) intake increased adipose and skeletal muscle peroxisome proliferator-activated receptor (PPAR) activity, reduced abdominal fat mass and affected PPAR transcription involved in glucose uptake/oxidation and fat oxidation (Seymour *et al.* 2011). Thus, these studies demonstrated that anthocyanins can modulate the gene expression of adipocytokines in humans and may have a distinct therapeutic advantage for the regulation of adipocyte function (Tsuda *et al.* 2005; Tsuda *et al.* 2006).




Figure 5: Schematic representation of anthocyanin mediated prevention of inflammation and fat metabolism in liver. Inflammation and cytokines leads to insulin resistance in diabetes and obesity. TNF- α signalling stimulates the production of inflammatory cytokines in liver through NF- κ B pathway. Anthocyanins from various plant sources can interact with the regulatory kinase molecule AMPK responsible for the overall metabolism and ATP production in the cell. AMPK is also responsible for the down regulation of inflammatory cytokine production. Anthocyanins stimulates AMPK pathway resulting in increased utilization of cellular glucose and fatty acids and decreasing inflammation in liver.

Increased lipid metabolism and utilization in the liver and other organs are key pathways through which fats are removed from the body. High fat diet feeding may induce fat accumulation in the liver and started steatosis to non-alcoholic fatty liver diseases. Microarray analysis of hepatic gene expression profiles indicated that PPAR signalling pathway, steroid biosynthesis, fatty acid metabolism and biosynthesis were modulated by sweet cherry (*P. avium*) anthocyanins in C57 BL/6 J mice fed with a high fat diet (Song *et al.* 2016a). Sweet cherry (*P. avium*) anthocyanins are also capable of reducing the hepatic steatosis in high fat diet fed mice (Wu *et al.* 2014; Song *et al.* 2016a). Mulberry (*M. alba*) anthocyanin extract was tested on hepatocytes cultured with high fatty acid. This extract enhanced fatty acid oxidation and suppressed fatty acid synthesis, which contributed to ameliorating lipid accumulation induced by oleic acid (OA) in human HepG2 as the cell model (Chang *et al.* 2013). This investigation

also identified that mulberry (*M. alba*) anthocyanin extract stimulated AMPK and inhibited acetyl coenzyme A carboxylase activities (Chang *et al.* 2013). Mulberry (*M. alba*) anthocyanin extract also attenuated the expression of sterol regulatory element-binding protein-1 (SREBP-1) and fatty acid synthase (FAS) in human HepG2 (Chang *et al.* 2013). Black elderberry (*S. nigra*) anthocyanins significantly lowered liver weights, serum TAG, serum monocyte chemo-attractant protein-1, serum insulin and TNF α , followed by the reduction of hepatic fatty acid synthase mRNA in the liver of high fat diet fed mice (Farrell *et al.* 2015). Based on this discussion a possible mechanism for inflammation and fat metabolism in liver has been proposed in **Figure 5**.

Table 4. Function of anthocyanin rich natural extracts and anthocyanins in the prevention of diabetes and obesity.

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Purple maize extract	RAW 264.7 macrophages and 3T3-L1 adipocytes	Pretreated with 1 mg/mL anthocyanin-rich water extracts	-Down-regulating pro-inflammatory mediator production in inflamed macrophages and adipocytes. -improving insulin sensitivity in insulin-resistant adipocytes	(Zhang <i>et al.</i> 2019a)
Cyanidin-3- <i>O</i> -glucoside-rich purple corn	Rat and human adipocytes	100 μ M extracts were used in rat and human cells	-Improve high fat diet-induced insulin resistance in rat and human.	(Tsuda <i>et al.</i> 2005; Tsuda <i>et al.</i> 2006)
delphinidin chloride	Eighty male (n=10, 8 groups) BALB/c mice were used in streptozotocin (STZ)-induced diabetes	100 mg delphinidin chloride/kg/24 h was intravenously injected for eight weeks	-Inhibited the protein glycation in diabetes mellitus and reduced the glycation rate of hemoglobin A1c.	(Gharib <i>et al.</i> 2013)
Delphinidin	HT-29 enterocyte-like human cell line and RF/J mice	Cell line was treated with 50 μ M delphinidin and mice were treated with 100 μ M delphinidin.	-Glucose absorption is inhibited in both mouse jejunum and a human enterocytic cell line in a free fatty acid receptor-dependent manner.	(Hidalgo <i>et al.</i> 2017)
Grape skin or whole grapes (<i>V. vinifera</i>), berries (<i>Rubus</i>)	<i>In vitro</i> α -glucosidase Inhibitory, insulin secretion assay were	0.5 mg/mL for α -glucosidase inhibitory assay.	-Enhance insulin secretion Inhibits COX-2 enzymes selectively -Have potent alpha-glucosidase inhibitory activity that suppress the increase in postprandial glucose level	(Matsui <i>et al.</i> 2001; Zhang <i>et al.</i> 2004; Griffiths and Smith 1972)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
and cherry (<i>Prunus</i>)	performed on INS-1 cells. <i>in vivo</i> was performed through male albino Wistar rats	<i>In vivo</i> oral study was performed via administration of 15 g/day not less than 2 weeks.	-Inhibitory activities for lens opacity Improve vision and prevent diabetic retinopathy.	
Pomegranate (<i>P. granatum</i>) juice	<i>In vitro</i> ABTS, DPPH and FRAP assay. 20 male participants Tested in rat model (n=9)	Soluble solids of juice value were ranged from 15.5 – 16.6%. 50 mL of pomegranate juice per day (which contain 1.5 mmol of total polyphenols). The dose of 21.42 mg/kg/day of total polyphenols orally by gavage for 6 weeks.	-Anthocyanins had anti-oxidative effects, which resulted in anti-oxidative effects on serum which might help reduce atherosclerosis development in these patients. -Decrease triglycerides and increase HDL-cholesterol levels in rats.	(Al-Awwadi <i>et al.</i> 2005; Gil <i>et al.</i> 2000; Rosenblat <i>et al.</i> 2006)
Lingon berry extract (<i>V. vitis-idaea</i>)	<i>In vitro</i> and <i>in vivo</i> studies HCD -induced hypercholesterolaemic	200 and 400 mg/ kg of body weight per day for	-Reduce the inflammatory cell infiltration and attenuate steatosis and hepatocellular fat deposit in the liver.	(Zhang <i>et al.</i> 2019c)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
	40 mice divided into four groups (10 mice per group)	10 weeks as dietary supplement.		
Anthocyanins and anthocyanosides from <i>Cornus</i> species	<i>In vitro</i> rodent pancreatic beta-cells and male Wistar rats with streptozotocin (STZ)-induced diabetes in 3 groups of 15 rats	Daily dosage of 40 mg/kg for 12 weeks.	-Protect pancreatic β -cells from glucose induced oxidative stress and improve and normalize capillary filtration of albumin.	(Cohen-Boulakia <i>et al.</i> 2000; Jayaprakasam <i>et al.</i> 2005)
Polymerized anthocyanin	40 male C57bl6/J mice divided into 4 groups induced nonalcoholic fatty liver disease.	400 mg/kg/day were given orally once a day for 12 weeks	-Effectively reduced TC and LDL-C. -Inhibited the activation of inflammatory pathways, depressing oxidative stress through increased antioxidant levels, and increasing β -oxidation to inhibit mitochondrial dysfunction.	(Fan <i>et al.</i> 2019)
Bark of the Indian banyan tree (<i>F. bengalensis</i>)	Moderately diabetic mice, rats and alloxan diabetic dogs	Different range of diet from 0.2-1.8 g/kg/day to different groups of animals for one month	-Significant hypoglycaemic, hypo-lipidemic and serum insulin-raising effects.	(Cherian <i>et al.</i> 1992; Augusti <i>et al.</i> 1994; Daniel <i>et al.</i> 2003)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Blackcurrant (<i>Ribes nigrum</i>) extract	OVX female Sprague-Dawley and sham surgery rats (3 groups, n = 9–10 rats/group)	Consumed 38 g/100 g blackcurrant extract (3%) orally for 3 months	<ul style="list-style-type: none"> -Decreased expression of hepatitis-related genes, such as tumour necrosis factor-α, <i>IL-6</i>, and <i>IL-1β</i>. -levels of serum triglycerides, total cholesterol, and LDL cholesterol decreased. 	(Nanashima <i>et al.</i> 2020)
Tart cherry extract	35 High fat diet fed mice were divided into 3 groups and	Consumed 60 mg/kg anthocyanin-rich tart cherry extract in a daily dosage for 6 weeks	<ul style="list-style-type: none"> -Failed to reverse the effects of the high fat diet on body weight and glucose tolerance. -Significantly reduced the leptin and IL-6 levels. -Significant enhancement in antioxidant capacity and SOD activity. 	(Nemes <i>et al.</i> 2019)
Sweet cherries (<i>P. auium</i>)	48 male C57BL/6 high-fat diet mice were divided into four groups	Consumed 2 mg/kg of body weight for 12 weeks	<ul style="list-style-type: none"> -Purified sweet cherries reduce the expression levels of IL-6 and TNFα genes and markedly increase the SOD and GPx activity. 	(Wu <i>et al.</i> 2014)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Cornelian Cherry (<i>Cornus mas</i>) and Japanese cornelian cherry (<i>C. officinalis</i>)	32 high-fat-fed insulin resistance C57BL/6 mice were divided into 4 groups. Streptozotocin induced diabetes rats model took 6 groups in each group contained average 10 rats and dose	C57BL/6 mice were fed with high-fat diet plus 1 g/kg of anthocyanins for 8 weeks Streptozotocin induced diabetic rats were treated with 50 mg/kg intravenous administration	-Body weight was decreased, normalized glucose intolerance, elevated circulating insulin, and a dramatic decrease in liver lipid.	(Yamahara <i>et al.</i> 1981; Jayaprakasam <i>et al.</i> 2006)
Blueberries V. <i>ashei</i> and yoghurt supplement	Diet induced obese C57BL/6 mice divided into eight groups (n = 10 for each group)	Consumed 6.4 g/kg of blueberries, 0.02 g/kg of cyanidin-3- <i>O</i> - β -glucoside and 3 g/kg of yoghurt body weight/ day during the 8 week treatment period	-Cyanidin-3- <i>O</i> - β -glucoside and the combination of peptides showed significant reduction of body fat and improved intraperitoneal glucose tolerance.	(Shi <i>et al.</i> 2019)
Chokeberry (<i>A. melanocarpa</i>) purple maize; (<i>Z. mays</i>)	72 male Wistar rats were divided into six separate groups (n = 12 each)	Supplemented orally with chokeberry juice 50 ml/kg/day or purple maize flour 50 g/kg/day for 16 weeks	-Reduced visceral adiposity index and total body fat mass; improved glucose tolerance and liver structure and function.	(Bhaswant <i>et al.</i> 2017)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	Reference
Anthocyanins	169 participants with dyslipidemia randomized, double-blind, placebo-controlled human trial	Two capsules (dose 40, 80, and 320 mg/day) twice daily preferably 30-min after breakfast and supper for 12 weeks	-320 mg/day anthocyanin supplementation reduced serum IL-6, TNF- α , MDA and urine 8-iso-PGF _{2α} and 8-OHdG than 80 mg/day and 40 mg/day anthocyanins. -Improved the anti-oxidative and anti-inflammatory capacity in a dose–response manner.	(Zhang <i>et al.</i> 2020a)
Purified anthocyanins	Randomized, double-blind, placebo-controlled human trial (Participants n = 111)	Orally five doses of anthocyanins (20, 40, 80, 160, or 320 mg/d) were asked to take once daily after meals for 14 days	-80 mg/day of anthocyanin showed the lowest baseline-adjusted fasting plasma glucose level. -8-iso-prostaglandin F _{2α} levels decreased with increasing anthocyanins dose. -Plasma interleukin-10 levels were negatively correlated with increasing anthocyanin.	(Guo <i>et al.</i> 2020)

Anthocyanins in Neurodegenerative Diseases

Anthocyanins have a strong antioxidant capacity, which can be very effective in several models of neurodegenerative diseases (Miller *et al.*, 2018; Stintzing *et al.* 2002; Shih *et al.* 2011; Traustadottir *et al.* 2009). They have a high oxygen radical absorbance capacity (ORAC) value, which is a major part of their neuroprotective function (Zafra-Stone *et al.* 2007; Zhu *et al.* 2010). In addition, anthocyanins act as antioxidants as they are able to directly trap free radicals, thus preventing ROS formation in affected cells. For example, anthocyanins can decrease the generation of ROS in *in vitro* models of alpha-beta peptide-induced toxicity, as well as in hydrogen peroxide injury (Shih *et al.* 2011; Hwang *et al.* 2012). Moreover, by using electron spin resonance spectroscopy it has been found that anthocyanins have a strong affinity to scavenge DPPH, alkyl, and hydroxyl free radicals in a dose dependent fashion (Hwang *et al.* 2012). Benefits of anthocyanins in neurodegenerative diseases are presented in **Table 5**.

Initiation of inflammatory gene expression and subsequent production of interleukins and pro-inflammatory cytokines is often recorded in neurodegeneration. Targeting these inflammatory processes may prove beneficial in limiting neuronal apoptosis associated with the disease. Anthocyanins display significant anti-inflammatory properties, given that they can inhibit various inflammatory biomarkers (e.g. interleukin-8 (IL-8)) (Zafra-Stone *et al.* 2007). As well as decreased IL-8 production, pomegranate (*P. granatum*) anthocyanins inhibit activation of NF- κ B (Rasheed *et al.* 2009) and inflammatory markers including activated p-NF- κ B (phosphorylated NF- κ B) (Rehman *et al.* 2017) which are responsible for the expression of several pro-inflammatory genes. The anthocyanins were shown to inhibit a number of other bio-molecules associated with the expression of several pro-inflammatory cytokines. Furthermore, cherry and blackberry (*R. fruticosus*) anthocyanins have been proven to be powerful COX-2 inhibitors, which is an important pro-inflammatory enzyme employed in the synthesis of prostacyclins (Saric *et al.* 2009; Zdarilova *et al.* 2010). At high concentrations (250 μ g/mL), anthocyanins have inhibited up to 95% of cyclooxygenase activity (Mulabagal *et al.* 2009). These findings indicate that anthocyanins may have a significant role in preventing inflammatory processes associated with neurodegenerative disease. A recent investigation also suggests that supplementation of blackcurrant (*Ribes nigrum*) anthocyanins resulted in increased cyclic glycine-proline (cGP) in the cerebrospinal fluid of patients suffering from Parkinson disease (Fan *et al.* 2018). Anthocyanins increased cGP, which is a neuropeptide that facilitates IGF-1 function in brains of the patients with Parkinson disease (Fan *et al.* 2018). Moreover, anthocyanin could also reduce cerebral oedema while reducing the matrix metalloproteinase-9 activity in reperfusion injury (Pan *et al.* 2018). In addition, NF- κ B and the NOD-like receptor pyrin domain-containing protein-3 inflammasome pathways were inhibited and TNF- α , IL-6 and interleukin-1 β levels were decreased by anthocyanin treatment (Pan *et al.* 2018). Anthocyanin extracts from blackcurrants (*R. nigrum*), blueberries (*V. corymbosum*) and hibiscus (*H. sabdariffa*) prevented dopaminergic cell death, microglial activation and amelioration of mitochondrial dysfunction in rotenone-induced cell culture model of Parkinson diseases (Strathearn *et al.* 2014). Another report showed that cyanidin 3-O-glucoside exerted a neuroprotective effect against ischemic stroke in mice (Min *et al.* 2011). Black soybean (*G. max*) anthocyanin extract showed significant U87 glioma cells survival when exposed to oxidative stress induced by oxygen-glucose deprivation (Kim *et al.* 2012).

An extract high in anthocyanins from blackcurrant (*R. nigrum*) and bilberry (*V. myrtillus*) showed beneficial effects and reduced behavioural abnormalities in a mouse model of Alzheimer's disease (Vepsäläinen *et al.* 2013). In addition, Qin, L. *et al.* (2013) suggested that cyanidin 3-*O*-glucoside rescued the cognitive impairments in beta-amyloid peptide-induced cognitive deficits in the rat model of Alzheimer's disease (Qin *et al.* 2013). Cyanidin 3-*O*- β -glucopyranoside was shown to improve cerebral glucose uptake, alleviate cognitive impairment and decrease fasting blood glucose levels in the APP^{swe}/PS1 Δ E9 mouse model of Alzheimer's disease (Song *et al.* 2016b). Moreover, black soybean (*G. max*) anthocyanins have been investigated against beta-amyloid induced neurotoxicity *in vitro* on cell line and *in vivo* in rat model. The results showed increased amyloid beta production in the nervous system which ultimately protected beta-amyloid-induced neurodegeneration (Badshah *et al.* 2015). Shah *et al.* (2015) also worked on black soybean (*G. max*) anthocyanins but used the ethanol-induced oxidative stress (Shah *et al.* 2015) and glutamate-induced oxidative stress (Shah *et al.* 2016) rat models. They found neuroprotection via inhibition of glutamate related neurotransmission, neuronal apoptosis (Shah *et al.* 2015) and protection against glutamate-induced AMPK induction, ROS production, neuroinflammation and neurodegeneration (Shah *et al.* 2016).

Anthocyanins also have the capacity to modulate cognitive and motor function, to enhance memory, and to possibly reduce age-related declines in neural function. Administration of isolated, semi-purified anthocyanins from purple sweet potato (*I. batatas*) improved cognitive performance in mice, and also inhibited peroxidation of lipids in rat brain tissues (Cho *et al.* 2003). Administration of blueberry extracts with significant anthocyanin content (but not purified pigments), led to reversal of age-related deficits in various neural and behavioural parameters (motor and memory functions) (Joseph *et al.* 1999). Further investigations demonstrated that anthocyanins were extremely bioavailable in endothelial cells, which was correlated with their prevention of neurodegenerative disorders and atherosclerosis (Youdim *et al.* 2000; Youdim *et al.* 2002). Anthocyanins improved memory and learning of rats with an estrogen deficit triggered by ovariectomy (Varadinova *et al.* 2009). *Pandanus amaryllifolius* (Pandan leaves) and *Z. mays* (purple waxy corn) extracts possess anthocyanins and showed cognitive enhancing effects, improved memory function, prevented oxidative stress and modulated cholinergic function in ovariectomized rats (Kirisattayakul *et al.* 2017). Another report suggests that supplementation with pure cyanidin 3-*O*-galactoside and blueberry extracts improves spatial memory and regulates hippocampal extracellular signal-regulated kinases expression in senescence-accelerated mice (Tan *et al.* 2014).

Table 5. Function of anthocyanin rich natural extracts and anthocyanins in neurodegenerative diseases.

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	References
Anthocyanins	<i>In vitro</i> models of alpha-beta peptide-induced toxicity and hydrogen peroxide injury. Inhibitory effect of berry anthocyanins on human gastric cancer cells MKN45	50 μ M of extract for ROS assay and $\alpha\beta$ peptide-induced toxicity, 30 μ L anthocyanin oligomers solution for DPPH, 2 mL of the sample solution for lipid peroxidation inhibition assay. 0.5% of berry extract were used to examine the inhibitory effect of berry anthocyanins on <i>Helicobacter pylori</i> -induced IL-8 production in gastric MKN 45 cells	-Decrease the generation of ROS. Scavenge DPPH, alkyl, and hydroxyl free radicals in a dose dependent fashion. -Anthocyanins display significant anti-inflammatory properties by inhibiting various inflammatory biomarkers, including IL-8.	(Hwang <i>et al.</i> 2012; Shih <i>et al.</i> 2011; Zafra-Stone <i>et al.</i> 2007)
Pomegranate (<i>P. granatum</i>)	KU812 cell line, Enzyme-linked immunosorbent assay, western blot analysis and transient transfection and luciferase activity assay	Pomegranate fruit extract (20–100 μ g/mL)	-Inhibit the activation of nuclear transcription NF κ B responsible for the expression of several pro-inflammatory genes. -Inhibit other bio-molecules responsible for the expression of pro-inflammatory cytokines.	(Rasheed <i>et al.</i> 2009)
Pomegranate extract	U-87 cells and rats (n=4)	Receiving 600 mg/kg/day oral administration	- Decreasing the MORs and cAMP protein levels in U-87 cells. -A significant decrease in cAMP responsive element binding protein (CREB) level and an	(Ridzwan <i>et al.</i> 2020)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	References
			increase in Brain-derived neurotrophic family (BDNF) as compared to rats treated with morphine.	
Bilberry anthocyanins	44 Alzheimer's disease model APP/PSEN1 mice (n=5-6).	20 mg/kg/day bilberry anthocyanins was given via gavage for three months	<ul style="list-style-type: none"> -Improves learning and memory abilities and reverses defects to cognitive functions. -Decreases serum and brain lipopolysaccharide (LPS) levels and increases fecal short-chain fatty acid content. -Downregulates the expression of inflammatory factors (TNF-α, NF-Kβ, IL-1β, IL-6, COX-2, iNOS and CD33) and chemokine receptor CX3CR1. -Decreases hippocampal neuroinflammatory responses, and induces phagocytosis of microglia to beta-amyloid protein plaques 	(Li <i>et al.</i> 2020)
Anthocyanin	Mice were subjected to laparotomy 7groups and each groups contains 12 mice.	50–100 mg/kg orally and 10 mg/kg intraperitoneal injection for 4 weeks before surgery and then once per day for 4 week after surgery	<ul style="list-style-type: none"> -Improved learning and memory in mice after surgery. -Significantly reduced neuroinflammation and microglia activation 	(Zhang <i>et al.</i> 2020b)
Anthocyanins	LPS-induced neurotoxicity, 15–20 mice/group divided	24 mg/kg/day for 2 weeks	-Prevented ROS production, inhibited neuroinflammation and neurodegeneration, and improved memory functions in LPS-treated mice.	(Khan <i>et al.</i> 2019)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	References
	into three groups. Mice injected with LPS		-Prevented neuroinflammation by lowering the levels of inflammatory markers (p-NF- κ B, TNF- α , and IL-1 β).	
Cherry (<i>Prunus cerasus</i>)	Male CBA/Hr mice 3 groups and 10 mice in each group	5 g per day for 14 days of commercial food pellets containing 4 mL of 10 and 50% dilution of original cherry juice	-Inhibits COX-2 responsible for pro-inflammatory enzyme required in the synthesis of prostacyclin	(Saric <i>et al.</i> 2009)
Purple sweet potato (<i>I. batatas</i>)	Male Sprague-Dawley (SD) rats for brain homogenate preparation and male ICR mice (4 groups)	25 mg/kg body weight orally once a day for 7 days	-Enhance cognitive performance. -Inhibits lipid peroxidation	(Cho <i>et al.</i> 2003)
Blueberry (<i>V. corymbosum</i>)	40 male Fischer 344 rats were divided into 4 groups	1.86% blueberry (w/v) as a supplement for 8 weeks	-Effective reversal of age-related deficits in various neural and behavioural parameters	(Joseph <i>et al.</i> 1999)
<i>Lycium ruthenicum</i> extract	d-galactose (d-gal)-treated rats were divided in 5 groups	50-200 mg/kg once daily for 7 weeks	-Anthocyanins reduced receptor for advanced glycation end products (RAGE) and suppressed oxidative stress caused by d-gal. - Anthocyanins suppressed microgliosis and astrogliosis. - Reduced the overexpression of nuclear factor kappa B (NF- κ B), interleukin-1 β (IL-1 β), cyclooxygenase-2 (COX-2), and tumour necrosis factor- α (TNF- α).	(Chen <i>et al.</i> 2019)

Food extracts containing anthocyanins	Model	Dose	Experimental Outcomes	References
			-Lowered C-jun N-terminal kinase (p-JNK), caspase-3 levels, and the B-cell lymphoma 2-associated X protein/B-cell lymphoma 2 (Bax/Bcl-2) ratio.	
Freeze-dried blueberry	13 men and 24 women, (60–75 years) were participated in a randomized, double-blind, placebo-controlled trial	24 g/day, equivalent to 1 cup of fresh blueberries for 90 days	On the California verbal learning test, participants in the blueberry group showed significantly fewer repetition errors which may improve cognition among older adults.	(Miller et al., 2018)
Whole frozen blueberries	94 participants were participated in a randomized, double blind, parallel groups, and placebo-controlled trial	Intake of 12.5 g/day orally for 12 to 24 weeks trial	Supplementation improved cognition.	(McNamara et al., 2018)

Anthocyanins in Cancer

Oxidative stress predominantly triggers the mutation in normal cellular DNA which may lead to the development of cellular proliferation and tumour formation (Aggarwal *et al.*, 2019). Moreover free radical generation may also change the cellular signalling cascade and contribute to the cellular differentiation and inflammation in tumour environment (Aggarwal *et al.*, 2019). The anthocyanins and related natural products may prevent the cancer and tumour development by several ways such as inhibiting oxidative stress mediated DNA mutation, modulates the cellular signalling processes, enhances the phase II reaction enzymes for detoxification, prevents angiogenesis, and inhibits cellular differentiation and proliferation (Lin *et al.*, 2017; Wang and Stoner, 2008). In both *in vitro* and *in vivo* research trials, anthocyanins have shown significantly reduced proliferation of cancer cells (Medic *et al.*, 2019) and inhibited tumour formation (Koide *et al.* 1997; Meiers *et al.* 2001; De-Xing 2003; Kang *et al.* 2003). Comparisons of the antiproliferative effects of anthocyanins on normal as well as cancer cells have surprisingly revealed that they selectively inhibit cancer cell growth while having insignificant effect on normal cell growth (Fakhri *et al.*, 2020; Matsumoto *et al.* 2001; Zhang *et al.* 2005). Moreover, anthocyanidins have a greater potential to inhibit cell proliferation than anthocyanins (Zhang *et al.* 2005; Hudlikar *et al.*, 2020). Key features of anthocyanin mediated anticancer effects are presented in **Table 6**.

The ability of anthocyanin to impede with carcinogenesis seems to be related to numerous potential mechanisms of action that include inhibition of cyclooxygenase enzymes and potent antioxidant potential (Reddy *et al.* 2005a). Previous literature reported that cyanidin 3-*O*-glucoside or peonidin 3-*O*-glucoside administration activated caspase-3, chromatin condensation and initiated cancer cell death (Chen *et al.* 2005). Anthocyanins also have been found to inhibit tumour formation by blocking activation of a mitogen-activated protein kinase pathway (Hou *et al.* 2004). This provides the first indication of a molecular basis for why anthocyanins display anti-carcinogenic properties. Fruit extracts from *Vaccinium* species (low bush blueberry, bilberry, cranberry, and lingonberry) having significant anthocyanin concentrations were effective against various stages of carcinogenesis (Bomser *et al.* 1996; Kandil *et al.* 2000; Kang *et al.* 2003; Smith *et al.* 2008). In addition, Seeram *et al.* (2006) found that extracts of *Vaccinium* species showed significant pro-apoptotic effects against human oral (KB, CAL-27), breast (MCF-7), colon (HT-29, HCT116) and prostate (LNCaP) tumour cell lines. Previous studies also suggested that malvidin inhibited AGS, HCT-116, NCI-H460, MCF-7 and SF-268 cell growth while pelargonidin inhibited AGS, HCT-116, NCI H460, MCF-7 and SF-268 cell growth (Zhang *et al.* 2005).

The anticancer activity of anthocyanins is linked to their phenolic structures. These effects have been verified *in vitro* using several cell culture systems including breast (Singletary *et al.* 2007; Olsson *et al.* 2004), colon (Parry *et al.* 2006; Renis *et al.* 2008), endothelial (Bagchi *et al.* 2004), leukemic (Feng *et al.* 2007), and liver cells (Meyers *et al.* 2003; Shih *et al.* 2007), as well as keratinocytes (Afaq *et al.* 2007). Anthocyanins have shown multiple anti-toxicant and anti-carcinogenic effects in various cell culture systems such as: directly scavenging (ROS), increasing the oxygen-radical absorbing capacity of cells, stimulating the expression of Phase II detoxification enzymes and reducing the formation of oxidative adducts in DNA. Other detoxification processes include decreasing lipid peroxidation,

inhibiting mutagenesis by environmental toxins and carcinogens, and reducing cellular proliferation by modulating signal transduction pathways. Anthocyanins have also been found to function by chelating metals and by direct binding to proteins in their anti-carcinogenic functions (Kong *et al.* 2003). Anthocyanins have also been proven to induce phase II antioxidant and detoxifying enzymes in cultured cells that contribute to its anti-carcinogenic properties (Shih *et al.* 2005).

In addition, apoptosis or programmed cell death plays a major role in the development and regulation of normal cellular function. Anthocyanin-rich extracts from berries (*Vaccinium* species) and grapes (*V. vinifera*) as well as several pure anthocyanidins and anthocyanins, have been found to exhibit pro-apoptotic effects in multiple cell types in *in vitro* studies (Martin *et al.* 2003; Olsson *et al.* 2004; Chen *et al.* 2005; Seeram *et al.* 2006; Afaq *et al.* 2007; Reddivari *et al.* 2007). This is via both intrinsic (mitochondrial) and extrinsic FAS pathways (Reddivari *et al.* 2007; Chang *et al.* 2005). Inflammation, on the other hand, has been shown to have role in the promotion of some cancer types in animals and probably in humans (Kwon *et al.* 2011). Abnormal up-regulation of two inflammatory proteins, NF- κ B and COX-2, is a common phenomenon in many cancers, and their inhibition can result in significant anti-carcinogenic effects (Martin *et al.* 2003; Chang *et al.* 2005). Anthocyanins can inhibit mRNA and/or protein expression levels of COX-2, NF- κ B and other various interleukins, and showed anti-inflammatory effects in multiple cell types *in vitro* (Huang *et al.* 2002; Afaq *et al.* 2005a; Reddy *et al.* 2005b; Rodrigo *et al.* 2006; Boivin *et al.* 2007).

Angiogenesis is the formation of new blood vessels from the existing vascular network and it is an important part of tumour growth and metastasis (Huang *et al.* 2006). Some of the strongest angiogenesis-activating molecules are the vascular endothelial growth factors (VEGF), whose expression is rapidly enhanced in developing tumours (Huang *et al.* 2006). The anti-angiogenic effects of anthocyanins have been demonstrated using cultured endothelial cells (Bagchi *et al.* 2004), oral cancer cells (Rodrigo *et al.* 2006) and mouse epidermal JB6 cells (Huang *et al.* 2006). Anthocyanins in all these cases have suppressed angiogenesis by several mechanisms. Firstly, anthocyanins cause inhibition of H₂O₂ and TNF- α -induced VEGF expression in epidermal keratinocytes (Bagchi *et al.* 2004). Secondly, anthocyanins showed an anti-angiogenesis effect by reducing VEGF and VEGF receptor expression in endothelial cells (Bagchi *et al.* 2004). In addition, mouse epidermal JB6 cells, when treated with an anthocyanin-rich black raspberry (*R. occidentalis*) extract, caused down-regulation of VEGF expression (Huang *et al.* 2006).

Proteolysis, an important and early invasion event, is the degradation of basement membrane collagen (Kelley *et al.* 2014). To degrade the extracellular matrix, barriers secrete proteolytic enzymes for successful invasion of tumour and stromal cells. Basement membrane degradation not only depends on the quantity of proteolytic enzymes present, but also on the balance between activated proteases and their naturally occurring inhibitors. Matrix metalloproteinases (MMP) and plasminogen activators are involved in the regulation of degrading the basement membranes (Brandstetter *et al.* 2001). Anthocyanin extracts from black rice (*O. sativa*), eggplant (*S. melongena*) and different berries have been investigated for their inhibition invasion of multiple cancer cell types. They were found to inhibit invasion of cancer cells by reducing the expression of urokinase-plasminogen activator (u-PA) and MMPs (Brandstetter *et al.* 2001).

Induction of cellular differentiation can be used to prevent and treat cancer through a cell-specific approach that will probably be less toxic than chemo/radiotherapy (Fimognari *et al.* 2004). *In vitro* treatment with anthocyanin (25–200 μ g/mL) in leukemic cells showed the reduction of nitro blue tetrazolium, (a functional marker for granulocyte/monocyte differentiation) and increased adherence of cells to plastic which indicated differentiation of the cells into a monocyte/macrophage-like phenotype (Fimognari *et al.* 2004). Anthocyanins treatment also initiated the naphthol AS-D chloroacetate activity which is a marker for granulocytic differentiation in leukemic cells. Additionally, anthocyanin treatment increased the number of α -naphthyl acetate esterase positive cells which also indicates a differentiation toward a monocytic/macrophagic lineage (Fimognari *et al.* 2004). Moreover, anthocyanins have also been found to induce differentiation in melanoma cells characterized by a significant increase in dendritic outgrowth along with a remodelling of the microtubular network (Serafino *et al.* 2004).

Table 6. Function of anthocyanin rich extracts and anthocyanins in the prevention of cancer.

Food extracts containing anthocyanin	Model	Dose	Experimental Outcomes	Reference
Anthocyanins (plant species particularly in berries and cherries)	Human colon cancer HT29 cells	50, 100, 200, or 400 mg/L of anthocyanins were used against colorectal cancer cell lines.	Anthocyanins promoted apoptosis of colorectal cancer cells (CRC) and inhibited growth of xenografted tumors. Mechanically, the PI3K/AKT/survivin pathway was targeted which enhanced the Bcl-2/Bax and caspase-dependent apoptotic pathways, ultimate result was impairment growth of CRC.	(Zhao et al., 2019)
Pomegranate (<i>P. granatum</i>) fruit extract and pure anthocyanin	Human epidermal keratinocytes, colon cancer cell. Female CD-1 mice were divided into 4 groups	Concentration ranges from 10 to 40 µg/mL 2 mg of extract repeated twice weekly up to the termination of the experiments at 30 weeks	Inhibiting UV-B–induced modulations of NF-κB and MAPK pathways and protecting cells against the adverse effects of UV-B radiation. Inhibiting Akt phosphorylation, COX-2 expression and NFκB DNA binding activity	(Afaq <i>et al.</i> 2005a; Afaq <i>et al.</i> 2005b; Afaq <i>et al.</i> 2007; Sharma <i>et al.</i> 2017)
Juice from strawberry (<i>Fragaria x ananassa</i>), raspberry (<i>R. idaeus</i>), black currant (<i>R. nigrum</i>), red currant (<i>R. rubrum</i>), white currant (<i>R. sativum</i>), gooseberry (<i>R. hirtellum</i>), high-bush blueberry (<i>V. corymbosum</i>), low-bush	Adenocarcinoma cell lines from stomach, mammary gland, prostatic and colorectal gland	Berry juice at 0, 10, 20, 30, 40 or 50 µg/mL.	Markedly inhibited TNF-induced expression of COX-2 and activation of nuclear transcription factor NFκB	(Boivin <i>et al.</i> 2007)

Food extracts containing anthocyanin	Model	Dose	Experimental Outcomes	Reference
blueberry (<i>V. angustifolia</i>), velvet leaf blueberry (<i>V. myrtilloides</i>), serviceberry (<i>Amelanchier sanguinea</i>), blackberry (<i>R. allegheniensis</i>) and sea buckthorn (<i>Hippophae rhamnoides</i>)				
Black raspberries (<i>Rubus occidentalis</i>)	Mouse epidermal cell line, human oral squamous and 30 colorectal 5-weeks-old male C57BL/6J mice (n=10)	Concentrations range from 1 to 100 µg/mL for epidermal cell line. 10 to 100 µg/mL concentration for human oral squamous cell. Orally 7.0 µmol/g/day of anthocyanins for 9 weeks	Inhibits benzo(a)pyrene diol-epoxide-induced activator protein 1 activation, NF-κB factor and VEGF transcription by targeting the phosphatidylinositol 3-Kinase/Akt pathway. Induced both terminal and apoptosis differentiation, suppressed nitric oxide synthase activity.	(Huang <i>et al.</i> 2002; Huang <i>et al.</i> 2006; Rodrigo <i>et al.</i> 2006; Zhang <i>et al.</i> 2018)
Cyanidin-3- <i>O</i> -rutinoside from black raspberry (<i>R. occidentalis</i>)	human leukaemia and lymphoma cell lines	50 to 120 µM or greater	Induced apoptosis by promoting p38 MAP kinase and JNK-mediated Bim phosphorylation.	(Feng <i>et al.</i> 2007)
Apple (<i>Malus domestica</i>), black raspberry (<i>R.</i>	Human keratinocytes, colon cancer cells	Four different concentrations were	Significant chelating capacities against Fe ²⁺ Increased cell oxygen-radical absorbing capacity.	(Bagchi <i>et al.</i> 2004; Long <i>et al.</i>

Food extracts containing anthocyanin	Model	Dose	Experimental Outcomes	Reference
<i>occidentalis</i>), black currant (<i>R. nigrum</i>), black chokeberries (<i>A. melanocarpa</i>), blueberries (<i>V. corymbosum</i>), chardonnay grape (<i>V. vinifera</i>), sea buckthorn (<i>H. rhamnoides</i>), plum (<i>Prunus domestica</i>), lingonberries (<i>V. vitis-idaea</i>), cherries (<i>P. avium</i>), and raspberries (<i>R. idaeus</i>)	HT29, breast cancer cells MCF-7 and thyroid HTh-7	used (0.025, 0.05, 0.25, and 0.5% of plant dry matter, final concentration 0.01-350 µg/mL) for colon cancer cells HT29 and breast cancer cells MCF-7. 10µg/mL for thyroid HTh-7 cells	Significantly inhibited breast cancer cells MCF-7 and colon cancer (HT-29) cell proliferation. Inhibited basal monocyte chemotactic protein-1 and inducible NFκB transcriptions through H ₂ O ₂ and TNFα-induced VEGF expression. Suppressed the activated Akt, mammalian rapamycin, and ribosomal protein S6 via reduced apoptosis and autophagy-dependent cell death.	2018; Olsson <i>et al.</i> 2004; Parry <i>et al.</i> 2006)
Pure cyanidin-3- <i>O</i> -glucopyranoside	Human leukaemia cell line	The concentrations of pure compound used was from 3.1 to 200.0 µg/mL	-Effect on protein kinase C, phosphatidylinositol 3-kinase and also induction of apoptosis and cytodifferentiation and to prevent and treat cancer.	(Fimognari <i>et al.</i> 2004)
Anthocyanins (cyanidin, delphinidin, malvidin, pelargonidin, peonidin and their derivatives)	Human gastric adenocarcinoma, mouse neuroblastoma Neuro-2A cell line and e rat hepatocyte Clone 9 cell line	Tested concentrations ranges from 0–200 µM or greater	-Malvidin treatment significantly increased p38 kinase expression and inhibited the extracellular signal-regulated kinases pathway, including mitogen-activation protein kinases, protein kinase c and phosphatidylinositol 3-kinase. -Stimulated the expression of Phase II detoxification enzymes regulated through phosphorylation by several protein kinases.	(Shih <i>et al.</i> 2005; Shih <i>et al.</i> 2007; Shih <i>et al.</i> 2011)

Food extracts containing anthocyanin	Model	Dose	Experimental Outcomes	Reference
			- Prevent amyloid- β -peptide-mediated neurodysfunction.	
Beetroot (<i>Beta vulgaris</i>)	MCF-7 (breast), HCT-116 (colon), AGS (stomach), CNS (central nervous system), and NCI-H460 (lung) tumour cell lines	Concentration ranges from 10 to 200 μ g/mL	-Anti-toxicant and anti-carcinogenic effects by decreasing lipid peroxidation, inhibition of COX-1 and COX-2 and decreasing of tumour cell growth.	(Reddy <i>et al.</i> 2005b)
Cyanidin-3- <i>O</i> -glucopyranoside and cyanidin chloride	Human colon cancer cells	Concentration ranges from 5 to 200 μ mol/L	-Effect on cell growth, directly scavenging reactive oxygen species (ROS) formation and cell cycle/stress proteins modification, including ataxia teleangectasia mutated protein. -Reducing the formation of oxidative adducts in DNA. Counteracting H ₂ O ₂ -induced DNA damage.	(Renis <i>et al.</i> 2008)
Cultivated strawberries	Liver cancer cells	Concentration ranges from 5 to 75 mg/mL	-Anti-toxicant and anti-carcinogenic effects by increasing the oxygen-radical absorbing capacity of cells and inhibiting cell proliferation.	(Meyers <i>et al.</i> 2003)
Black raspberries (<i>R. occidentalis</i>)	Mouse epidermal JB6 cell	concentrations ranging from 1 to 100 μ g/mL	-Down-regulation of VEGF expression resulting in anti-angiogenic effect.	(Huang <i>et al.</i> 2006)
Black rice (<i>O. sativa</i>)	<i>In vitro</i> enzyme inhibition assay	concentrations ranging from 50 to 200 μ g/mL	-Inhibited invasion of cancer cells by reducing the expression of MMP and urokinase-plasminogen activator (u-PA)	(Chen <i>et al.</i> 2006)

Food extracts containing anthocyanin	Model	Dose	Experimental Outcomes	Reference
Delphinidin	18 Four-week-old male C57BL/6Nhsd mice	15 mg/kg body weight, administered by oral gavage for 16 weeks.	Reduced triglyceride accumulation <i>in vitro</i> through the modulation of lipid metabolic gene expression but no effect on either metabolic alterations or histological abnormalities associated with HFHC diets	(Parry et al. 2018)

Anthocyanins in Visual Activity

Significantly improved visual activity can be achieved through administration of anthocyanins to humans and animals (**Table 7**), and their role has been well documented in improving night vision (Matsumoto *et al.* 2001; Nomi *et al.* 2019). Improvements in night vision adaptation occurred in humans following oral intake of black currant anthocyanosides (Nakaishi *et al.* 2001), and similar outcomes were recorded after administration of anthocyanins from bilberries (Muth *et al.* 2000). Regeneration of rhodopsin (a G-protein-coupled receptor localized in the retina of the eye) was stimulated by three anthocyanins from black currant and formation of regeneration intermediate was accelerated by cyanidin 3-*O*-rutinoside (Matsumoto *et al.* 2003). Anthocyanin-rich bilberry (*V. myrtillus*) extract prevented inflammation of endotoxin-induced uveitis using a mouse model (Miyake *et al.* 2012). A bilberry (*V. myrtillus*) extract rich in anthocyanins prevented the impairment of photoreceptor cell function, ameliorated the intracellular elevation of ROS and inhibited NF- κ B (Miyake *et al.* 2012). Thus, enhancement of rhodopsin regeneration has been proven to be one of the mechanisms by which anthocyanins improve visual activity.

A positive effect of anthocyanins on vision improvement was suggested by early clinical trials carried out in France and Italy (Rouher *et al.* 1972; Jayle *et al.* 1965). A controlled clinical trial of cyanoside chloride and heleniène on 31 patients suffering from functional disturbances of vision in the dark, reported that both compounds significantly improved photopic visual activity (Sole *et al.* 1984). In a German clinical trial, Difrarel®E (anthocyanosides and vitamin E) was given to 36 patients with progressive myopia for about 14 months. In around 50% of patients an increase in myopia was suppressed by approximately 50%, along with 29 patients showing stabilisation of fundus-alterations, and an overall improved and stable visual acuity was obtained (Politzer 1977). Anthocyanins from blackcurrant (*R. nigrum*) in the form of a concentrated extract powder were examined for their effects on asthenopia, an ophthalmological condition with nonspecific symptoms such as fatigue, red eyes, eye strain, pain in or around the eyes, blurred vision, headache and occasional double vision, and is a result of continuous exposure of the eye to video displays (Nakaishi *et al.* 2001). Oral administration of blackcurrant (*R. nigrum*) anthocyanins to such individuals at various doses decreased the dark adaptation threshold in a dose dependent fashion (Nakaishi *et al.* 2001). The photooxidation of pyridinium disretinoid A2E, an auto-fluorescence pigment accumulating in retinal pigment epithelial cells with age and also associated with some retinal disorders, was also found to be suppressed by scavenging singlet oxygen, upon administration of nine anthocyanins from bilberry (*V. myrtillus*) extracts (Jang *et al.* 2005). Black soybean (*G. max*) seed anthocyanins protected retinal neurons from *N*-methyl-*N*-nitrosourea-induced functional and structural damage (Paik *et al.* 2012). Anthocyanin from black soybean (*G. max*) protected human lens epithelial (HLE-B3) cells in H₂O₂-induced oxidative stress and prevented apoptosis (Mok *et al.* 2014).

The ocular distribution of blackcurrant (*R. nigrum*) anthocyanins in rats and rabbits after oral, intravenous and intraperitoneal administration was investigated (Matsumoto *et al.* 2006). Blackcurrant (*R. nigrum*) anthocyanins (BCA) were absorbed and distributed in ocular tissues as intact forms and passed through the blood-aqueous and blood-retinal barriers in both species. This study revealed that oral intake of anthocyanins or anthocyanin-rich extracts can be used to treat ophthalmological diseases such as glaucoma and myopia. Investigation of the effect of purified

high-dose anthocyanin oligomers on nocturnal visual function and clinical symptoms were carried out on 60 patients with asthenopia and refractive errors in both eyes. About 73% of patients were reported with improved symptoms (Lee *et al.* 2007). A randomized, placebo-controlled, double-masked trial showed that blackcurrant (*R. nigrum*) anthocyanins increased ocular blood flows during the two year trial in comparison with placebo-treated patients (Ohguro *et al.* 2012). However, significant changes were not observed in ocular or systemic conditions, including intraocular pressure, during the experiment (Ohguro *et al.* 2012). Another study suggests that bilberry (*V. myrtillus*) anthocyanins improved visual function in patients with normal tension glaucoma (Shim *et al.* 2012).

Flavonoids have a probable role in the prevention of diabetic cataracts (Varma and Kinoshita 1976). Flavonoids have also been found to delay or prevent the occurrence of cataracts in rat lenses perfused with a high-glucose solution or in diabetic rabbits (Varma and Kinoshita 1976; Laurens *et al.* 1985). Many naturally extracted or synthesized anthocyanin combinations with novel anti-cataract or anti-glaucoma activity have been reported in patents (Dilip and Tetsuya 2007). Reduction in the control of lens opacity was related to high levels of reducing sugars in *in vitro* and *in vivo* studies, i.e. the formation of experimental diabetic cataracts, which is a complication of diabetes occurring in about 10% of diabetic patients. It has been proven that caloric and food intake greatly influences the progression of diabetes and its complications (Thiraphatthanavong *et al.* 2014a). Inhibitory activities for lens opacity were shown by five anthocyanin monomers isolated from grape skin extract (Thiraphatthanavong *et al.* 2014a). Medium to high doses of *Z. mays* (purple waxy corn) extract, an important source of anthocyanins, reduced lens opacity along with a decreased MDA level, followed by improved experimental diabetic cataract in enucleated rat lenses (Thiraphatthanavong *et al.* 2014b).

Table 7. Function of anthocyanin rich extracts and anthocyanins in the improvement and protection of visual activity.

Food extracts containing anthocyanin	Model	Dose	Experimental Outcomes	Reference
Bilberry (<i>V. myrtillus</i>)	Human adult retinal pigment epithelial cells	100 μ M bilberry extract	-The photooxidation of pyridinium disretinoid A2E was suppressed by scavenging singlet oxygen.	(Jang <i>et al.</i> 2005)
Black currants (<i>R. nigrum</i>)	Human subjects with asthenopia. Thirty-five male Wistar rats and 16 Japan White male rabbits 6 groups (five groups consisted of 3 rabbits and the control group of one rabbit)	12.5, 20, and 50 mg/subject. Rats intraperitoneally received 500 mg blackcurrant anthocyanins powder per kg body weight (108 mg anthocyanins per kg body weight). rabbits received 92.6 mg/kg body weight of blackcurrant anthocyanins (anthocyanins 20 mg/ kg body weight) solution intravenously via an ear vein	-Significantly improved night vision -BCAs were absorbed and distributed in ocular tissues as intact forms and passed through the blood-aqueous and blood-retinal barriers, indicating that can used to treat myopia and glaucoma.	(Nakaishi <i>et al.</i> 2001) (Matsumoto <i>et al.</i> 2006; Lee <i>et al.</i> 2007)
Cyanoside chloride and heleniene	Human subjects with functional disturbances of vision in the dark. Human subjects with progressive myopia.	31 patients were treated with 200 mg of cyanoside chloride and heleniene thrice a day for at least 4 weeks.	-Significantly improved photopic visual acuity. -Cyaninoside chloride only improved visual functions related to mesopic and scotopic vision.	(Sole <i>et al.</i> 1984)

Toxicological aspects of anthocyanins

This review work showed that regular intake of anthocyanins rich colourful fruits and vegetables play an important role to maintain a healthy lifestyle which may protect against chronic diseases such as CVS, diabetes and cancer. Though no deficiency disorder has been reported which are associated with the lack of the consumption of anthocyanins (Wallace and Giusti, 2015). However, toxicological effects of anthocyanins and related food products need to be assessed scientifically to support the claims about its potential non-toxic effect. The daily intake of anthocyanins, established by the FAO/WHO expert committee, is 2.5 mg/kg per day, extracted from grape-skin (Wallace and Giusti, 2015). However, no recommended intake level of general anthocyanins for optimal health benefit or to avoid adverse effects is published yet. Low or no acute toxicity symptoms were reported for anthocyanin consumption, both *in vitro* and *in vivo* (animals and humans) models at usual dietary intake levels (Burton-Freeman et al., 2016; Harlinda et al., 2016; Khoo et al., 2017; Pojer et al., 2013). Moreover, a study on guinea pigs receiving 3 g/day of anthocyanins extract for 15 days followed by a washout period of one month reported no adverse effect developed in the animals (European Food Safety Authority, 2013). Toxicity testing report on *Lepisanthes alata* (Blume) *Leenh* fruits which are enriched with anthocyanins suggested that the fruits of the plant tested are safe for consumption (Anggraini et al., 2019). Furthermore, no negative effects developed due to the administration of very high doses of anthocyanin derivatives have been reported (Eker et al., 2019). In addition, the consumption of anthocyanins also increased the gut microbiota such as *Bifidobacterium spp.*, or *Lactobacillus spp.* which showed benefit in metabolic dysfunction (Eker et al., 2019).

Conclusion and future direction

The prevention of disease and wellbeing of health is an important aspect for a healthy life in modern society. Since the beginning of human history, nature has been considered as a provider of food and natural healing substances in the form of herbal plants. Even in modern medicine, drug development largely depends on the ancient knowledge of healing. Public health policy strongly encourages preventative health measures so as to reduce dependence on costly treatments. In this regard, anthocyanin serves as ‘potential drug’, with robust protective efficacy in several lifestyles related disorders. The majority of such attributes are due to its anti-inflammatory and antioxidant activities. There are many studies currently being conducted to further evaluate the health beneficial effects of this extraordinary pigment, as researchers, nowadays, are focusing on extracting health benefits from functional foods. Apart from direct antioxidant activity, a wide range of complex mechanisms have been proposed to explain most of the health promoting mechanisms of anthocyanins. Anthocyanin covers a large number of targets which include oxidants, xenobiotics, and excess of metals, radiations and pro-inflammatory factors. Furthermore, anthocyanin can interact directly with proteins and enzymes, which results into modulation in signalling pathways and related changes in cellular metabolism.

Overall, *in vitro* and *in vivo* results and clinical evidence indicates that anthocyanins could be a promising therapeutic agent against an extended range of diseases and disorders, which includes cardiovascular diseases, neurological impairments, diabetes, viral diseases, cancer, toxin-induced liver and kidney damage, inflammation and oxidative

radical-induced pathologies. While anthocyanins may have a low bioavailability, research indicates that the metabolites may be responsible for much of their health promoting properties. Available reports on cellular and animal models have provided sufficient clinical outcomes, which could influence researchers to study further and establish the anthocyanin based lead molecule for the benefits of mankind. The present study could assist in the very same process by providing a gross notion on the molecular mechanisms of anthocyanin action. However, lack of adequate knowledge about actual molecular mechanisms hinders the application of anthocyanin as a therapeutic drug for many lifestyle disorders. More studies are required to find the real activities of anthocyanins and their metabolites and the specific mechanisms that health promoting properties of anthocyanins are realised. We can hope to see anthocyanins contributing significantly in the manufacture of therapeutics in the near future and facilitate people to benefit from the gifts of nature rather than artificial products.

Acknowledgement

L. Nahar gratefully acknowledges the financial support of the European Regional Development Fund - Project ENOCH (No. CZ.02.1.01/0.0/0.0/16_019/0000868).

Conflict of Interest

Authors declared that there is no conflict of interest regarding this manuscript.

Author Contributions:

MAA, SDS and NS planned for the review. MAA, PI, MMR and NS made the literature collection and preliminary draft manuscript. FK, GEB and LN critically expanded the draft manuscript and checked the correctness of the references. MAA, NS and SDS made the tables and pictorial diagram. MAA, NS, GEB and SDS critically reviewed the whole manuscript. All author approved the final version of this manuscript.

References

- Abo El-Ella DM, Bishayee A (2019) Chapter 6 - The epigenetic targets of berry anthocyanins in cancer prevention. In: Bishayee A, Bhatia D (eds) *Epigenetics of Cancer Prevention*, vol 8. Academic Press, pp 129-148.
- Afaq F, Malik A, Syed D, Maes D, Matsui MS, Mukhtar H (2005a) Pomegranate fruit extract modulates UV-B-mediated phosphorylation of mitogen-activated protein kinases and activation of nuclear factor kappa B in normal human epidermal keratinocytes paragraph sign. *Photochem. Photobiol.* 81:38-45.
- Afaq F, Saleem M, Krueger CG, Reed JD, Mukhtar H (2005b) Anthocyanin- and hydrolyzable tannin-rich pomegranate fruit extract modulates MAPK and NF-kappaB pathways and inhibits skin tumorigenesis in CD-1 mice. *Int. J. Cancer.* 113:423-433.

- Afaq F, Syed DN, Malik A, Hadi N, Sarfaraz S, Kweon MH, Khan N, Zaid MA, Mukhtar H (2007) Delphinidin, an anthocyanidin in pigmented fruits and vegetables, protects human haccat keratinocytes and mouse skin against UVB-mediated oxidative stress and apoptosis. *J. Investig. Dermatol.* 127:222-232.
- Aggarwal V, Tuli HS, Varol A, Thakral F, Yerer MB, Sak K, Varol M, Jain A, Khan MA, Sethi G (2019) Role of reactive oxygen species in cancer progression: Molecular mechanisms and recent advancements. *Biomolecules* 9.
- Ahmadiani N, Sigurdson GT, Robbins RJ, Collins TM, Giusti MM (2019) Solid phase fractionation techniques for segregation of red cabbage anthocyanins with different colorimetric and stability properties. *Food Res. Int.* 120:688-696.
- Ahmet I, Spangler E, Shukitt-Hale B, Joseph JA, Ingram DK, Talan M (2009) Survival and cardioprotective benefits of long-term blueberry enriched diet in dilated cardiomyopathy following myocardial infarction in rats. *PLoS ONE.* 4:e7975.
- Akkarachiyasit S, Charoenlertkul P, Yibchok-Anun S, Adisakwattana S (2010) Inhibitory activities of cyanidin and its glycosides and synergistic effect with acarbose against intestinal alpha-glucosidase and pancreatic alpha-amylase. *Int. J. Mol. Sci.* 11:3387-3396.
- Alam MA (2019) Anti-hypertensive effect of cereal antioxidant ferulic acid and its mechanism of action. *Front Nutr.* 6, 121.
- Alam MA, Subhan N, Hossain H, Hossain M, Reza HM, Rahman MM, Ullah MO (2016) Hydroxycinnamic acid derivatives: a potential class of natural compounds for the management of lipid metabolism and obesity. *Nutr. Metab.* 13: 27.
- Al-Awwadi NA, Araiz C, Bornet A, Delbosc S, Cristol J-P, Linck N, Azay J, Teissedre P-L, Cros G (2005) Extracts enriched in different polyphenolic families normalize increased cardiac NADPH oxidase expression while having differential effects on insulin resistance, hypertension, and cardiac hypertrophy in high-fructose-fed rats. *J. Agric. Food Chem.* 53:151-157.
- Aloud BM, Raj P, McCallum J, Kirby C, Louis XL, Jahan F, Yu L, Hiebert B, Duhamel TA, Wigle JT *et al.* (2018) Cyanidin 3-*O*-glucoside prevents the development of maladaptive cardiac hypertrophy and diastolic heart dysfunction in 20-week-old spontaneously hypertensive rats. *Food Funct* 9:3466-3480.
- Anderson RA, Broadhurst CL, Polansky MM, Schmidt WF, Khan A, Flanagan VP, Schoene NW, *et al.* (2004) Isolation and characterization of polyphenol type-a polymers from cinnamon with insulin-like biological activity. *J. Agric. Food Chem.* 52:65-70.
- Anderson RA, Polansky MM (2002) Tea enhances insulin activity. *J. Agric. Food Chem.* 50:7182-7186.
- Andriambeloson E, Magnier C, Haan-Archipoff G, Lobstein A, Anton R, Beretz A, Stoclet JC, Andriantsitohaina R (1998) Natural dietary polyphenolic compounds cause endothelium-dependent vasorelaxation in rat thoracic aorta. *J. Nutr.* 128:2324-2333.
- Anggraini T, Wilma S, Syukri D, Azima F (2019) Total phenolic, anthocyanin, catechins, DPPH radical scavenging activity, and toxicity of *Lepisanthes alata* (Blume) *Leenh.* *Int. J. Food Sci.* 2019, 9703176.

- Aoki H, Kuze N, Kato Y (2002) Anthocyanins isolated from purple corn (*Zea mays* L.). *Foods and Food Ingredients J. Jpn.* 41-45.
- Aqil F, Gupta A, Munagala R, Jeyabalan J, Kausar H, Sharma RJ, Singh IP, Gupta RC (2012) Antioxidant and antiproliferative activities of anthocyanin/ellagitannin-enriched extracts from *Syzygium cumini* L. (Jamun, the Indian Blackberry). *Nutr. Cancer.* 64:428-438.
- Arevstrom L, Bergh C, Landberg R, Wu H, Rodriguez-Mateos A, Waldenborg M, Magnuson A, Blanc S, Frobert O (2019) Freeze-dried bilberry (*Vaccinium myrtillus*) dietary supplement improves walking distance and lipids after myocardial infarction: an open-label randomized clinical trial. *Nutr. Res.* 62:13-22.
- Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K, Shimomura I, Nakamura T, Miyaoka K *et al.* (2012) Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem. Biophys. Res. Commun.* 425:560-564.
- Atalay M, Gordillo G, Roy S, Rovin B, Bagchi D, Bagchi M, Sen CK (2003) Anti-angiogenic property of edible berry in a model of hemangioma. *FEBS Lett.* 544:252-257.
- Augusti KT, Daniel RS, Cherian S, Sheela CG, Nair CR (1994) Effect of leucopelargonin derivative from *Ficus bengalensis* Linn. on diabetic dogs. *Indian J. Med. Res.* 99:82-86.
- Aviram M, Rosenblat M, Gaitini D, Nitecki S, Hoffman A, Dornfeld L, Volkova N, Presser D, Attias J, Liker H *et al.* (2004) Pomegranate juice consumption for 3 years by patients with carotid artery stenosis reduces common carotid intima-media thickness, blood pressure and LDL oxidation. *Clin. Nutr.* 23:423-433.
- Azzini E, Vitaglione P, Intorre F, Napolitano A, Durazzo A, Foddai MS, Fumagalli A, Catasta G, Rossi L, Venneria E *et al.* (2010) Bioavailability of strawberry antioxidants in human subjects. *Br. J. Nutr.* 104:1165-1173.
- Badshah H, Kim TH, Kim MO (2015) Protective effects of anthocyanins against amyloid beta-induced neurotoxicity *in vivo* and *in vitro*. *Neurochem. Int.* 80:51-59.
- Bagchi D, Sen CK, Bagchi M, Atalay M (2004) Anti-angiogenic, antioxidant, and anti-carcinogenic properties of a novel anthocyanin-rich berry extract formula. *Biochemistry (Moscow)* 69:75-80.
- Bakuradze T, Tausend A, Galan J, Maria-Groh IA, Berry D, Tur JA, Marko D, Richling E (2019) Antioxidative activity and health benefits of anthocyanin-rich fruit juice in healthy volunteers. *Free Radic. Res.* 53 sup 1 1045-1055.
- Barnes MJ, Perry BG, Hurst RD, Lomiwes D (2020) Anthocyanin-rich new zealand blackcurrant extract supports the maintenance of forearm blood-flow during prolonged sedentary sitting. *Front.Nutr.* 74.
- Barton M, Cosentino F, Brandes RP, Moreau P, Shaw S, Luscher TF (1997) Anatomic heterogeneity of vascular aging: role of nitric oxide and endothelin. *Hypertension.* 30:817-824.
- Basu A, Rhone M, Lyons TJ (2010) Berries: emerging impact on cardiovascular health. *Nutr. Rev.* 68:168-177.
- Bhatt S, Nagappa AN, Patil CR (2020) Role of oxidative stress in depression. *Drug Discov. Today* 25, 1270-1276.
- Bell DR, Gochenaur K (2006) Direct vasoactive and vasoprotective properties of anthocyanin-rich extracts. *J. Appl. Physiol.* 100:1164-1170.
- Bell L, Williams CM (2019) A pilot dose–response study of the acute effects of haskap berry extract (*Lonicera caerulea* L.) on cognition, mood, and blood pressure in older adults. *Eur. J. Nutr.* 58:3325-3334.

- Bernatoniene J, Kopustinskiene DM (2018) The role of catechins in cellular responses to oxidative stress. *Molecules* 23, 965.
- Bertuglia S, Malandrino S, Colantuoni A (1995) Effects of the natural flavonoid delphinidin on diabetic microangiopathy. *Arzneimittel-Forsch* 45:481-485.
- Bhaswant M, Shafie SR, Mathai ML, Mouatt P, Brown L (2017) Anthocyanins in chokeberry and purple maize attenuate diet-induced metabolic syndrome in rats. *Nutr.* 41:24-31.
- Blesso CN (2019) Dietary anthocyanins and human health. *Nutrients*. 11.
- Boivin D, Blanchette M, Barrette S, Moghrabi A, Béliveau R (2007) Inhibition of cancer cell proliferation and suppression of TNF-induced activation of NF- κ B by edible berry juice. *Anticancer Res.* 27:937-948.
- Bomser J, Madhavi D, Singletary K, Smith M (1996) *In vitro* anticancer activity of fruit extracts from *Vaccinium* species. *Planta Medica*. 62:212-216.
- Bonesi M, Leporini M, Tenuta MC, Tundis R (2020) The Role of anthocyanins in drug discovery: Recent developments. *Curr. Drug Discov. Technol.* 17: 286-298.
- Borkowski T, Szymusiak H, Gliszczynska-Świgło A, Rietjens IMCM, Tyrakowska B (2005) Radical scavenging capacity of wine anthocyanins is strongly pH-dependent. *J. Agric. Food Chem.* 53:5526-5534.
- Bouayed J, Bohn T (2010) Exogenous antioxidants--double-edged swords in cellular redox state: health beneficial effects at physiologic doses versus deleterious effects at high doses. *Oxid. Med. Cell. Longev.* 3, 228-237.
- Brandstetter H, Grams F, Glitz D, Lang A, Huber R, Bode W, Krell H-W, Engh RA (2001) The 1.8-Å Crystal structure of a matrix metalloproteinase 8-barbiturate inhibitor complex reveals a previously unobserved mechanism for collagenase substrate recognition. *J. Biol. Chem.* 276:17405-17412.
- Broncel M, Kozirog M, Duchnowicz P, Koter-Michalak M, Sikora J, Chojnowska-Jezierska J (2010) *Aronia melanocarpa* extract reduces blood pressure, serum endothelin, lipid, and oxidative stress marker levels in patients with metabolic syndrome. *Med Sci Monit.* 16:CR28-34.
- Burton-Freeman B, Brzezinski M, Park E, Sandhu A, Xiao D, Edirisinghe I (2019) A selective role of dietary anthocyanins and flavan-3-ols in reducing the risk of type 2 diabetes mellitus: A review of recent evidence. *Nutr.* 11:841.
- Burton-Freeman B, Sandhu A, Edirisinghe I (2016) Chapter 35 - Anthocyanins, in: Gupta, R.C. (Ed.), *Nutraceuticals*. Academic Press, Boston, pp. 489-500.
- Carluccio Maria A, Siculella L, Ancora Maria A, Massaro M, Scoditti E, Storelli C, Visioli F, Distanto A, De Caterina R (2003) Olive oil and red wine antioxidant polyphenols inhibit endothelial activation. *Arterioscler. Thromb. Vasc. Biol.* 23:622-629.
- Chang J-J, Hsu M-J, Huang H-P, Chung D-J, Chang Y-C, Wang C-J (2013) Mulberry anthocyanins inhibit oleic acid induced lipid accumulation by reduction of lipogenesis and promotion of hepatic lipid clearance. *J. Agric. Food Chem.* 61:6069-6076.
- Chang Y-C, Huang H-P, Hsu J-D, Yang S-F, Wang C-J (2005) *Hibiscus* anthocyanins rich extract-induced apoptotic cell death in human promyelocytic leukemia cells. *Toxicol. Appl. Pharmacol.* 205:201-212.

- Chen L, Deng H, Cui H, Fang J, Zuo Z, Deng J, Li Y, Wang X, Zhao L (2017) Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget* 9:7204-7218.
- Chen L, Jiang B, Zhong C, Guo J, Zhang L, Mu T, Zhang Q, Bi X (2018) Chemoprevention of colorectal cancer by black raspberry anthocyanins involved the modulation of gut microbiota and SFRP2 demethylation. *Carcinogenesis* 39:471-481.
- Chen PN, Chu SC, Chiou HL, Chiang CL, Yang SF, Hsieh YS (2005a) Cyanidin 3-*O*-glucoside and peonidin 3-glucoside inhibit tumor cell growth and induce apoptosis in vitro and suppress tumor growth *in vivo*. *Nutr. Cancer* 53:232-243.
- Chen PN, Kuo WH, Chiang CL, Chiou HL, Hsieh YS, Chu SC (2006) Black rice anthocyanins inhibit cancer cells invasion via repressions of MMPs and u-PA expression. *Chem. Biol. Interact.* 163(3): 218-229.
- Chen QM, Maltagliati AJ (2018) Nrf2 at the heart of oxidative stress and cardiac protection. *Physiol. Genomics.* 50, 77-97.
- Chen S, Zhou H, Zhang G, Meng J, Deng K, Zhou W, Wang H, Wang Z, Hu N, Suo Y (2019) Anthocyanins from *Lycium ruthenicum* Murr. ameliorated d-galactose-induced memory impairment, oxidative stress, and neuroinflammation in adult rats. *J. Agric. Food Chem.* 67:3140-3149.
- Chen Y-F, Shibu M, Fan M, Chen M-C, Viswanadha V, Lin Y-L, Lai C-H, Lin K-H, Ho T-J, Kuo W-W *et al.* (2016) Purple rice anthocyanin extract protects cardiac function in STZ-induced diabetes rat hearts by inhibiting cardiac hypertrophy and fibrosis. *J. Nutr. Biochem.* 31:98-105.
- Cherian S, Kumar RV, Augusti KT, Kidwai JR (1992) Antidiabetic effect of a glycoside of pelargonidin isolated from the bark of *Ficus bengalensis* Linn. *Indian J. Biochem. Bio.* 29:380-382.
- Chiu L-W, Zhou X, Burke S, Wu X, Prior RL, Li L (2010) The purple cauliflower arises from activation of a MYB transcription factor. *Plant Physiol.* 154:1470-1480.
- Cho J, Kang JS, Long PH, Jing J, Back Y, Chung KS (2003) Antioxidant and memory enhancing effects of purple sweet potato anthocyanin and cordyceps mushroom extract. *Arch. Pharm. Res.* 26:821-825
- Chun OK, Chung S-J, Claycombe KJ, Song WO (2008) Serum C-reactive protein concentrations are inversely associated with dietary flavonoid intake in U.S. adults. *J. Nutr.* 138:753-760.
- Cines DB, Pollak ES, Buck CA, Loscalzo J, Zimmerman GA, Pober JS, Wick TM, Konkle BA, Schwartz BS *et al.* (1998) Endothelial cells in physiology and in the pathophysiology of vascular disorders. *Blood* 91:3527.
- Cohen-Boulakia F, Valensi PE, Boulahdour H, Lestrade R, Dufour-Lamartinie J-F, Hort-Legrand C, Behar A (2000) *In vivo* sequential study of skeletal muscle capillary permeability in diabetic rats: effect of anthocyanosides. *Metab.* 49:880-885.
- Cook MD, Sandu BAK, Joyce PJP (2020) Effect of New Zealand blackcurrant on blood pressure, cognitive function and functional performance in older adults. *J. Nutr. Gerontol. Geriatr.* 39:99-113.
- Cooke D, Schwarz M, Boocock D, Winterhalter P, Steward WP, Gescher AJ, Marczyklo TH (2006) Effect of cyanidin-3-*O*-glucoside and an anthocyanin mixture from bilberry on adenoma development in the ApcMin mouse model of intestinal carcinogenesis—Relationship with tissue anthocyanin levels. *Int. J. Cancer* 119:2213-2220.

- Curtis PJ, van der Velpen V, Berends L, Jennings A, Feelisch M, Umpleby AM, Evans M, Fernandez BO, Meiss MS, Minnion M *et al.* (2019) Blueberries improve biomarkers of cardiometabolic function in participants with metabolic syndrome—results from a 6-month, double-blind, randomized controlled trial. *AM J. Clin. Nutr.* 109:1535-1545.
- da Silva RFR, Barreira JCM, Heleno SA, Barros L, Calhelha RC, Ferreira ICFR (2019) Anthocyanin profile of elderberry juice: a natural-based bioactive colouring ingredient with potential food application. *Molecules.* 24:2359.
- Daniel RS, Devi KS, Augusti KT, Sudhakaran Nair CR (2003) Mechanism of action of antiatherogenic and related effects of *Ficus bengalensis* Linn. flavonoids in experimental animals. *Indian J. of Exp. Biol.* 41:296-303
- Daveri E, Cremonini E, Mastaloudis A, Hester SN, Wood SM, Waterhouse AL, Anderson M, Fraga CG, Oteiza PI (2018) Cyanidin and delphinidin modulate inflammation and altered redox signaling improving insulin resistance in high fat-fed mice. *Redox Biol.* 18:16-24.
- De-Xing H (2003) Potential mechanisms of cancer chemoprevention by anthocyanins. *Curr. Mol. Med.* 3:149-159.
- DeFuria J, Bennett G, Strissel KJ, Perfield JW, 2nd, Milbury PE, Greenberg AS, Obin MS (2009) Dietary blueberry attenuates whole-body insulin resistance in high fat-fed mice by reducing adipocyte death and its inflammatory sequelae. *J. Nutr.* 139:1510-1516.
- Demrow HS, Slane PR, Folts JD (1995) Administration of wine and grape juice inhibits in vivo platelet activity and thrombosis in stenosed canine coronary arteries. *Circulation.* 91:1182-1188.
- Di Castelnuovo A, Rotondo S, Iacoviello L, Donati MB, De Gaetano G (2002) Meta-analysis of wine and beer consumption in relation to vascular risk. *Circulation.* 105:2836-2844.
- Di Meo S, Reed TT, Venditti P, Victor VM (2016) Role of ROS and RNS sources in physiological and pathological conditions. *Oxid. Med. Cell. Longev.* 2016, 1245049-1245049.
- Dilip G, Tetsuya K (2007) Anthocyanins and anthocyanin-rich extracts: role in diabetes and eye function. *Asia Pac. J. Clin. Nutr.* 16:200-208.
- Dohadwala MM, Vita JA (2009) Grapes and cardiovascular disease. *J. Nutr.* 139:1788-1793.
- Eker ME, Aaby K, Budic-Leto I, Brncic SR, El SN, Karakaya S, Simsek S, Manach C, Wiczowski W, Pascual-Teresa S (2019) A Review of factors affecting anthocyanin bioavailability: Possible implications for the inter-individual variability. *Foods (Basel, Switzerland)* 9.
- Ellulu MS, Patimah I, Khaza'ai H, Rahmat A, Abed Y (2017) Obesity and inflammation: the linking mechanism and the complications. *Arch. Med. Sci.* 13:851-863.
- European Food Safety Authority (2013) Scientific opinion on the re-evaluation of anthocyanins (E 163) as a food additive, EFSA panel on food additives and nutrient sources added to Food (ANS). *EFSA J.* 11, 3145.
- Fakhri S, Khodamorady M, Naseri M, Farzaei MH, Khan H (2020) The ameliorating effects of anthocyanins on the cross-linked signaling pathways of cancer dysregulated metabolism. *Pharmacol. Res.* 159: 104895.
- Fallah AA, Sarmast E, Jafari T (2020) Effect of dietary anthocyanins on biomarkers of oxidative stress and antioxidative capacity: A systematic review and meta-analysis of randomized controlled trials. *J. Funct. Foods.* 68, 103912.

- Fan D, Alamri Y, Liu K, MacAskill M, Harris P, Brimble M, Dalrymple-Alford J, Prickett T, Menzies O, Laurenson A *et al.* (2018) Supplementation of blackcurrant anthocyanins increased cyclic glycine-proline in the cerebrospinal fluid of parkinson patients: potential treatment to improve insulin-like growth factor-1 function. *Nutr.* 10(6):714.
- Fan M, Choi Y-J, Tang Y, Bae SM, Yang HP, Kim E-K (2019) Efficacy and mechanism of polymerized anthocyanin from grape-skin extract on high-fat-diet-induced nonalcoholic fatty liver disease. *Nutr.* 11:2586.
- Farrell NJ, Norris GH, Ryan J, Porter CM, Jiang C, Blesso CN (2015) Black elderberry extract attenuates inflammation and metabolic dysfunction in diet-induced obese mice. *Br. J. Nutr.* 114:1123-1131.
- Felgines C, Talavera S, Gonthier M-P, Texier O, Scalbert A, Lamaison J-L (2003) Strawberry anthocyanins are recovered in urine as glucuro- and sulfoconjugates in humans. *J. Nutr.* 133:1296–1301.
- Feng R, Ni HM, Wang SY, Tourkova IL, Shurin MR, Harada H, Yin XM (2007) Cyanidin-3-*O*-rutinoside, a natural polyphenol antioxidant, selectively kills leukemic cells by induction of oxidative stress. *J. Biol. Chem.* 282:13468-13476.
- Fimognari C, Berti F, Nüsse M, Cantelli-Forti G, Hrelia P (2004) Induction of apoptosis in two human leukemia cell lines as well as differentiation in human promyelocytic cells by cyanidin-3-*O*- β -glucopyranoside. *Biochem. Pharmacol.* 67:2047-2056.
- Fitzpatrick DF, Hirschfield SL, Coffey RG (1993) Endothelium-dependent vasorelaxing activity of wine and other grape products. *AM J. Physiol-Heart C.* 265:H774-778.
- Fitzpatrick DF, Hirschfield SL, Ricci T, Jantzen P, Coffey RG (1995) Endothelium-dependent vasorelaxation caused by various plant extracts. *J. Cardiovasc. Pharmacol.* 26:90-95
- Folts JD (1998) Antithrombotic potential of grape juice and red wine for preventing heart attacks. *Pharm. Biol.* 36:21-27
- Fossen T, Andersen ØM (2003) Anthocyanins from red onion, *Allium cepa*, with novel aglycone. *Phytochem.* 62:1217-1220.
- Frank J, Kamal-Eldin A, Lundh T, Määtä K, Törrönen R, Vessby B (2002) Effects of dietary anthocyanins on tocopherols and lipids in rats. *J. Agric. Food Chem.* 50:7226-7230.
- Freedman JE, Parker C, Li L, Perlman JA, Frei B, Ivanov V, Deak LR, Iafra MD, Folts JD (2001) Select flavonoids and whole juice from purple grapes inhibit platelet function and enhance nitric oxide release. *Circulation.* 103:2792-2798.
- Fronad AD, Iuhas CI, Stirbu I, Leopold L, Socaci S, Andreea S, Ayvaz H, Andreea S, Mihai S, Diaconeasa Z, Carmen S (2019) Phytochemical characterization of five edible purple-reddish vegetables: anthocyanins, flavonoids, and phenolic acid derivatives. *Molecules.* 24:1536.
- Gabor M (1972) Pharmacologic effects of flavonoids on blood vessels. *Angiologica* 9:355-374.
- Gabrielska J, Oszmiański J (2005) Antioxidant activity of anthocyanin glycoside derivatives evaluated by the inhibition of liposome oxidation. *Zeitschrift für Naturforschung.* 60: 399-407.

- Garcia-Sanchez A, Miranda-Diaz AG, Cardona-Munoz EG (2020) The role of oxidative stress in physiopathology and pharmacological treatment with pro- and antioxidant properties in chronic diseases. *Oxid. Med. Cell. Longev.* 2020, 2082145.
- Ge Q, Ma X (2013) Composition and antioxidant activity of anthocyanins isolated from Yunnan edible rose (An ning). *Food Sci. and Human Wellness* 2:68-74.
- Gharib A, Faezizadeh Z, Godarzee M (2013) Treatment of diabetes in the mouse model by delphinidin and cyanidin hydrochloride in free and liposomal forms. *Planta Med*, 79(17): 1599-1604.
- Giampieri F, Tulipani S, Alvarez-Suarez JM, Quiles JL, Mezzetti B, Battino M (2012) The strawberry: composition, nutritional quality, and impact on human health. *Nutr.* 28:9-19.
- Gil MI, Tomas-Barberan FA, Hess-Pierce B, Holcroft DM, Kader AA (2000) Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J. Agric. Food Chem.* 48:4581-4589.
- Goncalves AC, Bento C, Silva BM, Silva LR (2017) Sweet cherries from Fundao possess antidiabetic potential and protect human erythrocytes against oxidative damage. *Food Res. Int.* 95:91-100.
- Gorinstein S, Caspi A, Libman I, Lerner HT, Huang D, Leontowicz H, Leontowicz M, Tashma Z, Katrich E, Feng S *et al.* (2006) Red grapefruit positively influences serum triglyceride level in patients suffering from coronary atherosclerosis: studies in vitro and in humans. *J. Agric. Food Chem.* 54:1887-1892.
- Graf D, Seifert S, Jaudszus A, Bub A, Watzl B (2013) Anthocyanin-rich juice lowers serum cholesterol, leptin, and resistin and improves plasma fatty acid composition in fischer rats. *PLoS ONE.* 8:e66690.
- Griffiths LA, Smith GE (1972) Metabolism of myricetin and related compounds in the rat. Metabolite formation *in vivo* and by the intestinal microflora *in vitro*. *Biochem. J.* 130:141-151.
- Grimes KL, Stuart CM, McCarthy JJ, Kaur B, Cantu EJ, Forester SC (2018) Enhancing the cancer cell growth inhibitory effects of table grape anthocyanins. *J.Food Sci.* 83:2369-2374.
- Guo Y, Zhang P, Liu Y, Zha L, Ling W, Guo H (2020) A dose-response evaluation of purified anthocyanins on inflammatory and oxidative biomarkers and metabolic risk factors in healthy young adults: A randomized controlled trial. *Nutr.* 74:110745.
- Harborne JB, Grayer RJ (1988) The anthocyanins. In: Harborne JB (ed) *The Flavonoids*. Springer, Boston. pp 1-20.
- Harborne JB, Williams CA (2001) Anthocyanins and other flavonoids. *Nat. Prod. Rep.* 18:310-333.
- Harlinda K, Anindya MR, Agmi SP, Rawan WK (2016) Antioxidant and toxicity properties of anthocyanin extracted from red flower of four tropical shrubs. *Nusantara Bioscience* 8, 135-140.
- He J, Magnuson BA, Lala G, Tian Q, Schwartz SJ, Giusti MM (2006) Intact anthocyanins and metabolites in rat urine and plasma after 3 months of anthocyanin supplementation. *Nutr. Cancer.* 54:3-12.
- Hidalgo J, Teuber S, Morera FJ, Ojeda C, Flores CA, Hidalgo MA, Nunez L, Villalobos C, Burgos, RA (2017) Delphinidin reduces glucose uptake in mice jejunal tissue and human intestinal cells lines through FFA1/GPR40. *Int. J. Mol. Sci.* 18(4).
- Horie K, Nanashima N, Maeda H (2019) Phytoestrogenic effects of blackcurrant anthocyanins increased endothelial nitric oxide synthase (eNOS) expression in human endothelial cells and ovariectomized rats. *Molecules* 24:1259.

- Hou D-X, Kai K, Li J-J, Lin S, Terahara N, Wakamatsu M, Fujii M, Young MR, Colburn N (2004) Anthocyanidins inhibit activator protein 1 activity and cell transformation: structure–activity relationship and molecular mechanisms. *Carcinogenesis*. 25:29-36.
- Hu W, Wang W, Ma Q, Liu T, Zhang J, Zhang J (2020) Blueberry anthocyanin-enriched extract ameliorates transverse aortic constriction-induced myocardial dysfunction via the DDAH1/ADMA/NO signaling pathway in mice. *Mol. Med. Rep.* 21:454-462.
- Hu T, Dong Y, He C, Zhao M, He Q (2020) The gut microbiota and oxidative stress in autism spectrum disorders (ASD). *Oxid. Med. Cell. Longev.* 2020, 8396708.
- Huang C, Huang Y, Li J, Hu W, Aziz R, Tang M-S, Sun N, Cassady J, Stoner GD (2002) Inhibition of benzo(a)pyrene diol-epoxide-induced transactivation of activated protein 1 and nuclear factor- κ B by black raspberry extracts. *Cancer Res.* 62:6857-6863
- Huang C, Li J, Song L, Zhang D, Tong Q, Ding M, Bowman L, Aziz R, Stoner GD (2006) Black raspberry extracts inhibit benzo(a)pyrene diol-epoxide–induced activator protein 1 activation and VEGF transcription by targeting the phosphatidylinositol 3-kinase/Akt pathway. *Cancer Res.* 66:581-587.
- Huang H, Jiang X, Xiao Z, Yu L, Pham Q, Sun J, Chen P, Yokoyama W, Yu LL, Luo YS, Wang TT (2016a) Red cabbage microgreens lower circulating low-density lipoprotein (LDL), liver cholesterol, and inflammatory cytokines in mice fed a high-fat diet. *J. Agric. Food Chem.* 64:9161-9171.
- Huang W, Yan Z, Li D, Ma Y, Zhou J, Sui Z (2018) Antioxidant and anti-inflammatory effects of blueberry anthocyanins on high glucose-induced human retinal capillary endothelial cells. *Oxid. Med. Cell. Longev.* 2018:10.
- Huang W, Zhu Y, Li C, Sui Z, Min W (2016b) Effect of blueberry anthocyanins malvidin and glycosides on the antioxidant properties in endothelial cells. *Oxid. Med. Cell. Longev.* 2016:10.
- Huang WY, Wang J, Liu YM, Zheng QS, Li CY (2014) Inhibitory effect of malvidin on TNF- α -induced inflammatory response in endothelial cells. *Eur. J. Pharmacol.* 723:67-72.
- Hudlikar R, Wu R, Cheng D, Kuo DH-C, Wang L, Peter R, Yin R, Li S, Kong A-N (2020) Anthocyanins and cancer prevention, in: Pezzuto JM, Vang O (Eds.), *Natural products for cancer chemoprevention: Single compounds and combinations*. Springer International Publishing, Cham, pp. 351-373.
- Hwang J-W, Kim E-K, Lee S-J, Kim Y-S, Moon S-H, Jeon B-T, Sung S-H, Kim E-T, Park P-J (2012) Antioxidant activity and protective effect of anthocyanin oligomers on H₂O₂-triggered G2/M arrest in retinal cells. *J. Agric. Food Chem.* 60:4282-4288.
- Ibrahim UK, Muhammad II, Salleh RM (2011) The effect of pH on color behavior of *Brassica oleracea* anthocyanin. *J. App. Sci.* 11:2406-2410.
- Ighodaro OM, Akinloye OA (2018) First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria J. Med.* 54, 287-293.
- Isaak CK, Petkau JC, Blewett H, O K, Siow YL (2017) Lingonberry anthocyanins protect cardiac cells from oxidative-stress-induced apoptosis. *Can. J. Physiol. Pharm.* 95:904-910.

- Jang YP, Zhou J, Nakanishi K, Sparrow JR (2005) Anthocyanins protect against A2E photooxidation and membrane permeabilization in retinal pigment epithelial cells. *Photochem. Photobiol.* 81:529-536.
- Jayaprakasam B, Olson LK, Schutzki RE, Tai M-H, Nair MG (2006) Amelioration of obesity and glucose intolerance in high-fat-fed C57BL/6 mice by anthocyanins and ursolic acid in cornelian cherry (*Cornus mas*). *J. Agric. Food Chem.* 54:243-248.
- Jayaprakasam B, Vareed SK, Olson LK, Nair MG (2005) Insulin secretion by bioactive anthocyanins and anthocyanidins present in fruits. *J. Agric. Food Chem.* 53:28-31.
- Jayle GE, Aubry M, Gavini H, Braccini G, De la Baume C (1965) Study concerning the action of anthocyanoside extracts of *Vaccinium myrtillus* on night vision. *Annales d'oculistique.* 198:556-562.
- Jennings A, Welch AA, Fairweather-Tait SJ, Kay C, Minihane AM, Chowienczyk P, Jiang B, Cecelja M, Spector T, Macgregor A, Cassidy A (2012) Higher anthocyanin intake is associated with lower arterial stiffness and central blood pressure in women. *AM J Clin Nutr.* 96:781-788.
- Jeong JW, Lee WS, Shin SC, Kim GY, Choi BT, Choi YH (2013) Anthocyanins downregulate lipopolysaccharide-induced inflammatory responses in BV2 microglial cells by suppressing the NF-kappaB and Akt/MAPKs signaling pathways. *Int. J. Mol. Sci.* 14:1502-1515.
- Jiang Y, Dai M, Nie W-J, Yang X-R, Zeng X-C (2017) Effects of the ethanol extract of black mulberry (*Morus nigra* L.) fruit on experimental atherosclerosis in rats. *J. Ethnopharmacol.* 200:228-235.
- Joo HK, Choi S, Lee YR, Lee EO, Park MS, Park KB, Kim C-S, Lim YP, Park J-T, Jeon BH (2018) Anthocyanin-rich extract from red chinese cabbage alleviates vascular inflammation in endothelial cells and Apo E^{-/-} mice. *Int. J. Mol. Sci.* 19:816.
- Joseph JA, Shukitt-Hale B, Denisova NA, Bielinski D, Martin A, McEwen JJ, Bickford PC (1999) Reversals of age-related declines in neuronal signal transduction, cognitive, and motor behavioral deficits with blueberry, spinach, or strawberry dietary supplementation. *J. Neurosci.* 19:8114-8121.
- Ju JH, Yoon HS, Park HJ, Kim MY, Shin HK, Park KY, Yang JO, Sohn MS, Do MS *et al.* (2011) Anti-obesity and antioxidative effects of purple sweet potato extract in 3T3-L1 adipocytes *in vitro*. *J. Med. Food.* 14:1097-1106.
- Kahkonen MP, Heinonen M (2003) Antioxidant activity of anthocyanins and their aglycons. *J. Agric. Food Chem.* 51:628-633.
- Kalt W, Hanneken A, Milbury P, Tremblay F (2010) Recent research on polyphenolics in vision and eye health. *J. Agric. Food Chem.* 58:4001-4007.
- Kandil FE, Song L, Pezzuto JM, Marley K, Seigler DS, Smith MAL (2000) Isolation of oligomeric proanthocyanidins from flavonoid-producing cell cultures. *In Vitro Cell Dev-Pl.* 36:492-500.
- Kang S-Y, Seeram NP, Nair MG, Bourquin LD (2003) Tart cherry anthocyanins inhibit tumor development in Apc (Min) mice and reduce proliferation of human colon cancer cells. *Cancer Lett.* 194:13-19.
- Kattoor AJ, Pothineni NVK, Palagiri D, Mehta JL (2017) Oxidative stress in atherosclerosis. *Curr. Atheroscler. Rep.* 19:42.

- Kay CD, Mazza G (2008) Bioactivity, absorption, and metabolism of anthocyanins. In: Daayf F, Lattanzio V (eds) Recent Advances in Polyphenol Research, vol 1. Wiley, pp 228-262
- Kay CD, Mazza G, Holub BJ (2005) Anthocyanins exist in the circulation primarily as metabolites in adult men. *J. Nutr.* 135:2582-2588.
- Kay CD, Mazza G, Holub BJ, Wang J (2007) Anthocyanin metabolites in human urine and serum. *Br. J. Nutr.* 91:933-942.
- Kelley LC, Lohmer LL, Hagedorn EJ, Sherwood DR (2014) Traversing the basement membrane *in vivo*: a diversity of strategies. *J. Cell. Biol.* 204:291-302.
- Keppler K, Humpf H-U (2005) Metabolism of anthocyanins and their phenolic degradation products by the intestinal microflora. *Bioorg. Med. Chem.* 13:5195-5205.
- Khan MS, Ali T, Kim MW, Jo MH, Chung JI, Kim MO (2019) Anthocyanins improve hippocampus-dependent memory function and prevent neurodegeneration via JNK/Akt/GSK3 β signaling in LPS-treated adult mice. *Mol. Neurobiol.* 56:671-687.
- Khoo HE, Azlan A, Tang ST, Lim SM (2017) Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food Nutr. Res.* 61:1361779.
- Kianbakht S, Hashem-Dabaghian F (2019) Antihypertensive efficacy and safety of *Vaccinium arctostaphylos* berry extract in overweight/obese hypertensive patients: A randomized, double-blind and placebo-controlled clinical trial. *Complement. Ther. Med.* 44:296-300.
- Kim SY, Wi H-R, Choi S, Ha TJ, Lee BW, Lee M (2015) Inhibitory effect of anthocyanin-rich black soybean testa (*Glycine max* (L.) Merr.) on the inflammation-induced adipogenesis in a DIO mouse model. *J. Funct. Foods.* 14:623-633.
- Kim YK, Yoon HH, Lee YD, Youn DY, Ha TJ, Kim HS, Lee JH (2012) Anthocyanin extracts from black soybean (*Glycine max* L.) protect human glial cells against oxygen-glucose deprivation by promoting autophagy. *Biomol. Ther.* 20:68-74.
- Kirisattayakul W, Wattanathorn J, Iamsaard S, Jittiwat J, Suriharn B, Lertrat K (2017) Neuroprotective and memory-enhancing effect of the combined extract of purple waxy corn cob and pandan in ovariectomized rats. *Oxid. Med. Cell. Longev.* 2017:12.
- Klatsky AL (2001) Commentary: Could abstinence from alcohol be hazardous to your health? *Int. J. Epidemiol.* 30:739-742.
- Koide T, Hashimoto Y, Kamei H, Kojima T, Hasegawa M, Terabe K (1997) Antitumor effect of anthocyanin fractions extracted from red soybeans and red beans *in vitro* and *in vivo*. *Cancer Biother Radio.* 12:277-280.
- Kong J-M, Chia L-S, Goh N-K, Chia T-F, Brouillard R (2003) Analysis and biological activities of anthocyanins. *Phytochem.* 64:923-933.
- Kuciel-Lewandowska J, Kasperczyk M, Bogut B, Heider R, Laber WT, Laber W, Paprocka-Borowicz M (2020) The impact of health resort treatment on the nonenzymatic endogenous antioxidant system. *Oxid. Med. Cell. Longev.* 2020, 8423105.

- Kulling ES, Rawel MH (2008) Chokeberry (*Aronia melanocarpa*) – a review on the characteristic components and potential health effects. *Planta Medica*. 74:1625-1634.
- Kurutas EB (2016) The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state. *Nutr. J.* 15, 71-71.
- Kusano S, Abe H, Tamura H (2001) Isolation of antidiabetic components from white-skinned sweet potato (*Ipomoea batatas* L.) *Biosci., Biotechnol. Biochem.* 65:109-114.
- Kwon Bc, Javed W, Elmqvist N, Yi JS (2011) Direct manipulation through surrogate objects. Paper presented at the Proceedings of the SIGCHI Conference on Human Factors in Computing Systems, Vancouver, BC, Canada.
- Lachin T, Reza H (2012) Anti-diabetic effect of cherries in alloxan induced diabetic rats. *Recent Pat. Endocr. Metab. Immune. Drug Discov.* 6:67-72.
- Lampe JW, Chang J-L (2007) Interindividual differences in phytochemical metabolism and disposition. *Seminars in Cancer Biol.* 17:347-353.
- Landrault N, Poucheret P, Azay J, Krosniak M, Gasc F, Jenin C, Cros G, Teissedre P-L (2003) Effect of a polyphenols-enriched chardonnay white wine in diabetic rats. *J. Agric. Food Chem.* 51:311-318.
- Laurens A, Gionobarber P, Sylla O, Gionobarber H (1985) Aldose reductase Inhibitors-A new therapeutic method for certain complications of diabetes. *Therapie.* 38:659-663.
- Lee C, Han D, Kim B, Baek N, Baik B-K (2013) Antioxidant and anti-hypertensive activity of anthocyanin-rich extracts from hullless pigmented barley cultivars. *Int. J. Food Sci. Technol.* 48:984-991.
- Lee J, Lee HK, Kim CY, Hong YJ, Choe CM, You TW, Seong GJ (2007) Purified high-dose anthocyanoside oligomer administration improves nocturnal vision and clinical symptoms in myopia subjects. *Br. J. Nutr.* 93:895-899.
- Lee M, Sorn SR, Park Y, Park HK (2016) Anthocyanin rich-black soybean testa improved visceral fat and plasma lipid profiles in overweight/obese Korean adults: A randomized controlled trial. *J. Med. Food.* 19:995-1003.
- Li J, Zhao R, Jiang Y, Xu Y, Zhao H, Lyu X, Wu T (2020) Bilberry anthocyanins improve neuroinflammation and cognitive dysfunction in APP/PSEN1 mice via the CD33/TREM2/TYROBP signaling pathway in microglia. *Food Funct.* 11:1572-1584.
- Li S, Wu B, Fu W, Reddivari L (2019) The anti-inflammatory effects of dietary anthocyanins against ulcerative colitis. *Int. J. Mol. Sci.* 20:2588.
- Liang Y-J, Shyu K-G, Wang B-W, Lai L-P (2006) C-reactive protein activates the nuclear factor- κ B pathway and induces vascular cell adhesion molecule-1 expression through CD32 in human umbilical vein endothelial cells and aortic endothelial cells. *J.Mol. Cell. Cardiol.* 40:412-420.
- Lietti A, Cristoni A, Picci M (1976) Studies on *Vaccinium myrtillus* anthocyanosides. I. Vasoprotective and antiinflammatory activity. *Arzneimittel-forschung.* 26:829-832.
- Lim S, Xu J, Kim J, Chen T.-Y, Su X, Standard J, Carey E, Griffin J, Herndon B, Katz B, Tomich J *et al.* (2013) Role of anthocyanin-enriched purple-fleshed sweet potato p40 in colorectal cancer prevention *Mol. Nutr. Food Res.* 57:1908-1917.
- Lin B-W, Gong C-C, Song H-F, Cui Y-Y (2017) Effects of anthocyanins on the prevention and treatment of cancer. *Br. J. Pharmacol.* 174, 1226-1243.

- Liu Y-Z, Wang Y-X, Jiang C-L (2017) Inflammation: the common pathway of stress-related diseases. *Frontiers in Human Neurosci.* 11: 316.
- Lobo V, Patil A, Phatak A, Chandra N (2010) Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn. Rev.* 4:118-126.
- Long H-L, Zhang F-F, Wang H-L, Yang W-S, Hou H-T, Yu J-K, Liu B (2018) Mulberry anthocyanins improves thyroid cancer progression mainly by inducing apoptosis and autophagy cell death. *Kaohsiung J. Med. Sci.* 34:255-262.
- Luna-Vital DA, Gonzalez de Mejia E (2018) Anthocyanins from purple corn activate free fatty acid-receptor 1 and glucokinase enhancing *in vitro* insulin secretion and hepatic glucose uptake. *PLoS ONE.* 13:e0200449.
- Luo Y, Fang J-L, Yuan K, Jin S-H, Guo Y (2019) Ameliorative effect of purified anthocyanin from *Lycium ruthenicum* on atherosclerosis in rats through synergistic modulation of the gut microbiota and NF- κ B/SREBP-2 pathways. *J. Funct. Foods.* 59:223-233.
- Ma Q (2013) Role of nrf2 in oxidative stress and toxicity. *Annu. Rev. Pharmacol. Toxicol.* 53, 401-426.
- Madhavi DL, Bomser J, Smith MAL, Singletary K (1998) Isolation of bioactive constituents from *Vaccinium myrtillus* (bilberry) fruits and cell cultures. *Plant Sci.* 131:95-103.
- Marco PH, Poppi RJ, Scarminio IS, Tauler R (2011) Investigation of the pH effect and UV radiation on kinetic degradation of anthocyanin mixtures extracted from *Hibiscus acetosella*. *Food Chem.* 125:1020-1027.
- Martin S, Giannone G, Andriantsitohaina R, Martinez MC (2003) Delphinidin, an active compound of red wine, inhibits endothelial cell apoptosis via nitric oxide pathway and regulation of calcium homeostasis. *Br. J. Pharmacol.* 139:1095-1102.
- Matsui T, Ebuchi S, Fukui K, Matsugano K, Terahara N, Matsumoto K (2004) Caffeoylsophorose, a new natural alpha-glucosidase inhibitor, from red vinegar by fermented purple-fleshed sweet potato. *Biosci. Biotech. Bioch.* 68:2239-2246.
- Matsui T, Ueda T, Oki T, Sugita K, Terahara N, Matsumoto K (2001) α -Glucosidase inhibitory action of natural acylated anthocyanins. 1. Survey of natural pigments with potent inhibitory activity. *J. Agric. Food Chem.* 49:1948-1951.
- Matsumoto H, Inaba H, Kishi M, Tominaga S, Hirayama M, Tsuda T (2001) Orally administered delphinidin 3-rutinoside and cyanidin 3-rutinoside are directly absorbed in rats and humans and appear in the blood as the intact forms. *J. Agric. Food Chem.* 49:1546-1551.
- Matsumoto H, Nakamura Y, Iida H, Ito K, Ohguro H (2006) Comparative assessment of distribution of blackcurrant anthocyanins in rabbit and rat ocular tissues. *Exp. Eye Res.* 83:348-356.
- Matsumoto H, Nakamura Y, Tachibanaki S, Kawamura S, Hirayama M (2003) Stimulatory effect of cyanidin 3-glycosides on the regeneration of rhodopsin. *J. Agric. Food Chem.* 51:3560-3563.
- Mauray A, Felgines C, Morand C, Mazur A, Scalbert A, Milenkovic D (2012) Bilberry anthocyanin-rich extract alters expression of genes related to atherosclerosis development in aorta of apo E-deficient mice. *Nutr. Metab. Cardiovas.* 22:72-80.
- Mazza G, Francis F (1995) Anthocyanins in grapes and grape products. *Critical Rev. in Food Sci. Nutr.* 35:341-371.

- Mazza G, Kay CD, Cottrell T, Holub BJ (2002) Absorption of anthocyanins from blueberries and serum antioxidant status in human subjects. *J. Agric. Food Chem.* 50:7731–7737.
- Mcghie T, Walton CM (2019) The bioavailability and absorption of anthocyanins: Towards a better understanding. *Mol. Nutr. Food Res.* 51:702-713.
- McNamara RK, Kalt W, Shidler MD, McDonald J, Summer SS, Stein AL, Stover AN, Krikorian R (2018) Cognitive response to fish oil, blueberry, and combined supplementation in older adults with subjective cognitive impairment. *Neurobiol Aging* 64: 147-156.
- Medic N, Tramer F, Passamonti S (2019) Anthocyanins in colorectal cancer prevention. A systematic review of the literature in search of molecular oncotargets. *Front. Pharmacol.* 10 (675).
- Meiers S, Kemény M, Weyand U, Gastpar R, von Angerer E, Marko D (2001) The anthocyanidins cyanidin and delphinidin are potent inhibitors of the epidermal growth-factor receptor. *J. Agric. Food Chem.* 49:958-962.
- Meyers KJ, Watkins CB, Pritts MP, Liu RH (2003) Antioxidant and antiproliferative activities of strawberries. *J. Agric. Food Chem.* 51:6887-6892.
- Milbury PE, Vita JA, Blumberg JB (2010) Anthocyanins are bioavailable in humans following an acute dose of cranberry juice. *J. Nutr.* 140:1099–1104.
- Miller MG, Hamilton DA, Joseph JA, Shukitt-Hale B (2018) Dietary blueberry improves cognition among older adults in a randomized, double-blind, placebo-controlled trial. *Euro. J. Nutr.* 57: 1169-1180.
- Min J, Yu S-W, Baek S-H, Nair KM, Bae O-N, Bhatt A, Kassab M, Nair MG, Majid A (2011) Neuroprotective effect of cyanidin-3-O-glucoside anthocyanin in mice with focal cerebral ischemia. *Neurosci. Lett.* 500:157-161.
- Mink PJ, Scrafford CG, Barraj LM, Harnack L, Hong C-P, Nettleton JA, Jacobs JDR (2007) Flavonoid intake and cardiovascular disease mortality: a prospective study in postmenopausal women. *AM J Clin Nutr.* 85:895-909.
- Miyake S, Takahashi N, Sasaki M, Kobayashi S, Tsubota K, Ozawa Y (2012) Vision preservation during retinal inflammation by anthocyanin-rich bilberry extract: cellular and molecular mechanism. *Lab. Invest.* 92:102-109.
- Mok JW, Chang DJ, Joo CK (2014) Antiapoptotic effects of anthocyanin from the seed coat of black soybean against oxidative damage of human lens epithelial cell induced by H₂O₂. *Curr. Eye Res.* 39:1090-1098.
- Montilla EC, Arzaba MR, Hillebrand S, Winterhalter P (2011) Anthocyanin composition of black carrot (*Daucus carota* ssp. *sativus* var. *atrorubens* Alef.) cultivars antonina, beta sweet, deep purple, and purple haze J. *Agric. Food Chem.* 59:3385-3390.
- Mulabagal V, Lang GA, DeWitt DL, Dalavoy SS, Nair MG (2009) Anthocyanin content, lipid peroxidation and cyclooxygenase enzyme inhibitory activities of sweet and sour cherries. *J. Agric. Food Chem.* 57:1239-1246.
- Munzel T, Camici GG, Maack C, Bonetti NR, Fuster V, Kovacic JC (2017) Impact of oxidative stress on the heart and vasculature: Part 2 of a 3-Part Series. *J. Am. Coll. Cardiol.* 70, 212-229.
- Muth ER, Laurent JM, Jasper P (2000) The effect of bilberry nutritional supplementation on night visual acuity and contrast sensitivity. *Altern. Med. Rev.* 5:164-173.

- Mykkänen OT, Huotari A, Herzig K-H, Dunlop TW, Mykkänen H, Kirjavainen PV (2014) Wild blueberries (*Vaccinium myrtillus*) alleviate inflammation and hypertension associated with developing obesity in mice fed with a high-fat diet. PLoS ONE. 9:e114790.
- Nahar, L and Sarker, S. D. (2019) Chemistry for Pharmacy Students: General, Organic and Natural Products Chemistry, 2nd Edition, Wiley & Sons, UK
- Nakaishi H, Matsumoto H, Tominaga S, Hirayama M (2001) Effects of black currant anthocyanoside intake on dark adaptation and VDT work-induced transient refractive alteration in healthy humans. Altern. Med. Rev. 6:553-562.
- Nakamura Y, Matsumoto H, Todoki K (2002) Endothelium-dependent vasorelaxation induced by black currant concentrate in rat thoracic aorta. Jpn J. Pharmacol. 89:29-35.
- Nanashima N, Horie K, Yamanouchi K, Tomisawa T, Kitajima M, Oey I, Maeda H (2020) Blackcurrant (*Ribes nigrum*) extract prevents dyslipidemia and hepatic steatosis in ovariectomized rats. Nutr. 12:1541.
- Naruszewicz M, Laniewska I, Millo B, Dłużniewski M (2007) Combination therapy of statin with flavonoids rich extract from chokeberry fruits enhanced reduction in cardiovascular risk markers in patients after myocardial infarction (MI). Atherosclerosis. 194:e179-e184.
- Nemes A, Homoki JR, Kiss R, Hegedus C, Kovacs D, Peitl B, Gal F, Stundl L, Szilvassy Z, Remenyik J (2019) Effect of anthocyanin-rich tart cherry extract on inflammatory mediators and adipokines involved in type 2 diabetes in a high fat diet induced obesity mouse model. Nutr. 11:1966.
- Nile SH, Park SW (2014) Edible berries: Bioactive components and their effect on human health. Nutr. 30:134-144.
- Nomi Y, Iwasaki-Kurashige K, Matsumoto H (2019) Therapeutic effects of anthocyanins for vision and eye Health. Molecules. 24: 3311.
- Nurfardilla SA, Saputri FC, Harahap Y (2019) Effects of *Hibiscus sabdariffa* calyces aqueous extract on the antihypertensive potency of captopril in the two-kidney-one-clip rat hypertension model. Evid. Based Complementary Altern. Med. 2019:9694212.
- Ohguro H, Ohguro I, Katai M, Tanaka S (2012) Two-year randomized, placebo-controlled study of black currant anthocyanins on visual field in glaucoma. Ophthalmologica. 228:26-35.
- Ojeda D, Jimenez-Ferrer E, Zamilpa A, Herrera-Arellano A, Tortoriello J, Alvarez L (2010) Inhibition of angiotensin convertin enzyme (ACE) activity by the anthocyanins delphinidin- and cyanidin-3-O-sambubiosides from *Hibiscus sabdariffa*. J. Ethnopharmacol. 127:7-10.
- Olsson ME, Gustavsson K-E, Andersson S, Nilsson Å, Duan R-D (2004) Inhibition of cancer cell proliferation in vitro by fruit and berry extracts and correlations with antioxidant levels. J. Agric. Food Chem. 52:7264-7271.
- Overall J, Bonney SA, Wilson M, Beermann A III, Grace MH, Esposito D, Lila MA, Komarnytsky S (2017) Metabolic effects of berries with structurally diverse anthocyanins. Int. J. Mol. Sci. 18:422.
- Paik SS, Jeong E, Jung SW, Ha TJ, Kang S, Sim S, Jeon JH, Chun MH, Kim IB, (2012) Anthocyanins from the seed coat of black soybean reduce retinal degeneration induced by N-methyl-N-nitrosourea. Exp. Eye Res. 97:55-62.

- Paixao J, Dinis TC, Almeida LM (2012) Malvidin-3-*O*-glucoside protects endothelial cells up-regulating endothelial NO synthase and inhibiting peroxynitrite-induced NF- κ B activation. *Chem. Biol. Interact.* 199:192-200.
- Pan Z, Cui M, Dai G, Yuan T, Li Y, Ji T, Pan Y (2018) Protective effect of anthocyanin on neurovascular unit in cerebral ischemia/reperfusion injury in rats. *Front. Neurosci.* 12: 947.
- Pandey KB, Rizvi SI (2009) Plant polyphenols as dietary antioxidants in human health and disease. *Oxid. Med. Cell. Longev.* 2:270-278.
- Pang L, Lian X, Liu H, Zhang Y, Li Q, Cai Y, Ma H, Yu X (2020) Understanding diabetic neuropathy: Focus on oxidative stress. *Oxid. Med. Cell. Longev.* 2020, 9524635.
- Parichatikanond W, Pinthong D, Mangmool S (2012) Blockade of the renin-angiotensin system with delphinidin, cyanin, and quercetin. *Planta Med.* 78:1626-1632.
- Parmar NS, Ghosh MN (1977) Protective effect of some bioflavonoids on the X-irradiation-induced increase in capillary permeability of rat intestine. *Indian J. Exp. Biol.* 15:311-313.
- Parra-Vargas M, Sandoval-Rodriguez A, Rodriguez-Echevarria R, Dominguez-Rosales JA, Santos-Garcia A, Armendariz-Borunda J (2018) Delphinidin ameliorates hepatic triglyceride accumulation in human hepg2 cells, but not in diet-induced obese mice. *Nutr.* 10 (8):1060.
- Parry J, Su L, Moore J, Cheng Z, Luther M, Rao JN, Wang J-Y, Yu, Liangli L (2006) Chemical compositions, antioxidant capacities, and antiproliferative activities of selected fruit seed flours. *J. Agric. Food Chem.* 54:3773-3778.
- Passeri V, Koes R, Quattrocchio FM (2016) New challenges for the design of high value plant products: stabilization of anthocyanins in plant vacuoles. *Front. Plant Sci.* 7: 153.
- Pereira SR, Pereira R, Figueiredo I, Freitas V, Dinis TCP, Almeida LM (2017) Comparison of anti-inflammatory activities of an anthocyanin-rich fraction from Portuguese blueberries (*Vaccinium corymbosum* L.) and 5-aminosalicylic acid in a TNBS-induced colitis rat model. *PLoS ONE.* 12:e0174116.
- Pergola C, Rossi A, Dugo P, Cuzzocrea S, Sautebin L (2006) Inhibition of nitric oxide biosynthesis by anthocyanin fraction of blackberry extract. *Nitric Oxide.* 15:30-39.
- Pojer E, Mattivi F, Johnson D, Stockley CS, (2013) The Case for Anthocyanin consumption to promote human health: A review. *Compr. Rev. Food Sci. Food Saf.* 12, 483-508.
- Politzer M (1977) Experiences in the medical treatment of progressive myopia (author's transl). *Klin. Monbl. Augenheilkd.* 171:616-619.
- Pospisil P, Prasad A, Rac M (2019) Mechanism of the formation of electronically excited species by oxidative metabolic processes: Role of reactive oxygen species. *Biomolecules* 9, 258.
- Poulose SM, Fisher DR, Larson J, Bielinski DF, Rimando AM, Carey AN, Schauss AG, Shukitt-Hale B (2012) Anthocyanin-rich Acai (*Euterpe oleracea* Mart.) fruit pulp fractions attenuate inflammatory stress signaling in mouse brain bv-2 microglial cells. *J. Agric. Food Chem.* 60:1084-1093.
- Prior RL, Wu X (2006) Anthocyanins: Structural characteristics that result in unique metabolic patterns and biological activities. *Free Radic. Res.* 40:1014–1028.
- Probst Y (2015) A review of the nutrient composition of selected *Rubus* berries. *Nutr. Food Sci.* 45:242-254.

- Putta S, Yarla NS, Kumar KE, Lakkappa DB, Kamal MA, Scotti L, Scotti MT, Ashraf GM, Rao, BSB, Kumari DS et. al. (2018) Preventive and therapeutic potentials of anthocyanins in diabetes and associated complications. *Curr. Med. Chem.* 25: 5347-5371.
- Qin L, Zhang J, Qin M (2013) Protective effect of cyanidin 3-*O*-glucoside on beta-amyloid peptide-induced cognitive impairment in rats. *Neurosci. Lett.* 534:285-288.
- Qin S, Sun D, Mu J, Ma D, Tang R, Zheng Y (2019) Purple sweet potato color improves hippocampal insulin resistance via down-regulating SOCS3 and galectin-3 in high-fat diet mice. *Behav. Brain Res.* 359:370-377.
- Ramirez-Tortosa C *et al.* (2001) Anthocyanin-rich extract decreases indices of lipid peroxidation and DNA damage in vitamin E-depleted rats. *Free Radic. Biol. Med.* 31:1033-1037.
- Rasheed Z, Akhtar N, Anbazhagan AN, Ramamurthy S, Shukla M, Haqqi TM (2009) Polyphenol-rich pomegranate fruit extract (POMx) suppresses PMACI-induced expression of pro-inflammatory cytokines by inhibiting the activation of MAP Kinases and NF-kappaB in human KU812 cells. *J. Inflamm.* 6:1.
- Reddivari L, Vanamala J, Chintharlapalli S, Safe SH, Miller JJC (2007) Anthocyanin fraction from potato extracts is cytotoxic to prostate cancer cells through activation of caspase-dependent and caspase-independent pathways. *Carcinogenesis.* 28:2227-2235.
- Reddy MK, Alexander-Lindo RL, Nair MG (2005a) Relative inhibition of lipid peroxidation, cyclooxygenase enzymes, and human tumor cell proliferation by natural food colors. *J. Agric. Food Chem.* 53:9268-9273.
- Reddy MK, Alexander-Lindo RL, Nair MG (2005b) Relative inhibition of lipid peroxidation, cyclooxygenase enzymes, and human tumor cell proliferation by natural food colors. *J. Agric. Food Chem.* 53:9268-9273.
- Rehman SU, Shah SA, Ali T, Chung JI, Kim MO (2017) Anthocyanins reversed d-galactose-induced oxidative stress and neuroinflammation mediated cognitive impairment in adult rats. *Mol. Neurobiol.* 54:255-271.
- Renis M, Calandra L, Scifo C, Tomasello B, Cardile V, Vanella L, Bei R, Fauci LL, Galvano F (2008) Response of cell cycle/stress-related protein expression and DNA damage upon treatment of Caco₂ cells with anthocyanins. *Br. J. Nutr.* 100:27-35.
- Ridzwan N, Jumli MN, Baig AA, Rohin MAK (2020) Pomegranate-derived anthocyanin regulates MORs-cAMP/CREB-BDNF pathways in opioid-dependent models and improves cognitive impairments. *J. Ayurveda. Integr. Med.* (in press). doi:<https://doi.org/10.1016/j.jaim.2019.12.001>.
- Rimm EB, Giovannucci EL, Willett WC, Colditz GA, Ascherio A, Rosner B, Stampfer MJ (1991) Prospective study of alcohol consumption and risk of coronary disease in men. *The Lancet* 338:464-468.
- Rodrigo KA, Rawal Y, Renner RJ, Schwartz SJ, Tian Q, Larsen PE, Mallery SR (2006) Suppression of the tumorigenic phenotype in human oral squamous cell carcinoma cells by an ethanol extract derived from freeze-dried black raspberries. *Nutr. Cancer.* 54:58-68.
- Rodriguez-Mateos A, Istas G, Boschek L, Feliciano RP, Mills CE, Boby C, Gomez-Alonso S, Milenkovic D, Heiss C (2019) Circulating anthocyanin metabolites mediate vascular benefits of blueberries: insights from randomized controlled trials, metabolomics, and nutrigenomics. *J. Gerontol A-Biol A.* 74:967-976.
- Rosenblat M, Hayek T, Aviram M (2006) Anti-oxidative effects of pomegranate juice (PJ) consumption by diabetic patients on serum and on macrophages. *Atherosclerosis.* 187:363-371.

- Rouher F, Sole P, Offret G, Bernard JA (1972) Demonstration by adaptoelectroretinography of the action of a cyanidin on nocturnal vision. *Archives D'ophtalmologie et Revue Generale D'ophtalmologie* 32:531-540.
- Roy M, Sen S, Chakraborti AS (2008) Action of pelargonidin on hyperglycemia and oxidative damage in diabetic rats: Implication for glycation-induced hemoglobin modification. *Life Sci.* 82:1102-1110.
- Sadilova E, Stintzing Florian C, Carle R (2006) Anthocyanins, colour and antioxidant properties of eggplant (*Solanum melongena* L.) and violet pepper (*Capsicum annuum* L.) peel extracts. *Z. für Naturforsch.* 61:527-535.
- Salehi B, Sharifi-Rad J, Cappellini F, Reiner Z, Zorzan D, Imran M, Sener B, Kilic M, El-Shazly M, Fahmy NM, et. al. (2020) The therapeutic potential of anthocyanins: Current approaches based on their molecular mechanism of action. *Front Pharmacol* 11: 1300-1300.
- Saric A, Sobocanec S, Balog T, Kusic B, Sverko V, Dragovic-Uzelac V, Levaj B, Cosic Z, Macak Safranko Z, Marotti T (2009) Improved antioxidant and anti-inflammatory potential in mice consuming sour cherry juice (*Prunus cerasus* cv. Maraska). *Plant Foods Hum. Nutr.* 64:231.
- Scarabelli TM, Mariotto S, Abdel-Azeim S, Shoji K, Darra E, Stephanou A, Chen-Scarabelli C, Marechal JD, Knight R, Ciampa A *et al.* (2008) Targeting STAT1 by myricetin and delphinidin provides efficient protection of the heart from ischemia/reperfusion-induced injury. *FEBS Lett.* 583:531-541.
- Seeram NP, Adams LS, Zhang Y, Lee R, Sand D, Scheuller HS, Heber D (2006) Blackberry, black raspberry, blueberry, cranberry, red raspberry, and strawberry extracts inhibit growth and stimulate apoptosis of human cancer cells *in vitro*. *J. Agric. Food Chem.* 54:9329-9339.
- Seeram NP, Schutzki R, Chandra A, Nair MG (2002) Characterization, quantification, and bioactivities of anthocyanins in *Cornus* species. *J. Agric. Food Chem.* 50:2519-2523.
- Serafino A, Sinibaldi-Vallebona P, Lazzarino G, Tavazzi B, Rasi G, Pierimarchi P, Andreola F, Moroni G, Galvano G, Galvano F *et al.* (2004) Differentiation of human melanoma cells induced by cyanidin-3-O- β -glucopyranoside. *FASEB J.* 18:1940-1942.
- Seymour E, Lewis SK, Urcuyo-Llanes DE, Tanone I, Kirakosyan A, Kaufman PB (2009) Regular tart cherry intake alters abdominal adiposity, adipose gene transcription, and inflammation in obesity-prone rats fed a high fat diet. *J. Med. Food.* 12:935-942.
- Seymour EM, Tanone II, Urcuyo-Llanes DE, Lewis SK, Kirakosyan A, Kondoleon MG, Kaufman PB, Bolling SF *et al.* (2011) Blueberry intake alters skeletal muscle and adipose tissue peroxisome proliferator-activated receptor activity and reduces insulin resistance in obese rats. *J. Med. Food.* 14:1511-1518.
- Shah SA, Amin FU, Khan M, Abid MN, Rehman SU, Kim TH, Kim MW, Kim MO (2016) Anthocyanins abrogate glutamate-induced AMPK activation, oxidative stress, neuroinflammation, and neurodegeneration in postnatal rat brain. *J. Neuroinflammation.* 13:286.
- Shah SA, Yoon GH, Kim MO (2015) Protection of the developing brain with anthocyanins against ethanol-induced oxidative stress and neurodegeneration. *Mol. Neurobiol.* 51:1278-1291.
- Sharma P, McClees SF, Afaq F (2017) Pomegranate for prevention and treatment of cancer: an update. *Molecules.* 22:177.

- Shi M, Mathai ML, Xu G, McAinch AJ, Su XQ (2019) The effects of supplementation with blueberry, cyanidin-3-*O*- β -glucoside, yoghurt and its peptides on obesity and related comorbidities in a diet-induced obese mouse model. *J. Funct. Foods*. 56:92-101.
- Shih P-H, Wu C-H, Yeh C-T, Yen G-C (2011) Protective effects of anthocyanins against amyloid β -peptide-induced damage in neuro-2A cells. *J. Agric. Food Chem*. 59:1683-1689.
- Shih P-H, Yeh C-T, Yen G-C (2005) Effects of anthocyanidin on the inhibition of proliferation and induction of apoptosis in human gastric adenocarcinoma cells. *Food Chem. Toxicol*. 43:1557-1566.
- Shih P-H, Yeh C-T, Yen G-C (2007) Anthocyanins induce the activation of phase-ii enzymes through the antioxidant response element pathway against oxidative stress-induced apoptosis. *J. Agric. Food Chem*. 55:9427-9435.
- Shim SH, Kim JM, Choi CY, Kim CY, Park KH (2012) *Ginkgo biloba* extract and bilberry anthocyanins improve visual function in patients with normal tension glaucoma. *J. Med. Food*. 15:818-823.
- Shimomura I, Funahashi T, Takahashi M, Maeda K, Kotani K, Nakamura T, Yamashita S, Miura M, Fukuda Y, Takemura K *et al.* (1996) Enhanced expression of PAI-1 in visceral fat: possible contributor to vascular disease in obesity. *Nat. Med*. 2:800-803.
- Singh PP, Chandra A, Mahdi F, Roy A, Sharma P (2010) Reconvene and reconnect the antioxidant hypothesis in human health and disease. *Indian J. Clin. Biochem*. 25, 225-243.
- Singletary KW, Jung KJ, Giusti M (2007) Anthocyanin-rich grape extract blocks breast cell DNA damage. *J. Med. Food*. 10:244-251.
- Skates E, Overall J, DeZego K, Wilson M, Esposito D, Lila MA, Komarnytsky S (2018) Berries containing anthocyanins with enhanced methylation profiles are more effective at ameliorating high fat diet-induced metabolic damage. *Food Chem. Toxicol*. 111:445-453.
- Skemiene K, Liobikas J, Borutaite V (2015) Anthocyanins as substrates for mitochondrial complex I - protective effect against heart ischemic injury. *FEBS J*. 282:963-971.
- Skrovankova S, Sumczynski D, Mlcek J, Jurikova T, Sochor J (2015) Bioactive compounds and antioxidant activity in different types of berries. *Int. J. Mol. Sci*. 16:24673-24706.
- Smith MAL, Marley KA, Seigler D, Singletary KW, Meline B (2008) Bioactive properties of wild blueberry fruits. *J. Food Sci*. 65:352-356.
- Sole P, Rigal D, Peyresblanques J (1984) Effects of cyaninoside chloride and Heleniene on mesopic and scotopic vision in myopia and night blindness. *J. FR Ophtalmol*. 7:35-39.
- Song H, Wu T, Xu D, Chu Q, Lin D, Zheng X (2016a) Dietary sweet cherry anthocyanins attenuates diet-induced hepatic steatosis by improving hepatic lipid metabolism in mice. *Nutr*. 32:827-833.
- Song N, Zhang L, Chen W, Zhu H, Deng W, Han Y, Guo J, Qin C (2016b) Cyanidin 3-*O*- β -glucopyranoside activates peroxisome proliferator-activated receptor- γ and alleviates cognitive impairment in the APPswe/PS1 Δ E9 mouse model. *Biochim. Biophys. Acta Mol. Basis Dis*. 1862:1786-1800.
- Speer H, D'Cunha NM, Alexopoulos NI, McKune AJ, Naumovski N (2020). Anthocyanins and human health-A focus on oxidative stress, inflammation and disease. *Antioxidants (Basel)* 9, 366.

- Stintzing FC, Stintzing AS, Carle R, Frei B, Wrolstad RE (2002) Color and antioxidant properties of cyanidin-based anthocyanin pigments. *J. Agric. Food Chem.* 50:6172-6181.
- Stoner GD, Sardo C, Apseloff G, Mullet D, Wargo W, Pound V (2005) Pharmacokinetics of anthocyanins and ellagic acid in healthy volunteers fed freeze-dried black raspberries daily for 7 days. *J. Clin. Pharmacol.* 45:1153–1164.
- Strathearn KE, Yousef GG, Grace MH, Roy SL, Tambe MA, Ferruzzi MG, Wu QL, Simon JE, Lila MA, Rochet JC (2014) Neuroprotective effects of anthocyanin- and proanthocyanidin-rich extracts in cellular models of Parkinsons disease. *Brain Res.* 1555:60-77.
- Su X, Griffin J, Xu J, Ouyang P, Zhao Z, Wang W (2019) Identification and quantification of anthocyanins in purple-fleshed sweet potato leaves. *Heliyon* 5:e01964.
- Sumner MD, Elliott-Eller M, Weidner G, Daubenmier JJ, Chew MH, Marlin R, Raisin CJ, Ornish D (2005) Effects of pomegranate juice consumption on myocardial perfusion in patients with coronary heart disease. *AM J.Cardiol.* 96:810-814.
- Sun H, Zhang P, Zhu Y, Lou Q, He S (2018) Antioxidant and prebiotic activity of five peonidin-based anthocyanins extracted from purple sweet potato (*Ipomoea batatas* (L.) Lam.). *Sci. Rep.* 8:5018.
- Tan L, Yang HP, Pang W, Lu H, Hu YD, Li J, Lu SJ, Zhang WQ, Jiang YG (2014) Cyanidin-3-*O*-galactoside and blueberry extracts supplementation improves spatial memory and regulates hippocampal ERK expression in senescence-accelerated mice. *Biomed. Environ. Sci.* 27:186-196.
- Thiraphatthanavong P, Wattanathorn J, Muchimapura S, Thukham-mee W, Lertrat K, Suriharn B (2014a) The combined extract of purple waxy corn and ginger prevents cataractogenesis and retinopathy in streptozotocin-diabetic rats. *Oxid. Med. Cell. Longev.* 2014:789406.
- Thiraphatthanavong P *et al.* (2014b) Preventive effect of *Zea mays* L. (purple waxy corn) on experimental diabetic cataract. *BioMed. Res. Int.* 2014:507435.
- Tian L, Tan Y, Chen G, Wang G, Sun J, Ou S, Chen W, Bai W (2019) Metabolism of anthocyanins and consequent effects on the gut microbiota. *Crit. Rev. Food Sci. Nutr.* 59:982-991.
- Toufektsian M-C, De Lorgeril M, Nagy N, Salen P, Donati MB, Giordano L (2008) Chronic dietary intake of plant-derived anthocyanins protects the rat heart against ischemia-reperfusion injury. *J. Nutr.* 138:747-752.
- Traustadottir T, Davies SS, Stock AA, Su Y, Heward CB, Roberts LJ, 2nd, Harman SM (2009) Tart cherry juice decreases oxidative stress in healthy older men and women. *J. Nutr.* 139:1896-1900.
- Tsuda T, Horio F, Uchida K, Aoki H, Osawa T (2003) Dietary cyanidin 3-*O*- β -D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. *J. Nutr.* 133:2125-2130.
- Tsuda T, Ueno Y, Kojo H, Yoshikawa T, Osawa T (2005) Gene expression profile of isolated rat adipocytes treated with anthocyanins. *Biochimica et Biophysica Acta (BBA)* 1733:137-147.
- Tsuda T, Ueno Y, Yoshikawa T, Kojo H, Osawa T (2006) Microarray profiling of gene expression in human adipocytes in response to anthocyanins. *Biochem. Pharmacol.* 71:1184-1197.

- Turck P, Fraga S, Salvador I, Campos-Carraro C, Lacerda D, Bahr A, Ortiz V, Hickmann A, Koetz M, Bello-Klein A *et al.* (2020) Blueberry extract decreases oxidative stress and improves functional parameters in lungs from rats with pulmonary arterial hypertension. *Nutr.* 70:110579.
- Turner K, Fischer JG, Grider A, Hargrove J, Tokarev J (2009) Anthocyanins increase antioxidant enzyme activity in ht-29 adenocarcinoma cells by martha. MsC thesis. Athens, Georgia, USA.
- Ulla A, Mohamed MK, Sikder B, Rahman AT, Sumi FA, Hossain M, Reza HM, Rahman GMS, Alam MA (2017) Coenzyme Q10 prevents oxidative stress and fibrosis in isoprenaline induced cardiac remodeling in aged rats. *BMC Pharmacol. Toxicol.* 18, 29-29.
- Ullah R, Khan M, Shah SA, Saeed K, Kim MO (2019) Natural antioxidant anthocyanins—A hidden therapeutic candidate in metabolic disorders with major focus in neurodegeneration. *Nutr.* 11:1195.
- Valensi P, Behar A, Attalah M, Cohen-Boulakia F, Pariès J, Attali JR (1998) Increased capillary filtration of albumin in diabetic patients-relation with gender, hypertension, microangiopathy, and neuropathy. *Metab.* 47:503-507.
- Valensi P, Cohen-Boulakia F, Attali JR, Behar A (1997) Changes in capillary permeability in diabetic patients. *Clin. Hemorheol. Micro.* 17:389-394.
- Valenza A, Bonfanti C, Pasini ME, Bellosa P (2018) Anthocyanins function as anti-inflammatory agents in a *Drosophila* model for adipose tissue macrophage infiltration. *BioMed. Res. Int.* 2018:6413172.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J (2007) Free radicals and antioxidants in normal physiological functions and human disease. *Int. J Biochem. Cell Biol.* 39, 44-84.
- van Velden DP, Mansvelt EPG, Troup GJ (2002) Red wines good, white wines bad? *Redox Rep.* 7:315-316.
- Varadinova MG, Docheva-Drenska DI, Boyadjieva NI (2009) Effects of anthocyanins on learning and memory of ovariectomized rats. *Menopause* 16:345-349.
- Varma SD, Kinoshita JH (1976) Inhibition of lens aldose reductase by flavonoids—Their possible role in the prevention of diabetic cataracts. *Biochem. Pharmacol.* 25:2505-2513.
- Vendrame S, Klimis-Zacas D (2015) Anti-inflammatory effect of anthocyanins via modulation of nuclear factor-kappaB and mitogen-activated protein kinase signaling cascades. *Nutr. Rev.* 73:348-358.
- Vepsäläinen S, Koivisto H, Pekkarinen E, Mäkinen P, Dobson G, McDougall GJ, Stewart D, Haapasalo A, Karjalainen RO, Tanila H *et al.* (2013) Anthocyanin-enriched bilberry and blackcurrant extracts modulate amyloid precursor protein processing and alleviate behavioral abnormalities in the APP/PS1 mouse model of Alzheimer's disease. *J. Nutr. Biochem.* 24:360-370.
- Wallace TC, Giusti MM (2015) Anthocyanins. *Advances in nutrition* (Bethesda, Md.) 6, 620-622.
- Wang L-S, Stoner GD (2008) Anthocyanins and their role in cancer prevention. *Cancer letters* 269, 281-290.
- Williamson G, Sies H, Heber D, Keen CL, Macdonald IA, Actis-Gorreta L, Momma TY, Ottaviani JJ, Holt RR, Schroeter H *et al.* (2009) Functional foods for health promotion: state-of-the-science on dietary flavonoids, Extended abstracts from the 12th Annual Conference on Functional Foods for Health Promotion, April 2009 *Nutr. Rev.* 67:736-743.

- Woodward G, Kroon P, Cassidy A, Kay C (2009) Anthocyanin stability and recovery: Implications for the analysis of clinical and experimental samples. *J. Agric. Food Chem.* 57:5271-5278.
- Worsztynowicz P, Napierała M, Białas W, Grajek W, Olkiewicz M (2014) Pancreatic α -amylase and lipase inhibitory activity of polyphenolic compounds present in the extract of black chokeberry (*Aronia melanocarpa* L.). *Process Biochem.* 49:1457-1463.
- Wu T, Jiang Z, Yin J, Long H, Zheng X (2016a) Anti-obesity effects of artificial planting blueberry (*Vaccinium ashei*) anthocyanin in high-fat diet-treated mice. *Int. J. Food Sci. Nutr.* 67:257-264.
- Wu T, Tang Q, Gao Z, Yu Z, Song H, Zheng X, Chen W (2013) Blueberry and mulberry juice prevent obesity development in C57BL/6 mice. *PLoS ONE.* 8:e77585.
- Wu T, Tang Q, Yu Z, Gao Z, Hu H, Chen W, Zheng X, Yu T (2014) Inhibitory effects of sweet cherry anthocyanins on the obesity development in C57BL/6 mice. *Int. J. Food Sci. Nutr.* 65:351-359.
- Wu T, Yang L, Guo X, Zhang M, Liu R, Sui W (2018) Raspberry anthocyanin consumption prevents diet-induced obesity by alleviating oxidative stress and modulating hepatic lipid metabolism. *Food Funct.* 9:2112-2120.
- Wu T, Yin J, Zhang G, Long H, Zheng X (2016b) Mulberry and cherry anthocyanin consumption prevents oxidative stress and inflammation in diet-induced obese mice. *Mol. Nutr. Food Res.* 60:687-694.
- Xu J-W, Ikeda K, Yamori Y (2004) Cyanidin-3-*O*-glucoside regulates phosphorylation of endothelial nitric oxide synthase. *FEBS Lett.* 574:176-180.
- Yamahara J, Mibu H, Sawada T, Fujimura H, Takino S, Yoshikawa M (1981) Biologically active principles of crude drugs. Antidiabetic principles of Corni Fructus in experimental diabetes induced by streptozotocin. *J. Pharm. Soc. Jpn.* 101:86-90.
- Yan F, Dai G, Zheng X (2016) Mulberry anthocyanin extract ameliorates insulin resistance by regulating PI3K/AKT pathway in HepG2 cells and db/db mice. *J. Nutr. Biochem.* 36:68-80.
- Yang Z, Zhai W (2010) Identification and antioxidant activity of anthocyanins extracted from the seed and cob of purple corn (*Zea mays* L.). *Innov. Food Sci. Emerg. Technol.* 11:169-176.
- Yaribeygi H, Sathyapalan T, Atkin SL, Sahebkar A (2020) Molecular mechanisms linking oxidative stress and diabetes mellitus. *Oxid. Med. Cell. Longev.* 2020, 8609213.
- Yawadio R, Tanimori S, Morita N (2007) Identification of phenolic compounds isolated from pigmented rices and their aldose reductase inhibitory activities. *Food Chem.* 101:1616-1625.
- You Q, Chen F, Wang X, Luo PG, Jiang Y (2011) Inhibitory effects of muscadine anthocyanins on alpha-glucosidase and pancreatic lipase activities. *J. Agric. Food Chem.* 59:9506-9511.
- Youdim KA, Martin A, Joseph JA (2000) Incorporation of the elderberry anthocyanins by endothelial cells increases protection against oxidative stress. *Free Radic. Biol. Med.* 29:51-60.
- Youdim KA, McDonald J, Kalt W, Joseph JA (2002) Potential role of dietary flavonoids in reducing microvascular endothelium vulnerability to oxidative and inflammatory insults. *J. Nutr. Biochem.* 13:282-288.
- Zafra-Stone S, Yasmin T, Bagchi M, Chatterjee A, Vinson JA, Bagchi D (2007) Berry anthocyanins as novel antioxidants in human health and disease prevention. *Mol. Nutr. Food Res.* 51:675-683.

- Zdarilova A, Svobodova AR, Chytilova K, Simanek V, Ulrichova J (2010) Polyphenolic fraction of *Lonicera caerulea* L. fruits reduces oxidative stress and inflammatory markers induced by lipopolysaccharide in gingival fibroblasts. *Food Chem. Toxicol.* 48:1555-1561.
- Zhang H, Guo J, Mao L, Li Q, Guo M, Mu T, Zhang Q, Bi X (2018) Up-regulation of miR-24-1-5p is involved in the chemoprevention of colorectal cancer by black raspberry anthocyanins. *Br. J. Nutr.* 122: 518-526.
- Zhang H, Xu Z, Zhao H, Wang X, Pang J, Li Q, Yang Y, Ling W (2020a) Anthocyanin supplementation improves anti-oxidative and anti-inflammatory capacity in a dose–response manner in subjects with dyslipidemia. *Redox Biol.* 32:101474.
- Zhang Q, Gonzalez de Mejia E, Luna-Vital D, Tao T, Chandrasekaran S, Chatham L, Juvik J, Singh V, Kumar D (2019a) Relationship of phenolic composition of selected purple maize (*Zea mays* L.) genotypes with their anti-inflammatory, anti-adipogenic and anti-diabetic potential. *Food Chem.* 289:739-750.
- Zhang X, Wei J, Tian J, Li N, Jia L, Shen W, Cui J (2019b) Enhanced anthocyanin accumulation of immature radish microgreens by hydrogen-rich water under short wavelength light. *Sci. Hortic.* 247:75-85.
- Zhang Y, Jayaprakasam B, Seeram NP, Olson LK, DeWitt D, Nair MG (2004) Insulin secretion and cyclooxygenase enzyme inhibition by cabernet sauvignon grape skin compounds. *J. Agric. Food Chem.* 52:228-233.
- Zhang Y, Meng Q, Yin J, Zhang Z, Bao H, Wang X (2020b) Anthocyanins attenuate neuroinflammation through the suppression of MLK3 activation in a mouse model of perioperative neurocognitive disorders. *Brain Res.* 1726:146504.
- Zhang Y, Vareed SK, Nair MG (2005) Human tumor cell growth inhibition by nontoxic anthocyanidins, the pigments in fruits and vegetables. *Life Sci.* 76:1465-1472.
- Zhang Z-c, Zhou Q, Huangfu G-y, Wu Y, Zhang J-l (2019c) Anthocyanin extracts of lingonberry (*Vaccinium vitis-idaea* (L.)) attenuate serum lipids and cholesterol metabolism in HCD-induced hypercholesterolaemic male mice. *Int. J. Food Sci. Technol.* 54:1576-1587.
- Zhao X, Feng P, He W, Du X, Chen C, Suo L, Liang M, Zhang N, Na A, Zhang Y (2019) The prevention and inhibition effect of anthocyanins on colorectal Cancer. *Curr. Pharm. Des.* 25, 4919-4927.
- Zhu F (2018) Anthocyanins in cereals: Composition and health effects. *Food Res. Int.* 109:232-249.
- Zhu F, Cai Y-Z, Yang X, Ke J, Corke H (2010) Anthocyanins, hydroxycinnamic acid derivatives, and antioxidant activity in roots of different chinese purple-fleshed sweetpotato genotypes. *J. Agric. Food Chem.* 58:7588-7596.
- Zhu Y, Sun H, He S, Lou Q, Yu M, Tang M, Tu L (2018) Metabolism and prebiotics activity of anthocyanins from black rice (*Oryza sativa* L.) *in vitro*. *PLoS ONE.* 13:e0195754.