



Size-susceptibility of *Cornu aspersum* exposed to the malacopathogenic nematodes *Phasmarhabditis hermaphrodita* and *P. californica*

Alex Grannell, James Cutler & Robbie Rae

To cite this article: Alex Grannell, James Cutler & Robbie Rae (2021) Size-susceptibility of *Cornu aspersum* exposed to the malacopathogenic nematodes *Phasmarhabditis hermaphrodita* and *P. californica*, *Biocontrol Science and Technology*, 31:11, 1149-1160, DOI: [10.1080/09583157.2021.1929072](https://doi.org/10.1080/09583157.2021.1929072)

To link to this article: <https://doi.org/10.1080/09583157.2021.1929072>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 23 May 2021.



Submit your article to this journal [↗](#)



Article views: 515



View related articles [↗](#)



View Crossmark data [↗](#)



RESEARCH ARTICLE



Size-susceptibility of *Cornu aspersum* exposed to the malacopathogenic nematodes *Phasmarhabditis hermaphrodita* and *P. californica*

Alex Grannell*, James Cutler* and Robbie Rae

School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, UK

ABSTRACT

Slugs and snails are agricultural pests usually controlled by chemical bait pellets however an alternative method is the commercially produced malacopathogenic nematode *Phasmarhabditis hermaphrodita*, which is sold across northern Europe. *P. hermaphrodita* can kill several slug species but is unable to affect larger pestiferous snails. Therefore, we examined whether the closely related species *Phasmarhabditis californica*, isolated from the U.K. could kill neonate or adult common garden snails (*Cornu aspersum*). In our first experiment, neonate *C. aspersum* were exposed to 1000 nematodes per tube containing three strains of *P. californica* (designated DMG0017, DMG0018 or DMG0019) (as well as the commercial formulation of *P. hermaphrodita* DMG0001) and feeding inhibition and survival were monitored over 14 days. All nematodes apart from *P. californica* (DMG0017) killed the snails and caused feeding inhibition. In a follow up experiment we exposed adult *C. aspersum* to two doses (30 and 90 nematodes per cm²) of *P. californica* (DMG0019) and *P. hermaphrodita* (DMG0001) and monitored survival, feeding inhibition and changes in weight over 21 days. Neither nematode species affected survival, feeding inhibition or weight of the adult snails. In summary, *P. californica* (DMG0018 and DMG0019) are as pathogenic as *P. hermaphrodita* (DMG0001) and could be used to target neonate snails, however, the use of malacopathogenic nematodes to control large adult snails continues to be problematic.

ARTICLE HISTORY

Received 23 February 2021

Accepted 8 May 2021

KEYWORDS

Parasitic nematodes;
gastropods; biological
control

1. Introduction

Many species of slugs and snails are highly pestiferous and affect crop production across the world (South, 1992). They are usually controlled by chemical bait pellets, such as metaldehyde, but due to its toxicity to non-target organisms (Homeida &

CONTACT Robbie Rae ✉ r.g.rae@ljmu.ac.uk School of Biological and Environmental Sciences, Liverpool John Moores University, Byrom Street, Liverpool L33AF, UK

*Both authors contributed equally.

Supplemental data for this article can be accessed at <https://doi.org/10.1080/09583157.2021.1929072>.

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Cooke, 1982) it has been banned in many European countries. An alternative to chemical control is the biological control agent *Phasmarhabditis hermaphrodita*, a parasitic nematode that can infect and kill several slug and snail species which has been formulated into a product called Nemaslug® available from BASF Agricultural Specialities (Wilson et al., 1993). *P. hermaphrodita* is applied to soil where it seeks slugs through attraction to mucus and faeces (Rae et al., 2006) and penetrates through a pore at the back of the mantle and kills the host in 4–21 days (Tan & Grewal, 2001a; Wilson et al., 1993). It is thought the nematodes vector the bacterium *Moraxella osloensis*, inside the slug where it proliferates and produces a lipopolysaccharide, which causes death to the slug (Tan & Grewal, 2001b, 2002). The nematodes reproduce on the cadaver and when the bacterial food supply is exhausted new dauer juveniles are produced that go in search of more hosts. *P. hermaphrodita* has been successfully shown to protect an array of crops from slug damage such as lettuce (Wilson et al., 1995), asparagus (Ester et al., 2003a), hostas (Grewal et al., 2001) and orchids (Ester et al., 2003b) (to name but a few).

P. hermaphrodita is not the only member of the 13 species (Tandingan De Ley et al., 2017) that encompass the *Phasmarhabditis* genus which can kill slugs. As well as *P. hermaphrodita*, *P. papillosa* (Laznik et al., 2020; McDonnell et al., 2020; Pieterse et al., 2017; Tandingan De Ley et al., 2020), *P. neopapillosa* (Hooper et al., 1999), *P. tawfiki* (Azzam & Tawfiki, 2003), *P. safricana* (Ross et al., 2018) and *P. californica* (McDonnell et al., 2020; Tandingan De Ley et al., 2020) have been shown to infect and kill slugs and snails. There is also a short report stating *P. bohémica*, *P. bonaquaense* and *P. apuliae* can also kill slugs (Nermut et al., 2019). One of the most frequently isolated species from the *Phasmarhabditis* genus is *P. californica*. It has been isolated from the U.K. (Andrus & Rae, 2019), the U.S.A. (McDonnell et al., 2020; Tandingan De Ley et al., 2016), New Zealand (Wilson et al., 2016) and recently Canada (Brophy et al., 2020). U.S. strains of *P. californica* have been shown to kill the grey field slug (*Deroceras reticulatum*) (McDonnell et al., 2020) and snails (*Theba pisana*) (Tandingan De Ley et al., 2020), however, there is nothing known about the pathogenic ability of *P. californica* isolated from the U.K. Three strains (designated DMG0017, DMG0018 and DMG0019) were isolated from a single snail (*Oxychilus draparnaudi*) in Pembrokeshire, Wales (SM809057) and have been kept in culture since 2014 on White traps (White, 1927) using rotting slug (*Limax flavus*), see Andrus and Rae (2019) for further details. Therefore, in order to investigate the pathogenicity of these strains we exposed *P. californica* (DMG0017, DMG0018 and DMG0019) to neonate and adult *Cornu aspersum*, a notorious pest species found across Europe and America (Guiller et al., 2012), and compared their effectiveness at causing mortality to the commercial strain of *P. hermaphrodita* (DMG0001). We also assessed whether the nematodes caused feeding inhibition in the snails (a common symptom of *P. hermaphrodita* infection in slugs – Glen et al., 2000) and would subsequently affect the growth and development of the snails by assessing differences in weight at the start and the end of the experiment. If *P. californica* killed *C. aspersum* it could be developed as a biological control agent to target pestiferous snails, rather than *P. hermaphrodita* which struggles to cause mortality to larger snails and slugs (Grimm, 2002; Speiser et al., 2001).

2. Materials and methods

2.1. Source of invertebrates

Neonate *C. aspersum* (0.16 ± 0.05 g, $n = 150$) and adult *C. aspersum* ‘gros gris’ (Supplementary Figure 1(A)) (mean weight 21.7 ± 0.33 g, $n = 180$) were purchased from online suppliers H and RH Escargot. Snails were kept at room temperature in terraria ($35 \times 23 \times 22$ cm) with ventilated lids containing moist coconut husk and fed lettuce and cucumber *ad libitum*. Snails were kept in moist conditions by liberally spraying tap water over the coconut husk every 2 days for 8 days prior to the experiment. *P. hermaphrodita* (DMG0001) were supplied by BASF Agricultural Specialities. Three strains of *P. californica* (DMG0017, DMG0018 and DMG0019) (Supplementary Figure 1(B)) were grown to the infective dauer stage using White traps (White, 1927) of dead *L. flavus* using methods by Andrus and Rae (2019).

2.2. Assessing the susceptibility of snails to *P. hermaphrodita* and *P. californica*

To assess the susceptibility of neonate snails a protocol by Cutler and Rae (2020) was adapted. Briefly, a piece of cotton wool was added to the bottom of fifteen 20 ml universal tubes. Nemaslug® (1–2 g) containing dauer stage *P. hermaphrodita* (DMG0001) was added to 45 ml of tap water, shaken vigorously for 2 min and then the numbers of alive *P. hermaphrodita* (DMG0001) in 20 μ l were quantified three times. This process was repeated with *P. californica* (DMG0017, DMG0018 and DMG0019) though samples were removed from the surrounding water from fresh White traps where dauer stage nematodes were present. One thousand dauer stage *P. hermaphrodita* (DMG0001) (in 1 ml of water) were added to five tubes using a 1000 μ l pipette and this process was repeated for *P. californica* (DMG0017, DMG0018 and DMG0019). Five tubes had the equivalent volume of water (1 ml) and no nematodes added and acted as the control. Two *C. aspersum* were added to each tube and the tubes were loosely sealed and stored in an incubator at 15°C. After 5 days exposure to the nematodes the snails were removed and allowed to feed on a 3.5 cm diameter piece of lettuce. Feeding was quantified on days 8 and 14 using methods by Cutler and Rae (2020). Briefly, remnants of the leaf discs were traced onto 1×1 mm² graph paper and their sized quantified by counting the number of 1 mm² squares. Survival of the snails was monitored every 2–3 days for 21 days. The experiment was repeated three times in total with 30 snails being exposed to each treatment.

To test the susceptibility of adult *C. aspersum* to *P. hermaphrodita* and *P. californica* we added approx. 100 g of pre-moistened coconut husk was added to 18 boxes ($35 \times 23 \times 22$ cm) to a depth of approx. 2 cm. To three boxes the recommended rate for field application (30 nematodes per cm²) of *P. hermaphrodita* (Rae et al., 2007) was applied. The same dose was used for three boxes for *P. californica* DMG0019 application. We chose the DMG0019 strain of *P. californica* as it consistently produces large numbers of dauer juveniles under lab conditions (data not shown). This set up was also repeated to examine the susceptibility of *C. aspersum* to a higher dose (90 nematodes per cm²) of *P. hermaphrodita* or *P. californica*. The other six boxes received water (10 ml) and acted as the control. Five discs of lettuce (3.5 cm diameter) were added to each box and the amount eaten was quantified every 2–3 days. The snails were weighed before

the start of the experiment and five *C. aspersum* were added to each box and stored at 15°C. The survival of the snails, as well as the amount eaten was quantified every 2–3 days for 21 days. After each time point the amount of food eaten was quantified it was replaced with fresh lettuce discs. The snails were weighed at the end of the experiment and the experiment was repeated twice.

2.3. Data analysis

Survival of neonate snails exposed to 0 or 1000 *P. hermaphrodita* (DMG0001), *P. californica* (DMG0017, DMG0018 and DMG0019) was analysed using a Log Rank test using OASIS (Yang et al., 2011). A Kruskal Wallis with Dunn's test was used to compare the amount eaten by neonate snails on day 8 and 14. The survival of adult snails exposed to 0, 30 or 90 *P. hermaphrodita* (DMG0001) and *P. californica* (DMG0019) per cm² on day 21 was compared using a Kruskal–Wallis test. The amount of lettuce eaten by adult snails exposed to 0, 30 or 90 *P. hermaphrodita* (DMG0001) and *P. californica* (DMG0019) per cm² on each recording day was compared using a Kruskal–Wallis test. The weight of the adult snails at the start and the end of the experiment was compared using a Student's *t* test. A One Way ANOVA was used to compare the mean weight of snails in each treatment on day 0 and 21. Normality was assessed using Kolmogorov–Smirnov tests. Statistical analysis was carried out using SPSS.

3. Results

3.1. Susceptibility of neonate *C. aspersum* to *P. hermaphrodita* and *P. californica*

There was a significant difference between the survival of untreated neonate *C. aspersum* compared to snails exposed to *P. hermaphrodita* (DMG0001) ($P = 0.0167$), *P. californica* (DMG0018) ($P = 0.0106$) and *P. californica* (DMG0019) ($P = 0.0105$) over 14 days (Figure 1). However, *P. californica* (DMG0017) ($P = 0.0809$) had no significant effect on the survival of neonate *C. aspersum*.

3.2. Feeding inhibition of neonate *C. aspersum* exposed to nematodes

After 8 and 14 days all nematodes had a significant effect on the amount of lettuce neonate *C. aspersum* ate compared to the untreated control ($P < 0.05$) (Figure 2(A,B)). There was no difference between the amount of lettuce eaten by *C. aspersum* exposed to *P. hermaphrodita* (DMG0001), or *P. californica* (DMG0017, DMG0018 and DMG0019) on day 8 or 14 ($P > 0.05$) (Figure 2(A,B)).

3.3. Susceptibility and feeding inhibition of adult *C. aspersum* to *P. hermaphrodita* and *P. californica*

After 21 days there was no significant difference between the survival of adult *C. aspersum* when exposed to 0, 30 or 90 *P. hermaphrodita* ($P > 0.05$) or *P. californica* DMG0019 ($P > 0.05$) per cm² (Figure 3(A,B)).




Figure 1. The mean percentage survival of neonate *Cornu aspersum* exposed to no dauer juvenile nematodes (black solid line), *Phasmarhabditis hermaphrodita* DMG0001 (black dotted line), *Phasmarhabditis californica* DMG0017 (grey solid line), *P. californica* DMG0018 (grey dashed line) and *P. californica* DMG0019 (black dash line) over 14 days. Bars represent \pm one standard error.

Similarly, there was no significant difference between the amount of lettuce eaten when adult *C. aspersum* were exposed to 0, 30 or 90 *P. hermaphrodita* ($P > 0.05$) or *P. californica* DMG0019 ($P > 0.05$) per cm^2 on any of the days (Figure 4(A,B)).

3.4. Differences in weight of adult *C. aspersum* exposed to nematodes

The snails lost weight over the course of the experiment as there was a significant difference between the weight of *C. aspersum* on day 0 and 21 when exposed to 0, 30 and 90 *P. hermaphrodita* ($P < 0.05$) and *P. californica* DMG0019 ($P < 0.05$) per cm^2 (Figure 5(A, B)). However, both species of nematode did not induce feeding inhibition as there was no significant difference between the weight of snails on day 21 after exposure to 0, 30 or 90 *P. hermaphrodita* ($P > 0.05$) or *P. californica* DMG0019 ($P > 0.05$) per cm^2 (Figure 5(A,B)).

4. Discussion

We have shown both *P. hermaphrodita* and *P. californica* were able to kill neonate *C. aspersum* but not adults. The lack of effect on adult snails is perhaps not surprising as previous research has shown *P. hermaphrodita* has little effect on adult *C. aspersum* (Glen et al., 1996), however, this was only known for *P. hermaphrodita* and not *P. californica*. Differences in the susceptibility of different stages of terrestrial gastropod to *Phasmarhabditis* nematodes have been reported previously. Both Grimm (2002) and Speiser et al. (2001) showed *P. hermaphrodita* could kill juvenile *Arion lusitanicus* but not larger specimens. Similar results have been observed with experiments with the giant




Figure 2. Feeding of neonate *C. aspersum* exposed to dauer juvenile *P. hermaphrodita* DMG0001, *P. californica* DMG0017, *P. californica* DMG0018 and *P. californica* DMG0019 on day 8 (A) and 14 (B). Bars represent \pm one standard error.

African snail (*Lissachatina fulica*). Specifically, McDonnell et al. (2018) showed that a US strain of *P. hermaphrodita* could kill >80% of neonate *L. fulica* after 31 days whilst Williams and Rae (2015) showed the commercial strain of *P. hermaphrodita* could not kill larger snails. However, using other *Phasmarhabditis* species has yielded potential for controlling larger gastropods. For example, recently *P. papillosa* has been shown to kill young adult and large adult *Arion vulgaris* (Laznik et al., 2020). As these species are problematic to control due to their high reproductive rate (South, 1992) and their resistance to *P. hermaphrodita* (Grimm, 2002; Speiser et al., 2001), the use of other *Phasmarhabditis* species warrants further investigation.




Figure 3. The mean percentage survival of adult *C. aspersum* exposed to 0 (black line), 30 (black dashed line) or 90 dauer juvenile *P. hermaphrodita* (grey line) (A) or *P. californica* DMG0019 (B) per cm² (grey line) over 21 days. Bars represent \pm one standard error.

In general, the pathogenic potential of *P. californica* is poorly researched. There has been no research on the pathogenic potential of *P. californica* isolated from the U.K. however, Tandingan De Ley et al. (2020) showed *P. californica* from the U.S. could kill the snail *T. pisana* (albeit with high doses of 150 nematodes per cm²) and was equally as pathogenic as *P. hermaphrodita* and *P. papillosa*. McDonnell et al. (2020) tested the infectivity of *P. hermaphrodita*, *P. californica* and *P. papillosa* to *D. reticulatum* and




Figure 4. Feeding of adult *C. aspersum* exposed to 0 (black line), 30 (black dashed line) or 90 dauer juvenile *P. hermaphrodita* (grey line) (A) or *P. californica* DMG0019 (B) per cm² (grey line) over 21 days. Bars represent \pm one standard error.

found *P. papillosa* was the most pathogenic. From our research it is clear that *P. californica* is pathogenic to neonate stages of *C. aspersum* but this is strain dependent. Specifically, both *P. californica* DMG0018 and DMG0019 were pathogenic to neonate *C. aspersum* but *P. californica* DMG0017 was not. Natural variation in pathogenicity of *Phasmarhabditis* nematodes towards other terrestrial gastropods has been reported. For example, Cutler and Rae (2020) reported that several wild strains of *P. hermaphrodita* were more virulent to slugs (*Deroceras invadens*) than the commercial strain and other wild strains. It should be noted that these *P. hermaphrodita* strains were isolated from separate slugs from different areas around Liverpool and are genetically distinct (Howe et al., 2020). The three strains of *P. californica* used in this study were




Figure 5. Weight of adult *C. aspersum* exposed to 0, 30 or 90 dauer juvenile *P. hermaphrodita* (A) or *P. californica* DMG0019 (B) per cm² on day 0 (black) and 21 (white). Bars represent \pm one standard error.

isolated from one single snail (*O. draparnaudi*) (Andrus & Rae, 2019). This highlights how strains isolated in close proximity to each other and phylogenetically similar (Howe et al., 2020) differ in their pathogenic abilities (Cutler & Rae, 2020) as well as host seeking abilities (Andrus et al., 2018). Therefore, in order to produce and develop a suitable biological control agent for slugs and snails, intraspecific variation of strains should be investigated.

It is clear that the ability to kill terrestrial gastropods is an evolutionary conserved trait across the genus *Phasmarhabditis* as from the 13 species *P. neopapillosa* (Hooper et al., 1999), *P. tawfiki* (Azzam & Tawfiki, 2003), *P. safricana* (Ross et al., 2018), *P. bohémica*,

P. bonaquaense and *P. apuliae* can also kill slugs (Nermut et al., 2019) (as well as *P. hermaphrodita* and *P. californica*). As these nematodes are easy to isolate, maintain (Andrus & Rae, 2019) and capable of killing a range of pestiferous gastropods, there is ample opportunity to explore the pathogenic potential of all these other species for commercialisation and formulation to make superior biological control agents. To begin with *P. californica* (particularly strains DMG0018 and DMG0019) would be worth investigating further as they show a clear pathogenic effect to neonate *C. aspersum*.

Acknowledgements

We are grateful to BASF Agricultural Specialities for funding AG and JC and Tom Goddard and Jack Shepherd for discussions.

Disclosure statement

No potential conflict of interest was reported by the author(s).

References

- Andrus, P., Ingle, O., Coleman, T., & Rae, R. (2018). Gastropod parasitic nematodes (*Phasmarhabditis* sp.) are attracted to hyaluronic acid in snail mucus by cGMP signalling. *Journal of Helminthology*, 94(42), e9. <https://doi.org/10.1017/S0022149X18000986>
- Andrus, P., & Rae, R. (2019). Development of *Phasmarhabditis hermaphrodita* (and members of the *Phasmarhabditis* genus) as new genetic model nematodes to study the genetic basis of parasitism. *Journal of Helminthology*, 93(3), 319–331. <https://doi.org/10.1017/S0022149X18000305>
- Azzam, K. M., & Tawfiki, M. F. S. (2003). Capability of the nematode *Phasmarhabditis tawfiki* Azzam to infect some snails, slugs and insects larvae in the laboratory. *Egyptian German Society Journal of Zoology*, 42, 27–32.
- Brophy, T., Howe, D. K., Denver, D. R., & Luong, L. T. (2020). First report of a gastropod parasitic nematode *Phasmarhabditis californica* (Nematoda: Rhabditidae) in Alberta, Canada. *Journal of Nematology*, 52, e2020–e2092. <https://doi.org/10.21307/jofnem-2020-092>
- Cutler, J., & Rae, R. (2020). Pathogenicity of wild and commercial *Phasmarhabditis hermaphrodita* exposed to the pestiferous slug *Deroceras invadens*. *Journal of Invertebrate Pathology*, 174, 107435. <https://doi.org/10.1016/j.jip.2020.107435>
- Ester, A., Rozen Van, K., & Hazendonk, A. (2003b). Efficacy of pesticides to control *Lehmannia valentiana* (Férussac) in orchids (*Cymbidium*) in greenhouse experiments. Slugs and Snails: Agricultural, Veterinary and Environmental Perspective. *British Crop Protection Council Symposium Proceedings*, 80, 89–94.
- Ester, A., Rozen Van, K., & Molendijk, L. P. G. (2003a). Field experiments using the rhabditid nematode *Phasmarhabditis hermaphrodita* or salt as control measures against slugs in green asparagus. *Crop Protection*, 22(5), 689–695. [https://doi.org/10.1016/S0261-2194\(03\)00003-6](https://doi.org/10.1016/S0261-2194(03)00003-6)
- Glen, D. M., Wilson, M. J., Hughes, L., Cargeeg, P., & Hajjar, A. (1996). Exploring and exploiting the potential of *Phasmarhabditis hermaphrodita* as a biocontrol agent for slugs. *Slugs and Snails: Agricultural, Veterinary and Environmental Perspectives, BCPC Symposium Proceedings*, 66, 271–280.
- Glen, D. M., Wiltshire, C. W., Hughes, L., Ester, A., Van Rozen, K., Castillejo, J., Iglesias, J., Speiser, B., Coupland, J., & Gwynn, R. (2000). The use of slug-parasitic nematodes and other techniques for control of slug and snail damage in horticultural crops: *Pests and Diseases. British Crop Protection Council (BCPC) Symposium Proceedings*, 345–350.

- Grewal, P. S., Grewal, S. K., Taylor, R. A. J., & Hammond, R. B. (2001). Application of molluscicidal nematodes to slug shelters: A novel approach to economic biological control of slugs. *Biological Control*, 22(1), 72–80. <https://doi.org/10.1006/bcon.2001.0958>
- Grimm, B. (2002). Effect of the nematode *Phasmarhabditis hermaphrodita* on young stages of the pest slug *Arion lusitanicus*. *Journal of Molluscan Studies*, 68(1), 25–28. <https://doi.org/10.1093/mollus/68.1.25>
- Guiller, A., Martin, M.-C., Hiraux, C., & Madec, L. (2012). Tracing the invasion of the Mediterranean land snail *Cornu aspersum aspersum* becoming an agricultural and garden pest in areas recently introduced. *PLoS ONE*, 7(12), e49674. <https://doi.org/10.1371/journal.pone.0049674>
- Homeida, A. M., & Cooke, R. G. (1982). Pharmacological aspects of metaldehyde poisoning in mice. *Journal of Veterinary Pharmacology and Therapeutics*, 5(1), 77–82. <https://doi.org/10.1111/j.1365-2885.1982.tb00500.x>
- Hooper, D. J., Wilson, M. J., Rowe, J. A., & Glen, D. M. (1999). Some observations on the morphology and protein profiles of the slug parasitic nematodes *Phasmarhabditis hermaphrodita* and *P. neopapillosa* (Nematoda: Rhabditidae). *Nematology*, 1(2), 173–182. <https://doi.org/10.1163/156854199508144>
- Howe, D. K., Ha, A. D., Colton, A., Tandingan De Ley, I., Rae, R. G., Ross, J., Wilson, M., Nermut, J., Zhao, Z., McDonnell, R. J., & Denver, D. R. (2020). Phylogenetic evidence for the invasion of a commercialized European *Phasmarhabditis hermaphrodita* lineage into North America and New Zealand. *PLoS One*, 15(8), e0237249. <https://doi.org/10.1371/journal.pone.0237249>
- Laznik, Ž, Majić, I., Trdan, S., Mala, A. P., Pieterse, A., & Ross, J. L. (2020). Is *Phasmarhabditis papillosa* (Nematoda: Rhabditidae) a possible biological control agent against the Spanish slug, *Arion vulgaris* (Gastropoda: Arionidae)? *Nematology*, 1, 1–9. <https://doi.org/10.1163/15685411-bja10063>
- McDonnell, R. J., Colton, A. J., Howe, D. K., & Denver, D. R. (2020). Lethality of four species of *Phasmarhabditis* (Nematoda: Rhabditidae) to the invasive slug, *Deroceras reticulatum* (Gastropoda: Agriolimacidae) in laboratory infectivity trials. *Biological Control*, 150. <https://doi.org/10.1016/j.biocontrol.2020.104349>
- McDonnell, R., Tandingan De Ley, T., & Paine, T. D. (2018). Susceptibility of neonate *Lissachatina fulica* (Achatinidae: Mollusca) to a US strain of the nematode *Phasmarhabditis hermaphrodita*. *Biocontrol, Science and Technology*, 28(11), 1091–1095. <https://doi.org/10.1080/09583157.2018.1514586>
- Nermut, J., Holley, M., & Půža, V. (2019). *Phasmarhabditis hermaphrodita* is not the only slug killing nematode. *Microbial and Nematode Control of Invertebrate Pests IOCS-WPRS Bulletin*, 150, 152–156.
- Pieterse, A., Tiedt, L. R., Malan, A. P., & Ross, J. L. (2017). First record of *Phasmarhabditis papillosa* (Nematoda: Rhabditidae) in South Africa, and its virulence against the invasive slug, *Deroceras panormitanum*. *Nematology*, 19(9), 1035–1050. <https://doi.org/10.1163/15685411-00003105>
- Rae, R. G., Robertson, J. F., & Wilson, M. J. (2006). The chemotactic response of *Phasmarhabditis hermaphrodita* (Nematoda: Rhabditida) to cues of *Deroceras reticulatum* (Mollusca: Gastropoda). *Nematology*, 8(2), 197–200. <https://doi.org/10.1163/156854106777998746>
- Rae, R., Verdun, C., Grewal, P. S., Robertson, J. F., & Wilson, M. J. (2007). Biological control of terrestrial molluscs using *Phasmarhabditis hermaphrodita*-progress and prospects. *Pest Management Science*, 63(12), 1153–1164. <https://doi.org/10.1002/ps.1424>
- Ross, J. L., Pieterse, A., Malan, A. P., & Ivanova, E. (2018). *Phasmarhabditis safricana* n. sp (Nematoda: Rhabditidae), a parasite of the slug *Deroceras reticulatum* from South Africa. *Zootaxa*, 4420(3), 391–404. <https://doi.org/10.11646/zootaxa.4420.3.5>
- South, A. (1992). *Terrestrial slugs: Biology, ecology and control*. Chapman and Hall.
- Speiser, B., Zaller, J. G., & Neudecker, A. (2001). Size-susceptibility of the pest slug *Deroceras reticulatum* and *Arion lusitanicus* to the nematode biocontrol agent *Phasmarhabditis hermaphrodita*. *Biocontrol*, 46(3), 311–320. <https://doi.org/10.1023/A:1011469730322>
- Tan, L., & Grewal, P. S. (2001a). Infection behaviour of the rhabditid nematode *Phasmarhabditis hermaphrodita* to the grey garden slug *Deroceras reticulatum*. *Journal of Parasitology*, 87(6), 1349–1354. [https://doi.org/10.1645/0022-3395\(2001\)087\[1349:IBOTRN\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2001)087[1349:IBOTRN]2.0.CO;2)

- Tan, L., & Grewal, P. S. (2001b). Pathogenicity of *Moraxella osloensis*, a bacterium associated with the nematode *Phasmarhabditis hermaphrodita*, to the slug *Deroceras reticulatum*. *Applied Environmental Microbiology*, 67(11), 5010–5016. <https://doi.org/10.1128/AEM.67.11.5010-5016.2001>
- Tan, L., & Grewal, P. S. (2002). Endotoxin activity of *Moraxella osloensis* against the grey garden slug, *Deroceras reticulatum*. *Applied and Environmental Microbiology*, 68(8), 3943–3947. <https://doi.org/10.1128/aem.68.8.3943-3947.2002>
- Tandingan De Ley, I., Holovachov, O., McDonnell, R. J., Bert, W., Paine, W., & De Ley, P. (2016). Description of *Phasmarhabditis californica* n. sp. and first report of *P. papillosa* (Nematoda: Rhabditidae) from invasive slugs in the USA. *Nematology*, 18(2), 175–193. <https://doi.org/10.1163/15685411-00002952>
- Tandingan De Ley, I., Schurkman, J., Wilen, C., & Dillman, A. R. (2020). Mortality of the invasive white garden snail *Theba pisana* exposed to three US isolates of *Phasmarhabditis* spp. (*P. hermaphrodita*, *P. californica*, and *P. papillosa*). *PLoS One*, 15(1), e0228244. <https://doi.org/10.1371/journal.pone.0228244>
- Tandingan De Ley, I. T., McDonnell, R. D., Paine, T. D., & De Ley, P. (2017). *Phasmarhabditis*: The slug and snail parasitic nematodes in North America. In M. M. M. Abd-Elgawad, T. H. Askary, & J. Coupland (Eds.), *Biocontrol agents: Entomopathogenic and slug parasitic nematodes* (pp. 560–578). CABI Publishing.
- White, G. F. (1927). A method for obtaining infective nematode larvae from cultures. *Science*, 66 (1709), 302–303. <https://doi.org/10.1126/science.66.1709.302-a>
- Williams, A. J., & Rae, R. (2015). Susceptibility of the Giant African snail (*Achatina fulica*) exposed to the gastropod parasitic nematode *Phasmarhabditis hermaphrodita*. *Journal of Invertebrate Pathology*, 127, 122–126. <https://doi.org/10.1016/j.jip.2015.03.012>
- Wilson, M. J., Glen, D. M., George, S. K., & Hughes, L. A. (1995). Biocontrol of slugs in protected lettuce using the rhabditid nematode *Phasmarhabditis hermaphrodita*. *Biocontrol, Science and Technology*, 5(2), 233–242. <https://doi.org/10.1080/09583159550039954>
- Wilson, M. J., Hughes, L. A., Hamacher, G. M., & Glen, D. M. (1993). The rhabditid nematode *Phasmarhabditis hermaphrodita* as a potential biological control agent for slugs. *Biocontrol Science and Technology*, 3(4), 503–511. <https://doi.org/10.1080/09583159309355306>
- Wilson, M. J., Wilson, D. J., Aalders, L. T., & Tourna, M. (2016). Testing a new low-labour method for detecting the presence of *Phasmarhabditis* spp. In slugs in New Zealand. *Nematology*, 18(8), 925–931. <https://doi.org/10.1163/15685411-00003005>
- Yang, J.-S., Nam, H.-J., Seo, M., Han, S. K., & Choi, Y. (2011). OASIS: Online application for the survival analysis of lifespan assays performed in aging research. *PLoS One*, 6(8), e23525. <https://doi.org/10.1371/journal.pone.0023525>