

Classical Period Floodwater farming and quarrying at Jabal Hamra Arlbieg in the Jordanian desert: economic support for the Faynan Orefield

Hannah Friedman¹, Russell Adams², James Anderson³, Patrick Byrne⁴, David Gilbertson⁵, John Grattan⁶, Keith Haylock⁶, Lindsay Holman², Chris O. Hunt^{4*} and Harry Toland⁶

1. Department of Classics, Modern Languages and Literatures, Texas Technical University, Lubbock TX 79409, USA

2. Department of Anthropology, University of Waterloo, Ontario N2L 3G1, Canada

3. North Island College, Vancouver Island, British Columbia V9W 8C1, Canada

4. School of Biological and Environmental Sciences, Liverpool John Moores University, Byrom Street, Liverpool L3 3AH, UK

5. School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK, and Department of Archaeology and Anthropology, Bournemouth University, Dorset BH12 5BB, UK.

6. Department of Geography and Earth Sciences, Aberystwyth University SY23 3DB, UK

* Corresponding author: C.O.Hunt@ljmu.ac.uk

Keywords: Floodwater farming; irrigation; olives; geochemistry; quarrying; desert varnish

Abstract

Recent research has shown that the Faynan Orefield was a scene of intensive metal production during the Classical Period, but the infrastructure supporting this activity is less well known. We present evidence for previously undetected floodwater-farming and quarrying at Jabal Hamra Arlbieg, which contributed to the economic life of the Faynan complex during Classical times. The site is located on an ancient route, south of the orefield. We describe the hydrological control features of the floodwater farm and a possible place of habitation. Pollen analysis suggests that olives and cereals were cultivated. Exposures of sediment sequences containing buried walls, and ceramics both within and upon these sediments, all indicate that the activity took place during Nabatean to Late Roman/Byzantine times. Two ancient quarried areas were distinguished from natural landforms by a combination of geomorphic properties and variations in the geochemistry of the Mn-Fe rich desert varnish on long-exposed and quarried rock surfaces. The site provided uncontaminated produce and building stone to the Faynan Orefield.

1. INTRODUCTION

Over the last 40 years, a series of projects have investigated the history and functioning of the great copper-production complex centred on the Khirbat Faynan, with its associated wall, channel and field networks mapped westward along the Wadi Faynan (Fig. 1; Barker et al., 2007; Hauptmann 2007). The Faynan Orefield (Figs. 1-3) was a key location of substantial Bronze Age into early mediaeval copper production (Adams 2002, 2003, Barker et al. 2007; Hauptmann 2007; Hunt et al. 2007; Hunt and el-Rishi 2010; Kind 1965; Knabb et al. 2016; Levy et al. 2001; 2002; 2003;

Weisgerber 2006) and a major source of environmental pollution (Grattan et al, 2007; 2016; Pyatt et al. 1999). The Faynan Orefield was one of the great, long-lasting industrial powerhouses of the ancient world, and its resource demand would have been substantial. Activities on the orefield were partially sustained by the organised development and exploitation of local ore, water, stone, plants, timber, animals and food. The orefield was, however, not self-sufficient: critical resources including charcoal and food were introduced along well-known routes from the inherently wetter and more biologically-productive Jordanian Plateau and from further afield. (Fig. 1: Baierle 1993; Baierle et al. 1989; Friedmann 2013; Horsfield and Conway 1930; Hunt et al. 2007; Hunt and el-Rishi 2010; Jouffroy-Bapicot et al. 2006; Rambeau 2010).

This account describes the nature, stratigraphy, palaeoecology, operation and economic roles of the newly discovered ancient runoff-harvesting floodwater farm and quarrying site termed here JHA, after the adjoining Jabal Hamra Arlbieg (also transliterated as Alrbieg and Arlbeig: 30°35'32"N 35°27'26"E). This site supported the metal mining and processing activity on the Faynan Orefield (Fig. 1, 2).

Fig. 1 and 2 hereabouts

2. LOCAL CONTEXT

The hinterlands of the Khirbat Faynan complex comprise three terrain types (Figs. 1-3). Eastward, the Mountains of Edom are arid, rugged, with a steep, fault-guided mountain front cut by deep gorges. The mountain-front has altitudinally-zoned vegetation. The vegetation zones are similar in many aspects to those reported nearby at the Jebel esh-Shara by Davies and Fall (2001). At high altitude there is dry, often rather open Mediterranean woodland characterised by *Pinus* (pines), *Cupressus* (cypress), *Quercus* (oak), *Juniperus* (junipers), *Pistacia* (lentisk/mastic tree), and *Olea* (oleaster) grading into shrub-steppe with *Artemisia* (wormwood), *Centaurea* (knapweeds), *Pistacia* and *Juniperus*. This passes downhill into Indo-Turanian steppe characterised by *Artemisia herba-alta* (wormwood). Some deeply-incised wadis contain spring-fed oases with *Palmae* (palms), *Salix* (willow), *Populus* (poplars) and *Pteropsida* (ferns), but in wadis with ephemeral flows there is sparse scrub of *Tamarix* (tamarisk), *Nerium* (oleander), *Retama* (broom) and *Phragmites* (reeds) (Baierle et al. 1989; Baierle 1993; Hunt et al. 2007).

A transitional terrain between the mountain front steppe and lowland deserts is characterised by very rapid changes of bedrock type, geomorphology, gradient, soils, runoff, vegetation and ecology. It contains massive Quaternary alluvial fans at the mouths of the gorges in the mountain front, and complex Pleistocene sequences associated with major wadis including the Wadi Faynan (McLaren et al. 2004). The Khirbat Faynan and many associated archaeological sites are located within this transitional zone (Barker et al. 2007). Vegetation is mostly highly-degraded steppe with ephemeral *Poaceae* (grasses), *Plantago* (plantains), *Caryophyllaceae* (sandworts), *Asteraceae* (thistles, ragworts and knapweeds) and some *Haloxylon* (Amaranthaceae) scrub.

A hyper-arid desertic lowland lies to the south and west of the transition zone, reaching far to the south in the Wadi Arabah. This terrain includes the Fidan Sand Sea, large wadis prone to violent floods, and isolated bedrock outliers surrounded by flatter terrain (el-Rishi et al. 2007; Raab'a 1994; Saqqa and Mohammed 2004). The desert lowlands are characterised by often very sparse Saharo-Sudanese vegetation of *Haloxylon* scrub and ephemeral desert grasses with very occasional *Acacia* (acacia) trees. Normally this vegetation supports a very limited agricultural economy; but grazing

pressures can be locally intense (Mohammed 1999). Ancient arable agriculture in this zone has not been reported previously.

JHA lies on the margin of the transition zone and the hyper-arid lowland in steppe-desert with some ephemeral grasses, abundant *Plantago* and Caryophyllaceae and isolated *Acacia*. The catchment of the Wadi al-Māk that runs past JHA is incised into the mountain-front.

Fig. 3 hereabouts

3. FLOODWATER FARMING

The role of water harvesting in supporting ancient and modern farming in arid lands is widely reported (e.g. Ashkenazi et al. 2012; Barker 2007; Barker et al. 1995; Beckers et al. 2012; Bruins 2012; De Vries 1987; Erikson-Gini 2012; Evenari et al. 1971, 1982; Gilbertson 1986; Gilbertson et al. 1994; Gilbertson and Kennedy 1984; Hunt and Gilbertson 1998; Kamash 2012; Lawton and Wilke 1979; Lüttge 2010; Oleson 2001, 2007; Ore and Bruins 2012; Oweis et al. 2012; Prinz 1996; Prinz and Malik 2002; Shanan 2000). Water-harvesting technology exploits ephemeral arid-zone runoff for arable agriculture, animals and people and has supported populations in drylands, sometimes for more than 5000 years.

With the exception of minerals, the distribution of potentially useful resources in the terrains around the Faynan Orefield is determined by the availability of water. Known local supplies of water (Barker et al. 2007) are shown in Table 1.

Table 1 about here

4. MATERIALS AND METHODS

Satellite-imagery was used to identify localities that could have supported ancient water-harvesting in the lowlands south and south east of the Khirbat Faynan complex. During the summers of 2009, 2011 and 2014 selective ground reconnaissance on foot and in vehicles covered approximately ca. 700 km² of these selected areas (Figs. 1-3). Once located, sites were mapped using dGPS. Exposures showing the stratigraphical relationships of walls were logged. Selected locations were excavated using single-context recording, drawn and sampled.

Desert varnish on exposed rock surfaces and sediment samples sieved at 2 mm were analysed chemically with p-XRF using a Niton XLt700 Analyser with low-power (1.0W) X-ray tube, Ag anode target and Peltier-cooled Si-Pin x-ray detector. Every effort was made to present the instrument perpendicular to the rock surface - the design of the instrument ensures a standardized working distance – and at least three replicate measurements using >30 second exposures at each sample point were made to allow for small-scale variation.

Samples for analysis of elemental composition were air-dried, disaggregated and sieved (<2 mm). Analysis of grain composition and major elements was carried out using a Phillips XL30 FEG SEM equipped with energy dispersive x-ray detector (Oxford Instruments X-Sight EDSATW). Sieved soil samples were mounted on carbon film-coated aluminium stubs. Images were collected using backscattered electron imaging (BSEI). Semi-quantitative information on elemental composition of individual grains was obtained using EDS. Trace element composition of soil samples was by ICP-MS following aqua-regia digestion. Blanks, duplicates and a certified reference material (CRM051) were used. Detection limits (ppb) of the ICP-MS were Pb 0.03, Zn 0.2, Cu 0.1 and Cd 0.05. High precision of

the ICP-MS control standard is reported (3%) and recover rates for the CRM were between 95% and 103%.

Air-dry samples for granulometry were sieved at 2 mm to remove coarse particles, then disaggregated in sodium hexametaphosphate 0.5% solution. Replicates were analysed in a LS13320 Meritic laser granulometer until the trace stabilized, then a further five replications per sample were made and the means determined.

Palynological analysis used the method of Hunt et al. (2007). Samples of 5 mg dry weight were disaggregated by boiling in sodium pyrophosphate 5% solution for 20 minutes, sieved through 100 μ m and on 6 μ m nominal nylon mesh to remove coarse clasts, fines and solutes, then gravity separated using the swirling technique to remove silt. The residues were mounted in glycerin gel with an admixed safranin stain and examined at 400 and 1000x magnification on an Olympus BH2 microscope equipped with phase and Nomarski interference-contrast. All particulate organic matter including the pollen and spores were enumerated and pollen and palynofacies diagrams were drawn using TILIA software (Grimm 1991) and the conventions of Hunt and Coles (1988).

5. THE DISTRIBUTIONS OF WALLS AND CHANNELS IN THE DESERTIC LOWLANDS

Locations of ancient water-harvesting and utilisation technology and other features of interest are shown in relation to the geomorphology and surface geology in Fig. 3.

Fragments of walls or lines of boulders, sometimes curving, occur on prehistoric surfaces, upon and within Late Quaternary dunes NE of the ancient metallurgical site at Barqa el-Hatiye (BeH). These are visibly parts of prehistoric settlements (Figs. 2,3: Adams 2003; Adams et al. 2010; Hauptmann 2007). They are not appropriately adapted to the ancient topography to manage water.

Long low barrages across active wadi-braidplains were found immediately east of modern Faynan Village/Rashayadah and across the Wadi Abu Dhubban west of Barqa el-Heteye (Figs. 1-3). These are modern constructions.

Wall-and-channel networks identified previously west of the Khirbat Faynan complex have spatial and topographic properties associated with runoff-farming (Figs. 2, 3; see Crook 1999; 2009; Grattan et al. 2007; Hunt and El Rishi 2010; Newsom et al. 2007). Some areas were supplied by an aqueduct from the lower Wadi Ghuwayr (Fig. 3). The new survey (Fig. 3) indicates that these and other smaller wall-networks nearby are located on Late Quaternary alluvial fans and fluvial sequences (McLaren et al. 2004).

Low, isolated walls, or groups of walls, sometimes orientated oblique to the ground slope and possibly associated with rainwater harvesting, were located on satellite-imagery and confirmed by ground survey on the rocky lower slopes of Umm 'Ishrin Sandstones at six isolated jabal within the desertic lowlands. All were less than 5 km from the mountain front and located to the north and south of Wadi Burūs (Fig. 3). Apart from JHA, the most complex of these small wall networks is on the east side of the Jabal Burūs close to where Wadi Burūs leaves the Jabal. The network at Jabal Burūs is rectilinear in overall plan, but includes small curvilinear walls (or circular enclosures) frequently one course high. These walls were undateable. They might have formed hillside micro-catchments, but no water-storage features were identified. Such enclosures, if augmented by

thorny branches, might be temporary gathering-points for animals grazing the marginal desertic lowlands during the winter vegetation-flush.

The clearest site of rainwater harvesting was detected at the foot of the north-facing slopes of the Jabal Hamra Arlbieg (Figs. 1 to 5). This hill is a geological and topographic outlier of the lower part of the Umm 'Ishrin Sandstone Formation (Raab'a 1994) that characterizes the mountain front, from which it is separated by the Malqa Fault. This location is Site BLS 87 of the Barqa' Landscape Survey (Adams et al. 2010).

6. FLOODWATER CONTROL AT JHA

Primary functional features of this floodwater farm are shown on Figs. 4-6 and Table 2. These are the characteristic properties of a floodwater diversion farm as defined by Beckers et al. (2012).

Table 2 hereabouts

Figs. 4-6 hereabouts

Various other walled enclosures, some very small, and of unclear origins and associations occur on the colluvium on the slopes that lead to the Jabal. Network 5 is an assemblage of fragments of walls and graves (GA-RC).

Bedforms in the flow routes shown in Fig. 4 indicate the wall-and-channel networks are still intermittently active. The locations and patterns of walls of wall-and-channel Networks 1, 2 and 3 and the outflow appear to reflect the distribution of part-buried fluvial bar-tops of likely Pleistocene or Early Holocene age (cf. McLaren et al. 2004). Alluvial sediments collected within distinct shallow sedimentary basins between these landforms and the Jabal, impounded by the wall-networks.

During the sustained heavy rainstorms of May 2014 (Farhan and Anbar 2014; Times of Israel Staff 2014, May 8) surface waters leaving this ancient floodwater farm through point 113 on Fig. 4 rejoined the flows within the unmanaged wadi-channels slightly to the north of the boundary wall. Their combined discharge was sufficient to create a very large lake covering ~40,000 m² (with much deposited silt and fine sands) that accumulated temporarily at the modern earthen barrage in the Wadi Abu Dhubban, approximately 15 km further west (Fig. 2).

7. CERAMICS

The ceramics comprise more than 2200 sherds found across JHA. There are small surface scatters of sherds and debris of the Early Bronze Age and Iron Age. Most sherds were of Nabatean-Roman age. Across JHA, potsherds of Nabatean and Roman age occur upon and within a distinctive widespread sand body (facies 2L and 2) associated with the construction and use of the wall-and-channel networks. Network 4 provided significant quantities of Nabatean Fine Ware sherds; one sherd of Nabatean Painted Fire Ware Dekophase 4 that dates to the 2nd and 3rd centuries C.E., unpainted Nabatean pottery, one 1st to 2nd century C.E. flanged rim jug (Schmid 2000; Gerber 2001; 360 fig. 1:17); one body sherd of unguentarium; one triangular rim cooking pot with hooked rim; and one with bulbous rim of the 3rd and 4th centuries C.E. Surface finds at JHA were of Late Roman/Byzantine date (4th through 6th centuries C.E.); and some distinctive Early Islamic blue-glazed pottery (8th to 10th centuries C.E.). A few sherds of Late Islamic date (15th through to 17th century C.E) occur at the floodwater entrance to JHA at the Wadi al Māk (Network 4) and near the

final exit-point of this impounded water (112, Fig. 4). The main development and use of the farm was thus in Nabatean-Roman times, with some continuity of use into the Late Roman/Byzantine and sporadic activity thereafter (Adams et al. 2010. Holman unpublished).

Ceramic evidence for domestic occupation was only found at BAS (features 1 to 9 - Fig. 4) where the assemblage was dominated by Nabatean and Early Roman fine-ware - mainly cooking or storage vessels. These finds include one triangular ribbed necked jar manufactured in Petra during the 1st to 3rd centuries AD (see Gerber 2008; 336, fig 22.15). Most pottery was made in Petra, although some was "Aqaba Ware" of Dolinka (2003: 79-90) from Aila (modern 'Aqaba).

8. STRATIGRAPHY AND SEDIMENTATION

Figs. 7-10 and Tables 3-6 summarise the lithostratigraphic and geochemical evidence at TR1 and TR3 (Network 1) and TR12 (Network 2: Fig. 4). The stratigraphy proved to be essentially similar in other trial trenches. Three widespread sedimentary facies (1, 2 and 3) and several more localised sub-facies were distinguished (Table 3).

[Tables 3-6 about here](#)

[Figs. 7, 8 about here](#)

The wall networks are partly built on the gravels of Facies 3 and the palaeosol 3r and thus postdate them. Some channels e.g. at 111 in Network 2 (Fig. 5) are cut into Facies 3 gravels. Sediments of facies 2, 2L and 2r abut some walls, but other walls were constructed on Facies 2 sands as they aggraded, for instance in TR1 and TR12 (Figs. 7-10, Tables 4-6). The channel-network is infilled with sediments of facies 2 and 2L. Facies 2 and 2L have granulometry and bedding suggestive of fluvial deposition (Fig. 9, Table 3). They contain potsherds of Nabatean-Roman age e.g. TR5 at Q1 (Table 6), suggesting that the farm was built and added to during the Nabatean-Roman interval as Facies 2, 2L and 2r accumulated. The possibility of greater antiquity cannot be eliminated for parts of the wall-and-channel networks, and parts of facies 2L and 2.

Facies 1 drapes over walls and earlier stratigraphic elements and thus post-dates them. It contains no indication of local human activity. The mixture of aeolian and fluvial deposition in this unit indicates a complex history, but there is no dating evidence other than stratigraphic position. Similar deposition occurs today in the region (Hunt et al. 2007; Touchan and Hughes 1999).

9. SEDIMENTARY HEAVY METALS

Archaeological evidence of past metallurgical activity was not detected upon Jabal Hamra Arlbieg, nor upon or within the alluvial sequences and wall-and-channel networks of JHA. Very occasionally, small clasts containing veins of green copper ores occur in the alluvial deposits. These were likely introduced by floodwaters from upstream sources within the Umm 'Ishrin Sandstone – this is also the situation today.

Detailed geochemical work in TR1 and TR3 sought to identify contamination from former local metallurgical activity (Figs. 9,10). Concentrations of heavy metals detected in Facies 3 are regarded as the local background because these coarse clastics are attributed to the Pleistocene or Early Holocene and pre-date the wall-and-channel network or any known ancient metallurgy in the area. Facies 2 and 2L, attributed to Nabatean-Roman times, also have low concentrations of metals. (cf.

Grattan et al. 2007, 2014, 2016). Facies 1 at TR1 had slightly higher, but still relatively low concentrations of cadmium, copper, lead, silver, tin and zinc. These concentrations are notably lower than those associated with Nabatean-Roman layers in the Faynan Orefield (Grattan et al. 2007, 2013, 2016 Knabb et al. 2016). Low concentrations of heavy metals in the alluvial sediments at JHA reflect natural processes. Plants and animals there would have been uncontaminated.

Fig. 9 hereabouts

Fig. 10 hereabouts

10. CROPS AND VEGETATION MANAGEMENT

The sediments at JHA are silty sands (Fig. 8) resulting from the application of irrigation waters. Analysis of major elements suggests predominance of Calcium, probably in the form of Calcium carbonate (Fig. 10). As such, the sediments would have accommodated considerable irrigation water, as long as drainage was impeded.

Pollen assemblages from TR3 (Fig. 11) are characterised by high frequencies of Caryophyllaceae (sandwort family), *Plantago* (plantains), and *Pinus* (pine) together with a wide diversity of herbs that are typical markers of the heavily-grazed dry steppe in the region today (Mohammed 1999; Hunt et al. 2007). These assemblages are consistent with intensive grazing in the past, as today.

Fig. 11 hereabouts

The pollen of plateau taxa remain broadly stable and in higher percentages than found in sites close to Khirbat Faynan (Hunt et al. 2007; Hunt and El-Rishi 2010). This suggests localised fuelwood extraction forest depletion for fuelwood on the plateau above the Khirbat Faynan did not extend as far south as JHA.

The percentages of *Olea* (olive) and *Cereal* (cereals, barley where identifiable) are large enough to suggest that these were grown at JHA. Both taxa are relatively low pollen producers, so the regional signatures of steppe and plateau taxa are stronger than those of the cultivated species. Olives, with their deep root systems, are likely to have been well-suited to the alluvial deposits (cf. Gilbertson et al. 1994:235, Fig. 13.6). The cereals may have been grown on the clay/silt-rich colluvium at the edges of the Jabal. The pollen evidence is incompatible with the site being used to grow forage, as suggested for the field systems at Khirbat Faynan by Hunt and el-Rishi (2010). Animals might, however, have been fed on crop waste, as occurred elsewhere during the Roman period (Hunt et al. 2001). The rapidly-decreasing *Acacia* percentages suggest that this thorny tree may have declined over time around JHA. Perhaps its wood was used as fuel locally or at the Khirbat Faynan complex (Hunt et al. 2007; Miller and Marston 2012).

The similarity of assemblages suggests that Facies 3a was laid down during the same episode of farming and immediately before Facies 2, and that Facies 1 at TR3 followed without a significant hiatus. The very high percentages of thermally mature material, derived from combustion, are typical of regional Nabatean-Roman period assemblages (Hunt and el-Rishi 2010; C.O. Hunt, unpublished data).

11. QUARRIED ROCKFACES

The frequency of ashlar-built construction in and around the Khirbat Faynan indicates that the ancient quarrying of bedrock was common in this region, but sources are not well-known. Ancient quarried rock faces were identified at Q1 and Q2 on the east-facing slopes at JHA adjacent to wall-and-channel networks 1 and 3 (Figs. 5, 12-14). A flat area in front of the more extensive quarried rockface at Q1 is bounded to the east by a semi-circular wall (Fig. 5). Virtually all the pottery from within this bounded area was manufactured from 1st to 3rd centuries CE in Petra. There is one Nabatean fine ware low-ring jug of the 1st Century C.E. The pottery is consistent with quarrying occurring during the 1st to 3rd centuries CE.

Figs. 12-14 hereabouts

It is important to distinguish between quarried rock surfaces and those produced by natural processes of rock failure in this rugged terrain. Bedrock-failures occur common in this region, and slab failures, especially down-dip, happen on Umm 'Ishrin Sandstone. Except for the major landslips mapped on such bedrocks in the Wadis Dana and Salawan by Barjous (1992), such events have not been observed at JHA, where bedding is ~horizontal. The differences between natural and quarried surfaces at Q1 are set out in Table 7, and Figs. 12 – 14.

Table 7 about here

At Q1, variations in the colour and extent of desert varnish (Cooke et al. 1993) led to the initial recognition of quarried rockfaces. The variations in the concentrations of Manganese (Mn) and Iron (Fe) are responsible for these colour differences – lighter quarried faces, and the dark un-quarried rock surfaces – are shown in Figs. 12-14). Petrological examination indicated that variations in colour and geochemistry did not reflect inherent variations in bedrock lithology. Natural hillslope rock surfaces, measured by p-XRF, varied between 6,385 to 17,543 ppm Mn and 8,812 to 18,839 ppm Fe. In contrast, vertical, pale-coloured quarried rock surfaces, where sandstone slabs had been prised away, had much lower concentrations (0 to 311 ppm Mn, and 2,820 to 8,903 ppm Fe). The volume of stone quarried from Q1 was estimated with the aid of a simple 3D model to be over 16,000 m³. In antiquity, the closest destination would have been the Khirbat Faynan.

12. HABITATION?

Sites with stone-built evidence for former human habitation at JHA are rare. Apart from the BAS with its accumulation of pottery sherds, including cooking ware of Nabatean and Early Roman age (Adams et al. 2010), there were only the assumed “footings” of one (possible) cobble-hut (101), and a few small indeterminate features on the northwest summit of the Jabal. Within the through-valley to the ESE of the Jabal, at wall-and-channel Network 5 there were some long linear fragments of walls that might mark former enclosures, as well as two zones nearby of recently robbed graves (GA/GC) with pottery fragments that suggest the Bronze Age or possibly Iron Age (Figs. 3 and 4). A circular enclosure of large boulders with dark desert varnish found at 107 to the north-east of the Wadi al Māk (Fig. 5) appears to be much older.

13. DISCUSSION

The topography, the wall-and-channel networks and associated alluvial deposits at JHA (Table 2) constitute a flood diversion scheme in the typology of Beckers et al. (2012). The networks were intended to introduce, stabilise, store and use both water and sandy/silty sediments and seem to have been well-adapted to this location. Precise contemporaneity or the progressive construction of all of the functioning parts of the farm cannot yet be demonstrated. Nevertheless, the apparent overall hydrological coherence points to its walls, channels and topographic relationships having a unity of design and purpose. The pattern of modern fluvial activity across the site suggests that the general hydrological regime that sustained JHA in antiquity continues today, long after its abandonment, as occurs on many ancient floodwater farms (Hunt and Gilbertson 1988). An advantage of this form of water control is its flexibility - waters in a year with little harvested runoff could be concentrated by selective blocking of the major channels into small areas to support a viable crop. In a high-runoff year, water could be distributed widely and allowed to exit the system. This type of floodwater farming is remarkably rugged and resilient in the face of often very adverse environments (Frasier 1983; Prinz 1996).

The general design of the JHA system and its spatial relationships between local terrain, walls and water flow recall those initially termed the “Nabataean System” of floodwater farming (Evenari and Koller 1956; Evenari et al. 1982), although these are now understood often to be older in origin. Whilst there is evidence for earlier human activity in the general vicinity of JHA, all the present information suggests the complex operated from Nabatean to late Roman or Byzantine times.

Cultivation focussed on olives, with some cereals. This reconstruction contrasts with the suggestion by Hunt and El-Rishi (2010) that the WF4 flood-water farm adjacent to Khirbat Faynan may have been largely for forage, but it chimes with the phytolith evidence of Knabb et al. (2016) which suggests the presence of tree crops in their Unit 7. Olives are resistant to episodic drought and fluvial erosion and can exploit deep soil moisture in coarser materials - impossible for herbaceous crops (Gilbertson et al. 1994). Tree crops would have provided essential shade for livestock in summer heat.

The site merits comparison with three other regional locations. In the Negev uplands, water management systems were often located within large through-wadis which contained significant water-retentive loessic soils. JHA differs with its impounded sandy alluvial infill sequences located among Pleistocene fluvial landforms in the distal parts of a massive alluvial fan, with very different topography and hydrology.

Qasr al-Tilah, 35-40 km further north, is also in transitional terrain – in this case where Wadi al-Tilah intersects with the northern end of the Wadi Arabah desert basin. This site is more elaborate than JHA, with an aqueduct, a reservoir, a fort or caravanserai, and a substantial network of raised agricultural fields on a grid pattern sustained by a network of water channels (now mostly destroyed by modern ploughing and bulldozing) but documented by early travellers and explorers, including Musil (1907: 209-214, Fig. 214 and 147ff); Frank (1934 213-215, Plate 29A and Plan 13); Alt (1935: 4); Glueck (1935: 12-17), more recently MacDonald (1992; SGNAS site 155). Although Hellenistic, Nabatean, early Roman and Byzantine pottery were collected from the site and its field system, most sherds dated to the Nabatean / early Roman period, as at JHA.

The floodwater farm at JHA has one (partial) parallel in the immediate area: wall networks WF4.1 and WF4.2 with “herring-bone patterns” near Khirbat Faynan (Fig. 2). Their modern land surfaces are likewise characterised by sherds of Nabatean - early Roman pottery (Creighton and Newsom 1999; Crook 1999, 2009; Gerber 1994; Kennedy and Bewley 2004; Newsom et al. 2007a). Nevertheless, the fundamental properties of their relationships with a far more subdued

topography, different surficial geology, soils/sediments, hydrological regime and primary water supplies all differ. These Faynan networks had much greater longevity than that at JHA.

The comparative sites mentioned above all differ from JHA because they are much larger. JHA is relatively small and does not seem to have supported many people. It is likely that it exported produce and building stone, rather than being a net importer of food, as was the Khirbat Faynan complex (Hunt and el-Rishi 2010). Export of materials would have been possible via the ancient road connecting the Khirbat Faynan with the Umm al Amad mines, which runs adjacent to the farm (Fig. 1). Food from JHA would have been healthy because heavy metal contamination of the site was very low, unlike the levels of contamination in the WF4 farming system adjacent to Khirbat Faynan (Pyatt et al. 1999; Grattan et al. 2007; Hunt & el-Rishi 2010; Perry et al. 2010) - although the ancient communities in the Faynan orefield may not have been aware of this. Certainly, it can be hypothesised that the yields per hectare at JHA would have been better than at WF4, where metal contamination still leads to stunted plants (Pyatt et al. 1999). It can therefore be concluded that the JHA farm was an important part of the support network of the Khirbat Faynan industrial complex.

14. CONCLUSIONS

Exploration south of Khirbat Faynan examined >700 km² of hot desert lowlands and the lower slopes of the Mountains of Edom and discovered JHA, a floodwater farm of the flood-diversion type that extended ≥1200 m². No other unambiguous indications of ancient floodwater farming or water-harvesting were detected in the survey area. The farm was constructed adjacent to the Wadi al Māk, from which it abstracted floodwaters. It was constructed in relatively sheltered topography at the foot of an isolated hill close to where Wadi al Māk left its gorge in the mountain front.

The patterns of inlets, channels, impluvium, walls and outflow at JHA relate to the local topography and hydrology, which in turn reflect the local surficial and bedrock geology. The construction and use of (at least) parts of its main wall-and-channel networks appears to have essentially contemporary. Walls and channels were built between and/or upon coarse fluvial Pleistocene or Early Holocene sediments and are interbedded with – and appear to have confined – a distinctive and widespread stratum of waterlain sands. Within the irrigated area, olives were grown on the alluvium, especially where underlain by coarse clastics. Cereals may have been cultivated on finer-grained sediments/soils. The farm appears to have been constructed during Nabatean-Roman times, with a continuation of use into the Late Roman/Byzantine. There is minor evidence for habitation at the site, suggesting that produce was sent elsewhere for consumption. Two areas of Umm 'Ishrin Sandstone quarried in antiquity were apparently associated with the Farm. It is possible that quarried ashlar went for construction at the Khirbat Faynan. The floodwater farm is immediately adjacent to an ancient route that links the Khirbat Faynan with the nearby copper mines at Umm al Amad in the Mountains of Edom. Both of these communities may have also been supported by the activities at JHA, as perhaps were transport and communications between them.

15. ACKNOWLEDGEMENTS

The authors thank the Department of Antiquities of Jordan, and especially the late Dr. Fawwaz Kraysheh for their support of the Barqa Landscape Project. The project was funded during the 2009 and 2010 seasons by a Standard Research Grant (No. 410-2008-0783) from the Social Sciences and Humanities Research Council of Canada to Russell Adams (PI, then at McMaster University), and

during the 2013 field season by a University of Waterloo/SSHRC 4A Award (SSHRC #410-2012-1462). The authors thank staff and student volunteers involved in the project. We thank the American Center for Oriental Research in Amman and its Director Dr. Barbara Porter and administrator Mrs. Kathy Nimri for their continued support and assistance, Professor David Kennedy and the Aerial Photographic Archive for Archaeology in the Middle East; and Antony Smith for the creation of the diagrams. Finally, we thank two anonymous reviewers for their constructive comments, which have improved this paper.

15. REFERENCES

- Adams, R. 2000. The Early Bronze Age III–IV transition in southern Jordan: evidence from Khirbat Hamra Idfan. In Philip, G. and Baird, D. (eds). *Ceramics and Change in the Early Bronze Age of the Southern Levant*, Sheffield, Sheffield Academic Press, 379–402.
- Adams, R. 2002. From factories to farms: the development of copper production at Faynan, southern Jordan, during the Early Bronze Age. In B. S. Ottaway and Wager, E.C. (eds.). *Metals and Society*. Oxford, British Archaeological Reports, International Series S1061; Archaeopress, 21–32.
- Adams, R. 2003. External influences at Faynan during the Early Bronze Age: a re-analysis of Building 1 at Barqa el-Hetiye, Jordan. *Palestine Exploration Quarterly* 135, 6–21.
- Adams, R.B, Anderson, J.D., Grattan, J.P., Gilbertson, D.D., Rouse, L., Friedman, H.A., Homan, M.M. and Toland, H. 2010. First Season of the Barqa Landscape Survey, South West Jordan. *Annual of the Department of Antiquities of Jordan* 54, 95-120.
- Alt, A. 1935. Aus der 'Araba II. Römische Kastelle und Strassen. *Zeitschrift des Deutschen Palästina-Vereins* 58, 1-59.
- Ashkenazi, E., Avni, Y. and Avni, G. 2012. A comprehensive characterization of ancient desert agricultural systems in the Negev Highlands of Israel. *Journal of Arid Environments* 86, 55-64.
- Baierle, H. U. 1993. *Vegetation und Flora im Sudwestlichen Jordanien*. Diss Bot 200. Stuttgart, Cramer.
- Baierle, H. U., Frey, W., Jagiella, C. and Kurschner, H. 1989. Die Brennstoffressourcen im Raum Fenan Wadi Araba, Jordanien und die bei der K pfererzverh ttung Verendeten Brennstoffe. In Hauptmann, A., Pernicka, E. and Wagner, G.A. (eds). *Arch ometallurgie der Alten Welt*, 213–22. Der Anschnitt Beiheft 7. Bochum, Selbstverlag des Deutschen Bergbau-Museums.
- Barjous, M.O. 1992. The Geology of the Ash Shawbak Area. Map Sheet No. 3151 111. Amman, Bulletin 19; Geology Directorate, Geological mapping Division, The Hashemite Kingdom of Jordan, Ministry of Energy and Mineral Resources, Natural Resources Authority, 82pp.
- Barker, G., Gilbertson, D. and Mattingly, D. (eds.). 2007. *Archaeology and Desertification: the Wadi Faynan Landscape Survey, Southern Jordan*. Oxford, Council for British Research in the Levant and Oxbow Books.
- Beckers, B., Berking, J. and Schott, B. 2012. Ancient Water Harvesting Methods in the Drylands of the Mediterranean and Western Asia. *eTopoi* 2, 145-164.
- Bienkowski, P. and Galor, K. 2006. (eds.). *Crossing the Rift. Resources, Routes, Settlement Patterns and Interaction in the Wadi Arabah*. Oxford: Council for British Research in the Levant/Oxbow.

- 459 Bruins, H.J. 2012. Ancient desert agriculture in the Negev and climate-zone boundary changes
460 during average, wet and drought years. *Journal of Arid Environments* 86, 28-42.
- 461 Bryan, K. 1929. Floodwater farming. *Geographical Review* 19, 444–56.
- 462 Cooke, R., Warren, A. and Goudie, A. 1993. *Desert Geomorphology*. London, UCL Press.
- 463 Creighton O.H. and Newson, P. 1999. Water management practices in the WF4 field system. In
464 Barker, G., Adams, R, Creighton, O. H., Crook, D., Gilbertson, D. D., Grattan, J. P., Hunt, C. O.,
465 Mattingly, D. J., McLaren, S. J., Mohammed, H. A., Newson, P., Palmer, C., Pyatt, F. B., Reynolds T. E.
466 G. and Tomber, R. *Environment and land use in the Wadi Faynan, southern Jordan: the third season*
467 *of geoarchaeology and landscape archaeology* (1998). *Levant* 31, 255–292.
- 468 Crook, D. 1999. Palaeohydrology. In Barker et al. *Levant* 31, 278-291.
- 469 Crook, D. 2009. Hydrology of the combined irrigation system in the Wadi Faynan, Jordan. *Journal of*
470 *Archaeological Science* 36(10), 2427-2436.
- 471 De Vries, B. 1987. The el-Lejjūn Water System. In Parker, S.T. (ed.). *The Roman Frontier in Central*
472 *Jordan: Interim Report on the Limes Arabicus Project 1980–1985*. Oxford, British Archaeological
473 *Reports International Series* 340 1, 399–428.
- 474 Dolinka, B. 2003. Nabataean Aila (Aquaba, Jordan) from a Ceramic Perspective. Oxford,
475 Archaeopress, *British Archaeological Reports International Series* 1116.
- 476 El-Rishi, H., Hunt, C., Gilbertson, D., Grattan, J., McLaren, S., Pyatt, B., Duller, G., Gillmore, G.,
477 Phillips, P. 2007. Chapter 3. The past and present landscapes of the Wadi Faynan:
478 geoarchaeological approaches and frameworks. In Barker, Graeme, Gilbertson, David and
479 Mattingly, David. (eds.). 2007. *Archaeology and Desertification: the Wadi Faynan Landscape*
480 *Survey, Southern Jordan*. Oxford, Council for British Research in the Levant & Oxbow Books, 59-95.
- 481 Engel, T. 1992. Nouvelle recherches sur les charbons du Chalcolithique, de l'âge de Bronze, et de
482 l'âge du Fer de Fenan / Wadi Araba, Jordanie. In J. L. Vernet (ed.). *Les Charbons de Bois, Les Anciens*
483 *Écosystèmes et le Rôle de l'Homme*, *Bulletin de la Société Botanique de France* 139; *Actualités*
484 *botaniques* 2/3/4 ; 553–63.
- 485 Engel, T. 1993. Charcoal remains from an Iron Age copper smelting slag heap at Feinan, Wadi
486 Arabah, Jordan. *Vegetation History and Archaeobotany* 2, 205–11.
- 487 Engel, T. and Frey, W. 1996. Fuel resources for copper smelting in antiquity in selected woodlands
488 in the Edom highlands to the Wadi Araba/Jordan. *Flora* 191, 29–39.
- 489 Erikson-Gini, T. 2012. Nabatean agriculture: myth and reality. *Journal of Arid Lands* 86, 50-54.
- 490 Evenari, M. and Koller, D. 1956. Ancient masters of the desert. *Scientific American* 194(4), 39-45.
- 491 Evenari, M., Shanan, L. and Tadmor, N. H. 1982 [1971]]. *The Negev: the Challenge of a Desert*. 2nd
492 edition. Cambridge, MA, Harvard University Press.
- 493 Farhan, Y. and Anbar, A. 2014. Fragile Landscape: Impact and Consequences of the May 2014 Flash-
494 flood Disaster in the Aqaba area, Southern Jordan. *Research Journal of Environmental and Earth*
495 *Sciences* 6(9), 451-465.
- 496 Frank, F. 1934. Aus der 'Araba I. Reiseberichte. *Zeitschrift des Deutschen Palästina-Vereins* 57, 191-
497 280.

- 498 Frasier, G.W. 1983. Water Harvesting for Collecting and Conserving Water Supplies. In Pathak, P.,
 499 El-Swaify, S.A., and Singh, S. (eds). Workshop on the Art and Management Alternatives for
 500 Optimising the Productivity of SAT Alfisols and related soils. 1-3 December 1983. ICRISAT Center,
 501 India, International Crop Research 67-77.
- 502 Frey, W., Gerlach, E. and Kurschner, H. 1991. Neolithische Holzkohlen aus dem Tell Wadi Fenan,
 503 Wadi Arabah, Jordanien und ihre interpretation. In Engel, T., Frey, W. and Kurschner, H. (eds.).
 504 Contributiones Selectae ad Floram et Vegetationem Orientis. Proceedings of the Third Symposium
 505 on Plant Life of Southwest Asia, Berlin 1990, Flora et Vegetatio Mundi 9. Stuttgart, Cramer, 121–32.
- 506 Friedman, H. 2013. Agriculture in the Faynan: Food Supply for Industry. In Bowman, A. and Wilson,
 507 A. (eds.). The Roman Agricultural Economy Organization, Investment, and Production. Oxford,
 508 Oxford University Press, 307-322.
- 509 Gerber, Y. 1994. Nabataean Coarse Ware Pottery. In Stucky, R.A., Gerber, Y., Kolb, B., and Schmid,
 510 S.G. 1994. Swiss-Lichtenstein Excavations at ez-Zantur in Petra 1993: the Fifth Campaign. Annual of
 511 the Department of Antiquities of Jordan 38, 271-292.
- 512 Gerber, Y. 2001. Selected Ceramic Deposits. In Fiema, Z.T. and Bikai, P.M. The Petra Church.
 513 Amman, American Center of Oriental Research Publication no. 3, 359-366.
- 514 Gerber, Y. 2008. Preliminary characterization of al-Humayma Ceramics. In Oleson, J.P., Reeves,
 515 M.B., Baker, G.S., de Bruijn, E., Gerber, Y., Nikolic, M. and Sherwood, A.N. Preliminary Report on
 516 Excavations at al-Humayma, Ancient Huwara, 2004 and 2005. Annual of the Department of
 517 Antiquities of Jordan 52, 309-342.
- 518 Gilbertson, D. D. (ed.). 1986. Runoff farming in rural arid lands. Applied Geography. "Volume 6 (1),
 519 182pp.
- 520 Gilbertson, D.D., Hunt, C.O., Fieller, N.R.J. and Barker, G.W.W. 1994. The environmental
 521 consequences and context of ancient floodwater farming in the Tripolitanian Pre-desert. In
 522 Millington, A.C. and Pye, K. (eds.). Environmental Change in Drylands: Biogeographical and
 523 Geomorphological Perspectives. Chichester, John Wiley and Sons, 229 - 251.
- 524 Gilbertson, D. and Kennedy, D. L. 1984. An archaeological reconnaissance of ancient water
 525 harvesting structures and 'wadi walls' in the Jordanian desert, north of Azraq oasis. Annual of the
 526 Department of Antiquities of Jordan 26, 151–62.
- 527 Glueck, N. 1935. Explorations in Eastern Palestine, II. The Annual of the American Schools of Oriental
 528 Research 15 (1934-1935). New Haven, CT: American Schools of Oriental Research.
- 529 Grattan, J.P., Adams, R.B., Friedman, H., Gilbertson, D., Haylock, K.I., Hunt, C.O. and Kent, M. 2016.
 530 The first polluted river? Repeated copper contamination of fluvial sediments associated with Late
 531 Neolithic human activity in southern Jordan. Science of the Total Environment 573, 247-257.
- 532 Grattan, J.P., Gilbertson, D.D., and Hunt, C.O. 2007. The local and global dimensions of
 533 metalliferous air pollution derived from a reconstruction of an eight thousand year record of copper
 534 smelting and mining at a desert-mountain frontier in southern Jordan. Journal of Archaeological
 535 Science 34, 83-110.
- 536 Grattan, J.P., Gilbertson, D.D. and Kent, M. 2013. Sedimentary metal-pollution signatures adjacent
 537 to the ancient centre of copper metallurgy of Khirbat Faynan in the desert of southern Jordan.
 538 Journal of Archaeological Science 40, 3834-3.

- 539 Grattan, J P, Gilbertson, D.D. and Waller, J.H. 2014. The geoarchaeology of “waste heaps” from the
 540 ancient mining and beneficiation of copper-rich ores in the Wadi Khalid in southern Jordan. *Journal*
 541 *of Archaeological Science*, 46, 428 – 433.
- 542 Grattan, J.P., Adams, R.B., Freidman, H., Gilbertson, D.D., Haylock, K.I., Hunt, C.O. and Kent, M.
 543 2016. The first polluted river? Repeated contamination of fluvial sediments associated with Late
 544 Neolithic human activity in southern Jordan. *Science of the Total Environment* 573, 247-257.
- 545 Grimm E.C. 1991. TILIA and TILIA*GRAPH. Illinois State Museum: Springfield.
- 546 Hashemite Kingdom of Jordan, Meteorological Department. 1988. Jordan: Climatological Data
 547 Handbook. Amman.
- 548 Hauptmann, A. 2007. The Archaeometallurgy of Copper. Evidence from the Faynan, Jordan.
 549 Heidelberg, Springer.
- 550 Haylock, K. 2011. Metalliferous properties of sediments in an Early Bronze Age settlement during
 551 the smelting and processing period in southern Jordan: the oldest anthropogenic polluted landscape.
 552 Unpublished dissertation, Institute of Geography and Earth Sciences, University of Aberystwyth,
 553 Wales.
- 554 Haylock, K. 2016. The Relationship between Iron Age Hill Forts, Roman Settlements and Metallurgy
 555 on the Atlantic Fringe. Unpublished PhD Dissertation, Department of Geography and Earth Sciences,
 556 Aberystwyth University.
- 557 Holman, L. (unpublished). Wadi Faynan Ceramics Report to Barqa’ Landscape Survey. 5pp.
- 558 Horsfield, G. and Conway, A. 1930. Historical and topographical notes on Edom. *Geographical*
 559 *Journal* 76(5), 369-390.
- 560 Hunt, C.O. 1985. Recent advances in pollen extraction techniques: a brief review. In Fieller, N.R.J.,
 561 Gilbertson, D.D. and Ralph, N.G.A. (eds.) *Palaeobiological Investigations*. British Archaeological
 562 Reports, International Series, Oxford, 266, 181-187.
- 563 Hunt, C. O. and Coles, G. M. 1988. The application of palynofacies analysis to geoarchaeology. in
 564 Slater, E. A. & Tate, J. O. (eds.) *Science and Archaeology*. British Archaeological Reports, British Series
 565 196, 473-484.
- 566 Hunt, C. O. and el-Rishi, H. 2010. Human activity in the ancient metal-smelting and farming complex
 567 in the Wadi Faynan, SW Jordan, at the desert margin in the Middle East. In Martini, I. P. and
 568 Chesworth, W. (eds.). *Landscapes and Societies: Selected Cases*. Heidelberg, Springer-Verlag, 121-
 569 134.
- 570 Hunt, C.O., el-Rishi, H.A., Gilbertson, D.D., Grattan, J.P., McLaren, S., Pyatt, F.B., Rushworth, G. and
 571 Barker, G.W. 2004. Early Holocene environments in the Wadi Faynan, Jordan. *The Holocene* 14(6),
 572 921-930.
- 573 Hunt, C.O. and Gilbertson, D.D. 1998. Context and impacts of ancient catchment management in
 574 Mediterranean countries: implications for sustainable resource use. In Johnson, P., Reid, I. and
 575 Wheater, H. (eds.). *Catchment management and resource assessment in dry areas*. Volume 2.
 576 *Hydrology in a Changing Environment*. Chichester, Wiley, 473-483.

- 577 Hunt, C.O, Gilbertson, D.D., and El-Rishi, H.A. 2007. An eight thousand year history of landscape,
578 climate and copper exploitation in the Middle East: the Wadi Faynan and the Wadi Dana National
579 Reserve in southern Jordan. *Journal of Archaeological Science* 34, 1306-1338.
- 580 Hunt, C. O., Rushworth, G., Gilbertson, D. D. & Mattingly, D. J. 2001. Romano- Libyan dryland animal
581 husbandry and landscape: pollen and palynofacies analyses of coprolites from a farm in the Wadi el-
582 Amud, Tripolitania. *Journal of Archaeological Science* 28, 351-363.
- 583 Jouffroy-Bapicot, I., Pulido, M., Baron, S., Galop, D., Monna, F., Lavoie, M., Ploquin, A., Petit, C., de
584 Beaulieu, J.-L., Richard, H. 2007. Environmental impact of early palaeometallurgy: pollen and
585 geochemical analysis. *Vegetation History and Archaeobotany* 16, 251–258.
- 586 Kamash, Z. 2012. Irrigation technology, society and environment in the Roman Near East. *Journal*
587 *of Arid Environments* 86, 65-74.
- 588 Kennedy, D. and Bewley, R. 2004. *Ancient Jordan from the Air*. London, Council for British Research
589 in the Levant, Oxford, Oxbow.
- 590 Kind, H.D. 1965. Antike Kupfergewinnung Rotem und Totem Meer. *Zeitschr Deutsch Palästina-*
591 *Verein* 81, 56-73.
- 592 Knabb, K.A., Erel, Y., Tirosh, O., Rittenour, T., Laparidou, S., Najjar, M. and Levy, T.E. 2016.
593 Environmental impacts of ancient copper mining and metallurgy: Multi-proxy investigation of
594 human-landscape dynamics in the Faynan valley, southern Jordan. *Journal of Archaeological Science*
595 74, 85-101.
- 596 Lawton, H. W. and Wilke, P. J. 1979. Ancient agricultural systems in dry regions. In A. E. Hall, G. H.
597 Cannell and H. W. Lawton (eds). *Agriculture in Semi-Arid Environments*. Ecological Studies 34.
598 Berlin: Springer-Verlag, 1–44.
- 599 Levy, T., Adams, R. B., Witten, A. J., Anderson, J., Arbel, Y., Kuah, S., Moreno, J., Lo, A. and Waggoner,
600 M. 2001. Early metallurgy, interaction, and social change: the Jabal Hamrat Fidan Jordan: research
601 design and 1998 archaeological survey: preliminary report. *Annual of the Department of Antiquities*
602 *of Jordan* 45, 159–87.
- 603 Levy, T., Adams, R. B, Hauptmann, A., Prange, M., Schmitt-Strecker and Najjar, M. 2002. Early
604 Bronze Age metallurgy: a newly discovered copper manufactory in southern Jordan. *Antiquity* 76,
605 425–37.
- 606 Levy, T., Adams, R. B., Anderson, J. D., Najjar, M., Smith, N., Arbel, Y., Soderbaum, L. and Muniz, M.
607 2003. An Iron Age landscape in the Edomite lowlands: archaeological surveys along the Wadi al-
608 Guwayb and Wadi al-Jariyeh, Jabal Hamrat Fidan, Jordan, 2002. *Annual of the Department of*
609 *Antiquities of Jordan* 47, 247–7.
- 610 Lucke, B., Roskin, J., Vanselow, K.A., Bruins, H.J., Abu-Jaber, N., Deckers, K., Lindauer, S., Porat, N.,
611 Reimer, P.J., Bäuml, R., Erickson-Gini, T., Kouki, P. 2019. Character, Rates, and Environmental
612 Significance of Holocene Dust Accumulation in Archaeological Hilltop Ruins in the Southern Levant.
613 *Geosciences* 9, 190.
- 614 Lüttge, U.R. 2010. Runoff-rainwater for Sustainable Desert Farming. In Ramawat, K.G. (ed.).
615 *Desert Plants: Biology and Biotechnology*, Heidelberg, Springer, 461-477.
- 616 MacDonald, B. 1992. *The Southern Ghors and Northeast 'Araba Archaeological Survey*. Sheffield:
617 J.R. Collis Publications.

- 618 McLaren, S. J. Gilbertson, D. D. Grattan, J. P. Hunt, C. O. Duller, G. and Barker, G. 2004. Quaternary
619 palaeogeomorphic evolution of the Wadi Faynan area, southern Jordan. *Palaeogeography,*
620 *Palaeoclimatology, Palaeoecology* 205, 131–54.
- 621 Miller, N.F. and Marston, J.M. 2012. Archaeological fuel remains as indicators of ancient west Asian
622 agropastoral and land-use systems. *Journal of Arid Environments* 86, 97-103.
- 623 Mohammed, H. 1999. The Holocene Palaeoenvironments of the Rift Margin in Southern Jordan.
624 Unpublished Ph.D thesis, University of Huddersfield.
- 625 Musil, A. 1907. *Arabia Petraea. I Moab; II Edom: Topographischer Reisebericht.* Vienna, Alfred
626 Hölder.
- 627 Newson, P., Barker, G., Daly, P., Gilbertson, D. 2007. Chapter 5. The Wadi Faynan field systems. In
628 Barker, Graeme, Gilbertson, David and Mattingly, David. (eds.). 2007. *Archaeology and*
629 *Desertification: the Wadi Faynan Landscape Survey, Southern Jordan.* Oxford, Council for British
630 Research in the Levant & Oxbow Books, 141-174.
- 631 Oleson, J. P. 2001. Water supply in Jordan through the ages. In MacDonald, B., Adams, R. and P.
632 Bienkowski, P. (eds.). *The Archaeology of Jordan.* Sheffield, Sheffield Academic Press, 603–619.
- 633 Oleson, J.P. 2007. Nabatean water supply, irrigation and agriculture: an overview. In Poltis, K.D.
634 (ed). *The World of the Nabateans.* Stuttgart, Franz Steiner Verlag, 217-249.
- 635 Ore, G. and Bruins, H.J. 2012. Design features of ancient agricultural terrace walls in the Negev
636 Desert: human-made geodiversity. *Land Degradation and Development* 23(4), 409-418.
- 637 Orloff, C.R. 2005. *The water supply and distribution system of the Nabataean city of Petra (Jordan).*
638 Cambridge
- 639 Oweis, T.Y., Prinz, D., and Hachum, A.Y. 2012. *Rainwater Harvesting for Agriculture in Dry Areas.*
640 London, CRC Press - Taylor and Francis Group.
- 641 Palmer, C., Gilbertson, D., el-Rishi, H., Hunt, C., Grattan J., McLaren, S., Pyatt, B. 2007. The Wadi
642 Faynan today: landscape, environment, people. In Barker, G., Gilbertson, D. and Mattingly, D.
643 (eds.). 2007. *Archaeology and Desertification: the Wadi Faynan Landscape Survey, Southern*
644 *Jordan.* Oxford, Council for British Research in the Levant and Oxbow Books, 25-58.
- 645 Perry, M.A., Coleman, D.S., Dettman, D.L., Grattan, J.P. and al-Shiyab, A.H. 2011. Condemned to
646 metallum? The origin and role of 4th-6th century A.D. Phoenician mining camp residents using multiple
647 chemical techniques. *Journal of Archaeological Science* 38, 558-569.
- 648 Prinz, D. 1996. Water Harvesting: past and future. In Pereira, L.S. (ed.). *The Sustainability of*
649 *Irrigated Agriculture.* Proceedings of the NATO Advanced Research Workshop, Vimeiro, 21 to 26
650 March 1994; Rotterdam, Balkema, 135-144.
- 651 Prinz, D. and Malik, A.H. 2002. *Runoff Farming.* WCA infoNET at IPTRID FAO
652 http://www.plantstress.com/articles/drought_m/runoff_farming.pdf
- 653 Pyatt, F.B., Barker, G.W., Birch, P., Gilbertson, D.D., Grattan, J.P., and Mattingly, D.J. 1999. King
654 Solomon's Miners - Starvation and Bioaccumulation? An Environmental Archaeological
655 Investigation in Southern Jordan. *Ecotoxicology and Environmental Safety* 43, 305-308.

- 656 Rabb'a, I. 1994. Geology of the Qurayqira Jabal Hamra Faddan, Map Sheet NO. 3051 II. Geological
657 Mapping Division Bulletin 28. Amman, Hashemite Kingdom of Jordan, Ministry of Energy and
658 Natural Resources Authority, Geology Directorate.
- 659 Rambeau, C.M.C. 2010. Palaeoenvironmental reconstruction in the Southern Levant: synthesis,
660 challenges, recent developments and perspectives. *Philosophical Transactions of the Royal Society*
661 *of London* 368 (1931), 5225-5248.
- 662 Saqqa, W. and Mohammed, A. 2004. Characterization of the aeolian terrain facies in the Wadi Araba
663 Desert, southwestern Jordan. *Geomorphology* 62, 63-87.
- 664 Schmid, S.G. 2000. Petra: Ez-Zantur II – Ergebnisse der Schweizerische-Lichtenstienischen
665 Ausgrabungen, 1988-1992. Teil I: Die Feinkeramik der Nabatäer: Typologie, Chronologie und
666 kulturhistorische Hintergründe. *Terra Archaeological Band II*. Mainz, Verlag Philipp von Zabern.
- 667 Shanan, L. 2000. Runoff, erosion and the sustainability of ancient irrigation systems in the Central
668 Negev Desert. In *The Hydrology-Geomorphology Interface: Rainfall, Sedimentation and Land Use*.
669 *Proceedings of the Jerusalem Conference May 1999, (IAHS Publication)*, 261, 75-106.
- 670 Times of Israel Staff 2014. Israel swept by rainiest May day on record. Times of Israel 8th May 2014
671 www.timesofisrael.com/highest-may-rainfall-ever-recorded-thursday/
- 672 Touchan, R. and Hughes, M.K. 1999. Dendrochronology in Jordan. *Journal of Arid Environments*
673 42(4), 291-303.
- 674 Weisberger, G. 2006. The mineral wealth of ancient Arabia and its use: I. Copper mining and
675 smelting at Feinan and Timna – a comparison and evaluation of techniques, production and
676 strategies. *Arabian Archaeology and Epigraphy* 17, 1-30.
- 677
- 678
- 679
- 680
- 681
- 682
- 683
- 684
- 685
- 686
- 687
- 688
- 689
- 690
- 691

CAPTIONS TO FIGURES

Fig. 1. The location of the floodwater farm at Jabal Hamra Arlbieg (JHA), the principal archaeological sites in the study area, and important routes of Classical times (after Barker et al. 2007 and Bienkowski and Galor 2006) and key topographic relationships.




Image derived from Google Earth showing location of Jabal Hamra Arlbieg and other key sites.



Fig. 3. Surface geological and geoarchaeological map, interpreted from satellite imagery and ground survey, Barjous (1992) and Raab'a (1994), showing the major floodwater farms and very small wall networks on isolated hills (outliers of mainly Cambrian bedrocks) in the desertic lowlands. This also shows (i) two modern barrages in the lowland desert drainage of the Wadi al Māk; and (ii) one across the Wadi Abu Dhubban that impounded a temporary lake >40,000 m² floored with carbonate-rich silts and sands as a result of the sustained violent rainstorm of May 2014 – much of its the feeder water from the Mountains of Edom having travelled close to or (lesser quantities) through the ancient floodwater farm at JHA (see Farhan and Anbar 2014; Times of Israel Staff 2014).




Fig. 4. The locations of the five wall-and-channel networks around Jabal Hamra Arlbieg (JHA) in relation to Quaternary alluvial fans, associated alluvial basins, major wadis, and major routes, interpreted from Google satellite imagery and ground survey, Barjous (1992) and Raab'a (1994).




Fig. 5. Stone-built structures, walls, wall-and-channel networks, ancient bedrock quarries (Q1 and the smaller unstudied Q2), “modern” ephemeral stream channels, ancient controlled water flow pathways, surface heights (italicised) derived from Google satellite imagery, and our precise topographic surveys in relation to the adjacent wadis around Jabal Hamra Arlbieg: revised from initial survey by Adams et al. (2010). CS3 and CS4 are levelled survey transects.




Fig. 6. The western part of floodwater farm JHA in a relatively sheltered basin on the northern side of the Jabal Hamra Arlbieg, photographed from immediately West of spot height 257 (metres above Jordanian Datum) on its northeast arm, looking West. The hilltop in the foreground is fractured Umm 'Ishrin Sandstone, overlooking Network 2 (to the right) and Network 1 (to the left). Largely infilled linear conduits conducted water from the bottom right into the networks of rectilinear enclosures and to the hydrological exit into the adjoining desert. [Image 100-9206].



Fig. 7. Schematic summary of the stratigraphy at TR12. Here, a palaeosol and wall running along one side of a channel overlies gravels attributed to Facies 3 and these are overlain by Facies 1. On the other side of the channel, the wall is embedded in Facies 2, suggesting that this is broadly equivalent in age to the palaeosol.




Fig. 8. Grain size analyses of sands from facies 3a, 2 and 1 at TR3. The strong peak in the medium sand is consistent with these sediments being locally derived from Umm 'Ishrin Sandstone, while the grain size distributions as a whole are consistent with localised aeolian and/or fluvatile deposition (cf. Lucke et al., 2019).




Fig. 9. Summary description of the alluvial lithostratigraphy at exposure TR1 and concentrations of selected heavy metals in the <2mm fractions of sediments, determined by p-XRF. Exposures TR2 to TR5 were broadly similar chemically.




Fig. 10. Major elements measured by EDS and heavy metals measured by ICP-MS at TR3.





Fig. 11. Pollen and palynofacies analysis of samples from TR3. Pollen is calculated as percentages of total pollen. Palynofacies is calculated as percentages of total particulate organic matter.




779 Fig. 12. Quarry Q1 in 2009 from the North. On the hillslope in the background the surface is fretted,
 780 with darker desert varnish and angular rock debris, while the quarried area (with the figures in the
 781 foreground) shows the pale brown colour of Umm 'Ishrin Sandstone, with little desert varnish,
 782 distinctive morphology and possible tool marks. In 2014, Test Pit TR5 was ~3 m behind investigator
 783 on the left side of the image




784

785 Fig. 13. Quarry Q1, from the north. Measurements are ppm of manganese (Mn) measured with p-
 786 XRF. Surfaces with desert varnish have higher values of Mn than do quarried surfaces.



787

788 Fig. 14. Quarry Q1 from the north. Measurements are ppm of iron (Fe) measured with p-XRF.
 789 Surfaces with desert varnish have higher values of Fe than do quarried surfaces.



790

791

792

793

794 TABLE CAPTIONS

795 Table 1. Sources of water in the terrains around the Faynan Orefield (after Barker et al. 2007)

Oases, springs and perennial streams in the lower parts of gorges, sometimes with water conducted by aqueduct, and stored in reservoirs;
Occasional microcatchment rainwater harvesting systems which are elsewhere associated with Nabatean "stopping points" on long-distance routes across the desert;
Very large networks of walls and channels that managed surface runoff and stream flow for agriculture (Fig. 3).

796

797

798 Table 2. Functional features of the floodwater farm at Jabal Hamra Arlbeig

Characteristics
Networks 1-3 lie in a sheltered alluvial basin within the north pointing arms of the Jabal;
a long, substantial wall and flow route (108), often built on Pleistocene linear fluvial bar-forms, separates the walls/conduits and sediments of Networks 1 – 3 from the unenclosed desert terrain;
three robust water-intakes (IT1 to IT3) abstracted torrential waters at the edge of Wadi al Māk (Fig.4);
channels (98, 99, 100) led flood waters into and through the more poorly defined, but also flood-scoured, wall-and-channel Network 4 (1-22);

water-impoundment area(s)/impluvia in wall networks walls and perhaps by a habitation site in Network 3 (98 and BAS/1-9) ponded water, allowing redistribution without inducing notable erosion;
onward distribution took place in robust channels with distinct cross-sections (93, 94, 95, perhaps 96), and a deliberately excavated flow route (111) through a Pleistocene fluvial-bar;
water and sediment would have been dispersed in larger and more elaborate wall-and-channel networks 1 (85-49) and 2 (23 – 92) sometimes with two inputs into an enclosure (70-87, 34-48). Presumably cultivation and/or grazing took place on the network alluvial infills. Excess water led (eventually) West-north west;
the outflow (112-113) into the lowland between the north-western arm of the Jabal and the end of the boundary wall (108). Such drainage is commonly considered to help mitigate any build-up of salts in soils/sediments or groundwater.

Table 3. Sedimentary facies identified at Jabal Hamra Arlbieg

Facies	Characteristics	Interpretation
3	Rounded to subangular sandy gravel, variably sorted, containing sandy lenses in places.	Braidplain sediments of later Pleistocene or early Holocene age (cf. Hunt et al. 2004; McLaren et al. 2004).
3a	Lenticular bodies of fine sands	Channel fills within braidplain sediments
3p	Reddened gravel with rootlet traces sometimes capping Facies 3	Pedogenically altered braidplain sediments
2r	Reddened sands with rootlets sometimes capping Facies 3.	Skeletal palaeosol
2	Well-sorted medium-fine sands infilling shallow basins in the surface of Facies 3 and water distribution channels. Bioturbation, deformation structures and Nabatean-Roman potsherds are sometimes present	Waterlain sediments. Bioturbation suggests that in some locations plants have episodically grown in these deposits before being covered with further sand. Deformation structures might suggest localised slumping. The presence of potsherds in these sediments suggests they accumulated during the Nabatean-Roman interval
2L	Thinly laminated, well-sorted carbonate-quartz medium to fine sands, with thin laminations, sometimes containing pebbly lenses. Typically infilling channels. Sometimes passing up into Facies 2	Waterlain sands with laminations from spreading flow. In channels pebbly lenses reflect episodes of scouring flow.
1a	Sandy gravel layer infilling a 'v' shaped feature in Facies 2	Gully fill – fluvial lag
1	Fine sands and silts overlying Facies 3 and 2.	Sediments of wind-dominated deposition with occasional surface wash

Table 4. Stratigraphy of trench TR1

Layer	Depth (m)	Sediments	Interpretation
3	>0.42	Rounded and sub-angular partly-imbricated fine sandy gravel, variably sorted	Braidplain fluvial /colluvial sediments
2	0.42-0.10	Well-sorted medium-fine sands	Flood and aeolian sediments
1a	0.12-0.10	Thin sandy gravel layer infilling a 'v' shaped feature	Fluvial lag
1	0.10-0.0	Fine sands and silts	Wind-dominated deposition with occasional surface wash

805 Table 5. Stratigraphy of trench TR2

Layer	Depth (m)	Sediments	Interpretation
3	0.40-0.33	Rounded and sub-angular partly-imbricated fine sandy gravel	Braidplain fluvial /colluvial sediments
3a	0.37-0.33	Lenticular body of fine sands (3a)	Channel-fill within braidplain sediments
2	0.33-0.09	Well-sorted medium-fine sands	Flood and aeolian sediments
1	0.09-0.0	Fine sands and silts	Wind-dominated deposition with occasional surface wash

806

807 Table 6. Stratigraphy of trench TR5 adjacent to Q1

Layer	Depth (m)	Sediments	Interpretation
2	0.50-0.36	Well-sorted medium-fine sands, with on its surface sherds of Nabatean/Early Roman wares	Flood and aeolian sediments, with traces of human activity on its surface
1	0.36-0.0	Fine sands and silts	Wind-dominated deposition with occasional surface wash

808

809 Table 7. Criteria used distinguish between quarried bedrock faces and original terrain on Umm
810 'Ishrin Sandstone at Jabal Hamra Arlbieg.

Criteria	Untouched, natural rock surface	Quarried bedrock to 8m above base of slope
Morphology	Rectilinear "gaps" typically only a few metres high or deep, associated with the particular strike and dip of the outcrop of a particular stratum, or small a group of strata.	Substantial arm-chair, basin-shaped, morphology with relatively steep sides, and steep and high backslope, and whose overall form bears only limited, if any relationship to thickness, strike and dip of the strata.
Slopes	Slopes of ~25°-35° throughout. There is no upper break of slope.	Slopes of 70° to almost vertical. The quarried face intersects with the original hillslope at a sharp angle.
Basal slope morphology	Basal slopes reduce rapidly in gradient at base, from c. 25°, to c. 5°.	The gradient remains near vertical, its base at the ground surface may be close to a right angle.
Basal slope materials	Basal slope formed of angular rock debris, typically with desert varnish, and fine-grained surficial sediments.	Exposure of bedrock, revealing ~horizontal to gently-dipping bedding planes and ~vertical joints.
Materials beyond the basal slope	Cobbles and boulders frequent and dipping downslope.	A quasi-horizontal surface often free of rock debris, of fine-grained alluvium, with isolated cobbles, stones, boulders, and angular slabs of bedrock that appear to have fallen from above on to the surface.
Desert varnish	Extensive desert varnish from dark brown to near black on 30-100% of rock surface and on some of the rock debris upon it.	Fresh rock surfaces, little or no desert varnish; where present brown to pale-brown.
Surface texture	Rock surface is fretted, rough, with pits or tafoni, with many sharp edges - the result of various forms of physical and chemical weathering. The micro-topography of the intersections between	Overall rock surface is often relatively smooth, but with a rectilinear "sharp" micro-topography where ashlar were removed, exploiting joints and bedding planes.

811

	bedding planes and the joint structure is frequently slightly rounded.	
--	------------------------------------------------------------------------	--