



## LJMU Research Online

**Heikura, IA, Stellingwerff, T and Areta, JL**

**Low energy availability in female athletes: From the lab to the field.**

<http://researchonline.ljmu.ac.uk/id/eprint/15167/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Heikura, IA, Stellingwerff, T and Areta, JL (2021) Low energy availability in female athletes: From the lab to the field. European Journal of Sport Science. ISSN 1536-7290**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

2

3 **Low Energy Availability in Female Athletes: From the Lab to**  
4 **the Field**

---

5

6

7 Ida A. Heikura<sup>1,2</sup>, Trent Stellingwerff<sup>1,2</sup>, Jose L. Areta<sup>3</sup>

8

9 <sup>1</sup>*Canadian Sport Institute - Pacific, Victoria, British Columbia, Canada;*

10 <sup>2</sup>*Exercise Science, Physical & Health Education, University of Victoria British Columbia, Canada;*

11 <sup>3</sup>*Research Institute for Sport and Exercise Sciences, Liverpool John Moores University*

12

13 **Journal:** European Journal of Sport Sciences

14

15 **Running title:** *Energy deficiency in females*

16

17 **Abstract:** 250 words

18 **Word count:** 5102 words

19 **Figure & tables:** 1 table and 2 figures

20 **References:** 60/60

21

22 **Address for correspondence:**

23 Ida Heikura

24 [iheikura@csipacific.ca](mailto:iheikura@csipacific.ca)

25

26

27 **ORCID:**

28 **Ida Heikura** - 0000-0002-1088-428X

29 **Trent Stellingwerff** – 0000-0002-4704-8250

30 **José L Areta** - 0000-0001-6918-1223

31 **Abstract**

32 Decades of laboratory research have shown impairments to several body systems after only 4-5 days of strictly  
33 controlled consistent low energy availability (LEA); where energy availability (EA) = Energy Intake (EI) –  
34 Exercise Energy Expenditure (EEE) / Fat-Free Mass. Meanwhile, cross-sectional reports exist on the  
35 interrelatedness of LEA, menstrual dysfunction and impaired bone health in females (the Female Athlete  
36 Triad). These findings have demonstrated that LEA is the key underpinning factor behind a broader set of  
37 health and performance outcomes, recently termed as Relative Energy Deficiency in Sport (RED-S). There is  
38 utmost importance of early screening and diagnosis of RED-S to avoid the development of severe negative  
39 health and performance outcomes. However, a significant gap exists between short-term laboratory studies and  
40 cross-sectional reports, or clinically field-based situations, of long-term/chronic LEA and no definitive,  
41 validated diagnostic tests for RED-S exist. This review aims to highlight methodological challenges related  
42 to the assessment of the components of EA equation in the field (e.g. challenges with EI and EEE measures).  
43 Due to the uncertainty of these parameters, we propose the use of more chronic “objective” markers of LEA  
44 (i.e. blood markers). However, we note that direct extrapolations of laboratory-based outcomes into the field  
45 are likely to be problematic due to potentially poor ecological validity and the extreme variability in most  
46 athlete’s daily EI and EEE. Therefore, we provide a critical appraisal of the scientific literature, highlighting  
47 research gaps, and a potential set of leading objective RED-S markers while working in the field.

48

49 **Keywords:** low energy availability, relative energy deficiency in sport, diagnosis, female athletes

50

51 **1. Introduction**

52 The associations between low energy availability (LEA)/eating disorders, menstrual dysfunction, and impaired  
53 bone health in female athletes were established more than three decades ago when it was discovered that female  
54 athletes reporting with menstrual dysfunction appeared to also suffer from low bone density and underlying  
55 eating disorders [1]. These early discoveries were followed by an official recognition of the condition as the  
56 female athlete triad (the Triad) in the 1990's [2], which was further updated in 2007 to emphasize that it was  
57 a continuum from health to disease and that the presence of any one of the three symptoms poses a risk for the  
58 female athlete [3], of which LEA is the underlying etiology. Energy availability (EA) is defined as dietary  
59 energy intake (EI) minus exercise energy expenditure (EEE) corrected for fat-free mass (FFM) [4]. Subsequent  
60 strictly controlled laboratory interventions in sedentary females established links between 4-5 days of an EA  
61 of <30 kcal/kg FFM/d and impaired endocrine marker concentrations, thus defining this as a threshold for LEA  
62 [4]. However, while the concept of EA is scientifically sound in well-controlled laboratory settings, applying  
63 it directly in the field is challenged by several methodological considerations, mainly to do with the large  
64 variability and poor accuracy of field-based EI and EEE assessments.

65 In 2014, these identified LEA impairments originally recognized from Triad research were known to extend  
66 across other body systems and functions, and both females and males, identified as Relative Energy Deficiency  
67 in Sport (RED-S) [5, 6]. Here, RED-S was defined as “impaired physiological function including, but not  
68 limited to, metabolic rate, menstrual function, bone health, immunity, protein synthesis, cardiovascular health  
69 caused by relative energy deficiency”[7]. Despite progress, there are still no validated tool(s) for early detection  
70 of LEA to prevent the development of more severe forms of RED-S.

71 Accordingly, this review will address some important differences and gaps between short-term (<7 days)  
72 laboratory-based investigations of LEA (Figure 1) and their application to the field where symptoms usually  
73 reflect underlying, medium-term (weeks to <3 months) or long-term (>3 months to years) LEA. Specifically,  
74 we will: 1) discuss the key methodological challenges of EA calculations; 2) summarize the current evidence  
75 of short-term, strictly controlled laboratory investigations on the effects of LEA, along with an evaluation of  
76 potential markers to use in the assessment of EA status; as well as 3) explore the challenges and risks of  
77 extrapolating laboratory-based findings to screening and monitoring of risk of LEA in the field. Although  
78 much progress has been made on LEA and RED-S in males [8], this review will focus primarily on females to  
79 leverage the extensive decades of Triad data, into the RED-S context. Taken together, we hope this review will  
80 illuminate significant progress in understanding of the impact of LEA and RED-S in females. We will also  
81 highlight our significant field-based experiences of working with athletes with LEA, to demonstrate the  
82 challenges in applying laboratory findings to the field, as well as explore future research opportunities.

83 **2. Methodology and challenges of the energy availability equation**

84 Broadly speaking, LEA status can be assessed: a) Directly, through algebraic calculations from dietary EI and  
85 EEE data; and b) Indirectly through assessment of potential symptoms indicative of short to long-term LEA.

86 **2.1 Assessment of EA in the lab and field: methodology and challenges**

87 Although the algebraic calculation of EA is simple  $((EI-EEE)/FFM)$ , the practical and validity challenges of  
88 accurately determining EI and EEE are often ignored, which can result in significant errors when assessing  
89 these in the field or with poor laboratory methods [4, 9].

90 EI can be estimated a number of ways, such as self-reports, interviews, and food frequency recalls [10]. All of  
91 these methods are prone to significant errors (up to  $\sim 600$  kcal/d [9]) due to factors such as  
92 under/over/misreporting, poor athlete compliance as well as differences in data entry between experts.  
93 Furthermore, intakes are often recorded over a brief (3-7 days) time span which acts as a compromise to  
94 maintain compliance, while allowing for increased reliability and relevance of data. From an athlete  
95 perspective, data should be collected across different types of training days, and both weekdays and weekends,  
96 to gain reliable information of the overall practices [10].

97 The assessment of EEE in the field is equally challenging. While doubly labeled water is typically considered  
98 a gold standard for estimating total daily energy expenditure (TDEE), it provides average values over a longer  
99 time-period (usually 1-3 weeks) and does not differentiate between EEE and TDEE. There are currently no  
100 gold standard measurements for the estimation of EEE. At best, the use of laboratory-based measures where  
101 heart rate (HR) is plotted against indirect calorimetry data [11] can be used. In general, HR monitors or power  
102 meters (cycling, [12]) are likely to be superior to the use of metabolic equivalents (METs, [13]), but even these  
103 have the potential for errors in the magnitude of  $\sim 100-600$  kcal/d [14]. Regardless, we note that net EEE  
104 describes the energy expended during exercise only, and the resting energy expenditure component should be  
105 subtracted from the overall EEE value prior to performing final EA calculations, based on the latest definition  
106 of EA [4, 5]. An additional challenge to the calculation of daily EEE is the consideration of what constitutes  
107 exercise versus activities of daily living (i.e. non-exercise activity thermogenesis (NEAT)). There is currently  
108 no consensus on when NEAT may become part of the EEE equation, and this is especially impactful for  
109 athletes who undertake exercise based commuting or physical labor employment situations. Finally, for  
110 optimal accuracy, FFM assessment requires specialized training and equipment (skinfold calipers, DXA, etc.),  
111 as well as careful preparation of the athlete (standardized measurement protocol) that are not readily available  
112 to all athletes [9]. Finally, we note that the assessment of the components of the EA equation requires  
113 specialized skills and training to optimize the accuracy and reliability of measures. For example, dietary  
114 analysis should ideally be done by a trained and experienced sports dietitian, assessment of EEE by an applied  
115 sports physiologist, while the measurement of body composition requires specialized training (for example,  
116 The International Society for the Advancement of Kinanthropometry certification or radiation safety training  
117 for use of Dual-energy X-ray Absorptiometry).

118 Considering the methodological challenges outlined above, it is not surprising that many primarily field-based  
119 studies have consistently shown that athletes divided into groups based on calculated EA show little differences  
120 in physiological outcomes related to LEA [15, 16], such as menstrual function [15, 17, 18], and that

121 impairments to physiological systems cannot be traced down to a single EA threshold [19]. Some of these  
122 mismatches may be explained by the fact that despite consistent findings in females in well-controlled  
123 laboratory studies, some of these outcomes may not apply to more elite athletes of all ages, sexes and training  
124 characteristics. Indeed, one significant gap in the literature is the lack of research around EA thresholds for  
125 adolescents or children. The adolescence period (~13 to 19 years of age) is a phase during which ~25% of  
126 adult total BMD is acquired during the 4-year period surrounding peak height velocity (puberty growth spurt),  
127 with ~90% of peak BMD being achieved by age 20 [20]. Further research is also required to better understand  
128 the impact of both gynecological age (age from first menses to current age) and accumulated eumenorrheic  
129 age (number of accumulated years with an eumenorrheic cycle) on EA thresholds and RED-S outcomes.  
130 Nevertheless, it is safe to assume that the EA thresholds for adolescents are likely to be very different to the  
131 ones applied to adults due to the different energetic requirements of growth, development and puberty. It is  
132 also possible that factors such as diet quality [21], timing of meals within-day and in relation to training [17],  
133 carbohydrate availability [22], the magnitude of changes in any of the EA components in any one time, overall  
134 training load and psychological stress may also impact on EA outcomes [9]. Furthermore, while studies on  
135 LEA have defined a threshold of EA below which impairments have been observed, we note that this threshold  
136 should be treated as a rough guideline while appreciating that several factors will eventually determine the  
137 threshold at/below which impairments occur and/or that impairments probably also occur at differing rates  
138 [23]. Overall, if one is to implement EA assessments, incredible care and gold standard methods should be  
139 implemented, which is challenging in most real-world situations. Accordingly, in the following sections we  
140 propose the use of objective LEA indicators for athlete screening and diagnosis (*section 3*).

## 141 ***2.2 Considerations of the time frame of EA assessment***

142 Most research has not considered the ecological validity of reporting daily EA over the short-term (e.g. within-  
143 day or <1 week) period as a true indicator of the EA over medium to long-term (weeks to months or years).  
144 Laboratory-based studies tend to implement a homogenous/consistent LEA over the course of 3-6 days [5].  
145 However, athletes show large natural fluctuations in EI and EEE (training or competition) within [24] and  
146 between days [12]. Preliminary analyses of unpublished data from our team show for the first time that in free-  
147 living professional male road cyclists in pre-season training, assessed over 7 consecutive days, there is a strong  
148 inverse relationship between daily EEE and EA ( $r^2 = -0.78$ ), with spontaneous daily EI failing to match the  
149 energy with increasing EEE resulting in very low EA on days with high EEE (*Areta JL, manuscript in*  
150 *preparation*). While these observations are in male cyclists, we believe that the same inverse relationship  
151 between EEE and EA would be evident in females and that EA is likely to vary between training days, largely  
152 as a function of EEE, as there are no reasons to believe there would be a sex-based difference for this  
153 phenomenon. This is in line with what we have previously shown in elite male cyclists, where alternating days  
154 (e.g. intermittent) of LEA (~15 kcal/kg FFM/d) on race days were reported in comparison to optimal EA (~57  
155 kcal/kg FFM/d) on days in-between racing [12]. Here, despite extreme LEA on race days, there was no effect  
156 on physique outcomes or concentrations of testosterone, T3, IGF-1 or cortisol over an 8-d period, perhaps due

157 to the fact that the average EA over this observation period was ~36 kcal/kg FFM/d. This suggests that brief  
158 intermittent periods of extreme energy deficits, at least in males, may be manageable if overall EA across a  
159 time-period of ~1 week remains within acceptable limits; however, these outcomes remain to be established  
160 in female athletes. This study also challenges the notion that matching EI to EEE within every 24 hr block is  
161 crucial, and instead suggests that perhaps it is more important to maintain an overall, optimal EA over a rolling  
162 average of days.

163 In contrast to this idea, some research shows that small to moderate within-day periods of LEA over a single  
164 24 hr observation period are associated with negative health symptoms despite adequate daily total EA [17,  
165 24]. For example, female athletes with a greater number of hours of energy deficit of >300 kcal and with the  
166 largest daily energy deficits reported to have greater body fat percentage compared to those with less severe  
167 energy deficits within-day [24]. These transient within day deficits were also correlated with lower RMR<sub>ratio</sub>  
168 and estradiol and higher cortisol concentrations in females with menstrual dysfunction [17]. Although over a  
169 single 24 hr period, these findings may suggest that EA should be matched within-day, perhaps around  
170 exercise, to maintain optimal health. However, since this data is cross-sectional and thus makes it impossible  
171 to draw conclusions on causality, as well as determine whether within-day EA deficiency was short-term or  
172 long-term, more research is warranted around this topic.

173 Furthermore, it is likely that the duration, depth and gradient of LEA all play a role in the “dose” of the LEA  
174 ([5]; Figure 2) and thus, the potential outcomes of an intervention. The challenge with strictly controlled  
175 laboratory studies is that it is often impossible to monitor athletes for longer time periods. As such, our current  
176 knowledge relies largely on extrapolations or field-based reports. Therefore, we recommend that the athlete  
177 aims to maintain an overall EA level that meets the specific goals and requirements of their individual situation.  
178 That said, we also suggest that athletes focus on the timing of meals in relation to exercise, as this may have  
179 an additive effect on athlete health.

### 180 **3. Measurement and interpretation of blood markers as an alternative option to assess LEA**

181 The early detection of LEA in athletes is imperative as it represents the gateway to long-term or repeated  
182 exposure to the deleterious outcomes of prolonged LEA. It is well established that long-term (>3 months to  
183 years) LEA can ultimately result in important negative clinical outcomes such as functional hypothalamic  
184 amenorrhea and low bone mineral density (BMD), as well as potentially several other impairments to health  
185 and performance [6, 7, 25]. However, identification of early markers is still in its infancy.

186 Clinical diagnosis of long-term LEA is typically determined through assessment of bone health and menstrual  
187 function [25], but these markers have important shortcomings. In relation to bone, indications of LEA  
188 (structural changes) are often detected only after months or years of long-term/chronic LEA, where the effects  
189 of low BMD and related stress fractures can be irreversible. In relation to menstrual function, alteration of  
190 normal menstrual bleeding—an easily identifiable clinical marker of prolonged LEA—lies at the end of a

191 continuum anteceded by luteal phase defects and anovulation [26, 27] and it may not become evident, or it can  
192 become masked, due to the use of oral contraceptives [28], despite LEA [19, 26, 29].

193 Given that there is currently no validated tool to integrate potential objective physiological markers for  
194 detection of short-and medium-term (days to weeks) LEA, this section aims to identify measurable parameters  
195 that have been shown to be sensitive and responsive to LEA in laboratory-based studies (Table 1). We believe  
196 that these parameters may allow for objective identification of athletes at risk of LEA independent of other  
197 stressors, and help overcoming the challenges of assessing EA in the field based on EI and EEE.

### 198 **3.1 Short-term effect of LEA on blood parameters**

199 Most laboratory-based studies in females have tested the causal effects of short-term (3-6 days) LEA, usually  
200 at <30 kcal/kg FFM/d, on a range of hormonal and metabolic parameters in blood [4, 23, 30]. Based on these  
201 studies, with a relatively brief exposure to LEA, there is a clear disruption in the hormonal milieu and  
202 metabolism, suggesting various parameters that could be used in clinical practice (Table 1).

203 *Leptin.* Leptin is an important regulator of energy metabolism and when reduced under normal levels, it exerts  
204 a modulatory effect peripherally on a range of tissues and centrally in the hypothalamus [31]. Circulating leptin  
205 has been shown to unequivocally be reduced with LEA [22, 23, 32-34]. An EA of 30 kcal/kg FFM/d for 5 days  
206 was sufficient to decrease circulating leptin by ~35%, and further decreases of ~70% were seen with an EA of  
207 10 kcal/kg FFM/d [23]. It must be kept in mind that many commercial laboratories do not routinely measure  
208 for leptin, but it is a parameter that may be available to be measured on request, but at high cost.

209 *Thyroid hormones.* Thyroid hormones modulate energy expenditure through central and peripheral pathways  
210 [35] and are the main regulator of resting metabolic rate (RMR) [36]. Circulating levels of thyroid hormones  
211 are regulated through integration of energy-sensing inputs in the hypothalamus [35]. While T4 has not been  
212 measured in most experimental research on EA, free and total T3 has been shown to consistently be  
213 downregulated with LEA [23, 32-34, 37-39]. A threshold EA of ~19-25 kcal/kg FFM/d for 5 days has been  
214 reported to decrease T3 in a dose-response study [40] and it appears that as little as 2 days of EA of ~11 kcal/kg  
215 FFM/d may reduce circulating T3 [36]. Group average reductions from baseline values have been shown to  
216 range from ~6% at an EA of 30 kcal/kg FFM/d [23] to ~20-25% at EA of 10-19 kcal/kg FFM/d [23, 32, 33,  
217 37, 39, 40]. On the contrary, some literature suggests that LEA may increase T4 [36, 40], which could be  
218 related to reduced conversion of T4 to T3.

219 *The hypothalamic-pituitary-ovarian axis.* The negative effects of LEA on female reproductive endocrinology  
220 and physiology have been thoroughly researched [41]. Available evidence suggests that LEA is the main cause  
221 of functional hypothalamic amenorrhea and disrupts normal endocrine responses during the early follicular  
222 phase, such as reducing pulse frequency and increased pulse amplitude of luteinizing hormone (LH) [23, 32,  
223 37, 39]. However, follicle stimulating hormone (FSH) and oestrogen (E2) morning values, which are important  
224 hormones in the clinical diagnosis functional hypothalamic amenorrhea [28] and clear candidates as early



225 markers of LEA, are not affected by short-term LEA [23, 32, 37-39]. Therefore, changes in these hormones  
226 may become evident only after prolonged (weeks to months) LEA, but more research should address if short-  
227 term LEA affects their normal concentration in the luteal phase. Additionally, about 50% of female athletes  
228 use hormonal birth control, which makes the use of these hormones as reference unsuitable [42].

229 *Insulin-like growth factor 1 (IGF-1) and growth hormone (GH)*. IGF-1 is an important hormone for protein  
230 synthesis and cell proliferation [43]. Pituitary release of GH exerts the majority of its effects through regulating  
231 IGF-1 release from the liver [44]. LEA induces a state of *GH resistance*, in which normal GH effect on liver  
232 IGF-1 release is impaired, resulting in increased circulating GH and reduced circulating IGF-1 [44]. LEA  
233 induces a state of GH resistance within days, making both GH and IGF-1 good candidates as clinical markers  
234 of short-term LEA. Three studies in females have reported GH concentrations above resting levels with 4-5  
235 days of EA  $\leq 20$  kcal/kg FFM/d [23, 32, 39]. The 24 h average GH concentration has been shown to increase  
236 by 23-120% with an EA of 10 to 20 kcal/kg FFM/d [23, 32], while morning values increased by ~26% with  
237 an EA ~13 kcal/kg FFM/d [39], with no effect with an EA of 30 kcal/kg FFM/d [23, 32]. IGF-1 has been  
238 shown to consistently be reduced with similar amounts of LEA in studies with female participants [23, 32-34,  
239 38, 39]. The reduction of morning circulating IGF-1 values has been on average ~34% for short-term EA 10-  
240 20 kcal/kg FFM/d [23, 32, 33, 38, 39], while no effect was seen with an EA of 30 kcal/kg FFM/d [23]. Based  
241 on the evidence, IGF-1 seems to have significant support to be used as a morning marker of short-term LEA,  
242 while GH is unclear. IGF-1 and GH are typically not routine tests but can be easily accessed on demand.

243 *Bone formation and resorption markers*. Low BMD and increased prevalence of stress fractures are the result  
244 of medium to long-term negative balance of bone matrix synthesis that is an outcome of an imbalance between  
245 bone resorption and bone formation. Acute changes in bone turnover can be detected by measuring blood  
246 circulating markers of bone resorption and bone formation that are sensitive to exercise and nutrition  
247 interventions [45] and change within days of LEA. The acute effect of LEA on markers of bone metabolism  
248 has been limited to a few studies which collectively seem to support markers of bone formation are more  
249 sensitive than on markers of bone resorption [33, 34, 46]. For example, bone formation marker carboxy-  
250 terminal propeptide of type 1 procollagen (P1CP) has been shown to decrease linearly in one study titrating  
251 EA from 30 to 10 kcal/kg FFM/d from 10 to 29% of baseline values, respectively [46]. It is beyond the scope  
252 of this review to elucidate all bone metabolism markers, and the interested reader is directed here [45].  
253 However, we should highlight that currently commercial labs do not offer analysis of these markers and  
254 validated established norms do not yet exist.

### 255 **3.2 Urine luteinizing hormone surge**

256 Functional hypothalamic amenorrhea sits at the end of a continuum of endocrine regulations anteceded by  
257 luteal phase defects and anovulation, which are observed in proportion to markers of energy conservation [26,  
258 27]. Although studies assessing the endocrine effects of LEA have not investigated changes in LH surge during  
259 the menstrual cycle, this represents another potential practical parameter that could be used in the field with

260 non-hormonal contraception users [47], before more severe menstrual disturbances become evident. It is  
261 important to highlight though, that one study failed to find a relationship between the level of energy deficit  
262 and severity of induced menstrual disturbances during three consecutive menstrual cycles in exercising women  
263 [48], making unclear how sensitive this parameter can be to LEA.

### 264 ***3.3 On using indirect markers to assess LEA in female athletes***

265 We believe that some of the biomarkers outlined above could potentially help in identifying short-term LEA  
266 in female athletes (Table 1). However, we note that changes in these markers may not exceed the threshold for  
267 clinical reference range. It is also unclear whether general population norms can be directly applied to the  
268 athletic population, but to date no athlete specific reference values have been established. Within-individual  
269 variations as well as assay reliability should be considered as sources of random error that must be taken into  
270 account to determine whether a change in a parameter represents a real and meaningful change indicative of  
271 LEA. Additionally, these markers may only indicate the acute presence of LEA, without currently having the  
272 capacity to distinguish exactly what level of LEA and/or for what duration. Potentially, markers that have been  
273 shown to be strongly related to menstrual dysfunction but have not shown changes in acute LEA studies —  
274 such as E2 and FSH— [5] could be investigated as markers of medium-long term exposure to LEA. Further  
275 complexity to the interpretation of these markers comes from the fact that some of the hormones (for example,  
276 GH and cortisol) may be sensitive to the effects of acute or chronic exercise [49]; it may therefore be  
277 challenging to determine whether abnormalities in a marker are due to exercise training or underlying LEA.  
278 Ideally, we recommend collection for these blood tests to be performed in the overnight fasted state, first thing  
279 in the morning prior to exercise and to have the last high-intensity (hard training) session at least 48 hr prior,  
280 to minimize the potential effects of exercise and normalize circadian rhythms on these markers. Finally, it is  
281 important to emphasize that the blood biomarkers are based on a range of well-controlled laboratory studies  
282 determining the effect of LEA **only** on very specific populations (often sedentary women) and need to be  
283 validated in wider athlete populations, specific timing of sample collection in relation to LEA exposure, and  
284 they may only be used as a broad guidance for further investigation of energetic status of athletes to avoid  
285 potential negative effects of prolonged LEA.

## 286 **4. Risks and flaws of extrapolating findings from the lab to the field – a identifying key differences** 287 **between the two different settings**

288 While strictly controlled, short-term laboratory investigations of LEA are essential to characterize the  
289 physiological outcomes of acute LEA, their applicability to the field is challenged by several factors, which  
290 will be outlined below.

### 291 ***4.1 Length of exposure***

292 Without exception, laboratory studies focusing on LEA implement very brief (3-5 days) protocols [11, 22, 23,  
293 33, 36, 38-40, 46]. Accordingly, these investigations are often unable inform us about the long-term effects of  
294 LEA. Even if a specific marker does not show a change after 5 days, it may be affected with increasing duration

295 of LEA (for example, total cholesterol and E2 appear to be influenced only after prolonged LEA; [23, 32, 37-  
296 39, 50]). Furthermore, some markers that have showed a change in short-term laboratory studies appear to be  
297 within the reference range in cross-sectional comparisons of amenorrheic vs eumenorrheic females (for  
298 example, insulin and IGF-1 rarely differ between the two [15, 16]). Indeed, long-term exposure seems to reduce  
299 the circulating level of at least some of these markers, even if the values remain within the clinical reference  
300 value [51]. Regardless, since athletes may spend a considerably longer time-period (often weeks to months;  
301  $\geq 10$  to 20 times longer than laboratory studies) in LEA [52], extrapolating observations from the lab to field  
302 settings appears inappropriate.

303 In addition, currently very little is known of the interactions between the magnitude of change required for a  
304 single biomarker (e.g. a decrease in P1NP or a decrease in T3) to induce a subsequent, meaningful functional  
305 impairment (e.g. with P1NP, a decrease in BMD and development of a stress fracture; or with the case of T3,  
306 a reduction in RMR). This is obviously a key consideration as an isolated change in a biomarker is irrelevant  
307 unless a functional outcome follows. Unfortunately, the literature lacks mechanistic data on the time-course or  
308 threshold for which a functional change can be expected. Nevertheless, an investigation in female physique  
309 competitors reported endocrine impairments (reductions in T3, E2, leptin and testosterone) as a result of a 4-  
310 month preparatory period (and LEA) for a competition but a rebound in most hormones except T3 and  
311 testosterone at 3-4 months post-competition [53]. The length of exposure is an important consideration as the  
312 reversal of impairments may be dependent on this factor [54]. Based on available literature, we hypothesize  
313 that reversal is rapid following brief (<1 to 3 months) periods of LEA [53], whereas there could be potential  
314 for the development for much slower reversal of biomarkers, and some irreversible symptoms (including  
315 persistent amenorrhea or low BMD), with increasing duration of LEA [15, 55, 56].

#### 316 **4.2 Ecological validity of daily EA and dietary macronutrient composition**

317 Another challenge in the application of laboratory-based results into the field is the use of sudden drops in EA  
318 (for example, from 45 to 15 kcal/kg FFM/d) that remain constantly low, without between-day fluctuations in  
319 EA. This is likely not representative of what the majority athletes may experience in the field, in which athletes  
320 may be exposed to LEA in different patterns (Figure 2). Most laboratory-based studies induce a  
321 consistent/homogeneous LEA (often  $\sim 10$ -20 kcal/kg FFM/d), while we have observed daily fluctuations in EA  
322 which can be inversely related to EEE [13; Areta JL, *manuscript in preparation*], which is line with the recent  
323 findings of fluctuation of EA across training/competition seasons being related to EEE [57]. These protocols  
324 are the opposite of what is currently considered best practice for weight loss for athletic populations, where  
325 some evidence supports the use of gradual, small changes in EI with weight-loss interventions [58].  
326 Meanwhile, more research is required to determine the optimal periodization of EA “dose” to prevent  
327 deleterious health outcomes, while optimizing potential performance outcomes (Figure 2). Furthermore,  
328 laboratory investigations tend to implement significantly lower EA values for the LEA intervention arms than  
329 what has been reported in the cross-sectional research [15, 50]. Therefore, it is possible that some of the  
330 changes reported in laboratory-based LEA investigations after only 5 days of LEA are more a result of a sudden

331 change and/or the application of an extreme LEA, or potentially, a combination of both, and may poorly  
332 translate into the field where more conservative approaches are likely to be implemented.

333 In addition to EI, the macronutrient composition deserves a brief discussion. Surprisingly and against the  
334 recommendations for optimized weight-loss approach to athletes [59], most studies on LEA have implemented  
335 a static macronutrient ratio for all levels of EA. This not only results in absolute reduction of EA, but also in a  
336 decrease in all macronutrients. Here, carbohydrates and proteins are of special interest as both have important  
337 roles in the management of the metabolic rate and physique outcomes during energy restriction. Indeed, studies  
338 have shown that some of the effects of LEA are mediated via reduced CHO availability [22], which appears  
339 to have effects on especially leptin and T3 concentrations [60, 61]. These in turn affect RMR, thus facilitating  
340 or suppressing metabolism during periods of LEA [62]. Meanwhile, increased protein intake in the face of  
341 LEA allows maintain increased muscle protein synthesis [63] lean mass and health during weight loss [59].  
342 Therefore, it could be argued that the poor emphasis on optimized CHO and protein intake, coupled with more  
343 extreme levels of LEA, in many LEA investigations may potentiate the negative outcomes and is likely to  
344 decrease the ecological validity and applicability of the results into the field practice when following current  
345 dietary recommendations.

346 Finally, strict laboratory-controlled investigations are often conducted in sedentary or only moderately trained  
347 individuals, where applications to the elite female athlete are troublesome. For example, there are genetic,  
348 anthropometric and physiological differences between untrained, trained and elite athletes [64]. In addition,  
349 and as mentioned earlier, elite athletes engage in high training loads which can be 2- to 4-fold greater compared  
350 to recreational athletes (for example, [65]).

## 351 **5. Summary and future research considerations**

352 Decades of laboratory research have shown that impairments to several body systems can be seen in as little  
353 as 4-5 days of strictly controlled LEA. Meanwhile, a large body of cross-sectional reports exists on the  
354 interrelatedness of eating disorders, menstrual dysfunction and impaired bone health in females. These findings  
355 have increased the awareness of LEA as the key underpinning factor behind a broader set of health and  
356 performance outcomes recognized in models of ‘Relative Energy Deficiency in Sport’ and the ‘Triad’.  
357 However, a significant gap exists between short-term laboratory studies and cross-sectional reports of long-  
358 term/chronic LEA; as such, no consistent and reliable tools exist to screen athletes for LEA. Here we have  
359 highlighted several methodological challenges related to the assessment of the parameters of EA equation on  
360 the field (namely, EI and EEE). In light of the uncertainty of these parameters, we propose the use of more  
361 “objective” parameters as ‘end point’ markers of LEA (i.e. blood markers). These suggestions are mostly based  
362 on short-term laboratory research which supports the use of several hormone markers in identifying LEA in  
363 female athletes. However, we note that direct extrapolations of laboratory-based outcomes into the field are  
364 likely to be problematic due to several challenges relating to ecological validity. Therefore, we provide a  
365 critical appraisal of the scientific literature and a set of potential set of tools to utilize on the field. Further work

366 is warranted to gain a more sophisticated understanding of the best biomarkers to use for assessment of  
367 medium- and long-term LEA. This includes the development of athlete specific standard ranges and an  
368 understanding of what constitutes a real meaningful change from baseline (if repeat measures are done) as well  
369 as of how to interpret isolated (single time point) measures. Future studies should aim to more systematically  
370 assess and define the best markers of LEA to detect early signs of LEA and thereby, avoid the long-term  
371 impairments to health and performance of the female athlete.


372 **Figure legends**

373 **Table 1.** Summary of parameters that are likely to provide an objective measure of short or moderate (days/weeks) exposure to low energy availability (LEA)  
 374 in females. For blood parameters assessed in a biochemical laboratory it is important to consider that within-individual changes should be considered provided  
 375 that values may be affected by LEA within individual without falling outside the accepted clinical range. Ideally, we recommend blood tests first thing in the  
 376 morning prior to exercise and to have the last high-intensity (hard training) session at least 48 hours prior, to minimize the potential effects of exercise and  
 377 normalize circadian rhythms on these markers. Overall, it is important to standardize measurements and consider the typical variation for an individual and error  
 378 of measurement to determine likelihood of real change. Rather than considering one parameter in isolation, it is likely a stronger indicator to note trends in  
 379 several parameters in combination.

Parameter	Sensitivity to LEA	Feasibility of test in real world	Best time of menstrual cycle to test	Ideal testing conditions	Refs
<b>Blood Leptin</b>	High	High; Not a routine biochemical assessment			[22, 23, 32-34]
<b>Blood IGF-1</b>	High	High	Concentrations independent of cycle phase? (studied in early follicular non-OC users)	Standardized morning fasted; completed in the same laboratory	[23, 32-34, 38, 39]
<b>Blood Bone formation and resorption markers</b>	Formation likely more sensitive than resorption	Low; Not a routine biochemical assessment			[33, 34, 46]
<b>Blood total and freeT3</b>	High	High			[23, 32-34, 36, 37, 39, 40]
<b>Urinary LH surge</b>	?	High; Affordable home kit	Between the follicular and luteal phase (non-OC users)	Collection of urine at 11-15 hr	[26, 27, 47]


380

381



382

383 **Figure 1.** Visual summary of the main differences between laboratory-based studies assessing the endocrine, metabolic and physiological effects of low energy  
 384 availability (LEA) in females (left) and the real-world environment (right) when working with athletes likely exposed to LEA. It is important to consider these  
 385 differences to understand the limitations and potential errors in extrapolating laboratory-based studies to field observations. These considerations put on evidence  
 386 that the typically considered LEA ‘threshold’ of 30 kcal/kg FFM/d—which has been determined in laboratory-based studies of ~5 days in duration— may not  
 387 necessarily represent a true indicator of LEA when assessed on the field. EA, energy availability; EI, energy intake; EEE, exercise energy expenditure; NEAT,  
 388 Non-exercise activity thermogenesis.



389

390 **Figure 2.** Visual representation of different ways in which low energy availability (LEA) or low carbohydrate availability (LCA) doses may be achieved in real-  
 391 world scenarios. While laboratory-based studies tend to induce single severe LEA (*black area*), it is likely that on the field there are different scenarios, such as  
 392 subtle LEA over prolonged time (*dashed area*) or a more heterogenous exposure to LEA during periods interspersed with adequate energy availability (*gray*  
 393 *areas*). In addition, the gradient of change in LEA (e.g. sudden vs gradual drops; *dashed line*) is likely to play a role.

394



395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445

## References

1. Drinkwater, B.L., et al., *Bone mineral content of amenorrheic and eumenorrheic athletes*. N Engl J Med, 1984. **311**(5): p. 277-81.
2. Yeager, K.K., et al., *The female athlete triad: disordered eating, amenorrhea, osteoporosis*. Med Sci Sports Exerc, 1993. **25**(7): p. 775-7.
3. Nattiv, A., et al., *American College of Sports Medicine position stand. The female athlete triad*. Med Sci Sports Exerc, 2007. **39**(10): p. 1867-82.
4. Loucks, A.B., B. Kiens, and H.H. Wright, *Energy availability in athletes*. J Sports Sci, 2011. **29 Suppl 1**: p. S7-15.
5. Areta, J.L., H.L. Taylor, and K. Koehler, *Low energy availability: history, definition and evidence of its endocrine, metabolic and physiological effects in prospective studies in females and males*. Eur J Appl Physiol, 2020.
6. Mountjoy, M., et al., *IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update*. Br J Sports Med, 2018. **52**(11): p. 687-697.
7. Mountjoy, M., et al., *The IOC consensus statement: beyond the Female Athlete Triad--Relative Energy Deficiency in Sport (RED-S)*. Br J Sports Med, 2014. **48**(7): p. 491-7.
8. Tenforde, A.S., et al., *Parallels with the Female Athlete Triad in Male Athletes*. Sports Med, 2016. **46**(2): p. 171-82.
9. Burke, L.M., et al., *Pitfalls of Conducting and Interpreting Estimates of Energy Availability in Free-Living Athletes*. Int J Sport Nutr Exerc Metab, 2018. **28**(4): p. 350-363.
10. Capling, L., et al., *Validity of Dietary Assessment in Athletes: A Systematic Review*. Nutrients, 2017. **9**(12).
11. Koehler, K., et al., *Low resting metabolic rate in exercise-associated amenorrhea is not due to a reduced proportion of highly active metabolic tissue compartments*. Am J Physiol Endocrinol Metab, 2016. **311**(2): p. E480-7.
12. Heikura, I.A., et al., *Alternate-Day Low Energy Availability During Spring Classics in Professional Cyclists*. Int J Sports Physiol Perform, 2019: p. 1233-1243.
13. Ainsworth, B.E., et al., *Compendium of physical activities: an update of activity codes and MET intensities*. Med Sci Sports Exerc, 2000. **32**(9 Suppl): p. S498-504.
14. O'Driscoll, R., et al., *How well do activity monitors estimate energy expenditure? A systematic review and meta-analysis of the validity of current technologies*. Br J Sports Med, 2020. **54**(6): p. 332-340.
15. Heikura, I.A., et al., *Low Energy Availability Is Difficult to Assess but Outcomes Have Large Impact on Bone Injury Rates in Elite Distance Athletes*. Int J Sport Nutr Exerc Metab, 2018. **28**(4): p. 403-411.
16. Koehler, K., et al., *Comparison of self-reported energy availability and metabolic hormones to assess adequacy of dietary energy intake in young elite athletes*. Appl Physiol Nutr Metab, 2013. **38**(7): p. 725-33.
17. Fahrenholtz, I.L., et al., *Within-day energy deficiency and reproductive function in female endurance athletes*. Scand J Med Sci Sports, 2018. **28**(3): p. 1139-1146.
18. Tornberg, A.B., et al., *Reduced Neuromuscular Performance in Amenorrheic Elite Endurance Athletes*. Med Sci Sports Exerc, 2017. **49**(12): p. 2478-2485.
19. Lieberman, J.L., et al., *Menstrual Disruption with Exercise Is Not Linked to an Energy Availability Threshold*. Med Sci Sports Exerc, 2018. **50**(3): p. 551-561.
20. Santos, L., K.J. Elliott-Sale, and C. Sale, *Exercise and bone health across the lifespan*. Biogerontology, 2017. **18**(6): p. 931-946.
21. Melin, A., et al., *Low-energy density and high fiber intake are dietary concerns in female endurance athletes*. Scand J Med Sci Sports, 2016. **26**(9): p. 1060-71.
22. Hilton, L.K. and A.B. Loucks, *Low energy availability, not exercise stress, suppresses the diurnal rhythm of leptin in healthy young women*. Am J Physiol Endocrinol Metab, 2000. **278**(1): p. E43-9.
23. Loucks, A.B. and J.R. Thuma, *Luteinizing hormone pulsatility is disrupted at a threshold of energy availability in regularly menstruating women*. J Clin Endocrinol Metab, 2003. **88**(1): p. 297-311.

- 446 24. Deutz, R.C., et al., *Relationship between energy deficits and body composition in elite female*  
447 *gymnasts and runners*. Med Sci Sports Exerc, 2000. **32**(3): p. 659-68.
- 448 25. Joy, E., et al., *2014 female athlete triad coalition consensus statement on treatment and return to*  
449 *play of the female athlete triad*. Curr Sports Med Rep, 2014. **13**(4): p. 219-32.
- 450 26. De Souza, M.J., et al., *High prevalence of subtle and severe menstrual disturbances in exercising*  
451 *women: confirmation using daily hormone measures*. Hum Reprod, 2010. **25**(2): p. 491-503.
- 452 27. De Souza, M.J., et al., *Severity of energy-related menstrual disturbances increases in proportion to*  
453 *indices of energy conservation in exercising women*. Fertil Steril, 2007. **88**(4): p. 971-5.
- 454 28. Gordon, C.M., et al., *Functional Hypothalamic Amenorrhea: An Endocrine Society Clinical Practice*  
455 *Guideline*. J Clin Endocrinol Metab, 2017. **102**(5): p. 1413-1439.
- 456 29. De Souza, M.J., et al., *Drive for thinness score is a proxy indicator of energy deficiency in exercising*  
457 *women*. Appetite, 2007. **48**(3): p. 359-67.
- 458 30. Elliott-Sale, K.J., et al., *Endocrine Effects of Relative Energy Deficiency in Sport*. Int J Sport Nutr Exerc  
459 Metab, 2018. **28**(4): p. 335-349.
- 460 31. Bluher, S. and C.S. Mantzoros, *Leptin in humans: lessons from translational research*. Am J Clin Nutr,  
461 2009. **89**(3): p. 991S-997S.
- 462 32. Loucks, A.B., *The response of luteinizing hormone pulsatility to 5 days of low energy availability*  
463 *disappears by 14 years of gynecological age*. J Clin Endocrinol Metab, 2006. **91**(8): p. 3158-64.
- 464 33. Papageorgiou, M., et al., *Bone metabolic responses to low energy availability achieved by diet or*  
465 *exercise in active eumenorrheic women*. Bone, 2018. **114**: p. 181-188.
- 466 34. Papageorgiou, M., et al., *Effects of reduced energy availability on bone metabolism in women and*  
467 *men*. Bone, 2017. **105**: p. 191-199.
- 468 35. McAninch, E.A. and A.C. Bianco, *Thyroid hormone signaling in energy homeostasis and energy*  
469 *metabolism*. Ann N Y Acad Sci, 2014. **1311**: p. 77-87.
- 470 36. Loucks, A.B. and R. Callister, *Induction and prevention of low-T3 syndrome in exercising women*. Am  
471 J Physiol, 1993. **264**(5 Pt 2): p. R924-30.
- 472 37. Loucks, A.B. and M. Verdun, *Slow restoration of LH pulsatility by refeeding in energetically disrupted*  
473 *women*. Am J Physiol, 1998. **275**(4): p. R1218-26.
- 474 38. Loucks, A.B. and E.M. Heath, *Dietary restriction reduces luteinizing hormone (LH) pulse frequency*  
475 *during waking hours and increases LH pulse amplitude during sleep in young menstruating women*. J  
476 Clin Endocrinol Metab, 1994. **78**(4): p. 910-5.
- 477 39. Loucks, A.B., M. Verdun, and E.M. Heath, *Low energy availability, not stress of exercise, alters LH*  
478 *pulsatility in exercising women*. J Appl Physiol (1985), 1998. **84**(1): p. 37-46.
- 479 40. Loucks, A.B. and E.M. Heath, *Induction of low-T3 syndrome in exercising women occurs at a threshold*  
480 *of energy availability*. Am J Physiol, 1994. **266**(3 Pt 2): p. R817-23.
- 481 41. Loucks, A.B., *Exercise Training in the Normal Female: Effects of Low Energy Availability on*  
482 *Reproductive Function*, in *Endocrinology of Physical Activity and Sport*, A.C. Hackney and N.W.  
483 Constantini, Editors. 2020, Springer International Publishing. p. 171-191.
- 484 42. Martin, D., et al., *Period Prevalence and Perceived Side Effects of Hormonal Contraceptive Use and*  
485 *the Menstrual Cycle in Elite Athletes*. Int J Sports Physiol Perform, 2018. **13**(7): p. 926-932.
- 486 43. Clemmons, D.R., *Role of IGF-I in skeletal muscle mass maintenance*. Trends Endocrinol Metab, 2009.  
487 **20**(7): p. 349-56.
- 488 44. Fazeli, P.K. and A. Klibanski, *Determinants of GH resistance in malnutrition*. J Endocrinol, 2014. **220**(3):  
489 p. R57-65.
- 490 45. Dolan, E., et al., *The Bone Metabolic Response to Exercise and Nutrition*. Exerc Sport Sci Rev, 2020.  
491 **48**(2): p. 49-58.
- 492 46. Ihle, R. and A.B. Loucks, *Dose-response relationships between energy availability and bone turnover*  
493 *in young exercising women*. J Bone Miner Res, 2004. **19**(8): p. 1231-40.
- 494 47. Khattab, A.F., F.A. Mustafa, and P.J. Taylor, *The use of urine LH detection kits to time intrauterine*  
495 *insemination with donor sperm*. Hum Reprod, 2005. **20**(9): p. 2542-5.

- 496 48. Williams, N.I., et al., *Magnitude of daily energy deficit predicts frequency but not severity of menstrual*  
497 *disturbances associated with exercise and caloric restriction*. Am J Physiol Endocrinol Metab, 2015.  
498 **308**(1): p. E29-39.
- 499 49. Wideman, L., et al., *Growth hormone release during acute and chronic aerobic and resistance*  
500 *exercise: recent findings*. Sports Med, 2002. **32**(15): p. 987-1004.
- 501 50. Melin, A., et al., *Energy availability and the female athlete triad in elite endurance athletes*. Scand J  
502 Med Sci Sports, 2015. **25**(5): p. 610-22.
- 503 51. De Souza, M.J., et al., *Luteal phase deficiency in recreational runners: evidence for a hypometabolic*  
504 *state*. J Clin Endocrinol Metab, 2003. **88**(1): p. 337-46.
- 505 52. Mathisen, T.F., et al., *Physical health and symptoms of relative energy deficiency in female fitness*  
506 *athletes*. Scand J Med Sci Sports, 2020. **30**(1): p. 135-147.
- 507 53. Hulmi, J.J., et al., *The Effects of Intensive Weight Reduction on Body Composition and Serum*  
508 *Hormones in Female Fitness Competitors*. Front Physiol, 2016. **7**: p. 689.
- 509 54. Cialdella-Kam, L., et al., *Dietary intervention restored menses in female athletes with exercise-*  
510 *associated menstrual dysfunction with limited impact on bone and muscle health*. Nutrients, 2014.  
511 **6**(8): p. 3018-39.
- 512 55. Ackerman, K.E., et al., *Fractures in Relation to Menstrual Status and Bone Parameters in Young*  
513 *Athletes*. Med Sci Sports Exerc, 2015. **47**(8): p. 1577-86.
- 514 56. Areta, J.L., *Case Study: Resumption of Eumenorrhea in Parallel With High Training Load After 4 Years*  
515 *of Menstrual Dysfunction: A 5-Year Follow-Up of an Elite Female Cyclist*. Int J Sport Nutr Exerc Metab,  
516 2020: p. 1-6.
- 517 57. Westerterp, K.R., *Seasonal variation in body mass, body composition and activity-induced energy*  
518 *expenditure: a long-term study*. Eur J Clin Nutr, 2020. **74**(1): p. 135-140.
- 519 58. Garthe, I., et al., *Effect of two different weight-loss rates on body composition and strength and*  
520 *power-related performance in elite athletes*. Int J Sport Nutr Exerc Metab, 2011. **21**(2): p. 97-104.
- 521 59. Hector, A.J. and S.M. Phillips, *Protein Recommendations for Weight Loss in Elite Athletes: A Focus on*  
522 *Body Composition and Performance*. Int J Sport Nutr Exerc Metab, 2018. **28**(2): p. 170-177.
- 523 60. Jenkins, A.B., et al., *Carbohydrate intake and short-term regulation of leptin in humans*. Diabetologia,  
524 1997. **40**(3): p. 348-51.
- 525 61. Spaulding, S.W., et al., *Effect of caloric restriction and dietary composition of serum T3 and reverse*  
526 *T3 in man*. J Clin Endocrinol Metab, 1976. **42**(1): p. 197-200.
- 527 62. Trexler, E.T., A.E. Smith-Ryan, and L.E. Norton, *Metabolic adaptation to weight loss: implications for*  
528 *the athlete*. J Int Soc Sports Nutr, 2014. **11**(1): p. 7.
- 529 63. Areta, J.L., et al., *Reduced resting skeletal muscle protein synthesis is rescued by resistance exercise*  
530 *and protein ingestion following short-term energy deficit*. Am J Physiol Endocrinol Metab, 2014.  
531 **306**(8): p. E989-97.
- 532 64. Al-Khelaifi, F., et al., *Metabolic GWAS of elite athletes reveals novel genetically-influenced*  
533 *metabolites associated with athletic performance*. Sci Rep, 2019. **9**(1): p. 19889.
- 534 65. Johansen, J.M., et al., *Improving Utilization of Maximal Oxygen Uptake and Work Economy in*  
535 *Recreational Cross-Country Skiers With High-Intensity Double-Poling Intervals*. Int J Sports Physiol  
536 Perform, 2020: p. 1-8.

537