



Coherence scale of magnetic fields generated in early-time forward shocks of GRBs

N. Jordana-Mitjans¹,¹★ C. G. Mundell¹, R. J. Smith², C. Guidorzi^{3,4,5}, M. Marongiu⁶, S. Kobayashi⁷, A. Gomboc⁷, M. Shrestha² and I. A. Steele²

¹Department of Physics, University of Bath, Claverton Down, Bath BA2 7AY, UK

²Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, UK

³Department of Physics and Earth Science, University of Ferrara, via Saragat 1, I-44122 Ferrara, Italy

⁴INFN – Sezione di Ferrara, Via Saragat 1, I-44122 Ferrara, Italy

⁵INAF – Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Piero Gobetti 101, I-40129 Bologna, Italy

⁶INAF – Osservatorio Astronomico di Cagliari – via della Scienza 5 – I-09047 Selargius, Italy

⁷Center for Astrophysics and Cosmology, University of Nova Gorica, Vipavska 13, 5000 Nova Gorica, Slovenia

Accepted 2021 April 4. Received 2021 March 19; in original form 2021 January 22

ABSTRACT

We report the earliest ever detection of optical polarization from a GRB forward shock (GRB 141220A), measured 129.5–204.3 s after the burst using the multicolour RINGO3 optical polarimeter on the 2-m fully autonomous robotic Liverpool Telescope. The temporal decay gradient of the optical light curves from 86 to ~ 2200 s post-burst is typical of classical forward shocks with $\alpha = 1.091 \pm 0.008$. The low-optical polarization $P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent (2σ) at mean time ~ 168 s post-burst is compatible with being induced by the host galaxy dust ($A_{V,HG} = 0.71 \pm 0.15$ mag), leaving low polarization intrinsic to the GRB emission itself – as theoretically predicted for forward shocks and consistent with previous detections of low degrees of optical polarization in GRB afterglows observed hours to days after the burst. The current sample of early-time polarization data from forward shocks suggests polarization from (a) the Galactic and host galaxy dust properties (i.e. $P \sim 1 - 3$ per cent), (b) contribution from a polarized reverse shock (GRB deceleration time, jet magnetization) or (c) forward shock intrinsic polarization (i.e. $P \leq 2$ per cent), which depends on the magnetic field coherence length-scale and the size of the observable emitting region (burst energetics, circumburst density).

Key words: magnetic fields – polarization – gamma-ray burst: individual: grb 141220A – ISM: jets and outflows.

1 INTRODUCTION

Gamma-ray bursts (GRBs) are the brightest flashes of γ -ray emission in the Universe. After the collapse of a massive star or coalescence of two compact stellar objects (Woosley 1993; Berger 2014; Abbott et al. 2017a, b), the accretion into new-born compact object powers ultrarelativistic jetted emission that – via internal dissipation processes – produces the highly variable and characteristic γ -ray prompt emission. In the fireball model framework, the ejecta is later on decelerated by the circumburst medium by a pair of external shocks – a reverse shock propagating back into the ejecta (Rees & Meszaros 1992; Sari & Piran 1999; Kobayashi 2000) and a forward shock propagating into the ambient medium – producing a long-lived afterglow that can be seen from X-rays to radio frequencies (e.g. Piran 1999; Mészáros 2002; Piran 2004).

During the first hundreds of seconds after the burst – for prompt and reverse shock emission – different polarimetric signatures are expected for competing jet models: unpolarized emission from weak, tangled magnetic fields for a baryonic jet (Rees & Meszaros 1994;

Medvedev & Loeb 1999) or highly polarized emission due to the presence of globally ordered magnetic fields (Granot & Königl 2003; Lyutikov, Pariev & Blandford 2003; Zhang & Kobayashi 2005; Komissarov et al. 2009). Currently, the discrepancy among the results of time-integrated γ -ray polarization studies of the prompt emission remains debated (Coburn & Boggs 2003; Rutledge & Fox 2004; Willis et al. 2005; Chattopadhyay et al. 2019; Kole et al. 2020; Gill & Granot 2021). Additionally, GRB 170114A time-resolved analysis suggests the evolution of the polarization degree and angle over a single pulse, reaching values of $P \sim 30$ per cent (Burgess et al. 2019). Early-time optical polarimetric studies of reverse shocks favour a mildly magnetized jet with primordial magnetic fields advected from the central engine (e.g. GRB 090102; Steele et al. 2009, GRB 120308A; Mundell et al. 2013). However, recent observations suggest that very energetic GRBs can be launched highly magnetized (GRB 190114C: Jordana-Mitjans et al. 2020).

Whilst polarimetry of the prompt emission and early afterglow determines the jet physics, polarimetric observations of forward shocks allow to test particle acceleration mechanisms (e.g. shock formation, magnetic turbulence), study the dust properties of GRBs' environments and resolve the large-scale geometry of jets at cosmological distances (Lazzati et al. 2003; Rossi et al. 2004). In the

* E-mail: N.Jordana@bath.ac.uk

afterglow framework, the forward shock is powered by shocked ambient medium and tangled magnetic fields are locally generated in shocks and amplified by plasma instabilities (e.g. via Weibel instability; Weibel 1959; Nishikawa et al. 2003; Silva et al. 2003; Medvedev et al. 2005). The magnetic field is randomly oriented in space with length-scales much smaller than the size of the observable region of the shock. Consequently, the emission is expected to be intrinsically unpolarized when the jet is on-axis (Medvedev & Loeb 1999). However, it can be significantly polarized if the random field is anisotropic (e.g. the averaged strength in the shock normal direction is stronger or weaker) and the line-of-sight runs almost along the jet edge. The second condition is satisfied when a jet break is observed (Ghisellini & Lazzati 1999; Sari 1999; Rossi et al. 2004). Additionally, the emission can show few percents of polarization due to differential dust extinction (Hiltner 1949; Lazarian 2007) along the Galactic line of sight and in the GRB environment (e.g. Lazzati et al. 2003; Klose et al. 2004).

Pioneering polarization studies of GRB afterglows found $P \sim 1$ –3 per cent hours to days after the onset of the GRB in the optical and near-infrared bands (e.g. Covino et al. 2004) – during the forward shock decay. The first polarization constraint was $P < 2.3$ per cent at 18 h post-burst for GRB 990123 (Hjorth et al. 1999). Observations of GRB 990510 registered steady levels of polarization at $P \sim 1.7$ per cent level from 18 to 21 h post-burst, hence validating synchrotron as the emitting mechanism (Covino et al. 1999; Wijers et al. 1999). The polarimetric monitoring of GRB 030329 afterglow ~ 0.5 –38 d post-burst measured low levels with significant variability around the light-curve jet break (Greiner et al. 2004) – expected for beamed ejecta. Polarization also behaved similarly during the achromatic break of GRB 091018 (Wiersema et al. 2012). Recent spectropolarimetric observations of GRB 191221B also detected low levels $P = (1.5 \pm 0.5)$ per cent for forward shock emission ~ 2.9 h after the burst (Buckley et al. 2020).

With the advance of robotic telescopes, the first early-time polarization constraint of an optical afterglow was made 203 s post-burst – during the afterglow onset of GRB 060418. The $P < 8$ per cent (2σ) suggested a significant contribution of unpolarized forward shock photons (Mundell et al. 2007a). GRB 091208B emission was interpreted as forward shock; the $P = (10.4 \pm 2.5)$ per cent detection during 149–706 s post-burst disfavoured the afterglow model and proposed other mechanisms for the amplification of the magnetic field at the front shock, e.g. magnetohydrodynamic instabilities (Uehara et al. 2012). GRB 131030A polarization degree remained steady at $P = (2.1 \pm 1.6)$ per cent level during 0.18 – 1.85 h post-burst and was associated to dust-induced polarization from the Galactic interstellar medium (ISM; King et al. 2014).

GRB 141220A optical light curve, presented in this paper, resembles those of late-time classical forward shocks detected hours to days after the prompt. In contrast, Liverpool Telescope (LT; Steele et al. 2004) observations indicate that the forward shock was dominating the total emission as soon as ~ 86 s after the burst. The afterglow was bright at the time of polarization observations – starting 129.5 s post-burst – hence providing good early-time constraints. GRB 141220A allows checking whether the early-time emission from forward shocks is intrinsically unpolarized as predicted in the fireball model and observed during late-time afterglows.

This paper is structured as follows. In Section 2, we detail the data analysis of GRB 141220A follow-up observations by the LT. In Section 3, we study the emission decay, the broad-band spectral properties of the burst and the polarization. In Section 4, we check closure relations for GRB 141220A, we discuss which mechanisms can explain the observed polarization levels and we compare GRB

141220A to other early-time forward shock measurements. In Section 5, we summarize our findings. Throughout this paper, we assume flat Lambda cold dark matter (Λ CDM) cosmology $\Omega_m = 0.32$, $\Omega_\Lambda = 0.68$, and $h = 0.67$, as reported by Planck Collaboration VI (2018a). We adopt the convention $F_\nu \propto t^{-\alpha} \nu^{-\beta}$, where α is the temporal index and β is the spectral index. The spectral index is related to the photon index like $\beta = \beta_{\text{PI}} - 1$. Uncertainties are quoted at 1σ confidence level unless stated otherwise.

2 OBSERVATIONS AND DATA REDUCTION

On 2014 December 20 at $T_0 = 06:02:52.7$ UT, the *Swift* Burst Alert Telescope (BAT; Cummings et al. 2014) was triggered by the detection of a pulse of γ -rays corresponding to the candidate GRB 141220A. The pulses lasted from $T_0 - 0.9$ s to $T_0 + 7.3$ s and consisted of one structured peak – centred at $\sim T_0 + 0.5$ s – followed by two fainter ones at $\sim T_0 + 3.5$ s and $\sim T_0 + 6.0$ s. The emission was detectable until ~ 30 s post-burst (see Fig. 1; Evans et al. 2009).

In the 15–350 keV band, BAT measured a peak energy $E_{\text{peak}} = 117 \pm 45$ keV with the 90 per cent of the burst fluence released during $T_{90} = 7.2 \pm 0.5$ s (Stamatikos et al. 2014). Both *Fermi* Gamma-ray Burst Monitor (GBM; Yu 2014) and *Konus-Wind* (KW; Golenetskii et al. 2014) detected the γ -ray prompt with similar T_{90} in the 50–300 keV and 20 keV–10 MeV energy range, respectively. KW detected the event up to ~ 4 MeV and measured a peak energy $E_{\text{peak}} = 139^{+10}_{-9}$ keV, isotropic energy $E_{\text{iso}} = (2.29 \pm 0.12) \times 10^{52}$ erg and a peak isotropic luminosity $L_{\text{iso}} = (2.89 \pm 0.04) \times 10^{52}$ erg s $^{-1}$ (Tsvetkova et al. 2017). Given the duration of the burst and the hardness ratio¹ $S(50\text{--}100\text{ keV})/S(25\text{--}50\text{ keV}) \sim 1.6$ (Lien et al. 2016), GRB 141220A is classified as a long-soft GRB type (Kouveliotou et al. 1993; Bromberg et al. 2013; Lien et al. 2016) – typically associated with a collapsar origin. This classification is further supported by $E_{\text{peak}}/T_{90} \sim 16$ keV s $^{-1}$ – the ratio is $E_{\text{peak}}/T_{90} \lesssim 50$ keV s $^{-1}$ for 99 per cent of long GRBs in BATSE sample (Shahmoradi & Nemiroff 2015).

A bright optical afterglow of 14.84 ± 0.17 mag was detected 86 s after BAT trigger at the GRB 141220A location by the 0.2-m SkyCam-Z telescope (see Section 2.2) – attached to the LT.

At 87.2 s after the burst, the *Swift* X-ray Telescope (XRT; Gompertz et al. 2014) also started observing the field. The XRT light curve has two distinct segments separated by a gap in observations: the first 57 s were done in Windowed Timing (WT) mode, whilst the steepest decay was observed in Photon Counting (PC) mode (see Fig. 1; Evans et al. 2009).

The 2-m LT reacted automatically to *Swift* alert (Guidorzi et al. 2006) with the three-band optical polarimeter and imager RINGO3 with observations starting at 129.5 s post-burst (see Section 2.1). LT observations consisted of 3×10 min epochs of RINGO3 instrument, followed by 6×10 s frames with the r band of the Optical Wide Field Camera (IO:O; see Section 2.3) and 7×10 min more with RINGO3.

At ~ 607 s post-burst, the 0.3-m *Swift* Ultraviolet/Optical Telescope (UVOT; Marshall & Cummings 2014) detected an optical counterpart of 17.30 ± 0.07 mag in the white band. At ~ 20.6 -min post-burst, the 10.4-m Gran Telescopio Canarias (GTC; de Ugarte Postigo et al. 2014b, a) detected GRB 141220A with 18.11 ± 0.05 mag in the r band and derived a spectral redshift of $z = 1.3195$.

¹The ratio between the fluences in the 50–100 and 25–50 keV energy bands.




Figure 1. The GRB 141220A multiwavelength light curves with *Swift* BAT, *Swift* XRT, LT SkyCam-Z *r*-equivalent band, LT RINGO3 *BV/R/I* bands, and LT IO:O *r* band. *Swift* observations are obtained from the web interface provided by Leicester University (Evans et al. 2009): BAT data are binned to S/N=7 and the absorbed 0.3–10 keV XRT light curve is converted into flux density at 1 keV. For completeness, we include the UV/optical observations reported in GCNs from UVOT (Marshall & Cummings 2014), GTC (de Ugarte Postigo et al. 2014b), and the MITSuME Akeno upper limits (Yano et al. 2014). The GCN observations do not include filter corrections. In the x-axis, T_0 corresponds to BAT trigger time; in the y-axis, the flux density is converted into RINGO3 *R* magnitude.

2.1 Calibration of RINGO3 *BV/R/I*-band observations

The RINGO3 instrument employs a spinning polaroid and two dichroic that split the light beam into three optical bands (Arnold et al. 2012). From 129.5 s post-burst, the source intensity was sampled every 1.15 s at eight polaroid angles by three fast-readout cameras. These 1.15 s exposures were automatically co-added by the telescope pipeline² into 10×1 -min integrations. Given RINGO3 instrumental configuration, the photometry is derived integrating the intensity of all eight frames (see Section 2.1.1) and the polarization degree and angle, measuring the relative intensity of the source at each of the eight polaroid positions (see Section 2.1.2). For both analyses, we use aperture photometry to derive the source flux with Astropy Photutils package (Bradley et al. 2016).

2.1.1 Photometric calibration of optical light curves

The signal-to-noise ratio (S/N) of the optical transient (OT) was high at the start of observations; the OT was detected at S/N \sim 30 in each of the first $\sim 25 \times 1.15$ s frames of the *BV* band. Given its brightness, we use the 1.15 s exposures instead of the 1-min integrations. We follow the same procedure as in Jordana-Mitjans et al. (2020) and we dynamically co-add the 1.15 s frames such that the OT reaches a minimum threshold of S/N = 10, which gives an ~ 0.4 – 0.1 mag photometric precision. *BV* band observations after ~ 550 s post-burst are the result of integrating frames; the *R/I* bands have lower S/N

and the frame co-adding starts at $\sim T_0 + 200$ s. The *BV* photometry after $\sim T_0 + 2000$ s is discarded because the OT fades beyond the S/N threshold; the source S/N is lower in the *R/I* bands and we only accept photometry up to ~ 1350 and ~ 1200 s post-burst, respectively. Integrations of lower S/N thresholds do not present statistically significant structure (within 3σ) in addition to the OT constant emission decay; frame binnings with thresholds S/N $\gg 10$ discard late-time photometry. We check the stability of RINGO3 during the OT observations with the stars in the field: an 11.5-mag star and three 15–19-mag stars. Using the OT binning for the photometry, the stars present an ~ 0.05 -mag deviation from the mean.

We calibrate the photometry in absolute flux in RINGO3 *BV/R/I* bandpass system following Jordana-Mitjans et al. (2020). Observations of three dereddened A0 type stars (HD 87112, HD 92573, HD 96781; Høg et al. 2000), including the GRB 141220A field, were scheduled via LT phase2UI³ using the same instrumental setup of the night of the burst. The observations were automatically dispatched during the nights of 2018 May 2–3. We correct for the mean Galactic extinction ($E_{B-V, MW} = 0.0128 \pm 0.0005$ is derived from a 5×5 arcmin² field statistic⁴; Schlegel, Finkbeiner & Davis 1998) and we use the RINGO3 magnitude-to-flux conversion from Jordana-Mitjans et al. (2020). This calibration adds ~ 0.05 -mag uncertainty to the photometry.

In Table 1 and Fig. 1, we present the photometry of GRB 141220A in RINGO3 *BV/R/I* bands. Note that magnitudes and

²<https://telescope.livjm.ac.uk/TelInst/Pipelines/>

³<http://telescope.livjm.ac.uk/PropInst/Phase2/>

⁴<https://irsa.ipac.caltech.edu/applications/DUST/>

Table 1. The GRB 141220A photometry in RINGO3 *BV/R/I* bands, SkyCam-Z *r*-equivalent band, and IO:O *r* band.

Band	t_{mid} (s)	$t_{\text{exp}}/2$ (s)	mag	mag _{err}	F_{ν} (Jy)	$F_{\nu \text{ err}}$ (Jy)
<i>BV</i>	130.1	0.6	15.73	0.08	1.86×10^{-3}	1.3×10^{-4}
<i>BV</i>	131.3	0.6	15.76	0.08	1.81×10^{-3}	1.3×10^{-4}
<i>BV</i>	132.4	0.6	15.78	0.08	1.77×10^{-3}	1.3×10^{-4}
<i>BV</i>	133.6	0.6	15.88	0.08	1.61×10^{-3}	1.2×10^{-4}
<i>BV</i>

t_{mid} corresponds to the mean observing time and t_{exp} to the length of the observation window. Magnitudes and flux density values are corrected for atmospheric and Galactic extinction. Table 1 is published in its entirety in the machine-readable format. A portion is shown here for guidance regarding its form and content.




Figure 2. The RINGO3 *BV/R/I* Stokes parameters (*q-u*) of the unpolarized standard stars (BD +32 3739, BD +33 2642, BD +28 4211, HD 14069, HD 109055, and G191-B2B) used for the determination of GRB 141220A instrumental polarization (circles). The red vertical line indicates GRB 141220A burst date (T_0) and the coloured horizontal lines correspond to the median *q-u* value per band.

fluxes are not corrected for the host galaxy extinction (see Section 3.3.2).

2.1.2 Calibration of optical polarization

We measure the flux of the OT at each of the eight rotor positions of the polaroid and, following Clarke & Neumayer (2002), we convert them to the Stokes parameters (*q-u*). The confidence levels of the Stokes parameters and the polarization degree and angle are determined with a Monte Carlo error propagation starting from 10^6 simulated flux values for each polaroid position.

RINGO3 instrumental polarization is subtracted in the Stokes parameters plane and the polarization angle is standardized to a reference (Słowikowska et al. 2016). As RINGO3 is regularly taking observations of standards stars, we choose observations taken during a period of ± 80 d on either side of the burst date T_0 (see Fig. 2). From 2014 October 1 to 2015 March 10, the following 9–11 mag unpolarized standards stars were observed with RINGO3: BD +32 3739, BD +33 2642, BD +28 4211, HD 14069, HD 109055, and G191-B2B. The instrumental polarization is determined with the median *q-u* of ~ 720 observations of the standards per RINGO3 band. We estimate an $\sim 3 \times 10^{-3}$ precision in the definition of the instrumental *q-u*, which could introduce no more than $P \sim 0.5$ per cent in polarization after their subtraction. We also check for any monotonic trend of the instrumental *q-u* during the $T_0 \pm 80$ d time-window; the Pearson’s correlation coefficients of each band are not significant with $|r| < 0.08$ and *p*-values > 0.1 .

Following Mundell et al. (2013) and Jordana-Mitjans et al. (2020), we co-add the 1.15-s frames of the first 10-min epoch so that the OT has maximum S/N. The *BV*-band provides the highest S/N for

the time window 129.5–204.3 s post-burst with polarization $P_{\text{BV}} = 2.8^{+2.0}_{-1.6}$ per cent at a 2σ confidence level. The OT is well detected at an average S/N ~ 60 in each of the eight polaroid position images. The remaining measurements of the *BV* band – and the *R/I* bands – are derived maximizing the S/N. The polarization angle is standardized to a reference using ~ 485 observations of 9–11.5-mag polarized standard stars taken during the $T_0 \pm 80$ d time-window: BD +25 727, BD +59 389, HD 155528, VI CYG 12, BD +64 106, and Hiltner 960 (Turnshek et al. 1990; Schmidt, Elston & Lupie 1992). GRB 141220A polarization measurements are presented in Table 2 with the polarization degree and the angle uncertainty quoted at 2σ confidence level.

We estimate RINGO3 depolarizing factor⁵ (D) during the $T_0 \pm 80$ d period using the ~ 485 observations of the $P = 4.5$ –9 per cent polarized standard stars and ~ 95 observations of the $P \sim 50$ per cent polarized planetary nebula of CRL 2688 (Shaw & Tarengi 1976). After *q-u* ellipticity corrections (Arnold 2017), RINGO3 depolarizing effect is negligible in the *BV* band ($D_{\text{BV}} = 1.00 \pm 0.01$) and small in the *R/I* bands ($D_{\{R,I\}} = 0.98 \pm 0.01, 0.94 \pm 0.01$). Because *R/I*-band polarization measurements are dominated by noise (resulting in upper limits), this correction is not applied to the polarization measurements of Table 2. We check the stability of RINGO3 instrument during observations using the bright star in the *R/I*-band field of view that has polarization $P_{\{R,I\}} = 0.22^{+0.13}_{-0.11}$ per cent, $0.23^{+0.18}_{-0.14}$ per cent. Using the OT binning, the polarization presents a $P_{\{R,I\}} \sim 0.07$ per cent deviation from the mean. We test the robustness of the polarization measurements with 1.5–3 FWHM photometric apertures for the OT and the bright star. There is no polarization variations within 1σ .

2.2 Photometric calibration of White-light SkyCam-Z observations

The SkyCam-Z telescope⁶ is attached to the LT mount and continuously monitors LT field of view with a $1 \times 1 \text{ deg}^2$ coverage in a white band – taking a single exposure of 10 s every minute. The SkyCam-Z started the first 10-s exposure of GRB 141220A field at 86 s post-burst, before RINGO3 observations. However, the LT telescope was still slewing towards the target at that time and the first frame has trails; we correct the flux offset of the first frame with the succeeding SkyCam-Z observation (at $T_0 + 145$ s) using 55 stars detected in both fields. Observations starting at $T_0 + 265$ s are co-added to reach a minimum threshold of S/N=5. We standardize the white-band photometry with the *r* band using five 8–12 mag stars from the SDSS DR12 catalogue (Alam et al. 2015). In Table 1 and Fig. 1, the photometry is corrected for the mean Galactic extinction ($A_r = 0.035 \pm 0.001$ mag) but not for host galaxy extinction (see Section 3.3.2).

2.3 Photometric calibration of *r*-band IO:O observations

The IO:O⁷ observations started at $T_0 + 2002$ s with the *r* filter. Given that the OT is detected at S/N ~ 100 per frame, we individually derive the OT flux. We standardize the photometry using ten 12–17 mag stars from the SDSS DR12 catalogue (Alam et al. 2015). In Table 1 and Fig. 1, we present the IO:O photometry in the *r* band.

⁵Note that P/D is the true polarization such that an instrument with no depolarization has $D = 1$.


⁶<https://telescope.livjm.ac.uk/TelInst/Inst/SkyCam/>

⁷<https://telescope.livjm.ac.uk/TelInst/Inst/IOO/>

Table 2. The GRB 141220A polarization observations with RINGO3 *BV/R/I* bands.

Band	t_{mid}	$t_{\text{exp}}/2$	S/N	q	q_{err}	u	u_{err}	P	P_{err}	θ	θ_{err}
	(s)	(s)						(per cent)	(per cent)	($^{\circ}$)	($^{\circ}$)
<i>BV</i>	166.9	37.4	60	-0.017	0.009	0.023	0.009	2.8	$^{+2.0}_{-1.6}$	63	20
<i>BV</i>	253.8	49.5	42	0.006	0.014	0.014	0.014	<4.5	—	—	—
<i>BV</i>	514.4	211.1	38	-0.008	0.015	0.014	0.015	<4.9	—	—	—
<i>BV</i>	1043.0	298.0	21	-0.009	0.026	0.008	0.026	<7.5	—	—	—
<i>BV</i>	1658.5	298.0	11	-0.042	0.051	0.022	0.051	<16.4	—	—	—
<i>R</i>	166.9	37.4	30	-0.006	0.018	-0.005	0.018	<5.2	—	—	—
<i>R</i>	282.0	77.7	23	-0.004	0.024	-0.005	0.024	<6.7	—	—	—
<i>R</i>	542.6	182.9	17	0.034	0.033	0.020	0.033	<11.4	—	—	—
<i>R</i>	1043.0	298.0	11	0.083	0.049	-0.005	0.049	<19.2	—	—	—
<i>R</i>	1658.5	298.0	7	0.007	0.085	-0.122	0.085	<31.4	—	—	—
<i>I</i>	166.9	37.4	23	-0.022	0.024	0.016	0.024	<8.1	—	—	—
<i>I</i>	282.0	77.7	17	-0.016	0.033	0.010	0.033	<9.8	—	—	—
<i>I</i>	542.5	182.9	14	-0.007	0.040	0.061	0.040	<14.9	—	—	—
<i>I</i>	1043.0	298.0	9	-0.086	0.063	-0.014	0.063	<22.8	—	—	—
<i>I</i>	1658.5	298.0	6	0.013	0.088	0.120	0.088	<32.0	—	—	—

^a P and θ uncertainties and upper limits are quoted at 2σ confidence level.

**Figure 3.** Temporal evolution of GRB 141220A polarization in RINGO3 *BV/R/I* bands with a colour map of the OT S/N. In the *BV* band, the black marker corresponds to the only polarization measurement.

The IO:O light curve is corrected for the mean Galactic extinction ($A_r = 0.035 \pm 0.001$ mag) but not for host galaxy extinction (see Section 3.3.2).

3 RESULTS

In this section, we present the time-resolved polarization measured in three optical bands (see Section 3.1), the optical light curves in three bands, and a white band (see Section 3.2), the spectral evolution in the optical bands (see Section 3.3.1) and the combined optical/X-rays spectral energy distributions (SEDs; see Section 3.3.2).

3.1 Polarization in three optical bands

We detect low polarization in RINGO3 *BV/R/I* bands since the start of observations at 129.5 s post-burst (see Fig. 3). The best constraint is the polarization measurement $P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent at 2σ confidence level, with polarization angle $\theta_{BV} = 63^{\circ} \pm 20^{\circ}$ (2σ). This measurement corresponds to the time window 129.5–204.3 s post-burst, i.e. 55.9–88.1 s after the burst in the GRB rest frame. Additionally, we use the permutation test detailed in Steele et al. 2017 to prove that the observed polarization is a measurement and not the

Table 3. Temporal (α) and spectral indexes (β) of GRB 141220A emission.

Index	Result	χ^2/dof	Fig.
α_{BV}	1.095 ± 0.005	657.7/535	—
α_r	1.09 ± 0.03	7.4/9	—
α_R	1.067 ± 0.009	203.7/220	—
α_I	1.105 ± 0.013	122.6/136	—
α_{opt}	1.091 ± 0.008	999.2/903	4
$\beta_{\text{X-rays}}^a$	1.1 ± 0.2	21.2/17	5
β_{opt}^a	0.6 ± 0.2	—	5
β_{opt}^{*b}	1.74 ± 0.11	26.2/31	4

^a The spectral indexes are linked as $\beta_{\text{X-rays}} = \beta_{\text{opt}} + 0.5$.

^b β_{opt}^* is not corrected for host galaxy extinction; β_{opt}^* is an upper limit of β_{opt} .

result of random noise in an unpolarized source. Consequently, we randomly reorder the eight flux values, corresponding to each angle position of the polaroid, to create a sample of 40 320 polarization values with the same S/N properties of the observed polarization. If the observed polarization lays within the simulated distribution ($p_t \lesssim 0.75$), it indicates that there is high probability of being the result of the random noise. The observed polarization $P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent belongs to the $p_t = 0.97$ upper quantile of the overall distribution; there is only 3 per cent probability that this measurement is consistent with zero polarization.

In the *BV* band, the polarization is further constrained at $P_{BV} < 5$ per cent (2σ) level until 726 s post-burst. Afterwards, the 2σ upper limits increase significantly with time due to the decreasing brightness of the OT, which cannot be compensated by integrating more frames. The *R/I* bands have less S/N and the observed polarization is within the instrument sensitivity, ranking $p_t < 0.75$ in the polarization permutation test. The best constraints in the *R/I* bands are the 2σ upper limits $P_{R,I} < 5.2$ per cent, 8.1 per cent during 129.5–204.3 s post-burst – consistent with the cotemporal *BV*-band measurement.

3.2 Optical light curves

The *BV/R/I* light curves are satisfactorily modelled with power laws that have compatible decay indexes (see Table 3). We note that the




Figure 4. The GRB 141220A *BV/r*-equivalent/*R/I* band light curves fitted with power laws of common slope $\alpha_{\text{opt}} = 1.091 \pm 0.008$. In the x -axis, T_0 corresponds to BAT trigger time. In the y -axis, the flux density is converted into RINGO3 *R* magnitude. The middle panel corresponds to the residuals of the fit. The bottom panel shows the evolution of the optical spectral index β_{opt}^* ; the dashed line is the mean. Note that β_{opt}^* is not corrected by the host galaxy extinction.

BV band has worse χ^2 statistics than the *R/I* bands due to smaller photometric uncertainties⁸ (higher S/N). The SkyCam-Z *White* and IO:O *r*-band observations are also fitted in a joint *r*-equivalent light curve. If we simultaneously fit all light curves with a single power law, the best-fitting model indicates that the emission was decaying as $\alpha_{\text{opt}} = 1.091 \pm 0.008$ since 86 s and up to ~ 2200 s after the burst (see Fig. 4).

3.3 Spectral evolution of the emission

3.3.1 Optical spectral index

Given RINGO3 simultaneous three-band imaging, we study the evolution of the optical spectral index. We follow Jordana-Mitjans et al. (2020) procedure and we co-add the 1.15-s frames of the lowest S/N band (*I* band) to reach a minimum threshold of S/N=20, which gives $\sim 0.05 - 0.07$ mag spectral precision through observations and discards photometry after ~ 700 s post-burst. We use the *I*-band binning to co-add the *BV/R*-band frames. We subtract the Galactic extinction ($E_{B-V, \text{MW}} = 0.0128 \pm 0.0005$; Schlegel et al. 1998) from the flux-calibrated photometry. For each temporal bin, we fit a power law to the three-band SED and we obtain the spectral index β_{opt}^* .

⁸The statistical uncertainty of the *BV* band is likely underestimated. For the first ~ 300 s of the light curve, it is smaller than the ~ 0.05 -mag deviation we measure in field stars.

The evolution of the optical spectral index is consistent with a mean $\beta_{\text{opt}}^* = 1.74 \pm 0.11$ ($\chi^2/\text{dof}=26.2/31$ and $p\text{-value}=0.7$); the optical emission does not suffer any significant spectral change during 130–725 s post-burst (see Fig. 4 bottom panel). Note that the spectral index β_{opt}^* is not corrected by the host galaxy extinction and, therefore, it is an upper limit of the true optical spectral index β_{opt} .

3.3.2 Broadband SED

To infer the optical to X-rays synchrotron spectrum, we analyse the broad-band SEDs using the time-resolved 0.3–10 keV XRT and RINGO3 observations. We extract the XRT spectra that have cotemporal RINGO3 observations using the web interface provided by Leicester University⁹ based on HEASoft (v. 6.22.1; Blackburn 1995). We then obtain a single RINGO3 photometric measurement per band by co-adding the 1.15-s frames that correspond to the XRT time interval. XRT has a gap of observations between WT and PC mode; consequently, the two broad-band SEDs correspond to the time intervals 129–148 and 1589–1957 s.

We model the combined optical/X-ray data with XSPEC (v. 12.9.1; Arnaud, Dorman & Gordon 1999) using χ^2 statistics. To ensure the Gaussian limit, we use the GRPPHA tool to group the XRT spectra such that there are at least 20 counts per bin. The models we employ are either power laws (powerlaw in XSPEC) or connected power laws (bknpower) that account for the rest-framed host galaxy and MW total hydrogen column absorption and dust extinction (phabs*zphabs*zdust*zdust). For both SEDs, we fix the MW dust extinction to $E_{B-V, \text{MW}} = 0.0128 \pm 0.0005$ (Schlegel et al. 1998), the total hydrogen column absorption to $N_{\text{H, MW}} = 1.33 \times 10^{20} \text{ cm}^{-2}$ (derived using the NHTot Swift tool¹⁰; Willingale et al. 2013) and the GRB redshift to $z = 1.3195$ (de Ugarte Postigo et al. 2014a). Given that the number of data bins is low compared to the number of parameters of the model, we further reduce the model complexity. We assume no temporal evolution for the host galaxy dust and total hydrogen absorption, and we also pair the optical and X-ray spectral indexes of both SEDs. Additionally, we assume an Small Magellanic Cloud (SMC) profile for the host galaxy dust absorption (Schady et al. 2007). For the broken power law model, we also link the optical and the X-rays indexes as $\beta_{\text{X-rays}} = \beta_{\text{opt}} + 0.5$, following closure relations for slow cooling synchrotron spectrum with the cooling break between the optical and X-rays bands (e.g. Sari, Piran & Narayan 1998; Gao et al. 2013). We leave the energy break (E_{break}) as a free parameter constrained between the optical and the X-ray bands. The data are best fitted with a broken power law model with $\chi^2/\text{dof} = 21.2/17$ and $p\text{-value} = 0.2$ (see Fig. 5). In comparison, a power law model yields worse χ^2 statistics ($\chi^2/\text{dof}=39.5/19$ and $p\text{-value}=0.004$). The best-fitting parameters for the broken power law model are a spectral index $\beta_{\text{opt}} = 0.6 \pm 0.2$ (i.e. $\beta_{\text{X-rays}} = 1.1 \pm 0.2$), energy breaks at $E_{\text{break},1} = 0.08^{+0.04}_{-0.03} \text{ keV}$ and $E_{\text{break},2} = 0.015 \pm 0.005 \text{ keV}$. In the host galaxy rest frame, the reddening is $E_{B-V, \text{HG}} = 0.24 \pm 0.06$ (i.e. absorption of $A_{V, \text{HG}} = 0.71 \pm 0.15 \text{ mag}$) and the total hydrogen absorption is $N_{\text{H, HG}} = (3 \pm 2) \times 10^{21} \text{ cm}^{-2}$. MW and Large Magellanic Cloud (LMC) dust profiles yield compatible results with $A_{V, \text{HG}} = 0.74 \pm 0.18 \text{ mag}$ and $A_{V, \text{HG}} = 0.76 \pm 0.16 \text{ mag}$, respectively.

⁹https://www.swift.ac.uk/user_objects/

¹⁰<https://www.swift.ac.uk/analysis/nhtot/>




Figure 5. The GRB 141220A broad-band SEDs with RINGO3 and XRT observations (Evans et al. 2009). The SEDs are best fitted with a broken power law model (solid line) that accounts for extinction in the optical and total hydrogen absorption in the X-rays (dashed line). The results of the fit are $\beta_{\text{opt}} = 0.6 \pm 0.2$, $\beta_{\text{X-rays}} = 1.1 \pm 0.2$, $E_{\text{break},1} = 0.08^{+0.04}_{-0.03}$ keV, $E_{\text{break},2} = 0.015 \pm 0.005$ keV, and in the host galaxy rest frame, $A_{V,\text{HG}} = 0.71 \pm 0.15$ mag and $N_{\text{H,HG}} = (3 \pm 2) \times 10^{21} \text{ cm}^{-2}$ ($\chi^2/\text{dof}=21.2/17$ and $p\text{-value}=0.2$). The error bars in the optical frequencies are the FWHM of RINGO3 *BV/R/I* bandpasses. The bottom panel corresponds to the residuals of the fit.

4 DISCUSSION

Here, we discuss the nature of the fireball, shock properties, and magnetic field structures in GRB 141220A, inferred from the measured properties of the observed light curves and optical polarization. In Section 4.1, we show that from the very start of our observations – beginning 86 s post-GRB – the afterglow properties are consistent with those exhibited by classical forward shocks observed in optical afterglows at late times – hours to days after a GRB. In Section 4.2, we discuss all possible contributions to the low polarization of $P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent (2σ) from 129.5 to 204.3 s post-burst and conclude that the polarization of the afterglow emission itself is very small. GRB 141220A polarization is expected from theoretical predictions but is at odds with the claim by Uehara et al. (2012) of polarized forward shock emission in GRB 091208B. In Section 4.3, we compare the multiwavelength properties of GRB 141220A and GRB 091208B, and put their properties into a wider context of other GRBs with dominant forward shocks emission and optical polarization data in the first hour after the burst.

4.1 A classical forward shock at early times

The optical emission of GRB 141220A decays with a temporal index $\alpha_{\text{opt}} = 1.091 \pm 0.008$ in the *BV/r*-equivalent/*R/I* bands during LT observations (86–2200 s post-burst), which is in agreement with $\alpha_r = 0.96 \pm 0.11$ reported from GTC observations at 20.6–38.7 min post-burst (de Ugarte Postigo et al. 2014a). There is also no evidence of additional spectral components in the optical as there is no colour evolution through RINGO3 bands (see Fig. 4 bottom panel).

The *Swift* XRT X-ray emission can be modelled with a broken power law¹¹ with $\chi^2/\text{dof} = 14.8/25$ and $p\text{-value}=0.9$. It consists of plateau-like emission ($\alpha_{\text{X-rays},1} = -0.25 \pm 0.18$) followed by a steepening of the light curve ($\alpha_{\text{X-rays},2} = 1.39 \pm 0.08$) at 214^{+16}_{-29} s post-burst, which we identify as the normal afterglow decay phase (Nousek et al. 2006).

The optical and the X-rays domains satisfy closure relations for interstellar medium (ISM) profile, slow cooling regime and with the cooling break frequency (ν_c) laying in between the bands ($\nu_{\text{opt}} < \nu_c < \nu_{\text{X-rays}}$; Sari et al. 1998; Gao et al. 2013). The expected decay rate difference $\Delta\alpha = \alpha_{\text{X-rays}} - \alpha_{\text{opt}} = 0.25$ is consistent with the need of a spectral break in the SED modelling (Section 3.3.2). The cooling frequency is expected to evolve like $\nu_c \propto t^{-\alpha_c}$ with $\alpha_c = 0.5$, which is compatible with $\alpha = 0.7 \pm 0.4$ derived from the two SEDs epochs. In closure relations conditions, the optical decay $\alpha_{\text{opt}} = 1.091 \pm 0.008$ corresponds to a typical electron index $p_{\text{CR}} = 2.45$ (e.g. Spitkovsky 2008; Curran et al. 2010), a X-rays temporal decay of $\alpha_{\text{X-rays,CR}} = 1.34$ and spectral indexes $\beta_{\text{opt,CR}} = 0.73$ and $\beta_{\text{X-rays,CR}} = 1.23$. These are in agreement with the ones measured: $\alpha_{\text{X-rays},2} = 1.39 \pm 0.08$ and $\beta_{\text{opt}} = 0.6 \pm 0.2$ (i.e. $\beta_{\text{X-rays}} = 1.1 \pm 0.2$).

4.2 What can contribute to the polarization of early-time forward shocks?

Forward shock emission was predicted theoretically to be unpolarized (Medvedev & Loeb 1999), as the external shock sweeps up circumburst material containing weak, tangled magnetic fields. This prediction was confirmed observationally with detections of low polarization $P \sim 1\text{--}3$ per cent in late-time afterglow measurements made hours to days after the burst (Covino et al. 1999; Hjorth et al. 1999; Wijers et al. 1999; Rol et al. 2000). These low polarizations were not thought to be intrinsic to the afterglow emission but to be associated with ISM dust or jet geometry effects (Covino et al. 2004). However, the early-time polarization detection of $P = (10.4 \pm 2.5)$ per cent at 149–706 s post-burst for GRB 091208B forward shock started a debate on the intrinsic polarization of forward shocks during the first $\sim 10^3$ s after the burst (Uehara et al. 2012). In contrast, our observations of GRB 141220A forward shock measure low polarization levels at early-times: $P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent (2σ) from 129.5 to 204.3 s post-burst and $P_{BV} < 5$ per cent (2σ) up to 726 s post-burst, which are comparable to late-time studies.

GRB 141220A is interesting for polarization observations at early times due to its brightness and a light-curve decay typical of late-time forward shocks ($\alpha_{\text{opt}} \sim 1$; e.g. Waxman 1997). The optical emission does not present statistically significant ‘bumps’ at early times (within 3σ) – usually associated with inhomogeneities in the jet or the ISM (Lazzati et al. 2002; Nakar & Piran 2003) – which reduces any issues regarding a possible break of the fireball symmetry that could introduce additional polarization signals (Granot & Königl 2003; Lazzati et al. 2003; Nakar & Oren 2004). We now discuss possible mechanisms that could produce the observed polarization at very early times.

4.2.1 Polarization caused by the jet geometry

Due to the relativistic beaming effect, only a small region of the physical size of the jet is visible ($1/\Gamma < \theta_j$). If the forward shock emission contains tangled magnetic fields in the plane of the shock

¹¹ A power law fit presents worse χ^2 statistics: $\chi^2/\text{dof} = 51.8/26$ and $p\text{-value} = 0.002$.




Figure 6. Left-hand panel: In grey, the GRBs duration (T_{90}) and the hardness ratio of *Swift* BAT sample (Kouveliotou et al. 1993; Lien et al. 2016). Right-hand panel: In grey, the GRBs fluence in the 10 keV–10 MeV band and the peak energy (E_{peak}) of KW sample (Tsvetkova et al. 2017, 2020). In colour, GRBs with early-time polarization measurements and significant forward shock contribution. We note that GRB 091208B fluence corresponds to the 8 keV–35 MeV energy range of *Fermi* GBM (Nava et al. 2012); GRB 100805A fluence and E_{peak} are derived from the 15–350 keV *Swift* BAT time-averaged spectrum and constrained fitting a Band function with the low-energy photon index fixed to $\alpha = -1$ and the high-energy photon index to $\beta < -2$ ($\chi^2/\text{dof}=11.7/9$ and $p\text{-value}=0.2$).

with a coherence length of the order of plasma skin depth scales, the polarization is cancelled out for a face-on jet (Medvedev & Loeb 1999). However, high polarization is expected when the spherical symmetry is broken. For a homogeneous jet, the polarization peaks before and after the jet break (Ghisellini & Lazzati 1999; Sari 1999) – when the ejecta has slowed down to $1/\Gamma \sim \theta_j$ (when on-axis) and the observer notices the edge of the jet (see also Rossi et al. 2004 for the case of structured jets).

For GRB 141220A, there are no signs of a steepening of the X-rays light curve up to $\sim 3 \times 10^4$ s post-burst. It is highly unlikely that we are detecting net polarization from a jet break during the first minutes after the explosion (Racusin et al. 2009) – requiring very small electron index. Additionally, edge-on/off-axis GRBs are predicted to have softer prompt emission, lower fluence and E_{peak} (e.g. Yamazaki, Ioka & Nakamura 2002; Yamazaki, Yonetoku & Nakamura 2003; Yamazaki, Ioka & Nakamura 2004; Beniamini & Nakar 2019; Gill, Granot & Kumar 2020); GRB 141220A is on the average of KW sample (see Fig. 6 right-hand side; Tsvetkova et al. 2017, 2020).

4.2.2 Post-shock induced polarization: the GRB environment

ISM dust can induce non-negligible levels of polarization to an unpolarized source via dichroic absorption (Serkowski, Mathewson & Ford 1975). The light is preferably absorbed when the electric field is parallel to the long axis of the dust grains. Because the short axis aligns with the magnetic field, the polarization traces the projected direction of the local magnetic field and the physical properties of dust grains.

The wavelength-dependent behaviour of polarization was characterized with MW stars by the Serkowski empirical relation $P = P_0 \exp[-K \ln^2(\lambda_{\text{max}}/\lambda)]$ (Serkowski et al. 1975; Whittet et al. 1992), where $\lambda_{\text{max}}(\mu\text{m}) = R_V/5.5$ is the wavelength at which the polarization is maximum, $K = 0.01 \pm 0.05 + (1.66 \pm 0.09)\lambda_{\text{max}}$ and $P_0 \lesssim 9 E_{B-V}$ is an upper limit on the induced polarization

that depends on the number of dust grains and the efficiency of the alignment. More recently, there have been indications that this limit is higher ($P_0 \lesssim 13 E_{B-V}$) at low column densities in the MW ($N_H < 5 \times 10^{20} \text{ cm}^{-2}$; Planck Collaboration XII 2018b; Panopoulou et al. 2019). Additionally, Fosalba et al. (2002) found that for the range $0.01 < E_{B-V} < 1$, the average polarization dependency with extinction can be fitted with $\langle P_0 \rangle = 3.5 E_{B-V}^{0.8}$.

The 11.5 mag star in the *R/I* band field-of-view presents polarization $P_{[R,I], \text{MW}} = 0.22^{+0.13}_{-0.11}$ per cent, $0.23^{+0.18}_{-0.14}$ per cent (2σ), which gives an estimate of the polarization induced by the Galactic ISM in the GRB 141220A line of sight. We note that the measured polarization agrees with the maximum expected $P_{[R,I], \text{MW}} \lesssim 0.29$ per cent (2σ) for $E_{B-V, \text{MW}} = 0.0128 \pm 0.0005$ at low column densities ($N_{H, \text{MW}} = 1.33 \times 10^{20} \text{ cm}^{-2}$). Using the Serkowski law, we determine that the polarization induced by the ISM in the *BV* band is $P_{BV, \text{MW}} = 0.23$ per cent.

In addition to the polarization induced by the Galactic dust, we have the contribution induced by the propagation of the light along the host galaxy ISM. GRB 141220A is within the top 30 per cent of the most extinguished GRBs of Covino et al. (2013) sample, with $A_{V, \text{HG}} = 0.71 \pm 0.15$ mag (i.e. $E_{B-V, \text{HG}} = 0.24 \pm 0.06$; Section 3.3.2). To check GRB 141220A host galaxy dust contribution to the intrinsic afterglow polarization, we assume that the redshifted Serkowski law is valid for the host galaxy SMC profile¹² with $P_0 \lesssim 9 E_{B-V}$ (e.g. Klose et al. 2004). Taking into account the dust-absorbed spectral slope of GRB 141220A (β_{opt}^* ; Section 3.3.1), we estimate that the maxi-

¹²SMC and LMC polarimetric studies found that the Serkowski law was still valid – with smaller λ_{max} on average in the SMC (Clayton, Martin & Thompson 1983; Clayton et al. 1996; Rodrigues et al. 1997) and polarimetric studies of M31 globular clusters measured $P_{\text{obs}} \sim [7.1, 15.3] E_{B-V}$ with a different $K(\lambda_{\text{max}})$ relationship (Clayton et al. 2004).

mum induced polarization in RINGO3 bands is¹³ $P_{\{BV,R,I\},\text{HG}} \lesssim 1.8$ per cent, 2.5 per cent, 2.8 per cent (2σ), with mean polarization $\langle P_{\{BV,R,I\},\text{HG}} \rangle \sim 0.6$ per cent, 0.8 per cent, 1.0 per cent.

As suggested in late-time (Lazzati et al. 2003; Covino et al. 2004; Wiersema et al. 2012) and early-time afterglow polarization studies (Jordana-Mitjans et al. 2020), differential light extinction from dust in the host galaxy can account for the observed polarization levels ($P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent). In Sections 4.2.3 and 4.2.4, we discuss the implications of a residual polarization intrinsic to GRB 141220A afterglow.

4.2.3 Contamination from a polarized reverse shock

During the early afterglow, we expect contributions from the reverse and the forward shock. GRB 141220A optical light curves present no signature of reverse shock emission and the temporal and spectral properties satisfy closure relations for forward shock emission (Section 4.1). The absence of a reverse shock at early times suggests that is peaking at lower frequencies (Mundell et al. 2007b; Laskar et al. 2013; Kopač et al. 2015) or high magnetization in the jet, which suppresses the afterglow emission of the reverse shock (Zhang & Kobayashi 2005).

The expected polarization of synchrotron emission is $P_0 = (p_{\text{CR}} + 1)/(p_{\text{CR}} + 7/3) = 72$ per cent for a homogeneous field in the slow cooling regime and for an electron power law index $p_{\text{CR}} = 2.45$ (inferred from closure relations in Section 4.1; Rybicki & Lightman 1979). However, if we take into account the depolarization caused by the rotation of the polarization vector of each element in a shell due to relativistic kinematic effects, the maximum polarization of the reverse shock is reduced to $P_0 \sim 50$ per cent (Lyutikov et al. 2003). Observationally, we have measured polarization in the range $P = 10\text{--}30$ per cent for mildly magnetized jets (GRB 090102, Steele et al. 2009; GRB 120308A, Mundell et al. 2013). Given the low polarization levels in GRB 141220A afterglow, we rule out significant contamination from the reverse shock.

The deceleration peak of the blastwave must have been at $t_{\text{dec}} \ll 86$ s after the GRB, well before SkyCam-Z observations; otherwise, it would affect the shape of the observed light curve. From the light-curve modelling,¹⁴ a $P \sim 30\text{--}50$ per cent polarized reverse shock contributing equally at the deceleration time (Mundell et al. 2007a, b) reproduces the measured polarization levels ($P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent) for peak times at $\lesssim 20\text{--}30$ s post-burst. We note that later deceleration times are permitted if the reverse shock is suppressed and/or less polarized than the expected (Laskar et al. 2019; Jordana-Mitjans et al. 2020).

4.2.4 Polarization from collisionless shocks

In matter-dominated jets, the magnetic fields do not influence the fireball dynamics but they still play a crucial role in efficient synchrotron emission and particle acceleration in collisionless shocks. The afterglow modelling usually indicates equipartition magnetic energy densities of $\epsilon_B \sim 10^{-6}\text{--}10^{-1}$ (Panaitescu & Kumar 2002; Beniamini et al. 2015; Wang et al. 2015). However, for the forward shock, we are left with the magnetic field of the circumburst medium

¹³MW and LMC profiles yield similar results: $P_{BV,\text{HG}} \lesssim 1.6$ per cent and $P_{BV,\text{HG}} \lesssim 1.5$ per cent, respectively.

¹⁴We fit the optical light curves with a reverse/forward shock model assuming ISM, slow cooling regime and the spectral configuration $\nu_{\text{m},r} \ll \nu_{\text{m},f} < \nu_{\text{opt}} < \nu_{\text{c},r} \leq \nu_{\text{c},f}$.

that has a few μG ($\epsilon_B \leq 10^{-11}$ for ISM; Medvedev & Loeb 1999). Therefore, in order to increase the magnetic energy density several orders of magnitudes, we need some mechanism to amplify the magnetic field that cannot be achieved by shock compression alone, e.g. via plasma instabilities (e.g. Weibel instability; Medvedev & Loeb 1999; Nishikawa et al. 2009) or by macroscopic turbulences due to preshock density inhomogeneities (Sironi & Goodman 2007; Inoue, Asano & Ioka 2011). In both cases, the magnetic field will have coherent length-scales (l_B) smaller than the observable emission area of the jet ($\theta_B \ll 1/\Gamma < \theta_j$), e.g.: $l_B \sim 10^5$ cm are expected for Weibel instability (Medvedev & Loeb 1999) and $l_B \sim 10^{13}$ cm for macroscopic turbulences (Sironi & Goodman 2007). Magneto-hydrodynamics simulations predict $P \leq 2$ per cent for Richtmyer–Meshkov instability (Inoue et al. 2011), which is in agreement with GRB 141220A forward shock polarization.

Observationally, we can estimate the coherent length-scale of the magnetic field assuming a polarization model with magnetic domains (Gruzinov & Waxman 1999; Uehara et al. 2012). If each patch contains ordered magnetic fields, the local polarization would be the theoretical expected for synchrotron emission when accounting for relativistic kinematic effects $P_0 \sim 50$ per cent (Lyutikov et al. 2003). Because the polarization angle is randomly oriented in each magnetic domain, the total observed polarization will be reduced by the number of visible patches (N) as $P = P_0/\sqrt{N}$. The observable region of the jet ($\sim R/\Gamma$) increases when the ejecta decelerates – where R is the emission radius and Γ is the Lorentz factor the blastwave. Therefore, more patches will be visible with time, which will cause random fluctuations in the polarization angle and the polarization degree will gradually decrease. Following Uehara et al. (2012) derivation, we estimate the coherent length-scales of the magnetic field as $l_B \sim R/(\Gamma\sqrt{N})$ given that $\Gamma \sim 260 E_0^{1/8} n_0^{-1/8} t_{\text{rest}}^{-3/8}$ and $R \sim 3.2 \times 10^{16} E_0^{1/4} n_0^{-1/4} t_{\text{rest}}^{1/4}$ (Sari 1997). Consequently,

$$l_B \sim 10^{14} \left(\frac{E_0}{10^{52} \text{ erg}} \right)^{1/8} \left(\frac{n_0}{1 \text{ cm}^{-3}} \right)^{-1/8} \left(\frac{t_{\text{rest}}}{1 \text{ s}} \right)^{5/8} P P_0^{-1} \text{ cm}, \quad (1)$$

where E_0 is the isotropic-equivalent kinetic energy, n_0 the circumburst medium density and t_{rest} is the rest-frame time after the burst. E_0 is related to the isotropic energy released in γ -rays as $E_{\text{iso}} = \eta E_0$; we adopt $\eta = 0.2$ for the radiative efficiency parameter. If we assume that the observed polarization $P_{BV} = 2.8^{+2.0}_{-1.6}$ per cent is intrinsic to the forward shock and $E_{\text{iso}} = (2.29 \pm 0.12) \times 10^{52}$ erg (Tsvetkova et al. 2017), the coherent length-scale of the magnetic field is $l_B \sim 10^{14} n_0^{-1/8}$ cm.

4.3 Polarization properties of forward shocks

Very early-time follow-up is still rare and dominated by small robotic telescopes, hence there is only a small sample of bright optical afterglows with polarization measurements (e.g. Steele et al. 2017). For those GRBs with polarization measurements, optical light curves consistent with a power law decay and with the emission interpreted as forward shock (at least 50 per cent of the total photons), we summarize their rest-frame properties in Fig. 7 and Table 4.

For all GRBs in Table 4 sample, the polarization observations are well after the end of prompt emission ($t_{\text{obs}}/T_{90} > 1$) and no contamination from prompt photons is expected (e.g. GRB 160625B, Troja et al. 2017; GRB 190114C, Jordana-Mitjans et al. 2020). We note that GRB 120327A optical light curve has a similar decay to GRB 141220A with $\alpha_{\text{opt}} \sim 1.2$ (Steele et al. 2017), GRB 091208B/GRB 100805A/GRB 131030A have shallower light curves with $\alpha_{\text{opt}} \sim 0.8$ (Uehara et al. 2012; King et al. 2014; Steele et al.




Figure 7. Rest-frame polarization observations in the optical of GRBs with light curves interpreted as early-time forward shocks. The vertical dotted lines correspond to the upper limits of the deceleration time (the earliest optical detection). The solid vertical line of GRB 060418 is the peak time of the fireball deceleration. *References* are as follows: GRB 060418 (Mundell et al. 2007a; Molinari et al. 2007), GRB 091208B (Nakajima et al. 2009; Uehara et al. 2012), GRB 100805A (Steele et al. 2017), GRB 120327A (Steele et al. 2017; Melandri et al. 2017), and GRB 131030A (Breeveld & Troja 2013; King et al. 2014).

2017) and GRB 060418 measurement was taken during the broad peak of the fireball deceleration (decaying later with $\alpha_{\text{opt}} \sim 1.3$; Mundell et al. 2007a; Molinari et al. 2007). Table 4 GRBs are characterized as long in the duration-hardness clustering of *Swift* BAT sample (Kouveliotou et al. 1993; Lien et al. 2016) – with GRB 141220A being the shortest and one of the hardest (see Fig. 6 left-hand panel). GRB 141220A fluence and peak energy is on the average of the *Swift* BAT sample (see Fig. 6 right-hand panel). They also follow the Amati relation ($E_{\text{iso}}-E_{\text{peak, rest}}$ correlation; Amati et al. 2002) within 1σ and 2σ of the scatter around the best-fitting line of Tsvetkova et al. (2020) and Nava et al. (2012), respectively. To derive the bulk Lorentz factor (Γ_{dec} ; Table 4), we assume ISM profile and thin shell regime and we use Sari & Piran (1999) derivation

$$\Gamma_{\text{dec}} = \left(\frac{3E_0}{32\pi m_p c^5 n_0} \right)^{1/8} t_{\text{dec, rest}}^{-3/8}, \quad (2)$$

where $t_{\text{dec, rest}}$ is the deceleration time of the fireball and m_p is the proton mass. The E_{iso} , $E_{\text{peak, rest}}$, and Γ_{dec} limits of Table 4 GRBs are compatible within 1σ of Ghirlanda et al. (2018) sample median.

GRB 120327A (Steele et al. 2017) and GRB 131030A (King et al. 2014) measured low values of polarization for forward shock emission, similar to our early-time result for GRB 141220A (see Fig. 7). Furthermore, King et al. (2014) also interpreted the polarization measurements to be dust-induced (from the MW). However, GRB 091208B polarization is significantly higher than for those GRBs (Uehara et al. 2012). GRB 091208B and GRB 141220A have similar high-energy properties (E_{iso} , E_{peak} , and fluence) but different polarization and afterglow decays (i.e. electron index; see Table 4). Uehara et al. (2012) measured $P = (10.4 \pm 2.5)$ per cent for GRB 091208B forward shock at a rest-frame time $t_{\text{rest}} \sim 207$ s, which corresponds to $l_{\text{B, grb09}} \sim 7 l_{\text{B, grb14}} (n_{0, \text{grb09}}/n_{0, \text{grb14}})^{-1/8}$ (see equation 1). This magnetic field length-scale predicts polarization $P \sim 20$ per cent for GRB 141220A at the time of observations – far in excess of that measured. Therefore, given that GRB 091208B and GRB 141220A have similar E_{iso} , either the magnetic field length-scale is almost an order of magnitude smaller in GRB 141220A or the circumburst medium density is 10^6 times greater in GRB 091208B

– such that Uehara et al. (2012) observed a smaller emitting region. However, we note that this would result in significant differences in both afterglows. Without the fine tuning of the microphysical parameters in

$$\nu_{\text{c, f}} = 6.3 \times 10^{15} n_0^{-1} (1+z)^{-1/2} (1+Y)^{-2} \left(\frac{\epsilon_B}{10^{-2}} \right)^{-3/2} \times \left(\frac{E_0}{10^{52} \text{ erg}} \right)^{-1/2} \left(\frac{t}{\text{day}} \right)^{-1/2} \text{ Hz} \quad (3)$$

and ignoring Synchrotron self-Compton effects (Y is the Inverse Compton parameter; e.g. Zhang et al. 2007), the cooling frequency would be already under the optical band at the start of GRB 091208B observations or over the X-ray band for GRB 141220A – inconsistent with the optical and X-ray data of both GRBs. Additionally, GRB 091208B high polarization was measured from a wide time-window and it is an average of the time-evolving polarization vector – favouring a more stable large-scale magnetic field component comparable to the reverse shock in GRB 090102 (Steele et al. 2009). This suggests that GRB 091208B afterglow could have contribution from a polarized reverse shock component (e.g. Section 4.2.3), be the emission of a long-lasting polarized reverse shock (Uhm & Beloborodov 2007) or less likely – because the light curve is consistent with ISM profile – be embedded in the magnetized wind from the progenitor (Granot & Königl 2003).

5 CONCLUSIONS

We used the 2-m fully autonomous LT to measure the early optical afterglow of *Swift* GRB141220A and its multicolour polarization. The polarization is already low $P_{\text{BV}} = 2.8^{+2.0}_{-1.6}$ per cent (2σ) at 129.5–204.3 s after the burst (i.e. 55.9–88.1 s in the GRB rest frame) in the RINGO3 BV band, and continues to be low $P_{\text{BV}} < 5$ per cent (2σ) to 727 s post-burst.

The temporal and spectral properties of the afterglow emission are consistent with a classical forward shock. We exclude alternative interpretations: a special viewing angle to an off-axis jet, which would require an anomalously low electron index, or an edge-on jet, for which achromatic breaks would be expected but are not observed.

Extinction through the Milky Way along the line-of-sight to GRB 141220A is low; the Galactic dust is expected to induce only $P \sim 0.2$ per cent polarization. We investigate different mechanisms that can contribute to the total polarization budget ($P_{\text{BV}} = 2.8^{+2.0}_{-1.6}$ per cent):

(i) The obscuration in the GRB host galaxy is high; we estimate that host galaxy dust could induce up to $P \lesssim 1.8$ per cent at most – consistent with our observations.

(ii) Reverse shocks are usually only detectable in the first minutes after the burst if bright – their afterglows have distinct temporal properties and fade more quickly than the slowly emerging forward shock radiation. Theoretical predictions that reverse shocks should be polarized ($P \sim 50$ per cent) have been confirmed observationally in mildly magnetized GRBs ($P = 10\text{--}30$ per cent). A reverse shock tail at the time of GRB 141220A polarization observations is very likely.

(iii) Forward shocks are theoretically predicted to be unpolarized or very lowly polarized with $P \leq 2$ per cent; late-time observations of forward shocks – when any primordial field has dissipated hours to a day after the burst – and GRB 141220A early-time observations confirm this.

Table 4. Rest-frame properties of GRBs with optical polarization measurements (P_{obs}) at early times ($t_{\text{obs, rest}}$) and afterglow optical decays (α_{opt}) interpreted as forward shock emission. Note that we only quote the first polarization measurement after the burst. We also state the rest-frame dust absorption for the host galaxy ($A_{\text{V, HG, rest}}$) and the MW ($A_{\text{V, MW}}$; Schlegel et al. 1998) and, the total hydrogen absorption for the rest-frame host galaxy ($N_{\text{H, HG, rest}}$) and the MW ($N_{\text{H, MW}}$; Willingale et al. 2013). We list the duration of the bursts in the 15–350 keV *Swift* BAT band ($T_{90, \text{rest}}$; Lien et al. 2016), the peak energy ($E_{\text{peak, rest}}$), the isotropic energy (E_{iso}), the deceleration time of the fireball ($t_{\text{dec, rest}}$) and the Lorentz factor at the deceleration time for ISM in thin shell regime (Γ_{dec} ; Sari & Piran 1999).

GRB	z	$t_{\text{obs, rest}}$ (s)	P_{obs} (per cent)	α_{opt}	$A_{\text{V, MW}}$	$A_{\text{V, HG, rest}}$	$N_{\text{H, MW}}$ (10^{21} cm^{-2})	$N_{\text{H, HG, rest}}$ (10^{21} cm^{-2})
060418	1.49	81.5–93.6	<8	1.28 ± 0.05	0.70	0.17 ± 0.02	1.59	$2.80^{+0.97}_{-0.85}$
091208B	1.063	72.2–342.2	10.4 ± 2.5	0.75 ± 0.02	0.16	$0.95^{+0.22}_{-0.20}$	0.58	$7.81^{+1.40}_{-1.21}$
100805A	1.85	49.2–112.3	<14	0.86 ± 0.04	0.58	–	2.64	$4^{+6}_{-4} a$
120327A	2.81	436.7–594.0	<4	1.22 ± 0.02	1.04	<0.03	2.66	10 ± 2
131030A	1.294	289.9–298.6	2.25 ± 1.65	0.78 ± 0.02	0.18	$0.01^{+0.03}_{-0.05}$	0.56	$4.6^{+1.5}_{-1.3}$
141220A	1.3195	55.9–88.1	$2.8^{+2.0}_{-1.6}$	1.091 ± 0.008	0.04	0.71 ± 0.15	0.13	3 ± 2
GRB		$T_{90, \text{rest}}$ (s)	$E_{\text{peak, rest}}$ (keV)		E_{iso} (10^{52} erg)		$t_{\text{dec, rest}}$ (s)	Γ_{dec}
060418		44 ± 19	535^{+77}_{-62}		14.7 ± 0.9		61 ± 4	198
091208B		7.2 ± 1.6	246 ± 15		1.97 ± 0.06		<37.8	>184
100805A		5.8 ± 0.7	148 ± 68^a		$1.2^{+0.8}_{-0.5} a$		<49.2	>157
120327A		16.7 ± 1.8	522^{+84}_{-61}		$19.1^{+1.9}_{-1.6}$		<69.8	>195
131030A		17.2 ± 1.6	450 ± 14		32.7 ± 1.3		<71.7	>206
141220A		3.1 ± 0.2	322^{+23}_{-21}		2.29 ± 0.12		<37.1	>189

References – GRB 060418 (Dupree et al. 2006; Mundell et al. 2007a; Schady et al. 2007; Molinari et al. 2007; Tsvetkova et al. 2020), GRB 091208B (Wiersema et al. 2009; Nakajima et al. 2009; Uehara et al. 2012; Schady et al. 2012; Nava et al. 2012), GRB 100805A (Oates et al. 2012; Steele et al. 2017), GRB 120327A (Perley & Tanvir 2012; D’Elia et al. 2014; Steele et al. 2017; Melandri et al. 2017; Tsvetkova et al. 2020), GRB 131030A (Breeveld & Troja 2013; de Ugarte Postigo et al. 2013; Xu et al. 2013; King et al. 2014; Littlejohns et al. 2015; Tsvetkova et al. 2017) and GRB 141220A (de Ugarte Postigo et al. 2014a; Tsvetkova et al. 2017; this work).

^aGRB 100805A $N_{\text{H, HG, rest}}$ is derived from *Swift* XRT 0.3–10 keV spectra and the E_{peak} and E_{iso} are computed from *Swift* BAT 15–350-keV spectra.

The nature of very early-time polarization properties of forward shocks, however, remains debated. Uehara et al. (2012) detected $P = (10.4 \pm 2.5)$ per cent in the first ~ 2.5 –11.8 min post-burst in GRB 091208B. They interpreted their result as the first detection of significant polarization from a forward shock. The high-energy properties (E_{iso} , E_{peak} , and fluence) of GRB 141220A and GRB 091208B are similar, so a direct comparison of their properties is valuable. The magnetic field length-scale of GRB 091208B forward shock implies $P \sim 20$ per cent polarization at the time of GRB 141220A observations, which does not agree with our observations. We find that it is not likely that GRB 091208B and GRB 141220A external mediums have extreme densities and we suggest larger coherent length-scales for the magnetic field of GRB 091208B, due to the presence of a large-scale magnetic field from the reverse shock.

Overall, this leads to the conclusion that forward shocks in GRBs are $P \leq 2$ per cent polarized in the first few minutes after the burst and coherence scales of the generated magnetic fields are consistent with theoretical predictions. Therefore, any significant polarization measurement at early times suggests reverse shock (or prompt) emission origin.

ACKNOWLEDGEMENTS

We thank the anonymous referee for the constructive comments that improved the paper. The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council. This work made use of data supplied by the UK *Swift* Science Data Centre at

the University of Leicester. The research leading to these results has received funding from the European Union’s Horizon 2020 Programme under the AHEAD project (grant agreement 654215). NJ and CGM acknowledge financial support from Mr Jim Sherwin and Mrs Hiroko Sherwin. CGM acknowledges support from the Science and Technology Facilities Council and the UK Research and Innovation (ST/N001265/1). AG acknowledges the financial support from the Slovenian Research Agency (grants P1-0031, I0-0033, J1-8136, J1-2460).

DATA AVAILABILITY

The data underlying this paper are available in the paper and in its online supplementary material.

REFERENCES

- Abbott B. P. et al., 2017a, *Phys. Rev. Lett.*, 119, 161101
- Abbott B. et al., 2017b, *ApJ*, 848, L12
- Alam S. et al., 2015, *ApJS*, 219, 12
- Amati L. et al., 2002, *A&A*, 390, 81
- Arnaud K., Dorman B., Gordon C., 1999, *Astrophysics Source Code Library*, record ascl:9910.005
- Arnold D., 2017, PhD thesis. Liverpool John Moores University
- Arnold D. M., Steele I. A., Bates S. D., Mottram C. J., Smith R. J., 2012, in McLean I. S., Ramsay S. K., Takami H., eds, *Ground-based and Airborne Instrumentation for Astronomy IV*. SPIE, Bellingham, p. 84462J
- Beniamini P., Nakar E., 2019, *MNRAS*, 482, 5430
- Beniamini P., Nava L., Duran R. B., Piran T., 2015, *MNRAS*, 454, 1073
- Berger E., 2014, *ARA&A*, 52, 43

- Blackburn J. K., 1995, in Shaw R. A., Payne H. E., Hayes J. J. E., eds, *Astronomical Data Analysis Software and Systems IV*, ASP Conference Series, Vol. 77, p. 367
- Bradley L. et al., 2016, *Astrophysics Source Code Library*, record ascl:1609.011
- Breeveld A. A., Troja E., 2013, *GRB Coord. Netw.*, 15414, 1
- Bromberg O., Nakar E., Piran T., Sari R., 2013, *ApJ*, 764, 179
- Buckley D. A. H. et al., 2020, preprint ([arXiv:2009.14081](https://arxiv.org/abs/2009.14081))
- Burgess J. M., Kole M., Berlato F., Greiner J., Vianello G., Produit N., Li Z. H., Sun J. C., 2019, *A&A*, 627, A105
- Chattopadhyay T. et al., 2019, *ApJ*, 884, 123
- Clarke D., Neumayer D., 2002, *A&A*, 383, 360
- Clayton G. C., Martin P. G., Thompson I., 1983, *ApJ*, 265, 194
- Clayton G. C., Green J., Wolff M. J., Zellner N. E. B., Code A. D., Davidsen A. F., WUPPE Science Team, HUT Science Team, 1996, *ApJ*, 460, 313
- Clayton G. C., Wolff M. J., Gordon K. D., Smith P. S., Nordsieck K. H., Babler B. L., 2004, *AJ*, 127, 3382
- Coburn W., Boggs S. E., 2003, *Nature*, 423, 415
- Covino S. et al., 1999, *A&A*, 348, L1
- Covino S. et al., 2013, *MNRAS*, 432, 1231
- Covino S., Ghisellini G., Lazzati D., Malesani D., 2004, in Feroci M., Frontera F., Masetti N., Piro L., eds, *Third Rome Workshop on Gamma-Ray Bursts in the Afterglow Era* ASP Conference Series, Vol. 312. Astronomical Society of the Pacific, San Francisco, p. 169
- Cummings J. R., D'Elia V., Gehrels N., Lien A. Y., Palmer D. M., Sbarufatti B., Siegel M. H., 2014, *GRB Coord. Netw.*, 17196, 1
- Curran P. A., Evans P. A., de Pasquale M., Page M. J., van der Horst A. J., 2010, *ApJ*, 716, L135
- D'Elia V. et al., 2014, *A&A*, 564, A38
- de Ugarte Postigo A., Thoene C. C., Sanchez-Ramirez R., Gorosabel J., Fynbo J. P. U., Tanvir N. R., Alvarez Iglesias C. A., 2013, *GRB Coord. Netw.*, 15408, 1
- de Ugarte Postigo A., Thoene C. C., Gorosabel J., Tanvir N., Fynbo J. P. U., Pesev P., Gomez Velarde G., Perez Valladares D., 2014a, *GRB Coord. Netw.*, 17198, 1
- de Ugarte Postigo A., Thoene C. C., Gorosabel J., Tanvir N. R., Fynbo J. P. U., Pesev P., Gomez Velarde G., Perez Valladares D., 2014b, *GRB Coord. Netw.*, 17200, 1
- Dupree A. K., Falco E., Prochaska J. X., Chen H. W., Bloom J. S., 2006, *GRB Coord. Netw.*, 4969, 1
- Evans P. A. et al., 2009, *MNRAS*, 397, 1177
- Fosalba P., Lazarian A., Prunet S., Tauber J. A., 2002, *ApJ*, 564, 762
- Gao H., Lei W.-H., Zou Y.-C., Wu X.-F., Zhang B., 2013, *New Astron. Rev.*, 57, 141
- Ghirlanda G. et al., 2018, *A&A*, 609, A112
- Ghisellini G., Lazzati D., 1999, *MNRAS*, 309, L7
- Gill R., Granot J., 2021, preprint ([arXiv:2101.06777](https://arxiv.org/abs/2101.06777))
- Gill R., Granot J., Kumar P., 2020, *MNRAS*, 491, 3343
- Golenetskii S. et al., 2014, *GRB Coord. Netw.*, 17207, 1
- Gompertz B. P. et al., 2014, *GRB Coord. Netw.*, 17203, 1
- Granot J., Königl A., 2003, *ApJ*, 594, L83
- Greiner J. et al., 2003, *Nature*, 426, 157
- Gruzinov A., Waxman E., 1999, *ApJ*, 511, 852
- Guidorzi C. et al., 2006, *Publ. Astron. Soc. Pac.*, 118, 288
- Hiltner W. A., 1949, *Science*, 109, 165
- Hjorth J. et al., 1999, *Science*, 283, 2073
- Høg E. et al., 2000, *A&A*, 355, L27
- Inoue T., Asano K., Ioka K., 2011, *ApJ*, 734, 77
- Jordana-Mitjans N. et al., 2020, *ApJ*, 892, 97
- King O. G. et al., 2014, *MNRAS*, 445, L114
- Klose S., Palazzi E., Masetti N., Stecklum B., Greiner J., Hartmann D. H., Schmid H. M., 2004, *A&A*, 420, 899
- Kobayashi S., 2000, *ApJ*, 545, 807
- Kole M. et al., 2020, *A&A*, 644, A124
- Komissarov S. S., Vlahakis N., Königl A., Barkov M. V., 2009, *MNRAS*, 394, 1182
- Kopač D. et al., 2015, *ApJ*, 806, 179
- Kouveliotou C., Meegan C. A., Fishman G. J., Bhat N. P., Briggs M. S., Koshut T. M., Paciesas W. S., Pendleton G. N., 1993, *ApJ*, 413, L101
- Laskar T. et al., 2013, *ApJ*, 776, 119
- Laskar T. et al., 2019, *ApJ*, 878, L26
- Lazarian A., 2007, *J. Quant. Spec. Radiat. Transf.*, 106, 225
- Lazzati D. et al., 2003, *A&A*, 410, 823
- Lazzati D., Rossi E., Covino S., Ghisellini G., Malesani D., 2002, *A&A*, 396, L5
- Lien A. et al., 2016, *ApJ*, 829, 7
- Littlejohns O. M. et al., 2015, *MNRAS*, 449, 2919
- Lyutikov M., Pariev V. I., Blandford R. D., 2003, *ApJ*, 597, 998
- Marshall F. E., Cummings J. R., 2014, *GRB Coord. Netw.*, 17201, 1
- Medvedev M. V., Loeb A., 1999, *ApJ*, 526, 697
- Medvedev M. V., Fiore M., Fonseca R. A., Silva L. O., Mori W. B., 2005, *ApJ*, 618, L75
- Melandri A. et al., 2017, *A&A*, 607, A29
- Mészáros P., 2002, *Annu. Rev. A&A*, 40, 137
- Molinari E. et al., 2007, *A&A*, 469, L13
- Mundell C. G. et al., 2007a, *Science*, 315, 1822
- Mundell C. G. et al., 2007b, *ApJ*, 660, 489
- Mundell C. G. et al., 2013, *Nature*, 504, 119
- Nakajima H., Yatsu Y., Mori Y. A., Endo A., Shimokawabe T., Kawai N., 2009, *GRB Coord. Netw.*, 10260, 1
- Nakar E., Oren Y., 2004, *ApJ*, 602, L97
- Nakar E., Piran T., 2003, *ApJ*, 598, 400
- Nava L. et al., 2012, *MNRAS*, 421, 1256
- Nishikawa K. I. et al., 2009, *ApJ*, 698, L10
- Nishikawa K. I., Hardee P., Richardson G., Preece R., Sol H., Fishman G. J., 2003, *ApJ*, 595, 555
- Nousek J. A. et al., 2006, *ApJ*, 642, 389
- Oates S. R., Page M. J., De Pasquale M., Schady P., Breeveld A. A., Holland S. T., Kuin N. P. M., Marshall F. E., 2012, *MNRAS*, 426, L86
- Panaiteescu A., Kumar P., 2002, *ApJ*, 571, 779
- Panopoulou G. V., Hensley B. S., Skolidis R., Blinov D., Tassis K., 2019, *A&A*, 624, L8
- Perley D. A., Tanvir N. R., 2012, *GRB Coord. Netw.*, 13133, 1
- Piran, 2004, *Rev. Mod. Phys.*, 76, 1143
- Piran T., 1999, *Phys. Rep.*, 314, 575
- Planck Collaboration et al., 2018a, *A&A*, 641, 67
- Planck Collaboration et al., 2018b, *A&A*, 641, 43
- Racusin J. L. et al., 2009, *ApJ*, 698, 43
- Rees M. J., Meszaros P., 1992, *MNRAS*, 258, 41
- Rees M. J., Meszaros P., 1994, *ApJ*, 430, L93
- Rodrigues C. V., Magalhães A. M., Coyne G. V., Piirola S. J. V., 1997, *ApJ*, 485, 618
- Rol E. et al., 2000, *ApJ*, 544, 707
- Rossi E. M., Lazzati D., Salmonson J. D., Ghisellini G., 2004, *MNRAS*, 354, 86
- Rutledge R. E., Fox D. B., 2004, *MNRAS*, 350, 1288
- Rybicki G. B., Lightman A. P., 1979, *Radiative processes in astrophysics*. Wiley, New York, NY
- Sari R., 1997, *ApJ*, 489, L37
- Sari R., 1999, *ApJ*, 524, L43
- Sari R., Piran T., 1999, *ApJ*, 520, 641
- Sari R., Piran T., Narayan R., 1998, *ApJ*, 497, L17
- Schady P. et al., 2007, *MNRAS*, 377, 273
- Schady P. et al., 2012, *A&A*, 537, A15
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
- Schmidt G. D., Elston R., Lupie O. L., 1992, *AJ*, 104, 1563
- Serkowski K., Mathewson D. S., Ford V. L., 1975, *ApJ*, 196, 261
- Shahmoradi A., Nemiroff R. J., 2015, *MNRAS*, 451, 126
- Shaw S. J., Tarengi M., 1976, *ApJ*, 204, L25
- Silva L. O., Fonseca R. A., Tonge J. W., Dawson J. M., Mori W. B., Medvedev M. V., 2003, *ApJ*, 596, L121
- Sironi L., Goodman J., 2007, *ApJ*, 671, 1858
- Stawikowska A., Krzeszowski K., Żejmo M., Reig P., Steele I., 2016, *MNRAS*, 458, 759
- Spitkovsky A., 2008, *ApJ*, 682, L5

- Stamatikos M. et al., 2014, GRB Coord. Netw., 17202, 1
- Steele I. A. et al., 2004, in Oschmann Jacobus M. J., ed., Proc. SPIE Conf. Ser. Vol. 5489, Ground-based Telescopes. SPIE, Bellingham, p. 679
- Steele I. A. et al., 2017, *ApJ*, 843, 143
- Steele I. A., Mundell C. G., Smith R. J., Kobayashi S., Guidorzi C., 2009, *Nature*, 462, 767
- Troja E. et al., 2017, *Nature*, 547, 425
- Tsvetkova A. et al., 2017, *ApJ*, 850, 161
- Tsvetkova A. et al., 2020, *AJ*, 908, 16
- Turnshek D. A., Bohlin R. C., Williamson R. L. I., Lupie O. L., Koornneef J., Morgan D. H., 1990, *AJ*, 99, 1243
- Uehara T. et al., 2012, *ApJ*, 752, L6
- Uhm Z. L., Beloborodov A. M., 2007, *ApJ*, 665, L93
- Wang X.-G. et al., 2015, *ApJS*, 219, 9
- Waxman E., 1997, *ApJ*, 485, L5
- Weibel E. S., 1959, *Phys. Rev. Lett.*, 2, 83
- Whittet D. C. B., Martin P. G., Hough J. H., Rouse M. F., Bailey J. A., Axon D. J., 1992, *ApJ*, 386, 562
- Wiersema K. et al., 2012, *MNRAS*, 426, 2
- Wiersema K., Tanvir N. R., Cucchiara A., Levan A. J., Fox D., 2009, GRB Coord. Netw., 10263, 1
- Wijers R. A. M. J. et al., 1999, *ApJ*, 523, L33
- Willingale R., Starling R. L. C., Beardmore A. P., Tanvir N. R., O'Brien P. T., 2013, *MNRAS*, 431, 394
- Willis D. R. et al., 2005, *A&A*, 439, 245
- Woosley S. E., 1993, *ApJ*, 405, 273
- Xu D., Fynbo J. P. U., Jakobsson P., Cano Z., Milvang-Jensen B., Malesani D., de Ugarte Postigo A., Hayes M., 2013, GRB Coord. Netw., 15407, 1
- Yamazaki R., Ioka K., Nakamura T., 2002, *ApJ*, 571, L31
- Yamazaki R., Yonetoku D., Nakamura T., 2003, *ApJ*, 594, L79
- Yamazaki R., Ioka K., Nakamura T., 2004, *ApJ*, 607, L103
- Yano Y. et al., 2014, GRB Coord. Netw., 17215, 1
- Yu H. F., 2014, GRB Coord. Netw., 17205, 1
- Zhang B. et al., 2007, *ApJ*, 655, 989
- Zhang B., Kobayashi S., 2005, *ApJ*, 628, 315

SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

Table1.txt

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the paper.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.