

**High intramuscular triglyceride turnover rates and the link to insulin sensitivity: Influence of obesity, type 2 diabetes and physical activity**

J.S. Barrett<sup>1</sup>, K.L. Whytock<sup>2</sup>, J.A. Strauss<sup>1</sup>, A.J.M. Wagenmakers<sup>1</sup>, S.O. Shepherd<sup>1</sup>

<sup>1</sup>Liverpool John Moores University, <sup>2</sup>Translational Research Institute, AdventHealth, Orlando, FL 32804, USA.

**Address for correspondence:**

Dr Sam Shepherd  
Research Institute for Sport & Exercise Sciences  
Liverpool John Moores University  
Tom Reilly Building  
Byrom Street  
Liverpool  
L3 3AF

## **Abstract**

Large intramuscular triglyceride (IMTG) stores in sedentary, obese individuals have been linked to insulin resistance, yet well-trained athletes exhibit high IMTG levels whilst maintaining insulin sensitivity. Contrary to previous assumptions, it is now known that IMTG content *per se* does not result in insulin resistance. Rather, insulin resistance is caused, at least in part, by the presence of high concentrations of harmful lipid metabolites, such as diacylglycerols and ceramides in muscle. Several mechanistic differences between obese sedentary individuals and their highly trained counterparts have been identified, that determine the differential capacity for IMTG synthesis and breakdown in these populations. In this review, we first describe the most up-to-date mechanisms by which a low IMTG turnover rate (both breakdown and synthesis) leads to the accumulation of lipid metabolites and results in skeletal muscle insulin resistance. We then explore current and potential exercise and nutritional strategies which target IMTG turnover in sedentary obese individuals, to improve insulin sensitivity. Overall, improving IMTG turnover should be an important component of successful interventions which aim to prevent the development of insulin resistance in the ever-expanding sedentary, overweight and obese populations.

## **Novelty Bullet points**

- A description of the most up-to-date mechanisms regulating turnover of the IMTG pool.
- An exploration of current and potential exercise/nutritional strategies to target and enhance IMTG turnover in obese individuals
- Overall, highlights the importance of improving IMTG turnover to prevent the development of insulin resistance

## **Key words**

Intramuscular triglyceride, perilipins, insulin resistance, exercise, skeletal muscle, type 2 diabetes

## Introduction

Physical inactivity combined with chronic over-consumption of an energy-dense diet causes expansion of adipose tissue depots around the body leading to obesity. The buffering capacity of adipose tissue can become impaired in obesity resulting in spill-over of circulating fatty acids (FA) and triglycerides into non-adipose tissues, such as the liver and skeletal muscle, leading to ectopic lipid deposition (Frayn, 2002). The delivery of excess lipid to skeletal muscle leads to accumulation of intramuscular triglyceride (IMTG) (Bachmann *et al.*, 2001; Chow *et al.*, 2014), which is characteristic of the obese and T2D states. Thus, high IMTG levels in sedentary obese individuals and T2D patients are associated with insulin resistance (Pan *et al.*, 1997; Kelley and Goodpaster, 2001; van Loon *et al.*, 2004). However, it is now known that a high IMTG content *per se* does not result in insulin resistance. Rather, insulin resistance is believed to be caused, at least in part, by the presence of harmful lipid metabolites, such as diacylglycerols (DAGs) and ceramides (Goodpaster *et al.*, 1997; Pan *et al.*, 1997; Forouhi *et al.*, 1999). Indeed, the accumulation of DAGs and ceramides has been shown to disrupt cell function, and specifically the capacity for insulin-stimulated glucose uptake into skeletal muscle via direct interference with the insulin signalling cascade (Yu *et al.*, 2002; Summers and Nelson, 2005; Chaurasia and Summers, 2015). However, defects in insulin signalling are not always observed in insulin resistant individuals (Meyer *et al.*, 2002; Hojlund *et al.*, 2003; Ramos *et al.*, 2021), and the precise mechanism by which lipid metabolites induce insulin resistance is far from certain. Consequently, the link between lipid metabolite accumulation and insulin resistance appears complex, and may be rooted in other factors such as lipid metabolite composition and subcellular localisation (extensively reviewed recently by (Bergman and Goodpaster, 2020). The association between IMTG accumulation and insulin resistance has also been disputed due to endurance trained athletes having a comparable or even higher IMTG content than obese individuals and T2D patients, whilst remaining highly insulin sensitive (Goodpaster *et al.*, 2001; van Loon *et al.*, 2003a). This phenomenon is now well known as the “athlete’s paradox” (Goodpaster *et al.*, 2001). The question of how endurance trained athletes exhibit similar IMTG content compared to obese individuals but are able to combine this with high levels of insulin sensitivity has been the subject of intense research in the last 20 years.

Although the mechanistic link between IMTG accumulation and insulin resistance is not yet fully established, the fundamental difference between endurance athletes when compared to obese individuals with or without T2D is their greater ability to utilise IMTG as a source of fuel during exercise (Schrauwen *et al.*, 2002; van Loon, 2004). Regular breakdown (lipolysis) of IMTG and oxidation of FA during exercise, coupled with elevated rates of FA uptake and IMTG (re)synthesis following exercise, creates a dynamic IMTG pool with a high turnover rate (van Loon, 2004; Moro, Bajpeyi and Smith, 2008). An attractive hypothesis is that regular IMTG turnover maintains insulin sensitivity by regulating the concentration and spatial distribution of lipid metabolites thereby ameliorating their impact on insulin signalling and cell function. However, exercise training appears to have little impact on insulin signalling (Christ-Roberts *et al.*, 2004; Frosig *et al.*, 2007) and does not always alter the concentration of lipid metabolites in muscle (Meyer *et al.*, 2002; Hojlund *et al.*, 2003). Rather, a greater ability to utilise IMTG in trained individuals leads to the hypothesis that the capacity to appropriately adjust FA storage and efficiently breakdown and oxidise FA in line with metabolic demand and FA availability is fundamental to improve insulin sensitivity. Because of this, focus has shifted to identifying the mechanisms that enable a high turnover rate of the IMTG pool in trained individuals in order to be able to create the optimal intervention in obese individuals and people with T2D and subsequently improve insulin sensitivity.

With this in mind, this review will first evaluate the differences in the storage and utilisation of IMTG between trained and more sedentary populations (i.e., obese and elderly individuals and those with T2D) to demonstrate that the dynamic nature of the IMTG pool in trained individuals is a crucial characteristic to the preservation of insulin sensitivity. Based on this information, we will then explore potential strategies to maximise IMTG turnover which could be implemented as interventions to improve insulin sensitivity in obese individuals and T2D patients.

### **Differences in IMTG storage between trained and sedentary individuals**

Although IMTG content itself has no mechanistic link to insulin resistance, in this context it is important to consider the fibre-specific distribution, subcellular location, and morphology of IMTG-containing lipid droplets (LD). In lean, healthy individuals there is a hierarchical distribution between the different fibre types with the majority of

IMTG being stored in type I fibres, followed by type IIa then type IIx fibres (Daemen, van Polanen and Hesselink, 2018; Whytock *et al.*, 2020). Within skeletal muscle, LDs are located either between the myofibrils (intermyofibrillar [IMF] LDs) or just beneath the surface membrane (subsarcolemmal [SS] LDs) (Nielsen *et al.*, 2017). Using transmission electron microscopy, it was recently shown that type I fibres of healthy males have small LD located in both the intermyofibrillar region and the subsarcolemmal region (Nielsen *et al.*, 2017), whereas type II fibres contain a similar number of LD in the intermyofibrillar region and the subsarcolemmal region but those in the subsarcolemmal region are ~20% larger in diameter (Nielsen *et al.*, 2017). Consequently, the size of subsarcolemmal LD in particular was associated with poorer insulin sensitivity, rather than LD number, at least in healthy untrained males (Nielsen *et al.*, 2017). Daemen *et al.*, (2018) extended these observations when comparing differences between trained individuals and patients with T2D. To this end, the elevated IMTG content in trained individuals was explained by a greater number of LD in the intermyofibrillar region of type I fibres, whereas individuals with T2D had a greater number of larger LD in the subsarcolemmal region of type II fibres (Daemen, van Polanen and Hesselink, 2018). It is important to note here that to date the majority of research investigating differences in LD location and morphology has been conducted in males, or without distinction between sex. Thus, differences in LD location and morphology over the lifespan and between sexes should be explored in future studies. Older adults have been shown to have larger LD, fewer mitochondria, and a lower proportion of LD in contact with mitochondria (Crane *et al.*, 2010), likely contributing to age-related decline in mitochondrial function and lipid metabolism.

Interestingly, 8 weeks of a high-calorie, high-fat diet induced insulin resistance in sedentary individuals and resulted in an increase in LD size rather than any changes in LD number (Covington *et al.*, 2017). More recently, we reported that a short-term (7-day) high-calorie, high-fat diet increased LD size and number in type I fibres in both the central and peripheral regions (Whytock *et al.*, 2020). In type IIa fibres LD size increased in both the SS and IMF region but only LD number increased in the SS region, whereas in type IIx fibres only SS-located LD increased in size with no changes in LD number observed (Whytock *et al.*, 2020). This suggests that changes in LD number and size may occur in a hierarchical manner based on both fibre type and subcellular region, at least in response to a high-calorie, high-fat diet. In contrast, a

combined weight loss and exercise training intervention in previously overweight or obese individuals resulted in a decrease in LD size concomitant with improved insulin sensitivity, even in the absence of a reduction in IMTG content (He, Goodpaster and Kelley, 2004). A large number of small LD located in the IMF region, as observed in healthy lean and trained individuals, creates a larger surface area to volume ratio, which is thought to be beneficial for the binding of proteins and lipolytic enzymes to the LD in order to liberate and release FA from the IMTG stored within. Moreover, LD are located in close proximity to mitochondria within skeletal muscle in healthy, trained individuals (Hoppeler *et al.*, 1999; Shaw, Jones and Wagenmakers, 2008), and exercise training in healthy or obese individuals increases the proportion of LD that are in contact with mitochondria (Tarnopolsky *et al.*, 2007; Shepherd *et al.*, 2017b). Together with a large number of small LDs, this adaptation likely creates an efficient means by which to channel FA liberated from IMTG within LD to the mitochondria for subsequent oxidation (Fig. 1). It is important to note here though that increased LD association with mitochondria does not necessarily mean the LD are utilised for oxidation, and this LD-mitochondria interaction may also support triacylglycerol synthesis and LD growth (Benador *et al.*, 2018; Benador *et al.*, 2019).

### **IMTG turnover in trained vs sedentary individuals.**

Cross-sectional comparisons between trained and untrained individuals confirm that endurance-trained individuals have a greater capacity to use IMTG as a substrate during exercise (Klein, Coyle and Wolfe, 1994; Coggan *et al.*, 2000). During moderate-intensity exercise in healthy individuals, IMTG-derived fatty acids contribute ~50% to total fat oxidation, with the remaining ~50% attributable to plasma FA (van Loon *et al.*, 2001). Serial muscle biopsies combined with microscopy-based analyses enable net changes in IMTG content to be determined and using this approach it is now known that IMTG utilisation preferentially occurs in type I fibres from IMTG-containing LD (van Loon *et al.*, 2003a; Shepherd *et al.*, 2013) that are located in the IMF region (Koh *et al.*, 2017; Jevons *et al.*, 2020). Moreover, in healthy individuals IMTG utilisation and FA oxidation during exercise is closely related to pre-exercise IMTG content (Shepherd *et al.*, 2013) whereby those with greatest IMTG stores have the greatest IMTG utilisation. Therefore, the high rate of IMTG utilisation observed in healthy, trained individuals must be matched by a large capacity for esterification and storage

of FA as IMTG following exercise. This has been illustrated in a recent study, which demonstrated that myotubes from athletic subjects have higher lipid turnover and lipid oxidation compared to those from sedentary individuals (Lund *et al.*, 2018). Specifically, myotubes from athletes exhibit higher rates of lipolysis and re-esterification of FA into the triacylglycerol (TAG) pool, indicating greater turnover of TAG stores. Importantly, higher complete oxidation and incomplete  $\beta$ -oxidation of FA in myotubes from the athletic population was also observed, suggesting they are able to more effectively rely on FA as a fuel source (Lund *et al.*, 2018). A greater accumulation of FA in myotubes derived from sedentary compared to athletic individuals led the authors to question whether the capacity for IMTG synthesis is downregulated in these individuals, and/or the capacity for lipid metabolite generation is upregulated (Lund *et al.*, 2018). The latter, of course, would consequently reduce insulin sensitivity.

Measuring the fractional synthesis rates (FSR) of IMTG in healthy individuals provides *in vivo* information on the rate of turnover of the IMTG pool. In this regard, IMTG FSR at rest in healthy individuals was first reported to be as high as  $\sim 3.4\%/h$ , suggesting that in this cohort complete turnover of the IMTG pool would occur in  $\sim 29$  h (Sacchetti *et al.*, 2004). Although Bergman *et al.* (2018) have since reported a lower resting IMTG FSR in trained individuals ( $\sim 1.56\%/h$ ), this was still more than 2-fold higher when compared to sedentary, lean individuals ( $\sim 0.61\%/h$ ) (Bergman *et al.*, 2018). Obese individuals have a lower IMTG FSR ( $\sim 0.42\%/h$ ) than the rates reported for lean, sedentary individuals, and the resting IMTG FSR for obese individuals with pre-diabetes is even lower ( $\sim 0.21\%/h$ ) (Perreault *et al.*, 2010). With these data, it is no surprise that Bergman *et al.* (2018) reported a positive correlation between IMTG FSR and insulin sensitivity at rest, along with a negative correlation between IMTG synthesis rates and the concentration of key lipid metabolites associated with insulin resistance. There is evidently a strong link between an individual's ability to breakdown and resynthesise IMTG and their level of insulin sensitivity.

As well as measuring IMTG FSR at rest, studies examining IMTG FSR during exercise alongside net changes in IMTG concentration provide further insight into the dynamics of the IMTG pool in trained and sedentary obese and T2D individuals. During 1 hour of moderate-intensity exercise, IMTG FSR is elevated compared to rest and while a

net reduction in IMTG content is observed in trained individuals, IMTG content remains unchanged in obese individuals and those with T2D (Bergman *et al.*, 2018). The latter finding is in line with previous studies measuring IMTG content in biopsies pre- and post-exercise which concluded that there is no net utilisation of IMTG in obese individuals and those with T2D (Kelley and Simoneau, 1994; Blaak and Wagenmakers, 2002). IMTG FSR is elevated during exercise in obesity and T2D (Bergman *et al.*, 2018), which could be due to the high circulating FFA concentrations often observed in these individuals (Axelsen *et al.*, 1999) supplying fatty acids for the synthesis of IMTG. In obese individuals and T2D patients though there is no net change in IMTG content during exercise, but for this to be true IMTG utilisation during exercise must be matched to IMTG FSR. Therefore, by combining pre- and post-exercise measures of IMTG content with estimates of IMTG FSR during exercise, it appears possible that obese and T2D individuals may utilise their IMTG stores, but this occurs in the absence of a net reduction in IMTG content (Bergman *et al.*, 2018), potentially due to replenishment of the IMTG stores from high circulating FFA concentrations (Axelsen *et al.*, 1999). It has also been reported that in individuals with normal glucose tolerance, IMTG FSR during exercise may be reduced, especially compared to individuals with prediabetes (Perreault *et al.*, 2010), obese individuals, and T2D patients (Bergman *et al.*, 2018) who all exhibit only very small changes in IMTG FSR during exercise. Thus, while there is not yet a consensus on how IMTG FSR is altered during exercise in trained, glucose tolerant individuals, these cross-sectional comparisons do highlight an inability to adjust IMTG FSR relative to metabolic demand in obese individuals with pre-diabetes and T2D. Importantly, a net reduction in IMTG content during exercise in endurance trained individuals will theoretically enable a greater capacity for uptake of plasma FFA and storage as IMTG in the post-exercise period. Without a net reduction in IMTG content during exercise in obese individuals, this may limit the capacity for FA's entering skeletal muscle following exercise to be stored as IMTG, and rather these FA's may instead be directed towards the generation of lipid metabolites.

## **Molecular mechanisms regulating IMTG turnover rate**

While the aim of this review is not to provide an in-depth account of the molecular mechanisms that regulate FA uptake and esterification, IMTG storage and breakdown,



it is pertinent that an up-to-date overview of these regulatory mechanisms is provided. For the former, the reader is directed to two excellent reviews (Badin, Langin and Moro, 2013; Lundsgaard, Fritzen and Kiens, 2018).

### *IMTG Synthesis*

Exogenous FA, derived from either adipose tissue or from the diet, are transported in the circulation and taken up into skeletal muscle to be stored as IMTG and/or oxidised as a fuel source. FA uptake into skeletal muscle is regulated primarily by FAT/CD36, although it is likely that this process is mediated by a series of transporter proteins reviewed in detail in (Schwenk *et al.*, 2010; Glatz and Luiken, 2018). Once in skeletal muscle, FA are converted to fatty acyl-CoA and directed to IMTG synthesis. Briefly, FA-CoA undergoes acylation catalysed by key enzymes glycerol-3-phosphate acyltransferase (GPAT), monoacylglycerol acyltransferase (MGAT) and diacylglycerol acyltransferase (DGAT) (Teodoro *et al.*, 2017), ultimately leading to the generation of TAG (or IMTG). The synthesised IMTG are then stored within LD and are a readily available fuel for healthy individuals.

Although the key enzymes that control IMTG synthesis have been identified, little is known about how they regulate this process in skeletal muscle. Following moderate-intensity exercise and a subsequent elevation of lipid availability (induced by an overnight lipid infusion), there is an increase in the protein expression of DGAT1 and GPAT1 (Schenk and Horowitz, 2007) and increased GPAT1 activity (Newsom *et al.*, 2011). Furthermore, overexpression of DGAT1 in rodents results in an increase in TAG content and a decrease in DAG (Liu *et al.*, 2007). However, GPAT1 and DGAT1 do not differ in expression between obese and lean individuals (Thrush *et al.*, 2009; Li *et al.*, 2011), and no differences are observed in DGAT1 mRNA expression between endurance trained, obese individuals and those with type 2 diabetes (Bergman *et al.*, 2018), nor in DGAT protein content (Amati *et al.*, 2011). Overall, there is currently no evidence to suggest that the expression or activity of DGAT or GPAT is impaired in obese and T2D individuals. As a result, it could be speculated that the machinery for IMTG synthesis is sufficient in all individuals, and it is the (as yet unknown) activation mechanism which is impaired in obesity and T2D.

### *IMTG lipolysis*

The reduced IMTG utilisation reported in obese individuals and those with T2D could be, at least partly, attributed to impaired rates of lipolysis. Indeed, when compared to lean individuals, obese individuals show impaired  $\beta_2$ -adrenergic-mediated stimulation of lipolysis in skeletal muscle (Blaak *et al.*, 2004). In skeletal muscle, the majority (~98%) of total TAG hydrolase activity (at least at rest) is regulated by adipose triglyceride lipase (ATGL) and hormone sensitive lipase (HSL). In this regard, it is important to note HSL protein content (Jocken *et al.*, 2007) and HSL phosphorylation (at Ser<sup>563</sup>, Ser<sup>555</sup> and Ser<sup>659</sup>) (Jocken *et al.*, 2008) are both lower in obese compared to lean individuals. However, it was reported that individuals with T2D actually had greater ATGL protein expression compared to lean and obese individuals (Badin *et al.*, 2011). Because ATGL may have a higher affinity to TAG (Haemmerle *et al.*, 2006), and HSL a higher affinity to DAG (Fredrikson *et al.*, 1981; Haemmerle *et al.*, 2002), it has been suggested that in obesity and T2D the imbalance between ATGL and HSL protein content favours DAG accumulation, and this contributes to the disruption of insulin signalling. Indeed, overexpression of ATGL in myotubes from lean, healthy, insulin-sensitive individuals induced DAG and ceramide accumulation, which was associated with reduced insulin-stimulated glycogen synthesis and reduced activation of IRS-1 and Akt (Badin *et al.*, 2011). Although, this imbalance was reported by Jocken *et al.*, (2008) with greater ATGL content and lower HSL content in obese individuals with T2D compared to lean (Jocken *et al.*, 2008), it was not evident in non-obese T2D, questioning its role in the development of insulin resistance. Moreover, it is now known that DAG that is derived from ATGL-mediated lipolysis is unable to activate the atypical PKC isoforms known to disrupt the insulin signalling cascade (Eichmann *et al.*, 2012). Additionally, Bergman *et al.*, (2018) has more recently shown that elevated IMTG content in obese individuals is not due to an imbalance between HSL and ATGL content, but more likely due to the specific species of ceramides present in obese individuals, and the subcellular location in which they are stored (Bergman *et al.*, 2018).

### *LD proteins and their regulation of IMTG turnover*

Surrounding a core of TAG and cholesterol esters, LDs have a phospholipid monolayer that is now known to be coated with numerous proteins which likely determines the functional role of each LD. The perilipin proteins (PLIN) are the group

of LD proteins most extensively investigated (Morales, Bucarey and Espinosa, 2017), with PLIN2, 3, 4 and 5 all being expressed in human skeletal muscle. Research conducted over the last decade has started to uncover a potential role for the PLIN proteins in both IMTG storage and lipolysis.

The role of PLIN in IMTG storage - It is evident from *in vitro* studies that the knockout of PLIN2 or PLIN5 in skeletal muscle compromises TAG storage (Bosma *et al.*, 2012; Gallardo-Montejano *et al.*, 2016). It makes sense then, that the overexpression of these PLIN isoforms results in quite the opposite, promoting TAG storage (Xu *et al.*, 2005; Bosma *et al.*, 2012; Gallardo-Montejano *et al.*, 2016). Similarly, the suppression of PLIN3 reduced LD maturation and TAG incorporation into IMTG stores in HeLa cells (Bulankina *et al.*, 2009), whereas in skeletal muscle myotubes augmenting PLIN3 gene expression increases IMTG content (Kleinert *et al.*, 2016). PLIN4 is purported to be the most abundant PLIN in skeletal muscle (Deshmukh *et al.*, 2015), yet its knockout in mice has no effect on skeletal muscle IMTG concentrations (Chen *et al.*, 2013), questioning the importance of this protein in IMTG storage.

Human biopsy studies demonstrate that PLIN2 protein expression is greater in trained versus sedentary individuals (Amati *et al.*, 2011; Shaw *et al.*, 2012; Shepherd *et al.*, 2013), females versus males (Shaw *et al.*, 2009; Peters *et al.*, 2012), and type 1 versus type 2 fibres (Shaw *et al.*, 2009), suggesting that PLIN2 is closely related to IMTG content in healthy individuals. The same observations also extend to PLIN3 (Peters *et al.*, 2012; Shepherd *et al.*, 2017b) and PLIN5 (Shepherd *et al.*, 2013; Shepherd *et al.*, 2017a; Shepherd *et al.*, 2017b; Daemen, van Polanen and Hesselink, 2018). Furthermore, when exercise training augments IMTG content, increases in PLIN2 (Shaw *et al.*, 2012; Shepherd *et al.*, 2013), PLIN3 (Shepherd *et al.*, 2017b) and PLIN5 (Peters *et al.*, 2012; Shepherd *et al.*, 2013) are also observed. Much less research has been conducted on PLIN4, although we recently showed that PLIN4 protein expression is greater in trained versus untrained individuals (Shepherd *et al.*, 2017b). Despite this, endurance training fails to augment PLIN4 mRNA or protein expression in healthy individuals (Peters *et al.*, 2012; Pourteymour *et al.*, 2015). Together, these data suggest that the expression of PLIN2, PLIN3 and PLIN5 is closely related to IMTG content, at least in healthy individuals or following a period of exercise training. This may be an important adaptation in order to support greater IMTG storage, especially

in the face of elevated FA availability and turnover. Indeed, in response to an acute lipid infusion (Shepherd *et al.*, 2017a) or 48 h of fasting (Gemmink *et al.*, 2016) (both conditions increase FA availability) in trained individuals, there appears to be a redistribution of the pre-existing PLIN2, PLIN3 and PLIN5 protein pool (which could be from either LD-bound or non-LD-bound) to the expanded LD pool. Importantly, this redistribution was not apparent in sedentary individuals (Shepherd *et al.*, 2017a) and the capacity to redistribute PLIN5 to maintain coverage of the expanded LD pool was associated with a greater maintenance of insulin sensitivity (Gemmink *et al.*, 2016). More recently, we showed that in elite triathletes, post-exercise increases in IMTG content occurred *prior* to a redistribution of the PLIN (2,3,5) protein pool (Jevons *et al.*, 2020). Taken together, this suggests that the PLIN proteins do not play a direct role in IMTG synthesis but coating of LD with PLINs may be an important adaptation which supports IMTG storage.

*The role of PLIN in IMTG breakdown* - Research has also focused on the potential role of PLIN2, PLIN3 and PLIN5 in supporting IMTG breakdown and utilisation during exercise. A role for the PLIN proteins in TAG breakdown stems from evidence showing that PLIN2, PLIN3 and PLIN5 can interact with the key lipolytic enzymes ATGL and HSL (Anthonen *et al.*, 1998; Granneman *et al.*, 2011; Macpherson *et al.*, 2013). Moreover, both PLIN2 and PLIN5 are thought to suppress lipolysis at rest by preventing the interaction between ATGL and CGI-58, whereas this inhibition is relieved permitting ATGL to interact with CGI-58 upon lipolytic stimulation (Wang *et al.*, 2011; Macpherson *et al.*, 2013). PLIN3 knockout in myotubes results in a reduction of lipid oxidation (Covington *et al.*, 2014). However, in response to an endurance exercise bout PLIN3 expression is positively correlated to whole-muscle homogenate palmitate oxidation rates as well as whole-body cumulative fat oxidation (Covington *et al.*, 2014). Recently, AMPK phosphorylation of PLIN3 was shown to bring about conformational changes to PLIN3 that expose the C-terminus and promote LD dispersion to facilitate lipolysis (Zhu *et al.*, 2019). This new data potentially underpins the relationship between PLIN3 and lipolysis.

In human studies, we initially showed that LD labelled with PLIN2 or PLIN5 are preferentially broken down in lean, sedentary individuals during 1 h of moderate-intensity exercise (Shepherd *et al.*, 2012; Shepherd *et al.*, 2013). However, when

assessed following six weeks of endurance training or sprint interval training, only LD labelled with PLIN5 were preferentially targeted for breakdown in an equivalent bout of exercise (Shepherd *et al.*, 2013). More recently, we have shown that during more prolonged (4 h) of moderate-intensity exercise in elite triathletes, this preferential use of PLIN labelled LD is not apparent. The use of LD not labelled with PLIN proteins could be attributed to a very high rate of IMTG turnover compared to sedentary individuals, especially during such prolonged exercise. Nevertheless, it does appear that PLIN5 plays a key functional role regulating IMTG breakdown, since 1 h of moderate-intensity exercise led to a redistribution of HSL specifically to LD labelled with PLIN5 (Whytock *et al.*, 2018).

#### *FA as signalling molecules*

It is now beginning to be understood that FA play a crucial role in skeletal muscle adaptation to exercise, by acting as ligands for peroxisome proliferator-activated receptor (PPAR)  $\alpha$  and  $\delta$  to support transcription of genes involved in lipid metabolism (Banner *et al.*, 1993). An extensive review of this topic is beyond the scope of the current paper (readers are directed to (Funai and Semenkovich, 2011) but it is pertinent to briefly consider the importance of FA as signalling molecules in the context of adaptation. In humans, suppression of lipolysis with nicotinic acid prior to exercise resulted in reduced mRNA expression of Peroxisome proliferator-activated receptor (PPAR) gamma coactivator 1-alpha (PGC-1 $\alpha$ ), PPAR $\alpha$  and PPAR $\delta$ , demonstrating a role for FA availability in exercise-induced gene expression (Watt *et al.*, 2004). In contrast, treatment with a PPAR $\delta$  agonist resulted in a dose-dependent increase in skeletal muscle FA oxidation in mice, as well as increased expression of mRNA encoded for proteins involved in FA catabolism, such as  $\beta$ -oxidation enzymes, FA transport proteins and uncoupling proteins (Tanaka *et al.*, 2003).

The source of the FA that act as ligands and activate PPAR $\alpha$  and PPAR $\delta$  is not yet confirmed but could be linked to the activation of intramuscular lipases acting upon IMTG. In support, ATGL-mediated hydrolysis of triacylglycerol promotes activation of PGC-1 $\alpha$  and PPAR $\alpha$  signalling in order to upregulate mitochondrial biogenesis. Moreover, ATGL-mediated lipolysis activates SIRT-1, the protein responsible for the deacetylation, and therefore activation, of PGC-1 $\alpha$  (Khan *et al.*, 2015). Until recently, the mechanism by which ATGL-mediated lipolysis promotes mitochondrial biogenesis

was unknown. However, Najt *et al.* (2020) recently reported that monounsaturated FA are able to activate SIRT-1, thereby enhancing PGC-1 $\alpha$ /PPAR $\alpha$  signalling. Importantly, these monounsaturated FA were derived from intracellular LDs (Najt *et al.*, 2020), permitting speculation that FA liberated from IMTG stored in LD may play a role in promoting mitochondrial biogenesis, although evidence for this is not yet available. Najt *et al.* (2020) also identified a role for PLIN5 as a FA binding protein, which is able to bind LD-derived monounsaturated FA and transport them to the nucleus, at least in response to cAMP/PKA-mediated lipolytic stimulation in hepatocytes. This observation is consistent with that of Gallardo-Monejano *et al.* (2016), who reported that fasting-induced lipolysis stimulates PKA-mediated phosphorylation of PLIN5 followed by its translocation to the nucleus (Gallardo-Montejano *et al.*, 2016). Here, PLIN5 interacts with SIRT-1 and PGC-1 $\alpha$  to increase transcription of proteins involved in FA catabolism. The translocation of PLIN5 was also shown for the first time to influence the transcriptional regulation of mitochondrial respiration and mitochondrial biogenesis (Gallardo-Montejano *et al.*, 2016). Taken together, it appears that intracellular LDs are more than a source of FA for oxidation but may play a key role in the regulation of mitochondrial biogenesis and the FA catabolism programme. Theoretically then, enhancing the utilisation and turnover of the IMTG pool could be one strategy to stimulate mitochondrial biogenesis and increase the capacity for fat oxidation. We will now explore potential strategies that could be used to induce these adaptations and subsequently enhance insulin sensitivity and discuss whether these have application in clinical populations.

## **Strategies to improve IMTG turnover**

### *Exercise training*

It is well known that endurance exercise training is a powerful stimulus to augment oxidative capacity and IMTG utilisation during exercise (Baldwin *et al.*, 1972; Kiens *et al.*, 1993; Phillips *et al.*, 1996; Bergman *et al.*, 1999; van Loon, 2004). Following endurance training there is an increase in the number of IMTG-containing LDs that are in direct contact with mitochondria (Tarnopolsky *et al.*, 2007; Devries *et al.*, 2013; Shepherd *et al.*, 2017a), which together with expansion of the mitochondrial network enhances the total capacity for FA  $\beta$ -oxidation (Granata, Jamnick and Bishop, 2018). In sedentary lean and obese individuals endurance training also augments the

expression of proteins that regulate IMTG breakdown and LD dynamics, including ATGL (Alsted *et al.*, 2009), PLIN2 (Shaw *et al.*, 2012; Shepherd *et al.*, 2013), PLIN3 (Shepherd *et al.*, 2017a) and PLIN5 (Shepherd *et al.*, 2017a). Notably, the increased expression of the PLIN proteins occurs primarily in type I fibres, which may explain why following a period of endurance training the increase in IMTG utilisation is also predominantly in type I fibres (Van Proeyen *et al.*, 2011a; Van Proeyen *et al.*, 2011b; Shepherd *et al.*, 2013). Importantly, the augmented use of IMTG following endurance training is related to an improvement in insulin sensitivity (Van Proeyen *et al.*, 2011a; Van Proeyen *et al.*, 2011b; Shepherd *et al.*, 2013).

Other forms of exercise training can also increase the capacity for IMTG utilisation during exercise. For example, 6 weeks of sprint interval training in sedentary individuals enhances IMTG utilisation during a single bout of moderate-intensity exercise, to a similar degree as endurance training (Shepherd *et al.*, 2013; Scribbans *et al.*, 2014) at least in lean, sedentary individuals. Underpinning this, several different forms of high intensity (or sprint) interval training have been shown to enhance skeletal muscle oxidative capacity (Little *et al.*, 2010; MacInnis and Gibala, 2017; Astorino and Schubert, 2018), and we have also reported an increased expression of PLIN2 and PLIN5 following sprint interval training (Shepherd *et al.*, 2013). Interestingly, in lean, sedentary individuals 6 weeks of whole-body resistance training also increases IMTG utilisation during a single bout of moderate-intensity exercise (Shepherd *et al.*, 2014), although the net changes in IMTG content during exercise following training are typically less than when compared to endurance or sprint interval training (Shepherd *et al.*, 2013). Given that resistance training can enhance mitochondrial content and oxidative capacity (Tang, Hartman and Phillips, 2006; Balakrishnan *et al.*, 2010; Pesta *et al.*, 2011) as well as resting IMTG content (Shepherd *et al.*, 2014), this finding is perhaps not unexpected. Moreover, because both high intensity interval training and resistance training improve insulin sensitivity in sedentary (Ishii *et al.*, 1998) and obese individuals (Croymans *et al.*, 2013; Ryan *et al.*, 2020), it is tempting to speculate that an enhanced capacity for IMTG utilisation during exercise could, at least in part, contribute to this effect. Although this is yet to be investigated directly, high intensity interval training in obese individuals augments several adaptations that would support greater IMTG turnover, including increased mitochondrial content (Gibala *et al.*, 2006; Burgomaster *et al.*, 2008; Larsen *et al.*, 2015; Chrois *et al.*, 2020) and mitochondrial

interaction with LDs, increased HSL and CD36 content (Talanian *et al.*, 2010) and greater protein expression of PLIN2, PLIN3, and PLIN5 (Shepherd *et al.*, 2013; Shepherd *et al.*, 2017b).

As discussed above, exercise stimulates IMTG utilisation in lean, healthy individuals (Shepherd *et al.*, 2013; Scribbans *et al.*, 2014), and to a lesser extent in those with obesity and T2D (Shepherd *et al.*, 2017a; Bergman *et al.*, 2018). However, a poor capacity to simultaneously reduce the rate of IMTG synthesis during exercise in individuals with obesity and T2D results in a minimal to zero decrease in IMTG content post-exercise (van Loon *et al.*, 2004). The result of this is a limited capacity for FA's entering skeletal muscle following exercise to be stored as IMTG and interfere with insulin signalling. Thus, in sedentary and obese individuals, additional strategies to exercise alone may be required to augment IMTG utilisation (or turnover) in order to create a net decrease in IMTG content post-exercise. In this context, the question then arises as to whether the insulin sensitising-effect of regular exercise training can be enhanced, by manipulating the conditions under which exercise is undertaken in order to maximise IMTG utilisation and obtain a post-exercise decrease in IMTG content (Fig. 2).

#### *Anti-lipolytic drug therapy*

During prolonged exercise there is a progressive decline in IMTG oxidation rate which is inversely related to the concomitant increase in plasma free fatty acid (FFA) concentrations (Romijn *et al.*, 1993; Romijn *et al.*, 1995; van Loon *et al.*, 2003a). Thus, it is purported that elevated plasma FFA concentrations may suppress IMTG utilisation during exercise. Pharmacological inhibition of adipose tissue lipolysis, via the anti-lipolytic agent Acipimox, before and during exercise abolishes the progressive rise in plasma FFA during exercise and results in enhanced IMTG oxidation in lean, healthy individuals (van Loon *et al.*, 2005a). Individuals with obesity and T2D exhibit elevated circulating plasma FFA and triglyceride concentrations, which is linked to the development of insulin resistance (Boden, 2003), and therefore could also be part of the mechanism by which exercise-induced IMTG utilisation is suppressed in these individuals. In support, inhibition of adipose tissue lipolysis before and during exercise in obese T2D patients increases IMTG oxidation at rest, during 60 minutes of moderate-intensity exercise, and for several hours' post-exercise (van Loon *et al.*,



2005b). Moreover, this was accompanied by a superior rate of glycogen oxidation, and greater post-exercise insulin sensitivity (van Loon et al., 2005b).

Several studies show that short-term Acipimox treatment (250 mg, two-to-three times per day) for up to 2 weeks can reduce fasting plasma free fatty acids and increase insulin sensitivity and glucose control in obese and T2D individuals (Bajaj *et al.*, 2005; Daniele *et al.*, 2014; Phielix *et al.*, 2014; van de Weijer *et al.*, 2015). Furthermore, 8 weeks of Acipimox treatment lowers plasma free fatty acids, cholesterol and triglyceride concentrations in obese individuals and T2D patients (Crepaldi *et al.*, 1988; Stuyt, Kleinjans and Stalenhoef, 1998). Lower plasma free fatty acid levels reduce the availability of FA for uptake into skeletal muscle, potentially minimising the accumulation of lipid in this tissue. Despite these positive changes to blood lipids and insulin sensitivity, longer term Acipimox treatment results in a rebound rise in fasting plasma FFA (Fulcher *et al.*, 1992; Vaag and Beck-Nielsen, 1992; Saloranta *et al.*, 1993) and both hepatic and skeletal muscle insulin sensitivity is unchanged (Makimura *et al.*, 2016). Therefore, while chronic treatment with Acipimox does not seem feasible, combining exercise with anti-lipolytic therapy may represent an effective strategy to augment insulin sensitivity in individuals with obesity and T2D.

#### *Fasted exercise*

As high plasma FFA concentrations suppress IMTG utilisation during exercise, a simple strategy to reduce FFA availability would be to feed carbohydrate, since carbohydrate ingestion increases plasma insulin levels and subsequently suppresses circulating FFA through insulin-mediated inhibition of HSL in adipose tissue (Watt *et al.*, 2004). However, insulin also suppresses HSL activity in skeletal muscle (Enoksson *et al.*, 1998), and therefore carbohydrate ingestion would theoretically lead to a decrease in IMTG utilisation. The overnight fasted state is characterised by low plasma insulin concentrations, and therefore skeletal muscle lipolytic enzyme activity remains functional (Horowitz et al., 1997; Arkinstall et al., 2001). Indeed, in healthy individuals, two hours of moderate-intensity cycling in the overnight-fasted compared to carbohydrate-fed state led to greater IMTG utilisation in type 1 fibres (De Bock *et al.*, 2005). Similar results were obtained in overweight/obese males, although in this population IMTG utilisation was more pronounced in both type 1 and type 2 fibres following fasted compared to postprandial exercise (Edinburgh *et al.*, 2020). It should

be noted though that when the exercise duration is extended to 3 h, there appears to be no effect of carbohydrate feeding before (Fell et al., 2021) and/or during (Stellingwerff et al., 2007; Fell et al., 2021) exercise on IMTG utilisation. In this case, it is possible that the duration of exercise overrides the inhibitory effect of carbohydrate feeding on IMTG utilisation. Nevertheless, it is worth noting that limiting glucose availability by exercising in the fasted state encourages skeletal muscle cells to increase transcriptional activities of factors that upregulate the fatty acid oxidation programme and thereby induce metabolic adaptations for efficient lipid oxidation (Canto et al., 2010). Six weeks of endurance training in lean, healthy individuals in a fasted state was more effective for increasing skeletal muscle oxidative capacity, CD36 and FATBP<sub>m</sub> content, and net IMTG breakdown during a single exercise bout compared to undertaking the same training in a fed condition (Van Proeyen et al., 2011b). Moreover, 6 weeks of endurance training in a fasted state in overweight/obese individuals augmented skeletal muscle remodelling of phospholipids (Edinburgh et al., 2020). Thus, fasted exercise appears to augment changes in skeletal muscle phospholipids, by reducing saturated FFA, that correlate with improved post-prandial insulinemia.

Whether these beneficial adaptations to fasted exercise can enhance the insulin-sensitising effect of exercise training for obese individuals and individuals with T2D has been investigated in two studies to date. Edinburgh et al., (2020) reported increased oral glucose sensitivity in overweight/obese individuals completing 6 weeks of training in the fasted state. Additionally, 12-weeks of endurance training in males diagnosed with T2D and randomised to exercising in an overnight fasted state or after breakfast, saw greater improvements in HbA1c when an exercise programme is completed in the fed state compared to the fasted state. There was no difference in the ability to reduce fat mass, increase fat oxidation or improve HDL concentrations between nutritional strategies (Verboven et al., 2020). While fasted exercise typically enhances IMTG utilisation, it is not yet clear whether this translates into greater benefits to insulin sensitivity and other markers of cardiometabolic health.

*Training with low muscle glycogen availability*

The “train-low” paradigm has gained interest over the last decade, and typically consists of performing an initial bout of high intensity exercise (to reduce muscle glycogen concentrations) before a second exercise bout is performed several hours later or the following morning. Importantly, the ingestion of carbohydrate is restricted between sessions to prevent muscle glycogen resynthesis, such that the second exercise bout is commenced with low muscle glycogen concentrations. Under these conditions, rates of whole-body fat oxidation are augmented (Hansen *et al.*, 2005; Yeo *et al.*, 2008b; Morton *et al.*, 2009; Hulston *et al.*, 2010) and while there is no direct evidence for an increased IMTG utilisation when exercising with low muscle glycogen, one would speculate that this does occur. If true, this strategy could prove to be applicable for individuals with metabolic disease. Note though that research in this area to date is limited to highly trained males.

Importantly, in trained males systematically commencing exercise with reduced muscle glycogen availability augments the activation of signalling proteins (Baar and McGee, 2008) leading to elevated gene expression of mitochondrial proteins (Bartlett, Hawley and Morton, 2015). The signalling pathways stimulated by a reduced muscle glycogen availability have been explained in detail elsewhere (Hawley *et al.*, 2018; Hearris *et al.*, 2018; Impey *et al.*, 2018). Briefly, low muscle glycogen availability stimulates greater activity of AMPK and p38MAPK (Wojtaszewski *et al.*, 2003; Chan *et al.*, 2004), which in turn leads to activation and translocation of p53 and PGC-1 $\alpha$  to the nucleus and mitochondria (Bartlett *et al.*, 2013; Andrade-Souza *et al.*, 2019). Here, these proteins help regulate the transcription of key mitochondrial proteins and those involved in mitochondrial fusion and fission. Exercising under conditions of reduced muscle glycogen availability also enhances circulating FFA concentrations, which in turn activates the nuclear transcription factor PPAR $\delta$  (Philp *et al.*, 2013), to upregulate the expression of proteins linked to lipid metabolism, including  $\beta$ -HAD (Yeo *et al.*, 2008b; Hulston *et al.*, 2010), HSL (Arkininstall *et al.*, 2004), and the FA transport proteins, FATBP and CD36 (Arkininstall *et al.*, 2004; De Bock *et al.*, 2008; Lane *et al.*, 2015). With reduced CHO oxidation, there is a concomitant increase in total lipid oxidation (Hearris *et al.*, 2019) and step wise increases in AMPK activation showing that “train-low” provides a potent stimulus for promoting endurance adaptation.

Studies which incorporate training sessions that are commenced with low muscle glycogen availability over several weeks report increased activity and content of the mitochondrial proteins citrate synthase,  $\beta$ -HAD, and SDH (Hansen et al., 2005; Yeo et al., 2008a; Morton et al., 2009), alongside elevations in whole-body fat oxidation (Yeo et al., 2008a), and result in a greater contribution of IMTG to total energy expenditure during moderate-intensity exercise in well-trained cyclists (Hulston et al., 2010). Increasing oxidative enzyme capacity supports greater fat oxidation, which in athletic populations, is a key consideration for improving substrate utilisation and promoting glycogen sparing, thereby enhancing performance. From a clinical perspective, manipulating carbohydrate availability around exercise is also of relevance, since the adaptations outlined above would likely contribute to an improved IMTG turnover and underpin an increase in insulin sensitivity. Of course, using a prior exercise session to create a stimulus of low muscle glycogen availability (i.e., the 'traditional' train-low model) is challenging in sedentary, overweight/obese individuals, who likely will not perform exercise of sufficient intensity and/or duration to reduce glycogen below the 'threshold' required to stimulate the signalling responses mentioned above. Therefore, whilst effective, the traditional train-low model may only be adopted in overweight/obese individuals with high motivation and who have access to a specialist support network.

#### *Low carbohydrate, high fat (LCHF) diet*

Using diet may be a more appropriate strategy to generate a state of low muscle glycogen availability in sedentary overweight/obese individuals with or without T2D. To target fat oxidation, macronutrient intake can be manipulated by either a non-ketogenic low-carbohydrate high-fat diet (where fat supplies 60-65% fat), or by a ketogenic low-carbohydrate high-fat diet (where fat supplies 75-80% of daily intake) (Burke et al., 2020). The purpose of a ketogenic diet is to induce fasting-like effects and leads to the production of ketone bodies, which can provide an additional substrate for oxidative energy production. Moreover, a ketogenic diet therefore leads to increases in whole-body fat oxidation rates, and a subsequent reduction in whole-body carbohydrate oxidation, which can be explained entirely by a decrease in muscle glycogen utilization (Starling et al., 1997; Burke et al., 2000; Helge et al., 2001). Unlike

the above methods of glycogen manipulation, a ketogenic diet is typically a chronic strategy to maximise fat as a source of fuel (Burke, 2021).

The specific adaptations to a ketogenic diet, and the timeline of these adaptive responses, are controversial (Lindseth, 2017; Burke *et al.*, 2020), but in terms of exercise capacity, they seem to be related to increased delivery, uptake and oxidation of free fatty acids in skeletal muscle (see detailed review by (Burke, 2021). Adaptations to a ketogenic diet in trained athletes can occur as quickly as within 5 days but are often accompanied by feelings of fatigue due to exposure to this extreme change in diet (Burke *et al.*, 2017). Over a longer period (12 weeks) though, it was recently reported that a LCHF ketogenic diet combined with exercise increased mitochondrial respiratory control ratio, ATP production and muscle triglyceride content (Miller *et al.*, 2020). Whilst there is still little data on changes in mitochondrial proteins, LD proteins or IMTG content in human skeletal muscle in response to a ketogenic diet, there is a robust increase in whole-body fat oxidation.

A non-ketogenic LCHF diet also enhances whole-body fat oxidation studies investigating the mechanisms for greater oxidative capacity due to high fat diet typically investigate this short-term using a non-ketogenic LCHF diet. Following 5 days of a high fat diet, increases in whole-body fat oxidation occurred, despite reduced skeletal muscle mitochondrial respiration (Leckey *et al.*, 2018). Importantly, a 12% increase in FAT/CD36 protein was observed, which suggests an increased capacity for FA uptake (Leckey *et al.*, 2018). Therefore, it is likely that increased whole-body fat oxidation following a non-ketogenic LCHF diet is the result of increased transport and delivery of FA to skeletal muscle. To date, no study has detailed the mechanisms underpinning the increase in fat utilisation in human skeletal muscle, yet previous studies has shown that short-term exposure (~5 days) to a high fat diet increases IMTG content (Yeo *et al.*, 2008a), HSL content (Stellingwerff *et al.*, 2006), and the protein abundance of FAT/CD36 (Cameron-Smith *et al.*, 2003) and carnitine palmitoyl transferase (Goedecke *et al.*, 1999). These adaptations demonstrate an adaptive response which could improve the capacity of the exercising muscle to increase the breakdown and resynthesis of IMTG stores in response to long-term ingestion of a LCHF diet. Future studies should then investigate whether an improved turnover of fat

at a whole-body and muscle level contribute to the improved glycaemic control reported following a LCHF diet (Ahmed et al., 2020).

Whilst the above addresses the potential impact of chronic changes in diet on IMTG turnover, it should also be noted that macronutrient intake in the hours and days following an exercise bout can impact IMTG stores. For example, traditional sports nutrition guidelines suggest the consumption of a high carbohydrate diet following exercise in order to support glycogen resynthesis (Ivy, 1991). However, when diets high in carbohydrate, and therefore low in fat (<15% energy from fat), are consumed post-exercise, IMTG resynthesis is shown to be substantially impaired (Decombaz et al., 2000; Decombaz et al., 2001; Larson-Meyer, Newcomer and Hunter, 2002). Even when the diet is made up of ~24% energy from fat (which could be considered quite typical of a high carbohydrate diet), IMTG repletion over the subsequent 48 h following exercise is still impaired (van Loon et al., 2003b). To date, work in this area has only used healthy, trained individuals who have high rates of IMTG utilisation during exercise resulting in a net decrease in IMTG content post-exercise. Given that a net decrease in IMTG content following exercise appears to be absent in obese individuals or T2D patients, a high-fat diet in the period following exercise could theoretically contribute to aberrant storage of IMTG and the generation of lipid metabolites. Future research is required to examine this though.

#### *Weight loss and Calorie restriction (CR) on IMTG utilisation*

Weight loss achieved via a restriction in caloric intake (but where the relative macronutrient contribution remains the same) is a powerful strategy to improve insulin sensitivity in overweight and obese individuals with or without T2D (Moore et al., 2000). Notably, diet-induced weight loss often results in a decrease in IMTG content (Goodpaster et al., 2000; Dube et al., 2011), and this occurs alongside reductions in the concentrations DAGs and ceramides in muscle (Dube et al., 2011), suggesting a remodelling of the intramuscular lipid pool.

Six months of CR increased markers of mitochondrial biogenesis and mitochondrial DNA and reduced oxidative stress markers in overweight individuals (Civitarese et al., 2007), but key enzymes of the TCA cycle,  $\beta$ -oxidation and electron transport chain

were unchanged. Over 16-weeks, CR has shown to increase citrate synthase activity (Menshikova *et al.*, 2017) and reduce skeletal muscle lipid content (Goodpaster *et al.*, 2000), but does not alter mitochondria volume or enzymes from the beta oxidation pathway (Menshikova *et al.*, 2017). Even when similar weight loss is achieved, a CR intervention alone does not achieve improvements in mitochondrial content or electron transport chain enzyme activity, whereas a CR plus exercise intervention does (Toledo *et al.*, 2008). CR has also been shown to reduce IMTG in skeletal muscle, as well as mRNA of genes involved in lipogenesis and FA transport yet showed no change in mitochondrial content or mRNA genes involved in mitochondrial biogenesis (Sparks *et al.*, 2017).

A combined intervention of exercise and CR in an athletic population increases IMTG content (Nadeau *et al.*, 2006), and skeletal muscle oxidative capacity (Pruchnic *et al.*, 2004; Nadeau *et al.*, 2006). Conversely, weight loss induced by a combination of exercise and CR in obese individuals either decreases (Rabol *et al.*, 2009), or does not change IMTG content, but leads to reductions in LD size and increases mitochondrial content (He, Goodpaster and Kelley, 2004). This suggests that a combined exercise and weight loss intervention remodels the LD pool such that the ability to utilise fat as a fuel source and regularly turnover IMTG would be enhanced, rather than simply reducing the amount of IMTG in skeletal muscle. Considering long-term CR interventions fail to improve oxidative capacity, a combined intervention of CR and exercise to reduce lipid metabolites and improve mitochondrial oxidative capacity, is likely the most powerful strategy to increase IMTG utilisation and improve skeletal muscle lipid turnover in individuals at risk of T2D.

## Summary

It is now widely agreed that the absolute level of IMTG is unrelated to insulin resistance, but rather a high rate of turnover of the IMTG pool appears to be fundamental to the preservation of insulin sensitivity. Thus, in the trained state IMTG represents a highly dynamic lipid pool within muscle that can be adjusted relative to metabolic demand. Chronic exercise training creates a stimulus of regular breakdown and resynthesis of the IMTG pool, reducing the potential for accumulation of toxic lipid metabolites and therefore the risk of skeletal muscle insulin resistance. In obese

individuals and people with T2D there is a collective imbalance between the rate of FA uptake into muscle, esterification and storage, IMTG breakdown and oxidation. Despite many years of research, the molecular mechanisms underlying this imbalance are yet to be fully understood. The discovery that the PLIN proteins may play a role regulating LD dynamics has provided additional understanding over the last decade, but efforts to identify the mechanisms responsible for the low turnover of the IMTG pool in obesity and T2D should continue to form the basis of future work. Exercise training studies demonstrate that improved IMTG turnover is a key adaptation contributing to improved insulin sensitivity at both the skeletal muscle and whole-body level. Importantly, there appears to be potential for the use of pharmacological or nutritional strategies to maximise the insulin-sensitising effect of exercise interventions. Future work is now required to test these interventions in the longer term in clinical populations, with consideration for how these interventions can be successfully adopted in the real world.

## Author statements

Competing interests: The authors declare there are no competing interests.

Contributor's statement: All authors have approved the final version of the manuscript and agree to be accountable for all aspects of the work. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

## References

Ahmed, S.R., Bellamkonda, S., Zilbermint, M., Wang, J. and Kalyani, R.R. (2020) Effects of the low carbohydrate, high fat diet on glycemic control and body weight in patients with type 2 diabetes: experience from a community-based cohort. *BMJ Open Diabetes Res Care*, 8 (1). 10.1136/bmjdr-2019-000980.

Alsted, T.J., Nybo, L., Schweiger, M., Fledelius, C., Jacobsen, P., Zimmermann, R., Zechner, R. and Kiens, B. (2009) Adipose triglyceride lipase in human skeletal muscle is upregulated by exercise training. *Am J Physiol Endocrinol Metab*, 296 (3), E445-453. 10.1152/ajpendo.90912.2008.

Amati, F., Dube, J.J., Alvarez-Carnero, E., Edreira, M.M., Chomentowski, P., Coen, P.M., Switzer, G.E., Bickel, P.E., Stefanovic-Racic, M., Toledo, F.G. and Goodpaster, B.H. (2011) Skeletal muscle triglycerides, diacylglycerols, and ceramides in insulin



789 resistance: another paradox in endurance-trained athletes? *Diabetes*, 60 (10), 2588-  
790 2597. 10.2337/db10-1221.

791  
792 Andrade-Souza, V.A., Ghiarone, T., Sansonio, A., Silva, K.A.S., Tomazini, F.,  
793 Arcoverde, L., Fyfe, J., Perri, E., Saner, N. and Kuang, J. (2019) Exercise twice-a-day  
794 potentiates skeletal muscle signalling responses associated with mitochondrial  
795 biogenesis in humans, which are independent of lowered muscle glycogen content.  
796 *bioRxiv*, 547489.

797  
798 Anthonsen, M.W., Ronnstrand, L., Wernstedt, C., Degerman, E. and Holm, C. (1998)  
799 Identification of novel phosphorylation sites in hormone-sensitive lipase that are  
800 phosphorylated in response to isoproterenol and govern activation properties in vitro.  
801 *J Biol Chem*, 273 (1), 215-221. 10.1074/jbc.273.1.215.

802  
803 Arkinstall, M.J., Bruce, C.R., Nikolopoulos, V., Garnham, A.P. and Hawley, J.A. (2001)  
804 Effect of carbohydrate ingestion on metabolism during running and cycling. *J Appl*  
805 *Physiol* (1985), 91 (5), 2125-2134. 10.1152/jappl.2001.91.5.2125.

806  
807 Arkinstall, M.J., Tunstall, R.J., Cameron-Smith, D. and Hawley, J.A. (2004) Regulation  
808 of metabolic genes in human skeletal muscle by short-term exercise and diet  
809 manipulation. *Am J Physiol Endocrinol Metab*, 287 (1), E25-31.  
810 10.1152/ajpendo.00557.2003.

811  
812 Astorino, T.A. and Schubert, M.M. (2018) Changes in fat oxidation in response to  
813 various regimes of high intensity interval training (HIIT). *Eur J Appl Physiol*, 118 (1),  
814 51-63. 10.1007/s00421-017-3756-0.

815  
816 Axelsen, M., Smith, U., Eriksson, J.W., Taskinen, M.R. and Jansson, P.A. (1999)  
817 Postprandial hypertriglyceridemia and insulin resistance in normoglycemic first-  
818 degree relatives of patients with type 2 diabetes. *Ann Intern Med*, 131 (1), 27-31.  
819 10.7326/0003-4819-131-1-199907060-00006.

820  
821 Baar, K. and McGee, S. (2008) Optimizing training adaptations by manipulating  
822 glycogen. *European journal of sport science*, 8 (2), 97-106.

823  
824 Bachmann, O.P., Dahl, D.B., Brechtel, K., Machann, J., Haap, M., Maier, T.,  
825 Loviscach, M., Stumvoll, M., Claussen, C.D., Schick, F., Haring, H.U. and Jacob, S.  
826 (2001) Effects of intravenous and dietary lipid challenge on intramyocellular lipid  
827 content and the relation with insulin sensitivity in humans. *Diabetes*, 50 (11), 2579-  
828 2584. 10.2337/diabetes.50.11.2579.

829  
830 Badin, P.M., Langin, D. and Moro, C. (2013) Dynamics of skeletal muscle lipid pools.  
831 *Trends Endocrinol Metab*, 24 (12), 607-615. 10.1016/j.tem.2013.08.001.

832  
833 Badin, P.M., Louche, K., Mairal, A., Liebisch, G., Schmitz, G., Rustan, A.C., Smith,  
834 S.R., Langin, D. and Moro, C. (2011) Altered skeletal muscle lipase expression and  
835 activity contribute to insulin resistance in humans. *Diabetes*, 60 (6), 1734-1742.  
836 10.2337/db10-1364.

837

Bajaj, M., Suraamornkul, S., Romanelli, A., Cline, G.W., Mandarino, L.J., Shulman, G.I. and DeFronzo, R.A. (2005) Effect of a sustained reduction in plasma free fatty acid concentration on intramuscular long-chain fatty Acyl-CoAs and insulin action in type 2 diabetic patients. *Diabetes*, 54 (11), 3148-3153. 10.2337/diabetes.54.11.3148.

Balakrishnan, V.S., Rao, M., Menon, V., Gordon, P.L., Pilichowska, M., Castaneda, F. and Castaneda-Sceppa, C. (2010) Resistance training increases muscle mitochondrial biogenesis in patients with chronic kidney disease. *Clin J Am Soc Nephrol*, 5 (6), 996-1002. 10.2215/CJN.09141209.

Baldwin, K.M., Klinkerfuss, G.H., Terjung, R.L., Mole, P.A. and Holloszy, J.O. (1972) Respiratory capacity of white, red, and intermediate muscle: adaptative response to exercise. *Am J Physiol*, 222 (2), 373-378. 10.1152/ajplegacy.1972.222.2.373.

Banner, C.D., Gottlicher, M., Widmark, E., Sjoval, J., Rafter, J.J. and Gustafsson, J.A. (1993) A systematic analytical chemistry/cell assay approach to isolate activators of orphan nuclear receptors from biological extracts: characterization of peroxisome proliferator-activated receptor activators in plasma. *J Lipid Res*, 34 (9), 1583-1591. <https://www.ncbi.nlm.nih.gov/pubmed/8228641>

Bartlett, J.D., Hawley, J.A. and Morton, J.P. (2015) Carbohydrate availability and exercise training adaptation: too much of a good thing? *Eur J Sport Sci*, 15 (1), 3-12. 10.1080/17461391.2014.920926.

Bartlett, J.D., Louhelainen, J., Iqbal, Z., Cochran, A.J., Gibala, M.J., Gregson, W., Close, G.L., Drust, B. and Morton, J.P. (2013) Reduced carbohydrate availability enhances exercise-induced p53 signaling in human skeletal muscle: implications for mitochondrial biogenesis. *Am J Physiol Regul Integr Comp Physiol*, 304 (6), R450-458. 10.1152/ajpregu.00498.2012.

Benador, I.Y., Veliova, M., Liesa, M. and Shirihai, O.S. (2019) Mitochondria Bound to Lipid Droplets: Where Mitochondrial Dynamics Regulate Lipid Storage and Utilization. *Cell Metab*, 29 (4), 827-835. 10.1016/j.cmet.2019.02.011.

Benador, I.Y., Veliova, M., Mahdavian, K., Petcherski, A., Wikstrom, J.D., Assali, E.A., Acin-Perez, R., Shum, M., Oliveira, M.F., Cinti, S., Sztalryd, C., Barshop, W.D., Wohlschlegel, J.A., Corkey, B.E., Liesa, M. and Shirihai, O.S. (2018) Mitochondria Bound to Lipid Droplets Have Unique Bioenergetics, Composition, and Dynamics that Support Lipid Droplet Expansion. *Cell Metab*, 27 (4), 869-885 e866. 10.1016/j.cmet.2018.03.003.

Bergman, B.C., Butterfield, G.E., Wolfel, E.E., Casazza, G.A., Lopaschuk, G.D. and Brooks, G.A. (1999) Evaluation of exercise and training on muscle lipid metabolism. *Am J Physiol*, 276 (1), E106-117. 10.1152/ajpendo.1999.276.1.E106.

Bergman, B.C. and Goodpaster, B.H. (2020) Exercise and Muscle Lipid Content, Composition, and Localization: Influence on Muscle Insulin Sensitivity. *Diabetes*, 69 (5), 848-858. 10.2337/dbi18-0042.

- Bergman, B.C., Perreault, L., Strauss, A., Bacon, S., Kerege, A., Harrison, K., Brozinick, J.T., Hunerdosse, D.M., Playdon, M.C., Holmes, W., Bui, H.H., Sanders, P., Siddall, P., Wei, T., Thomas, M.K., Kuo, M.S. and Eckel, R.H. (2018) Intramuscular triglyceride synthesis: importance in muscle lipid partitioning in humans. *Am J Physiol Endocrinol Metab*, 314 (2), E152-E164. 10.1152/ajpendo.00142.2017.
- Blaak, E.E., Schiffelers, S.L., Saris, W.H., Mensink, M. and Kooi, M.E. (2004) Impaired beta-adrenergically mediated lipolysis in skeletal muscle of obese subjects. *Diabetologia*, 47 (8), 1462-1468. 10.1007/s00125-004-1471-y.
- Blaak, E.E. and Wagenmakers, A.J. (2002) The fate of [U-(13)C]palmitate extracted by skeletal muscle in subjects with type 2 diabetes and control subjects. *Diabetes*, 51 (3), 784-789. 10.2337/diabetes.51.3.784.
- Boden, G. (2003) Effects of free fatty acids (FFA) on glucose metabolism: significance for insulin resistance and type 2 diabetes. *Exp Clin Endocrinol Diabetes*, 111 (3), 121-124. 10.1055/s-2003-39781.
- Bosma, M., Hesselink, M.K., Sparks, L.M., Timmers, S., Ferraz, M.J., Mattijssen, F., van Beurden, D., Schaart, G., de Baets, M.H., Verheyen, F.K., Kersten, S. and Schrauwen, P. (2012) Perilipin 2 improves insulin sensitivity in skeletal muscle despite elevated intramuscular lipid levels. *Diabetes*, 61 (11), 2679-2690. 10.2337/db11-1402.
- Bulankina, A.V., Deggerich, A., Wenzel, D., Mutenda, K., Wittmann, J.G., Rudolph, M.G., Burger, K.N. and Honing, S. (2009) TIP47 functions in the biogenesis of lipid droplets. *J Cell Biol*, 185 (4), 641-655. 10.1083/jcb.200812042.
- Burgomaster, K.A., Howarth, K.R., Phillips, S.M., Rakobowchuk, M., Macdonald, M.J., McGee, S.L. and Gibala, M.J. (2008) Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *J Physiol*, 586 (1), 151-160. 10.1113/jphysiol.2007.142109.
- Burke, L.M. (2021) Ketogenic low-CHO, high-fat diet: the future of elite endurance sport? *J Physiol*, 599 (3), 819-843. 10.1113/JP278928.
- Burke, L.M., Angus, D.J., Cox, G.R., Cummings, N.K., Febbraio, M.A., Gawthorn, K., Hawley, J.A., Minehan, M., Martin, D.T. and Hargreaves, M. (2000) Effect of fat adaptation and carbohydrate restoration on metabolism and performance during prolonged cycling. *J Appl Physiol* (1985), 89 (6), 2413-2421. 10.1152/jappl.2000.89.6.2413.
- Burke, L.M., Ross, M.L., Garvican-Lewis, L.A., Welvaert, M., Heikura, I.A., Forbes, S.G., Mirtschin, J.G., Cato, L.E., Strobel, N., Sharma, A.P. and Hawley, J.A. (2017) Low carbohydrate, high fat diet impairs exercise economy and negates the performance benefit from intensified training in elite race walkers. *J Physiol*, 595 (9), 2785-2807. 10.1113/JP273230.
- Burke, L.M., Sharma, A.P., Heikura, I.A., Forbes, S.F., Holloway, M., McKay, A.K.A., Bone, J.L., Leckey, J.J., Welvaert, M. and Ross, M.L. (2020) Crisis of confidence averted: Impairment of exercise economy and performance in elite race walkers by

ketogenic low carbohydrate, high fat (LCHF) diet is reproducible. *Plos One*, 15 (6), e0234027. 10.1371/journal.pone.0234027.

Cameron-Smith, D., Burke, L.M., Angus, D.J., Tunstall, R.J., Cox, G.R., Bonen, A., Hawley, J.A. and Hargreaves, M. (2003) A short-term, high-fat diet up-regulates lipid metabolism and gene expression in human skeletal muscle. *Am J Clin Nutr*, 77 (2), 313-318. 10.1093/ajcn/77.2.313.

Canto, C., Jiang, L.Q., Deshmukh, A.S., Matak, C., Coste, A., Lagouge, M., Zierath, J.R. and Auwerx, J. (2010) Interdependence of AMPK and SIRT1 for metabolic adaptation to fasting and exercise in skeletal muscle. *Cell Metab*, 11 (3), 213-219. 10.1016/j.cmet.2010.02.006.

Chan, M.H., McGee, S.L., Watt, M.J., Hargreaves, M. and Febbraio, M.A. (2004) Altering dietary nutrient intake that reduces glycogen content leads to phosphorylation of nuclear p38 MAP kinase in human skeletal muscle: association with IL-6 gene transcription during contraction. *FASEB J*, 18 (14), 1785-1787. 10.1096/fj.03-1039fje.

Chaurasia, B. and Summers, S.A. (2015) Ceramides - Lipotoxic Inducers of Metabolic Disorders. *Trends Endocrinol Metab*, 26 (10), 538-550. 10.1016/j.tem.2015.07.006.

Chen, W., Chang, B., Wu, X., Li, L., Sleeman, M. and Chan, L. (2013) Inactivation of Plin4 downregulates Plin5 and reduces cardiac lipid accumulation in mice. *Am J Physiol Endocrinol Metab*, 304 (7), E770-779. 10.1152/ajpendo.00523.2012.

Chow, L.S., Mashek, D.G., Austin, E., Eberly, L.E., Persson, X.M., Mashek, M.T., Seaquist, E.R. and Jensen, M.D. (2014) Training status diverges muscle diacylglycerol accumulation during free fatty acid elevation. *Am J Physiol Endocrinol Metab*, 307 (1), E124-131. 10.1152/ajpendo.00166.2014.

Christ-Roberts, C.Y., Pratipanawatr, T., Pratipanawatr, W., Berria, R., Belfort, R., Kashyap, S. and Mandarino, L.J. (2004) Exercise training increases glycogen synthase activity and GLUT4 expression but not insulin signaling in overweight nondiabetic and type 2 diabetic subjects. *Metabolism*, 53 (9), 1233-1242. 10.1016/j.metabol.2004.03.022.

Chrois, K.M., Dohlmann, T.L., Sogaard, D., Hansen, C.V., Dela, F., Helge, J.W. and Larsen, S. (2020) Mitochondrial adaptations to high intensity interval training in older females and males. *Eur J Sport Sci*, 20 (1), 135-145. 10.1080/17461391.2019.1615556.

Civitarese, A.E., Carling, S., Heilbronn, L.K., Hulver, M.H., Ukropcova, B., Deutsch, W.A., Smith, S.R., Ravussin, E. and Team, C.P. (2007) Calorie restriction increases muscle mitochondrial biogenesis in healthy humans. *PLoS Med*, 4 (3), e76. 10.1371/journal.pmed.0040076.

Coggan, A.R., Raguso, C.A., Gastaldelli, A., Sidossis, L.S. and Yeckel, C.W. (2000) Fat metabolism during high-intensity exercise in endurance-trained and untrained men. *Metabolism*, 49 (1), 122-128. 10.1016/s0026-0495(00)90963-6.

- Covington, J.D., Galgani, J.E., Moro, C., LaGrange, J.M., Zhang, Z., Rustan, A.C., Ravussin, E. and Bajpeyi, S. (2014) Skeletal muscle perilipin 3 and coatomer proteins are increased following exercise and are associated with fat oxidation. *Plos One*, 9 (3), e91675. 10.1371/journal.pone.0091675.
- Covington, J.D., Johannsen, D.L., Coen, P.M., Burk, D.H., Obanda, D.N., Ebenezer, P.J., Tam, C.S., Goodpaster, B.H., Ravussin, E. and Bajpeyi, S. (2017) Intramyocellular Lipid Droplet Size Rather Than Total Lipid Content is Related to Insulin Sensitivity After 8 Weeks of Overfeeding. *Obesity (Silver Spring)*, 25 (12), 2079-2087. 10.1002/oby.21980.
- Crane, J.D., Devries, M.C., Safdar, A., Hamadeh, M.J. and Tarnopolsky, M.A. (2010) The effect of aging on human skeletal muscle mitochondrial and intramyocellular lipid ultrastructure. *J Gerontol A Biol Sci Med Sci*, 65 (2), 119-128. 10.1093/gerona/glp179.
- Crepaldi, G., Avogaro, P., Descovich, G.C., Di Perri, T., Postiglione, A., Sirtori, C.R., Strano, A., Ventura, S. and Musatti, L. (1988) Plasma lipid lowering activity of acipimox in patients with type II and type IV hyperlipoproteinemia. Results of a multicenter trial. *Atherosclerosis*, 70 (1-2), 115-121. 10.1016/0021-9150(88)90105-0.
- Croymans, D.M., Paparisto, E., Lee, M.M., Brandt, N., Le, B.K., Lohan, D., Lee, C.C. and Roberts, C.K. (2013) Resistance training improves indices of muscle insulin sensitivity and beta-cell function in overweight/obese, sedentary young men. *J Appl Physiol* (1985), 115 (9), 1245-1253. 10.1152/jappphysiol.00485.2013.
- Daemen, S., Gemmink, A., Brouwers, B., Meex, R.C.R., Huntjens, P.R., Schaart, G., Moonen-Kornips, E., Jorgensen, J., Hoeks, J., Schrauwen, P. and Hesselink, M.K.C. (2018) Distinct lipid droplet characteristics and distribution unmask the apparent contradiction of the athlete's paradox. *Mol Metab*, 17, 71-81. 10.1016/j.molmet.2018.08.004.
- Daemen, S., van Polanen, N. and Hesselink, M.K.C. (2018) The effect of diet and exercise on lipid droplet dynamics in human muscle tissue. *J Exp Biol*, 221 (Pt Suppl 1). 10.1242/jeb.167015.
- Daniele, G., Eldor, R., Merovci, A., Clarke, G.D., Xiong, J., Tripathy, D., Taranova, A., Abdul-Ghani, M. and DeFronzo, R.A. (2014) Chronic reduction of plasma free fatty acid improves mitochondrial function and whole-body insulin sensitivity in obese and type 2 diabetic individuals. *Diabetes*, 63 (8), 2812-2820. 10.2337/db13-1130.
- De Bock, K., Derave, W., Eijnde, B.O., Hesselink, M.K., Koninckx, E., Rose, A.J., Schrauwen, P., Bonen, A., Richter, E.A. and Hespel, P. (2008) Effect of training in the fasted state on metabolic responses during exercise with carbohydrate intake. *J Appl Physiol* (1985), 104 (4), 1045-1055. 10.1152/jappphysiol.01195.2007.
- De Bock, K., Richter, E.A., Russell, A.P., Eijnde, B.O., Derave, W., Ramaekers, M., Koninckx, E., Leger, B., Verhaeghe, J. and Hespel, P. (2005) Exercise in the fasted state facilitates fibre type-specific intramyocellular lipid breakdown and stimulates glycogen resynthesis in humans. *J Physiol*, 564 (Pt 2), 649-660. 10.1113/jphysiol.2005.083170.

- Decombaz, J., Fleith, M., Hoppeler, H., Kreis, R. and Boesch, C. (2000) Effect of diet on the replenishment of intramyocellular lipids after exercise. *Eur J Nutr*, 39 (6), 244-247. 10.1007/s003940070002.
- Decombaz, J., Schmitt, B., Ith, M., Decarli, B., Diem, P., Kreis, R., Hoppeler, H. and Boesch, C. (2001) Postexercise fat intake repletes intramyocellular lipids but no faster in trained than in sedentary subjects. *Am J Physiol Regul Integr Comp Physiol*, 281 (3), R760-769. 10.1152/ajpregu.2001.281.3.R760.
- Deshmukh, A.S., Murgia, M., Nagaraj, N., Treebak, J.T., Cox, J. and Mann, M. (2015) Deep proteomics of mouse skeletal muscle enables quantitation of protein isoforms, metabolic pathways, and transcription factors. *Mol Cell Proteomics*, 14 (4), 841-853. 10.1074/mcp.M114.044222.
- Devries, M.C., Samjoo, I.A., Hamadeh, M.J., McCready, C., Raha, S., Watt, M.J., Steinberg, G.R. and Tarnopolsky, M.A. (2013) Endurance training modulates intramyocellular lipid compartmentalization and morphology in skeletal muscle of lean and obese women. *J Clin Endocrinol Metab*, 98 (12), 4852-4862. 10.1210/jc.2013-2044.
- Dube, J.J., Amati, F., Toledo, F.G., Stefanovic-Racic, M., Rossi, A., Coen, P. and Goodpaster, B.H. (2011) Effects of weight loss and exercise on insulin resistance, and intramyocellular triacylglycerol, diacylglycerol and ceramide. *Diabetologia*, 54 (5), 1147-1156. 10.1007/s00125-011-2065-0.
- Edinburgh, R.M., Bradley, H.E., Abdullah, N.F., Robinson, S.L., Chrzanowski-Smith, O.J., Walhin, J.P., Joannis, S., Manolopoulos, K.N., Philp, A., Hengist, A., Chabowski, A., Brodsky, F.M., Koumanov, F., Betts, J.A., Thompson, D., Wallis, G.A. and Gonzalez, J.T. (2020) Lipid Metabolism Links Nutrient-Exercise Timing to Insulin Sensitivity in Men Classified as Overweight or Obese. *J Clin Endocrinol Metab*, 105 (3). 10.1210/clinem/dgz104.
- Eichmann, T.O., Kumari, M., Haas, J.T., Farese, R.V., Jr., Zimmermann, R., Lass, A. and Zechner, R. (2012) Studies on the substrate and stereo/regioselectivity of adipose triglyceride lipase, hormone-sensitive lipase, and diacylglycerol-O-acyltransferases. *J Biol Chem*, 287 (49), 41446-41457. 10.1074/jbc.M112.400416.
- Enoksson, S., Degerman, E., Hagstrom-Toft, E., Large, V. and Arner, P. (1998) Various phosphodiesterase subtypes mediate the in vivo antilipolytic effect of insulin on adipose tissue and skeletal muscle in man. *Diabetologia*, 41 (5), 560-568. 10.1007/s001250050947.
- Fell, J.M., Hearn, M.A., Ellis, D.G., Moran, J.E.P., Jevons, E.F.P., Owens, D.J., Strauss, J.A., Cocks, M., Louis, J.B., Shepherd, S.O. and Morton, J.P. (2021) Carbohydrate improves exercise capacity but does not affect subcellular lipid droplet morphology, AMPK and p53 signalling in human skeletal muscle. *J Physiol*, 599 (11), 2823-2849. 10.1113/JP281127.

- Forouhi, N.G., Jenkinson, G., Thomas, E.L., Mullick, S., Mierisova, S., Bhonsle, U., McKeigue, P.M. and Bell, J.D. (1999) Relation of triglyceride stores in skeletal muscle cells to central obesity and insulin sensitivity in European and South Asian men. *Diabetologia*, 42 (8), 932-935. 10.1007/s001250051250.
- Frayn, K.N. (2002) Adipose tissue as a buffer for daily lipid flux. *Diabetologia*, 45 (9), 1201-1210. 10.1007/s00125-002-0873-y.
- Fredrikson, G., Stralfors, P., Nilsson, N.O. and Belfrage, P. (1981) Hormone-sensitive lipase from adipose tissue of rat. *Methods Enzymol*, 71 Pt C, 636-646. 10.1016/0076-6879(81)71076-0.
- Frosig, C., Rose, A.J., Treebak, J.T., Kiens, B., Richter, E.A. and Wojtaszewski, J.F. (2007) Effects of endurance exercise training on insulin signaling in human skeletal muscle: interactions at the level of phosphatidylinositol 3-kinase, Akt, and AS160. *Diabetes*, 56 (8), 2093-2102. 10.2337/db06-1698.
- Fulcher, G.R., Walker, M., Catalano, C., Agius, L. and Alberti, K.G. (1992) Metabolic effects of suppression of nonesterified fatty acid levels with acipimox in obese NIDDM subjects. *Diabetes*, 41 (11), 1400-1408. 10.2337/diab.41.11.1400.
- Funai, K. and Semenkovich, C.F. (2011) Skeletal muscle lipid flux: running water carries no poison. *Am J Physiol Endocrinol Metab*, 301 (2), E245-251. 10.1152/ajpendo.00152.2011.
- Gallardo-Montejano, V.I., Saxena, G., Kusminski, C.M., Yang, C., McAfee, J.L., Hahner, L., Hoch, K., Dubinsky, W., Narkar, V.A. and Bickel, P.E. (2016) Nuclear Perilipin 5 integrates lipid droplet lipolysis with PGC-1alpha/SIRT1-dependent transcriptional regulation of mitochondrial function. *Nat Commun*, 7, 12723. 10.1038/ncomms12723.
- Gemmink, A., Bosma, M., Kuijpers, H.J., Hoeks, J., Schaart, G., van Zandvoort, M.A., Schrauwen, P. and Hesselink, M.K. (2016) Decoration of intramyocellular lipid droplets with PLIN5 modulates fasting-induced insulin resistance and lipotoxicity in humans. *Diabetologia*, 59 (5), 1040-1048. 10.1007/s00125-016-3865-z.
- Gibala, M.J., Little, J.P., van Essen, M., Wilkin, G.P., Burgomaster, K.A., Safdar, A., Raha, S. and Tarnopolsky, M.A. (2006) Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *J Physiol*, 575 (Pt 3), 901-911. 10.1113/jphysiol.2006.112094.
- Glatz, J.F.C. and Luiken, J. (2018) Dynamic role of the transmembrane glycoprotein CD36 (SR-B2) in cellular fatty acid uptake and utilization. *J Lipid Res*, 59 (7), 1084-1093. 10.1194/jlr.R082933.
- Goedecke, J.H., Christie, C., Wilson, G., Dennis, S.C., Noakes, T.D., Hopkins, W.G. and Lambert, E.V. (1999) Metabolic adaptations to a high-fat diet in endurance cyclists. *Metabolism*, 48 (12), 1509-1517. 10.1016/s0026-0495(99)90238-x.

- Goodpaster, B.H., He, J., Watkins, S. and Kelley, D.E. (2001) Skeletal muscle lipid content and insulin resistance: evidence for a paradox in endurance-trained athletes. *J Clin Endocrinol Metab*, 86 (12), 5755-5761. 10.1210/jcem.86.12.8075.
- Goodpaster, B.H., Thaete, F.L., Simoneau, J.A. and Kelley, D.E. (1997) Subcutaneous abdominal fat and thigh muscle composition predict insulin sensitivity independently of visceral fat. *Diabetes*, 46 (10), 1579-1585. 10.2337/diacare.46.10.1579.
- Goodpaster, B.H., Theriault, R., Watkins, S.C. and Kelley, D.E. (2000) Intramuscular lipid content is increased in obesity and decreased by weight loss. *Metabolism*, 49 (4), 467-472. 10.1016/s0026-0495(00)80010-4.
- Granata, C., Jamnick, N.A. and Bishop, D.J. (2018) Training-Induced Changes in Mitochondrial Content and Respiratory Function in Human Skeletal Muscle. *Sports Med*, 48 (8), 1809-1828. 10.1007/s40279-018-0936-y.
- Granneman, J.G., Moore, H.P., Mottillo, E.P., Zhu, Z. and Zhou, L. (2011) Interactions of perilipin-5 (Plin5) with adipose triglyceride lipase. *J Biol Chem*, 286 (7), 5126-5135. 10.1074/jbc.M110.180711.
- Haemmerle, G., Lass, A., Zimmermann, R., Gorkiewicz, G., Meyer, C., Rozman, J., Heldmaier, G., Maier, R., Theussl, C., Eder, S., Kratky, D., Wagner, E.F., Klingenspor, M., Hoefler, G. and Zechner, R. (2006) Defective lipolysis and altered energy metabolism in mice lacking adipose triglyceride lipase. *Science*, 312 (5774), 734-737. 10.1126/science.1123965.
- Haemmerle, G., Zimmermann, R., Hayn, M., Theussl, C., Waeg, G., Wagner, E., Sattler, W., Magin, T.M., Wagner, E.F. and Zechner, R. (2002) Hormone-sensitive lipase deficiency in mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *J Biol Chem*, 277 (7), 4806-4815. 10.1074/jbc.M110355200.
- Hansen, A.K., Fischer, C.P., Plomgaard, P., Andersen, J.L., Saltin, B. and Pedersen, B.K. (2005) Skeletal muscle adaptation: training twice every second day vs. training once daily. *J Appl Physiol* (1985), 98 (1), 93-99. 10.1152/japplphysiol.00163.2004.
- Hawley, J.A., Lundby, C., Cotter, J.D. and Burke, L.M. (2018) Maximizing Cellular Adaptation to Endurance Exercise in Skeletal Muscle. *Cell Metab*, 27 (5), 962-976. 10.1016/j.cmet.2018.04.014.
- He, J., Goodpaster, B.H. and Kelley, D.E. (2004) Effects of weight loss and physical activity on muscle lipid content and droplet size. *Obes Res*, 12 (5), 761-769. 10.1038/oby.2004.92.
- Hearris, M.A., Hammond, K.M., Fell, J.M. and Morton, J.P. (2018) Regulation of Muscle Glycogen Metabolism during Exercise: Implications for Endurance Performance and Training Adaptations. *Nutrients*, 10 (3). 10.3390/nu10030298.
- Hearris, M.A., Hammond, K.M., Seaborne, R.A., Stocks, B., Shepherd, S.O., Philp, A., Sharples, A.P., Morton, J.P. and Louis, J.B. (2019) Graded reductions in preexercise



muscle glycogen impair exercise capacity but do not augment skeletal muscle cell signaling: implications for CHO periodization. *J Appl Physiol* (1985), 126 (6), 1587-1597. 10.1152/jappphysiol.00913.2018.

Helge, J.W., Watt, P.W., Richter, E.A., Rennie, M.J. and Kiens, B. (2001) Fat utilization during exercise: adaptation to a fat-rich diet increases utilization of plasma fatty acids and very low density lipoprotein-triacylglycerol in humans. *J Physiol*, 537 (Pt 3), 1009-1020. 10.1111/j.1469-7793.2001.01009.x.

Hojlund, K., Staehr, P., Hansen, B.F., Green, K.A., Hardie, D.G., Richter, E.A., Beck-Nielsen, H. and Wojtaszewski, J.F. (2003) Increased phosphorylation of skeletal muscle glycogen synthase at NH2-terminal sites during physiological hyperinsulinemia in type 2 diabetes. *Diabetes*, 52 (6), 1393-1402. 10.2337/diabetes.52.6.1393.

Hoppeler, H., Billeter, R., Horvath, P.J., Leddy, J.J. and Pendergast, D.R. (1999) Muscle structure with low- and high-fat diets in well-trained male runners. *Int J Sports Med*, 20 (8), 522-526. 10.1055/s-1999-8838.

Horowitz, J.F., Mora-Rodriguez, R., Byerley, L.O. and Coyle, E.F. (1997) Lipolytic suppression following carbohydrate ingestion limits fat oxidation during exercise. *Am J Physiol*, 273 (4), E768-775. 10.1152/ajpendo.1997.273.4.E768.

Hulston, C.J., Venables, M.C., Mann, C.H., Martin, C., Philp, A., Baar, K. and Jeukendrup, A.E. (2010) Training with low muscle glycogen enhances fat metabolism in well-trained cyclists. *Med Sci Sports Exerc*, 42 (11), 2046-2055. 10.1249/MSS.0b013e3181dd5070.

Impey, S.G., Hearris, M.A., Hammond, K.M., Bartlett, J.D., Louis, J., Close, G.L. and Morton, J.P. (2018) Fuel for the Work Required: A Theoretical Framework for Carbohydrate Periodization and the Glycogen Threshold Hypothesis. *Sports Med*, 48 (5), 1031-1048. 10.1007/s40279-018-0867-7.

Ishii, T., Yamakita, T., Sato, T., Tanaka, S. and Fujii, S. (1998) Resistance training improves insulin sensitivity in NIDDM subjects without altering maximal oxygen uptake. *Diabetes Care*, 21 (8), 1353-1355. 10.2337/diacare.21.8.1353.

Jevons, E.F.P., Gejl, K.D., Strauss, J.A., Ortenblad, N. and Shepherd, S.O. (2020) Skeletal muscle lipid droplets are resynthesized before being coated with perilipin proteins following prolonged exercise in elite male triathletes. *Am J Physiol Endocrinol Metab*, 318 (3), E357-E370. 10.1152/ajpendo.00399.2019.

Jocken, J.W., Langin, D., Smit, E., Saris, W.H., Valle, C., Hul, G.B., Holm, C., Arner, P. and Blaak, E.E. (2007) Adipose triglyceride lipase and hormone-sensitive lipase protein expression is decreased in the obese insulin-resistant state. *J Clin Endocrinol Metab*, 92 (6), 2292-2299. 10.1210/jc.2006-1318.

Jocken, J.W., Smit, E., Goossens, G.H., Essers, Y.P., van Baak, M.A., Mensink, M., Saris, W.H. and Blaak, E.E. (2008) Adipose triglyceride lipase (ATGL) expression in

human skeletal muscle is type I (oxidative) fiber specific. *Histochem Cell Biol*, 129 (4), 535-538. 10.1007/s00418-008-0386-y.

Kelley, D.E. and Goodpaster, B.H. (2001) Skeletal muscle triglyceride. An aspect of regional adiposity and insulin resistance. *Diabetes Care*, 24 (5), 933-941. 10.2337/diacare.24.5.933.

Kelley, D.E. and Simoneau, J.A. (1994) Impaired free fatty acid utilization by skeletal muscle in non-insulin-dependent diabetes mellitus. *J Clin Invest*, 94 (6), 2349-2356. 10.1172/JCI117600.

Khan, S.A., Sathyanarayan, A., Mashek, M.T., Ong, K.T., Wollaston-Hayden, E.E. and Mashek, D.G. (2015) ATGL-catalyzed lipolysis regulates SIRT1 to control PGC-1alpha/PPAR-alpha signaling. *Diabetes*, 64 (2), 418-426. 10.2337/db14-0325.

Kiess, B., Essen-Gustavsson, B., Christensen, N.J. and Saltin, B. (1993) Skeletal muscle substrate utilization during submaximal exercise in man: effect of endurance training. *J Physiol*, 469, 459-478. 10.1113/jphysiol.1993.sp019823.

Klein, S., Coyle, E.F. and Wolfe, R.R. (1994) Fat metabolism during low-intensity exercise in endurance-trained and untrained men. *Am J Physiol*, 267 (6 Pt 1), E934-940. 10.1152/ajpendo.1994.267.6.E934.

Kleinert, M., Parker, B.L., Chaudhuri, R., Fazakerley, D.J., Serup, A., Thomas, K.C., Krycer, J.R., Sylow, L., Fritzen, A.M., Hoffman, N.J., Jeppesen, J., Schjerling, P., Ruegg, M.A., Kiess, B., James, D.E. and Richter, E.A. (2016) mTORC2 and AMPK differentially regulate muscle triglyceride content via Perilipin 3. *Mol Metab*, 5 (8), 646-655. 10.1016/j.molmet.2016.06.007.

Koh, H.E., Nielsen, J., Saltin, B., Holmberg, H.C. and Ortenblad, N. (2017) Pronounced limb and fibre type differences in subcellular lipid droplet content and distribution in elite skiers before and after exhaustive exercise. *J Physiol*, 595 (17), 5781-5795. 10.1113/JP274462.

Lane, S.C., Camera, D.M., Lassiter, D.G., Areta, J.L., Bird, S.R., Yeo, W.K., Jeacocke, N.A., Krook, A., Zierath, J.R., Burke, L.M. and Hawley, J.A. (2015) Effects of sleeping with reduced carbohydrate availability on acute training responses. *J Appl Physiol* (1985), 119 (6), 643-655. 10.1152/jappphysiol.00857.2014.

Larsen, S., Danielsen, J.H., Sondergaard, S.D., Sogaard, D., Vigelsoe, A., Dybbøe, R., Skaaby, S., Dela, F. and Helge, J.W. (2015) The effect of high-intensity training on mitochondrial fat oxidation in skeletal muscle and subcutaneous adipose tissue. *Scand J Med Sci Sports*, 25 (1), e59-69. 10.1111/sms.12252.

Larson-Meyer, D.E., Newcomer, B.R. and Hunter, G.R. (2002) Influence of endurance running and recovery diet on intramyocellular lipid content in women: a <sup>1</sup>H NMR study. *Am J Physiol Endocrinol Metab*, 282 (1), E95-E106. 10.1152/ajpendo.2002.282.1.E95.

- Leckey, J.J., Hoffman, N.J., Parr, E.B., Devlin, B.L., Trewin, A.J., Stepto, N.K., Morton, J.P., Burke, L.M. and Hawley, J.A. (2018) High dietary fat intake increases fat oxidation and reduces skeletal muscle mitochondrial respiration in trained humans. *FASEB J*, 32 (6), 2979-2991. 10.1096/fj.201700993R.
- Li, M., Paran, C., Wolins, N.E. and Horowitz, J.F. (2011) High muscle lipid content in obesity is not due to enhanced activation of key triglyceride esterification enzymes or the suppression of lipolytic proteins. *Am J Physiol Endocrinol Metab*, 300 (4), E699-707. 10.1152/ajpendo.00316.2010.
- Lindseth, I. (2017) Methodological issues question the validity of observed performance impairment of a low carbohydrate, high fat diet. *J Physiol*, 595 (9), 2989. 10.1113/JP273990.
- Little, J.P., Safdar, A., Wilkin, G.P., Tarnopolsky, M.A. and Gibala, M.J. (2010) A practical model of low-volume high-intensity interval training induces mitochondrial biogenesis in human skeletal muscle: potential mechanisms. *J Physiol*, 588 (Pt 6), 1011-1022. 10.1113/jphysiol.2009.181743.
- Liu, L., Zhang, Y., Chen, N., Shi, X., Tsang, B. and Yu, Y.H. (2007) Upregulation of myocellular DGAT1 augments triglyceride synthesis in skeletal muscle and protects against fat-induced insulin resistance. *J Clin Invest*, 117 (6), 1679-1689. 10.1172/JCI30565.
- Lund, J., Helle, S.A., Li, Y., Lovsletten, N.G., Stadheim, H.K., Jensen, J., Kase, E.T., Thoresen, G.H. and Rustan, A.C. (2018) Higher lipid turnover and oxidation in cultured human myotubes from athletic versus sedentary young male subjects. *Sci Rep*, 8 (1), 17549. 10.1038/s41598-018-35715-7.
- Lundsgaard, A.M., Fritzen, A.M. and Kiens, B. (2018) Molecular Regulation of Fatty Acid Oxidation in Skeletal Muscle during Aerobic Exercise. *Trends Endocrinol Metab*, 29 (1), 18-30. 10.1016/j.tem.2017.10.011.
- MacInnis, M.J. and Gibala, M.J. (2017) Physiological adaptations to interval training and the role of exercise intensity. *J Physiol*, 595 (9), 2915-2930. 10.1113/JP273196.
- Macpherson, R.E., Vandenboom, R., Roy, B.D. and Peters, S.J. (2013) Skeletal muscle PLIN3 and PLIN5 are serine phosphorylated at rest and following lipolysis during adrenergic or contractile stimulation. *Physiol Rep*, 1 (4), e00084. 10.1002/phy2.84.
- Makimura, H., Stanley, T.L., Suresh, C., De Sousa-Coelho, A.L., Frontera, W.R., Syu, S., Braun, L.R., Looby, S.E., Feldpausch, M.N., Torriani, M., Lee, H., Patti, M.E. and Grinspoon, S.K. (2016) Metabolic Effects of Long-Term Reduction in Free Fatty Acids With Acipimox in Obesity: A Randomized Trial. *J Clin Endocrinol Metab*, 101 (3), 1123-1133. 10.1210/jc.2015-3696.
- Menshikova, E.V., Ritov, V.B., Dube, J.J., Amati, F., Stefanovic-Racic, M., Toledo, F.G.S., Coen, P.M. and Goodpaster, B.H. (2017) Calorie Restriction-induced Weight Loss and Exercise Have Differential Effects on Skeletal Muscle Mitochondria Despite

Similar Effects on Insulin Sensitivity. *J Gerontol A Biol Sci Med Sci*, 73 (1), 81-87.  
10.1093/gerona/glw328.

Meyer, M.M., Levin, K., Grimmsmann, T., Beck-Nielsen, H. and Klein, H.H. (2002) Insulin signalling in skeletal muscle of subjects with or without Type II-diabetes and first degree relatives of patients with the disease. *Diabetologia*, 45 (6), 813-822.  
10.1007/s00125-002-0830-9.

Miller, V.J., LaFountain, R.A., Barnhart, E., Sapper, T.S., Short, J., Arnold, W.D., Hyde, P.N., Crabtree, C.D., Kackley, M.L., Kraemer, W.J., Villamena, F.A. and Volek, J.S. (2020) A ketogenic diet combined with exercise alters mitochondrial function in human skeletal muscle while improving metabolic health. *Am J Physiol Endocrinol Metab*, 319 (6), E995-E1007. 10.1152/ajpendo.00305.2020.

Moore, L.L., Visionsi, A.J., Wilson, P.W., D'Agostino, R.B., Finkle, W.D. and Ellison, R.C. (2000) Can sustained weight loss in overweight individuals reduce the risk of diabetes mellitus? *Epidemiology*, 11 (3), 269-273. 10.1097/00001648-200005000-00007.

Morales, P.E., Bucarey, J.L. and Espinosa, A. (2017) Muscle Lipid Metabolism: Role of Lipid Droplets and Perilipins. *J Diabetes Res*, 2017, 1789395.  
10.1155/2017/1789395.

Moro, C., Bajpeyi, S. and Smith, S.R. (2008) Determinants of intramyocellular triglyceride turnover: implications for insulin sensitivity. *Am J Physiol Endocrinol Metab*, 294 (2), E203-213. 10.1152/ajpendo.00624.2007.

Morton, J.P., Croft, L., Bartlett, J.D., Maclaren, D.P., Reilly, T., Evans, L., McArdle, A. and Drust, B. (2009) Reduced carbohydrate availability does not modulate training-induced heat shock protein adaptations but does upregulate oxidative enzyme activity in human skeletal muscle. *J Appl Physiol* (1985), 106 (5), 1513-1521.  
10.1152/japplphysiol.00003.2009.

Nadeau, K.J., Ehlers, L.B., Aguirre, L.E., Moore, R.L., Jew, K.N., Ortmeyer, H.K., Hansen, B.C., Reusch, J.E. and Draznin, B. (2006) Exercise training and calorie restriction increase SREBP-1 expression and intramuscular triglyceride in skeletal muscle. *Am J Physiol Endocrinol Metab*, 291 (1), E90-98.  
10.1152/ajpendo.00543.2005.

Najt, C.P., Khan, S.A., Heden, T.D., Witthuhn, B.A., Perez, M., Heier, J.L., Mead, L.E., Franklin, M.P., Karanja, K.K., Graham, M.J., Mashek, M.T., Bernlohr, D.A., Parker, L., Chow, L.S. and Mashek, D.G. (2020) Lipid Droplet-Derived Monounsaturated Fatty Acids Traffic via PLIN5 to Allosterically Activate SIRT1. *Mol Cell*, 77 (4), 810-824 e818.  
10.1016/j.molcel.2019.12.003.

Newsom, S.A., Schenk, S., Li, M., Everett, A.C. and Horowitz, J.F. (2011) High fatty acid availability after exercise alters the regulation of muscle lipid metabolism. *Metabolism*, 60 (6), 852-859. 10.1016/j.metabol.2010.08.004.

Nielsen, J., Christensen, A.E., Nellemann, B. and Christensen, B. (2017) Lipid droplet size and location in human skeletal muscle fibers are associated with insulin sensitivity. *Am J Physiol Endocrinol Metab*, 313 (6), E721-E730. 10.1152/ajpendo.00062.2017.

Pan, D.A., Lillioja, S., Kriketos, A.D., Milner, M.R., Baur, L.A., Bogardus, C., Jenkins, A.B. and Storlien, L.H. (1997) Skeletal muscle triglyceride levels are inversely related to insulin action. *Diabetes*, 46 (6), 983-988. 10.2337/diab.46.6.983.

Perreault, L., Bergman, B.C., Hunerdosse, D.M., Playdon, M.C. and Eckel, R.H. (2010) Inflexibility in intramuscular triglyceride fractional synthesis distinguishes prediabetes from obesity in humans. *Obesity (Silver Spring)*, 18 (8), 1524-1531. 10.1038/oby.2009.454.

Pesta, D., Hoppel, F., Macek, C., Messner, H., Faulhaber, M., Kobel, C., Parson, W., Bartscher, M., Schocke, M. and Gnaiger, E. (2011) Similar qualitative and quantitative changes of mitochondrial respiration following strength and endurance training in normoxia and hypoxia in sedentary humans. *Am J Physiol Regul Integr Comp Physiol*, 301 (4), R1078-1087. 10.1152/ajpregu.00285.2011.

Peters, S.J., Samjoo, I.A., Devries, M.C., Stevic, I., Robertshaw, H.A. and Tarnopolsky, M.A. (2012) Perilipin family (PLIN) proteins in human skeletal muscle: the effect of sex, obesity, and endurance training. *Appl Physiol Nutr Metab*, 37 (4), 724-735. 10.1139/h2012-059.

Phielix, E., Jelenik, T., Nowotny, P., Szendroedi, J. and Roden, M. (2014) Reduction of non-esterified fatty acids improves insulin sensitivity and lowers oxidative stress, but fails to restore oxidative capacity in type 2 diabetes: a randomised clinical trial. *Diabetologia*, 57 (3), 572-581. 10.1007/s00125-013-3127-2.

Phillips, S.M., Green, H.J., Tarnopolsky, M.A., Heigenhauser, G.F., Hill, R.E. and Grant, S.M. (1996) Effects of training duration on substrate turnover and oxidation during exercise. *J Appl Physiol* (1985), 81 (5), 2182-2191. 10.1152/jappl.1996.81.5.2182.

Philp, A., MacKenzie, M.G., Belew, M.Y., Towler, M.C., Corstorphine, A., Papalamprou, A., Hardie, D.G. and Baar, K. (2013) Glycogen content regulates peroxisome proliferator activated receptor- partial differential (PPAR- partial differential) activity in rat skeletal muscle. *Plos One*, 8 (10), e77200. 10.1371/journal.pone.0077200.

Pourteymour, S., Lee, S., Langleite, T.M., Eckardt, K., Hjorth, M., Bindesboll, C., Dalen, K.T., Birkeland, K.I., Dreven, C.A., Holen, T. and Norheim, F. (2015) Perilipin 4 in human skeletal muscle: localization and effect of physical activity. *Physiol Rep*, 3 (8). 10.14814/phy2.12481.

Pruchnic, R., Katsiaras, A., He, J., Kelley, D.E., Winters, C. and Goodpaster, B.H. (2004) Exercise training increases intramyocellular lipid and oxidative capacity in older adults. *Am J Physiol Endocrinol Metab*, 287 (5), E857-862. 10.1152/ajpendo.00459.2003.

Rabøl, R., Svendsen, P.F., Skovbro, M., Boushel, R., Haugaard, S.B., Schjerling, P., Schrauwen, P., Hesselink, M.K., Nilas, L., Madsbad, S. and Dela, F. (2009) Reduced skeletal muscle mitochondrial respiration and improved glucose metabolism in nondiabetic obese women during a very low calorie dietary intervention leading to rapid weight loss. *Metabolism*, 58 (8), 1145-1152. 10.1016/j.metabol.2009.03.014.

Ramos, P.A., Lytle, K.A., Delivanis, D., Nielsen, S., LeBrasseur, N.K. and Jensen, M.D. (2021) Insulin-Stimulated Muscle Glucose Uptake and Insulin Signaling in Lean and Obese Humans. *J Clin Endocrinol Metab*, 106 (4), e1631-e1646. 10.1210/clinem/dgaa919.

Romijn, J.A., Coyle, E.F., Sidossis, L.S., Gastaldelli, A., Horowitz, J.F., Endert, E. and Wolfe, R.R. (1993) Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am J Physiol*, 265 (3 Pt 1), E380-391. 10.1152/ajpendo.1993.265.3.E380.

Romijn, J.A., Coyle, E.F., Sidossis, L.S., Zhang, X.J. and Wolfe, R.R. (1995) Relationship between fatty acid delivery and fatty acid oxidation during strenuous exercise. *J Appl Physiol* (1985), 79 (6), 1939-1945. 10.1152/jappl.1995.79.6.1939.

Ryan, B.J., Schleh, M.W., Ahn, C., Ludzki, A.C., Gillen, J.B., Varshney, P., Van Pelt, D.W., Pitchford, L.M., Chenevert, T.L., Gioscia-Ryan, R.A., Howton, S.M., Rode, T., Hummel, S.L., Burant, C.F., Little, J.P. and Horowitz, J.F. (2020) Moderate-Intensity Exercise and High-Intensity Interval Training Affect Insulin Sensitivity Similarly in Obese Adults. *J Clin Endocrinol Metab*, 105 (8). 10.1210/clinem/dgaa345.

Sacchetti, M., Saltin, B., Olsen, D.B. and van Hall, G. (2004) High triacylglycerol turnover rate in human skeletal muscle. *J Physiol*, 561 (Pt 3), 883-891. 10.1113/jphysiol.2004.075135.

Saloranta, C., Taskinen, M.R., Widen, E., Harkonen, M., Melander, A. and Groop, L. (1993) Metabolic consequences of sustained suppression of free fatty acids by acipimox in patients with NIDDM. *Diabetes*, 42 (11), 1559-1566. 10.2337/diab.42.11.1559.

Schenk, S. and Horowitz, J.F. (2007) Acute exercise increases triglyceride synthesis in skeletal muscle and prevents fatty acid-induced insulin resistance. *J Clin Invest*, 117 (6), 1690-1698. 10.1172/JCI30566.

Schrauwen, P., van Aggel-Leijssen, D.P., Hul, G., Wagenmakers, A.J., Vidal, H., Saris, W.H. and van Baak, M.A. (2002) The effect of a 3-month low-intensity endurance training program on fat oxidation and acetyl-CoA carboxylase-2 expression. *Diabetes*, 51 (7), 2220-2226. 10.2337/diabetes.51.7.2220.

Schwenk, R.W., Holloway, G.P., Luiken, J.J., Bonen, A. and Glatz, J.F. (2010) Fatty acid transport across the cell membrane: regulation by fatty acid transporters. *Prostaglandins Leukot Essent Fatty Acids*, 82 (4-6), 149-154. 10.1016/j.plefa.2010.02.029.

- Scribbans, T.D., Edgett, B.A., Vorobej, K., Mitchell, A.S., Joannis, S.D., Matusiak, J.B., Parise, G., Quadrilatero, J. and Gurd, B.J. (2014) Fibre-specific responses to endurance and low volume high intensity interval training: striking similarities in acute and chronic adaptation. *Plos One*, 9 (6), e98119. 10.1371/journal.pone.0098119.
- Shaw, C.S., Jones, D.A. and Wagenmakers, A.J. (2008) Network distribution of mitochondria and lipid droplets in human muscle fibres. *Histochem Cell Biol*, 129 (1), 65-72. 10.1007/s00418-007-0349-8.
- Shaw, C.S., Shepherd, S.O., Wagenmakers, A.J., Hansen, D., Dendale, P. and van Loon, L.J. (2012) Prolonged exercise training increases intramuscular lipid content and perilipin 2 expression in type I muscle fibers of patients with type 2 diabetes. *Am J Physiol Endocrinol Metab*, 303 (9), E1158-1165. 10.1152/ajpendo.00272.2012.
- Shaw, C.S., Sherlock, M., Stewart, P.M. and Wagenmakers, A.J. (2009) Adipophilin distribution and colocalization with lipid droplets in skeletal muscle. *Histochem Cell Biol*, 131 (5), 575-581. 10.1007/s00418-009-0558-4.
- Shepherd, S.O., Cocks, M., Meikle, P.J., Mellett, N.A., Ranasinghe, A.M., Barker, T.A., Wagenmakers, A.J.M. and Shaw, C.S. (2017a) Lipid droplet remodelling and reduced muscle ceramides following sprint interval and moderate-intensity continuous exercise training in obese males. *Int J Obes (Lond)*, 41 (12), 1745-1754. 10.1038/ijo.2017.170.
- Shepherd, S.O., Cocks, M., Tipton, K.D., Ranasinghe, A.M., Barker, T.A., Burniston, J.G., Wagenmakers, A.J. and Shaw, C.S. (2012) Preferential utilization of perilipin 2-associated intramuscular triglycerides during 1 h of moderate-intensity endurance-type exercise. *Exp Physiol*, 97 (8), 970-980. 10.1113/expphysiol.2012.064592.
- Shepherd, S.O., Cocks, M., Tipton, K.D., Ranasinghe, A.M., Barker, T.A., Burniston, J.G., Wagenmakers, A.J. and Shaw, C.S. (2013) Sprint interval and traditional endurance training increase net intramuscular triglyceride breakdown and expression of perilipin 2 and 5. *J Physiol*, 591 (3), 657-675. 10.1113/jphysiol.2012.240952.
- Shepherd, S.O., Cocks, M., Tipton, K.D., Witard, O.C., Ranasinghe, A.M., Barker, T.A., Wagenmakers, A.J. and Shaw, C.S. (2014) Resistance training increases skeletal muscle oxidative capacity and net intramuscular triglyceride breakdown in type I and II fibres of sedentary males. *Exp Physiol*, 99 (6), 894-908. 10.1113/expphysiol.2014.078014.
- Shepherd, S.O., Strauss, J.A., Wang, Q., Dube, J.J., Goodpaster, B., Mashek, D.G. and Chow, L.S. (2017b) Training alters the distribution of perilipin proteins in muscle following acute free fatty acid exposure. *J Physiol*, 595 (16), 5587-5601. 10.1113/JP274374.
- Sparks, L.M., Redman, L.M., Conley, K.E., Harper, M.E., Yi, F., Hodges, A., Eroshkin, A., Costford, S.R., Gabriel, M.E., Shook, C., Cornnell, H.H., Ravussin, E. and Smith, S.R. (2017) Effects of 12 Months of Caloric Restriction on Muscle Mitochondrial Function in Healthy Individuals. *J Clin Endocrinol Metab*, 102 (1), 111-121. 10.1210/jc.2016-3211.

- Starling, R.D., Trappe, T.A., Parcell, A.C., Kerr, C.G., Fink, W.J. and Costill, D.L. (1997) Effects of diet on muscle triglyceride and endurance performance. *J Appl Physiol* (1985), 82 (4), 1185-1189. 10.1152/jappl.1997.82.4.1185.
- Stellingwerff, T., Boon, H., Gijsen, A.P., Stegen, J.H., Kuipers, H. and van Loon, L.J. (2007) Carbohydrate supplementation during prolonged cycling exercise spares muscle glycogen but does not affect intramyocellular lipid use. *Pflugers Arch*, 454 (4), 635-647. 10.1007/s00424-007-0236-0.
- Stellingwerff, T., Spriet, L.L., Watt, M.J., Kimber, N.E., Hargreaves, M., Hawley, J.A. and Burke, L.M. (2006) Decreased PDH activation and glycogenolysis during exercise following fat adaptation with carbohydrate restoration. *Am J Physiol Endocrinol Metab*, 290 (2), E380-388. 10.1152/ajpendo.00268.2005.
- Stuyt, P.M., Kleinjans, H.A. and Stalenhoef, A.F. (1998) Tolerability and effects of high doses acipimox as additional lipid-lowering therapy in familial hypercholesterolemia. *Neth J Med*, 53 (5), 228-233. 10.1016/s0300-2977(98)00076-x.
- Summers, S.A. and Nelson, D.H. (2005) A role for sphingolipids in producing the common features of type 2 diabetes, metabolic syndrome X, and Cushing's syndrome. *Diabetes*, 54 (3), 591-602. 10.2337/diabetes.54.3.591.
- Talanian, J.L., Holloway, G.P., Snook, L.A., Heigenhauser, G.J., Bonen, A. and Spriet, L.L. (2010) Exercise training increases sarcolemmal and mitochondrial fatty acid transport proteins in human skeletal muscle. *Am J Physiol Endocrinol Metab*, 299 (2), E180-188. 10.1152/ajpendo.00073.2010.
- Tanaka, T., Yamamoto, J., Iwasaki, S., Asaba, H., Hamura, H., Ikeda, Y., Watanabe, M., Magoori, K., Ioka, R.X., Tachibana, K., Watanabe, Y., Uchiyama, Y., Sumi, K., Iguchi, H., Ito, S., Doi, T., Hamakubo, T., Naito, M., Auwerx, J., Yanagisawa, M., Kodama, T. and Sakai, J. (2003) Activation of peroxisome proliferator-activated receptor delta induces fatty acid beta-oxidation in skeletal muscle and attenuates metabolic syndrome. *Proc Natl Acad Sci U S A*, 100 (26), 15924-15929. 10.1073/pnas.0306981100.
- Tang, J.E., Hartman, J.W. and Phillips, S.M. (2006) Increased muscle oxidative potential following resistance training induced fibre hypertrophy in young men. *Appl Physiol Nutr Metab*, 31 (5), 495-501. 10.1139/h06-026.
- Tarnopolsky, M.A., Rennie, C.D., Robertshaw, H.A., Fedak-Tarnopolsky, S.N., Devries, M.C. and Hamadeh, M.J. (2007) Influence of endurance exercise training and sex on intramyocellular lipid and mitochondrial ultrastructure, substrate use, and mitochondrial enzyme activity. *Am J Physiol Regul Integr Comp Physiol*, 292 (3), R1271-1278. 10.1152/ajpregu.00472.2006.
- Teodoro, B.G., Sampaio, I.H., Bomfim, L.H., Queiroz, A.L., Silveira, L.R., Souza, A.O., Fernandes, A.M., Eberlin, M.N., Huang, T.Y., Zheng, D., Neufer, P.D., Cortright, R.N. and Alberici, L.C. (2017) Long-chain acyl-CoA synthetase 6 regulates lipid synthesis and mitochondrial oxidative capacity in human and rat skeletal muscle. *J Physiol*, 595 (3), 677-693. 10.1113/JP272962.



1582

1583 Thrush, A.B., Brindley, D.N., Chabowski, A., Heigenhauser, G.J. and Dyck, D.J.  
1584 (2009) Skeletal muscle lipogenic protein expression is not different between lean and  
1585 obese individuals: a potential factor in ceramide accumulation. *J Clin Endocrinol*  
1586 *Metab*, 94 (12), 5053-5061. 10.1210/jc.2008-2565.

1587

1588 Toledo, F.G., Menshikova, E.V., Azuma, K., Radikova, Z., Kelley, C.A., Ritov, V.B. and  
1589 Kelley, D.E. (2008) Mitochondrial capacity in skeletal muscle is not stimulated by  
1590 weight loss despite increases in insulin action and decreases in intramyocellular lipid  
1591 content. *Diabetes*, 57 (4), 987-994. 10.2337/db07-1429.

1592

1593 Vaag, A.A. and Beck-Nielsen, H. (1992) Effects of prolonged Acipimox treatment on  
1594 glucose and lipid metabolism and on in vivo insulin sensitivity in patients with non-  
1595 insulin dependent diabetes mellitus. *Acta Endocrinol (Copenh)*, 127 (4), 344-350.  
1596 10.1530/acta.0.1270344.

1597

1598 van de Weijer, T., Phielix, E., Bilet, L., Williams, E.G., Ropelle, E.R., Bierwagen, A.,  
1599 Livingstone, R., Nowotny, P., Sparks, L.M., Pagliarunga, S., Szendroedi, J., Havekes,  
1600 B., Moullan, N., Pirinen, E., Hwang, J.H., Schrauwen-Hinderling, V.B., Hesselink,  
1601 M.K., Auwerx, J., Roden, M. and Schrauwen, P. (2015) Evidence for a direct effect of  
1602 the NAD<sup>+</sup> precursor acipimox on muscle mitochondrial function in humans. *Diabetes*,  
1603 64 (4), 1193-1201. 10.2337/db14-0667.

1604

1605 van Loon, L.J. (2004) Use of intramuscular triacylglycerol as a substrate source during  
1606 exercise in humans. *J Appl Physiol* (1985), 97 (4), 1170-1187.  
1607 10.1152/japplphysiol.00368.2004.

1608

1609 van Loon, L.J., Greenhaff, P.L., Constantin-Teodosiu, D., Saris, W.H. and  
1610 Wagenmakers, A.J. (2001) The effects of increasing exercise intensity on muscle fuel  
1611 utilisation in humans. *J Physiol*, 536 (Pt 1), 295-304. 10.1111/j.1469-  
1612 7793.2001.00295.x.

1613

1614 van Loon, L.J., Koopman, R., Manders, R., van der Weegen, W., van Kranenburg,  
1615 G.P. and Keizer, H.A. (2004) Intramyocellular lipid content in type 2 diabetes patients  
1616 compared with overweight sedentary men and highly trained endurance athletes. *Am*  
1617 *J Physiol Endocrinol Metab*, 287 (3), E558-565. 10.1152/ajpendo.00464.2003.

1618

1619 van Loon, L.J., Koopman, R., Stegen, J.H., Wagenmakers, A.J., Keizer, H.A. and  
1620 Saris, W.H. (2003a) Intramyocellular lipids form an important substrate source during  
1621 moderate intensity exercise in endurance-trained males in a fasted state. *J Physiol*,  
1622 553 (Pt 2), 611-625. 10.1113/jphysiol.2003.052431.

1623

1624 van Loon, L.J., Manders, R.J., Koopman, R., Kaastra, B., Stegen, J.H., Gijsen, A.P.,  
1625 Saris, W.H. and Keizer, H.A. (2005a) Inhibition of adipose tissue lipolysis increases  
1626 intramuscular lipid use in type 2 diabetic patients. *Diabetologia*, 48 (10), 2097-2107.  
1627 10.1007/s00125-005-1889-x.

1628

1629 van Loon, L.J., Schrauwen-Hinderling, V.B., Koopman, R., Wagenmakers, A.J.,  
1630 Hesselink, M.K., Schaart, G., Kooi, M.E. and Saris, W.H. (2003b) Influence of  
1631 prolonged endurance cycling and recovery diet on intramuscular triglyceride content

in trained males. *Am J Physiol Endocrinol Metab*, 285 (4), E804-811. 10.1152/ajpendo.00112.2003.

van Loon, L.J., Thomason-Hughes, M., Constantin-Teodosiu, D., Koopman, R., Greenhaff, P.L., Hardie, D.G., Keizer, H.A., Saris, W.H. and Wagenmakers, A.J. (2005b) Inhibition of adipose tissue lipolysis increases intramuscular lipid and glycogen use in vivo in humans. *Am J Physiol Endocrinol Metab*, 289 (3), E482-493. 10.1152/ajpendo.00092.2005.

Van Proeyen, K., Szlufcik, K., Nielens, H., Deldicque, L., Van Dyck, R., Ramaekers, M. and Hespel, P. (2011a) High-fat diet overrules the effects of training on fiber-specific intramyocellular lipid utilization during exercise. *J Appl Physiol* (1985), 111 (1), 108-116. 10.1152/japplphysiol.01459.2010.

Van Proeyen, K., Szlufcik, K., Nielens, H., Ramaekers, M. and Hespel, P. (2011b) Beneficial metabolic adaptations due to endurance exercise training in the fasted state. *J Appl Physiol* (1985), 110 (1), 236-245. 10.1152/japplphysiol.00907.2010.

Verboven, K., Wens, I., Vandenabeele, F., Stevens, A.N., Celie, B., Lapauw, B., Dendale, P., LJC, V.A.N.L., Calders, P. and Hansen, D. (2020) Impact of Exercise-Nutritional State Interactions in Patients with Type 2 Diabetes. *Med Sci Sports Exerc*, 52 (3), 720-728. 10.1249/MSS.0000000000002165.

Wang, H., Sreenivasan, U., Hu, H., Saladino, A., Polster, B.M., Lund, L.M., Gong, D.W., Stanley, W.C. and Sztalryd, C. (2011) Perilipin 5, a lipid droplet-associated protein, provides physical and metabolic linkage to mitochondria. *J Lipid Res*, 52 (12), 2159-2168. 10.1194/jlr.M017939.

Watt, M.J., Southgate, R.J., Holmes, A.G. and Febbraio, M.A. (2004) Suppression of plasma free fatty acids upregulates peroxisome proliferator-activated receptor (PPAR) alpha and delta and PPAR coactivator 1alpha in human skeletal muscle, but not lipid regulatory genes. *J Mol Endocrinol*, 33 (2), 533-544. 10.1677/jme.1.01499.

Whytock, K.L., Parry, S.A., Turner, M.C., Woods, R.M., James, L.J., Ferguson, R.A., Stahlman, M., Boren, J., Strauss, J.A., Cocks, M., Wagenmakers, A.J.M., Hulston, C.J. and Shepherd, S.O. (2020) A 7-day high-fat, high-calorie diet induces fibre-specific increases in intramuscular triglyceride and perilipin protein expression in human skeletal muscle. *J Physiol*, 598 (6), 1151-1167. 10.1113/JP279129.

Whytock, K.L., Shepherd, S.O., Wagenmakers, A.J.M. and Strauss, J.A. (2018) Hormone-sensitive lipase preferentially redistributes to lipid droplets associated with perilipin-5 in human skeletal muscle during moderate-intensity exercise. *J Physiol*, 596 (11), 2077-2090. 10.1113/JP275502.

Wojtaszewski, J.F., MacDonald, C., Nielsen, J.N., Hellsten, Y., Hardie, D.G., Kemp, B.E., Kiens, B. and Richter, E.A. (2003) Regulation of 5'AMP-activated protein kinase activity and substrate utilization in exercising human skeletal muscle. *Am J Physiol Endocrinol Metab*, 284 (4), E813-822. 10.1152/ajpendo.00436.2002.

Xu, G., Sztalryd, C., Lu, X., Tansey, J.T., Gan, J., Dorward, H., Kimmel, A.R. and Londos, C. (2005) Post-translational regulation of adipose differentiation-related protein by the ubiquitin/proteasome pathway. *J Biol Chem*, 280 (52), 42841-42847. 10.1074/jbc.M506569200.

Yeo, W.K., Lessard, S.J., Chen, Z.P., Garnham, A.P., Burke, L.M., Rivas, D.A., Kemp, B.E. and Hawley, J.A. (2008a) Fat adaptation followed by carbohydrate restoration increases AMPK activity in skeletal muscle from trained humans. *J Appl Physiol* (1985), 105 (5), 1519-1526. 10.1152/jappphysiol.90540.2008.

Yeo, W.K., Paton, C.D., Garnham, A.P., Burke, L.M., Carey, A.L. and Hawley, J.A. (2008b) Skeletal muscle adaptation and performance responses to once a day versus twice every second day endurance training regimens. *J Appl Physiol* (1985), 105 (5), 1462-1470. 10.1152/jappphysiol.90882.2008.

Yu, C., Chen, Y., Cline, G.W., Zhang, D., Zong, H., Wang, Y., Bergeron, R., Kim, J.K., Cushman, S.W., Cooney, G.J., Atcheson, B., White, M.F., Kraegen, E.W. and Shulman, G.I. (2002) Mechanism by which fatty acids inhibit insulin activation of insulin receptor substrate-1 (IRS-1)-associated phosphatidylinositol 3-kinase activity in muscle. *J Biol Chem*, 277 (52), 50230-50236. 10.1074/jbc.M200958200.

Zhu, J., Xu, M., Liu, Y., Zhuang, L., Ying, K., Liu, F., Liu, D., Ma, W. and Songyang, Z. (2019) Phosphorylation of PLIN3 by AMPK promotes dispersion of lipid droplets during starvation. *Protein Cell*, 10 (5), 382-387. 10.1007/s13238-018-0593-9.


**Figure 1. A representation of the subcellular location of LD in skeletal muscle in insulin-resistant, obese individuals and insulin-sensitive, trained individuals.**

Insulin-sensitive, trained individuals typically exhibit a large number of small LD which are primarily located in the intermyofibrillar region of type I fibres. Moreover, the number of LD is two-to-three-fold greater in type I compared to type II fibres in trained individuals. In contrast, insulin-resistant, obese individuals tend to exhibit LD that are much large in size compared to insulin-sensitive, trained individuals. Furthermore, in insulin-resistant individuals a higher proportion of LD appear to be present in the subsarcolemmal region compared to trained individuals. This is especially true in type II fibres, where the number and size of SS LD is ~two-fold greater compared to trained individuals (Daemen et al., 2018). Thus, the muscle of trained individuals is characterised by a large number of small LD located in the intermyofibrillar region of type I fibres, whereas the muscle of insulin-resistant obese individuals is characterised by large LDs stored in the subsarcolemmal region of type II fibres.


**Figure 2. Schematic overview of the distribution of LD and mitochondria in skeletal muscle in insulin-resistant, obese individuals and insulin-sensitive, trained individuals.**

Endurance, resistance, or high intensity interval exercise training promotes a shift towards a more structured network of LD and mitochondria; that is, more LD are labelled with PLIN proteins (PLIN2, 3, 4 or 5, but indicated as one by the small orange ellipse) and are located in close proximity to the mitochondria. This collective adaptation to exercise training enables more efficient breakdown and oxidation of IMTG stored in LD. This is important in order to create a net decrease in IMTG following exercise, which subsequently increases the capacity for IMTG synthesis and storage in the hours following exercise. Thus, following training the overall turnover rate of the IMTG pool is enhanced, which reduces the risk of generating and accumulating toxic lipid metabolites that would otherwise contribute to the development of insulin resistance. In obesity and T2D there is a poor capacity to reduce the rate of IMTG synthesis during exercise and therefore generate a net decrease in IMTG content post-exercise. Additional strategies to exercise alone may therefore be required to create a post-exercise decrease in IMTG content in obese individuals. We propose several co-strategies (anti-lipolytic drug therapy, fasted exercise, training with low muscle glycogen, a LCHF diet or WL/CR) that target an improved rate of IMTG turnover, and therefore maximize the insulin-sensitising effects of exercise training alone in obese individuals. Future work is now required to test these interventions in the longer term in clinical populations, with consideration for how these interventions can be successfully adopted in the real world.

Insulin-sensitive, trained individual





Insulin-resistant, obese individual



Type I fibers


Type II fibers



IMF

SS

Insulin-resistant, obese individual



Exercise Training

+ Anti-lipolytic drug therapy  
+ Fasted state  
+ Low glycogen state  
+ LCHF diet  
+ WL/CR

?

Insulin-sensitive, trained individual

