

**Should I stay or should I go?**  
**How activity synchronization affects fission decisions**

**Laura Busia<sup>1</sup>, Colleen M. Schaffner<sup>2</sup>, Filippo Aureli<sup>3 4</sup>**

<sup>1</sup> Independent researcher

<sup>2</sup> Psychology Department, Adams State University, Alamosa, CO, USA

<sup>3</sup> Instituto de Neuroetologia, Universidad Veracruzana, Xalapa, Veracruz, Mexico

<sup>4</sup> Research Centre in Evolutionary Anthropology and Palaeoecology, Liverpool John Moores University, Liverpool, UK

**Abstract (max 200 words)**

Group-living animals need to deal with conflicting interests to maintain cohesion. When the costs of doing so outweigh the benefits, the group may (temporarily) split into two or more subgroups. Conflicting interests can concern the activity to pursue or the direction of travel. Temporary group separation is a common feature in species with a high degree of fission-fusion dynamics. We investigated the role activity synchronization played in fission decisions in a spider monkey group living in the Otoh Ma'ax Yetel Kooch Nature Reserve, Yucatan, Mexico. For 21 months, we recorded every fission event occurring in the followed subgroup, as well as the subgroup activity. We classified the activity as “synchronized” when at least the 75% of subgroup members performed the same activity (resting, foraging, socializing or travelling); otherwise, we classified it as “non-synchronized”. We found that fission events occurred more often when the activity was non-synchronized. In addition, when the activity was synchronized, fission events occurred more often when spider monkeys were travelling than when they engaged in other subgroup activities. Our findings highlight the role of conflicting interests over the activity to pursue and travel direction on fission decisions.

**Key words:** activity synchronization; fission-fusion dynamics; decision making process; Ateles.

**Introduction**

Most studies on decision-making processes in group-living animals focus on collective

decisions, i.e. when a group of individuals coordinate their behaviour [1], to maintain the benefits of group living (e.g. safety from predation: [2,3]). Collective decisions are usually studied when individuals try to reach a consensus on when to move [4,5], where  
40 to move and where to eat [6-10].

To stay in the same group, members need to compromise their different needs and synchronize activities. If the cost of synchronization outweighs the benefits, the lack of consensus may result in a (temporary) break-up of the group [11-13]. For example, red deer (*Cervus elaphus*) are rarely in mixed-sex groups outside the breeding  
45 season probably due to the difference in the length of the foraging-resting bouts between males and females [14,15]. Furthermore, consensus needs to be reached not only for the activity, but also for the travel direction. The lack of consensus over travel direction may result in a leader change during the travelling bout [16], a delay in the decision-making process [17], or the temporary break-up of the group [9,18,19].

In several species temporary formation of subgroups occurs several times a day. This is the case for species with a high degree of fission-fusion dynamics [20], in which subgroups frequently change their size and composition [21]. The decision to fission into smaller subgroups is a trade-off between the need to reduce feeding competition [22-26] and the need to decrease the risk of predation [25,27,28]. However, other  
50 ecological and social factors may affect fission decisions, such as food characteristics [29], the quality of social relationships [30-32], the location within the home range [33], and the degree of human disturbance [34]. Here we investigate whether the degree of synchronization of subgroup members' activity could affect individual fission decisions in wild Geoffroy's spider monkeys (*Ateles geoffroyi*), a species with a high degree of  
55 fission-fusion dynamics [35].  
60

We focus on fission decisions related to conflicting interests over the activity to pursue and travel direction. We have two predictions. First, when the degree of activity synchronization among subgroup members is lower, we expect fission to be more likely (Prediction 1). Individuals may also face conflicting interests over travel direction based  
65 on differences in the location to be reached. Thus, our second prediction is when the activity is highly synchronized, we expect a higher likelihood of fission when subgroup members travel rather than when they engage in other activities (Prediction 2).

## Methods

### 70 *Field site and study subjects*

The field site is near the Punta Laguna lake within the natural protected area of Otoch Ma'ax Yetel Kooch, Yucatan Peninsula, Mexico (20°38' N, 87°38' W, [36]). We studied 22 adult and subadult individuals of a group of Geoffroy's spider monkeys living in the protected area (6 adult males, 10 adult females, 1 subadult male, 5 subadult females).

75 The study group has been the focus of a continuous long-term project since 1997 [37]. All monkeys are fully habituated to human observers and individually recognized by facial features, scars and fur coloration.

#### *Data collection*

80 Data were collected from January 2013 to September 2014. While following a subgroup, the first author continuously recorded any change in each subgroup member's activity in a digital voice recorder with the help of field assistants spread in the area occupied by the subgroup, so that every subgroup member could be monitored (see Supplementary information for a detailed description of data collection). We considered  
85 the following four activities: resting (individual in a stationary position either lying, sitting or hanging), foraging (individual actively searching, manipulating and/or ingesting food items), social (individual involved in social interactions, e.g., grooming) and travelling (individual moving together with other subgroup members in the same direction).

90 Subgroup membership was continuously updated as we recorded the identity of every member of the initially encountered subgroup and all changes due to fission and fusion events. An individual was considered part of the followed subgroup if it was <30 m from a subgroup member according to a chain rule established for this study site ([38], see [39] for the concept of the chain rule). Fission was defined as individuals  
95 from the followed subgroup separating from one another into different subgroups and was recorded when one or more individuals were not seen within 30 m from any member of the followed subgroup for 30 min [40].

#### *Data analyses*

100 To test Prediction 1, we considered the subgroup activity as synchronized when at least 75% of subgroup members performed the same activity [41]. Otherwise, we considered it non-synchronized. We tested Prediction 1 in two ways. The first consisted of considering the number of fission events occurring during a given day, depending on whether the subgroup activity was synchronized or non-synchronized. We used a

generalized linear mixed model (GLMM) in which the dependent variable was the number of fission events occurring when the activity was synchronized or non-synchronized during a given day. Activity duration was entered with the “offset” function. Activity synchronization (synchronized or non-synchronized) was the predictor variable, whereas the maximum subgroup size recorded during the duration of the activity was entered as a control variable, as the number of individuals in the subgroup may affect the likelihood of activity synchronization [42], and thus fission events [43]. We included the month in which the data were collected as a random factor to control for multiple data points during the same month.

The second way we tested Prediction 1 was to consider the number of fission events depending on the time spent in synchronized activity during each observation hour. We used a GLMM in which the dependent variable was the number of fission events occurring in an hour and the time spent in a synchronized activity during that hour as the predictor variable. We also included the maximum subgroup size recorded during the observation hour and the subgroup type (only males, only females or mixed-sex subgroup) as control variables because they may affect the likelihood of fission events, as individuals are more often in same-sex subgroups [44]. Given the frequent variation in subgroup composition, we could include the subgroup type as control variable only in this model because of the shorter temporal scale (i.e. 1 hour in this model instead of 1 day in the previous model). When subgroup composition type changed during an observation hour, the subgroup composition type occurring for most of the time was selected (we excluded the few cases in which more than one subgroup composition type occurred for most of the time). We included the day in which the data were collected as a random factor to control for multiple data points during the same day.

To test Prediction 2, only periods with synchronized activities were considered. We used a GLMM in which the dependent variable was the number of fission events that occurred during the same activity type in each observation day and activity duration was entered with the “offset” function. The type of activity (travelling or non-travelling) was the predictor variable, the maximum subgroup size recorded during the duration of the activity was entered as a control variable, and the month was entered as a random factor (Data available: see [45])

We ran all the GLMMs using the lme4 package [46] in R (version 3.6.0, R Core Team 2019). We compared full models with null models, which included only the

random factor and the control variables, using a likelihood ratio test with the function  
 140 anova [47]. We checked model assumptions using the “performance” package [48].

## Results

Prediction 1 that fission would be more likely when the activity of subgroup members  
 was not synchronized was supported. The rate of fission events was higher when the  
 145 subgroup activity was non-synchronized compared to when it was synchronized at the  
 temporal unit of observation day (Table 1, Fig 1). A similar result was found at the  
 temporal unit of observation hour (Table 2).

**Table 1:** Results of the GLMM showing the association between the number of fission events (dependent  
 150 variable) and whether the subgroup activity was synchronized or non-synchronized throughout the  
 observation day.

Fixed effects	Estimate	Std. Error	z value	p
(Intercept)	-4.74479	0.15885	-29.869	<0.0001
Activity synchronization	-0.55263	0.11839	-4.668	<0.0001
Maximum subgroup size	0.01427	0.01569	0.909	0.363

*The model was significantly different from the null model (likelihood ratio test:  $N = 416$ ;  $\chi^2 = 73.056$ ;  $p < 0.0001$ ). The negative estimate of activity synchronization means that there were fewer fission events  
 155 when the activity was synchronized than when the activity was non-synchronized.*

**Table 2:** Results of the GLMM showing the association between the number of fission events (dependent  
 variable) and the time spent in synchronized subgroup activity in each observation hour.

Fixed effects	Estimate	Std. Error	z value	p
(Intercept)	-0.669354	0.217579	-3.076	0.002
Time in synchronized activity	-0.009286	0.003516	-2.641	0.008
Maximum subgroup size	0.017826	0.020084	0.888	0.375
Subgroup type (only males)	-0.126567	0.204421	-0.619	0.536
Subgroup type (mixed)	0.318309	0.151351	2.103	0.036

*The model was significantly different from the null model (likelihood ratio test:  $N = 682$ ;  $\chi^2 = 6.613$ ;  $p < 0.010$ ). The categories of Subgroup type “only males” and “mixed” were compared with “only  
 160 females”.*

(FIG 1 here)

Prediction 2 that fission would be more likely when the subgroup activity was travelling  
 165 compared to non-travelling was also supported (Table 3, Fig 2).

**Table 3:** Results of the GLMM showing the association between the number of fission events (dependent variable) and the time of travelling throughout the day.

Fixed effects	Estimate	Std. Error	z value	p
(Intercept)	-5.12511	0.19544	-26.223	< 2e-16
Activity type	-0.38841	0.14141	-2.747	0.0060
Maximum subgroup size	0.02466	0.01938	1.272	0.2033

*The model was significantly different from the null model (likelihood ratio test:  $N = 432$ ;  $\chi^2 = 55.994$ ;  $p < 0.0001$ ). The negative estimate of Activity type means that there were fewer fission events when the activity was non-travelling than when the activity was travelling.*

(Fig 2 here)

## Discussion

Overall, we found that subgroup activity played an important role in influencing fission decisions in wild spider monkeys. We found a higher fission rate when the subgroup activity was non-synchronized compared to when it was synchronized (Prediction 1).

When activity was synchronized, we found a higher fission rate when the subgroup was travelling compared to when subgroup members engaged in other activities (Prediction 2).

We found that the fission rate was higher when the subgroup activity was not synchronized than when it was synchronized at two temporal scales: several hours up to the entire observation day and each observation hour, supporting Prediction 1, although a note of caution is warranted due to the effect of subgroup size may have on activity synchronization [42]. Whereas most studies on activity synchronization focus on the need for individuals to forage and subsequently travel or fission depending on group cohesion level [13,14], our findings suggest that the lack of activity synchronization is sufficient to promote subgroup fission. As individuals of species with a high degree of fission-fusion dynamics do not need to reach a consensus on the activity to pursue [11,13], the difference in the activities performed (e.g., an individual already foraged and needs to rest) may promote fissioning from other subgroup members.

Another possible complementary explanation is related to the quality of social relationships between group members, such as affiliation and compatibility [49]. The relation between synchrony and affiliation is evident in humans [50] and other animals [51-53]. Spider monkeys' tendency to associate in subgroups with the most compatible group members [34] likely results in more shared activities. Thus, non-synchronized

subgroup activity may be a predictor of the presence of subgroup members with social  
200 relationship characterized by low compatibility, which may lead to fission.

Prediction 2 was also supported. Our findings showed that fission rates were  
higher when the subgroup activity was travelling than when subgroup members were  
engaged in other activities. Given that we compared the likelihood of subgroup fission  
during travelling bouts with the likelihood of subgroup fission during other  
205 synchronized activities, we interpret this result as the outcome of conflicting interests  
over the travel direction rather than over the next activity to pursue, which may occur  
during any of the synchronized activities. In cases of conflict over travel direction,  
individuals of some species adopt strategies to maintain group cohesion [16,17],  
whereas others fission into subgroups [9,18,19]. Thus, it is likely that spider monkeys  
210 adopt the latter strategy. Conflict over travel direction is likely due to three reasons:  
differing nutritional needs [29]; engaging in different activities (e.g., feeding vs  
sleeping); and, males and females experience different travel costs [54] and therefore  
may not travel as far as males [55].

Overall, our study contributes to understanding fission decisions, highlighting  
215 the important role played by activity synchronization in species with a high degree of  
fission-fusion dynamics. Staying together when group members experience conflict  
over activity to pursue and travel direction is not the only option, as individuals can  
temporarily fission from other group members. Our findings support this view,  
providing further evidence about fission being a mechanism for conflict management  
220 [56].

### **Acknowledgements**

We thank Anthony R. Denice for his outstanding contribution in data collection,  
Augusto Canul, Eulogio Canul, Juan Canul and Macedonio Canul for their valuable  
225 assistance during fieldwork and Sandra Smith for her overall support. We are grateful to  
Gabriel Ramos-Fernández and Laura Vick for sharing the management of the long-term  
project. We are also indebted to Chester Zoo and The National Geographic Society for  
financially supporting the long-term project. We thank the Consejo Nacional por la  
Ciencia y la Tecnología (CONACyT) for LB's PhD studentship (CVU n° 490429) and  
230 for equipment (n°I0101/152/2014 C-133/2014). We conducted this research with  
permission from CONANP and SEMARNAT (SGPA/DGVS/00910/13 and  
SGPA/DGVS/02716/14).

## Reference list

- 235      **1.** Couzin ID, Krause J. 2003. Self-organization and collective behavior in  
vertebrates. In *Advances in the Study of Behavior* (Eds. Slater PJB, Rosenblatt JS,  
Snowdown CT, Roper TJ) pp 1-67 San Diego, Usa: Academic Press.
- 2.** Ioannou CC. 2017. Grouping and predation. *Encyclopedia of evolutionary  
psychological science*, 1-6.
- 240      **3.** Ioannou CC, Rocque F, Herbert-Read JE, Duffield C, Firth JA. 2019. Predators  
attacking virtual prey reveal the costs and benefits of leadership. *Proc. Natl Acad.  
Sci.* 116, 8925-8930 (doi: [10.1073/pnas.1816323116](https://doi.org/10.1073/pnas.1816323116)).
- 4.** King AJ, Sueur C. 2011. Where next? Group coordination and collective decision  
making by primates. *Int. J. Primatol.* 32, 1245-1267 (doi: [10.1007/s10764-011-9526-7](https://doi.org/10.1007/s10764-011-9526-7)).
- 245      **5.** Walker RH, King AJ, McNutt JW, Jordan NR. 2017. Sneeze to leave: African  
wild dogs (*Lycaon pictus*) use variable quorum thresholds facilitated by sneezes in  
collective decisions. *Proc R Soc B* 284, 20170347 (doi: [10.1098/rspb.2017.0347](https://doi.org/10.1098/rspb.2017.0347)).
- 6.** Prins H. 1996. *Ecology and behaviour of the African buffalo: social inequality  
and decision making* (Vol. 1). Springer Science & Business Media.
- 250      **7.** Sueur C, Petit O. 2008. Organization of group members at departure is driven by  
social structure in Macaca. *Int J Primatol* 29, 1085-1098 (doi: [10.1007/s10764-008-9262-9](https://doi.org/10.1007/s10764-008-9262-9)).
- 8.** Petit O, Gautrais J, Leca JB, Theraulaz G, Deneubourg JL. 2009. Collective  
decision-making in white-faced capuchin monkeys. *Proc R Soc B.* 276, 3495–3503  
(doi: [10.1098/rspb.2009.0983](https://doi.org/10.1098/rspb.2009.0983)).
- 255      **9.** Merkle JA, Sigaud M, Fortin D. 2015. To follow or not? How animals in fusion–  
fission societies handle conflicting information during group decision-making.  
*Ecol. Lett.* 18, 799-806 (doi: [10.1111/ele.12457](https://doi.org/10.1111/ele.12457)).
- 260      **10.** Palacios-Romo TM, Castellanos F, Ramos-Fernandez G. 2019. Uncovering the  
decision rules behind collective foraging in spider monkeys. *Anim. Behav.* 149, 121-  
133 (doi: [10.1016/j.anbehav.2019.01.011](https://doi.org/10.1016/j.anbehav.2019.01.011)).
- 11.** Conradt L, Roper TJ. 2007. Democracy in animals: the evolution of shared  
group decisions. *Proc R Soc B.* 274, 2317-2326 (doi: [10.1098/rspb.2007.0186](https://doi.org/10.1098/rspb.2007.0186)).
- 265      **12.** Sueur C. 2012. Viability of decision-making systems in human and animal  
groups. *J. Theor. Biol.* 306, 93-103 (doi: [10.1016/j.jtbi.2012.04.020](https://doi.org/10.1016/j.jtbi.2012.04.020)).



13. Sueur C, King AJ, Conradt L, Kerth G, Lusseau D, Mettke-Hofmann C et al. 2011. Collective decision-making and fission–fusion dynamics: a conceptual framework. *Oikos* 120, 1608-1617 (doi: [10.1111/j.1600-0706.2011.19685.x](https://doi.org/10.1111/j.1600-0706.2011.19685.x)).
- 270 14. Conradt L. 1998. Could asynchrony in activity between the sexes cause intersexual segregation in ruminant?. *Proc. R. Soc. B.* 265, 1359-1368 (doi: [10.1098/rspb.1998.0442](https://doi.org/10.1098/rspb.1998.0442)).
15. Ruckstuhl KE. 1998. Foraging behaviour and sexual segregation in bighorn sheep. *Anim. Behav.* 56, 99–106 (doi: [10.1006/anbe.1998.0745](https://doi.org/10.1006/anbe.1998.0745)).
- 275 16. Van Belle S, Estrada A, Garber PA. 2013. Collective group movement and leadership in wild black howler monkeys (*Alouatta pigra*). *Behav Ecol. Sociobiol.* 67, 31-41 (doi: 10.1007%2Fs00265-012-1421-5).
17. Strandburg-Peshkin A, Farine DR, Couzin ID, Crofoot MC. 2015. Shared decision-making drives collective movement in wild baboons. *Science* 348, 1358-280 1361 (doi: 10.1126/science.aaa5099).
18. Patriquin KJ, Leonard ML, Broders HG, Garroway CJ. 2010. Do social networks of female northern long-eared bats vary with reproductive period and age? *Behav. Ecol. Sociobiol.* 64, 899-913 (doi: 10.1007%2Fs00265-010-0905-4).
19. Kerth G. 2010. Group decision-making in fission–fusion societies. *Behav. Process* 84, 662-663 (doi: [10.1098/rspb.2006.3647](https://doi.org/10.1098/rspb.2006.3647)).
- 285 20. Kummer H. 1971. *Primate societies: Group techniques of ecological adaptation*. Chicago: Aldine.
21. Aureli F, Schaffner CM, Boesch C, Bearder SK, Call J, Chapman CA et al. 2008. Fission-fusion dynamics: new research frameworks. *Curr. Anthropol.* 49, 627-654 290 (doi: 10.1086/586708).
22. Chapman CA, Wrangham RW, Chapman LJ. 1995. Ecological constraints on group size: an analysis of spider monkey and chimpanzee subgroups. *Behav. Ecol. Sociobiol* 36, 59-70 (doi: 10.1007/s00265-004-0781-x).
23. Lehmann J, Boesch C. 2004. To Fission or to Fusion: Effects of Community 295 Size on Wild Chimpanzee (*Pan troglodytesverus*) Social Organisation. *Behavioral Ecology and Sociobiology* 56, 207-216.
24. Smith JE, Memenis SK, Holekamp KE. 2007. Rank-related partner choice in the fission–fusion society of the spotted hyena (*Crocuta crocuta*). *Behav. Ecol. Sociobiol* 61, 753-765 ([10.1007/s00265-006-0305-y](https://doi.org/10.1007/s00265-006-0305-y)).

- 300     **25.** Schaffner CM, Rebecchini L, Ramos-Fernandez G, Vick LG, Aureli F 2012. Cope with the negative consequences of hurricanes through changes in diet, Activity Budget, and Fission–Fusion Dynamics. *Int J. Primatol.* 33, 922-936 (doi: [10.1007/s10764-012-9621-4](https://doi.org/10.1007/s10764-012-9621-4)).
- 305     **26.** Smith-Aguilar SE, Ramos-Fernández G, Getz WM. 2016. Seasonal changes in socio-spatial structure in a group of free-living spider monkeys (*Ateles geoffroyi*). *PloS One* 11, e0157228 (doi: [10.1371/journal.pone.0157228](https://doi.org/10.1371/journal.pone.0157228)).
- 310     **27.** Heithaus MR, Dill LM. 2002. Food availability and tiger shark predation risk influence bottlenose dolphin habitat use. *Ecology* 83, 480-491 (doi: [10.2307/2680029](https://doi.org/10.2307/2680029)).
- 315     **28.** Link A, Di Fiore A. 2013. Effects of predation risk on the grouping patterns of white-bellied spider monkeys (*Ateles belzebuth belzebuth*) in Western Amazonia. *Am. J. Phys Anthropol.* 150: 579–590 (doi: [10.1002/ajpa.22230](https://doi.org/10.1002/ajpa.22230)).
- 320     **29.** Busia L, Schaffner CM, Rothman JM, Aureli F. 2016. Do fruit nutrients affect subgrouping patterns in wild spider monkeys (*Ateles geoffroyi*)?. *Int. J. Primatol* 37, 738-751 (doi: [10.1007/s10764-016-9935-8](https://doi.org/10.1007/s10764-016-9935-8)).
- 325     **30.** Lusseau D. 2007. Why are male social relationships complex in the Doubtful Sound bottlenose dolphin population?. *PLoS One* 2, e348 (doi: [10.1371/journal.pone.0000348](https://doi.org/10.1371/journal.pone.0000348)).
- 330     **31.** Tokuda M, Boubli JP, Izar P, Strier KB. 2012. Social cliques in male northern muriquis *Brachyteles hypoxanthus*. *Curr. Zool* 58, 342-352 (doi: [10.1093/czoolo/58.2.342](https://doi.org/10.1093/czoolo/58.2.342)).
- 335     **32.** Busia L, Schaffner CM, Aureli F. 2017. Relationship quality affects fission decisions in wild spider monkeys (*Ateles geoffroyi*). *Ethology* 123, 405-411 (doi: [10.1111/eth.12609](https://doi.org/10.1111/eth.12609)).
- 340     **33.** Wallace RB. 2008. Towing the party line: territoriality, risky boundaries and male group size in spider monkey fission–fusion societies. *Am. J. Primatol.* 70, 271-281 (doi: [10.1002/ajp.20484](https://doi.org/10.1002/ajp.20484)).
- 345     **34.** Bond ML, Lee DE, Ozgul A, König B. 2019. Fission–fusion dynamics of a megaherbivore are driven by ecological, anthropogenic, temporal, and social factors. *Oecologia* 191, 335-347 (doi: [10.1007/s00442-019-04485-y](https://doi.org/10.1007/s00442-019-04485-y)).
- 350     **35.** Aureli F, Schaffner CM 2008. Social interactions, social relationships and the social system of spider monkeys. In *Spider monkeys: behavior, ecology and evolution of the genus Ateles* (Ed Campbell CJ) pp 236-265. Cambridge: Cambridge

University Press.

- 335 **36.** Ramos-Fernández G, Ayala-Orozco B 2003. Population size and habitat use of  
spider monkeys at Punta Laguna, Mexico. In *Primates in Fragments* (Eds. Marsh  
LK, Chapman CA), pp. 191-209). New York: Springer US
- 37.** Ramos-Fernández G, Aureli F, Schaffner CM, Vick LG. 2018. Ecología,  
comportamiento y conservación de los monos araña (*Ateles geoffroyi*): 20 años de  
340 estudio en Punta Laguna, México. In *La primatología en Latinoamérica* (Eds  
Urbani B, Kowalewski M, Grassetto Teixeira de Cunha R, de la Torre S, Cortez-  
Ortiz L), pp 531-544. Venezuela: Ediciones IVIC.
- 38.** Ramos-Fernandez G. 2005. Vocal communication in a fission-fusion society: do  
spider monkeys stay in touch with close associates? *Int. J. Primatol.* 26, 1077- 1092  
345 (doi: [10.1007/s10764-005-6459-z](https://doi.org/10.1007/s10764-005-6459-z)).
- 39.** Croft DP, James R, Krause J. 2008. *Exploring animal social networks*.  
Princeton, NJ: Princeton University Press.
- 40.** Rebecchini L, Schaffner CM, Aureli F 2011. Risk is a component of  
social relationships in spider monkeys. *Ethology* 117, 691–699 (doi: [10.1111/j.1439-  
350 0310.2011.01923.x](https://doi.org/10.1111/j.1439-0310.2011.01923.x)).
- 41.** Van Belle S 2015. Female participation in collective group defense in black  
howler monkeys (*Alouatta pigra*). *Am. J. Primatol.* 77, 595-604 (doi:  
[10.1002/ajp.22380](https://doi.org/10.1002/ajp.22380)).
- 42.** Gautrais J, Michelena P, Sibbald A, Bon R, Deneubourg JL. 2007. Allelomimetic  
355 synchronization in Merino sheep. *Anim. Behav.* 74, 1443-1454 (doi:  
[10.1016/j.anbehav.2007.02.020](https://doi.org/10.1016/j.anbehav.2007.02.020)).
- 43.** Conradt L, Roper TJ. 2000. Activity synchrony and social cohesion: a fission-  
fusion model. *Proc. R. Soc. B* 267, 2213-2218 (doi: [10.1098/rspb.2000.1271](https://doi.org/10.1098/rspb.2000.1271)).
- 44.** Hartwell KS, Notman H, Bonenfant C, Pavelka MS. 2014. Assessing the  
360 occurrence of sexual segregation in spider monkeys (*Ateles geoffroyi yucatanensis*),  
its mechanisms and function. *Int. J. Primatol* 35, 425-444 (doi: [10.1007/s10764-  
013-9746-0](https://doi.org/10.1007/s10764-013-9746-0)).
- 45.** Busia L, Schaffner CM, Aureli F. 2021. Activity synchronization and fission  
decisions , Dryad, Dataset, <https://doi.org/10.5061/dryad.ttdz08kzr>.
- 365 **46.** Bates D, Maechler M, Bolker B, Walker S 2014. lme4: Linear mixed-effects  
models using Eigen and S4. R package version 1.1-7. [http://CRAN.R-project.  
org/package= lme4](http://CRAN.R-project.org/package=lme4).

47. Dobson AJ, Barnett AG 2008. *An introduction to generalized linear models*. Boca Raton, FL: Chapman & Hall/CRC.
- 370 48. Nakagawa S, Johnson PC, Schielzeth H. 2017. The coefficient of determination  $R^2$  and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *J. R. Soc. Interface* 14, 20170213 (doi: [10.1098/rsif.2017.0213](https://doi.org/10.1098/rsif.2017.0213)).
- 375 49. Cords M, Aureli F. 2000. Reconciliation and relationship quality. In *Natural conflict resolution* (Eds Aureli F, De Waal F). pp. 177–198. California: University of California Press.
50. Tarr B, Launay J, Dunbar RIM. 2016 Silent disco: dancing in synchrony leads to elevated pain thresholds and social closeness. *Evol. Hum. Behav.* 37, 343–349. (doi:10.1016/j.evolhumbehav.2016.02.004).
- 380 51. Connor RC, Smolker R, Bejder L. 2006. Synchrony, social behaviour and alliance affiliation in Indian Ocean bottlenose dolphins, *Tursiops aduncus*. *Anim. Behav.* 72, 1371–1378 (doi: [10.1016/j.anbehav.2006.03.014](https://doi.org/10.1016/j.anbehav.2006.03.014)).
52. Senigaglia V, de Stephanis R, Verborgh P, Lusseau D. 2012. The role of synchronized swimming as affiliative and anti-predatory behavior in long-finned pilot whales. *Behav. Process.* 91, 8–14 (doi: [10.1016/j.beproc.2012.04.011](https://doi.org/10.1016/j.beproc.2012.04.011)).
- 385 53. Duranton C, Gaunet F. 2016. Behavioural synchronization from an ethological perspective: Short overview of its adaptive values. *Adapt Behav.* 24, 181–191 (doi: [10.1177/1059712316644966](https://doi.org/10.1177/1059712316644966)).
54. Williams JM, Liu HY, Pusey AE. 2002. Costs and benefits of grouping for female chimpanzees at Gombe. In: *Behavioural Diversity in Chimpanzees and Bonobos* (Eds. Boesch C, Hohmann G, Marchant L), pp 192–203. Cambridge: Cambridge University Press.
- 390 55. Shimooka Y. 2005. Sexual differences in ranging of *Ateles belzebuth belzebuth* at La Macarena, Colombia. *Int. J Primatol.* 26, 385–406 (doi: [10.1007/s10764-005-2930-0](https://doi.org/10.1007/s10764-005-2930-0)).
- 395 56. Aureli F, Fraser ON, Schaffner CM, Schino G. 2012. The regulation of social relationships. In *The Evolution of Primate Societies* (Eds. Silk JB, Kappeler PM, Mitani JC, Call J, Palombit RA), pp 531–551. Chicago: University of Chicago Press.