



LJMU Research Online

Beigrezaei, S, Mazidi, M, Davies, IG, Salehi-Abargouei, A, Ghayour-Mobarhan, M and Khayatzadeh, SS

The association between dietary behaviors and insomnia among adolescent girls in Iran

<http://researchonline.ljmu.ac.uk/id/eprint/16255/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Beigrezaei, S, Mazidi, M, Davies, IG, Salehi-Abargouei, A, Ghayour-Mobarhan, M and Khayatzadeh, SS (2022) The association between dietary behaviors and insomnia among adolescent girls in Iran. *Sleep Health*, 8 (2). pp. 195-199. ISSN 2352-7218

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk


<http://researchonline.ljmu.ac.uk/>



Contents lists available at ScienceDirect

Sleep Health

Journal of the National Sleep Foundation

journal homepage: sleephealthjournal.org

Dietary behavior and insomnia

The association between dietary behaviors and insomnia among adolescent girls in Iran

Sara Beigrezaei, PhD^{a,b,#}, Mohsen Mazidi, PhD^{c,d,#}, Ian G. Davies, PhD^e,
 Amin Salehi-Abargouei, PhD^{a,b}, Majid Ghayour-Mobarhan, MD, PhD^{f,**},
 Sayyed Saeid Khayyatzaeh, PhD^{a,b,*}

^a Nutrition and Food Security Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran^b Department of Nutrition, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran^c Department of Twin Research and Genetic Epidemiology, King's College London, London, UK^d Medical Research Council Population Health Research Unit, University of Oxford, Oxford, UK. Clinical Trial Service Unit and Epidemiological Studies Unit (CTSU), Nuffield Department of Population Health, University of Oxford, Oxford, UK^e Research Institute of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK^f Metabolic Syndrome Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Keywords:

Dietary behavior
 Spicy food
 Breakfast
 Meal
 Insomnia

ABSTRACT

Background: Insomnia is associated with a poor quality of life and increased risk of somatic and social problems. The aim of current study was to investigate the relationship between dietary behaviors and insomnia in Iranian adolescent girls.

Methods: This cross-sectional study was performed among 988 girls aged 12–18 years. A questionnaire was used to determine dietary behaviors in nine domains. To assess insomnia, a validated Iranian version of the Insomnia Severity Index was applied. Multivariable logistic regression examined the association between dietary behaviors and insomnia in crude and adjusted models.

Results: Highest adherence to regular meal consumption was related to the lowest odds of insomnia (odds ratio [OR]: 0.44, 95% confidence interval [CI]: 0.24–0.81). Compared with individuals who consumed breakfast never or once a week, those who always consumed breakfast had a lower likelihood of insomnia (OR: 0.56, 95% CI: 0.36–0.88). These associations remained significant in all adjusted models. Subjects who ate spicy food every day had 4.73 times greater odds of insomnia than individuals who never ate spicy food (OR: 4.73, 95% CI: 1.09–20.56). After controlling for age, menstruation, parent death, parents' divorce and parents' (mother and father) employment status, this relationship remained (OR: 4.59, 95% CI: 1.05–20.10); however, the association was no longer significant after controlling for the other covariates. No significant relationship was found between other dietary habits and insomnia for the unadjusted or adjusted models.

Conclusion: Lowest rates of insomnia were found among participants who had the lowest frequency of eating spicy foods and the highest frequency of eating breakfast and eating regular meals. Further prospective studies are required to confirm these findings.

© 2021 National Sleep Foundation. Published by Elsevier Inc. All rights reserved.

Introduction

Sleep disorders are associated with an increased risk of chronic health conditions, including cardiovascular disease, endocrine disruption, and psychological disturbance.^{1,2} Insomnia is a common sleep

disorder often associated with poor quality of life and increased risk of somatic and social problems.^{3,4} Adolescence is a critical period for an almost 3-fold increase in prevalence of insomnia, particularly among girls.⁵ In puberty, cultural transitions, brain maturation, biological, and psychological changes predispose adolescents to sleep problems.^{6,7} Along with psychological and physiological disorders, diet and lifestyle are the main determinants for sleep quality and quantity.⁸ Findings from 2 epidemiological studies suggest that a higher dietary intake of fish, fruits, and vegetables improves sleep quality, whereas a high dietary intake of carbohydrates is related to poor sleep health.^{3,9–11}

Dietary behavior is considered a principal component of nutrition-related lifestyle, which includes the regularity of breakfast, main

*Corresponding author: Sayyed Saeid Khayyatzaeh, PhD, Nutrition and Food Security Research Center, Shahid Sadoughi University of Medical Sciences, Shohadaye gomnam BLD. ALEM square, Yazd, Iran. Tel.: +983531492228.**Corresponding author: Majid Ghayour-Mobarhan, MD, PhD, Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Azadi square, Mashhad, Iran. Tel.: +985133223822.

E-mail address: khayyatzaeh@yahoo.com (S.S. Khayyatzaeh).

Equally as first author.

meal, and snack consumption; the quality of food (eg, fried and spicy food); and sufficient chewing.¹² Poor dietary behaviors are associated with an increased risk of obesity, type 2 diabetes, metabolic syndrome, cardiovascular disease, gastrointestinal diseases and mental health problems.^{13–15} These noncommunicable diseases are all associated with poor sleep outcomes.^{16,17} Although some studies have investigated the association between diet-related practices and overall sleep health, limited data are available about the relationship between dietary behavior profiles and chronic insomnia. Detrimental effects of some unhealthy dietary behavior, including energy-dense food consumption at night and frequent breakfast skipping, on poor sleep outcomes have been reported.^{3,13} Additionally, some studies have indicated that poor oral health is associated with sleep problems. A cross-sectional study reported a positive association between gingival inflammation and sleep problems,¹⁸ and a cohort study showed that individuals with fewer than 10 teeth were at a higher risk of sleep disorders than adults with at least 20 teeth.¹⁹

To the best of our knowledge, no published studies have investigated the relationship between dietary behavior and insomnia in Iranian adolescents. Given the importance of dietary behavior and adverse consequences of poor sleep health, we explored the association between dietary behavior profiles and chronic insomnia in Iranian adolescent girls.

Materials and methods

Study population and design

This cross-sectional study was performed in January 2015 in the cities of Mashhad and Sabzevar, Iran. The population consisted of 988 female students aged 12–18 years recruited using random cluster sampling from schools in these cities. Exclusion criteria included individuals with autoimmune disease, cancer, metabolic bone disease, hepatic or renal failure, cardiovascular disorders, malabsorption or thyroid, parathyroid, adrenal diseases, and anorexia nervosa or bulimia. Additionally, participants were excluded if they took, within the last 6 months, any medication categorized as anti-inflammatory, antidepressant, antidiabetic, or antiobesity; vitamin D or calcium supplements; or hormone therapy.²⁰ The Ethics Committee of Mashhad University of Medical Sciences, Mashhad, Iran, approved the study, and all participants and their parents provided written informed consent.

Dietary behavior assessment

To assess dietary behaviors during the past month, we used a dietary behavior questionnaire, with the following questions: How many main meals do you eat per day? (1/ 2/ 3); How many times do you consume snacks throughout a day? (Never/ 1–2/ ≥ 3 times); Do you have consistent meal consumption timing? (Never/ sometimes/ almost/ always); How many days do you eat breakfast during a week? (Never/ 1 day/ 2–4 days/ 5–6 days/ every day); How much do you chew your food? (Low/ moderate/ high); Can you chew any food? (Yes/ only soft food/ no); Do you drink fluids with your meals? (Never/ sometimes/ almost/ always); How many times do you eat fried foods during a week? (Never/ 1–3 in week/ 4–6 in week/ every day); How many times do you eat fried foods during a week? (Never/ 1–3 in week/ 4–6 in week/ every day).²⁰

Insomnia assessment

To evaluate insomnia, we used the Insomnia Severity Index (ISI) questionnaire, which has 7 questions with scores ranging between 0 and 4, stratified into 5 categories: 0 (None), 1 (Mild), 2 (Moderate), 3

(Severe), and 4 (Very Severe). The total ISI score ranges from 0 to 28. Insomnia is defined as total ISI score was >7 .²¹

Covariate measurements

General information including age, menstruation status, parental death or divorce, and father's and mother's occupation was collected by face-to-face interview using a standard questionnaire. Physical activity was assessed through a validated modifiable activity questionnaire²² and provided as metabolic equivalents in hours per week. Weight and height were measured using standard protocols, and the mean of 2 measurements was reported. Body mass index (BMI) was calculated as weight (kg) divided by square of the height (m^2). Dietary intake was evaluated during face-to-face interview by a valid and reliable food frequency questionnaire with 147 food items.²³ Reported dietary intakes were converted to grams and entered into the Nutritionist IV software. To calculate daily nutrient intake values for each participant, the US Department of Agriculture's national nutrient databank was used.²⁴

Statistical analyses

The normality of data was assessed using the Kolmogorov-Smirnov test. To compare continuous and categorical variables between subjects with vs. without insomnia, independent samples t-test and chi-square tests were used, respectively. *P* values $<.05$ were used to compare the distribution of quantitative or qualitative variables across categories of dietary habits or insomnia status. Logistic regression was applied to evaluate the relationship between dietary behavior and insomnia prevalence in crude and adjusted models. Adjustments for age, menstruation, parents' death, parents' divorce, and father's and mother's employment status were performed in Model I. Physical activity, BMI and energy intake were additionally adjusted in the Model II. Bonferroni correction was applied to all models to adjust for multiple comparisons. The significance level was assessed in 2 steps: traditional cutoff of *P* values $<.05$ and Bonferroni correction of *P* value $<.006$. To analyze the data, the Statistical Package for Social Sciences (SPSS) (version 23.0, SPSS Inc, Chicago, IL) was used.

Results

The general characteristics of the study participants are summarized in Table 1. The prevalence of insomnia was 17%. There were no significant differences for age, BMI percentile, or physical activity between individuals with and without insomnia. The prevalence of menstruation and parental death or divorce were significantly higher in subjects with insomnia.

Dietary intakes are summarized in Table 2. There were no significant differences for dietary intakes between individuals with and without insomnia in the crude or energy-adjusted models. The distribution of insomnia prevalence across categories of dietary behaviors is provided in Table 3. Significant differences were observed for insomnia prevalence across frequency of main meal intake; regular meal consumption (consistent meal consumption timing); and frequencies of breakfast, fried food, and spicy food consumption ($P <.05$; Table 3). However, there was no difference for insomnia prevalence by rate of chewing food, snack consumption, or chewing sufficiency.

For the crude and multivariable-adjusted models, the odds ratios for the association between dietary behavior and insomnia are presented in Table 4. The highest adherence to regular meal consumption was related to lowest odds of insomnia (odds ratio [OR]: 0.44, 95% confidence interval [CI]: 0.24–0.81). This association remained significant after adjustments for potential confounding variables in Model I (OR: 0.47, 95% CI: 0.25–0.87) and Model II (OR: 0.46, 95% CI: 0.22–0.95). Compared with individuals who never consumed

Table 1
General characteristics of study participants with and without insomnia

Variables	Insomnia		P value*	
	No (n = 814)	Yes (n = 174)		
Age (y)	14.52 ± 1.52	14.71 ± 1.55	.097	
Menstruation (yes) (%)	87.7	92.9	.041	
Parents' death (yes) (%)	3.7	4.0	.007	
Parents' divorce (yes) (%)	3.7	7.1	.001	
Father employment status (employed) (%)	99.1	99	.600	
Mother employment status (employed) (%)	5.9	7.1	.461	
Body mass index percentile (%)	<25	41.9	42.8	.885
	25-50	28	27	
	50-85	10.5	9	
	≥85	19.7	21.2	
Physical activity (MET.h/week)	45.35 ± 3.57	45.56 ± 3.62	.444	

* Obtained from independent samples t-test for continuous variables and chi-squared test for categorical variables.

breakfast or consumed breakfast once a week, those who always consumed breakfast had a lower chance of insomnia (OR: 0.56, 95% CI: 0.36-0.88). This significant relationship was observed in all adjusted models; Model I (OR: 0.57, 95% CI: 0.36-0.89) and Model II (OR: 0.47, 95% CI: 0.27-0.80). Subjects who ate spicy food every day had 4.73 times greater odds for insomnia than the individual who never ate spicy food (OR: 4.73, 95% CI: 1.09-20.56). This significant association was observed only in the first adjusted model (OR: 4.59, 95% CI: 1.05-20.10). In crude and adjusted models, there was no significant relationship with insomnia for other dietary habits, including main meal intake, snack consumption, rate of food chewing, intake of fluid with meal, consumption of fried foods, and chewing sufficiency. After applying the Bonferroni correction, the inverse association between highest adherence to breakfast consumption and odds of insomnia remained significant only in the final adjusted model (*P* value < .005).

Discussion

The aim of the present study was to investigate the association between dietary behaviors and the prevalence of insomnia in

Table 2
Energy and dietary intakes of study participants between subjects with and without insomnia

	Insomnia		P value*	
	No (n = 578)	Yes (n = 174)		
Energy intake (Kcal)	2693.3 ± 833.8	2733.9 ± 843.2	.577	
Carbohydrate (% energy)	54.8 ± 7.34	54.79 ± 7.5	.956	
Protein (% energy)	13.7 ± 2.2	13.5 ± 2.2	.344	
Fat (% energy)	33.78 ± 7.8	34 ± 2.2	.708	
Cholesterol (mg)	Crude	233.3 ± 127.8	251.2 ± 166	.131
	Energy adjusted	235.6 ± 114.8	250.8 ± 151.5	.224
Potassium (mg)	Crude	3717 ± 1362	3754 ± 1403	.763
	Energy adjusted	3740 ± 777	3721 ± 712	.774
Sodium (mg)	Crude	4163 ± 1803	4206 ± 1661	.777
	Energy adjusted	4172 ± 1386	4160 ± 1235	.921
Sucrose (gr)	Crude	22.2 ± 14.2	22.07 ± 14.2	.881
	Energy adjusted	22.9 ± 13.07	22.5 ± 13.01	.683
Vitamin A (mcg)	Crude	587.5 ± 593.5	610.7 ± 415.6	.630
	Energy adjusted	590.4 ± 576.9	605.9 ± 368.5	.738
Vitamin C (mg)	Crude	95.8 ± 63.5	94.5 ± 56.2	.814
	Energy adjusted	191.1 ± 63.5	189.8 ± 56.2	.814
Folic acid (mcg)	Crude	606.5 ± 197.6	600 ± 214.2	.753
	Energy adjusted	656 ± 176.13	638.4 ± 152.7	.200
Calcium (mg)	Crude	1130 ± 488.5	1182 ± 530.6	.256
	Energy adjusted	1137.9 ± 368.7	1172 ± 410.8	.311

* Obtained from independent samples t-test.

adolescent girls. Eating breakfast most days of the week and regular meal consumption were related to the lowest odds of insomnia, while the highest frequency of spicy food intake increased the chance of insomnia. To the best of our knowledge, this is the first study to investigate dietary behaviors and insomnia prevalence.

We found that the highest adherence to regular meal consumption was associated with decreased odds of insomnia. Wehrens et al reported that meal timing alters human molecular clocks and circadian rhythms, which may influence sleep-wake schedules.²⁵ In addition, previous studies have reported that irregular meal timing can produce a disruption in the circadian system that might lead to

Table 3
The distribution of insomnia prevalence across categories of dietary behaviors

Type of dietary habit	Insomnia (%)		P value*	
	No	Yes		
Main meal intake	1 time	4 (28)	3.2 (7)	.033
	2 times	25.4 (177)	34.4 (75)	
	3 times	70.6 (492)	62.4 (136)	
Snack consumption	Never	5.2 (36)	7.1 (15)	.093
	1-2	58.3 (402)	55.2 (117)	
	3-5	32 (221)	29.2 (62)	
	>5	4.5 (31)	8.5 (18)	
Regular meal consumption	Never	9.7 (68)	12 (26)	.007
	Sometimes	38 (265)	46.1 (100)	
	Almost	30.2 (211)	30 (65)	
Rate of food chewing	Always	22.1 (154)	23.7 (26)	.247
	Low	11.4 (80)	15.5 (34)	
	Moderate	78.5 (552)	74 (162)	
Breakfast consumption	High	10.1 (71)	10.5 (23)	.013
	Never or 1 day	14.6 (101)	18.7 (40)	
	2-4 day	25.8 (179)	29.9 (64)	
Intake of fluids with meal	5-6 day	14.9 (103)	19.2 (41)	.117
	Every day	81.8 (310)	32.2 (69)	
	Never	5.5 (38)	6.9 (15)	
Consumption of fried food	Sometimes	35.6 (248)	30 (65)	.033
	Almost	29.7 (207)	26.3 (57)	
	Always	29.3 (204)	36.9 (80)	
Consumption of spicy food	Never	3.5 (24)	1.4 (3)	.008
	1-3 in week	67.2 (467)	59.8 (131)	
	4-6 in week	24.6 (171)	33.3 (73)	
	Every day	4.7 (33)	5.5 (12)	
Chewing sufficiency	Never	3.1 (22)	0.9 (2)	.093
	1-3 in week	28.8 (202)	25.3 (55)	
	4-6 in week	39.6 (278)	34.1 (74)	
Only soft foods	Every day	28.5 (200)	39.6 (86)	.093
	Yes	96.9 (678)	93.6 (204)	
	No	2.7 (19)	5.5 (12)	
No	0.4 (3)	0.9 (2)		

* Obtained from chi-squared test.

Table 4
Multivariable-adjusted odds ratio of the association between dietary behaviors and insomnia

Type of dietary habit	Crude	Model I	Model II
Main meal intake			
1 time	1	1	1
2 times	1.69 (0.71-4.05)	1.72 (0.72-4.14)	3.16 (0.91-11.01)
3 times	1.11 (0.47-2.58)	1.15 (0.49-2.72)	2.23 (0.65-7.63)
<i>P</i> trend	.085*	.128	.578
Snack consumption			
Never	1	1	1
1-2	0.70 (0.37-1.32)	0.69 (0.36-1.32)	0.63 (0.31-1.28)
3-5	0.67 (0.37-1.31)	0.67 (0.34-1.33)	0.62 (0.29-1.32)
>5	1.39 (0.6-3.22)	1.52 (0.64-3.59)	1.39 (0.52-3.74)
<i>P</i> trend	.526	.410	.600
Regular meal consumption			
Never	1	1	1
Sometimes	0.98 (0.60-1.64)	0.102 (0.61-1.70)	1.187 (0.66-2.15)
Almost	0.80 (0.47-1.37)	0.83 (0.48-1.43)	0.90 (0.42-1.48)
Always	0.44 (0.24-0.81)	0.47 (0.25-0.87)	0.46 (0.22-0.95)
<i>P</i> trend	.002	.003	.002
Rate of food chewing			
Low	1	1	1
Moderate	0.69 (0.44-1.07)	0.71 (0.45-1.10)	0.63 (0.38-1.05)
High	0.76 (0.41-1.41)	0.78 (0.42-1.45)	0.57 (0.27-1.22)
<i>P</i> trend	.308	.353	.111
Breakfast consumption			
Never or 1 day	1	1	1
2-4 day	0.90 (0.56-1.43)	0.90 (0.56-1.44)	0.91 (0.53-1.58)
5-6 day	1.00 (0.60-1.60)	0.99 (0.59-1.67)	0.90 (0.50-1.64)
Every day	0.56 (0.36-0.88)	0.57 (0.36-0.89)	0.47 (0.27-0.80)
<i>P</i> trend	.005	.007	.001#
Intake of fluids with meal			
Never	1	1	1
Sometimes	0.66 (0.34-1.28)	0.67 (0.35-1.31)	0.53 (0.25-1.16)
Almost	0.70 (0.36-1.35)	0.72 (0.36-1.40)	0.60 (0.28-1.33)
Always	0.99 (0.51-1.9)	1.01 (0.52-1.95)	0.81 (0.38-1.76)
<i>P</i> trend	.154	.147	.293
Consumption of fried foods			
Never	1	1	1
1-3 in week	2.24 (0.66-7.59)	2.12 (0.62-7.24)	1.39 (0.40-4.89)
4-6 in week	3.41 (0.99-11.7)	3.42 (0.99-11.82)	2.40 (0.66-8.61)
Every day	2.91 (0.74-11.45)	2.78 (0.69-11.12)	2.21 (0.51-9.61)
<i>P</i> trend	.011	.007	.009
Consumption of spicy foods			
Never	1	1	1
1-3 in week	2.99 (0.68-13.13)	2.87 (0.65-12.70)	1.58 (0.34-7.33)
4-6 in week	2.93 (0.67-12.73)	2.85 (0.65-12.50)	1.58 (0.34-7.30)
Every day	4.73 (1.09-20.56)	4.59 (1.05-20.10)	2.25 (0.49-1.40)
<i>P</i> trend	.003	.003	.088
Chewing sufficiency			
Yes	1	1	1
Only soft foods	2.10 (1.00-4.39)	2.19 (1.30-4.66)	2.00 (0.815-4.92)
No	2.21 (0.37-13.35)	2.07 (0.34-12.59)	2.38 (0.39-14.95)
<i>P</i> trend	.042	.001	.090

Model I: adjusted for age, menstruation, parent's death, parent's divorce, parent's (mother and father) employment status.

Model II: additionally, adjusted for physical activity, BMI and energy intake.

* *P* for trend based on logistic regression.

Significant after Bonferroni correction.

unhealthy consequences such as glucose intolerance and obesity-related factors.^{26,27}

In the current study, more than 62% of individuals who consumed only one or 2 main meals per day did not consume breakfast. Skipping breakfast appears to be an unhealthy and frequent dietary behavior observed in many populations, especially young people and adolescents.¹⁴ Breakfast is considered to be the "most important meal of the day,"²⁸ and a main indicator of a healthy lifestyle.²⁹ Indeed, skipping breakfast is a growing problem, especially among adolescent girls.³⁰ Our findings showed the highest frequency of breakfast consumption was associated with the lowest chance of

insomnia among adolescent girls. These results are consistent with previous studies that reported an association between the lack of regular breakfast consumption and sleep disorders in adolescents.^{14,31} Skipping breakfast is related to lower sleep duration and higher sleep disturbances,^{32,33} and adolescents with a greater frequency of breakfast intake have a lower likelihood of insomnia.¹⁴ Arakawa et al.³⁴ reported that skipping breakfast was directly associated with later bedtimes among 3754 high school students. Also, lower stress, anxiety and depression levels were observed among individuals who consumed breakfast on a daily basis.³⁵

A greater frequency of spicy food consumption was associated with increased odds of insomnia in our sample. A trial on a group of young healthy men revealed a clear relationship between spicy meals and sleep disturbances.³⁶ Indeed, the effect of spicy foods on sleep may be mediated through 2 mechanisms. Firstly, isothiocyanates and capsaicin directly induce oxygen consumption through stimulation of heat-producing tissues.³⁷ Second, capsaicin is concentrated in central and peripheral nervous tissue and causes brain reflexes by direct stimulation of sensory C fibers that relate to poor sleep.³⁸ There is strong evidence that individuals who have higher intake of spicy foods experience more intense symptoms of functional gastrointestinal disorders (FGIDs),^{39,40} an important risk factor for sleep-related disorders.^{41,42}

Some important points should be considered in interpreting the results of our study. We cannot determine the direction of the associations revealed in our cross-sectional design. Previous studies have demonstrated that sleep quality and sleep duration may directly modulate hormones involved in appetite and energy balance, and these, in turn, might alter dietary intakes and food choices.^{43,44} Ghrelin is one of the strongest appetite hormones, while leptin is commonly considered a major anorexigenic hormone. Several previous reports indicated ghrelin and leptin levels were altered in sleep disturbances.⁴⁵⁻⁴⁷ Sleep is the most important behavior that aligns the circadian clock with other human behaviors, such as eating. Individuals with poor sleep often tend to follow irregular sleep schedules, which could determine whether they would have regular meal timing during the day or how often they would skip breakfast.^{48,49} In a study on adolescents in Taiwan, adequate sleep duration was related to healthy dietary behavior such as eating breakfast every day.⁵⁰

To the best of our knowledge, our study is the first to examine the relationship between adherence to specific dietary behaviors and prevalence of insomnia. It was carried out on a large sample population of adolescent girls, and we considered multiple confounding factors. Nevertheless, some limitations should be considered. First, the cross-sectional nature of the study is a major limitation. Second, the dietary behavior and insomnia data relied on self-reported questionnaires, which may be susceptible to misreporting. The current study was performed only on adolescent girls, limiting generalizability; and we did not account for potential confounding by parental income or food insecurity.

Conclusion

In summary, the lowest rates of insomnia were found among participants who had the lowest frequency of eating spicy foods and the highest frequency of eating breakfast and eating regular meals. We are unable to explore any significant associations between insomnia and other dietary habits, including main meal intake, snack consumption, rate of food chewing, chewing sufficiency, fluid intake with meals and consumption of fried foods. Further studies, particularly of a prospective nature, are required to examine the associations between adherence to specific dietary behaviors and the risk of sleep-related disorders.

Authors' contributions

The paper was drafted by S.B. with contributions from all authors. S.S.K., A.S.A., and M.G.M. designed the study. M.M. and S.S.K.H. participated in field implementation and sampling. M.G.M. and S.S.K. involved in clinical examination and patient confirmation. S.B., M.M., and I.D. contributed to statistical analyses. M.G.M., S.S.K.H., and I.D. supervised the study. Final linguistic revision was done by I.D. All authors contributed to the development of, read and approved the final version of the manuscript.

Declaration of conflict of interest

The authors have declared no conflicts of interest.

Funding

This study was financially supported by Mashhad University of Medical Sciences, Mashhad, grant number 931188.

Acknowledgments

The authors are grateful to all study participants, volunteers, and study personnel.

References

- de Zambotti M, Goldstone A, Colrain IM, et al. Insomnia disorder in adolescence: Diagnosis, impact, and treatment. *Sleep Med Rev.* 2019;39:12–24.
- Sivertsen B, Lallukka T, Salo P, et al. Insomnia as a risk factor for ill health: results from the large population-based prospective HUNT study in Norway. *J Sleep Res.* 2014;23(2):124–132.
- Katagiri R, Asakura K, Kobayashi S, et al. Low intake of vegetables, high intake of confectionary, and unhealthy eating habits are associated with poor sleep quality among middle-aged female Japanese workers. *J Occup Health.* 2014;56(5):359–368.
- Johnson EO, Roth T, Schultz L, et al. Epidemiology of DSM-IV insomnia in adolescence: lifetime prevalence, chronicity, and an emergent gender difference. *Pediatrics.* 2006;117(2):e247–e256.
- Zhang J, Chan NY, Lam SP, et al. Emergence of sex differences in insomnia symptoms in adolescents: a large-scale school-based study. *Sleep.* 2016;39(8):1563–1570.
- Dahl RE, Lewin DS. Pathways to adolescent health sleep regulation and behavior. *J Adolesc Health.* 2002;31(6):175–184.
- Campbell IG, Grimm KJ, de Bie E, et al. Sex, puberty, and the timing of sleep EEG measured adolescent brain maturation. *Proc Natl Acad Sci USA.* 2012;109(15):5740–5743.
- Khan MK, Faight EL, Chu YL, et al. Is it nutrients, food items, diet quality or eating behaviours that are responsible for the association of children's diet with sleep? *J Sleep Res.* 2017;26(4):468–476.
- Tanaka E, Yatsuya H, Uemura M, et al. Associations of protein, fat, and carbohydrate intakes with insomnia symptoms among middle-aged Japanese workers. *J Epidemiol.* 2013;23(2):132–138.
- Pahlavani N, Khayyat-zadeh SS, Banazadeh V, et al. Adherence to a Dietary Approach to Stop Hypertension (DASH)-style in relation to daytime sleepiness. *Nat Sci Sleep.* 2020;12:325.
- Rostami H, Khayyat-zadeh SS, Tavakoli H, et al. The relationship between adherence to a Dietary Approach to Stop Hypertension (DASH) dietary pattern and insomnia. *BMC Psychiatry.* 2019;19(1):1–7.
- Sun J, Yi H, Liu Z, et al. Factors associated with skipping breakfast among inner Mongolia medical students in China. *BMC Public Health.* 2013;13(1):42.
- Kapantais E, Chala E, Kaklamanou D, et al. Breakfast skipping and its relation to BMI and health-compromising behaviours among Greek adolescents. *Public Health Nutr.* 2011;14(1):101–108.
- Otsuka Y, Kaneita Y, Itani O, et al. Association between unhealthy dietary behaviors and sleep disturbances among Japanese adolescents: a nationwide representative survey. *Sleep Biol Rhythms.* 2019;17(1):93–102.
- Shin A, Lim S-Y, Sung J, et al. Dietary intake, eating habits, and metabolic syndrome in Korean men. *J Am Diet Assoc.* 2009;109(4):633–640.
- Grandner MA, Jackson NJ, Pak VM, et al. Sleep disturbance is associated with cardiovascular and metabolic disorders. *J Sleep Res.* 2012;21(4):427–433.
- Sharma N, Lee J, Youssef I, et al. Obesity, cardiovascular disease and sleep disorders: insights into the rising epidemic. *J Sleep Disord Ther.* 2017;6(1).
- Carra MC, Schmitt A, Thomas F, et al. Sleep disorders and oral health: a cross-sectional study. *Clin Oral Investig.* 2017;21(4):975–983.
- Koyama S, Aida J, Cable N, et al. Sleep duration and remaining teeth among older people. *Sleep Med.* 2018;52:18–22.
- Khayyat-zadeh SS, Mirmoosavi SJ, Fazeli M, et al. High-dose vitamin D supplementation is associated with an improvement in several cardio-metabolic risk factors in adolescent girls: a nine-week follow-up study. *Ann Clin Biochem.* 2018;55(2):227–235.
- Morin CM. *Insomnia: Psychological Assessment and Management.* New York: Guilford Press; 1993.
- Delshad M, Ghanbarian A, Ghaleh NR, et al. Reliability and validity of the modifiable activity questionnaire for an Iranian urban adolescent population. *Int J Prev Med.* 2015;6(1):3.
- Hosseini Esfahani F, Asghari G, Mirmiran P, et al. Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the Tehran Lipid and Glucose Study. *J Epidemiol.* 2010;20(2):150–158.
- Pehrsson P, Haytowitz D, Holden J, et al. USDA's national food and nutrient analysis program: food sampling. *J Food Compos Anal.* 2000;13(4):379–389.
- Wehrens SM, Christou S, Isherwood C, et al. Meal timing regulates the human circadian system. *Curr Biol.* 2017;27(12):1768–1775.e3.
- St-Onge M-P, Ard J, Baskin ML, et al. Meal timing and frequency: implications for cardiovascular disease prevention: a scientific statement from the American Heart Association. *Circulation.* 2017;135(9):e96–e121.
- Lopez-Minguez J, Gómez-Abellán P, Garaulet M. Timing of breakfast, lunch, and dinner. Effects on obesity and metabolic risk. *Nutrients.* 2019;11(11):2624.
- Mahoney CR, Taylor HA, Kanarek RB, et al. Effect of breakfast composition on cognitive processes in elementary school children. *Physiol Behav.* 2005;85(5):635–645.
- Nedeltsheva AV, Kilkus JM, Imperial J, et al. Sleep curtailment is accompanied by increased intake of calories from snacks. *Am J Clin Nutr.* 2009;89(1):126–133.
- Lundqvist M, Vogel NE, Levin L-A. Effects of eating breakfast on children and adolescents: A systematic review of potentially relevant outcomes in economic evaluations. *Food Nutr Res.* 2019;63.
- Al-Hazzaa H, Musaiger A, Abahussain N, et al. Lifestyle correlates of self-reported sleep duration among Saudi adolescents: a multicentre school-based cross-sectional study. *Child Care Health Dev.* 2014;40(4):533–542.
- Weiss A, Xu F, Storfer-Isser A, et al. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep.* 2010;33(9):1201–1209.
- Ohida T, Osaki Y, Doi Y, et al. An epidemiologic study of self-reported sleep problems among Japanese adolescents. *Sleep.* 2004;27(5):978–985.
- Arakawa M, Taira K, Tanaka H, et al. A survey of junior high school students' sleep habit and lifestyle in Okinawa. *Psychiatry Clin Neurosci.* 2001;55(3):211–212.
- Richards G, Smith AP. Breakfast and energy drink consumption in secondary school children: breakfast omission, in isolation or in combination with frequent energy drink use, is associated with stress, anxiety, and depression cross-sectionally, but not at 6-month follow-up. *Front Psychol.* 2016;7:106.
- Edwards SJ, Montgomery IM, Colquhoun EQ, et al. Spicy meal disturbs sleep: an effect of thermoregulation? *Int J Psychophysiol.* 1992;13(2):97–100.
- Henry C, Emery B. Effect of spiced food on metabolic rate. Human nutrition. *Clin Nutr.* 1986;40(2):165–168.
- Donnerer J, Amann R, Schuligoi R, et al. Absorption and metabolism of capsaicinoids following intragastric administration in rats. *Naunyn-Schmiedeberg's Arch Pharmacol.* 1990;342(3):357–361.
- Saneei P, Sadeghi O, Feizi A, et al. Relationship between spicy food intake and chronic uninvestigated dyspepsia in Iranian adults. *J Dig Dis.* 2016;17(1):28–35.
- Khayyat-zadeh SS, Kazemi-Bajestani SMR, Mirmoosavi SJ, et al. Dietary behaviors in relation to prevalence of irritable bowel syndrome in adolescent girls. *J Gastroenterol Hepatol.* 2018;33(2):404–410.
- Hyun MK, Baek Y, Lee S. Association between digestive symptoms and sleep disturbance: a cross-sectional community-based study. *BMC Gastroenterol.* 2019;19(1):1–6.
- Lacy BE, Everhart K, Crowell MD. Functional dyspepsia is associated with sleep disorders. *Clin Gastroenterol Hepatol.* 2011;9(5):410–414.
- Motivala SJ, Tomiyama AJ, Ziegler M, et al. Nocturnal levels of ghrelin and leptin and sleep in chronic insomnia. *Psychoneuroendocrinology.* 2009;34(4):540–545.
- Taheri S, Lin L, Austin D, et al. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med.* 2004;1(3):e62.
- Wren AM, Small CJ, Abbott CR, et al. Ghrelin causes hyperphagia and obesity in rats. *Diabetes.* 2001;50(11):2540–2547.
- Korbonits M, Goldstone AP, Gueorguiev M, et al. Ghrelin—a hormone with multiple functions. *Front Neuroendocrinol.* 2004;25(1):27–68.
- Schmid SM, Hallschmid M, Jauch-Chara K, et al. A single night of sleep deprivation increases ghrelin levels and feelings of hunger in normal-weight healthy men. *J Sleep Res.* 2008;17(3):331–334.
- Huang T, Mariani S, Redline S. Sleep irregularity and risk of cardiovascular events: the multi-ethnic study of atherosclerosis. *J Am Coll Cardiol.* 2020;75(9):991–999.
- Bei B, Wiley JF, Trinder J, et al. Beyond the mean: A systematic review on the correlates of daily intraindividual variability of sleep/wake patterns. *Sleep Med Rev.* 2016;28:108–124.
- Chen M-Y, Wang EK, Jeng Y-J. Adequate sleep among adolescents is positively associated with health status and health-related behaviors. *BMC Public Health.* 2006;6(1):1–8.