



Systematic review

A systematic review of interventions to increase physical activity and reduce sedentary behaviour following bariatric surgery

Jennifer D. James^{a,*}, Wendy Hardeman^b, Mark Goodall^c, Helen Eborall^d,
Victoria S. Sprung^e, Laura J. Bonnett^f, John P.H. Wilding^a

^a Department of Cardiovascular and Metabolic Medicine, Institute of Life Course and Medical Sciences, University of Liverpool, UK

^b School of Health Sciences, University of East Anglia, UK

^c Institute of Population Health Sciences, University of Liverpool, UK

^d Usher Institute, University of Edinburgh, UK

^e Sport and Exercise Sciences, Liverpool John Moores University, UK

^f Institute of Translational Medicine, University of Liverpool, UK

Abstract

Background Bariatric surgery promotes weight loss and improves co-morbid conditions, with patients who are more physically active having better outcomes. However, levels of physical activity and sedentary behaviour often remain unchanged following surgery.

Objectives To identify interventions and components thereof that are able to facilitate changes in physical activity and sedentary behaviour.

Eligibility Physical activity and/or sedentary behaviour must have been measured, pre and post intervention, in patients who have undergone bariatric surgery.

Study appraisal and synthesis methods : Four databases were searched with key-words. Two researchers conducted paper screening, data extraction and risk-of-bias assessment.

Results Twelve studies were included; eleven were randomised. Two were delivered presurgery and ten postsurgery; five found positive effect. Moderate to vigorous physical activity increased in three studies, two of which also found a significant increase in step count. The fourth found a significant increase in strenuous activity and the fifth a significant increase in metabolic equivalent of task/day and reduced time spent watching television.

Limitations Meta-analysis could not be conducted due to heterogeneity of outcomes and the tools used.

Conclusion and implications of key findings This review has identified interventions and components thereof that were able to provoke positive effect. However, intervention and control conditions were not always well described particularly in terms of behaviour change techniques and the rationale for their use.

Systematic review registration number PROSPERO (CRD42019121372)

Keywords: Physical activity; Sedentary behaviour; Bariatric surgery; Physiotherapy

* Corresponding author at: University of Liverpool, Clinical Sciences Centre, Aintree Site, Liverpool University Hospitals NHS Foundation Trust, Liverpool, L9 7AL, UK.

E-mail address: jennifer.james@liverpool.ac.uk (J.D. James).

URL: @Physio_JJ (J.D. James).

<https://doi.org/10.1016/j.physio.2021.10.002>

0031-9406/© 2021 The Authors. Published by Elsevier Ltd on behalf of Chartered Society of Physiotherapy. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contribution of the Paper

- Identifies interventions and behaviour change techniques that have been associated with increased physical activity in patients following bariatric surgery.
- Highlights the need for detailed reporting of control and intervention conditions, particularly with reference to behaviour change techniques and the rationale for their use.
- Highlights the need for research that focuses on ‘how’ to facilitate changes in physical activity and sedentary behaviour, rather than ‘what’ the physiological consequences of these changes are. This is particularly important for physiotherapists, who often lead on behaviour change interventions that aim to increase physical activity e.g. pre and rehabilitation, so that they can evidence base their clinical practice.

© 2021 The Authors. Published by Elsevier Ltd on behalf of Chartered Society of Physiotherapy. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Background

Systematic reviews and meta-analyses have identified that bariatric surgery is the most effective intervention for obesity, producing greater weight loss (mean difference -26 kg, 95% confidence interval $(-31$ to $-21)$ and a higher remission rate of diabetes (relative risk 22.1, 3.2–154.3), compared to non-surgical treatment [1] based on two years follow up. However, surgery is not a panacea and weight regain and recurrence of complications e.g. diabetes, can and does occur [2–4]. Consequently, there is an increasing focus on postsurgical lifestyle interventions, targeting physical activity [PA] to help patients to maintain the benefits of surgery [5]. PA is a behaviour, defined as “any bodily movement produced by skeletal muscles that uses energy” [6], and its benefits are well established; PA is protective against chronic conditions such as coronary heart disease and type 2 diabetes [7]. Current recommendations are for at least 150 minutes of moderate to vigorous intensity physical activity [MVPA] per week for adults [7–9], although there are no specific guidelines for patients following bariatric surgery to optimise weight and health status beyond these [10]. Despite this, there is some evidence that exercise (including strength training) following surgery may be beneficial in preserving non-fat mass [11], promote greater weight and fat mass loss, and improve physical fitness [12,13].

Sedentary behaviour [SB], sits on the PA continuum and is a separate and distinct behaviour of interest. It is defined as “any waking behavior characterized by an energy expenditure ≤ 1.5 metabolic equivalents (METs), while in a sitting, reclining or lying posture” [14]. This is different to inactivity, which occurs when an individual does not meet the PA guidelines. Thus it is possible to be both physically active and sedentary if one is able to achieve 150 minutes of MVPA per week through structured physical activity and also spends ‘prolonged periods’ of time sitting, e.g. due to a desk based job [15]. SB is of increasing interest to research and clinical communities and warrants attention in this review as it is associated with chronic health conditions including cardiovascular disease, type 2 diabetes and some cancers [16]. It is

particularly relevant to this patient group who are at increased risk of these conditions due to their obesity, which is likely to persist after bariatric surgery [17].

A recent systematic review [18] of prospective studies with at least 10 adults, ≥ 18 years undergoing bariatric surgery in which PA or physical function was measured pre and postsurgery, found an overall trend for PA to increase up-to 12 months following surgery. Step count increased between 3–6 months, but was accompanied with a reduction in objectively measured MVPA [18], meaning that 90% of patients were still not sufficiently active, according to the guidelines, and 24–29% (depending on the measure used to assess PA) became less active after surgery [6,19]. A recent meta-analysis of studies, which objectively measured SB found that patients were no less sedentary following bariatric surgery [20].

Given the potential benefits of increased PA and reduced SB, we aimed to synthesise the evidence for interventions that targeted either or both behaviours, in the context of bariatric surgery.

In behaviour change interventions (where specific behaviours such as PA and SB are targeted), behaviour change techniques (BCTs) are the ‘active ingredients’, defined as “observable, replicable and irreducible component of an intervention designed to alter or redirect causal processes that regulate behavior” [21]. Therefore, we expected these to be included in studies in this review. An example of a BCT according to the BCT taxonomy v1 (BCTTv1) [21], is ‘goal setting’, defined as ‘set or agree a goal defined in terms of the behaviour to be achieved’ e.g. ‘agree a daily walking goal (e.g. 3 miles) with the person and reach agreement about the goal’. As BCTs are the ‘active ingredients’ of behaviour change interventions, we decided to extract these as part of this review.

Studies were eligible for inclusion if they measured PA pre and postintervention using either self-reported or objective tools. This was a pragmatic decision as an initial scoping search identified that both were used and we did not want to exclude relevant studies that might have offered valuable information regarding intervention conditions.

Objectives

The aim of our study was to identify interventions and components thereof that would be able to facilitate changes in these behaviours after bariatric surgery. This information would be particularly useful for clinicians who understand the benefits in changing these behaviours, but are unsure of how best to support patients with this.

Our specific objectives were to synthesise the efficacy of the interventions and their components: characteristics, mode of delivery (groups or individual), outcome measures and behaviour change techniques [BCTs]. The review followed a preregistered protocol PROSPERO (CRD42019121372) and PRISMA guidelines see Fig. 1.

Data sources

Four databases were searched; Medline, CINAHL, Scopus and PsychINFO with search terms that related to the population, intervention and outcome (date of searches 23/11/2018–04/12/2018). In view of the time between searches and submission for publication, the same search strings were re-run in the databases for publications from 2018 to 2020 to update the review (final search 23/05/2020). Search terms included MeSH headings and key words based on bariatric surgery (such as bariatric, weight loss) and more specific terms that specified the surgery of interest and their synonyms (i.e. gastric bypass, roux-en y gastric bypass). Keywords associated with intervention and outcome such as programme, intervention and physical activity were searched after which the individual search strings were combined.

Two additional papers were identified during peer review. JJ screened all abstracts and titles, and VS a 10% random sample of these. All papers selected for full text review were independently screened for inclusion by JJ and VS who agreed on the papers to be included; reference lists of these were also reviewed for additional potential studies, which were subjected to the same processes.

Study selection

Any interventions that sought to increase PA or reduce SB were included, however, where the focus was principally on physiological effects, these were not, as these will be of less clinical value to clinicians whose focus is on how to support patients to change these behaviours.

Unlike previous reviews, ours focused on two target behaviours (PA and SB) and intervention efficacy in consideration with these, measured after bariatric surgery. We also considered components of the interventions — BCTs with a view to providing clinicians with the evidence for their use in addition to the effect on PA and SB.

Prospective primary research studies were included if they used either self-reported or objectively measured PA or SB, measured at baseline and at least one other time point after

surgery; the intervention may have commenced either prior to or following surgery. Single arm, pilot and feasibility studies were eligible for inclusion, as they may have offered valuable evidence to meet the objectives of this review. Where papers referenced earlier publications about the same study, these were retrieved and relevant data extracted.

There were no date limits to the search. Papers not published in English were excluded as were thesis and conference proceedings.

Data extraction and synthesis

A data extraction table was developed using the PICOD framework; it was piloted, reviewed and refined. Data extracted included study and participant characteristics, and intervention components e.g. supervised exercise. Other data extracted included mode of delivery, BCTs, duration, outcomes, and outcome measures; attendance, engagement and retention (as reported), intervention fidelity, and control group components. JJ and VS extracted data independently, after which they met to reach agreement without the need for intervention of a third reviewer. Study authors were contacted for additional information when required. JJ and WH independently extracted BCTs in all study arms using the BCCTv1 [21]. The confidence for their presence was indicated with + (probably present) and ++ (definitely present), after which they met to discuss and agree final codes.

All studies were assessed using the Cochrane Risk of Bias 2 tool [22] which considers risk of bias [ROB] in randomised studies in five separate domains: randomisation, deviations from the intended interventions, missing outcome data, measurement of the outcome and selection of the reported result. In this tool, the domain with the highest ROB judgement dictates the overall risk. As one of the studies was a single arm non-randomised pilot, only four domains of this tool were relevant [23]. ROB was independently assessed by JJ and LB; and by JJ and VS for studies retrieved in the updated search and from peer review of the manuscript.

Results

From a total of 3884 identified articles, forty-eight were assessed for full text review. Thirty-six were excluded due to: no postoperative measures of PA ($n = 11$), discussion paper or non-peer reviewed ($n = 8$), not an intervention study ($n = 6$), no baseline measures of PA ($n = 4$), PA not measured ($n = 3$), article not in English ($n = 3$) and, intervention did not seek to increase PA ($n = 1$). Data was extracted from twelve unique interventions (see Fig. 1); seven from the original search, three from the updated search and two obtained as a result of peer review.

Eight of the studies were randomised trials [24–31], one was quasi-experimental [32], two were randomised feasi-




Fig. 1. PRISMA flow diagram.

bility studies [33,34] and one was a single arm pilot study [23].

Eleven studies compared the intervention against a control group, the composition of which varied and included either generic PA input or advice [25,26,29–33], dietary focus [24,34], or were not explicitly described [27,28]. Sample size ranged from 10 [23] to 259 participants [24] and all participants were either considering or had undergone surgery. Participants in nine studies were mostly female (57–92%) [25,27–34]; the final three comprised females only [23,24,26]. Six of the studies had a mix of surgical procedures, the most common, were roux en-y-gastric bypass [RYGB] and sleeve gastrectomy [23,27,29,30,33,34]. Four of these included participants who had undergone gastric band surgery; this represented only a small proportion of the surgical types [27,29,33,34]. The remaining six studies comprised a single surgical type: RYGB [25,28,31,35] sleeve gastrectomy [32] or vertical banded gastroplasty [26]. See Table 1 for study design and baseline characteristics.

All studies measured levels of PA after surgery. This was stated as an intervention aim or hypothesis in six studies [25,26,29,30,32,33] and was measured in all twelve. SB was measured as an outcome in eight studies [24–28,31–33] one of which hypothesised that SB would reduce as a result of the intervention [32]. See Table 2 for details of the intervention and control conditions.

Nine of the twelve interventions included supervised activity or exercise, which was delivered by: PA specialists [30], exercise physiologist [31], qualified gym instructor [27], physiotherapist and exercise specialist [23], exercise and sports science professionals [32], physiotherapists [25], clinical dietitian or masters students in clinical nutrition [28] or was unspecified [33,34]. The duration and frequency of supervised exercise sessions ranged from once weekly for up-to 60 minutes [23,31,34] to four times weekly for 90 minutes [32]. This was delivered either on a one-to-one basis [27,34], in groups [23,28,30,32], a combination [25,33] or was not reported [31]. In these nine studies, participants were either expected or supported to change their PA behaviour outside of the sessions.

The final three interventions were counselling or discussion based; in two, participants were asked to increase MVPA and steps [29] or to increase their PA aiming for American College of Sports Medicine guidelines (150 minutes of MVPA per week) [26]. The final intervention involved participants discussing PA behaviours, focusing on the difficulties that might occur and how these might be managed in the future [35].

The duration of the intervention varied; the shortest was six weeks, delivered prior to surgery [29,37], whilst the longest was three years and was delivered during standard follow up visits [26].

Table 1

Study design and baseline characteristics. GB Gastric band, RYGB Roux en y gastric bypass. BMI data, mean (standard deviation), BMI data Baillot *et al.* 2016, 2018; median (25th and 75th quartiles).

Reference	Type of study	Research design	Total sample size	Control/intervention	% Female	Surgical type	BMI (kg/m ²) at baseline
Baillot <i>et al.</i> 2016, 2018 [30,36]	Evaluation	RCT	30	Control Intervention	75% 85%	RYGB 73% gastric sleeve 27% RYGB 62% gastric sleeve 38%	47.8 (40.3 to 54.0) 44.8 (42.1 to 53.0)
Bond <i>et al.</i> 2017 [29]	Sub-sample of evaluation study	RCT	36	Control Intervention	79% 91%	Total sample; RYGB <i>n</i> = 18 (50%), gastric banding <i>n</i> = 16 (44%), gastric sleeve <i>n</i> = 2 (6%)	44.4 (7.1) 46.7 (7.1)
Carnero <i>et al.</i> 2017 [31]	Evaluation	RCT	128	Control Intervention	86% 91%	All RYGB	44.4 (7.5) 45.8 (7.4)
Coleman <i>et al.</i> 2017 [33]	Feasibility	RCT	51	Control Intervention	84% 85%	RYGB <i>n</i> = 17 (68%), gastric sleeve <i>n</i> = 7 (24%), lap band <i>n</i> = 1 (4%) RYGB <i>n</i> = 21 (81%), gastric sleeve <i>n</i> = 4 (15%), lap band <i>n</i> = 1 (4%)	33.1 (5.8) 32.7 (5.8)
Hanvold <i>et al.</i> 2019 [28]	Evaluation	RCT	165	Control Intervention	76% 73%	All RYGB	31.0 (4.8) 30.8 (4.9)
Herring <i>et al.</i> 2017 [27]	Evaluation	RCT	24	Control Intervention	92% 92%	RYGB 33%, gastric sleeve 67%, RYGB 33%, gastric sleeve 58%, gastric band 8%	38.2 (6.1) 39.4 (4.3)
Jassil <i>et al.</i> 2015 [23]	Pilot	Single arm	10	Intervention	100%	Data provided for 8 participants only. RYGB <i>n</i> = 2, gastric sleeve <i>n</i> = 6 All sleeve gastrectomy	38.5 (7.2) 43.1 (4.5) 43.8 (5.3)
Jiménez-Loaisa <i>et al.</i> 2020 [32]	Evaluation	Quasi-experimental	40	Control Intervention	73% 82%	All vertical banded gastroplasty	Whole sample 49.5 (7.5)
Papalazarou <i>et al.</i> 2010 [26]	Evaluation	RCT	30	Control Intervention	100%	All RYBG	40.7 (4.6) 40.8 (4.5)
Sellberg <i>et al.</i> 2019 [24]	Evaluation	RCT	259	Control Intervention	100%	GB 67%, RYGB 33% GB 71%, RYGB 29%	41.0 (3.7) 47.3 (10.0)
Shah <i>et al.</i> 2011 [34]	Feasibility	RCT	33	Control Intervention	92% 90%	All RYGB	34.1 (5.4) 33.3 (6.2)
Stolberg <i>et al.</i> 2018 [25]	Evaluation	RCT	60	Control Intervention	~75% ~65%		

Table 2
Intervention and control conditions; attendance, engagement and retention.

Reference	Timing relative to surgery	Control	Intervention	Duration	Attendance, engagement and retention (as reported)
Baillot <i>et al.</i> 2018 [30]	Pre	Individual counselling every 6–8 weeks presurgery for at least 6 months, and postsurgery at months 3, 6, 9, and 12 with a dietitian and PA specialist. Plus, optional access to an educational group for PA, nutrition and psychological input related to weight management.	Three weekly 80-minute sessions: 10 minutes of warm-up, 30 minutes of endurance activity (at 55 to 85% of the heart reserve), 20 to 30 minutes of strength exercises with small equipment (dumbbells, elastic bands, medicine balls and sticks) and 10 minutes of a cool-down period, with monthly aqua gym session, until 2 weeks before surgery.	33 (8) weeks before surgery (range 27–51 weeks).	Intervention participants attended a median of 70 (45–90%) of the total recommended exercise sessions (3×/week) from the baseline of the PreSET until 2 weeks before surgery. Of the 30 participants randomised to either the intervention or control, 25 completed the study. 80 participants were randomised, 36 had surgery from which 31 (86%) completed postoperative follow up.
Bond <i>et al.</i> 2017 [29]	Pre	Participants were advised to begin exercising but did not receive any specific PA prescription, recommendations or strategies to facilitate this.	Six consecutive weekly individual face to face counselling sessions. PA was logged, monitored and a pedometer provided. Goals were set to increase bout-related walking minutes and steps per day relative to baseline. Counselling sessions reviewed self-monitoring records, goal progression, problem solving, teaching behaviour change strategies and developing action plans.	6 weeks.	
Carnero <i>et al.</i> 2017 [31]	Post	Participants received health education in 6 session, held once monthly which included lectures, discussion and information on topics including medication, nutrition and upper body stretching. Participants reported their PA habits at these sessions.	Participants received the same information as the control group, plus intervention. Weeks 1–4 participants exercise for as long as feasible (10–15 mins) at 60–70% of maximal heart rate. This progressed over 3 months to 120 minutes/week for the final three months of the intervention. Exercises included cycling or walking at home (stationary bike or treadmill or outdoors). Participants were encouraged to exercise for 30 minutes per day.	6 months	128 participants randomised: 66 to the intervention group and 62 to the control group. Retention was 91% ($n = 54$) in the intervention group and 95% ($n = 59$) in the control group. From a total of 96 participants, 23 were excluded from analysis due to insufficient PA data.
Coleman <i>et al.</i> 2017 [33]	Post	Weight assessment and phone calls which comprised counselling to encourage regular MVPA, although this did not contain and standardized recommendations. Phone calls and monitoring took place within the first two weeks post operatively, then at 2 and 6 months, and annually thereafter.	There were 2 phases; intervention and maintenance. The intervention phase comprised twice weekly 60-min group exercise classes comprising strength, flexibility and aerobic activities, plus at least 3 days per week of self-directed exercise. Daily pedometer with recording of steps and activities and weekly telephone counselling. The 6 months maintenance phase comprise once weekly exercise classes and once monthly telephone counselling.	12 months	Participation in the intervention was limited for ten out of the 25 (40%) assigned to the group due to a pre-existing condition. Of the remaining participants, 44% developed a condition during the programme which limited their participation with the intervention. Of 51 participants randomised to either the intervention or control, 43 completed the 12 months follow up

Table 2 (Continued)

Reference	Timing relative to surgery	Control	Intervention	Duration	Attendance, engagement and retention (as reported)
Hanvold <i>et al.</i> 2019 [28]	Post	Three follow up consultations in the first year with a dietitian or doctor, with an annual review thereafter for the next four years.	Sixteen group meetings comprising 12–15 participants lasting 2 hours. Participants were advised to reduce their sedentary time and to undertake ≥ 75 minutes/week higher intensity activity. Sessions included a 30-minute PA session with various activities including Nordic walking, climbing stairs, and strength training (weather dependent). An activity coach guided participants in ‘the use of Nordic walking and use of pedometer.	24 months	Attendance at the intervention varied from 35–84%, with the average number of sessions being attended was 8 (4) (out of a possible 16). Eight participants withdrew from the intervention group ($n = 85$), and fifteen withdrew from the control group ($n = 80$). Of 165 participants randomised, 142 completed the 24 months follow up.
Herring <i>et al.</i> 2017 [27]	Post	Participants in the control group continued with usual follow-up care. After their 12-week assessment, the control group also received the discharge advice session discussing the same topics. All participants were given an example exercise programme and progression (for example, home-based exercise, walking, swimming), along with the diet information sheet	Three sixty-minute gym sessions/week, of moderate intensity aerobic and resistance training for 12 weeks. Aerobic exercise training typically lasted 45 minutes; the first exercise programme was 35 minutes with a longer warm-up period and progressed to 45 min by the end of week 2. The warm up period reduced as participants fitness improved. Exercise sessions were progressive and personalised.	12 weeks	Participants in the intervention group attended a mean of 34 (3) sessions (out of a possible 36), equating to a 95% adherence. Of 24 participants randomised, 21 completed follow up.
Jassil <i>et al.</i> 2015 [23]	Post	No control group	Eight weekly sessions comprising 60 minutes of exercise followed by a 60-minutes group discussion on lifestyle education and nutritional-behavioural change sessions. The intervention was designed and delivered by a physiotherapist and exercise specialist and intensity of exercise varied according to individual functional capacity and increased progressively each week. Participants were encouraged to exercise to BORG 13 based on Borg’s Rate of Perceived Exertion scale.	8 weeks	Two patients attended all the sessions and the other patients attended seven ($n = 1$), six ($n = 1$), five ($n = 2$), four ($n = 1$), and three sessions ($n = 1$), respectively. The median number of sessions attended was six. Eight out of 10 patients completed follow up.
Jimenez-Loasisa <i>et al.</i> 2020 [32]	Post	Participants in the control group were given usual care recommendations focused on trying to maintain an active lifestyle after surgery (for weight loss and maintenance), but no specific advice was given.	The frequency of the intervention increased throughout the 6-month intervention period, beginning with 2 sessions/week for the first two months, then three sessions/week in the following two months, to four sessions/week for the final two. Sessions in the first two months were 60 minutes, increasing to 90 minutes in the following 4 months. The intervention involved cardiorespiratory and strengthening exercises, and sessions such as ‘body expression, dance, directed activities (aerobic, spinning etc.), beach and pool activities, core training, trekking and traditional Spanish games’. Participants were provided with home exercises which did not require ‘large resources or joining a gym’.	6 months	The attendance rate for the intervention group was 80% (68.1–88.9%). Retention for the intervention and control group at 13 months postsurgery (final assessment) was 89 ($n = 17$) and 75% ($n = 15$) respectively.

Table 2 (Continued)

Reference	Timing relative to surgery	Control	Intervention	Duration	Attendance, engagement and retention (as reported)
Papalazarou <i>et al.</i> 2010 [26]	Post	Postoperative dietetic assessment every week for the first three months, reducing to every other week for three months, and then monthly for six months. Participants were reviewed every three months in the second year, and every six in the third (total of 30 sessions). During these sessions general information was provided on adopting PA.	A patient-centred collaborative approach was used with behaviour modification techniques such as self-monitoring, self-evaluation, goal setting, reinforcement, stimulus control, and relapse prevention. Every session included nutritional education, dietary intake and physical activity with information provided on increasing PA. The intervention was delivered when participants attended for their usual care appointments.	36 months	All 30 participants (100%) attended all sessions and follow up.
Sellberg <i>et al.</i> 2018 & 2019 [24,35]	Post	Consultation with a dietitian, nurse or surgeon to discuss medical complications, weight loss and postsurgery diet, this took place ‘a few weeks, 6 months, 1 and 2 years after surgery’.	Weekly group sessions over a period of four weeks, comprising 4 different topics, the first of which concerned physical activity. Participants were encouraged to discuss how they wanted to behave in future specific situations, and the aim was to provide participants with strategies to manage future difficulties with PA behavior.	4 weeks	156 participants were allocated to the intervention group. Of these $n = 53$ did not attend any sessions, $n = 15$ attended 1, $n = 18$ attended 2, $n = 37$ attended 3 and $n = 33$ attended 4. Of 259 participants, 203 completed follow up.
Shah <i>et al.</i> 2011 [34]	Post	Participants from both groups had individual behavioural therapy focusing on stimulus control, eating behaviour and stress management.	Participants were asked to exercise on at least 5 days/week and attend supervised exercise sessions 1–2×/week. In addition, participants had behavioural therapy relating to exercise every 2 weeks, which included goal setting, self-monitoring, cognitive–behavioural strategies, and problem solving and relapse prevention.	12 weeks	33 participants were randomised; 12 to the control group and 21 to the intervention group. Four of control group provided baseline data only; three withdrew because they wanted to be in the intervention group. The fourth did not have time to participate. Five participants in the intervention group did not complete the study; four provided baseline and follow up data at 6 weeks but not at 12 weeks, and one provided data only at baseline. Time was given as the reason for dropping out of the study.
Stolberg <i>et al.</i> 2018 [25]	Post	Participants were given the clinics standard information about the importance of being physically active after RYGB.	Forty minutes of exercise, twice weekly for 26 consecutive weeks. The sessions comprised moderate intensity endurance and resistance exercises and were supervised by a physiotherapist. Participants also had free access to fitness centres during the intervention.	26 weeks	Nineteen of 32 participants (59%) allocated to the intervention group attended $\geq 50\%$ of the planned training sessions (deemed to be acceptable attendance). Out of the 60 participants randomised, 42 completed follow up.

See Table 2 for attendance, engagement and retention details.

BCTs to increase PA could be identified in all intervention, but not all control groups. The number of BCTs in intervention conditions varied from four [25] to nineteen [29,37]. The five most common were: Instruction on how to perform the behaviour (BCT 4.1) ($n = 10$), demonstration of the behaviour (BCT 6.1) ($n = 10$), goals setting behaviour (BCT 1.1) ($n = 9$), behavioural practice/rehearsal (BCT 8.1) ($n = 9$) and action planning (BCT 1.4) ($n = 7$). One BCT, prompts and cues (BCT 7.1) was identified as being used to reduce SB in the intervention group, although SB was not measured as an outcome in the study [29].

Information regarding control group conditions ($n = 11$) was generally lacking, (see Table 2) making it difficult to extract details. From the eleven control groups, five had identifiable BCTs: goal setting (BCT 1.1) ($n = 3$), information about health consequences (BCT 5.1) ($n = 2$), problem solving (BCT 1.2) ($n = 1$) and self-monitoring behaviour (BCT 2.3) ($n = 1$), with two being the maximum number of identifiable BCTs identified in any one group [27,32] (see Table 3 for BCTs identified in each of the studies).

None of the authors powered their studies to detect a change in PA or SB, specifically in the time period following bariatric surgery although postoperative PA, SB or both, were measured by all in this review. Six studies presented aims or hypothesis related to PA, five of which sought to assess if the intervention would increase this [24,25,30,32,33] with one identifying bout related MVPA and steps specifically as outcomes of interest [29]. The remaining six measured PA as one of their study outcomes [23,26–28,31,34].

Self-report and objective measures were used to assess PA; four studies used self-report only [23,26,28,30], four used self-report and objective tools [25,27,33,34] and four used objective measures only [24,29,31,32] (see Table 4 for outcome details).

Of the eight studies that used self-report, validated questionnaires were used in five [26–28,30,34], whilst the others either adapted or used questions from existing questionnaires [23,25,33]. Accelerometers were used in six studies and were worn either on the arm, at the mid-axillary line at the superior aspect of the iliac crest, or the hip [24,25,27,29,31,32]; pedometers were used in two [33,34].

SB and sitting time was measured in eight studies, three used self-report [26,28,33], and five measured this objectively via accelerometry [25,27,31,32,35].

Five of the interventions resulted in statistically significant improvements in PA compared to the control group or baseline (where there was no control). Bond *et al.* [29] found that intervention participants had greater increases than the control group in bout related MPVA minutes/day at postintervention ($P = 0.016$) but not at follow up (six months postsurgery) ($P = 0.15$). Steps/day also differed between the groups at postintervention ($P = 0.031$) and follow up ($P = 0.024$).

Herring *et al.* [27] found a mean increase in MVPA in the intervention group between baseline and follow up, this decreased in the control group, resulting in a significant difference between the two ($P = 0.043$), however the difference was no longer significant 12 weeks thereafter, following a period of no intervention. Jassil *et al.* [23] reported a significant increase in strenuous PA minutes/week from a baseline of zero to 44 (SD 49) ($P < 0.05$) and a non-statistically significant reduction ($P = 0.310$) in MVPA following eight weeks of intervention compared to baseline (no control).

Shah *et al.* [34] reported a significant increase in steps in the intervention group from ~5500 to almost 10000 steps/day. There was a statistically significant group by week interaction ($P = 0.03$) and within group change in the intervention group ($P < 0.001$). Self-reported time spent in moderate PA increased by more than three times for the intervention group but remained the same in the control group when compared to baseline. Again, there was a significant group by week interaction ($P = 0.02$) and within group change in the intervention ($P < 0.001$) but not in the control group. There were no changes in self-reported light PA in either of the groups, and time spent in vigorous PA remained at zero for both groups at all time points.

Postintervention, Papalazarou *et al.* [26] reported a significant increase in PA ($P = 0.001$) and a reduction in SB measured via time spent watching (hours/day) ($P = 0.039$), compared to the control group. The remaining seven studies reported that there were no significant differences between the intervention and control group following the intervention on either PA or SB.

None of the authors commented on fidelity of delivery i.e. whether the intervention was delivered as planned; although one did report that instructors undertook training prior to delivering the intervention, which included peer review and feedback [32].

Attendance, engagement and retention varied across the studies, as did the level of detail provided by authors. Baillot *et al.* [30] delivered their intervention presurgery reporting a mean duration of 32.6 weeks, with variation between participants due to anticipated surgical dates. Median participant attendance was 70 (45–90%) of the total sessions. Of an initial 30 participants (15 in each group), 25 completed the study with 14 and 12 in the intervention and control group respectively. Bond *et al.* [37] randomised 80 participants, with 42 and 38 in the intervention and control group respectively. Thirty-three out of 40 (83%) allocated to the intervention completed all six sessions. Of the 80 who were randomised, 36 went on to have surgery ($n = 22$ in intervention, $n = 14$ in control), with final follow up data available for 31.

Carnero *et al.* [31] randomised 66 patients to the intervention group, six of whom did not complete the intervention; five reported time constraints with the sixth lost to follow up. Of the 60 participants in this group, 14 were excluded from analysis due to insufficient PA data.

Coleman *et al.* [33], randomised 51 participants to the control or intervention. In the first six months, intervention

Table 3
BCTs identified in each study.

Reference	Control/Intervention	BCT identified and confidence (+, ++) All coded for PA unless indicated. No BCTs for SB were identified.
Baillot <i>et al.</i> 2018 [30]	Control Intervention	None 2.4 Self-monitoring of outcomes of behaviour + 2.6 Bio feedback ++ 3.1 Social support unspecified + 4.1 Instruction ++ 8.1 Behavioural practice rehearsal ++ 8.7 Graded tasks ++ 11.2 Reducing negative emotions ++
Bond <i>et al.</i> 2017 [29]	Control Intervention	1.1 Goal setting ++ 1.1 Goal setting behaviour ++ 1.2 Problem solving + 1.4 Action planning ++ 1.5 Review goals behaviour + 1.8 Behavioural contract+ 2.2 Feedback on behaviour ++ 2.3 Self-monitoring behaviour + 3.1 Social support unspecified ++ 4.1 Information on how to do the behaviour+ 5.3 Information about health consequences ++ 6.1 Demonstration of behaviour + 7.1 Prompts and cues+ (PA & SB) 8.2 Behaviour substitution + 8.7 Graded tasks ++ 9.2 Pros and cons + 11.2 Reducing negative emotions + 12.5 Adding objects to the environment ++ 14.7 Reward incompatible behaviour +
Carnero <i>et al.</i> 2017 [31]	Control Intervention	2.3 Self-monitoring behaviour + 1.4 Action planning ++ 2.1 Monitoring of behaviour by others without feedback ++ 2.2 Feedback on behaviour ++ 2.3 Self-monitoring of behaviour ++ 2.6 Biofeedback ++ 4.1 Instruction on how to perform the behaviour ++ 6.1 Demonstration of the behaviour ++ 8.1 Behavioural practice/rehearsal ++ 8.7 Graded tasks ++
Coleman <i>et al.</i> 2017 [33]	Control Intervention	None 1.1 Goal setting ++ 1.4 Action planning ++ 2.2 Feedback on behaviour + 2.3 Self-monitoring behaviour + 3.1 Social support unspecified + 4.1 Instruction ++ 6.1 Demonstration of the behaviour (or modelling) ++ 8.1 Behavioural practice ++ 12.5 Adding objects to the environment ++
Hanvold <i>et al.</i> 2019 [28]	Control Intervention	None 1.1 Goal setting ++ 1.4 Action planning ++ 2.3 Self-monitoring ++ 4.1 Instruction on how to perform the BH + 5.1 Information about health consequences + 5.3 Information about social and environmental consequences ++ 8.1 Behavioural goals (PA & SB) ++ 8.3 Habit formation + 12.5 Adding objects to the environment ++
Herring <i>et al.</i> 2017 [27]	Control Intervention	1.1 Goal setting behaviour + 1.2 Problem solving + 1.1 Goal setting behaviour ++ 1.2 Problem solving ++ 4.1 How to do behaviour + 8.1 Behavioural practice/ rehearsal ++ 8.7 Graded tasks ++
Jassil <i>et al.</i> 2015 [23]		1.1 Goal setting behaviour ++ 1.4 Action planning ++ 1.5 Review of behavioural goals ++ 2.1 Monitoring of behaviour without feedback + 2.3 Self-monitoring behaviour ++ 4.1 Instruction on how to perform the behaviour ++ 8.1 Behavioural practice ++ 8.7 Graded tasks ++ 12.5 Adding objects to the environment ++
Jimenez-Loaisa <i>et al.</i> 2020 [32]	Control Intervention	1.1 Goal setting BH ++ 5.1 Information about health consequences ++ 1.1 Goal setting BH ++ 2.2 Feedback on BH ++ 4.1 Instruction on how to perform the BH ++ 5.1 Information about health consequences ++ 8.1 Behavioural practice/ rehearsal ++ 10.4 Social reward +
Papalazarou <i>et al.</i> 2010 [26]	Control Intervention	None 1.1 Goal setting behaviour ++ 1.2 Problem solving ++ 1.7 Review outcome goals ++ 2.4 Self-monitoring outcome ++ 10.9 Self-reward++
Sellberg <i>et al.</i> 2018 & 2019 [24,35]	Control Intervention	None 1.1 Goal setting BH ++ 1.2 Problem solving ++ 1.4 Action planning ++ 8.3 Habit formation + 11.2 Reduce negative emotions + 13.3 Incompatible beliefs +
Shah <i>et al.</i> 2011 [34]	Control Intervention	None 1.1 Goal setting behaviour ++ 1.2 Problem solving ++

Table 3 (Continued)

Reference	Control/Intervention	BCT identified and confidence (+, ++) All coded for PA unless indicated. No BCTs for SB were identified.
		1.3 Goal setting outcome ++ 1.4 Action planning ++ 2.1 Monitoring of behaviour by others without feedback ++ 2.3 Self-monitoring of behaviour ++ 2.7 Feedback on outcome of behaviour ++ 4.1 Instruction on how to perform the behaviour ++ 6.1 Demonstration of the behaviour ++ 8.1 Behavioural practice/ rehearsal ++ 8.7 Graded tasks ++
Stolberg <i>et al.</i> 2018 [25]	Control Intervention	5.1 Information about health consequences+ 2.1 Monitoring of behaviour by others without feedback 4.1 Instruction on how to perform the behaviour ++ 8.1 Behavioural practice ++ 12.5 Adding objects to the environment +

participants attended 56% of all classes offered (average of one class per week) and reported exercising 3.1 (1.7) days per week outside of this. Engagement was limited for some, who were unable to undertake the intervention as planned. Ten (40%) were limited by a pre-existing condition (not detailed) and of the remaining participants, 15 (44%) developed a condition, which limited their participation. During the ‘maintenance phase’ in the following six months, attendance reduced to 32% of classes/week and participants reduced their exercise outside of the classes to 2.9 (± 1.8) days a week. In this later period, contact with the counsellor increased from 69% in the first six months to 93%. Postintervention measures were completed by 23 and 21 participants from the control and intervention group respectively with final follow up data for 23 and 20 participants respectively. Hanvold *et al.* [28] randomised 165 participants; 85 to the intervention and 80 to the control group. Average intervention session attendance varied from 35–84%, (8 ± 4 sessions from a possible 16 over 2-years). Follow up measures were available for 77 and 65 participants in the intervention and control groups respectively.

Herring *et al.* [27] randomised 24 participants equally. Over the 12-week intervention period, mean attendance was 34.2 sessions out 36 (95%). Final follow up data was available for 21 participants ($n = 11$ and $n = 10$ in intervention and control respectively).

Jassil *et al.* [23] recruited a single cohort of 10 patients; eight completed the intervention over an eight-week period, median attendance was six out of eight sessions (75%). Jiménez-Loaisa *et al.* [32] randomised forty participants equally, and reported an average intervention attendance rate of 80% (range 68–89%). Follow up data was available for 17 and 15 participants in the intervention and control groups respectively.

Sellberg *et al.* [24] provided a comprehensive breakdown for participant attendance. One hundred and fifty-six out of 259 were randomised to the intervention group; $n = 33$ attended zero sessions, $n = 13$ attended one, $n = 13$ attended two, $n = 31$ attended 3 and $n = 30$ attended four. One hundred and twenty intervention participants, and 83 out of 103 from the control group completed the study follow up.

Shah *et al.* [34] recruited 33 participants, with 21 randomised to the intervention and 12 to the control group. Four withdrew from the control group after baseline measures, three because they wanted to be in the other group, and one due to time constraints. From the 21 participants in the intervention group, five withdrew with four providing follow up data.

Stolberg *et al.* [25] randomised 60 participants resulting in 32 in the intervention and 28 in the control group. In the 26-week intervention period, 19 out of 32 participants (59.4%) attended $\geq 50\%$ of planned exercise sessions (considered to be compliant). Data for 27 and 25 participants in the intervention and control group respectively was available for analysis at 12 months, reducing to 22 and 20 for intervention and control group respectively at 24-month measures.

Papalazarou *et al.* [26] reported 100% attendance at each of the sessions, which was delivered when participants attended for their usual care and was attributed to the intervention being intense and delivered on a 1:1 basis (correspondence with author). ROB was assessed using the Cochrane ROB2 tool [22] (see Table 5), because there are no other suitable tools for behavioural interventions evaluated in randomised studies. All studies were judged to have ‘some concerns’. The reasons for this included the absence of a published protocol ahead of the study, lack of masking of participants and the use of self-reported tools to measure PA or SB.

Discussion

This review identified twelve studies and found mixed evidence about intervention effects on increasing PA and/or reducing SB following bariatric surgery. Detail regarding intervention and control conditions in particular varied, and was often lacking in regards to chosen BCTs and the rationale for their use. This is a significant limitation; by definition BCTs were expected in all studies included in this review as they are the ‘active ingredients’ [21] in behaviour change interventions, and all the studies included in this review sought to change at least one of the two target behaviours.

Table 4

Outcomes, follow up and results.

Reference	PA measurement tool and units	Post intervention or follow up period	Results
Baillot <i>et al.</i> 2018 [30]	Self-report, IPAQ-SF: METs	Follow up 1 year following surgery (intervention presurgery)	Changes in the self-reported levels of PA were not significantly different between groups. According to accelerometry, the intervention group had a greater number of steps, light and moderate PA compared to the control group one-year postsurgery; however, there was no objective baseline measure of this and the authors acknowledge poor concordance between self-report and objective measures.
Bond <i>et al.</i> 2017 [29]	Objective, accelerometer: MVPA mins and steps/day	Follow up 6 months after surgery (intervention presurgery).	The intervention group had a greater mean increase in bout related MVPA of 22.0 (mins/day) increasing from 4.3 to 26.3 (minutes/day) ($P = 0.016$), compared to the control group whose mean MVPA increased by 1.0 (minutes/day) from a baseline of 10.4 (minutes/day). At follow up, mean MVPA in the intervention group was 28.7 minutes/day, and 18.5 in the control groups; statistical significance was no longer maintained ($P = 0.15$). The intervention group increased their mean steps/day by 2793 from a baseline of 5163 to 7950 (corrected $P = 0.031$), which slightly reduced at follow up to 7870 ($P = 0.024$) but was still statistically significant compared to the control group who increased their steps from a baseline of 5069 to 5601 at postintervention but reduced to 5087 at follow up.
Carnero <i>et al.</i> 2017 [31]	Objective, accelerometer: METs, steps	Post intervention: For the last week of the intervention which was 6 months in duration.	Both groups significantly increased their steps per day and time spent in light, moderate and vigorous PA and reduced their sedentary time, but there was no difference between the groups.
Coleman <i>et al.</i> 2017 [33]	Self-report via questionnaire with measures of sedentary activity and questions from Behavioural Risk Factor Surveillance survey: MVPA and sedentary activity minutes/day. Pedometer: steps/day	Post intervention (after 6 months of intervention), repeated 6 months after the maintenance phase.	No differences between the groups in self-reported PA or steps/day.
Hanvold <i>et al.</i> 2019 [28]	Self-report, 7-day physical activity recall questionnaire: time spent on low, moderate and high PA.	Post intervention.	No differences between the groups in changes in PA level or time spent on different activities.
Herring <i>et al.</i> 2017 [27]	Self-report, IPAQ-SF: METs. Objective, accelerometer: sedentary, light PA, MVPA and step count	Post intervention immediately following the intervention and follow up 12 weeks thereafter.	Post intervention, MVPA increased by 10.5 (SD 9.2) (mins/day) in the intervention from a baseline of 28.3 (SD 24.0) (minutes/day) ($P = 0.043$). MVPA in the control group reduced by -1.5 (SD 14.5) from a baseline of 29.7 (SD 18.7) mins/day). At follow up, mean MVPA in the intervention group was 7.5 (SD 19.8) mins/day above baseline. This reduced by -3.4 (SD 16.2) in the control group from the baseline, and the difference between the groups was no longer statistically significant.
Jassil <i>et al.</i> 2015 [23]	Self-report, questionnaire adapted from National Audit of Cardiac Rehabilitation: MVPA minutes	Post intervention; one month after the intervention.	Authors report mean time spent on strenuous activity was 44 (49) minutes, from a baseline of zero ($P < 0.05$), no significant differences to moderate intensity PA.
Jiménez-Loasia <i>et al.</i> 2020 [32]	Objective, accelerometer: time spent in sedentary, light, moderate to vigorous PA.	Post intervention; six months after the intervention and again six months thereafter.	No increases in PA in the intervention group, at any time point when compared with the control group.
Papalazarou <i>et al.</i> 2010 [26]	Self-report, Harokopio Physical Activity Questionnaire: METs and time spent watching television (hours/day)	Post intervention	PA increased from 1.26 (SEM = 0.01) to 1.62 (SEM = 0.04) in the intervention group and from 1.30 (SEM = 0.20) to 1.34 (SEM = 0.03) in the control group; statistically significant difference between the groups ($P = 0.001$). SB reduced in the intervention group from a baseline of 2.8 (SEM = 0.28) (hours/day) to 0.80 (SEM = 0.14) and from 2.46 (SEM = 0.35) to 2.30 (SEM = 0.33), statistically significant difference between the groups ($P = 0.039$)

Table 4 (Continued)

Reference	PA measurement tool and units	Post intervention or follow up period	Results
Sellberg <i>et al.</i> 2019 [24]	Objective, accelerometer: sedentary, light and moderate-vigorous PA.	1-year post intervention	No effects of the intervention were found.
Shah <i>et al.</i> 2011 [34]	Self-report, Seven-day physical activity recall Objective, pedometer steps	Mid-intervention (6 weeks) and postintervention.	Self-reported PA increased by more than three times in the intervention group but was unchanged in the control group. Significant group by week interaction in the intervention group ($P=0.02$) and within group change ($P<0.001$). Steps/day increased from about 5500 at baseline to nearly 10,000 in the intervention group, and only slightly in the control group. Group by week interaction ($P=0.03$) and within group change in the intervention group were statistically significant ($P<0.001$).
Stolberg <i>et al.</i> 2018 [25]	Self-report, Recent Physical Activity Questionnaire + authors added 5 questions to home activities section. Objective, accelerometer: METs	Postintervention and follow up 12 months after the intervention	No significant difference between the groups at post intervention or follow up.

Table 5

ROB for each of the studies. * refers to judgement for objective and self-report measures respectively.

Reference	Domain 1: ROB from randomisation process	Domain 2: ROB due to deviations from intended intervention	Domain 3: ROB due to missing outcome data	Domain 4: ROB in measurement of the outcome	Domain 5: ROB in selection of the reported result	Overall ROB judgement according to guidance
Baillot <i>et al.</i> 2018 [30]	Low	Low	Low	High	Some concerns	High
Bond <i>et al.</i> 2017 [29]	Low	Some concerns	Low	Low	Some concerns	Some concerns
Carnero <i>et al.</i> 2017 [31]	Low	Some concerns	Low	Low	Low	Some concerns
Coleman <i>et al.</i> 2017 [33]	Low	High	Some concerns	Low/ High *	Some concerns	High
Hanvold <i>et al.</i> 2019 [28]	Low	Some concerns	Low	High	Some concerns	High
Herring <i>et al.</i> 2017 [27]	Low	Some concerns	Low	Low/High*	Some concerns	Some/high*
Jassil <i>et al.</i> 2015* [23]	N/A	Some concerns	Low	High	Some concerns	High
Jimenez-Loaisa <i>et al.</i> 2020 [32]	Low	Some concerns	Low	Low	Some concerns	Some concerns
Papalazarou <i>et al.</i> 2010 [26]	Low	Low	Low	High	Some concerns	High
Stolberg <i>et al.</i> 2018 [25]	Low	Some concerns	Low	Low/High*	Some concerns	Some/high*
Shah <i>et al.</i> 2011 [34]	Low	Some concerns	Low	Low/ High	Some concerns	Some/high*
Sellberg <i>et al.</i> 2019 [24]	Low	Some concerns	Low	Low	Some concerns	Some concerns

The lack of detail regarding control conditions made it difficult to extract BCTs and as a result limited our ability to draw comparisons between conditions. Five out of the twelve studies reported evidence for a positive effect, but this needs to be interpreted with caution as studies were not statistically powered to detect changes in PA or SB after bariatric surgery. It is this specific time period that is of interest in this review in line with a research recommendation by NICE,

which seeks to identify if postoperative intervention programmes improve outcomes including weight and weight loss maintenance following surgery [5]. Only one study found a significant reduction in SB, measured as time watching television [26].

Data was extracted for both the intervention and control groups, although they were not always fully described, which necessitated contact with the study authors. Use of the

TIDier checklist would have been helpful and should be considered by researchers, as it would more confidently enable researchers and clinicians to appraise the evidence, decide upon its relevance, and replicate effective interventions [38].

Components of the interventions varied; some involved repeated supervised exercise sessions over a prolonged period, whilst others were counselling based. Regardless, all of the interventions directed participants to increase their PA outside of the sessions, thereby requiring participants to change their behaviour.

Authors stated that they used ‘behavioural modification techniques’ [26], ‘the most enduring strategies for behaviour change’ [33] and employed ‘self-regulatory behaviour change techniques’ [23], but the target behaviour to which these strategies or techniques related to was not always clear.

BCTs associated with initiation of PA were present in four of the five interventions that showed positive effect: demonstration of the behaviour [23,27,29], behavioural practice/rehearsal [23,27] and graded tasks [23,27,29,34]. BCTs associated with maintenance of PA [39], were present in five of the interventions with significant positive effects: action planning (BCT 1.4) [23,29,34], how to perform the behaviour (BCT 4.1) [23,27,29,34] prompts and cues (BCT 7.1) [29], behavioural practice and rehearsal (BCT 8.1) [23,27,34], graded tasks (BCT 8.7) [23,27,29,34] and self-reward (BCT 10.9) [26]. Thus, this review provides preliminary support the use of these BCTs from primary research in this patient group.

Of the seven studies that measured SB [24–28,32,33] only one found a reduction following the intervention [26]. However, as BCTs could not be coded for this target behaviour, this review is unable to add to the evidence base for interventions to reduce SB after bariatric surgery. Drawing from the wider literature, meta-analytical evidence suggests that promising BCTs may include: self-monitoring [40], social comparison, problem solving, demonstration of the behaviour, goal setting (behaviour), behaviour substitution, and habit reversal [41].

Future research that seeks to change PA or SB should ensure that the rational for and descriptions of BCTs are clear. The latter could be achieved by using the BCTT[v1] [21]. Researchers should also state whether the intervention aims to support initiation or maintenance of PA or SB as there is evidence that effective BCTs may differ [39].

Both objective and self-reported tools were used to measure PA and SB. In eight of the 12 studies questionnaires were used. Compared with objective measuring tools, questionnaires are cheap, more easily available, reliable, but their validity has been questioned as they are susceptible to bias [20]. Where self-reported and objectively measured PA has been compared, validity has been described as ‘moderate at best’ [42,43] and is supported by a study in this review [27]; there was a significant difference between the groups according to accelerometer data but not the IPAQ-SF. It is important to note that self-reported outcomes allow only a proxy measure of time spent being physically active, whereas objective

tools quantify PA [44], but this must be balanced against available clinical resources.

We found that interventions that required the greatest time commitments in terms of participant attendance and contact hours showed lower session attendance [30] and engagement [33]. Conversely interventions with positive effect were either relatively short in duration six [29], eight [23] or 12 weeks [27,34] or combined with participants’ usual care appointments [26] over a longer period of time. Sellberg *et al.* [24] reflected on participant attendance and suggested that time commitments required for the intervention (4 sessions of 1.5 hours duration) might not be suited to this patient group and alternative delivery methods should be considered. Time was cited by other authors as a factor in participant withdrawal [31,34] as was the unacceptability of randomisation to control conditions [27,34] and these issues should be considered in future research. It might be more appropriate to randomise sites rather than individuals to reduce the risk of participant withdrawal for this reason. Participant burden must be reduced as much as possible with alternative modes of delivery considered. For example, publicly available online platforms e.g. Skype or Zoom could be used. Although these are non-traditional methods of delivering interventions, their use has increased during the Covid-19 pandemic as they ensure ‘social distance’, and it would also help to protect this patient group who are at greater risk of serious disease from Covid-19 [45]. Both of these issues are illustrated by primary qualitative research with participants from cardiac rehabilitation, which found that reasons for participants’ disengagement can be due to their perception of the programme and intervention characteristics, including their [in]convenience [46].

A limitation of the studies in this review are that none reported on fidelity — the extent to which interventions were delivered as intended. As a result, it can be challenging to confidently attribute positive effect or lack of thereof to the intervention as this could be due to [sub-optimal] delivery rather than the intervention itself [47–49]. Strategies to promote treatment fidelity such as assessment of adherence to intervention protocols in behavioural interventions and assessment of fidelity mean that validity and reliability could be more confidently assured [50]. This is particularly important in interventions where there is potential for variation, for example in multicentre studies where there are different facilitators delivering the intervention. Future complex intervention studies which seek to change behaviour should include fidelity assessments to assess if the intervention has been delivered as planned and if participants have the necessary skills to be able to implement the intervention even if it is delivered faithfully [50,51].

Although every study was deemed to have at least some concerns for bias, this does need to be seen in the context of interventions where either it is not possible to mask participants to their allocation, or because the participants themselves are the assessor of the outcome due to the use of self-reported outcome measures [52]. The ROB judgements

in this review are consistent with those in other behaviour change studies, and a reflection of the tool used which is more suited to bio-medical trials, where variables can be controlled to provide internal validity to infer causality and are not necessarily reflective of real-life situations [52,53]. The lack of a valid and reliable risk of bias tool for behaviour change interventions evaluated in randomised studies meant that the team chose the Cochrane Risk of Bias ROB2 tool [22]. Although this tool is not ideally suited to behaviour change research, it was able to highlight that future studies could reduce their risk of bias by using objective rather than self-reported measures which would help to strengthen evidence of efficacy and effectiveness.

Strengths and limitations

A strength of this review is that the methods were robust; authors were contacted as required to obtain missing or additional information and two members of the research team executed each stage; screening, data extraction and ROB assessment. Rather than focusing solely on the efficacy of the interventions on PA and SB, this review also considered how healthcare professionals might facilitate changes in the two target behaviours using BCTs. This is particularly valuable to clinicians whose role, either as specialists or as part of a wider public health strategy is to support patients to increase PA and/or reduce SB.

A limitation is that grey literature was not searched and articles that were not published in English were excluded. Meta-analysis could not be conducted due to the heterogeneity of the outcomes and their measurements used in this review.

Conclusion and implications of key findings

In conclusion, this review identified evidence from five interventions that were able to provoke increases in PA but only one demonstrated a reduction in SB. Importantly none of these studies were powered with PA or SB as the primary outcome measure, measured postsurgery. It is this specific time-period, which is the focus of the review as there is evidence that changes in the two target behaviours of PA and SB after surgery might have a positive impact on patients' postsurgical and longer-term outcomes.

Details regarding intervention and control conditions in particular varied and the choice of and rationale for BCTs was not always clear. This is an important finding from the review; there is robust evidence for the physiological effects of increasing PA and reducing SB, which clinicians can use as evidence to support the aim of their treatment plans, but a lack of evidence for how to facilitate changes in these two target behaviours using BCTs. Any study that seeks to change a target behaviour should be explicit with regards to the rationale, chosen BCTs and whether the aim is to initiate or maintain

behaviour change. Therefore, there is a need for high-quality studies evaluating theory and evidence-based interventions to promote PA and reduce SB, that clinicians can use to help patients to change these target behaviours and optimise their postsurgical outcomes.

Funding: Jennifer James is funded by a National Institute for Health Research (NIHR) [ICA Clinical Doctoral Research Fellowship] for this research project. This publication presents independent research. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Conflict of interest: JJ, WH, VSS, LJB, HE, MG have no declarations to declare.

JW reports grants from National Institute for Health Research, during the conduct of the study; grants, personal fees and other from AstraZeneca, other from Astellas, personal fees and other from Boehringer Ingelheim, personal fees and other from Mundipharma, personal fees and other from Napp, grants, personal fees and other from Novo Nordisk, other from Lilly, other from Sanofi, grants, personal fees and other from Takeda, other from Wilmington Healthcare, other from Rhythm Pharmaceuticals, outside the submitted work.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.physio.2021.10.002>.

References

- [1] Gloy VL, Briel M, Bhatt DL, Kashyap SR, Schauer PR, Mingrone G, et al. Bariatric surgery versus non-surgical treatment for obesity: a systematic review and meta-analysis of randomised controlled trials. *BMJ* 2013;347:f5934.
- [2] Magro DO, Geloneze B, Callejas F, Pareja JC. Long-term weight regain after gastric bypass: a 5-year prospective study. *Obes Surg* 2008;648–51.
- [3] Cooper TC, Simmons EB, Webb K, Burns JL, Kushner RF. Trends in weight regain following roux-en-Y gastric bypass (RYGB) bariatric surgery. *Obes Surg* 2015;25(8):1474–81.
- [4] DiGiorgi M, Rosen DJ, Choi JJ, Milone L, Schroppe B, Olivero-Rivera L, et al. Re-emergence of diabetes after gastric bypass in patients with mid- to long-term follow-up. *Surg Obes Relat Dis* 2010;6(3):249–53.
- [5] NICE. Obesity: identification, assessment and management (CG189); 2014.
- [6] WHO. Global recommendations of physical activity for health. World Health Organisation; 2011.
- [7] Officers UCM. Physical Activity Guidelines; 2019.
- [8] Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011;43(7):1334–59.

- [9] World Health O. WHO Guidelines on physical activity and sedentary behavior; 2020.
- [10] Mechanick JI, Youdim A, Jones DB, Timothy Garvey W, Hurley DL, Molly McMahon M, *et al.* Clinical practice guidelines for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient — 2013 update: Cosponsored by American Association of Clinical Endocrinologists, the Obesity Society, and American Society for Metabolic & Bariatric Surgery. *Surg Obes Relat Dis* 2013;9(2):159–91.
- [11] Hassannejad A, Khalaj A, Mansournia MA, Rajabian Tabesh M, Alizadeh Z. The effect of aerobic or aerobic-strength exercise on body composition and functional capacity in patients with BMI ≥ 35 after bariatric surgery: a randomized control trial. *Obes Surg* 2017;27:2792–801.
- [12] Bellicha A, Ciangura C, Poitou C, Portero P, Oppert JM. Effectiveness of exercise training after bariatric surgery—a systematic literature review and meta-analysis. *Obes Rev* 2018;19(11):1544–56.
- [13] Ren Z-Q, Lu G-D, Zhang T-Z, Xu Q. Effect of physical exercise on weight loss and physical function following bariatric surgery: a meta-analysis of randomised controlled trials. *BMJ open* 2018;8(10):e023208.
- [14] Tremblay MS, Aubert S, Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE, *et al.* Sedentary Behavior Research Network (SBRN) — Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act* 2017;14(1):75.
- [15] van der Ploeg HP, Hillsdon M. Is sedentary behaviour just physical inactivity by another name? *Int J Behav Nutr Phys Act* 2017;14(1):142.
- [16] Owen N, Salmon J, Koohsari MJ, Turrell G, Giles-Corti B. Sedentary behaviour and health: mapping environmental and social contexts to underpin chronic disease prevention. *Br J Sports Med* 2014;48(3):174–7.
- [17] Strain GW, Kolotkin RL, Dakin GF, Gagner M, Inabnet WB, Christos P, *et al.* The effects of weight loss after bariatric surgery on health-related quality of life and depression. *Nutr Diabetes* 2014;4(9), e132-e.
- [18] Herring LY, Stevinson C, Davies MJ, Biddle SJ, Sutton C, Bowrey D, *et al.* Changes in physical activity behaviour and physical function after bariatric surgery: a systematic review and meta-analysis. *Obes Rev* 2016;17(3):250–61.
- [19] King WC, Hsu JY, Belle SH, Courcoulas AP, Eid GM, Flum DR, *et al.* Pre- to postoperative changes in physical activity: report from the longitudinal assessment of bariatric surgery-2 (LABS-2). *Surg Obes Relat Dis* 2012;8(5):522–32.
- [20] Barbosa CGR, Verlengia R, Ribeiro AGSV, de Oliveira MRM, Crisp AH. Changes in physical activities patterns assessed by accelerometry after bariatric surgery: a systematic review and meta-analysis. *Obes Med* 2019;13:6–12.
- [21] Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, *et al.* The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* 2013;46(1):81–95.
- [22] Revised Cochrane risk-of-bias tool for randomized trials (RoB 2) (version 9 October 2018). [Internet]. 2018 [cited 3rd March 2019]. Available from: <https://www.riskofbias.info/welcome/rob-2-0-tool>.
- [23] Jassil FC, Manning S, Lewis N, Steinmo S, Kingett H, Lough F, *et al.* Feasibility and impact of a combined supervised exercise and nutritional-behavioral intervention following bariatric surgery: a pilot study. *J Obes* 2015;2015:693829.
- [24] Sellberg F, Possmark S, Willmer M, Tynelius P, Berglind D. One-year follow-up of a dissonance-based intervention on quality of life, well-being, and physical activity after Roux-en-Y gastric bypass surgery: a randomized controlled trial. *Surg Obes Relat Dis* 2019;15(10):1731–7.
- [25] Stolberg CR, Mundbjerg LH, Bladbjerg E-M, Funch-Jensen P, Gram B, Juhl CB. Physical training following gastric bypass: effects on physical activity and quality of life—a randomized controlled trial. *Qual Life Res* 2018;27(12):3113–22.
- [26] Papalazarou A, Yannakoulia M, Kavouras SA, Komesidou V, Dimitriadis G, Papakonstantinou A, *et al.* Lifestyle intervention favorably affects weight loss and maintenance following obesity surgery. *Obesity* (Silver Spring, Md) 2010;18:1348–53.
- [27] Herring LY, Stevinson C, Carter P, Biddle SJH, Bowrey D, Sutton C, *et al.* The effects of supervised exercise training 12–24 months after bariatric surgery on physical function and body composition: a randomised controlled trial. *Int J Obes* 2017;41:909–16.
- [28] Hanvold SE, Vinknes KJ, Loken EB, Hjartaker A, Klungsoyr O, Birkeland E, *et al.* Does lifestyle intervention after gastric bypass surgery prevent weight regain? A randomized clinical trial. *Obes Surg* 2019;29(11):3419–31.
- [29] Bond DS, Thomas JG, Vithiananthan S, Unick J, Webster J, Royce GD, *et al.* Intervention-related increases in preoperative physical activity are maintained 6-months after Bariatric surgery: results from the bari-active trial. *Int J Obes* 2017;41(2005):467–70.
- [30] Baillet A, Vallée C-A, Mampuya WM, Dionne IJ, Comeau E, Méziat-Burdin A, *et al.* Effects of a pre-surgery supervised exercise training 1 year after bariatric surgery: a randomized controlled study. *Obes Surg* 2018;28(4):955–62.
- [31] Carnero EA, Dubis GS, Hames KC, Jakicic JM, Houmard JA, Coen PM, *et al.* Randomized trial reveals that physical activity and energy expenditure are associated with weight and body composition after RYGB. *Obesity* (Silver Spring, Md) 2017;25:1206–16.
- [32] Jiménez-Loaisa A, González-Cutre D, Beltrán-Carrillo VJ, Alcaraz-Ibáñez M. Changes in bariatric patients' physical activity levels and health-related quality of life following a postoperative motivational physical activity intervention. *Obes Surg* 2020;30:2302–12.
- [33] Coleman KJ, Caparosa SL, Nichols JF, Fujioka K, Koebnick C, McCloskey KN, *et al.* Understanding the capacity for exercise in post-bariatric patients. *Obes Surg* 2017;27:51–8.
- [34] Shah M, Snell PG, Rao S, Adams-Huet B, Quittner C, Livingston EH, *et al.* High-volume exercise program in obese bariatric surgery patients: a randomized, controlled trial. *Obesity* (Silver Spring) 2011;19(9):1826–34.
- [35] Sellberg F, Possmark S, Ghaderi A, Naslund E, Willmer M, Tynelius P, *et al.* A dissonance-based intervention for women post roux-en-Y gastric bypass surgery aiming at improving quality of life and physical activity 24 months after surgery: study protocol for a randomized controlled trial. *BMC Surg* 2018;18:25.
- [36] Baillet A, Mampuya WM, Dionne IJ, Comeau E, Meziat-Burdin A, Langlois M-F. Impacts of supervised exercise training in addition to interdisciplinary lifestyle management in subjects awaiting bariatric surgery: a randomized controlled study. *Obes Surg* 2016;26:2602–10.
- [37] Bond DS, Vithiananthan S, Thomas JG, Trautvetter J, Unick JL, Jakicic JM, *et al.* Bari-active: a randomized controlled trial of a preoperative intervention to increase physical activity in bariatric surgery patients. *Surg Obes Relat Dis* 2015;11:169–77.
- [38] Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, *et al.* Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ Br Med J* 2014;348:g1687.
- [39] Howlett N, Trivedi D, Troop NA, Chater AM. Are physical activity interventions for healthy inactive adults effective in promoting behavior change and maintenance, and which behavior change techniques are effective? A systematic review and meta-analysis. *Transl Behav Med* 2019;9(1):147–57.
- [40] Compennolle S, DeSmet A, Poppe L, Crombez G, De Bourdeaudhuij I, Cardon G, *et al.* Effectiveness of interventions using self-monitoring to reduce sedentary behavior in adults: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act* 2019;16(1):63.
- [41] Brierley ML, Chater AM, Smith LR, Bailey DP. The effectiveness of sedentary behaviour reduction workplace interventions on cardiometabolic risk markers: a systematic review. *Sports Med* 2019;49(11):1739–67.

- [42] Helmerhorst H, Brage S, Warren J, Besson H, Ekelund E. A systematic review of reliability and objective criterion-related validity of physical activity questionnaires. *Int J Behav Nutr Phys Act* 2012;9(103).
- [43] Lee PH, Macfarlane DJ, Lam TH, Stewart SM. Validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF): a systematic review. *Int J Behav Nutr Phys Act* 2011;8:115.
- [44] Troiano RP, McClain JJ, Brychta RJ, Chen KY. Evolution of accelerometer methods for physical activity research. *Br J Sports Med* 2014;48(13):1019–23.
- [45] Sattar N, McInnes IB, McMurray JJ. Obesity a risk factor for severe COVID-19 infection: multiple potential mechanisms. *Circulation* 2020.
- [46] Resurreccion DM, Motrico E, Rubio-Valera M, Mora-Pardo JA, Moreno-Peral P. Reasons for dropout from cardiac rehabilitation programs in women: a qualitative study. *PLoS One* 2018;13(7):e0200636.
- [47] Sekhon M, Cartwright M, Francis JJ. Acceptability of health care interventions: a theoretical framework and proposed research agenda. *Br J Health Psychol* 2018;23(3):519–31.
- [48] Hardeman W, Michie S, Fanshawe T, Prevost AT, McLoughlin K, Kinmonth AL. Fidelity of delivery of a physical activity intervention: predictors and consequences. *Psychol Health* 2008;23(1):11–24.
- [49] Bellg AJ, Borrelli B, Resnick B, Hecht J, Minicucci DS, Ory M, *et al.* Enhancing treatment fidelity in health behavior change studies: best practices and recommendations from the NIH Behavior Change Consortium. *Health Psychol* 2004;23(5):443–51.
- [50] Borrelli B, Sepinwall D, Ernst D, Bellg AJ, Czajkowski S, Breger R, *et al.* A new tool to assess treatment fidelity and evaluation of treatment fidelity across 10 years of health behavior research. *J Consult Clin Psychol* 2005;73(5):852.
- [51] Rixon L, Baron J, McGale N, Lorencatto F, Francis J, Davies A. Methods used to address fidelity of receipt in health intervention research: a citation analysis and systematic review. *BMC Health Serv Res* 2016;16(1):663.
- [52] Tarquinio C, Kivits J, Minary L, Coste J, Alla F. Evaluating complex interventions: perspectives and issues for health behaviour change interventions. *Psychol Health* 2015;30(1):35–51.
- [53] de Bruin M. Risk of bias in randomised controlled trials of health behaviour change interventions: evidence, practices and challenges. *Psychol Health* 2015;30(1):1–7.

Available online at www.sciencedirect.com

ScienceDirect