



LJMU Research Online

Rivera-Caravaca, JM, Buckley, BJR, Harrison, SL, Fazio-Eynullayeva, E, Underhill, P, Marin, F and Lip, GYH

Direct-acting oral anticoagulants use prior to COVID-19 diagnosis and associations with 30-day clinical outcomes

<http://researchonline.ljmu.ac.uk/id/eprint/16825/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Rivera-Caravaca, JM, Buckley, BJR, Harrison, SL, Fazio-Eynullayeva, E, Underhill, P, Marin, F and Lip, GYH (2021) Direct-acting oral anticoagulants use prior to COVID-19 diagnosis and associations with 30-day clinical outcomes. Thrombosis Research. 205. pp. 1-7. ISSN 0049-3848

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Direct-acting Oral Anticoagulants Use Prior to COVID-19 Diagnosis and Associations with 30-day Clinical Outcomes

José Miguel Rivera-Caravaca^{1,2}, Benjamin J.R. Buckley^{1,3}, Stephanie L. Harrison^{1,3}, Elnara Fazio-Eynullayeva⁴, Paula Underhill⁵, Francisco Marín², Gregory Y.H.Lip^{1,3,6}

1. Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest Hospital, Liverpool, United Kingdom.
2. Department of Cardiology, Hospital Clínico Universitario Virgen de la Arrixaca, University of Murcia, Instituto Murciano de Investigación Biosanitaria (IMIB-Arrixaca), CIBERCV, Murcia, Spain.
3. Cardiovascular and Metabolic Medicine, Institute of Life Course and Medical Sciences, University of Liverpool, Liverpool, United Kingdom.
4. TriNetX LLC., Cambridge, MA, United States.
5. TriNetX LLC., London, United Kingdom.
6. Department of Clinical Medicine, Aalborg University, Aalborg, Denmark.

Address for correspondence:

Prof. Gregory Y.H. Lip

Tel.: 0151 794 9020; E-mail: gregory.lip@liverpool.ac.uk

Abstract

Background: It is unclear if direct-acting oral anticoagulants (DOACs) use before hospitalization due to COVID-19 diagnosis would potentially impact the severity and clinical outcomes thereafter. We compared 30-day hospitalization/re-hospitalization and clinical outcomes between patients on chronic DOAC therapy and patients not on oral anticoagulation (OAC) therapy at time of COVID-19 diagnosis.

Methods: We used data from TriNetX, a global federated health research network. Patients aged ≥ 18 years who were treated with DOACs at time of COVID-19 diagnosis between 20 January 2020 and 28 February 2021 were included, and matched with patients not on OAC therapy from the same period. All patients were followed-up at 30-days after COVID-19 diagnosis. The primary outcomes were all-cause mortality, hospitalization/re-hospitalization, venous thromboembolism (VTE) and intracranial hemorrhage (ICH).

Results: 738,423 patients were included. After propensity score matching (PSM), 26,006 patients remained in the study (13,003 on DOACs; 13,003 not on OAC). DOAC-treated patients (mean age 67.1 ± 15.4 years, 52.2% male) had higher relative risks (RRs) and lower 30-days event-free survival as compared to patients not on OAC for all-cause mortality (RR 1.27, 95% CI 1.12-1.44; Log-Rank test $p=0.010$), hospitalization/re-hospitalization (RR 1.72, 95% CI 1.64-1.82; Log-Rank test $p<0.001$) and VTE (RR 4.51, 95% CI 3.91-5.82; Log-Rank test $p<0.001$), but not for ICH (RR 0.90, 95% CI 0.54-1.51; Log-Rank test $p=0.513$).

Conclusion: In COVID-19 patients, previous DOAC therapy at time of diagnosis was not associated with improved clinical outcomes or lower hospitalization/re-hospitalization rate compared to patients not taking OAC therapy.

Keywords: Coronavirus disease 2019; SARS-CoV-2; thrombosis; anticoagulant; direct-acting oral anticoagulants; bleeding; mortality.

Introduction

The novel Coronavirus Disease 2019 (COVID-19), caused by the Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), is a respiratory infection which may present with a broad picture of symptoms ranging from mild upper respiratory symptoms to acute respiratory distress syndrome and severe pneumonia (1).

From the early stage of the pandemic, some patients with COVID-19 present with multi-organ involvement, including a higher risk of thrombosis (2). Indeed, COVID-19 can induce changes in the clotting system, platelet activation and artery dysfunction, that ultimately lead to vascular inflammation, a hypercoagulable state and endothelial dysfunction (3, 4). As a result, thromboembolic complications are not rare in these patients, with an estimated VTE incidence of 25% (2, 5-7). Thus, anticoagulation is now established as one of the main therapies for the management of COVID-19 patients (8-10).

Direct-acting oral anticoagulants (DOACs) have demonstrated appropriate efficacy/safety profile in different settings; however, previous studies have reported drug-drug interactions of DOACs with some antiviral treatments used in COVID-19 patients, which may even increase plasma levels of DOACs (11-13). However, it is unclear if chronic DOAC therapy before the diagnosis of COVID-19 would potentially impact the severity and clinical outcomes after COVID-19 diagnosis, since most of the evidence in relation to anticoagulation and COVID-19 refers to the in-hospital setting.

In the present study, we aimed to compare hospitalization and clinical outcomes 30-days after COVID-19 diagnosis between patients on chronic DOAC therapy at time of COVID-19 diagnosis and patients not on OAC therapy in this time frame, using a propensity score matching (PSM) approach.

Methods

We used data from TriNetX, a global federated health research network with real-time updates of anonymised electronic medical records (EMRs), predominately in the United States (US). The network is comprised of healthcare organisations (HCOs), including academic medical centres, specialty physician practices and community hospitals. Approximately, 18 million adult patients had a visit in a TriNetX HCO during 2020.

For this study, patients within the TriNetX research network with COVID-19 between 20 January 2020 and 28 February 2021, were included. The patients were stratified by whether they had (or not) a recording of receiving DOACs in the one-year period prior to COVID-19 recorded in their EMRs. The DOAC group was composed by patients with COVID-19 aged ≥ 18 years who received either dabigatran, apixaban, rivaroxaban or edoxaban between the above period, and should be on DOAC therapy for at least one year before COVID-19 diagnosis and remained on this therapy at COVID-19 diagnosis. Patients cannot have concomitant anticoagulant therapy (oral or parenteral, ie. warfarin, acenocoumarol, phenprocoumon, desirudin, defibrotide, argatroban, betrixaban, lepirudin, fondaparinux, heparin, bivalirudin, enoxaparin, dalteparin, tirofiban, and eptifibatide). The non-OAC group included patients with COVID-19 aged ≥ 18 years between the same period who were not on any anticoagulant therapy (either oral or parenteral).

The start date was chosen as 20 January 2020 because COVID-19 was first confirmed in the US on this date, and the TriNetX network is predominately US-based (14). COVID-19 was identified using criteria provided by TriNetX based on Centres for Disease Control and Prevention (CDC) coding guidelines (15). COVID-19 status was determined using codes in EMRs or a positive test result identified with COVID-19-specific laboratory codes. Specifically, COVID-19 was identified by one or more of the

International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) codes or a positive laboratory test result (using LOINCs) in the EMRs of the patients (Supplementary Table 1). The searches were run in TriNetX on 6 of May 2021, which allowed for at least 30-days of follow-up for all participants from the time all conditions (COVID-19 and DOAC or COVID-19 and no OAC) were fulfilled. When the searches were run, there were 60 participating HCOs within the TriNetX research network.

The following data were also available from the patient EMRs: baseline demographics, comorbidities (e.g. hypertension, coronary artery disease, diabetes mellitus, heart failure, previous stroke, peripheral vascular disease, prior gastrointestinal haemorrhage and prior intracerebral haemorrhage, using ICD-10 codes) and medication use (e.g. antiplatelets, beta-blockers, calcium channel blockers, anti-arrhythmics, angiotensin-converting enzyme inhibitors, angiotensin receptor blockers and diuretics).

Follow-up and clinical outcomes

All patients were followed-up at 30-days after COVID-19 diagnosis. Clinical outcomes recorded as primary included all-cause mortality, hospitalization/re-hospitalization (if the patient was already hospitalized at time of COVID-19 diagnosis), venous thromboembolism (VTE) and intracranial hemorrhage (ICH). The secondary outcomes were the composite of ischemic stroke/transient ischemic attack(TIA)/systemic embolism (SE); the composite of ICH/gastrointestinal bleeding; myocardial infarction; and the composite of any thrombotic or thromboembolic event. Further details about the ICD-10-CM codes used for the identification of every outcome are included in Supplementary Table 2.

Ethical issues

As a federated network, research studies using TriNetX do not require ethical approval. To comply with legal frameworks and ethical guidelines guarding against data re-identification, the identity of participating HCOs and their individual contribution to each dataset are not disclosed. The TriNetX platform only uses aggregated counts and statistical summaries of de-identified information. No Protected Health Information or Personal Data is made available to the users of the platform.

Statistical analysis

Continuous variables were expressed as mean and standard deviation (SD), and tested for differences with independent-sample *t* tests. Categorical variables were expressed as absolute frequencies and percentages, and tested for differences with chi-squared test.

The TriNetX platform was used to run 1:1 PSM using logistic regression. The platform uses 'greedy nearest-neighbour matching' with a caliper of 0.1 pooled standard deviations and difference between propensity scores ≤ 0.1 . We assessed covariate balance between groups using standardised mean differences (SMDs). Any baseline characteristic with a SMD between cohorts lower than 0.1 is considered well matched (16).

Risk Ratios (RR) with 95% confidence intervals (CI) for 30-days outcomes were calculated following PSM. Kaplan-Meier survival curves were also produced with Log-Rank tests after PSM. No imputations were made for missing data. Two-sided p-values < 0.05 were accepted as statistically significant. Statistical analysis was performed using the TriNetX Analytics function in the online research platform.

Results

An overall cohort of 738,423 patients with COVID-19 was included. Of these, 13,229 (6,899 [52.15%] males, mean age 67.30±15.43 years) were on DOAC therapy at the time of COVID-19 diagnosis, and 725,194(315,061[43.45%] males, mean age 45.50±18.10 years) were not on OAC therapy at this time period.

Patients from the cohort with previous DOAC therapy were older and had a significantly higher proportion of males, non-Hispanic/Latino patients and comorbidity (Table 1). After PSM, there were 26,006 patients who remained in the study, 13,003 individuals included in each cohort (1:1), and both were well-balanced on age, gender and ethnicity, apart from all the included comorbidities (Table 1).

Comparisons of clinical outcomes in propensity score matched populations

After PSM, the risks for all outcomes except ICH were significantly higher in the DOAC cohort, as summarized in the Figure 1. Regarding the primary outcomes, the RR for all-cause mortality was 1.27 (95% CI 1.12-1.44) in the DOAC population, with decreased survival probability (Log-Rank test p=0.010) (Figure 2). There was also a higher risk of hospitalization/re-hospitalization (RR 1.72, 95% CI 1.64-1.82) and a higher risk of VTE (RR 4.51, 95% CI 3.91-5.82) in previous DOAC users, with Kaplan-Meier analysis showing that 30-day survival probability was lower in these patients (Log-Rank tests p<0.001) (Figure 2). The risk of ICH was not significantly different between both cohorts (RR 0.90, 95% CI 0.54-1.51; Log-Rank test p=0.513).

In terms of secondary outcomes, the RR of the composite ischemic stroke/TIA/SE was 1.59 (95% CI 1.31-1.92; Log-Rank test p<0.001). Similarly, there were significantly higher risks of ICH/gastrointestinal bleeding (RR 1.27, 95% CI 1.02-1.59), myocardial infarction (RR 1.49, 95% CI 1.19-1.86), and the composite of

thrombotic and thromboembolic events (RR 2.81, 95% CI 2.55-3.11). Kaplan-Meier analysis also demonstrated that patients from the DOAC cohort had poorer survival probabilities for myocardial infarction and the composite outcome (Log-Rank tests $p=0.004$ and $p<0.001$; respectively), but not for ICH/gastrointestinal bleeding (Log-Rank test $p=0.120$) (Supplementary Figure 1).

Sensitivity analyses

In order to investigate further the clinical outcomes, we performed two sensitivity analyses.

In the first one, previous atrial fibrillation (AF) or VTE were not used for adjusting the PSM. Thus, 26,466 patients remained in the study after PSM; 13,233 patients in the previous DOAC cohort and 13,233 patients in the cohort without previous OAC. The matched cohorts were balanced for all other variables shown in the main analysis, but not in terms of AF (50.22% vs. 18.02%; $p<0.001$; SMD = 0.722) and VTE (20.09% vs. 7.27%; $p<0.001$; SMD = 0.380). Even so, the risk for all-cause mortality was higher in the DOAC cohort (RR 1.65, 95% CI 1.45-1.88), as well as the risk for hospitalization/re-hospitalization (RR 1.89, 95% CI 1.79-2.00), and VTE (RR 9.14, 95% CI 7.55-11.06) (Log-Rank tests $p<0.001$). However, there were no difference in terms of ICH between cohorts (RR 1.40, 95% CI 0.79-2.48; Log-Rank test $p=0.362$). KM curves for these outcomes are shown in Supplementary Figure 2.

Another sensitivity analysis was performed accounting for the severity of COVID-19 infection. This analysis showed 25,978 patients overall after PSM (12,989 on previous DOAC:12,989 not on previous OAC), and both cohorts were balanced in terms of hospitalization at inclusion (34.79% vs. 34.40%; $p=0.506$; SMD = 0.008), intensive care unit (ICU) stay at inclusion (13.74% vs. 13.00%; $p=0.083$; SMD = 0.022)

and mechanical respiratory assistance requirements at inclusion (5.27% vs. 4.67%; $p=0.026$; SMD = 0.028). The results were similar to those found for the main analysis, with a significantly increased risk for all-cause mortality (RR 1.35, 95% CI 1.19-1.53), hospitalization/re-hospitalization (RR 1.58, 95% CI 1.50-1.66), and VTE (RR 4.39, 95% CI 3.82-5.05) (Log-Rank tests $p<0.001$) in the DOAC cohort, without differences regarding ICH risk (RR 0.82, 95% CI 0.49-1.36; Log-Rank test $p=0.308$) (Supplementary Figure 3).

Discussion

In the present study including a large cohort of patients with COVID-19, we demonstrate that chronic DOAC therapy before COVID-19 diagnosis was not associated with improved clinical outcomes or hospitalization at 30-days, compared to non-OAC therapy, after adjusting for comorbidities using PSM.

As far as we are aware, the present study represents the largest cohort of patients investigating the efficacy and safety of previous DOAC use before COVID-19 diagnosis. Importantly, our analysis is balanced by PSM and included both outpatients and inpatients. Although our results confirm some previous observations, the topic remains controversial and reinforces the hypothesis that OAC-treated patients are particularly vulnerable and still have an inherent pro-inflammatory state.

COVID-19 increases the risk of both arterial and venous thrombotic complications. For this reason, anticoagulation is one of the target therapies in the management of patients with COVID-19. Thus, the American Society of Hematology guidelines on the use of anticoagulation for thromboprophylaxis in patients with COVID-19 suggests using prophylactic-intensity over intermediate- or therapeutic-intensity anticoagulation for patients with COVID-19–related acute illness who do not have suspected or confirmed VTE (17). Recently, the INSPIRATION trial supported this, showing that among COVID-19 patients admitted to the ICU, intermediate-dose prophylactic anticoagulation, did not result in a significant difference in the composite outcome of venous or arterial thrombosis, treatment with extracorporeal membrane oxygenation, or mortality within 30 days, compared with standard-dose prophylactic anticoagulation (18). Indeed, an interim analysis of combined data from the REMAP-CAP, ATTACC, and ACTIV4a trials showed that inpatients with severe disease, receiving prophylactic anticoagulation at therapeutic doses did not reduce the

requirement for organ support (19). These results reinforced the hypothesis that more aggressive anticoagulation therapy does not have much benefit as expected.

On the other hand, a recent real-world observational study demonstrated that early initiation of prophylactic anticoagulation compared with no anticoagulation among COVID-19 patients admitted to hospital was associated with a decreased risk of 30-day mortality and no increased risk of serious bleeding events (20); however, the frequency of DOAC use was low in this study. A large cohort study simulating an intention-to-treat trial analyzed the effect of anticoagulation therapy on mortality, showing that patients with moderate or severe illness benefited from anticoagulation in the first 48 hours of hospitalization, with a similar efficacy for apixaban compared to enoxaparin in decreasing mortality (21). Unfortunately, most of the evidence thus far is focused on *hospitalized* patients (and in-hospital outcomes) and there are limited data about the role of previous antithrombotic therapies before COVID-19 diagnosis on the progression and prognosis of the disease. In particular, information about COVID-19 patients who are not yet hospitalized is scarce.

One small study in an Italian cohort of elderly patients with COVID-19 concluded that chronic DOAC intake was an independently associated with a decreased mortality risk (HR 0.38, 95% CI 0.17-0.58; $p=0.010$) (22). Similarly, another study in Italy showed that elderly patients with COVID-19 on chronic OAC treatment for AF had lower all-cause mortality rate ratio compared to their PSM non-anticoagulated counterpart (23). A further cohort study including hospitalized COVID-19 patients from Germany showed that pre-existing therapy with both DOACs or VKAs was associated with a lower risk for the primary endpoint (all-cause mortality or need for invasive or non-invasive ventilation or extracorporeal membrane oxygenation) (24). Similarly, in COVID-19 patients initially admitted in medical wards of French hospitals, previous

OAC with VKA or DOACs significantly decreased ICU admission or in-hospital mortality (25).

However, Sivaloganathan *et al.* demonstrated that patients taking antithrombotic therapy (anticoagulant or antiplatelet agents) at the time of infection with COVID-19, did not have a significantly different mortality risk to those patients not taking these drugs (26). A further recent study showed no difference in the risk of acute respiratory distress syndrome at admission or death during hospitalization, between COVID-19 patients treated with antiplatelets or anticoagulants pre-admission, compared to those untreated (27). Likewise, anticoagulant use pre-COVID-19 diagnosis was not associated with a decreased risk for all-cause mortality, mechanical ventilation or hospital admission in a retrospective study from the New York City health system, suggesting that previous anticoagulant use did not protect against development of severe COVID-19 (28). Data from a preliminary analysis of the HOPE COVID-19 Registry evidenced a significantly lower survival and higher mortality risk in COVID-19 patients on OAC therapy at hospital admission compared to patients without prior OAC at admission (29). Other study showed that chronic anticoagulants or antiplatelets use was not associated with a lower risk of any primary outcome in COVID-19 patients, including hospitalization, mortality, VTE, emergency department visit, ICU stay, invasive ventilator use or any outcome (OR anticoagulants vs. none 1.21, 95% CI 0.93-1.56) (30). Finally, a nationwide register-based cohort study in Sweden demonstrated that ongoing DOAC use at the time of SARS-CoV-2 infection was not associated with reduced risk of COVID-19 hospitalization nor the composite of ICU admission or death due to COVID-19 (31).

The published data thus far indicate that the evidence for DOACs in this context is controversial, and the present study (being the largest series) shows that previous

chronic DOAC therapy at time of COVID-19 diagnosis was not associated with improved clinical outcomes or lower hospitalization, and instead, an increased risk of mortality, hospitalization/re-hospitalization and VTE. The reasons underlying these observations are complex. Even despite PSM adjusting, patients on DOACs may have increased risk for other reasons, including a baseline pro-inflammatory state. OAC has the potential to reduce this by blocking inflammation hypercoagulability activation in the setting of COVID-19 (32, 33), but at the same time the SARS-Cov-2 enhances abnormal coagulation markers (3). In its severe form, COVID-19 induces a cytokine storm that predispose to thrombotic disease due to excessive inflammation, platelet activation, endothelial dysfunction, and blood stasis (2, 34, 35). We could hypothesize that the presence of SARS-CoV-2 infection may increase the already-high thrombotic risk of these patients on DOACs, and a pathophysiological explanation for the observed VTE risk and lower survival on our study.

Currently, there are several ongoing clinical trials in the context of COVID-19, in both inpatients and outpatients (36). However, patients included in randomized clinical trials (RCTs) are usually more selected, so results from observational studies like this one are still necessary and highly encouraged in order to support and complement RCTs, as well as to provide real-world evidence.

Limitations

We should acknowledge limitations in relation to this study. The main limitation is that the data were collected from the HCO EMRs and some health conditions may be underreported. Recording of ICD codes in EMR may vary by factors including age, comorbidities, severity of illness, length of in-hospital stay, and in-hospital mortality. Further residual confounding may include lifestyle factors such as alcohol consumption and physical activity, which were not available. In addition, the use of ICD codes for

the identification of COVID-19 cases also requires validation studies in terms of sensitivity and specificity. Furthermore, propensity scores are a method used to balance covariates, but in observational studies propensity scores are estimated and therefore there is no certainty that the propensity score was 100% accurate. We also could not determine if there was any impact of attending different HCOs because of data privacy restrictions. We examined all deaths of the included patients captured within the TriNetX network; however, deaths outside of the participating HCOs are not well captured. Finally, our main objective was to investigate the impact of prior DOAC therapy on short-term prognosis after COVID-19 diagnosis. For this reason we did not take into account the anticoagulation therapy once patients were diagnosed of COVID-19, since our interest was on the previous use of DOACs (or not). We acknowledge that this may imply a bias, but the large cohort and the sensitivity analyses performed aim to make these results valid and generalizable.

Conclusion

In patients with COVID-19, previous chronic DOAC therapy at time of COVID-19 diagnosis was not associated with improved clinical outcomes or lower hospitalization rate compared to COVID-19 patients not taking OAC therapy, even after adjusting for comorbidities.

Figure legends

Figure 1. Forest plot of odds ratios for outcomes in the DOAC population after propensity score matching.

TIA = transient ischemic attack; SE = systemic embolism; ICH = intracranial hemorrhage.

Figure 2. Comparison of survival curves for the primary outcomes between patients on DOACs or not at COVID-19 diagnosis after propensity score matching.

Purple line = Prior DOAC use.

Green line = Not prior OAC use.

Acknowledgements

Conflict of interest statement

JMR-C has received a grant from Sociedad Española de Trombosis y Hemostasia (grant for short international training stays 2020), and the First Contact Initiative Grant 2020 from the European Society of Cardiology Council on Basic Cardiovascular Science. GYHL: Consultant and speaker for BMS/Pfizer, Boehringer Ingelheim and Daiichi-Sankyo. No fees are received personally. Other authors report no conflict of interest.

Funding

The authors received no specific funding for this work.

References

1. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239-42.
2. Bikdeli B, Madhavan MV, Jimenez D, Chuich T, Dreyfus I, Driggin E, et al. COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-up. *J Am Coll Cardiol*. 2020.
3. Violi F, Pastori D, Cangemi R, Pignatelli P, Loffredo L. Hypercoagulation and Antithrombotic Treatment in Coronavirus 2019: A New Challenge. *Thromb Haemost*. 2020;120(6):949-56.
4. Del Sole F, Farcomeni A, Loffredo L, Carnevale R, Menichelli D, Vicario T, et al. Features of severe COVID-19: A systematic review and meta-analysis. *Eur J Clin Invest*. 2020;50(10):e13378.
5. Piazza G, Campia U, Hurwitz S, Snyder JE, Rizzo SM, Pfeferman MB, et al. Registry of Arterial and Venous Thromboembolic Complications in Patients With COVID-19. *J Am Coll Cardiol*. 2020;76(18):2060-72.
6. McBane RD, 2nd, Torres Roldan VD, Niven AS, Pruthi RK, Franco PM, Linderbaum JA, et al. Anticoagulation in COVID-19: A Systematic Review, Meta-analysis, and Rapid Guidance From Mayo Clinic. *Mayo Clin Proc*. 2020;95(11):2467-86.
7. Zhang C, Shen L, Le KJ, Pan MM, Kong LC, Gu ZC, et al. Incidence of Venous Thromboembolism in Hospitalized Coronavirus Disease 2019 Patients: A Systematic Review and Meta-Analysis. *Front Cardiovasc Med*. 2020;7:151.

8. Zhai Z, Li C, Chen Y, Gerotziapas G, Zhang Z, Wan J, et al. Prevention and Treatment of Venous Thromboembolism Associated with Coronavirus Disease 2019 Infection: A Consensus Statement before Guidelines. *Thromb Haemost.* 2020;120(6):937-48.
9. Spyropoulos AC, Levy JH, Ageno W, Connors JM, Hunt BJ, Iba T, et al. Scientific and Standardization Committee communication: Clinical guidance on the diagnosis, prevention, and treatment of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost.* 2020;18(8):1859-65.
10. Bikdeli B, Madhavan MV, Gupta A, Jimenez D, Burton JR, Der Nigoghossian C, et al. Pharmacological Agents Targeting Thromboinflammation in COVID-19: Review and Implications for Future Research. *Thromb Haemost.* 2020.
11. Liverpool-Drug-Interaction-Group. Detailed recommendations for interactions with experimental COVID-19 therapies 2020 [updated 09 April 2020; . Available from: <https://www.covid19-druginteractions.org/>.
12. Mueck W, Kubitzka D, Becka M. Co-administration of rivaroxaban with drugs that share its elimination pathways: pharmacokinetic effects in healthy subjects. *Br J Clin Pharmacol.* 2013;76(3):455-66.
13. Testa S, Prandoni P, Paoletti O, Morandini R, Tala M, Dellanoce C, et al. Direct oral anticoagulant plasma levels' striking increase in severe COVID-19 respiratory syndrome patients treated with antiviral agents: The Cremona experience. *J Thromb Haemost.* 2020;18(6):1320-3.
14. Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H, et al. First Case of 2019 Novel Coronavirus in the United States. *N Engl J Med.* 2020;382(10):929-36.

15. Centers-for-Disease-Control-and-Prevention. ICD-10-CM official coding guidelines - supplement coding encounters related to COVID-19 coronavirus outbreak 2020 [Available from: <https://www.cdc.gov/nchs/data/icd/ICD-10-CM-Official-Coding-Gudance-Interim-Advice-coronavirus-feb-20-2020.pdf>].
16. Haukoos JS, Lewis RJ. The Propensity Score. *Jama*. 2015;314(15):1637-8.
17. Cuker A, Tseng EK, Nieuwlaat R, Angchaisuksiri P, Blair C, Dane K, et al. American Society of Hematology 2021 guidelines on the use of anticoagulation for thromboprophylaxis in patients with COVID-19. *Blood Adv*. 2021;5(3):872-88.
18. Sadeghipour P, Talasaz AH, Rashidi F, Sharif-Kashani B, Beigmohammadi MT, Farrokhpour M, et al. Effect of Intermediate-Dose vs Standard-Dose Prophylactic Anticoagulation on Thrombotic Events, Extracorporeal Membrane Oxygenation Treatment, or Mortality Among Patients With COVID-19 Admitted to the Intensive Care Unit: The INSPIRATION Randomized Clinical Trial. *JAMA*. 2021;325(16):1620-30.
19. ATTACC A-aR-CmR. Results of interim analysis 2021 [Available from: <https://www.attacc.org/presentations>].
20. Rentsch CT, Beckman JA, Tomlinson L, Gellad WF, Alcorn C, Kidwai-Khan F, et al. Early initiation of prophylactic anticoagulation for prevention of coronavirus disease 2019 mortality in patients admitted to hospital in the United States: cohort study. *BMJ*. 2021;372:n311.
21. Billett HH, Reyes-Gil M, Szymanski J, Ikemura K, Stahl LR, Lo Y, et al. Anticoagulation in COVID-19: Effect of Enoxaparin, Heparin, and Apixaban on Mortality. *Thromb Haemost*. 2020.

22. Rossi R, Coppi F, Talarico M, Boriani G. Protective role of chronic treatment with direct oral anticoagulants in elderly patients affected by interstitial pneumonia in COVID-19 era. *Eur J Intern Med.* 2020;77:158-60.
23. Denas G, Gennaro N, Ferroni E, Fedeli U, Lorenzoni G, Gregori D, et al. Reduction in all-cause mortality in COVID-19 patients on chronic oral anticoagulation: A population-based propensity score matched study. *Int J Cardiol.* 2020.
24. Fröhlich GM, Jeschke E, Eichler U, Thiele H, Alhariri L, Reinthaler M, et al. Impact of oral anticoagulation on clinical outcomes of COVID-19: a nationwide cohort study of hospitalized patients in Germany. *Clin Res Cardiol.* 2021:1-10.
25. Chocron R, Galand V, Cellier J, Gendron N, Pommier T, Bory O, et al. Anticoagulation prior to hospitalization is a potential protective factor for COVID-19: insight from a French multicenter cohort study. *J Am Heart Assoc.* 2021:e018288.
26. Sivaloganathan H, Ladikou EE, Chevassut T. COVID-19 mortality in patients on anticoagulants and antiplatelet agents. *Br J Haematol.* 2020;190(4):e192-e5.
27. Russo V, Di Maio M, Attena E, Silverio A, Scudiero F, Celentani D, et al. Clinical impact of pre-admission antithrombotic therapy in hospitalized patients with COVID-19: A multicenter observational study. *Pharmacol Res.* 2020;159:104965.
28. Tremblay D, van Gerwen M, Alsen M, Thibaud S, Kessler A, Venugopal S, et al. Impact of anticoagulation prior to COVID-19 infection: a propensity score-matched cohort study. *Blood.* 2020;136(1):144-7.
29. Rivera-Caravaca JM, Núñez-Gil IJ, Vivas D, Viana-Llamas MC, Uribarri A, Becerra-Muñoz VM, et al. Clinical profile and prognosis in patients on oral anticoagulation before admission for COVID-19. *Eur J Clin Invest.* 2020:e13436.

30. Ho G, Dusendang JR, Schmittiel J, Kavecansky J, Tavakoli J, Pai A. Association of chronic anticoagulant and antiplatelet use on disease severity in SARS-COV-2 infected patients. *J Thromb Thrombolysis*. 2021:1-6.
31. Flam B, Wintzell V, Ludvigsson JF, Mårtensson J, Pasternak B. Direct oral anticoagulant use and risk of severe COVID-19. *J Intern Med*. 2020.
32. Connors JM, Levy JH. Thromboinflammation and the hypercoagulability of COVID-19. *J Thromb Haemost*. 2020;18(7):1559-61.
33. Paranjpe I, Fuster V, Lala A, Russak AJ, Glicksberg BS, Levin MA, et al. Association of Treatment Dose Anticoagulation With In-Hospital Survival Among Hospitalized Patients With COVID-19. *J Am Coll Cardiol*. 2020;76(1):122-4.
34. Levi M, Thachil J, Iba T, Levy JH. Coagulation abnormalities and thrombosis in patients with COVID-19. *Lancet Haematol*. 2020;7(6):e438-e40.
35. Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*. 2020;395(10229):1033-4.
36. Talasaz AH, Sadeghipour P, Kakavand H, Aghakouchakzadeh M, Kordzadeh-Kermani E, Van Tassell BW, et al. Recent Randomized Trials of Antithrombotic Therapy for Patients With COVID-19: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2021;77(15):1903-21.