



LJMU Research Online

Gee, I, Semple, S, Watson, A and Crossfield, A

Nearly 85% of tobacco smoke is invisible - a confirmation of previous claims

<http://researchonline.ljmu.ac.uk/id/eprint/1698/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Gee, I, Semple, S, Watson, A and Crossfield, A (2013) Nearly 85% of tobacco smoke is invisible - a confirmation of previous claims. TOBACCO CONTROL, 22 (6). ISSN 0964-4563

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

85% of Tobacco Smoke is Invisible – a confirmation of previous claims.

Ivan L Gee¹, Sean Semple^{2,3}, Adrian Watson⁴ and Andrea Crossfield⁵.

Author Affiliations:

¹ Faculty of Health and Applied Social Sciences, Liverpool John Moores University, Liverpool, UK.

² Scottish Centre for Indoor Air, Population Health, Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, UK.

³ Scottish Centre for Indoor Air, Institute of Occupational Medicine, Edinburgh, UK.

⁴ School of Science and the Environment, Manchester Metropolitan University, Manchester, UK.

⁵ Tobacco Free Futures, Manchester, UK.

Correspondence to:

Dr Ivan Gee, Faculty of Health and Applied Social Sciences, Liverpool John Moore's University, Henry Cotton Building, 15-21 Webster St, Liverpool, L3 2ET. UK.

Email: i.l.gee@ljamu.ac.uk; Tel: 0151 231 4300.

Contributors: the original concept for the article came from Andrea Crossfield, when researching the provenance of the claim that 85% of tobacco smoke is invisible. Ivan Gee led both the data analysis and the drafting of the paper with significant contributions from Sean Semple and Adrian Watson. All contributors reviewed and edited the final manuscript. Ivan Gee is the guarantor for the article.

Keywords: Secondhand smoke, particulate matter, tobacco advocacy, tobacco campaigns.

Word Count: 650.

Tobacco control campaigns frequently make use of statistical and scientific information to inform the public and policy makers about the dangers of tobacco smoke. It is clearly crucial for the credibility of tobacco control programmes that accurate, scientifically valid information is used in these campaign materials and Siegel [1] highlights the importance of ensuring that the evidence base of these materials is scientifically sound.

In 2002 the British Medical Association produced a report on passive smoking that indicated that:

“Almost 85 per cent of second-hand smoke is in the form of invisible, odourless gases” [2].

This referenced a US National Research Council report [3] as the source of this information but this report does not directly explain the 85% figure. At no point does the US report suggest that 85% of second-hand smoke or environmental tobacco smoke (ETS) is invisible.

However, despite the lack of unambiguous scientific evidence, the suggestion that 85% of tobacco smoke is invisible has been widely reported as fact in authoritative reports as well as numerous campaign materials, both in the UK and internationally. These include the influential Royal College of Physicians' document on 10 reasons to make public places smokefree[4], successful TV campaigns[5] and the Tobacco Control Resource Centre[6].

We have examined the available data on tobacco smoke compositions and believe that a justification of the 85% figure can be made and we present details of this in this letter.

In order to determine how much of tobacco smoke is 'invisible' it is necessary to obtain estimates of typical particle phase concentrations (the visible components) and gaseous concentrations (the invisible fraction). This gas to particle ratio can then be used as an estimate of the ratio between

invisible and visible components. Many studies of tobacco smoke composition have been conducted but not all of these report both particulate and gaseous materials. Nelson et. al.[7], working for RJ Reynolds tobacco company have conducted numerous detailed chemical studies of environmental tobacco smoke using environmental chambers. They report a range of gaseous and particle concentrations in ETS for cigarettes from 11 countries.

Of the gaseous components contained in tobacco smoke, nitrogen (N_2) oxygen (O_2) and carbon dioxide (CO_2) predominate but carbon monoxide (CO) is by far the largest contributing pollutant that has potential health impacts. Total volatile organic compounds represent the second largest potentially harmful gaseous component, but because they are a mixture of unspecified individual compounds and reported in micrograms/L, they cannot be converted to the same units as ETS particle measurements. As such, we chose to use CO alone to estimate the gas to particle ratio; using CO only results in more conservative estimates.

Particles are reported by Nelson et al as RSP (respirable suspended particles) using an impactor with a $3.5\ \mu\text{m}$ cut off point. For tobacco smoke particle size distributions RSP is a good estimate of the total suspended particulate (TSP) content [8]. RSP concentrations ranged from $1285\ \mu\text{g}/\text{m}^3$ to $1740\ \mu\text{g}/\text{m}^3$, with the average RSP across cigarettes from all 11 countries being $1545\ \mu\text{g}/\text{m}^3$. Average CO concentrations ranged from $5.6\ \mu\text{L}/\text{L}$ to $7.3\ \mu\text{L}/\text{L}$ and across all 11 countries averaged at $6.6\ \mu\text{L}/\text{L}$.

CO concentrations in $\mu\text{L}/\text{L}$ were converted to $\mu\text{g}/\text{m}^3$ based on a molecular weight of 28 and 1 mole of an ideal gas occupying 24.45 L at standard ambient temperature and pressure (SATP) [9] allowing for the proportion of CO by mass in ETS to be calculated. Table 1 shows the RSP and CO concentrations with the calculated contribution from CO to the mass of ETS aerosol for the lowest, highest and average ETS concentrations as presented by the Nelson data.

Table 1: Carbon Monoxide (CO) and Respirable Suspended Particle (RSP) concentrations in environmental tobacco smoke (ETS) together with the calculated % contribution of CO to ETS aerosol mass.

| | CO ($\mu\text{L/L}$)[7] | CO ($\mu\text{g}/\text{m}^3$) | RSP ($\mu\text{g}/\text{m}^3$) [7] | % contribution of CO to ETS |
|----------------------------|---------------------------|---------------------------------|--------------------------------------|-----------------------------|
| Lowest concentrations | 5.6 | 6412 | 1285 | 83.3 |
| Highest concentrations | 7.3 | 8359 | 1740 | 82.8 |
| Average ETS concentrations | 6.6 | 7578 | 1545 | 83.1 |

These estimates suggest that approximately 83% of tobacco smoke is in a gaseous form that is not visible, supporting the previously reported estimates of 85%. We also note that recent evidence would suggest that most ETS particulate matter is of less than 100nm diameter [10] and, as such, likely to be ‘invisible’ to the naked eye, further increasing the proportion of ETS by mass that would not be visible to smokers and non-smokers alike. We therefore consider it reasonable for tobacco control campaigns to continue to report that 83% of the harmful components within ETS are invisible.

Footnotes

Funding: this research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors

Competing Interests: none

References

1. Siegel M. Is the tobacco control movement misrepresenting the acute cardiovascular health effects of secondhand smoke exposure? An analysis of the scientific evidence and commentary

on the implications for tobacco control and public health practice. *Epidemiol Perspect Innov.* 2007; 4: 12.

2. British Medical Association (BMA) Towards smokefree public places BMA London 2002: 1.
3. National Research Council. Environmental tobacco smoke: measuring exposures and assessing health effects. National Academic Press, Washington 1986
4. Royal College of Physicians (RCP). Tobacco smoke pollution: the hard facts. 10 Reasons to make public places smokefree. RCP, London: 2006: 7.
5. Smokefree Resource Centre. Smokefree campaigns: the invisible killer.
<http://smokefree.nhs.uk/resources/news/campaigns/invisible-killer/> (accessed 02.02.2012)
6. Tobacco Control Resource Centre. TCRC Newsletter 2003. http://www.tobacco-control.org/tcrc_Web_Site/Pages_tcrc/News/News_Letters.htm (accessed 02.02.2012)
7. Nelson PR, Conrad FW, Kelly SP et al. Composition of environmental tobacco smoke (ETS) from international cigarettes and determination of ETS-RSP:particulate marker ratios. *Environ. Int.* 1997;23:47-52.
8. Pankow JF, Mader BT, Isabelle LM, et. al. Conversion of Nicotine in Tobacco Smoke to Its Volatile and Available Free-Base Form through the Action of Gaseous Ammonia. *Environ. Sci. Technol.* 1997; 31:2428-2433.
9. Bettelheim FA, Brown WH. Campbell MK and Farrell SO (2008) Introduction to general, organic and biochemistry, Chapter 5 pp148.
10. Becquemin MH, Bertholon JF, Attoui M, et al. Airway deposition of nanoparticles from second hand cigarette smoke. *Rev Mal Respir.* 2010;27(5):441-8.