

LJMU Research Online

Alawsi, MA, Zubaidi, S, Al-Bdairi, NSS, Al-Ansari, N and Hashim, KS

Drought Forecasting: A Review and Assessment of the Hybrid Techniques and Data Pre-processing

https://researchonline.ljmu.ac.uk/id/eprint/17159/

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Alawsi, MA, Zubaidi, S, Al-Bdairi, NSS, Al-Ansari, N and Hashim, KS ORCID logoORCID: https://orcid.org/0000-0001-9623-4060 (2022) Drought Forecasting: A Review and Assessment of the Hybrid Techniques and Data Pre-processing. Hvdrologv. 9 (7). ISSN 2306-5338

LJMU has developed LJMU Research Online for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

http://researchonline.ljmu.ac.uk/





Drought Forecasting: A Review and Assessment of the Hybrid Techniques and Data Pre-Processing

Mustafa A. Alawsi ^{1,2}, Salah L. Zubaidi ², Nabeel Saleem Saad Al-Bdairi ², Nadhir Al-Ansari ^{3,*} and Khalid Hashim ^{4,5}

- ¹ Department of Building and Construction Techniques Kut Technical Institute, Middle Technical University, Wasit 52001, Iraq; mustafa.abdulgani@mtu.edu.iq
- ² Department of Civil Engineering, Wasit University, Wasit 52001, Iraq; salahlafta@uowasit.edu.iq (S.L.Z.); nsaleem@uowasit.edu.iq (N.S.S.A.-B.)
- ³ Department of Civil Environmental and Natural Resources Engineering, Lulea University of Technology, 971 87 Lulea, Sweden
- ⁴ Built Environment and Sustainable Technologies (BEST) Research Institute, Liverpool John Moores University, Liverpool L3 3AF, UK; k.s.hashim@ljmu.ac.uk
- ⁵ Department of Environment Engineering, Babylon University, Babylon 51001, Iraq
- * Correspondence: nadhir.alansari@ltu.se

Abstract: Drought is a prolonged period of low precipitation that negatively impacts agriculture, animals, and people. Over the last decades, gradual changes in drought indices have been observed. Therefore, understanding and forecasting drought is essential to avoid its economic impacts and appropriate water resource planning and management. This paper presents a recent literature review, including a brief description of data pre-processing, data-driven modelling strategies (i.e., univariate or multivariate), machine learning algorithms (i.e., advantages and disadvantages), hybrid models, and performance metrics. Combining various prediction methods to create efficient hybrid models has become the most popular use in recent years. Accordingly, hybrid models have been increasingly used for predicting drought. As such, these models will be extensively reviewed, including preprocessing-based hybrid models, and hybridisation of components combination-based with preprocessing-based hybrid models. In addition, using statistical criteria, such as RMSE, MAE, NSE, MPE, SI, BIC, AIC, and AAD, is essential to evaluate the performance of the models.

Keywords: data pre-processing; drought; hybrid models; machine learning; performance metrics

1. Introduction

Drought has become more common all over the world in the last few decades because of climate change and global warming [1]. It is one of the most expensive and least understood natural disasters. It may have devastating consequences on agriculture, water supply, ecosystems, public health, watershed health, and the economy. Each year, the United States suffers losses of six to eight billion dollars as a result of droughts [2]. Yan et al. [3] mentioned that drought is also a major natural disaster in China, with its frequency reaching 70% in certain places during the summer. Moreover, it causes enormous socioeconomic losses, especially in agriculture. According to the United Nations (2011), drought conditions in the Horn of Africa in 2011 resulted in significant starvation in Kenya, Ethiopia, Djibouti, and Somalia [4]. Therefore, it is defined as a complex and poorly understood phenomenon because of the complexity of its contributing factors and its underlying driven factors [5]. It is usually caused by a reduction in rainfall over a long time [6]. Furthermore, in some cases, it occurs due to abnormalities in evapotranspiration and temperature [7]. As a result, it has a long-lasting effect over broad regions, lasting months or even years. It affects food production and has detrimental impacts on the economic performance of large areas or whole nations [8,9].



Citation: Alawsi, M.A.; Zubaidi, S.L.; Al-Bdairi, N.S.S.; Al-Ansari, N.; Hashim, K. Drought Forecasting: A Review and Assessment of the Hybrid Techniques and Data Pre-Processing. *Hydrology* **2022**, *9*, 115. https://doi.org/10.3390/ hydrology9070115

Academic Editor: Saeed Golian

Received: 5 May 2022 Accepted: 24 June 2022 Published: 26 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). Climate change has a considerable challenge for human beings in the future since it has the potential to cause more radical global occurrences with enormous socio-economic consequences [10,11]. As a result of growing international attention from governments and scholars, many resources have been devoted to developing measures to mitigate the effects of climate change and reverse its consequences. It is projected that the impact of climate change will likely increase in the future due to the global warming effect, which will increase the evaporation of water. This issue will most likely result in more acute drought occurrences [12,13]. Additionally, climate change affects the ecology in several ways, including but not limited to, changing precipitation patterns that may cause drought several consequences (i.e., depletion of freshwater supplies, population growth, the effect of climate change and global warming, as well as the increasing frequency and intensity of droughts). These consequences are either short-term catastrophes or long-term changes in the climate system [14,15].

Forecasting drought is essential for water supply management, irrigated agriculture, recreational tourism, environmental monitoring, and the environment's health [16,17]. Fung et al. [18], and Anshuka et al. [19] reviewed various models and techniques that have been applied in previous research to forecast drought. No global approach can outperform all models in all areas of study; thus, it is necessary to evaluate each case separately, assessing the performance of each technique or the combination of different approaches in each research field [20].

Traditional models propose that the relationship between dependent and independent (variables) is linear and can be unsuitable for solving practical application issues [21]. As a result, the complex character and non-linearity of the drought process require algorithms that can simulate non-linear time series data. Therefore, artificial intelligence (AI) algorithms in drought forecasting have received important attention [20]. Moreover, various studies have indicated that AI algorithms outperform traditional methods [22,23]. These AI algorithms are, for example, artificial neural networks (ANNs) [24], support vector machines (SVMs) [25], random forests [26], and the adaptive neuro-fuzzy inference system (ANFIS) [27].

AI has also been used for forecasting droughts by employing a multilayer perceptron network for stochastic synthesis of daily rainfall time series that preserve the Hurst coefficient and the autocovariance structure of all time scales [28]. This statistical property is very important in water management applications because it is related to "the tendency of wet years to cluster into multi-year wet periods or of dry years to cluster into multi-year drought periods" [29].

The hybrid model is developed by combining several different methods; one of these approaches serves as the primary model, while the others serve as pre-or post-processing methods [30]. Various studies have demonstrated that hybrid methods are better than standalone methods. Therefore, several studies recommended using hybrid models to increase forecast precision [22,31,32]. Recently, hybrid prediction models have become more popular than standalone models in hydrology. Figure 1 presents the percentage of studies on hybrid ML models that were used to predict drought indices over the last four years.

Different types of hybrid models have been created and successfully used to increase prediction accuracy [33,34]. Therefore, the hybrid models are classified into three categories:

(1) The pre-processing-based hybrid models (PBH): data pre-processing techniques have an efficient effect on improving the quality of data and selecting the best number of predictors that lead to enhancing the precision of the forecasting model [35]. Numerous studies have been conducted by combining the model with data pre-processing to predict drought, e.g., Zhang et al. [22], Belayneh et al. [23], Mathivha et al. [36], and Djerbouai and Souag-Gamane [37].

(2) The parameter optimisation-based hybrid models (OBH): The primary idea behind the OBH models is to use optimisation algorithms to identify the optimal parameters of the models [34], and combine the models with nature-inspired algorithms for hydrological drought prediction such as Adnan, et al. [32], Nabipour et al. [20], Banadkooki et al. [38], and Kisi et al. [39].

(3) Hybridisation of components combination-based with preprocessing-based hybrid models (HCPH): Combining two models that have the remarkable performance of prediction models with preprocessing-based hybrid models [34]. Few studies employed this type of hybrid model, such as Khan et al. [31] and Wu et al. [40].



Figure 1. A percentage of studies on hybrid ML models used to predict drought indices over the last four years.

Based on our knowledge, three recent review papers have investigated drought forecasting. Anshuka et al. [19] focused on the Standardised Precipitation Index (SPI), the most widely used drought indicator that the World Meteorological Organisation recommends, and a meta-analysis was conducted to test the suitability of data-driven models for predicting SPI. The main goal of the second review paper is to give researchers a quick overview of the models' principles and historical uses. This would keep them from overlooking a possible choice of models and save them a lot of time on the problem [18]. Sundararajan et al. [8] offered a literature survey using machine learning in drought prediction, performance metrics, drought indices, and datasets. Therefore, in addition to the previous literature review, this paper focuses on data pre-processing, data-driven modelling strategies (i.e., univariate or multivariate), machine learning algorithms (i.e., advantages and disadvantages), hybrid models, and performance metrics. The general framework of drought prediction and evaluation is shown in Figure 2.



Figure 2. The general framework of drought prediction and evaluation.

2. Drought

Drought is an unforecastable natural calamity that may cause significant threats such as water shortages and ecosystem deterioration. In addition, water is a vital resource for agriculture and many other human activities. Thus, a water shortage may negatively impact agricultural productivity and threaten food security. Water shortage is a significant concern in different parts of the world [32,41]. Erhardt and Czado [42] mentioned that there are different types of droughts:

- Hydrological (water supplies are dwindling);
- Meteorological (shortage of rainfall);
- Agricultural (soil moisture deficiency);
- Groundwater (decreased levels, discharge, and recharge of groundwater); and
- Socio-economic (an excess demand for commodities due to water scarcity), which is driven by several variables.

The prediction of drought is difficult because of its variability. Thus, it is vital to employ robust and accurate predicting models [38,43]. Drought forecasting is imperative since it allows for early warnings and hence more effective risk management. Different climatic variables are used as inputs to the forecasting model. The technique given here may be beneficial for decision-makers in reducing its consequences and developing effective strategies for water resource management on a monthly or seasonal timeline [32,44]. Agricultural production, the economy, and the environment are all negatively affected by their impacts. For example, from 1999 to 2000, a significant drought in Central and South-West Asia affected 60 million people and resulted in an estimated \$4.29 billion in economic losses [21,45].

Drought indicators are a scale of hydrological, meteorological, agricultural, or socioeconomic variables that may be used to reference probable its-related stress or deficiency; its' indices attempt to determine the extent of the drought's reach [46,47], and it is considered an effective instruments for making decisions rather than relying on raw data [48]; despite the existence of several indices, there is no widely accepted index for all kinds of droughts that take a wide diversity of drought producing factors into consideration [49,50]. The indices include the Standardized Precipitation Index (SPI) [51], Standardized Precipitation Evapotranspiration Index (SPEI) [5,52], Groundwater Resource Index (GRI) [38], Standard Index of Annual Precipitation (SIAP) [53], Precipitation Index (PI) [13], Standardized Hydrological Drought Index (SHDI) [20], Reconnaissance Drought Index (RDI) [54], Streamflow Drought Index (SDI) [55], and Palmer Drought Severity Index (PDSI) [56]. Tables 1–4 present the classification of SPI, PI, SIAP, and PDSI indices.

Table 1. Classification of drought based on SPI [57].

SPI Values	Class
>2	Extremely wet
1.5 to 1.99	Very wet
1.0 to 1.49	Moderately wet
-0.99 to 0.99	Near normal
-1.0 to -1.49	Moderately dry
-1.5 to -1.99	Severely dry
<-2	Extremely dry

Table 2. Classification of drought based on PI [13].

PI Values	Drought Description
>1.0	Extremely wet
0.75 to 1.0	Very wet
0.5 to 0.75	Moderately wet
0.5 to -0.5	Normal
-0.5 to -0.75	Moderate drought
-0.75 to -1.0	Severe drought
<-1.0	Extreme drought

Table 3. Classification of drought based on SIAP [53].

SIAP Values	Classes of Drought Intensity
>0.84	Extremely wet
0.52 to 0.84	Wet
-0.52 to 0.52	Normal
-0.52 to -0.84	Drought
<-0.84	Extreme drought

PDSI Values	Category
$PDSI \leq -4$	Extreme drought
$-4 < PDSI \le -3$	Severe drought
$-3 < PDSI \le -2$	Moderate drought
$-2 < PDSI \leq -1$	Mild drought
$PDSI \ge -1$	No drought

Table 4. Classification of drought based on PDSI [58].

3. Data-Driven Modelling Strategies

_

Data-driven modelling strategies generally consist of univariate and/or multivariate models [59]. Based on previous studies, Table 5 summarises the comprehensive details of two strategies: univariate (i.e., input-itself-output) and multivariate (i.e., more than one input variable and another output). For instance, a univariate model might just use a drought index (same time series) as input and output, while a multivariate model would utilise several parameters (e.g., rainfall, temperature, evaporation, etc.) as input and a drought index as output. When data needed for forecasting are available, such as precipitation, temperature, evaporation, and other factors influencing drought, these data can be used as input variables. When only drought index data are available, predicting drought index changes must rely only on drought index series with a specific number of time lags [59,60]. The table demonstrates that most previous studies applied a univariate strategy, and a few used a multivariate method.

Table 5. Different input strategies.

Ref.	Туре	Input Data Parameter	Output Parameter
		Twelve multivariate datasets (derived from statistically	
[61]	Multivariate	significant lagged combinations of precipitation, temperature, and humidity)	SPI1, SPI 3, SPI 6, SPI 12
[31]	Multivariate	Rainfall data series with SPI Lag	$SPI_{(t+1)}$
[35]	Multivariate	antecedent SPIs and antecedent accumulated monthly rainfall	SPI 3, SPÍ 6
[20]	Multivariate	SHDI Lag, SPI Lag, and precipitation.	SHDI 1, SHDI 3, SHDI 6.
[53]	Multivariate	Rainfall and SIAP Lag Water level and SWSI Lag	SIAP SWSI
[62]	Univariate	SPI Lag	SPI 6, SPI 12
[58]	Univariate	PDSI Lag	PDSI $(t+1)$
[63]	Univariate	MSPI Lag	$MSPI_{(t+1)}$
[64]	Univariate	SPI Lag	$SPI_{(t+1)}$
[65]	Univariate	EDI Lag	$EDI_{(t+1)}$
[39]	Univariate	SPI Lag	SPI 3, SPÌ 6, SPI 9, SPI 12
[66]	Univariate	EDI Lag	$EDI_{(t+1)}$
[67]	Univariate	sc-PDSI Lag	sc-PDSI $(t+1)$ $(t+3)$ $(t+6)$
	** • • ·		GRI 6, GRI 12,
[38]	Univariate	GRI Lag	GRI 24
[32]	Univariate	SPI Lag	SPI 3, SPI 6, SPI 9, SPI12
[25]	Univariate	PDSI Lag	PDSI (t)
[68]	Univariate	SPI Lag	SPI 3
[37]	Univariate	SPI Lag	SPI 3, SPI 6, SPI 12
[= 4]	Linizzaniata	DDI Lag	RDI 6, RDI 9,
[34]	Univariate	KDI Lag	RDI 12
[52]	Univariato	SDELL ag	SPEI 1, SPEI 3,
[32]	Ulivallate	JI LI Lag	SPEI 6
			sc-PDSI 1,
[27]	Univariate	sc-PDSI Lag	sc-PDSI 3,
			sc-PDSI 6

4. Machine Learning Algorithms

Machine learning (ML) algorithms are an implementation of Artificial Intelligence (AI). ML algorithms are designed with the ability to learn from previous experiences and perform new tasks. These algorithms are typically divided into supervised learning and unsupervised learning. Supervised learning employs labelled training data to produce output data based on past experience. Unsupervised learning is challenging since the system only has unlabeled data, yet it works independently to find the information [8]. Several machine learning algorithms are employed in prediction, such as artificial neural networks (ANNs) [53], adaptive neuro-fuzzy inference system (ANFIS) [39], support vector regression (SVR) [69], and random forest (RF) [61], gradient boosting (GB) [70], k-Nearest Neighbour (k-NN) [71], and decision tree (DT) [72].

Table 6 shows the advantages and disadvantages of the most popular machine learning algorithms.

Model Type	Advantages	Disadvantages	Ref.
ANN	-Ability to simulate and predict non-stationary and non-linear time series.	-Sometimes, ANNs have issues forecasting unstable and non-stationary time series. If data pre-processing does not apply, the ANN will be unable to forecast and solve issues.	[48,73]
ANFIS	-Use the fuzzy logic and neural network in a single model to increase efficiency.	-It needs a lot of training data to create a precise model, and these data may not be available every time.	[27,74]
RF	-Accuracy in modelling improves as the number of trees increases. -Ability to process large datasets involving several features	-Applying the model with a large number of trees causes a slow training process.	[26,75]
SVR	-It has flexibility for multiple options due to the availability of different kernel functions.	-It needs effective parameter optimisation to provide more accurate predictions.	[75,76]

Table 6. Advantages and disadvantages of stand-alone machine learning models.

5. Data Pre-Processing

Despite the high accuracy of different models in estimating hydroclimate parameters (for example, rainfall), these models may have limitations when dealing with hydrological time series that are often non-stationary and cover a wide variety of scales, ranging from a few minutes to several decades. Data pre-processing may be necessary to overcome comparable defects and problems [14,77]. According to Maier and Dandy [78], it is essential to pre-process data suitably before using it in ANN. These strategies are necessary to ensure that all data is treated equally in the learning model. Data pre-processing is vital for the majority of hydrologic time series to attain higher prediction performance [79]. It has been effectively utilised in various fields of study, e.g., monthly rainfall forecasting [80], urban water demand prediction [81], irrigation water prediction [82], and drought forecasting [83]. Data pre-processing includes normalisation, cleaning, and selecting the best model input [84]. It can be divided into three parts.

5.1. Normalisation

Screening continuous factors for normality is an essential early stage in the analysis, and the histograms are one of the most important ways of assessing normality [85]. In ML approaches, data curation is a necessary pre-processing step, beginning with data normali-

sation to restrict the input value range [86]. As a result, converting continuous variables is essential for creating time series that are normally or close to normally distributed [87]. Therefore, data normalisation seeks to obtain the identical range of values for each model's inputs and produces a time series normally or nearly normally distributed [88]. The natural logarithm was utilised to normalise the data to decrease the impact of multicollinearity between input variables [89]. In addition, Z-score normalisation is a traditional standardisation approach that uses the mean (μ) and standard deviation (σ) to standardise parameters [90]; it is calculated using Equation (1).

$$Z - score = \frac{Xi - \mu}{\sigma} \tag{1}$$

where *x* is the actual value of the dataset parameter *i*.

Moreover, Min-Max normalisation (*Xnorm*) is one of the most common and widely used data normalisation methods. It is a method that linearly transforms variables, where min and max represent the minimum and maximum values of x [91], as shown in Equation (2).

$$Xnorm = \frac{x - \min(x)}{\max(x) - \min(x)}$$
(2)

Moreover, Decimal scaling (Xi') is moving the decimal point of the variable's values to achieve the normalised value. The number of decimal points moved depends on the maximum absolute of value of the variables [92]; it is computed by the Equation (3).

$$Xi' = \frac{Xi}{10^k} \tag{3}$$

k is the smallest integer such that $Max(\lfloor x \rfloor) < 1$.

5.2. Cleaning

Data cleaning techniques include treating the outliers and pre-treatment signals [87]. An outlier is a case in which one variable has an extreme value that leads to distorted statistics. The box and whisker method was employed to identify outliers, which were then treated. This method significantly improves the accuracy of the suggested prediction model [93]. Additionally, there are various noise components in each time series, and the pre-treatment signal approaches (i.e., Wavelet [94], Empirical Mode Decomposition [95], Singular Spectrum Analysis [96], etc.) are the most effective methods to denoise the original time series by analysing them into multiple components [88]. As shown in Equation (4), A time series (T) could be analysed into a trend (X), stochastic (Y), seasonal (Z), and noise (N) [30].

$$\Gamma = X + Y + Z + N \tag{4}$$

5.3. A Selecting Appropriate Descriptors

Selecting appropriate predictors to simulate hydrological variables is difficult, specifically in non-linear hydrologic systems [68,97]. In general, choosing the optimal model input is one of the most significant steps in data pre-processing and developing an acceptable forecast model [87]. When the data-driven modelling strategy is univariate, different methods are used to choose the best antecedent lags scenario, e.g., the Mutual Information technique [88]. On the other hand, various approaches are employed when the strategy is multivariate, such as dimensionality reduction, which can be achieved by retaining input features that have large variances and discarding those terms that have small variances (i.e., Principal Component Analysis (PCA) is a feature selection approach that can reduce the model's dimensionality without impacting its performance.) [86]. Additionally, variance inflation factor and tolerance approaches are used to determine potential multicollinearity and exclude inputs from ML algorithms.) [14,93,97]. Moreover, automated weighting techniques can be used, such as Bayesian and frequentist algorithms [98]. According to

Tabachnick and Fidell [85], the required sample size for the prediction model is based on the number of predictors (independent variables), as shown in Equation (5) (e.g., if the number of predictors is four variables, the model needs 104 + 4 = 108 cases).

$$N \ge 104 + V \tag{5}$$

where: N: sample size.

V: number of predictors.

Based on previous studies reviewed in this research, Table 7 summarises data preprocessing. Most of these studies used one or two data pre-processing steps; furthermore, several studies did not employ the best model input. Besides, a few studies employed all the data pre-processing steps.

 Table 7. Summary of data preprocessing.

Authors	Normalisation	Cleaning	Best Model Input
Danandeh Mehr et al. [99]	Yes	No	Yes
Ali et al. [61]	Yes	Yes	Yes
Aghelpour et al. [58]	Yes	No	No
Aghelpour et al. [63]	Yes	No	No
Safavi et al. [64]	Yes	Yes	No
Fung et al. [100]	Yes	Yes	No
Zhang et al. [101]	No	Yes	Yes
Khan et al. [31]	No	Yes	No
Pham et al. [35]	Yes	Yes	Yes
Banadkooki et al. [38]	Yes	No	Yes
Adnan et al. [32]	Yes	No	No
Nabipour et al. [20]	Yes	No	No
Mohamadi et al. [68]	Yes	No	Yes
Djerbouai and Souag-Gamane [37]	Yes	Yes	No
Wu et al. [40]	Yes	Yes	No
Soh et al. [52]	Yes	Yes	No
Başakın et al. [27]	Yes	Yes	No
Das et al. [62]	Yes	Yes	No
Belayneh et al. [23]	Yes	Yes	No
Kisi et al. [39]	Yes	No	Yes

6. Hybrid Models

Hybrid models are a relatively new class of hydrological models that have grown in popularity in recent decades. Fung et al. [18] mentioned the first drought forecasting hybrid model in the hydrology field in 2007. One or more strategies are combined to form a hybrid model; one of these techniques works as the main model, while the others are used as pre-processing or post-processing approaches [88]. Employing hybrid models in drought prediction can be divided into three categories, namely: pre-processing-based hybrid models (PBH), parameter optimisation-based hybrid models (OBH), and hybridisation of components combination-based with pre-processing-based hybrid models (HCPH) [34]. Different studies applied the hybrid models, as shown in Figure 3 and Table 8.



Figure 3. Classification of hybrid models that were employed for forecasting drought indices.

6.1. Pre-Processing-Based Hybrid Models (PBH)

Data pre-processing is important for the majority of hydrologic time series to attain higher prediction performance [79]. In pre-processing-based hybrid models, the input data is pre-processed by different methods, as mentioned in Section 5.

A drought forecasting model that contains singular spectrum analysis (SSA) and a single least square support vector machine (LSSVM) model. Monthly Standardised Precipitation Index (SPI) data for the Tseng-Wen reservoir catchment in southern Taiwan (1975–2015). The findings indicate that SSA-LSSVM2 (LSSVM-based model combining with SSA utilising the antecedent SPI as input) outperforms LSSVM1 (LSSVM-based approach using antecedent SPI as an input). Additionally, when the performance of SSA-LSSVM2 and SSA-LSSVM3 (SSA-LSSVM-based approach using antecedent cumulative monthly rainfall as an input) is compared, it is determined that SSA-LSSVM3 is the best-suited model for SPI drought forecasting [35]. Two model types, namely: Artificial Neural Network (ANN) and Support Vector Regression (SVR). Moreover, the Wavelet Packet Transform (WPT) approach was used for pre-processing data. The Standardised Precipitation Index (SPI) from 1985 to 2013 in India was used to build and assess the model. The findings indicate that hybrid strategies (i.e., WPT-ANN and WPT-SVR) outperform single machine learning approaches (i.e., ANN and SVR). For the majority of grid sites, the hybrid WPT-ANN model outperformed the WPT-SVR model [62].

An artificial neural network (ANN) model forecasts hydrological and meteorological drought indices. These indices contain the Standardised Water Storage Index (SWSI) and the Standard Index of Annual Precipitation (SIAP). Additionally, the study applied the wavelet (W) for denoising the raw data. The data of the Langat River Basin, which is situated in central Malaysia, over 30 years (1986–2016) was employed to conduct the study. The results demonstrate that the W method increases the quality of raw data. Moreover, the combined model (W-ANN) can simulate the SWSI and SIAP indices with R equal to 0.940 and 0.973, respectively [53].

The Discrete Wavelet Transform (DWT) tool combines separately with k-Nearest Neighbour (kNN), Fuzzy and Support Vector Machine (SVM) approach tools to increase the accuracy of drought predictions using the Palmer drought severity index (PDSI). The models employ data from Turkey's Marmara area from 1960 to 2016. This research demonstrates that combined, WkNN, W-Fuzzy, and W-SVM models outperform stand-alone models (i.e., KNN, Fuzzy, and SVM models) in prediction performance. The prediction performance of the W-Fuzzy model is a little better than that of the W-KNN and W-SVM models [25].

A hybrid wavelet-neural network (W-ANN) model to simulate the standard precipitation index (SPI 6, SPI 12) in the north of the Haihe River Basin, China, from 1960 to 2010. The hybrid model was compared with the ANN and ARIMA methods. The outcomes reveal that W-ANN is the best suited based on several performance metrics [22].

The combined Wavelet (W) with neural networks (W-ANN) technique to forecast the standard precipitation index (SPI) at three-time scales (SPI 3, SPI 6, and SPI 12) in Algeria's Algerois basin (1936–2008). The combined model was compared with the standalone, ANN, and two traditional stochastic models (ARIMA and SARIMA). The results indicated that the wavelet method improves data quality and W-ANN offers the best performance based on different statistical criteria (i.e., NSE, RMSE, and MAE) [37].

The two machine learning models are support vector regression (SVR) and artificial neural networks (ANNs) with and without using wavelet transformations (WT) to predict the Standard Precipitation Index (SPI) for the long term. Moreover, these techniques were compared with the traditional model (ARIMA). The data for Awash River Basin, Ethiopia, from 1970 to 2005, were employed to build and assess the models. The statistical indicators, RMSE, MAE, and R², demonstrate that the WT-ANN technique outperformed all other models in predicting SPI 12 and SPI 24 [102].

The wavelet transform approach was adopted to denoise the raw data. Two machine learning models (ANN and SVR) and one conventional model (ARIMA) were used to forecast the drought indices. The data for Awash River Basin, Ethiopia, from 1970 to 2005, were employed to simulate short-term drought indices (SPI 3 and SPI 6). Several performance indicators (RMSE, MAE, and R²) were utilised to compare the predicted outcomes of the data-driven models. The WT method demonstrate a capability to clean raw data from noise and the hybrid models (WT-ANN and WT-SVR) outperformed standalone models (ANN, SVR, and ARIMA). Additionally, WT-ANN delivers the best performance for forecast drought [23].

An adaptive neuro-fuzzy inference system (ANFIS) with and without empirical mode decomposition (EMD) to forecast the self-calibrated Palmer Drought Severity Index (sc-PDSI). EMD analyses time series into sub-series and applies ANFIS to each sub-series to forecast the drought index from 1900 to 2016 for Adana, Turkey. The statistical indicators,

NSE and MSE, demonstrate that the hybrid EMD-ANFIS model is better than the ANFIS model when used alone [27].

Three models were used, namely: support vector regression (SVR), artificial neural network (ANN), and hybrid Wavelet and artificial neural network (W-ANN), to predict SPI 3 and SPI 12 in Ethiopia's Awash River Basin from 1970 to 2005. All models' performances were assessed using MAE, RMSE, and R². The results demonstrate that wavelet neural networks (W-ANN) are the best technique for drought prediction [24].

6.2. Parameter Optimisation-Based Hybrid Models (OBH)

New evolutionary algorithms were used to speed up the convergence of the soft computing models and increase their accuracy [68]. In hydrological research, bio-inspired optimisation approaches have been effectively used to improve models' abilities [54]. The parameters of ML models were determined by using optimisation algorithms [38]. Various types of metaheuristic algorithms are used to solve such optimisation problems. The metaheuristic algorithms emulated mathematical methods by following natural phenomena, such as a physical annealing process, animal behaviour, biological evolutionary process, etc. [103].

The ANN models with three meta-heuristic optimisation algorithms, namely the salp swarm algorithm (SSA), particle swarm optimisation (PSO), and genetic algorithm (GA), to predict the groundwater resource index (GRI). The study area is the Yazd plain, Iran, at different timescales (6, 12, and 24 months) (180 months). The GRI was modelled using five input scenarios: GRI-1 lagged, GRI-2 lagged, GRI-3 lagged, GRI-4 lagged, and GRI-5 month-lagged. The result revealed that hybrid models PSO-ANN (5), SSA-ANN (5), and GA-ANN (5) exceed other hybrid ANN models. Furthermore, a lower mean absolute error (MAE) was found in ANN-SSA models using an input scenario (5) [38].

The random vector functional link (RVFL) combined with the particle swarm optimisation (PSO), genetic algorithm (GA), grey wolf optimisation (GWO), social spider optimisation (SSO), salp swarm algorithm (SSA), and hunger games search algorithm (HGS) to predict SPI (SPI_3, SPI_6, SPI_9, and SPI_12). The data was collected from three stations in Bangladesh for 30 years. The results reveal that the HGS algorithm performed better than the other algorithms and significantly increased the RVFL method's accuracy in drought forecasting [32].

A hybridisation of artificial neural networks (ANN) with novel nature-inspired optimisation algorithms, includes the grasshopper optimization algorithm (GOA), salp swarm algorithm (SSA), biogeography-based optimisation (BBO), and particle swarm optimization (PSO) for simulating short-term hydrological drought, from October 1963 to September 2017. The Standardised Hydrological Drought Index (SHDI) was utilised to predict drought under scenarios of SHDI 1, SHDI 3, and SHDI 6 in the Dez dam in the southwestern region of Iran. According to the findings, the hybridised model outperformed the traditional ANN. Overall; PSO outperformed the other optimisation techniques in terms of performance [20].

The adaptive neuro-fuzzy inference system (ANFIS) combined with different metaheuristic algorithms, including the genetic algorithm (GA), ant colony algorithm (ACO), particle swarm optimisation (PSO), and butterfly optimisation algorithm (BOA) to forecast different time scales of SPI (SPI_3, SPI_6, SPI_9, and SPI_12). Monthly precipitation data from 1985 to 2015 in Iran were used in this study. The results demonstrate that hybrid models are better than the standalone model. In addition, the ANFIS-PSO method achieves the highest degree of accuracy [39].

Four types of machine learning methods were employed to predict SPI 3 in Iran, using data from 1980 to 2014. These methods are the adaptive neuro-fuzzy interface system (ANFIS), radial basis function neural network (RBFNN), multilayer perceptron (MLP), and support vector machine (SVM). The above methods were hybridised with different metaheuristic algorithms, including a nomadic people algorithm (NPA), the slap swarm algorithm (SSA), the bat algorithm (BA), and the krill algorithm. Generally, single SVM,

RBFNN, ANFIS, and MLP models performed less well than hybrid models. Among all the hybrid models, the ANFIS–NPA method performed the best based on statistical criteria [68].

6.3. Hybridisation of Components Combination-Based with Preprocessing-Based Hybrid Models (HCPH)

The components combination-based hybrid models (CBH) models aim to improve the prediction accuracy by combining the remarkable abilities of single predicting models [33]. The classic CBH models typically consist of linear and nonlinear components and are simulated by statistical and intelligent models [104]. CBH models are combined with preprocessing-based hybrid models (PBH) to create HCPH models to discard indeterminacy and extract useful information from raw data [34].

There are three scenarios to use ANN for predicting drought. The first one uses ANN as a standalone model. Second a combined model includes WT-ANN. Finally, a novel technique contains WT-ARIMA-ANN. This study analysed 30-year rainfall data for Malaysia's Langat River Basin from 1986 to 2016. The WT-ARIMA-ANN technique offers the best scenario to forecast drought [31].

The model wavelet-ARIMA long-short-term memory (WT-ARIMA-LSTM) was used to estimate precipitation and predicate drought events through the China Z-Index (CZI) based on annual precipitation. The data used are monthly from 1967 to 2017 for three stations in China. The hybrid technique (WT-ARIMA-LSTM) was compared separately with the standalone models ARIMA and LSTM. The results presented that the hybrid model had greater prediction accuracy than the ARIMA and LSTM at varying training and test sets [40].

The WT method to clean raw data and ANFIS and hybrid ARIMA-ANN models were applied to forecast SPEI (SPEI 1, SPEI 3, and SPEI 6) in the Langat River Basin in Malaysia from 1976 to 2015. The models were evaluated using RMSE, MAE, R²_{adj}, Nash-Sutcliffe Coefficient of Efficiency (NSE), and Willmott's Index of Agreement. The hybrid technique (WT-ARIMA-ANN) forecasts the drought index better for both the short and mid-term [52].

The following is a summary of findings from a review of several articles:

- 1. Numerous studies have demonstrated that combined metaheuristics with ML models outperform single-model approaches.
- 2. Combining decomposition techniques with machine learning (ML) models may be used to increase the performance of ML models.
- 3. The wavelet method has been proven to be successful in denoising raw data, increasing the results' accuracy.

Region	Size of Data	Model	Best Model	Performance Metric
India	1901–2015	SVR-HHO, SVR-PSO	SVR-HHO	RMSE, MAE, COC, NSE, WI
Turkey	1975–2010	ANFIS, SVM, ANN, W-ANFIS, W-SVM, W-ANN	W-ANFIS	R ² , RMSE, K–S test
Bangladesh	30 years	RVFL, RVFL- (PSO, GA, GWO, SSO, SSA, HGS)	RVFL-HGS	RMSE, MAE, NSE, R ²
Iran	1960–2018	SVM-DA, ARMA, RBFNN, SVM	SVM-DA	RMSE, NRMSE, WI, R, MAE
Taiwan	1975–2015	LSSVM1, SSA-LSSVM2, SSA-LSSVM3	SSA-LSSVM3	RMSE, MAE, R
Iran	1974–2018	SVR, SVR-FA, SVR-WOA, W-SVR	W-SVR	RMSE, MAE, WI, NSE
	RegionIndiaTurkeyBangladeshIranTaiwanIran	RegionSize of DataIndia1901-2015Turkey1975-2010Bangladesh30 yearsIran1960-2018Taiwan1975-2015Iran1974-2018	RegionSize of DataModelIndia1901–2015SVR-HHO, SVR-PSOTurkey1975–2010ANFIS, SVM, ANN, W-ANFIS, W-SVM, W-ANNBangladesh30 yearsGA, GWO, SSO, SSA, HGS)Iran1960–2018SVM-DA, ARMA, RBFNN, SVM LSSVM1, Taiwan1975–2015Iran1975–2015SSA-LSSVM2, SSA-LSSVM3 SVR, SVR-FA, SVR-WOA, W-SVR	RegionSize of DataModelBest ModelIndia1901–2015SVR-HHO, SVR-PSOSVR-HHOTurkey1975–2010ANFIS, SVM, ANN, W-ANFIS, W-SVM, W-ANNW-ANFISBangladesh30 yearsGA, GWO, SSO, SSA, HGS)RVFL-HGSIran1960–2018SVM-DA, ARMA, RBFNN, SVMSVM-DATaiwan1975–2015SSA-LSSVM2, SSA-LSSVM3 SVR, SVR-FA, W-SVRSVR-SVR

Table 8. Summary of studies applying machine learning models for forecasting drought indices.

Authors	Region	Size of Data	Model	Best Model	Performance Metric
Altunkaynak and Jalilzadnezam- abad [25]	Turkey	1960–2016	Fuzzy, kNN SVM, W-Fuzzy, W-kNN, W-SVM	W-Fuzzy	MSE, CE, R ²
Wu et al. [40]	China	1967–2017	wavelet-ARIMA- LSTM, ARIMA, LSTM	wavelet-ARIMA- LSTM	RMSE, MAE, R ²
Malik et al. [65]	India	1901–2015	SVR–GWO, SVR–SHO	SVR-GWO	MAE, RMSE, NSE, WI, R
Danandeh Mehr et al. [106]	Turkey	1971–2016	WPGP, AR1, GP, RF	WPGP	NSE, RMSE
Banadkooki et al. [38]	Iran	15 years	ANN–SSA, ANN–PSO, ANN-GA	ANN-SSA	NSE, RMSE, MAE
Xu et al. [107]	China	1951–2017	ARIMA, SVR, LSTM, ARIMA-SVR, LS-SVR, ARIMA-LSTM	ARIMA-LSTM	NSE, MSE, MAE, RMSE
Alquraish et al. [108]	Saudi Arabia	1968- 2019	ARIMA, ARIMA–GA, HMM HMM-GA, ARIMA–GA–ANN	ARIMA-GA- ANN	RMSE, R ² , NSE, MAD
Nabipour et al. [20]	Iran	1963–2017	ANN-GOA, ANN-SSA, ANN-BBO, ANN-PSO, ANN	ANN-PSO	RMSE, R ²
Das et al. [62]	India	1985–2013	ANN, SVR, WPT -ANN, WPT-SVR	WPT -ANN	R ² , RMSE, MAE
Xu et al. [109]	China	1951–2017	ARIMA, ARIMA–SVR	ARIMA-SVR	RMSE, MAE, R ² , NSE
Danandeh Mehr et al. [99]	Turkey	1971–2016	ENN, ENN-SA	ENN-SA	NSE, MXL, RMSE, BIC
Khan et al. [31]	Malaysia	1986–2016	ANN, wavelet-ANN, Wavelet-ARIMA- ANN	Wavelet- ARIMA-ANN	R, RMSE, R ²
Mohamadi et al. [68]	Iran	1980–2014	ANFIS, ANFIS-NPA, MLP–NPA, RBFNN-NPA, SVM–NPA	ANFIS-NPA	NSE, RMSE, MAE, PBIAS, R ²
Özger et al. [67]	Turkey	116 years	M5, ANFIS, SVM, W-ANFIS, W-SVM, W-M5, EMD-ANFIS, EMD-SVM, EMD-M5	W-ANFIS	MSE, NSE, R ²
Aghelpour et al. [63]	Iran	59 years	ANFIS, ANFIS-ACO, ANFIS-GA,	ANFIS-ACO	RMSE, MAE, WI
Başakın et al. [27]	Turkey	1900–2016	ANFIS-PSO ANFIS, EMD-ANFIS	EMD-ANFIS	MSE, NSE
Fung et al. [100]	Malaysia	1976–2015	W–BS–SVR, multi- input-W–F–SVR, weighted-W-F–SVR	Weighted-W–F– SVR	RMSE, R ² , MAE
Kisi et al. [39]	Iran	1985–2015	ANFIS, ANFIS-PSO, ANFIS-GA, ANFIS-BOA, ANFIS-ACOR	ANFIS-PSO	RMSE, MAE, IA
Ali et al. [61]	Pakistan	1981–2015	MEMD-SA-RF, KRR, RF MEMD-SA-KRR	MEMD-SA-RF	MSE, R, RMSE

Table 8. Cont.

Authors	Region	Size of Data	Model	Best Model	Performance Metric
Zhang et al. [101]	China	1979–2016	ARIMA, W-ANN, SVM	ARIMA	R ² , MSE, NSE, K–S
Khan et al. [53]	Malaysia	1986-2016	ANN, W-ANN	W-ANN	R, RMSE
Safavi et al. [64]	Iran	1969–2009	W-SVM, CS-SVM, SVM	W-SVM	R ² , RMSE
Soh et al. [52]	Malaysia	1976–2015	Wavelet-ARIMA- ANN, W-ANFIS	Wavelet-ARIMA- ANN	R ² _{adj} , RMSE, MAE, NSE
Zhang et al. [22]	China	1960–2010	ARIMA, ANN, W-ANN	W-ANN	K–S, R ² , Kendall rank correlation
Djerbouai and Souag-Gamane [37]	Algeria	1936–2008	ANN, W-ANN, ARIMA, SARIMA	W-ANN	NSE, RMSE, MAE
Deo et al. [110]	Australia	1916–2012	ELM, ANN, LSSVR, W-ANN, W-LSSVR, W-ELM	W-ELM	R ² , WI, NSE, RMSE, MAE, P _{dv}
Belayneh et al. [23]	Ethiopia	1970–2005	ARIMA, ANN, SVR, W-SVR, W-ANN	W-ANN	RMSE, MAE, R ²

Table 8. Cont.

7. Performance Metrics

In a drought forecast, it is necessary to evaluate the overall performance and capacity of the prediction model. The criteria used to assess the efficacy of the forecasting model are essential since it impacts the decision to select the best model or scenario [111]. The following are the most applied criteria used in the previous studies:

Root mean square error (RMSE) [112]; Mean absolute error (MAE) [113]; Determination coefficient (R²) [114]; The correlation coefficient (R) [115];

Nash-Sutcliffe-efficiency (NSE) [116]; Mean percentage error (MPE) [117];

Scatter index (SI) [118];

Bayesian information criterion (BIC) and Akaike information criterion (AIC) [119]; Absolute average deviation (AAD) [120].

7.1. Mean Absolute Error

The mean absolute error (MAE) is a statistical term that refers to the total of the absolute variations between predicted and measured [113]. The formula for calculating MAE is in the following Equation (6):

$$MAE = \frac{1}{N} \sum_{i=1}^{N} |Ri - Pi|$$
(6)

where: Pi: Represents predicted value;

Ri: Represent real value;

- \overline{R}_i : Represent mean of real value;
- \overline{P}_i : Represent mean of predicted value;
- N: Represents the total number;
- *i*: Represents a single data.

7.2. Root Mean Squared Error

The root mean squared error (RMSE) is the average squared variation between the predicted and actual output. It finds the data concentration around the best fit line [112,121].

It is employed whenever the error is significantly non-linear; this is a good indicator of the accuracy of a forecast [8].

$$RMSE = \sqrt{\sum_{i}^{N} \frac{(Ri - Pi)^2}{N}}$$
(7)

7.3. Determination Coefficient

The training-validation data set and the testing data set are being used. The coefficient of determination (R^2) is a statistical measurement of the correlation between the observed and expected values in a given situation. Using R^2 values, which range between 0 and 1, the complete relationship between the data set and the line drawn across them is shown by 1, and no relation to the type between the data and the line drawn through them is indicated by 0 [114,122].

$$R^{2} = \left(\frac{\sum_{i=1}^{N} (Ri - \overline{Ri}) (Pi - \overline{Pi})}{\sqrt{\sum (Ri - \overline{Ri})^{2} \sum (Pi - \overline{Pi})^{2}}}\right)^{2}$$
(8)

7.4. Nash-Sutcliffe Efficiency

In 1970, Nash and Sutcliffe designed the Nash-Sutcliffe efficiency (NSE). Additionally, it evaluates the accuracy of hydrological models [8,116]. The NSE is sensitive to variations between forecasts and observations that are both additive and proportional in magnitude. As a result, since NSE squares the values of paired differences, it has a disproportionate sensitivity to extreme values [37].

$$NSE = 1 - \left\{ \frac{\sum_{i=1}^{N} (Ri - Pi)^2}{\sum_{i=1}^{N} (Ri - \overline{Ri})^2} \right\}$$
(9)

7.5. Mean Percentage Error (MPE)

The mean percentage error (MPE) is the percentage deviation between the measured and predicted [117], as shown in Equation (10).

$$MPE = \frac{1}{N} \sum_{i=1}^{N} \frac{Pi - Ri}{Ri}$$
(10)

7.6. Scatter Index (SI)

The Scatter Index (*SI*) is dimensionless and an indicator of a model's overall relative accuracy. The model accuracy was considered poor if $SI \ge 30\%$, fair if $20\% \le SI < 30\%$, good if $10\% \le SI < 20\%$, and excellent if SI < 10% [117,118].

$$SI = \frac{RMSE}{\overline{R}i} \times 100 \tag{11}$$

7.7. Bayesian Information Criterion (BIC) and Akaike Information Criterion (AIC)

BIC and AIC are select model metrics, in which a conventional assessment measure is modified based on the number of data points used for calibration, m, and the number of free parameters in each model, p [119]. Equations (12) and (13) are employed to calculate the Bayesian information criterion (BIC) and the Akaike information criterion (AIC).

$$BIC = m.\ln(RMSE) + p.\ln(m)$$
(12)

$$AIC = m.\ln(RMSE) + 2p \tag{13}$$

7.8. Absolute Average Deviation (AAD)

AAD is the statistical metric that measures the deviation between the model predictions and the experimental results. Generally, the AAD between forecasted and observed findings is preferred to be as small as possible [123].

$$ADD(\%) = \left\{\frac{1}{N}\sum_{i=1}^{N}\frac{Ri-Pi}{Ri}\right\} \times 100$$
(14)

8. Future Research

Banadkooki et al. [38] proposed that future research may consider hybrid ANN models and the influence of climate change on drought modelling. Moreover, various preprocessing strategies for identifying the optimal input scenario for the models might be considered. Adnan et al. [32] applied RVFL, RVFL-PSO, RVFL-GA, RVFL-GWO, RVFL-SSO, RVFL-SSA, and RVFL-HGS to forecast SPI. The recommendation is to make a comparison between the algorithms and models under investigation and other metaheuristic algorithms and/or models. Aghelpour et al. [63] recommended that future studies employ optimisation algorithms such as Firefly, Gray Wolf, Krill herd, etc., to improve ANFIS's ability to forecast MSPI. Pham et al. [35] proposed that oceanic/atmospheric circulation parameters may be used as model inputs for the SSA-LSSVM method in addition to historical precipitation. Moreover, the influence of other data pre-processing techniques (e.g., discrete wavelet transform) on the LSSVM's efficiency might be investigated further. Ahmadi et al. [54] evaluated a single SVR along with the hybrid SVR-WOA, SVR-FA, and W-SVR to predicate the reconnaissance drought index (RDI). The recommendation is that different AI models may be combined with various bio-inspired optimisation techniques and wavelet analysis to develop other kinds of hybrid models.

Moreover, based on analysis of the previous studies in this field, hybrid models can be improved by the following:

- Using various data pre-treatment techniques, such as singular spectrum analysis (SSA) and empirical mode decomposition (EMD).
- 2. It is suggested to employ a multivariate strategy.
- 3. The selection of input variables is critical and influences the performance and accuracy of a model's output. As a result, it is recommended that more efforts should be put into determining the optimal input variable combination scenario. Hence, it is also recommended that other methods should be used to determine the inputs, such as feature selection methods, feature extraction methods, and dimensionality reduction methods.
- 4. The use of hybrid metaheuristic algorithms and machine learning techniques in drought predicting has grown considerably in recent years. Nevertheless, there is still room for enhancement concerning drought prediction.
- Applying the hybridisation of pre-processing-based with parameter optimisationbased hybrid models (i.e., including both PBH and OBH).

9. Conclusions

This research has reviewed the literature on usually applied data-driven strategies concerning forecast performance and precision. The analysis demonstrated that the performance of various machine learning, probabilistic, and time-series models is typically consistent and comparable.

Various factors affect the prediction technique's performance and accuracy; choice and comparison of techniques were conducted based on data pre-processing, an appropriate timescale, data-driven modelling strategies (univariate or multivariate), and a metaheuristic algorithm combined with the model. Recent research proved that separate classical approaches no longer provide the most accurate findings. Accordingly, hybrid models are the most effective tools that must be used to increase the accuracy of drought predictions. A comprehensive hybrid model incorporates both pre-processing and metaheuristic algorithm

techniques. Thus, a key strength of the present study was it represents a comprehensive examination of all the above factors.

Most of the data on drought is used to develop non-linear predictive models. For this type of data, models that incorporate only proven effective variables are more accurate than models that incorporate all available data without testing variables' efficiency. Consequently, in future research, the efficiency of the variables should be tested (best model input) before using all of the data as an input to the prediction models and applying normalisation and cleaning. Moreover, although significant advances in hybrid model techniques have been made recently, no new techniques, among others, have emerged as the best predicting method. Therefore, drought prediction remains a research problem, which leaves room for researchers to enhance hybrid techniques for specific applications.

Author Contributions: Conceptualization, M.A.A. and S.L.Z.; methodology, M.A.A., S.L.Z. and N.A.-A.; investigation, M.A.A.; resources, N.A.-A. and K.H.; writing—original draft preparation, M.A.A.; writing—review and editing, M.A.A., S.L.Z. and N.S.S.A.-B.; visualization, N.S.S.A.-B. and K.H.; supervision, S.L.Z.; project administration, S.L.Z.; funding acquisition, N.A.-A. All authors have read and agreed to the published version of the manuscript.

Funding: The APC was funded by Lulea University of Technology.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

AAD	Absolute Average Deviation
ACO	Ant Colony Optimization
AIC	Akaike Information Criterion
ANFIS	Adaptive Neuro-Fuzzy Inference System
ANN	Artificial neural network
AR1	Autoregressive
ARIMA	Autoregressive integrated moving average
ARMA	Autoregressive Moving Average
ARIMA-GA	Auto-regressive integrated moving average-genetic algorithm
BBO	Biogeography-Based Optimisation.
BIC	Bayesian Information Criterion
BOA	Butterfly Optimization Algorithm
BS	Boosting
CANFIS	Co-active neuro fuzzy inference system
CE	Coefficient of Efficiency
CS	Cuckoo Search
DA	Dragonfly Algorithm
EDI	Effective Drought Index
ELM	Extreme learning machine
EMD	Empirical Mode Decomposition
ENN	Elman Neural Network
F	Fuzzy
FA	Firefly Algorithm
GA	Genetic Algorithm
GOA	Grasshopper Optimisation Algorithm.
GP	Genetic programming
GRI	Groundwater Resource Index
GWO	Grey Wolf Optimizer
HGS	Hunger Games Search algorithm
HHO	Harris Hawks Optimization
HMM-GA	Hidden Markov model-genetic algorithm
IA	Index of Agreement
KA	Krill Algorithm
kNN	k- Nearest Neighbour
KRR	Kernel Ridge Regression

VC	Kalmagaran Smirnan
K-S	
LSSVM	Least Square Support Vector Machine
LSSVMI	LSSVM based model using antecedent SPI as input
LSSVR	Least squares support vector regression
LSTM	Long Short-Term Memory
LS-SVR	Least square- Support Vector Regression
M5	Model Tree
MAE	Mean Absolute Error
MAD	Mean absolute deviation
MEMD	Multivariate Empirical Mode Decomposition
MLP	Multilaver Perceptron
MLR	Multiple linear regression
MPE	Mean Percentage Error
MSPI	Multivariate Standardized Precipitation Index
MXI	Maximum Likelihood
NIAL	Nomadic People Algorithm
NDMCE	Normalized Past Maan Squared Freer
NCE	Normalized Root Mean Squared Error
INJE DJ	Parameters were deviation
Pav	Percentage peak deviation
PBIAS	Percent Bias
PDSI	Palmer Drought Severity Index
PSO	Particle Swarm Optimisation
R	Correlation Coefficient
\mathbb{R}^2	Coefficient of Determination
R ² _{adj}	Adjusted Coefficient of Determination
RBFNN	Radial Basis Function Neural Network
RDI	Reconnaissance drought index
RF	Random Forest
RMSE	Root Mean Square Error
RVFL	Random Vector Functional Link
SA	Simulated Annealing optimization algorithm.
SARIMA	Seasonal Autoregressive Integrated Moving Average
sc-PDSI	Self-calibrated Palmer Drought Severity Index
SHDI	Standardised Hydrological Drought Index
SHO	Spotted Hyena Optimizer
SI	Scotter Index
SIAD	Standard Index of Annual Draginitation
SIAI	Standardized Proginitation Evanetranenization Index
SPEI	Standardized Precipitation Index
SSA	Singular spectrum analysis
SSA	Saip swarm algorithm
SSA-LSSVM2	The LSSVM-based model coupling with SSA using antecedent SPI as input.
SSA-LSSVM3	The SSA-LSSVM-based model using antecedent accumulated monthly rainfall
	as input was developed and compared to SSALSSVM2.
SSO	Social Spider Optimization
SVM	Support Vector Machine
SVR	Support Vector Regression
SWSI	Standardized Water Storage Index
W	Wavelet
WANN	Wavelet Artificial Neural Network
W-BS-SVR	Wavelet-Boosting-Support Vector Regression
W-F-SVR	Wavelet-Fuzzy-Support Vector Regression
WI	Willmott's Index
WOA	Whale Optimization Algorithm
WPGP	Wavelet packet-genetic programming
WPT	Wavelet Packet Transform
WSVM	Wavelet-Support Vector Machine

References

- 1. Jasim, A.I.; Awchi, T.A. Regional meteorological drought assessment in Iraq. Arab. J. Geosci. 2020, 13, 284. [CrossRef]
- 2. Zhang, Q.; Li, Q.; Singh, V.P.; Shi, P.; Huang, Q.; Sun, P. Nonparametric Integrated Agrometeorological Drought Monitoring: Model Development and Application. *J. Geophys. Res. Atmos.* **2018**, *123*, 73–88. [CrossRef]
- 3. Yan, N.; Wu, B.; Boken, V.K.; Chang, S.; Yang, L. A drought monitoring operational system for China using satellite data: Design and evaluation. *Geomat. Nat. Hazards Risk* 2014, 7, 264–277. [CrossRef]
- Wanders, N.; Van Lanen, H.A.J. Future discharge drought across climate regions around the world modelled with a synthetic hydrological modelling approach forced by three general circulation models. *Nat. Hazards Earth Syst. Sci.* 2015, 15, 487–504. [CrossRef]
- 5. Mulualem, G.M.; Liou, Y.-A. Application of Artificial Neural Networks in Forecasting a Standardized Precipitation Evapotranspiration Index for the Upper Blue Nile Basin. *Water* **2020**, *12*, 643. [CrossRef]
- Dorjsuren, M.; Liou, Y.-A.; Cheng, C.-H. Time Series MODIS and in Situ Data Analysis for Mongolia Drought. *Remote Sens.* 2016, 8, 509. [CrossRef]
- Luo, L.; Apps, D.; Arcand, S.; Xu, H.; Pan, M.; Hoerling, M. Contribution of temperature and precipitation anomalies to the California drought during 2012–2015. *Geophys. Res. Lett.* 2017, 44, 3184–3192. [CrossRef]
- Sundararajan, K.; Garg, L.; Srinivasan, K.; Kashif Bashir, A.; Kaliappan, J.; Pattukandan Ganapathy, G.; Kumaran Selvaraj, S.; Meena, T. A Contemporary Review on Drought Modeling Using Machine Learning Approaches. *Comput. Modeling Eng. Sci.* 2021, 128, 447–487. [CrossRef]
- 9. Park, S.; Im, J.; Han, D.; Rhee, J. Short-Term Forecasting of Satellite-Based Drought Indices Using Their Temporal Patterns and Numerical Model Output. *Remote Sens.* 2020, 12, 3499. [CrossRef]
- 10. Yuan, X.-C.; Sun, X.; Lall, U.; Mi, Z.-F.; He, J.; Wei, Y.-M. China's socioeconomic risk from extreme events in a changing climate: A hierarchical Bayesian model. *Clim. Change* **2016**, *139*, 169–181. [CrossRef]
- 11. Tsutsui, H.; Sawada, Y.; Onuma, K.; Ito, H.; Koike, T. Drought Monitoring over West Africa Based on an Ecohydrological Simulation (2003–2018). *Hydrology* **2021**, *8*, 155. [CrossRef]
- 12. Trenberth, K.E.; Dai, A.; van der Schrier, G.; Jones, P.D.; Barichivich, J.; Briffa, K.R.; Sheffield, J. Global warming and changes in drought. *Nat. Clim. Change* 2013, *4*, 17–22. [CrossRef]
- 13. Alsumaiei, A.A.; Alrashidi, M.S. Hydrometeorological Drought Forecasting in Hyper-Arid Climates Using Nonlinear Autoregressive Neural Networks. *Water* **2020**, *12*, 2611. [CrossRef]
- 14. Zubaidi, S.L.; Dooley, J.; Alkhaddar, R.M.; Abdellatif, M.; Al-Bugharbee, H.; Ortega-Martorell, S. A Novel approach for predicting monthly water demand by combining singular spectrum analysis with neural networks. *J. Hydrol.* **2018**, *561*, 136–145. [CrossRef]
- Zubaidi, S.L.; Kot, P.; Alkhaddar, R.M.; Abdellatif, M.; Al-Bugharbee, H. Short-Term Water Demand Prediction in Residential Complexes: Case Study in Columbia City, USA. In Proceedings of the 2018 11th International Conference on Developments in eSystems Engineering (DeSE), Cambridge, UK, 2–5 September 2018; pp. 31–35. [CrossRef]
- 16. Malik, A.; Kumar, A. Meteorological drought prediction using heuristic approaches based on effective drought index: A case study in Uttarakhand. *Arab. J. Geosci.* 2020, *13*, 276. [CrossRef]
- 17. Han, J.; Singh, V.P. Forecasting of droughts and tree mortality under global warming: A review of causative mechanisms and modeling methods. *J. Water Clim. Change* **2020**, *11*, 600–632. [CrossRef]
- 18. Fung, K.F.; Huang, Y.F.; Koo, C.H.; Soh, Y.W. Drought forecasting: A review of modelling approaches 2007–2017. *J. Water Clim. Change* **2020**, *11*, 771–799. [CrossRef]
- 19. Anshuka, A.; van Ogtrop, F.F.; Willem Vervoort, R. Drought forecasting through statistical models using standardised precipitation index: A systematic review and meta-regression analysis. *Nat. Hazards* **2019**, *97*, 955–977. [CrossRef]
- Nabipour, N.; Dehghani, M.; Mosavi, A.; Shamshirband, S. Short-Term Hydrological Drought Forecasting Based on Different Nature-Inspired Optimization Algorithms Hybridized With Artificial Neural Networks. *IEEE Access* 2020, *8*, 15210–15222. [CrossRef]
- 21. Agana, N.A.; Homaifar, A. EMD-Based Predictive Deep Belief Network for Time Series Prediction: An Application to Drought Forecasting. *Hydrology* **2018**, *5*, 18. [CrossRef]
- 22. Zhang, Y.; Li, W.; Chen, Q.; Pu, X.; Xiang, L. Multi-models for SPI drought forecasting in the north of Haihe River Basin, China. *Stoch. Environ. Res. Risk Assess.* 2017, *31*, 2471–2481. [CrossRef]
- 23. Belayneh, A.; Adamowski, J.; Khalil, B. Short-term SPI drought forecasting in the Awash River Basin in Ethiopia using wavelet transforms and machine learning methods. *Sustain. Water Resour. Manag.* **2015**, *2*, 87–101. [CrossRef]
- 24. Belayneh, A.; Adamowski, J. Standard Precipitation Index Drought Forecasting Using Neural Networks, Wavelet Neural Networks, and Support Vector Regression. *Appl. Comput. Intell. Soft Comput.* **2012**, 2012, 1–13. [CrossRef]
- 25. Altunkaynak, A.; Jalilzadnezamabad, A. Extended lead time accurate forecasting of palmer drought severity index using hybrid wavelet-fuzzy and machine learning techniques. *J. Hydrol.* **2021**, *601*, 126619. [CrossRef]
- 26. Dikshit, A.; Pradhan, B.; Alamri, A.M. Short-Term Spatio-Temporal Drought Forecasting Using Random Forests Model at New South Wales, Australia. *Appl. Sci.* 2020, *10*, 4254. [CrossRef]
- Başakın, E.E.; Ekmekcioğlu, Ö.; Özger, M. Drought prediction using hybrid soft-computing methods for semi-arid region. Modeling Earth Syst. Environ. 2020, 7, 2363–2371. [CrossRef]

- 28. Rozos, E.; Dimitriadis, P.; Bellos, V. Machine Learning in Assessing the Performance of Hydrological Models. *Hydrology* **2021**, *9*, 5. [CrossRef]
- 29. Koutsoyiannis, D. Hydrologic persistence and the Hurst phenomenon. In *Water Encyclopedia, Vol. 4, Surface and Agricultural Water;* Lehr, J.H., Keeley, J.W., Lehr, J.K., Kingery, T.B., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2005; Chapter 1.
- Zubaidi, S.L.; Ortega-Martorell, S.; Kot, P.; Alkhaddar, R.M.; Abdellatif, M.; Gharghan, S.K.; Ahmed, M.S.; Hashim, K. A Method for Predicting Long-Term Municipal Water Demands Under Climate Change. *Water Resour. Manag.* 2020, 34, 1265–1279. [CrossRef]
- Khan, M.M.H.; Muhammad, N.S.; El-Shafie, A. Wavelet based hybrid ANN-ARIMA models for meteorological drought forecasting. J. Hydrol. 2020, 590, 125380. [CrossRef]
- 32. Adnan, R.M.; Mostafa, R.R.; Islam, A.R.M.T.; Gorgij, A.D.; Kuriqi, A.; Kisi, O. Improving Drought Modeling Using Hybrid Random Vector Functional Link Methods. *Water* **2021**, *13*, 3379. [CrossRef]
- 33. Hajirahimi, Z.; Khashei, M. Hybrid structures in time series modeling and forecasting: A review. *Eng. Appl. Artif. Intell.* 2019, *86*, 83–106. [CrossRef]
- Hajirahimi, Z.; Khashei, M. Hybridization of hybrid structures for time series forecasting: A review. Artif. Intell. Rev. 2022. [CrossRef]
- Pham, Q.B.; Yang, T.-C.; Kuo, C.-M.; Tseng, H.-W.; Yu, P.-S. Coupling Singular Spectrum Analysis with Least Square Support Vector Machine to Improve Accuracy of SPI Drought Forecasting. *Water Resour. Manag.* 2021, 35, 847–868. [CrossRef]
- Mathivha, F.; Sigauke, C.; Chikoore, H.; Odiyo, J. Short-Term and Medium-Term Drought Forecasting Using Generalized Additive Models. Sustainability 2020, 12, 4006. [CrossRef]
- 37. Djerbouai, S.; Souag-Gamane, D. Drought Forecasting Using Neural Networks, Wavelet Neural Networks, and Stochastic Models: Case of the Algerois Basin in North Algeria. *Water Resour. Manag.* **2016**, *30*, 2445–2464. [CrossRef]
- Banadkooki, F.B.; Singh, V.P.; Ehteram, M. Multi-timescale drought prediction using new hybrid artificial neural network models. *Nat. Hazards* 2021, 106, 2461–2478. [CrossRef]
- 39. Kisi, O.; Docheshmeh Gorgij, A.; Zounemat-Kermani, M.; Mahdavi-Meymand, A.; Kim, S. Drought forecasting using novel heuristic methods in a semi-arid environment. *J. Hydrol.* **2019**, *578*, 124053. [CrossRef]
- 40. Wu, X.; Zhou, J.; Yu, H.; Liu, D.; Xie, K.; Chen, Y.; Hu, J.; Sun, H.; Xing, F. The Development of a Hybrid Wavelet-ARIMA-LSTM Model for Precipitation Amounts and Drought Analysis. *Atmosphere* **2021**, *12*, 74. [CrossRef]
- 41. Mishra, D.; Goswami, S.; Matin, S.; Sarup, J. Analyzing the extent of drought in the Rajasthan state of India using vegetation condition index and standardized precipitation index. *Modeling Earth Syst. Environ.* **2021**, *8*, 601–610. [CrossRef]
- Erhardt, T.M.; Czado, C. Standardized drought indices: A novel univariate and multivariate approach. J. R. Stat. Soc. Ser. C Appl. Stat. 2017, 67, 643–664. [CrossRef]
- 43. Emadodin, I.; Reinsch, T.; Taube, F. Drought and Desertification in Iran. Hydrology 2019, 6, 66. [CrossRef]
- 44. Nnaji, G.A.; Clark, C.J.; Chan-Hilton, A.B.; Huang, W. Drought prediction in Apalachicola–Chattahoochee–Flint River Basin using a semi-Markov model. *Nat. Hazards* **2016**, *82*, 267–297. [CrossRef]
- 45. Zhang, R.; Chen, Z.Y.; Xu, L.J.; Ou, C.Q. Meteorological drought forecasting based on a statistical model with machine learning techniques in Shaanxi province, China. *Sci. Total Environ.* **2019**, *665*, 338–346. [CrossRef]
- 46. Yihdego, Y.; Vaheddoost, B.; Al-Weshah, R.A. Drought indices and indicators revisited. *Arab. J. Geosci.* 2019, 12, 69. [CrossRef]
- Ali, M.; Deo, R.C.; Downs, N.J.; Maraseni, T. Multi-stage committee based extreme learning machine model incorporating the influence of climate parameters and seasonality on drought forecasting. *Comput. Electron. Agric.* 2018, 152, 149–165. [CrossRef]
- Ali, Z.; Hussain, I.; Faisal, M.; Nazir, H.M.; Hussain, T.; Shad, M.Y.; Mohamd Shoukry, A.; Hussain Gani, S. Forecasting Drought Using Multilayer Perceptron Artificial Neural Network Model. *Adv. Meteorol.* 2017, 2017, 5681308. [CrossRef]
- 49. Kaur, A.; Sood, S.K. Cloud-Fog based framework for drought prediction and forecasting using artificial neural network and genetic algorithm. *J. Exp. Theor. Artif. Intell.* **2019**, *32*, 273–289. [CrossRef]
- Kaur, A.; Sood, S.K. Artificial Intelligence-Based Model For Drought Prediction and Forecasting. Comput. J. 2020, 63, 1704–1712. [CrossRef]
- 51. Poornima, S.; Pushpalatha, M. Drought prediction based on SPI and SPEI with varying timescales using LSTM recurrent neural network. *Soft Comput.* **2019**, *23*, 8399–8412. [CrossRef]
- Soh, Y.W.; Koo, C.H.; Huang, Y.F.; Fung, K.F. Application of artificial intelligence models for the prediction of standardized precipitation evapotranspiration index (SPEI) at Langat River Basin, Malaysia. *Comput. Electron. Agric.* 2018, 144, 164–173. [CrossRef]
- Khan, M.; Muhammad, N.; El-Shafie, A. Wavelet-ANN versus ANN-Based Model for Hydrometeorological Drought Forecasting. Water 2018, 10, 998. [CrossRef]
- 54. Ahmadi, F.; Mehdizadeh, S.; Mohammadi, B. Development of Bio-Inspired- and Wavelet-Based Hybrid Models for Reconnaissance Drought Index Modeling. *Water Resour. Manag.* 2021, 35, 4127–4147. [CrossRef]
- Shamshirband, S.; Hashemi, S.; Salimi, H.; Samadianfard, S.; Asadi, E.; Shadkani, S.; Kargar, K.; Mosavi, A.; Nabipour, N.; Chau, K.-W. Predicting Standardized Streamflow index for hydrological drought using machine learning models. *Eng. Appl. Comput. Fluid Mech.* 2020, 14, 339–350. [CrossRef]
- 56. Tufaner, F.; Ozbeyaz, A. Estimation and easy calculation of the Palmer Drought Severity Index from the meteorological data by using the advanced machine learning algorithms. *Environ. Monit. Assess.* **2020**, *192*, 576. [CrossRef] [PubMed]

- Freitas, A.A.; Drumond, A.; Carvalho, V.S.B.; Reboita, M.S.; Silva, B.C.; Uvo, C.B. Drought Assessment in São Francisco River Basin, Brazil: Characterization through SPI and Associated Anomalous Climate Patterns. *Atmosphere* 2021, 13, 41. [CrossRef]
- Sun, X.; Lai, P.; Wang, S.; Song, L.; Ma, M.; Han, X. Monitoring of Extreme Agricultural Drought of the Past 20 Years in Southwest China Using GLDAS Soil Moisture. *Remote Sens.* 2022, 14, 1323. [CrossRef]
- 59. Zhang, Z.; Zhang, Q.; Singh, V.P. Univariate streamflow forecasting using commonly used data-driven models: Literature review and case study. *Hydrol. Sci. J.* 2018, 63, 1091–1111. [CrossRef]
- O'Reilly, C.; Moessner, K.; Nati, M. Univariate and Multivariate Time Series Manifold Learning. *Knowl.-Based Syst.* 2017, 133, 1–16.
 [CrossRef]
- Ali, M.; Deo, R.C.; Maraseni, T.; Downs, N.J. Improving SPI-derived drought forecasts incorporating synoptic-scale climate indices in multi-phase multivariate empirical mode decomposition model hybridized with simulated annealing and kernel ridge regression algorithms. *J. Hydrol.* 2019, 576, 164–184. [CrossRef]
- 62. Das, P.; Naganna, S.R.; Deka, P.C.; Pushparaj, J. Hybrid wavelet packet machine learning approaches for drought modeling. *Environ. Earth Sci.* 2020, *79*, 221. [CrossRef]
- Aghelpour, P.; Bahrami-Pichaghchi, H.; Kisi, O. Comparison of three different bio-inspired algorithms to improve ability of neuro fuzzy approach in prediction of agricultural drought, based on three different indexes. *Comput. Electron. Agric.* 2020, 170, 105279. [CrossRef]
- 64. Safavi, H.R.; Sharghi, S.; Komasi, M. Wavelet and cuckoo search-support vector machine conjugation for drought forecasting using Standardized Precipitation Index (case study: Urmia Lake, Iran). *J. Hydroinform.* **2018**, 20, 975–988. [CrossRef]
- 65. Malik, A.; Tikhamarine, Y.; Souag-Gamane, D.; Rai, P.; Sammen, S.S.; Kisi, O. Support vector regression integrated with novel meta-heuristic algorithms for meteorological drought prediction. *Meteorol. Atmos. Phys.* **2021**, *133*, 891–909. [CrossRef]
- Malik, A.; Tikhamarine, Y.; Sammen, S.S.; Abba, S.I.; Shahid, S. Prediction of meteorological drought by using hybrid support vector regression optimized with HHO versus PSO algorithms. *Environ. Sci. Pollut. Res. Int.* 2021, 28, 39139–39158. [CrossRef] [PubMed]
- 67. Özger, M.; Başakın, E.E.; Ekmekcioğlu, Ö.; Hacısüleyman, V. Comparison of wavelet and empirical mode decomposition hybrid models in drought prediction. *Comput. Electron. Agric.* **2020**, *179*, 105851. [CrossRef]
- Mohamadi, S.; Sammen, S.S.; Panahi, F.; Ehteram, M.; Kisi, O.; Mosavi, A.; Ahmed, A.N.; El-Shafie, A.; Al-Ansari, N. Zoning map for drought prediction using integrated machine learning models with a nomadic people optimization algorithm. *Nat. Hazards* 2020, 104, 537–579. [CrossRef]
- 69. Mokhtarzad, M.; Eskandari, F.; Jamshidi Vanjani, N.; Arabasadi, A. Drought forecasting by ANN, ANFIS, and SVM and comparison of the models. *Environ. Earth Sci.* 2017, *76*, 729. [CrossRef]
- 70. Sofos, F.; Stavrogiannis, C.; Exarchou-Kouveli, K.K.; Akabua, D.; Charilas, G.; Karakasidis, T.E. Current Trends in Fluid Research in the Era of Artificial Intelligence: A Review. *Fluids* **2022**, *7*, 116. [CrossRef]
- 71. Lee, T.R.; Wood, W.T.; Phrampus, B.J. A Machine Learning (kNN) Approach to Predicting Global Seafloor Total Organic Carbon. *Glob. Biogeochem. Cycles* **2019**, *33*, 37–46. [CrossRef]
- 72. Winkler, D.; Haltmeier, M.; Kleidorfer, M.; Rauch, W.; Tscheikner-Gratl, F. Pipe failure modelling for water distribution networks using boosted decision trees. *Struct. Infrastruct. Eng.* **2018**, *14*, 1402–1411. [CrossRef]
- 73. Bayatvarkeshi, M.; Mohammadi, K.; Kisi, O.; Fasihi, R. A new wavelet conjunction approach for estimation of relative humidity: Wavelet principal component analysis combined with ANN. *Neural Comput. Appl.* **2018**, *32*, 4989–5000. [CrossRef]
- 74. Maiti, S.; Tiwari, R.K. A comparative study of artificial neural networks, Bayesian neural networks and adaptive neuro-fuzzy inference system in groundwater level prediction. *Environ. Earth Sci.* **2013**, *71*, 3147–3160. [CrossRef]
- 75. Ibrahim, K.S.M.H.; Huang, Y.F.; Ahmed, A.N.; Koo, C.H.; El-Shafie, A. A review of the hybrid artificial intelligence and optimization modelling of hydrological streamflow forecasting. *Alex. Eng. J.* **2022**, *61*, 279–303. [CrossRef]
- 76. Parveen, N.; Zaidi, S.; Danish, M. Support vector regression (SVR)-based adsorption model for Ni (II) ions removal. *Groundw. Sustain. Dev.* 2019, 9, 100232. [CrossRef]
- 77. Nourani, V.; Molajou, A.; Uzelaltinbulat, S.; Sadikoglu, F. Emotional artificial neural networks (EANNs) for multi-step ahead prediction of monthly precipitation; case study: Northern Cyprus. *Theor. Appl. Climatol.* **2019**, *138*, 1419–1434. [CrossRef]
- 78. Maier, H.R.; Dandy, G.C. Neural networks for the prediction and forecasting of water resources variables: A review of modelling issues and applications. *Environ. Model. Softw.* **2000**, *15*, 101–124. [CrossRef]
- 79. Unnikrishnan, P.; Jothiprakash, V. Daily rainfall forecasting for one year in a single run using Singular Spectrum Analysis. *J. Hydrol.* **2018**, *561*, 609–621. [CrossRef]
- 80. Ouyang, Q.; Lu, W. Monthly Rainfall Forecasting Using Echo State Networks Coupled with Data Preprocessing Methods. *Water Resour. Manag.* 2017, 32, 659–674. [CrossRef]
- 81. Seo, Y.; Kwon, S.; Choi, Y. Short-Term Water Demand Forecasting Model Combining Variational Mode Decomposition and Extreme Learning Machine. *Hydrology* **2018**, *5*, 54. [CrossRef]
- Zhang, J.; Li, H.; Shi, X.; Hong, Y. Wavelet-Nonlinear Cointegration Prediction of Irrigation Water in the Irrigation District. Water Resour. Manag. 2019, 33, 2941–2954. [CrossRef]
- 83. So, J.-M.; Lee, J.-H.; Bae, D.-H. Development of a Hydrological Drought Forecasting Model Using Weather Forecasting Data from GloSea5. *Water* **2020**, *12*, 2785. [CrossRef]

- Zubaidi, S.; Al-Bugharbee, H.; Ortega-Martorell, S.; Gharghan, S.; Olier, I.; Hashim, K.; Al-Bdairi, N.; Kot, P. A Novel Methodology for Prediction Urban Water Demand by Wavelet Denoising and Adaptive Neuro-Fuzzy Inference System Approach. *Water* 2020, 12, 1628. [CrossRef]
- 85. Tabachnick, B.G.; Fidell, L.S. Using Multivariate Statistics; Pearson: Boston, MA, USA, 2013.
- 86. Sofos, F.; Karakasidis, T.E. Nanoscale slip length prediction with machine learning tools. Sci. Rep. 2021, 11, 12520. [CrossRef]
- 87. Zubaidi, S.L.; Gharghan, S.K.; Dooley, J.; Alkhaddar, R.M.; Abdellatif, M. Short-Term Urban Water Demand Prediction Considering Weather Factors. *Water Resour. Manag.* 2018, *32*, 4527–4542. [CrossRef]
- Zubaidi, S.L.; Ortega-Martorell, S.; Al-Bugharbee, H.; Olier, I.; Hashim, K.S.; Gharghan, S.K.; Kot, P.; Al-Khaddar, R. Urban Water Demand Prediction for a City That Suffers from Climate Change and Population Growth: Gauteng Province Case Study. *Water* 2020, 12, 1885. [CrossRef]
- 89. Behboudian, S.; Tabesh, M.; Falahnezhad, M.; Ghavanini, F.A. A long-term prediction of domestic water demand using preprocessing in artificial neural network. *J. Water Supply: Res. Technol.*—AQUA **2014**, 63, 31–42. [CrossRef]
- Malek, N.H.A.; Wan Yaacob, W.F.; Md Nasir, S.A.; Shaadan, N. Prediction of Water Quality Classification of the Kelantan River Basin, Malaysia, Using Machine Learning Techniques. *Water* 2022, 14, 1067. [CrossRef]
- Jain, S.; Shukla, S.; Wadhvani, R. Dynamic selection of normalization techniques using data complexity measures. Expert Syst. Appl. 2018, 106, 252–262. [CrossRef]
- 92. Akilli, A.; Atil, H. Evaluation of Normalization Techniques on Neural Networks for the Prediction of 305-Day Milk Yield. *Turk. J. Agric. Eng. Res.* **2020**, *1*, 354–367. [CrossRef]
- Zubaidi, S.L.; Hashim, K.; Ethaib, S.; Al-Bdairi, N.S.S.; Al-Bugharbee, H.; Gharghan, S.K. A novel methodology to predict monthly municipal water demand based on weather variables scenario. J. King Saud Univ.-Eng. Sci. 2022, 34, 163–169. [CrossRef]
- Zhu, S.; Lu, H.; Ptak, M.; Dai, J.; Ji, Q. Lake water-level fluctuation forecasting using machine learning models: A systematic review. *Environ. Sci. Pollut. Res. Int.* 2020, 27, 44807–44819. [CrossRef] [PubMed]
- Le, J.A.; El-Askary, H.M.; Allali, M.; Sayed, E.; Sweliem, H.; Piechota, T.C.; Struppa, D.C. Characterizing El Niño-Southern Oscillation Effects on the Blue Nile Yield and the Nile River Basin Precipitation using Empirical Mode Decomposition. *Earth Syst. Environ.* 2020, *4*, 699–711. [CrossRef]
- Le, H.D.; Le, T.N.; Wang, J.W.; Liang, Y.S. Singular Spectrum Analysis for Background Initialization with Spatio-Temporal RGB Color Channel Data. *Entropy* 2021, 23, 1644. [CrossRef]
- Malik, A.; Kumar, A.; Singh, R.P. Application of Heuristic Approaches for Prediction of Hydrological Drought Using Multi-scalar Streamflow Drought Index. *Water Resour. Manag.* 2019, 33, 3985–4006. [CrossRef]
- 98. Paulson, N.H.; Zomorodpoosh, S.; Roslyakova, I.; Stan, M. Comparison of statistically-based methods for automated weighting of experimental data in CALPHAD-type assessment. *Calphad* **2020**, *68*, 101728. [CrossRef]
- 99. Danandeh Mehr, A.; Vaheddoost, B.; Mohammadi, B. ENN-SA: A novel neuro-annealing model for multi-station drought prediction. *Comput. Geosci.* 2020, 145, 104622. [CrossRef]
- Fung, K.F.; Huang, Y.F.; Koo, C.H. Coupling fuzzy–SVR and boosting–SVR models with wavelet decomposition for meteorological drought prediction. *Environ. Earth Sci.* 2019, 78, 693. [CrossRef]
- Zhang, Y.; Yang, H.; Cui, H.; Chen, Q. Comparison of the Ability of ARIMA, WNN and SVM Models for Drought Forecasting in the Sanjiang Plain, China. *Nat. Resour. Res.* 2019, 29, 1447–1464. [CrossRef]
- 102. Belayneh, A.; Adamowski, J.; Khalil, B.; Ozga-Zielinski, B. Long-term SPI drought forecasting in the Awash River Basin in Ethiopia using wavelet neural network and wavelet support vector regression models. *J. Hydrol.* **2014**, *508*, 418–429. [CrossRef]
- 103. Lee, K.S.; Geem, Z.W. A new meta-heuristic algorithm for continuous engineering optimization: Harmony search theory and practice. *Comput. Methods Appl. Mech. Eng.* 2005, 194, 3902–3933. [CrossRef]
- 104. Khashei, M.; Hajirahimi, Z. A comparative study of series arima/mlp hybrid models for stock price forecasting. *Commun. Stat.-Simul. Comput.* 2018, 48, 2625–2640. [CrossRef]
- 105. Taylan, E.D.; Terzi, Ö.; Baykal, T. Hybrid wavelet–artificial intelligence models in meteorological drought estimation. *J. Earth Syst. Sci.* **2021**, *130*, 38. [CrossRef]
- Danandeh Mehr, A.; Safari, M.j.S.; Nourani, V. Wavelet Packet-Genetic Programming: A New Model for Meteorological Drought Hindcasting. *Tek. Dergi* 2021, 32, 11029–11050. [CrossRef]
- Xu, D.; Zhang, Q.; Ding, Y.; Zhang, D. Application of a hybrid ARIMA-LSTM model based on the SPEI for drought forecasting. Environ. Sci. Pollut. Res. Int. 2022, 29, 4128–4144. [CrossRef] [PubMed]
- Alquraish, M.; Abuhasel, K.A.; Alqahtani, A.S.; Khadr, M. SPI-Based Hybrid Hidden Markov–GA, ARIMA–GA, and ARIMA–GA–ANN Models for Meteorological Drought Forecasting. *Sustainability* 2021, 13, 12576. [CrossRef]
- Xu, D.; Zhang, Q.; Ding, Y.; Huang, H. Application of a Hybrid ARIMA–SVR Model Based on the SPI for the Forecast of Drought—A Case Study in Henan Province, China. J. Appl. Meteorol. Climatol. 2020, 59, 1239–1259. [CrossRef]
- Deo, R.C.; Tiwari, M.K.; Adamowski, J.F.; Quilty, J.M. Forecasting effective drought index using a wavelet extreme learning machine (W-ELM) model. *Stoch. Environ. Res. Risk Assess.* 2016, 31, 1211–1240. [CrossRef]
- 111. Elavarasan, D.; Vincent, D.R.; Sharma, V.; Zomaya, A.Y.; Srinivasan, K. Forecasting yield by integrating agrarian factors and machine learning models: A survey. *Comput. Electron. Agric.* **2018**, 155, 257–282. [CrossRef]
- 112. Dikshit, A.; Pradhan, B.; Alamri, A.M. Long lead time drought forecasting using lagged climate variables and a stacked long short-term memory model. *Sci. Total Environ.* **2021**, 755, 142638. [CrossRef]

- Tian, Y.; Xu, Y.P.; Wang, G. Agricultural drought prediction using climate indices based on Support Vector Regression in Xiangjiang River basin. Sci. Total Environ. 2018, 622–623, 710–720. [CrossRef]
- 114. Park, H.; Kim, K.; Lee, D.k. Prediction of Severe Drought Area Based on Random Forest: Using Satellite Image and Topography Data. *Water* **2019**, *11*, 705. [CrossRef]
- 115. Abbasi, A.; Khalili, K.; Behmanesh, J.; Shirzad, A. Drought monitoring and prediction using SPEI index and gene expression programming model in the west of Urmia Lake. *Theor. Appl. Climatol.* **2019**, *138*, 553–567. [CrossRef]
- Mouatadid, S.; Raj, N.; Deo, R.C.; Adamowski, J.F. Input selection and data-driven model performance optimization to predict the Standardized Precipitation and Evaporation Index in a drought-prone region. *Atmos. Res.* 2018, 212, 130–149. [CrossRef]
- 117. Li, M.-F.; Tang, X.-P.; Wu, W.; Liu, H.-B. General models for estimating daily global solar radiation for different solar radiation zones in mainland China. *Energy Convers. Manag.* **2013**, *70*, 139–148. [CrossRef]
- 118. Shiri, J.; Kişi, Ö. Comparison of genetic programming with neuro-fuzzy systems for predicting short-term water table depth fluctuations. *Comput. Geosci.* 2011, 37, 1692–1701. [CrossRef]
- Dawson, C.W.; Abrahart, R.J.; See, L.M. HydroTest: A web-based toolbox of evaluation metrics for the standardised assessment of hydrological forecasts. *Environ. Model. Softw.* 2007, 22, 1034–1052. [CrossRef]
- 120. Mazaheri, H.; Ghaedi, M.; Ahmadi Azqhandi, M.H.; Asfaram, A. Application of machine/statistical learning, artificial intelligence and statistical experimental design for the modeling and optimization of methylene blue and Cd(ii) removal from a binary aqueous solution by natural walnut carbon. *Phys. Chem. Chem. Phys.* 2017, 19, 11299–11317. [CrossRef]
- Achour, K.; Meddi, M.; Zeroual, A.; Bouabdelli, S.; Maccioni, P.; Moramarco, T. Spatio-temporal analysis and forecasting of drought in the plains of northwestern Algeria using the standardized precipitation index. *J. Earth Syst. Sci.* 2020, 129, 42. [CrossRef]
- 122. Adamowski, J.; Fung Chan, H.; Prasher, S.O.; Ozga-Zielinski, B.; Sliusarieva, A. Comparison of multiple linear and nonlinear regression, autoregressive integrated moving average, artificial neural network, and wavelet artificial neural network methods for urban water demand forecasting in Montreal, Canada. *Water Resour. Res.* **2012**, *48*, 1–14. [CrossRef]
- 123. Ait Belale, R.; M'Hamdi Alaoui, F.E.; Chhiti, Y.; Sahibeddine, A.; Munoz Rujas, N.; Aguilar, F. Study on the thermophysical properties of waste cooking oil biodiesel fuel blends with 1-butanol. *Fuel* **2021**, *287*, 119540. [CrossRef]