



LJMU Research Online

Jalil, B, Naser, AY, Prieto, JM and Heinrich, M

Herbal supplements in Jordan: a cross-sectional survey of pharmacists' perspectives and knowledge

<http://researchonline.ljmu.ac.uk/id/eprint/17365/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Jalil, B, Naser, AY, Prieto, JM and Heinrich, M (2022) Herbal supplements in Jordan: a cross-sectional survey of pharmacists' perspectives and knowledge. BMJ Open, 12 (7). e057405-e057405. ISSN 2044-6055

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

BMJ Open Herbal supplements in Jordan: a cross-sectional survey of pharmacists' perspectives and knowledge

Banaz Jalil ,¹ Abdallah Y Naser ,² Jose M Prieto,³ Michael Heinrich¹

To cite: Jalil B, Naser AY, M Prieto J, *et al.* Herbal supplements in Jordan: a cross-sectional survey of pharmacists' perspectives and knowledge. *BMJ Open* 2022;**12**:e057405. doi:10.1136/bmjopen-2021-057405

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057405>).

Received 15 September 2021
Accepted 05 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Pharmacognosy and Phytotherapy, UCL School of Pharmacy, 29 - 39 Brunswick Sq, London, UK

²Department of Applied Pharmaceutical Sciences and Clinical Pharmacy, Isra University, Amman, Jordan

³School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool, UK

Correspondence to

Dr Banaz Jalil; b.jalil@ucl.ac.uk

ABSTRACT

Objectives Pharmacists are ideal partners for engaging with the needs and expectations of patients. They can play a vital role by providing information and supplying herbal medicines. In some community settings, pharmacists are also the main first point of care. This study explored Jordanian community pharmacists' perspectives and knowledge of herbal medicines available in pharmacies.

Design A cross-sectional study using an online survey was developed, and it was distributed via social media platforms. The one-way analysis of variance (ANOVA) test was used to compare the mean knowledge scores between different demographic groups. Multiple linear regression analysis was used to identify predictors of herbal medicines knowledge.

Setting Jordanian community pharmacies.

Participants 401 Jordanian community pharmacists.

Results Herbal supplements are sold in practically all pharmacies (98.5%). Slimming aids (14.7%), followed by sexual and sports enhancements (14%) and maintaining general health (12.1%) were most requested by Jordanian customers. While supplements for maintaining general health (12%), followed by slimming aids (11.4%) and skin conditions (9.3%) were most recommended by Jordanian pharmacists. 63.1% were not aware of potential herb–drug interactions, 95.6% did not receive complaints from customers about herbal medicines and 41.2% would not report adverse reactions to the national pharmacovigilance services. The mean knowledge score for knowledge of use, regulation, adverse reactions, and drug interactions was 3.7 (SD: 0.7), 3.5 (SD: 0.8), 3.6 (SD: 0.8), and 3.6 (SD: 0.8) (out of 5), respectively. ANOVA test showed that total pharmacists' knowledge scores significantly differed based on the length of time practising pharmacy ($p < 0.05$).

Conclusion This study highlights some key concerns relating to recommendations, awareness and reporting of herbal medicines among Jordanian community pharmacists. Pharmacists need enhanced education to provide objective and evidence-based information on the benefits–risks of herbal medicines. Future studies need to be carried out to confirm whether our findings are transferable to other Middle Eastern countries.

INTRODUCTION

Herbal products are used as supplements with preventive health benefits, as herbal medicines, in cosmetics and in a wide range of other products which are of

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study surveyed community pharmacists' perception, recommendation, and knowledge of the use and safety of herbal supplements. It presents the first study conducted in the Middle East assessing pharmacists' general awareness, experience with customers and views on reporting herbal supplements-related adverse reactions.
- ⇒ Community pharmacists are frontline healthcare professionals regularly asked to advise customers about herbal supplements.
- ⇒ Data collections are based on self-reporting from the pharmacist rather than direct observation. Being a self-reported questionnaire, response bias is likely.
- ⇒ Future qualitative studies seeking community pharmacists' perception and knowledge of herbal supplements may be needed to gauge concerns and complement our quantitative study results in the wider Middle East.

medical-pharmaceutical relevance. Large sections of the population in low/middle-income countries still rely on traditional practitioners, medicinal plants and herbal medicines for their primary healthcare,^{1 2} with the increasing importance of commercially available supplements and medical products. In the Middle East, diverse legislative approaches and classifications of herbal medicines were implemented in response to the public interest, market value and the potential safety issues of unregulated herbal supplements. In a global report by the WHO (2019), Jordan was reported as not currently having a national policy on the use of traditional medicine/complementary medicine (TM/CM); however, laws and regulations which had been established in 2001 were updated in 2016.³ In addition, there are no independent national research institutes focusing on TM/CM or herbal medicines. The regulations of herbal medicines are partly the same as those for conventional pharmaceuticals. In Jordan, herbal medicines categorised as prescription medications



are sold in pharmacies (ie, with medical claims), which require a prescription, while herbal supplements sold in pharmacies and outlets as over-the-counter (OTC) products (ie, with claims on health benefits and nutrient content⁴) are dispensed by a pharmacist without needing a prescription.

In recent years, there has been an increased global interest in vitamins and herbal supplements. In 2017, their global market exceeded US\$100 billion⁵ and in 2021, the Europe market was worth US\$19.96 billion and is estimated to reach US\$26.71 billion by 2026.⁶ However, this widespread use of herbal supplements might be associated with health risks related to medicinal plants' compositions, their inappropriate use/dosage or the quality of the products (eg, adulteration and contamination).⁷⁻¹⁰ Moreover, due to the common use of these products in patients with comorbidities, interactions with prescription and OTC medications are possible, leading to a decrease in the effectiveness of pharmacological treatments or toxic manifestations.¹¹ The health consequences of these interactions were widely reported in both case reports and clinical studies.¹²⁻²⁰ Patients taking anticancer, cardiovascular and immunosuppressant medications were predominantly affected by these interactions.²¹ With the lack of awareness in identifying and reporting these reactions among the public and healthcare professionals, this risk could be amplified. Among healthcare professionals, pharmacists are ideal partners for engaging with the needs and expectations of patients, and they could play a vital role by providing information and supplying herbal supplements.^{10 22-32} In addition, healthcare professionals' perceptions and attitudes of these products may influence the patients' self-care decisions.

In Middle Eastern countries, such as Saudi Arabia, Jordan, Qatar and Kuwait, several studies reported relatively positive attitudes towards the role of community pharmacists. For example, in Jordan, most participants (62.7%) reported their willingness to seek pharmacists' advice when the condition was not so severe, requiring the visit of the physician.³³ In Saudi Arabia, 43% acknowledged the pharmacists' role in solving medication-related problems, 34% considered the pharmacist a healthcare provider, and 44.6% felt that the pharmacist is indispensable and an active part of the healthcare system.³⁴ In Qatar, most patients agreed that community pharmacists should provide them with the directions for use of the medication (93%) and advise them about the treatment of minor health conditions (79%).³⁵ In addition, treatment of minor health conditions, consultation on OTC medications and accuracy checking of prescription medications were among the expected duties of community pharmacists.³⁶

While some publications have described the current pharmacy education in the Middle East,³⁷⁻³⁹ no recent studies were identified on the education in pharmacognosy and phytochemistry in the pharmacy curriculum in the Middle East. However, in 2017, a study in Jordan

reported that Jordanian universities have one or two pharmacognosy and phytochemistry courses in the clinical phytotherapy domain for the undergraduate BPharm and PharmD degree levels.⁴⁰ The PharmD degree level, which two public universities provide, also incorporates an additional course on the chemistry of medicinal plants with their approved therapeutic indications, dosage and precautions.⁴⁰

With the increased use of herbal medicines in Jordan, primarily as self-medication for infertility, diabetes, dyslipidaemia and hypertension,⁴¹⁻⁴⁸ several significant adverse reactions and herb-drug interactions are poorly understood problems. This is exacerbated by a lack of understanding, reporting and awareness of these reactions among healthcare professionals, particularly pharmacists. To provide appropriate patient advice, the pharmacist must have good knowledge of herbal medicines, particularly those available in pharmacies.

Several studies conducted in the Middle East measure pharmacists' attitudes, knowledge and views of herbal medicines⁴⁹⁻⁵¹ or pharmacists' awareness of adverse reactions and herb-drug interactions of specific herbal supplements.^{42 52-56} Another theme is the Jordanian pharmacists' and herbalists' opinions on education related to herbal medicines⁴⁰ and the prevalence of and awareness of the use of herbal medicines among pharmacy staff.⁵⁷ No studies focused on Jordanian pharmacists' perspective and knowledge of herbal medicines or on assessing the general awareness, experiences with customers and views on reporting of related adverse reactions. Therefore, this study aims to assess Jordanian pharmacists' knowledge of use, safety, awareness and willingness to report adverse reactions associated with herbal medicine use.

METHODS

Study design and population

Between May and July 2020, a cross-sectional online survey was conducted using social media platforms such as Facebook and Twitter, which were used to distribute the questionnaire link since social media are a valuable tool for conducting this type of data collection.⁵⁸ The study aims and objectives were clearly explained at the beginning of the survey. In addition, the participants were informed that participation in the study was completely voluntary. The study population included only pharmacists working at community pharmacies in Jordan.

Sample size calculation

Of the 24747 licensed pharmacists working in Jordan, 7525 are estimated to work in community pharmacies. The minimum recommended sample size for our survey study was calculated using the Raosoft sample size calculator ($n = N \times x / (n - 1) E^2 + x$), where n is the sample size required, N is the population size, x is the CI, which was considered to be 95%, and E is the margin of error, which is 5%.

The minimum recommended sample size was calculated to be 366. Therefore, in our study, the questionnaire link was kept open until 467 responses were collected to ensure that sufficient eligible participants would be included.

Patient and public involvement

The aim of this study was to explore Jordanian community pharmacists' perspectives and knowledge of herbal medicines available in pharmacies. The target population of this study was community pharmacists. At the design of the study, we decided that the input of pharmacists and their associations was the most important basis for developing the study. Although there was no direct involvement of patients or the public in the design, or conduct, or reporting, or dissemination plans of our study, the outcomes could potentially benefit the public by enhancing the safe and evidence-based use of herbal medicines. The results of this study will be freely available to participants via publication.

Development of survey questionnaire

The survey questionnaire was developed based on extensive literature review to gather pharmacists' views, knowledge, and practices in relation to the herbal medicines and supplements available in pharmacies^{22–24 59 60} and from a previous survey study conducted in another Middle Eastern country.⁶¹ The online questionnaire comprised five main sections with 19 questions covering the following components:

- A. Participants' demographic data, such as age, gender, the university from where they graduated, qualifications, place of practice, length of time practising pharmacy, and degree or further training on herbal medicine.
- B. Five questions covering uses, recommendations of herbal medicines and pharmacists' advice on these products. The response for this part was provided in the form of 'yes and no' for the advice question and in the form of a 7-point rank order: 1=most requested/recommended; 7=least requested/recommended for the use and recommendation questions.
- C. Two questions designed to determine views and recommendations of herbal medicines in minor or long-term conditions. The response for this part was provided in the form of 'yes', 'no' and one answer option to select from (eg, herbal medicine, OTC medication, both or others).
- D. Four questions, including self-assessed knowledge about uses, regulations, adverse reactions and drug interactions of herbal medicines. The response for this part of the questionnaire was provided in the form of a 5-point Likert scale that ranged from excellent to terrible, with the value of 1 assigned to the response 'terrible' and 5 to the response 'excellent'. Based on the use of Likert scale, the maximum obtainable self-assessed knowledge score for the four dimensions is 20.

- E. Three questions based on pharmacists' practices towards experiences, awareness and reporting of adverse reactions relating to herbal medicines. The response for this part was provided in the form of 'yes' and 'no'.

The questionnaire took, on average, 10–15 min to complete. To minimise any misunderstanding between the terms of 'herbal medicine/supplements' and 'OTC' medication/drugs by the participants, the terms were clearly defined on the first page of the questionnaire. The questionnaires' content was reviewed, and it was piloted with five pharmacists who are native Arabic speakers and are practising pharmacy in the Middle East (eg, Jordan, Iraq, Kuwait, Lebanon and Saudi Arabia) and those whose first language is not English. This was explored to test the clarity and the logical flow of the questionnaire and since the questionnaire was not in Arabic, which is the first language in the country of research (ie, Jordan).

Statistical analysis

The SPSS software, V.26, was used to analyse the results. Descriptive statistics were used to describe demographic data, using percentages and frequencies to express categorical variables. Kolmogorov-Smirnov, Shapiro-Wilk tests and histogram were used to check the normality of the data, which showed that the data are normally distributed. Pharmacists' knowledge scores were interpreted as a continuous scale based on the scale midpoint (10 out of 20), where scores above the midpoint identified better knowledge about herbal medicines. The one-way analysis of variance (ANOVA) test was used to compare the mean knowledge scores between different demographic groups. Multiple linear regression analysis was used to identify predictors of herbal medicine knowledge.

RESULTS

Participants' characteristics

A total of 467 responses were obtained from distributing the survey link through social media. Of those, 58 were excluded as they did not meet one or more of the inclusion criteria (ie, Jordanian pharmacists currently practising pharmacy in other Middle Eastern countries). In addition, a further eight responses were also excluded as the participants only provided the demographic information and no data on the survey as such, leaving a total of 401 surveys for analysis. These 401 are practising community pharmacy in Jordan (for demographic details, see [table 1](#)).

The uses and advice on herbal supplements

Nearly all (98.5%) of the participants reported the availability of herbal supplements in their community pharmacies. Three hundred and eight (76.8%) of the participants reported that their customers ask for herbal supplements with those promoted were for weight loss (slimming aids) (14.7%), followed by supplements for sexual and sports enhancements (14%) and maintaining general health (12.1%) being customers' most commonly

**Table 1** Characteristics of study participants (n=401)

Characteristics	Frequency (%)
Gender	
Male	223 (55.6)
Female	178 (44.4)
Age (years)	
21–30	258 (64.3)
31–40	94 (23.4)
41–50	38 (9.5)
51–60	11 (2.7)
Over 61	0
Qualifications (highest level)	
Bachelor's in pharmacy or diploma	379 (94.5)
Masters	18 (4.5)
PharmD	3 (0.7)
PhD qualifications	1 (0.2)
Institution/university they graduated from (n=380)	
Governmental universities	
University of Jordan	96 (24.3)
Jordan University of Science and Technology	20 (5.1)
The Hashemite University	5 (1.3)
Mutah University	4 (1.0)
Al-Balqa Applied University	3 (0.8)
Private universities	
Al-Isra University	112 (28.4)
Al-Zaytoonah University	46 (11.6)
University of Petra	23 (5.8)
Al-Ahliyya Aman University	22 (5.6)
Al-Zarqa Private University	12 (3.0)
Applied Science Private University	7 (1.8)
Middle East University	7 (1.8)
Philadelphia University in Jordan	5 (1.3)
Amman Training Centre	5 (1.3)
Jerash University	3 (0.8)
Others	9 (2.7)
Non-Jordanian universities (n=15)	
One University	9 (2.3)
Al-Khawarizmi University	2 (0.5)
Al-Quds University	2 (0.5)
Al-Karak University	2 (0.5)
Length of time practising pharmacy (years)	
1–10	351 (87.5)
11–20	36 (9.0)
21–30	12 (3.0)
Over 31	2 (0.5)
Degree/further training on herbal medicines	
Yes degree/further training	13 (3.2)
No degree/further training	388 (96.8)

requested supplements. Other categories were supplements promoted to treat chronic and acute health conditions, skin conditions, relieve stress and dissatisfaction with prescription medications and were the least ranked categories requested by Jordanian customers (n=298) (table 2).

Three hundred and nine (78.2%) of the participants said they would recommend herbal supplements to their customers (n=300) with supplements promoted for maintaining general health (12%), followed by slimming aids (for weight loss) (11.4%) and skin conditions (9.3%) being study participants' most commonly recommended. In addition, supplements for sexual and sports enhancements (9.1%) ranked fourth among the categories. Other categories were supplements promoted to treat chronic and acute health conditions, dissatisfaction with prescription medications and relief of stress and were ranked the least recommended categories by the participants (table 3). In general, there are some differences with respect to what is being sold versus what is being recommended by the participants. As for advice, 376 (96.9%) participants reported advising customers about the safety of herbal medicines (n=388).

Views and recommendations of herbal supplements

Half of the participants (50.4%; n=194) would recommend OTC medications for minor health conditions, while roughly one-third (35.1%; n=135) said they would recommend herbal medicines. A total of 14.3% (n=55) of the participants said their recommendation is based on the minor health condition and 0.3% for others (n=385).

When participants were asked about their recommendations to customers with long-term conditions, 83.6% (n=321) said they would not recommend the concomitant use of herbal medicines and prescription medications. In comparison, 16.4% (n=63) said they would recommend the concomitant use of herbal medicines and prescription medications (n=384).

Knowledge of herbal supplements

Among those who completed the knowledge of use of herbal medicines question (n=386), 63.2% reported 'good' knowledge, while for knowledge of regulation (n=385), 56.4% claimed to have 'good' knowledge on regulation. Furthermore, on knowledge of adverse reactions, 56.1% said they acquire 'good' knowledge, and for knowledge of herb–drug interactions (n=384), 56.8% said they have 'good' knowledge (table 4).

The mean knowledge score for the participants was 14.2 (SD: 2.8) out of 20 (which is equal to 71.0% out of the total score). The mean knowledge score for knowledge of use, knowledge of regulation, knowledge of adverse reactions, and knowledge of drug interactions was 3.7 (SD: 0.7), 3.5 (SD: 0.8), 3.6 (SD: 0.8), and 3.6 (SD: 0.8) (out of 5), respectively. ANOVA test showed that total pharmacists' knowledge scores significantly differed based on length of time practising pharmacy (p<0.05), indicating those with 11–20 years of experience had the highest

Table 2 Categories of herbal supplements as requested by the Jordanian customers (n=298)

	Most requested	2nd choice	3rd choice	4th choice	5th choice	6th choice	Least requested
Maintenance of general health	12.1%	20.8%	22.1%	19.8%	13.1%	6.7%	5.4%
Dissatisfaction with prescription medications	5.7%	11.4%	21.1%	21.8%	13.1%	13.8%	13.1%
For treatment of acute & chronic health conditions	5.4%	9.0%	18.1%	22.7%	10.0%	17.7%	17.1%
To relieve stress	6.7%	16.4%	17.1%	18.4%	15.7%	11.4%	14.4%
For weight loss	14.7%	17.7%	21.7%	18.7%	12.7%	5.7%	9.0%
For sexual & sports enhancement	14.0%	21.4%	21.7%	14.4%	14.7%	6.4%	7.4%
For skin conditions	5.4%	12.7%	19.1%	19.1%	15.4%	13.4%	15.1%

Measured on a 7-point rank order: 7=least requested; 1=most requested.

knowledge scores, followed by those with 21–30 years of experience, then those with 1–10 years of experience. Multiple linear regression analysis showed that having a PhD qualification is associated with a lower knowledge score ($p<0.001$) (table 5).

Awareness and reporting of adverse reactions related to herbal supplements

Out of all 385 participants who completed the question, 243 (63.1%) of participants were not aware of interactions between herbal medicines and prescription medications. In contrast, only 142 (36.9%) were aware of these possible interactions. Very few participants provided examples of herbs with the potential of drug interactions (n=4), such as St John's wort, Gingko biloba and warfarin with vitamin K supplements.

In addition, 386 (95.6%) of participants did not come across any adverse reactions, while 17 (4.4%) claimed to have encountered adverse reactions associated with herbal medicines (n=385). Limited examples were provided about adverse reactions encountered by the participants, such as hypoglycaemia, allergic reactions and fertility issues with herbal medicine (n=3).

When participants were asked about their view of reporting the adverse reactions of herbal medicines to

the national pharmacovigilance services in the country (n=388), 228 participants (58.8%) would report adverse reactions from herbal medicines to the national pharmacovigilance services, while 160 (41.2%) would not consider reporting it.

DISCUSSIONS

The prevalence of herbal medicine use continues to grow globally, including in the Middle East^{62–64} and specifically among the Jordanian population.^{45 65} This, combined with the prevalence of chronic diseases,⁶⁶ the concomitant use of herbal medicines and prescription medications⁶⁷ and a lack of knowledge about herb–drug interactions among Jordanians, poses a poorly understood challenge to the healthcare professionals.⁴⁴

Our study investigates customers' use and pharmacists' recommendation of herbal supplements, as well as pharmacists' knowledge, willingness to advise, awareness of safety and their views of reporting herbal-related adverse reactions in Jordan. It presents one of a few studies conducted in the Middle East assessing pharmacists' perspectives and knowledge of herbal medicines available in pharmacies.^{42 49 50 52 53 55 56} This study reinforces that

Table 3 Categories of herbal supplements as recommended by the participants (n=300)

	Most recommended	2nd choice	3rd choice	4th choice	5th choice	6th choice	Least recommended
Maintenance of general health	12.0%	19.1%	19.7%	20.7%	16.4%	7.0%	5.0%
Dissatisfaction with prescription medications	5.0%	11.3%	18.0%	23.3%	14.7%	12.3%	15.3%
For treatment of acute & chronic health conditions	4.7%	10.7%	18.3%	17.3%	13.7%	16.7%	18.7%
To relieve stress	5.3%	17.3%	19.0%	21.3%	12.7%	10.3%	14.0%
For weight loss	11.4%	19.8%	19.5%	17.4%	12.8%	8.4%	10.7%
For sexual & sports enhancement	9.1%	17.8%	22.6%	17.2%	14.5%	8.1%	10.8%
For skin conditions	9.3%	10.0%	21.0%	20.0%	12.7%	12.0%	15.0%

Measured on a 7-point rank order: 7=least recommended; 1=most recommended.

**Table 4** Participants' self-reported knowledge on use (n=386), regulations (n=385), adverse reactions (n=387) and drug interactions (n=384)

	Knowledge of the use of herbal medicines	Knowledge of regulation of herbal medicines	Knowledge of adverse reactions of herbal medicines	Knowledge of drug interaction of herbal medicines
Excellent	5.7%	5.2%	6.5%	6.5%
Good	63.2%	56.4%	56.1%	56.8%
Average	22.5%	25.2%	25.1%	24.5%
Poor	7.8%	11.9%	11.9%	11.7%
Terrible	0.8%	1.3%	0.5%	0.5%

Measured on a 5-point Likert scale: 5=excellent; 4=good; 3=average; 2=poor; 1=terrible.

pharmacists worldwide have frontline role in providing advice to customers about these products.^{10 22–32}

Our study shows that herbal supplements are readily available in practically all pharmacies surveyed. This correlates with an increased demand from the public and pharmacists' recommendation to use supplements promoted for slimming and sexual and sports enhancements. The consumption of these supplements without medical advice or proper follow-up is a concern due to the nature of the therapeutic claims and lack of information provided with these products.⁶⁸ Moreover, the current regulatory framework may not ensure their quality and safety,^{10 68 69} including a high risk of adulteration with undeclared synthetic substances, usually consisting of the addition of prescription or banned medications.^{61 70–73} This clearly poses a health risk to the customers. In line with these findings, the use of a diverse range of slimming aids among Jordanians was reported,⁷⁴

with the majority reporting the use of these supplements in the 3 months prior to the survey.⁷⁴ According to Abdel-Qader *et al*,⁶⁵ Jordanians' use of herbal supplements is for chronic diseases and obesity, while sexual enhancers accounted for the lowest share.⁶⁵ In Jordan, limited pharmaceutical options for weight management exist (the sole registered pharmaceutical agent is orlistat). Treatment is quite expensive compared with individual income levels. Furthermore, orlistat has multiple side effects that restrict its usability among its users, including gastrointestinal upset and stomach and back pain. The proportion of people who use sexual enhancers is much higher than indicated, and thus research focusing on the use of sexual enhancers was suggested.⁶⁵

Pharmacists' self-reported knowledge score on the herbal medicine uses, their regulations, potential adverse reactions and drug interactions of herbal medicines is 'good', with the total knowledge score being higher on

Table 5 Multiple linear regression analysis

Model	Unstandardised coefficients		Standardised coefficients		Sig.	95.0% CI for B	
	B	SE	Beta	t		Lower bound	Upper bound
(Constant)	14.629	3.796		3.854	0.000	7.164	22.093
Age							
31–40 years	0.536	0.344	0.083	1.558	0.120	–0.141	1.213
41–50 years	–0.435	0.570	–0.046	–0.763	0.446	–1.555	0.686
51–60 years	0.630	1.195	0.038	0.527	0.598	–1.719	2.979
Qualifications (highest level)							
Bachelor's in pharmacy or diploma	0.550	1.582	0.046	0.348	0.728	–2.561	3.661
Masters	0.674	1.705	0.051	0.395	0.693	–2.680	4.027
PhD qualifications	–14.148	3.080	–0.260	–4.593	0.000***	–20.205	–8.091
Length of time practising pharmacy (years)							
1–10	0.149	2.864	0.016	0.052	0.958	–5.483	5.781
11–20	–0.453	2.788	–0.028	–0.162	0.871	–5.934	5.029
21–30	–1.208	2.891	–0.145	–0.418	0.676	–6.893	4.476
Gender (female)	–0.468	0.277	–0.084	–1.689	0.092	–1.012	0.077
Degree/further training on herbal medicines (no)	0.329	1.123	0.015	0.293	0.770	–1.880	2.538

***P<0.001.

use than the one on their regulations, potential adverse reactions and herb–drug interactions. Similar findings were reported from studies conducted in Kuwait, Saudi Arabia, Palestine, Iraq, Lebanon and the United Arab Emirates,^{15 24–27 30 31} which showed better knowledge of the herbal medicine uses among pharmacists,^{49 50 53 55 56} in comparison with their knowledge of the potential adverse reactions and herb–drug interactions.^{49 50 53 55 56} In addition, we found a significant inverse ($p < 0.05$) correlation between self-reported ‘total pharmacists’ knowledge’ and ‘years of experience’. A possible explanation for this might be that newly graduated pharmacists had taken additional relevant courses about herbal medicines during their undergraduate studies. In support of this, two recent studies conducted in Jordan and Palestine exploring pharmacists’ knowledge reported similar findings relating to knowledge scores and years of experience.^{50 57} In contrast, other studies in the Middle East reported higher knowledge scores among those who received education/training/qualification on herbal medicines,^{49 52} or pharmacists’ knowledge was influenced by the place of work (ie, government or private sector),⁵⁶ which was not in our study. On the other hand, studies conducted in other parts of the world showed that the pharmacists’ knowledge increased with an increase in years of experience.^{22 75–77} Clearly, pharmacists gain more knowledge over time. Therefore, further research is needed to gain better understanding of whether this difference is influenced by differences in pharmacists’ education and training across different geographical locations.

Pharmacists’ knowledge, their experience of customer-reporting adverse reactions and their awareness of these reactions could play a vital role in identifying and reporting these reactions. More than half of the participants responding ($n=243$) were not aware of potential herb–drug interactions, and an even larger majority ($n=386$) did not receive complaints from the customers about herbal medicines. In contrast, more than a quarter of the Jordanian customers experienced unpleasant symptoms from herbal medicines, including constipation, diarrhoea, nausea/vomiting, palpitation, sweating, headache, fever and allergic reactions.⁶⁵ A possible reason for this variation in experience between Jordanian pharmacists and customers in our and other studies could be linked to these supplements being purchased from herbal shops and online rather than from pharmacies. Hence, the adverse reactions may have been reported back to the herbalists rather than to the pharmacists or not reported at all. In fact, herbal shops were cited by Abdel-Qader *et al*⁶⁵ as Jordanians’ first choice to purchase herbal supplements, followed by pharmacies as their second choice.⁶⁵ Another explanation for this variation could be the pharmacists’ lack of training relating to the knowledge of and awareness to identify adverse reactions.³¹ Generally, findings across the literature investigating pharmacists’ education related to herbal medicines were limited or non-existent, particularly in the Middle East. A previous

Jordanian cross-sectional study found that none of the surveyed pharmacists received guidelines on dietary herbal supplements during their formal training before graduation.⁷⁸

Participants’ lack of awareness and experience with customers reporting complaints linked to herbal medicines were coupled with more than a quarter not willing to report the adverse reactions to the national pharmacovigilance centre. Adverse reactions from using herbal medicines could be anything from a minor adverse reaction caused by the medicinal plants’ composition, such as heartburn caused by peppermint, to life-threatening reactions like cancer, hepatotoxicity or nephrotoxicity caused by *Aristolochia* spp.^{79–82} In addition, the adverse reaction of herbal medicines could be caused in situations of low-quality herbal medicines or the misidentified, adulterated or contaminated herbs being used.^{80 83–85} Based on the national pharmacovigilance database in Jordan, between 2010 and 2014, most medications implicated in adverse reactions were antineoplastics (37.6%), followed by immunomodulators (14.1%), antibiotics (10.3%) and analgesics (6.6%).⁸⁶ Although, in the past years, there has been a fourfold increase in adverse reactions reported, no reports were received for adverse reactions related to herbal medicines. This highlights some concerns related to the identification and recognition of these reactions leading to under-reporting of such reactions⁸⁷ by the public or pharmacists. Several barriers, such as lack of time, lack of knowledge about herbal medicines, the complexity of the reporting process and pharmacists’ perception of their professional roles, could also be contributing to the issue of under-reporting.⁸⁸

Our findings reinforce the importance of improving the public’s and pharmacists’ awareness and ability in recognising and reporting adverse reactions after consumption of herbal medicines. It also contributes to the important global debate on how to improve pharmacists’ knowledge and awareness of the safety of herbal medicines available in pharmacies.^{25–27 29 32 49 50 52–56 60} This, in turn, will impact the public’s awareness and confidence and, thus in, safety associated with the use of such preparations.

There remain some important limitations. The survey link was completed online and circulated via social media. Snowballing cannot ascertain a representative distribution or completion of the survey. This might have affected the generalisability of our findings as we might have missed some of the targeted population. Although the formal pharmacy education in Jordan is in the English language, the survey questions were in English and not in the Arabic language, which might have affected the participants’ understanding of the survey items. Also, data collection is based on self-reporting by the participants, therefore, response bias is likely. Furthermore, although the survey had a good response rate from those who completed bachelor’s degree in pharmacy, there were very few responses from those who have PharmD or PhD qualifications. Therefore, our findings should be interpreted carefully.

Implications of findings to practice

The undergraduate pharmacy core curriculum in Jordan should be encouraged to include more formal academic training about the pharmacological and toxicological effects of herbal medicines, as well as on the detection, understanding and reporting of adverse reactions. Although participants generally reported a good self-perceived knowledge of herbal medicines, there is still a need to improve the breadth and scope of this education, particularly in areas of safety, awareness and reporting related to potential adverse reactions. Continuous professional development in these areas will also be warranted. This will help give pharmacists first-hand experience of what is expected in practice. In addition, it can provide Jordanian pharmacists with good knowledge about the uses and the safety of herbal medicines available in pharmacies. Hence, future pharmacists could provide more objective and evidence-based information on the risks and benefits of herbal medicines to customers. This improved knowledge will be important to enhance public safety.

In conclusion, the current study has revealed some specific problems relating to awareness, identification and reporting of potential adverse reactions related to herbal medicines among Jordanian pharmacists. In addition, the uses and recommendations of slimming aids and sexual and sports enhancement products are likely to be a customer health risk due to possible adulteration in these supplements. The limited research on awareness and reporting of potential adverse reactions of herbal medicines among pharmacists has wider implications on practice and customers' safety, particularly in the Middle East. In the pharmacy curriculum, the safety of herbal medicines should be covered more thoroughly and reinforced through continuous professional development. Hence, future pharmacists can provide evidence-based information and advise customers more effectively about the safe use of herbal supplements. Furthermore, raising awareness and educating the public about the risks and benefits of herbal supplements available on the market will also be warranted.

Contributors This study is a part of BJ's PhD dissertation (2021) at the UCL School of Pharmacy. BJ and MH developed the concept for this study and the main tool. JMP primarily supervised most of JB's PhD project and contributed to the method development and interpretation. AYN contributed to developing country-specific elements of the questionnaire, its dissemination in Jordan and statistical analysis. All authors contributed to the writing and approved the manuscript.

Funding This work was (partially) supported by the Schlumberger Foundation Faculty for the Future programme awarded to BJ (award/grant number is not applicable).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval This study involves human participants and the ethical approvals for this study were obtained from the Jordanian Ministry of Health (Jordan reference no. mba/ethics committee/2064), Jordanian Pharmacists Association and UCL Research Ethics Committee (University College London, UK reference no. 9059/002).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. No additional data available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Banaz Jalil <http://orcid.org/0000-0002-1535-3744>

Abdallah Y Naser <http://orcid.org/0000-0001-8440-7446>

REFERENCES

- 1 Harvey AL. Natural products in drug discovery. *Drug Discov Today* 2008;13:894–901.
- 2 Williamson EM. Drug interactions between herbal and prescription medicines. *Drug Saf* 2003;26:1075–92.
- 3 World Health Organization. *WHO global report on traditional and complementary medicine 2019*. Geneva, 2019.
- 4 World Health Organization. *Who global report on traditional and complementary medicine 2019*. World Health organization. Report No.: 9241515430 2019.
- 5 Asmelashe Gelayee D, Binega Mekonnen G, Asrade Atnafe S, et al. Herbal medicines: personal use, knowledge, attitude, dispensing practice, and the barriers among community pharmacists in Gondar, Northwest Ethiopia. *Evid Based Complement Alternat Med* 2017;2017:6480142.
- 6 Forecast MD. Europe herbal supplements market, 2021. Available: <https://www.marketdataforecast.com/market-reports/europe-herbal-supplements-market>
- 7 Azaizeh H, Saad B, Khalil K, et al. The state of the art of traditional Arab herbal medicine in the eastern region of the Mediterranean: a review. *Evid Based Complement Alternat Med* 2006;3:229–35.
- 8 Ernst E. Adulteration of Chinese herbal medicines with synthetic drugs: a systematic review. *J Intern Med* 2002;252:107–13.
- 9 Booker A, Heinrich M. Value chains of botanical and herbal medicinal products: a European perspective. *HerbalGram* 2016;112:40–5.
- 10 Heinrich M. Quality and safety of herbal medical products: regulation and the need for quality assurance along the value chains. *Br J Clin Pharmacol* 2015;80:62–6.
- 11 Cuzzolin L, Zaffani S, Benoni G. Safety implications regarding use of phytomedicines. *Eur J Clin Pharmacol* 2006;62:37–42.
- 12 Anlamert W, Sermappasak P, Yokubol D, et al. Pomelo enhances cyclosporine bioavailability in healthy male Thai volunteers. *J Clin Pharmacol* 2015;55:377–83.
- 13 Showande SJ, Adegbolagun OM, Igbinoba SI, et al. In vivo pharmacodynamic and pharmacokinetic interactions of *Hibiscus sabdariffa* calyces extracts with simvastatin. *J Clin Pharm Ther* 2017;42:695–703.
- 14 Cheng Y-Y, Hsieh C-H, Tsai T-H. Concurrent administration of anticancer chemotherapy drug and herbal medicine on the perspective of pharmacokinetics. *J Food Drug Anal* 2018;26:S88–95.
- 15 Wilson V, Maulik SK. Herb-Drug interactions in neurological disorders: a critical appraisal. *Curr Drug Metab* 2018;19:443–53.
- 16 Choi S, Oh D-S, Jerng UM. A systematic review of the pharmacokinetic and pharmacodynamic interactions of herbal medicine with warfarin. *PLoS One* 2017;12:e0182794.
- 17 Asher GN, Corbett AH, Hawke RL. Common herbal dietary supplement-drug interactions. *Am Fam Physician* 2017;96:101–7.
- 18 Sprouse AA, van Breemen RB. Pharmacokinetic interactions between drugs and botanical dietary supplements. *Drug Metab Dispos* 2016;44:162–71.
- 19 Awortwe C, Bruckmueller H, Cascorbi I. Interaction of herbal products with prescribed medications: a systematic review and meta-analysis. *Pharmacol Res* 2019;141:397–408.
- 20 Babos MB, Heinan M, Redmond L, et al. Herb-Drug interactions: worlds intersect with the patient at the center. *Medicines* 2021;8:44.
- 21 Awortwe C, Makiwane M, Reuter H, et al. Critical evaluation of causality assessment of herb-drug interactions in patients. *Br J Clin Pharmacol* 2018;84:679–93.
- 22 Kwan D, Boon HS, Hirschhorn K, et al. Exploring consumer and pharmacist views on the professional role of the pharmacist with respect to natural health products: a study of focus groups. *BMC Complement Altern Med* 2008;8:40.

- 23 Nelson MV, Bailie GR, Areny H. Pharmacists' perceptions of alternative health approaches--a comparison between U.S. and British pharmacists. *J Clin Pharm Ther* 1990;15:141-6.
- 24 Kwan D, Hirschhorn K, Boon HUS. And Canadian pharmacists' attitudes, knowledge, and professional practice behaviors toward dietary supplements: a systematic review. *BMC Complement Altern Med* 2006;6:1-10.
- 25 Nathan JP, Cicero LA, Koumis T, et al. Availability of and attitudes toward resources on alternative medicine products in the community pharmacy setting. *J Am Pharm Assoc* 2005;45:734-9.
- 26 Dolder C, Lacro J, Dolder N, et al. Pharmacists' use of and attitudes and beliefs about alternative medications. *Am J Health Syst Pharm* 2003;60:1352-7.
- 27 Clauson KA, McQueen CE, Shields KM, et al. Knowledge and attitudes of pharmacists in Missouri regarding natural products. *Am J Pharm Educ* 2003;67:41.
- 28 Naidu S, Wilkinson JM, Simpson MD. Attitudes of Australian pharmacists toward complementary and alternative medicines. *Ann Pharmacother* 2005;39:1456-61.
- 29 Koh H-L, Teo H-H, Ng H-L. Pharmacists' patterns of use, knowledge, and attitudes toward complementary and alternative medicine. *J Altern Complement Med* 2003;9:51-63.
- 30 Semple SJ, Hotham E, Rao D, et al. Community pharmacists in Australia: barriers to information provision on complementary and alternative medicines. *Pharm World Sci* 2006;28:366-73.
- 31 Ng JY, Tahir U, Dhaliwal S, JY N, Barriers DS. Barriers, knowledge, and training related to pharmacists' counselling on dietary and herbal supplements: a systematic review of qualitative studies. *BMC Health Serv Res* 2021;21:499.
- 32 Waddington F, Naunton M, Kyle G, et al. A systematic review of community pharmacist therapeutic knowledge of dietary supplements. *Int J Clin Pharm* 2015;37:439-46.
- 33 Wazaify M, Al-Bsoul-Younes A, Abu-Gharbieh E, et al. Societal perspectives on the role of community pharmacists and over-the-counter drugs in Jordan. *Pharm World Sci* 2008;30:884-91.
- 34 Al-Arifi MN. Patients' perception, views and satisfaction with pharmacists' role as health care provider in community pharmacy setting at Riyadh, Saudi Arabia. *Saudi Pharm J* 2012;20:323-30.
- 35 El Hajj MS, Salem S, Mansoor H. Public's attitudes towards community pharmacy in Qatar: a pilot study. *Patient Prefer Adherence* 2011;5:405.
- 36 El Hajj MS, Mekki R, Elkaffash R, et al. Public attitudes towards community pharmacy in Arabic speaking Middle Eastern countries: a systematic review. *Res Social Adm Pharm* 2021;17:1373-95.
- 37 Kheir N, Zaidan M, Younes H, et al. Pharmacy education and practice in 13 middle Eastern countries. *Am J Pharm Educ* 2008;72:133.
- 38 Al-Wazaify M, Matowe L, Albsoul-Younes A, et al. Pharmacy education in Jordan, Saudi Arabia, and Kuwait. *Am J Pharm Educ* 2006;70:18.
- 39 Al-Ghananeem AM, Malcom DR, Shammas S, et al. A call to action to transform pharmacy education and practice in the Arab world. *Am J Pharm Educ* 2018;82:7014.
- 40 Basheti IA, Elayeh ER, Al Natour DB DB, et al. Opinions of pharmacists and herbalists on herbal medicine use and receiving herbal medicine education in Jordan. *Trop J of Pharm Res* 2017;16:689-96.
- 41 Nathan PJ, Lu K, Gray M, et al. The neuropharmacology of L-theanine(N-ethyl-L-glutamine): a possible neuroprotective and cognitive enhancing agent. *J Herb Pharmacother* 2006;6:21-30.
- 42 Khader Y, Sawair FA, Ayoub A, et al. Knowledge and attitudes of lay public, pharmacists, and physicians toward the use of herbal products in North Jordan. *J Altern Complement Med* 2008;14:1186-7.
- 43 Wazaify M, Afifi FU, El-Khateeb M, et al. Complementary and alternative medicine use among Jordanian patients with diabetes. *Complement Ther Clin Pract* 2011;17:71-5.
- 44 Wazaify M, Alawwa I, Yasein N, et al. Complementary and alternative medicine (cam) use among Jordanian patients with chronic diseases. *Complement Ther Clin Pract* 2013;19:153-7.
- 45 Issa RA, Basheti IA. Herbal medicine use by people in Jordan: exploring beliefs and knowledge of herbalists and their customers. *J Biol Sci* 2017;17:400-9.
- 46 Bardaweel SK, Shehadeh M, Suaifan GA, et al. Complementary and alternative medicine utilization by a sample of infertile couples in Jordan for infertility treatment: clinics-based survey. *BMC Complement Altern Med* 2013;13:1-7.
- 47 Afifi FU, Wazaify M, Jabr M, et al. The use of herbal preparations as complementary and alternative medicine (cam) in a sample of patients with cancer in Jordan. *Complement Ther Clin Pract* 2010;16:208-12.
- 48 El-Dahiyat F, Rashrash M, Abuhamdah S, et al. Herbal medicines: a cross-sectional study to evaluate the prevalence and predictors of use among Jordanian adults. *J Pharm Policy Pract* 2020;13:2.
- 49 Hijazi MA, Shatila H, El-Lakany A, et al. Beliefs, practices and knowledge of community pharmacists regarding complementary and alternative medicine: national cross-sectional study in Lebanon. *BMJ Open* 2019;9:e025074.
- 50 Shraim NY, Shawahna R, Sorady MA, et al. Community pharmacists' knowledge, practices and beliefs about complementary and alternative medicine in Palestine: a cross-sectional study. *BMC Complement Altern Med* 2017;17:429.
- 51 Mohammed SI, Al- Shadedi MI, Kasim AA. Knowledge, use and recommendation of Iraqi pharmacist toward complementary and alternative medicine. *IJPS* 2020;29:88-93.
- 52 Alnaim L, Almazrou S, Alsunbul M, et al. Pharmacist's knowledge and attitudes towards complementary and alternative medicine in Saudi Arabia: A cross-sectional study. *Adv Integr Med* 2018;5:96-102.
- 53 Khodour MR, Kurdi M, Hallak HO, et al. Pharmacists' knowledge, attitudes and practices towards herbal remedies in West Bank, Palestine. *Int Arch Med* 2016.
- 54 Alsayari A, Almghaslah D, Khaled A, et al. Community pharmacists' knowledge, attitudes, and practice of herbal medicines in Asir region, Kingdom of Saudi Arabia. *Evid Based Complement Alternat Med* 2018;2018:1568139.
- 55 Abahussain NA, Abahussain EA, Al-Oumi FM. Pharmacists' attitudes and awareness towards the use and safety of herbs in Kuwait. *Pharm Pract* 2007;5:125-9.
- 56 Fahmy SA, Abdu S, Abuelkhair M. Pharmacists' attitude, perceptions and knowledge towards the use of herbal products in Abu Dhabi, United Arab Emirates. *Pharm Pract* 2010;8:109-15.
- 57 Naky Y. The prevalence, attitude and awareness of herbal medicine products use among pharmacy practitioner in Jordan. *Pharm J* 2019;11.
- 58 Wilson RE, Gosling SD, Graham LT. A review of Facebook research in the social sciences. *Perspect Psychol Sci* 2012;7:203-20.
- 59 Freymann H, Rennie T, Bates I, et al. Knowledge and use of complementary medicine and alternative medicine among British undergraduate pharmacy students. *Pharm World Sci* 2006;28:13-18.
- 60 Bouldin AS, Smith MC, Garner DD, et al. Pharmacy and herbal medicine in the US. *Soc Sci Med* 1999;49:279-89.
- 61 Jalil BT. Herbal supplements in the middle East (Iraq and Jordan): regulation, quality and safety of and development of a method to detect common adulterants. Doctoral thesis (Ph.D), (University College London) 2021 <https://discovery.ucl.ac.uk/id/eprint/10128779/>
- 62 Kamel FO, Magadmi RM, Hagraas MM, et al. Knowledge, attitude, and beliefs toward traditional herbal medicine use among diabetics in Jeddah Saudi Arabia. *Complement Ther Clin Pract* 2017;29:207-12.
- 63 Awad A, Al-Shaye D. Public awareness, patterns of use and attitudes toward natural health products in Kuwait: a cross-sectional survey. *BMC Complement Altern Med* 2014;14:105.
- 64 Hwang JH, Kim Y-R, Ahmed M, et al. Use of complementary and alternative medicine in pregnancy: a cross-sectional survey on Iraqi women. *BMC Complement Altern Med* 2016;16:191.
- 65 Abdel-Qader DH, Albassam A, Ismael NS, et al. Herbal medicine use in the Jordanian population: a nationally representative cross-sectional survey. *J Pharm Pharmacog Res* 2020;8:525-36.
- 66 Mubarak FM, Froelicher ES, Jaddou HY, et al. Hypertension among 1000 patients with type 2 diabetes attending a national diabetes center in Jordan. *Ann Saudi Med* 2008;28:346-51.
- 67 Thiab S, Barakat M, Al-Qudah Raja'a, et al. The perception of Jordanian population towards concomitant administration of food, beverages and herbs with drugs and their possible interactions: a cross-sectional study. *Int J Clin Pract* 2021;75:e13780.
- 68 Raynor DK, Dickinson R, Knapp P, et al. Buyer beware? does the information provided with herbal products available over the counter enable safe use? *BMC Med* 2011;9:94.
- 69 Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol* 2014;4:177.
- 70 Tucker J, Fischer T, Upjohn L, et al. Unapproved pharmaceutical ingredients included in dietary supplements associated with us food and drug administration warnings. *JAMA Netw Open* 2018;1:e183337-e.
- 71 Ekar T, Kreft S. Common risks of adulterated and mislabeled herbal preparations. *Food Chem Toxicol* 2019;123:288-97.
- 72 Venhuis BJ, de Kaste D. Towards a decade of detecting new analogues of sildenafil, tadalafil and vardenafil in food supplements: a history, analytical aspects and health risks. *J Pharm Biomed Anal* 2012;69:196-208.



- 73 Rocha T, Amaral JS, Oliveira MBPP. Adulteration of dietary supplements by the illegal addition of synthetic drugs: a review. *Compr Rev Food Sci Food Saf* 2016;15:43–62.
- 74 Akour A, Kasabri V, Bulatova N, *et al.* Patterns and perceived efficacy of herbal medicine for weight loss and maintenance: a cross-sectional survey from Jordan. *Eur J Integr Med* 2020;35:101086.
- 75 Duraz AY, Khan SA, Knowledge KSA. Knowledge, attitudes and awareness of community pharmacists towards the use of herbal medicines in muscat region. *Oman Med J* 2011;26:451.
- 76 Oshikoya KA, Oreagba IA, Ogunleye OO, *et al.* Herbal medicines supplied by community pharmacies in Lagos, Nigeria: pharmacists' knowledge. *Pharm Pract* 2013;11:219.
- 77 Tahir M, Upadhyay DK, Iqbal MZ, *et al.* Knowledge of the use of herbal medicines among community pharmacists and reporting their adverse drug reactions. *J Pharm Bioallied Sci* 2020;12:436.
- 78 Shilbayeh SA. Exploring knowledge and attitudes towards counselling about vitamin supplements in Jordanian community pharmacies. *Pharm Pract* 2011;9:242.
- 79 Newall CA, Anderson LA, Phillipson JD. *Herbal medicines. A guide for health-care professionals.* The Pharmaceutical Press, 1996.
- 80 Shaw D. Toxicological risks of Chinese herbs. *Planta Med* 2010;76:2012–8.
- 81 Heinrich M, Chan J, Wanke S, *et al.* Local uses of Aristolochia species and content of nephrotoxic aristolochic acid 1 and 2--a global assessment based on bibliographic sources. *J Ethnopharmacol* 2009;125:108–44.
- 82 Michl J, Ingrouille MJ, Simmonds MSJ, *et al.* Naturally occurring aristolochic acid analogues and their toxicities. *Nat Prod Rep* 2014;31:676–93.
- 83 Booker A, Frommenwiler D, Reich E, *et al.* Adulteration and poor quality of Ginkgo biloba supplements. *J Herb Med* 2016;6:79–87.
- 84 Booker A, Agapouda A, Frommenwiler DA, *et al.* St John's wort (*Hypericum perforatum*) products - an assessment of their authenticity and quality. *Phytomedicine* 2018;40:158–64.
- 85 Booker A, Jalil B, Frommenwiler D, *et al.* The authenticity and quality of *Rhodiola rosea* products. *Phytomedicine* 2016;23:754–62.
- 86 Alsbou M, Abdeen G, Batarseh A, *et al.* Analysis of the National pharmacovigilance database in Jordan (2010-2014). *Biomed Pharmacol J* 2017;10:319–28.
- 87 Barnes J, Mills SY, Abbot NC, *et al.* Different standards for reporting ADRs to herbal remedies and conventional OTC medicines: face-to-face interviews with 515 users of herbal remedies. *Br J Clin Pharmacol* 1998;45:496–500.
- 88 Walji R, Boon H, Barnes J, *et al.* Reporting natural health product related adverse drug reactions: is it the pharmacist's responsibility? *Int J Pharm Pract* 2011;19:383–91.