JAMA Pediatrics | Original Investigation

Association of Food and Nonalcoholic Beverage Marketing With Children and Adolescents' Eating Behaviors and Health A Systematic Review and Meta-analysis

Emma Boyland, PhD; Lauren McGale, PhD; Michelle Maden, PhD; Juliet Hounsome, PhD; Angela Boland, PhD; Kathryn Angus; Andrew Jones, PhD

IMPORTANCE There is widespread interest in the effect of food marketing on children; however, the comprehensive global evidence reviews are now dated.

OBJECTIVE To quantify the association of food and nonalcoholic beverage marketing with behavioral and health outcomes in children and adolescents to inform updated World Health Organization guidelines.

DATA SOURCES Twenty-two databases were searched (including MEDLINE, CINAHL, Web of Science, Embase, and The Cochrane Library) with a publication date limit from January 2009 through March 2020.

STUDY SELECTION Preferred Reporting Items for Systematic Reviews and Meta-Analyses reporting guidelines were followed. Inclusion criteria were primary studies assessing the association of food marketing with specified outcomes in children and adolescents (aged 0-19 years). Exclusion criteria were qualitative studies or those on advertising of infant formula. Of 31 063 articles identified, 96 articles were eligible for inclusion in the systematic review, and 80 articles in the meta-analysis (19 372 participants).

DATA EXTRACTION AND SYNTHESIS Two reviewers independently extracted data. Random-effects models were used for meta-analyses; meta-regressions, sensitivity analyses, and *P* curve analyses were also performed. Where appropriate, pooling was conducted using combining *P* values and vote counting by direction of effect. Grading of Recommendations Assessment, Development, and Evaluation was used to judge certainty of evidence.

MAIN OUTCOMES AND MEASURES Critical outcomes were intake, choice, preference, and purchasing. Important outcomes were purchase requests, dental caries, body weight, and diet-related noncommunicable diseases.

RESULTS Participants totaled 19 372 from 80 included articles. Food marketing was associated with significant increases in intake (standardized mean difference [SMD], 0.25; 95% CI, 0.15-0.35; P < .001), choice (odds ratio, 1.77; 95% CI, 1.26-2.50; P < .001), and preference (SMD, 0.30; 95% CI, 0.12-0.49; P = .001). Substantial heterogeneity (all >76%) was unexplained by sensitivity or moderator analyses. The combination of P values for purchase requests was significant but no clear evidence was found for an association of marketing with purchasing. Data on dental health and body weight outcomes were scarce. The certainty of evidence was graded as very low to moderate for intake and choice, and very low for preference and purchasing.

CONCLUSIONS AND RELEVANCE In this systematic review and meta-analysis, food marketing was associated with increased intake, choice, preference, and purchase requests in children and adolescents. Implementation of policies to restrict children's exposure is expected to benefit child health.

JAMA Pediatr. 2022;176(7):e221037. doi:10.1001/jamapediatrics.2022.1037 Published online May 2, 2022.

Supplemental content

Author Affiliations: Department of Psychology, University of Liverpool, Liverpool, United Kingdom (Boyland, McGale, Jones); Department of Psychology, Edge Hill University, Ormskirk, United Kingdom (McGale); Liverpool Reviews and Implementation Group, University of Liverpool, Liverpool, United Kingdom (Maden, Hounsome, Boland); Institute for Social Marketing & Health, University of Stirling, Stirling, Scotland (Angus).

Corresponding Author:

Emma Boyland, PhD, Department of Psychology, University of Liverpool, Bedford Street South, Liverpool L69 7ZA, United Kingdom (eboyland@ liverpool.ac.uk). lobal trends show substantial increases in obesity among children in recent decades.¹ This has serious implications for morbidity and mortality given that childhood obesity tracks into adulthood² and excess weight is an important risk factor for noncommunicable disease (NCD).³ Changes in global systems are key drivers of rising obesity, specifically growth in the production of affordable, highly processed foods that are effectively marketed.⁴

Food and/or nonalcoholic beverage (hereafter referred to as *food*) marketing that largely promotes products high in fat, sugar, and/or salt (HFSS) is prevalent across television, digital media, outdoor spaces, and sport. Children and adolescents are particularly vulnerable to the effects of food marketing given their immature cognitive and emotional development, peer-group influence, and high exposure. In the pathway linking exposure to HFSS food marketing with behavioral and health effects is complex but associations meet the criteria for a causal relationship. HFSS food marketing also negatively affects numerous child rights, including the right to the enjoyment of the highest attainable standard of health, the right to adequate food, and the right to privacy.

Implementation of the World Health Organization (WHO) Set of Recommendations on the Marketing of Foods and Nonalcoholic Beverages to Children¹⁴ has been inconsistent. ¹³ The underpinning evidence review¹⁵ largely predated the internet as a major marketing platform¹⁶ and there is more than a decade of new research to consider. Although its conclusions are corroborated by more recent reviews and meta-analyses, ¹⁷⁻²¹ these are also limited to television advertising and dated digital marketing forms (eg, advergaming), including selective outcomes, such as intake, and lack assessment of evidential value or certainty. Therefore, WHO commissioned the current research to inform the development of updated recommendations to restrict food marketing to children.

Methods

We conducted a systematic review and a series of metaanalyses following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline. The WHO Nutrition Guidance Expert Advisory Group Subgroup on Policy Actions formulated the research question and identified the critical and important outcomes to be captured (eAppendix 1 in the Supplement). The terms marketing, exposure, and power were used as defined by WHO. The protocol was preregistered in the PROSPERO database in May 2019 (CRD42019137993).

Search Strategy and Selection Criteria

We considered primary studies (randomized clinical trial [RCT] or nonrandomized study [NRS]) for inclusion if they assessed the association of food marketing with specified outcomes in children (aged 0-19 years). Exclusion criteria comprised qualitative designs and studies assessing the effect of advertising for infant formula or of marketing strategies outside of WHO's definition. Critical outcomes comprised food intake, choice,

Key Points

Question What is the association between food marketing (compared with less or no food marketing) and eating behavior and health in children and adolescents across the extant literature?

Findings In this systematic review and meta-analysis of 96 studies (64 randomized clinical trials, 32 nonrandomized studies), food marketing was associated with significant increases in food intake, choice, preference, and purchase requests. There was no clear evidence of associations with purchasing, and little evidence on dental health or body weight outcomes.

Meaning Results support the implementation of policies to restrict children's exposure to food marketing.

preference, and purchasing (by, or on behalf of, children). Important outcomes were purchase requests (by children to a caregiver), dental caries and erosion, body weight, body mass index (BMI) and obesity, and diet-related NCDs (including validated surrogate indicators). Outcomes are defined in eAppendix 1 in the Supplement.

Searches were conducted in April 2019 and updated in March 2020 by an information specialist (M.M.). Data were analyzed in December 2020. Searches were limited to studies added to databases from January 1, 2009 (the previous global review included evidence to December 2008).¹⁵ We searched MEDLINE, CINAHL, Web of Science, Embase, ERIC, The Cochrane Library (CDSR, CENTRAL), Business Source Complete, EconLit, Emerald, JSTOR, HMIC, Advertising Education Forum, The Campbell Library, Database of Promoting Health Effectiveness Reviews (DoPHER), Healthevidence.org, TRIP, IRIS, Global Index Medicus, KOREAMED, Communication & Mass Media Complete, Academic Search Complete, and Index to Legal Periodicals & Books Full Text (H.W. Wilson). Targeted searches of Google and Google Scholar were undertaken. The search strategy is provided in the eAppendix 1 in the Supplement. All searches were peer reviewed (checked for accuracy by 3 researchers [E.B., L.M., K.A.] and a WHO librarian).

These searches were supplemented by (1) handsearching reference lists of retrieved systematic reviews and eligible studies, (2) contact with topic experts, (3) forward and backward citation searching of included studies, and (4) a WHO evidence call for data.²⁴ No language restrictions were applied.

Two reviewers (E.B., L.M., J.H., M.M.) independently screened studies against the inclusion criteria, assessing titles and abstracts to identify potentially relevant studies, then reviewing full texts. Titles and abstracts of articles not in English were screened using Google Translate, then researchers proficient in both languages translated the full texts for review. For multiple publications from the same cohort, we used data from the main contrast (food marketing vs no, less, or less powerful marketing) or the biggest sample. Disagreement was resolved through consensus and, if necessary, consulting a third reviewer. The search and screening processes were combined for this and a parallel review on the effectiveness of food marketing policies (Prospero identifier: CRD42019132506).

Quality Assessment

We used Risk of Bias 2 to assess bias in RCTs and the Newcastle-Ottawa Scale to assess quality of the NRS. Bias assessments were conducted by one reviewer and independently checked by a second (E.B., L.M., J.H, M.M.).

Statistical Analysis

Two reviewers (E.B., L.M.) independently extracted data using prepiloted forms. Study authors were contacted, if necessary, to provide data. Where data were only available in a figure, we used WebPlotDigitizer (version 4.3) for extraction.²⁵

For studies with multiple interventions, we extracted data from all relevant interventions and the control group or most relevant comparator intervention. For studies with interventions comprising different levels of the same marketing exposures, we selected the largest (eg, most advertisements) as the exposure arm to maximize identification of effects. Relevant outcome measures and effect estimates were extracted. Where more than 1 eligible effect measure was available, we extracted the most comprehensive measure (eg, overall intake rather than of a single item) or prioritized the unhealthy categories.

Cochrane recommendations were followed for the synthesis. ²⁶ Meta-analysis was used where studies were sufficiently homogenous. Where meta-analysis was not possible, we selected the most appropriate synthesis method available: combining *P* values using Fisher method or vote counting by direction of effect (eAppendix 1 in the Supplement).

For meta-analyses, random-effects restricted maximum likelihood estimator analyses were conducted using the metafor package in R (version 4.1.3; R Foundation for Statistical Computing). The I^2 (inconsistency) statistic was used to assess heterogeneity, with a value of I^2 less than 50% indicating substantial heterogeneity. We undertook leave-one-out, trim and fill analyses, graphical displays of heterogeneity (GOSH), and Egger regression test to examine bias. 29 We examined any influential cases with a difference in beta score more than $1.^{30}$

When appropriate, we conducted subgroup (moderation) analyses by study design (RCT vs NRS), marketing manipulation type (exposure vs power), and marketing channel (television vs digital vs packaging). Within RCTs, we examined whether risk of bias scores (low vs medium) moderated the association (not possible for the preference outcome owing to the small number of data points), and within NRSs we conducted meta-regressions to examine if scores on the NOS were associated with the effect. For diet and choice outcomes, we examined whether mean age of children in the sample or BMI z score of the sample was associated with effect size using meta-regression (not possible for the preference outcome owing to the small number of data points). To examine evidential value, we conducted P curve analyses using the dmetar function in \mathbb{R} .

We used Grading of Recommendations Assessment, Development, and Evaluation (GRADE)³² to judge the certainty of evidence as high, moderate, low, or very low (eAppendix 1 in the Supplement). Research team certainty assessments were revised where necessary following discussion with the WHO Nutrition Guidance Expert Advisory Group Subgroup on Policy Actions.

Results

A total of 31 063 titles were assessed for eligibility and 28 682 were ineligible (**Figure 1**). Of 2381 full-text articles assessed, 96 studies were included in the systematic review and 80 in the meta-analyses. Study characteristics are provided in the Supplement (eAppendix 2 in the Supplement). Pooled critical outcome data for food intake, choice, and preference are summarized in the **Table**. Overall forest plots are shown in **Figure 2**, **Figure 3**, and **Figure 4**. Forest plots for subgroup analyses, GOSH, and *P* curve plots are in eAppendix 4 in the Supplement.

Data relating to other outcomes, bias assessments, and all GRADE tables are in the Supplement (eAppendices 3, 5, and 6 in the Supplement). No relevant studies were identified with the diet-related NCDs outcome.

For food intake, 46 studies (in 43 articles) were identified (31 RCTs, 33-61 8 observational NRSs, 62-69 and 7 experimental NRSs⁷⁰⁻⁷⁵). Pooled analyses of data from 41 studies (42 effect sizes) found that food marketing was associated with a significant increase in intake (standardized mean difference [SMD], 0.25; 95% CI, 0.15-0.35; z = 4.77; $I^2 = 77.2\%$; P < .001; Figure 2). The association was robust to sensitivity analyses and GOSH analyses demonstrated that across 100 000 iterations of the analyses the pooled effect SMD was approximately 0.24 (eAppendix 4 in the Supplement). There was no statistical evidence that study design ($\chi^2 = 1.75$; P = .19), marketing manipulation type (χ^2 = 0.39; P = .53), or marketing channel ($\chi^2 = 0.71$; P = .70) significantly moderated the effect sizes. A meta-regression of mean age of children in the studies (mean [range] age, 8.6 years [4.1-13.6]) on the effect size was not significant ($\beta = -0.02$; 95% CI, -0.071 to 0.252; P = .35). There was no association between BMI z scores (mean [range], 1.01 [0.01-2.30]) and the effect size (β = 0.20; 95% CI, -0.136 to 0.534; P = .24). The P curve continuous test for evidential value was significant (z = 8.226; P < .001), indicating a true effect, as the distributions of *P* values were more frequent at *P* less than .01 compared with *P* of approximately .05. Of the 5 studies not included in the pooled analyses, 3 found associations of food marketing on intake 53,62,68 and 2 found no association. 64,75 The certainty of evidence for RCTs was moderate (affected by unexplained high heterogeneity), and for NRSs was very low (observational studies have a lower starting position within the GRADE assessment and certainty was downgraded owing to the imprecision of the effect size estimates from these studies).

For food choice, 37 studies (in 36 articles) were identified (27 RCTs^{59,76-96} and 10 experimental NRSs^{72,97-100}). Pooled analyses of data from 27 studies found that food marketing was significantly associated with food choice (odds ratio [OR], 1.77; 95% CI, 1.26-2.50; $z=3.27; I^2=77.5\%; P<.001$; Figure 3). Specifically, food marketing exposure was associated with increased odds of 1.77 times greater choice of the test item(s), irrespective of whether the test item was unhealthy or healthy. However, we note that only 3 of 27 effect sizes^{85,87,99} reported on choice of healthier items specifically and only 1 of those did so within a study design in which the marketing ex-




Table. Pooled Associations of Food Marketing Compared With No Marketing, Less Food Marketing, or Less Powerful Food Marketing With Critical Outcomes

Study type	No. of	Participants, No./total No. (%)			GRADE certainty				
	studies	Experimental arm Control arm		Effect size (95% CI)	of evidence				
Nonrandomized studies			Firmental arm Control arm Effect size (95% CI) of evidence (5/8436 (50.3) 4191/8436 (49.7) SMD, 0.34 (0.12-0.57) ^a Very low (416 (62.7) 155/416 (37.3) OR, 0.56 0.05-5.99) ^a Very low (5/2908 (50.1) 1452/2908 (49.9) SMD, 0.20 (0.10-0.30) ^a Moderate						
Intake	11	4245/8436 (50.3)	4191/8436 (49.7)	SMD, 0.34 (0.12-0.57) ^a	Very low				
Choice	5	261/416 (62.7)	155/416 (37.3)	OR, 0.56 0.05-5.99) ^a	Very low				
Preference	4	1010/1972 (51.2)	962/1972 (48.8)	SMD, 0.21 (0.07-0.35)b	Very low				
Randomized trials									
Intake	30	1456/2908 (50.1)	1452/2908 (49.9)	SMD, 0.20 (0.10-0.30) ^a	Moderate				
Choice	22	1916/3838 (49.9)	1922/3838 (50.1)	OR, 1.97 1.46-2.66) ^a	Moderate				
Preference	8	894/1802 (49.6)	908/1802 (50.4)	SMD, 0.38 (0.03-0.72) ^{b,c}	Very low				

Abbreviations: GRADE, Grading of Recommendations Assessment, Development, and Evaluation; OR, odds ratio; SMD, standardized mean difference.

posure itself was for healthier food. ⁸⁵ The association was robust to sensitivity analyses and GOSH analyses demonstrated that across 100 000 iterations of the analyses the pooled effect OR was approximately 1.70 (eAppendix 4 in the Supplement). There was no statistical evidence that study design ($\chi^2 = 3.01; P = .08$), marketing manipulation type ($\chi^2 = 0.012; P = .91$), or marketing channel ($\chi^2 = 0.02, P = .99$) significantly moderated the effect sizes. A meta-regression of mean age of children in the studies (mean [range] age, 8.76 [4.0-11.8] years) on the effect size was not significant ($\beta = -0.08; 95\%$ CI, -0.345 to 0.178; P = .53). The continuous test for evidential value was significant (z = 8.287; P < .001). Ten stud-

ies were not included in the pooled analysis; of these 8 found an association of food marketing with food choice (of which 7 were in the direction of greater choice of test items with food marketing exposure^{74,101-106} while 1 found greater choice of test items in the control condition¹⁰⁷) and 2 found no association.^{104,108} Supplementary analysis of 3 of these studies^{103,104} that used a crossover design with binary outcomes showed a nonsignificant pooled OR of 3.45 (95% CI, 0.97-12.43). The certainty of evidence for RCTs was moderate (unexplained high heterogeneity), and for NRSs was very low (observational studies, risk of bias, and imprecision of the effect size estimates).

^a The high heterogeneity of the pooled effect size (≥50%) is unexplained by sensitivity analyses (although the association did not change direction or significance).

^b High heterogeneity (although the association did not change direction or significance).

^c Sensitivity analyses demonstrated that there was variability in the association when individual studies were removed but these analyses do not provide a public health-relevant explanation for the heterogeneity so downgrading for heterogeneity is still appropriate within the GRADE assessment.

Figure 2. Forest Plot of Intake Data From Eligible Studies

Study	Standardized mean difference (95% CI)
Aerts and Smith, ⁷⁰ 2019; study 1	0.21 (-0.05 to 0.47)
Aerts and Smith, ⁷⁰ 2019; study 2	0.40 (0.02 to 0.77)
Anderson et al, ³³ 2015; overweight	0.24 (-0.25 to 0.74)
Anderson et al, 33 2015; normal weight	0.04 (-0.50 to 0.58)
Anschutz et al, ³⁵ 2010	-0.14 (-0.58 to 0.30)
Anschutz et al, 34 2009	0.04 (-0.32 to 0.40)
Boyland et al, ⁷¹ 2013	0.64 (0.21 to 1.07)
Brown et al, ⁷² 2017	-0.31 (-0.68 to 0.06)
Coates et al, ³⁶ 2019	0.05 (-0.34 to 0.44)
Coates et al, ³⁷ 2019	0.63 (0.26 to 1.01)
Dovey et al, ⁷³ 2011	1.18 (0.90 to 1.47)
Emond et al, ⁶³ 2019	0.57 (0.36 to 0.78)
Emond et al, ³⁸ 2016	0.53 (0.02 to 1.05)
Folkvord et al, ³⁹ 2013	0.69 (0.34 to 1.04)
Folkvord et al, ⁴⁰ 2017; Dutch sample	0.69 (0.30 to 1.09)
Folkvord et al, 40 2017; Spanish sample	-0.01 (-0.31 to 0.29)
Folkvord et al, ⁴¹ 2016	-0.04 (-0.52 to 0.45)
Folkvord et al, ⁴² 2014	0.50 (0.15 to 0.85)
Folkvord et al, ⁴³ 2015	0.49 (0.07 to 0.90)
Forman et al, ⁴⁴ 2009	0.01 (-0.41 to 0.44)
Gilbert-Diamond et al, ⁴⁵ 2016	0.07 (-0.23 to 0.37)
Gregori et al, ⁴⁸ 2017	
	-0.32 (-0.81 to 0.17)
regori et al, ⁴⁷ 2014	-0.03 (-0.24 to 0.19)
regori et al, ⁴⁶ 2013	0.03 (-0.33 to 0.39)
larris et al, ⁵⁰ 2018	-0.18 (-0.59 to 0.23)
arris et al, ⁵¹ 2012	0.14 (-0.25 to 0.53)
larris et al, ⁴⁹ 2009	0.29 (-0.07 to 0.65)
Keller et al, ⁵² 2012; boys	0.05 (-0.34 to 0.44)
Keller et al, ⁵² 2012; girls	0.50 (0.08 to 0.93)
Kelly et al, ⁶⁵ 2015	0.50 (0.23 to 0.76)
Leonard et al, ⁷⁴ 2019	-0.06 (-0.57 to 0.46)
Lorenzoni et al, ⁵⁴ 2017	0.98 (-0.06 to 2.01)
Lorenzoni et al, ⁵⁵ 2017	-0.21 (-1.02 to 0.59)
Masterson et al, ⁵⁶ 2019	-0.02 (-0.29 to 0.26)
McGale et al, ⁵⁷ 2020	0.77 (0.13 to 1.40)
Minaker et al, ⁶⁶ 2011	0.16 (0.09 to 0.23)
Norman et al, ⁵⁸ 2018	0.27 (0.06 to 0.47)
Olafsdottir et al, ⁶⁷ 2014	0.34 (0.22 to 0.46)
Silva et al, ⁶⁹ 2020	0.07 (-0.17 to 0.31)
Smith et al, ⁵⁹ 2020	-0.04 (-0.48 to 0.40)
Smits and Vandebosch, 60 2012	0.53 (0.00 to 1.06)
Vecchio et al, ⁶¹ 2019	0.49 (-1.13 to 2.12)
Pooled effect	0.25 (0.15 to 0.35)
. 55155 5.756	0.23 (0.13 to 0.33)

For food preference, 20 studies (in 19 articles) were identified (12 RCTs^{53,77,79,94,104,109-114} and 8 experimental NRSs^{97,103,106,115-119}). Pooled analyses of data from 12 studies found that food marketing was significantly associated with increased food preference (SMD, 0.30; 95% CI, 0.12-0.49; Z=3.21, $I^2=90.0\%$; P=.001; Figure 4). The association was robust to sensitivity analyses and GOSH analyses demonstrated that across 100 000 iterations of the analyses the pooled effect SMD was approximately 0.53 (eAppendix 4 in the Supplement). There was no statistical evidence that study design (χ^2 [1] = 0.19; P=.67), marketing manipulation type ($\chi^2=0.44$; P=.51), or marketing channel ($\chi^2=1.29$; P=.53) significantly moderated the effect sizes. The continuous test for

evidential value was significant (z=5.504; P<.01). Eight studies were not able to be included in the pooled analysis, of which 6 found an association of food marketing with preference^{2,53,103,104,106,112} and 2 found no association. When studies with crossover designs and binary outcomes^{103,104,106,112} were analyzed separately, there was a significant association of marketing with preference (OR, 3.49; 95% CI, 2.03-6.22; z=4.40; P<.001). The certainty of evidence for both RCTs and NRSs was very low (inconsistency, imprecision).

For food purchasing, 5 studies (1 RCT, ¹²⁰ 1 experimental NRSs, ¹²¹ and 3 observational NRSs^{66,122,123}) were identified. All 3 observational NRSs (moderate quality to high quality) found

Figure 3. Forest Plot of Choice Data From Eligible Studies

Agante and Pascoal, ⁹⁷ 2019 Arendt et al, ⁷⁶ 2015 Boyland et al, ⁷⁷ 2015 Brown et al, ⁷² 2017 Bruce et al, ⁷⁸ 2016 Dois and Anante, ⁷⁹ 2011 Dixon et al, ⁸¹ 2014 Dixon et al, ⁸⁰ 2017	1.53 (0.48 to 2.58) 0.81 (0.04 to 1.58) 0.12 (-1.10 to 1.35) 1.12 (-1.80 to 4.04) 0.21 (-1.34 to 1.75) 1.36 (0.81 to 1.90) 0.11 (-0.46 to 0.68)			_		-	
Boyland et al, ⁷⁷ 2015 Brown et al, ⁷² 2017 Bruce et al, ⁷⁸ 2016 Dois and Anante, ⁷⁹ 2011 Dixon et al, ⁸¹ 2014	0.12 (-1.10 to 1.35) 1.12 (-1.80 to 4.04) 0.21 (-1.34 to 1.75) 1.36 (0.81 to 1.90)			-	_		
Brown et al, ⁷² 2017 Bruce et al, ⁷⁸ 2016 Dois and Anante, ⁷⁹ 2011 Dixon et al, ⁸¹ 2014	1.12 (-1.80 to 4.04) 0.21 (-1.34 to 1.75) 1.36 (0.81 to 1.90)						
Bruce et al, ⁷⁸ 2016 Dois and Anante, ⁷⁹ 2011 Dixon et al, ⁸¹ 2014	0.21 (-1.34 to 1.75) 1.36 (0.81 to 1.90)			-	_		
Dois and Anante, ⁷⁹ 2011 Dixon et al, ⁸¹ 2014	1.36 (0.81 to 1.90)		_				-
Dixon et al, ⁸¹ 2014			-				
,	0.11 / 0.46 +0.0 (0)						
Dixon et al, ⁸⁰ 2017	U.11 (-U.46 to U.68)			-			
	0.11 (-0.39 to 0.60)			-			
Ferguson et al, ⁸³ 2012	2.26 (0.57 to 3.95)			-	-		
Ferguson et al, ⁸² 2014	0.76 (-1.10 to 2.61)			-		-	
Gatou et al, ⁸⁴ 2016	-0.13 (-1.05 to 0.79)			-			
Hobin et al, ⁸⁵ 2012	1.01 (0.52 to 1.50)			-	-		
Hudson and Elliott, ⁹⁸ 2013	0.45 (-0.28 to 1.18)			-	_		
Matthes and Naderer, ⁸⁶ 2015	1.28 (0.24 to 2.32)			-	-		
McDarby et al, ⁸⁷ 2018	0.91 (0.13 to 1.69)			-	_		
Naderer et al, ⁸⁹ 2018	0.27 (-0.75 to 1.29)				_		
Naderer et al, ⁸⁸ 2018	0.80 (0.04 to 1.56)			_			
Naderer et al, ⁹⁰ 2018	2.14 (1.44 to 2.83)				-	_	
Pempek and Calvert, 99 2009	-5.29 (-7.56 to -3.02)	 					
Putnam et al, ⁹¹ 2018	0.70 (-0.24 to 1.64)			-			
Royne et al, ⁹² 2017	0.61 (-0.94 to 2.15)			-			
Smith et al, ⁵⁹ 2020	2.00 (0.98 to 3.01)				-	_	
Tarabashkina et al, ⁹³ 2016	0.18 (-0.25 to 0.60)			-			
Jeda et al, ¹⁰⁰ 2014	-1.07 (-2.55 to 0.41)						
Ulger et al, ⁹⁵ 2008	-0.50 (-1.37 to 0.36)		-	-			
Uribe and Fuentes-García, ⁹⁶ 2015	0.67 (0.16 to 1.18)			-	_		
Toomey and Francis, ⁹⁴ 2013	0.04 (-0.88 to 0.96)				-		
Pooled effect	0.57 (0.23 to 0.92)			•			

an association between food marketing and purchasing (2 effects of public health harm, ^{66,122} 1 of public health benefit¹²³). The RCT (with some concerns of bias)¹²⁰ and moderate-quality experimental NRS¹²¹ found no association. The proportion of studies that found clear association of potential public health harm (1 of 4) was 25% (95% CI, 1.3%-78.1%). The proportion of studies that found unclear associations of potential public health harm (1 of 4) was 25% (95% CI, 1.3%-78.1%). The proportion of studies that showed any association (clear or unclear) of public health harm (2 of 5) was 40% (95% CI, 7.3%-83.0%). The certainty of evidence for both RCTs and NRSs was very low (risk of bias, inconsistency, imprecision).

For purchase requests, 6 studies (5 RCTs 60,81,110,111,114 and 1 observational NRS 68) were identified. The combination of P values was statistically significant in both model iterations (eAppendix 1 in the Supplement) suggesting evidence of food marketing associations with this outcome. The certainty of evidence for RCTs was moderate (risk of bias), and for NRS was very low (observational studies, risk of bias).

For dental caries, 2 observational NRSs were identified. A moderate-quality study found a clear association of public health harm 124 and a high-quality study found no association. 69 The proportion of studies that showed any association (clear or unclear) with public health harm (1 of 2) was 50% (95% CI,

9%-90.5%). The certainty of evidence was very low (risk of bias, inconsistency, indirectness).


Very little evidence was available on the association between food marketing and body weight or BMI. This review identified a single, moderate-quality observational NRS with no significant associations. The certainty of evidence was very low (risk of bias, indirectness). No studies were found with relevant data on diet-related NCDs or validated surrogate indicators.

Discussion

In this study, food marketing exposure was associated with increases in children's food intake, choice of and preference toward test items, and purchase requests. There was little evidence to support associations with food purchasing by or on behalf of children, while data relating to dental health and body weight outcomes were scarce. No studies were found for the diet-related NCDs or validated surrogate indicators outcome.

The effect sizes from the pooled analyses were small for intake and preference, moderate to large for choice, and robust to sensitivity analyses. *P* curve analyses demonstrated significant evidential value, indicative of a lack of selective re-

Figure 4. Forest Plot of Preference Data From Eligible Studies



porting or P hacking. These findings are largely consistent with, and build on, previous findings, 15,17,18,20,21 although there are some discrepancies. For example, Russell et al 20 identified a moderating effect of BMI, such that children with overweight or obesity consumed an average of 45.6 kilocalories more than children with healthy weight following exposure to food advertisements. That type of subgroup analysis was not possible here owing to a lack of appropriate data reported in the studies (of the 5 effect sizes included in each group, 20 2 took place pre-2009, so were excluded here).

Strengths and Limitations

A strength of the present review is that it has linked diverse formats of food marketing exposure (including newer digital forms, such as social media influencer marketing) to a range of behavioral and health outcomes. Other analyses have reported on a single format of marketing exposure (eg, screenbased) and fewer than 3 outcomes. The certainty of evidence for critical outcomes was most frequently rated as very low or moderate, which could be regarded as a limitation. However, as has been described previously, 125 this reflects the nature of the GRADE criteria. GRADE prioritizes RCT data with clinical outcomes and requires certainty to be downgraded where there is unexplained heterogeneity, even where results are consistent between RCTs and NRSs and show similar findings to previous reviews, as here. The substantial observed heterogeneity, also consistent with previous meta-analyses, 17,18,20 was unexplained by sensitivity analyses, or subgroup analyses on overall study design (although for intake and choice outcomes, variability was reduced when only RCTs were included), marketing manipulation, marketing format, study quality, participant age, or BMI. Therefore, this heterogeneity is likely a consequence of the large number of studies and more nuanced differences in study design (eg, stimulus types and outcome measurement). Substantial variability in outcome measurement is acceptable in meta-analysis but has implications for heterogeneity and therefore GRADE assessments.126

This study has limitations. As with the previous WHO review, ¹⁵ much of the evidence lies at the proximal end of the spectrum (relative to a hierarchy of food marketing effects ¹¹) with data available on food intake, choice, and preference outcomes, but far less for the more distal outcomes (body weight and NCDs). Intake studies tend to measure immediate or short-term intake (directly following exposure to the marketing stimulus), rather than assessing diet across the day or longer term. Research gaps at the distal end likely reflect the substantial methodological challenge of conducting such studies, given that weight gain (or development of diet-related NCDs) typically occurs gradually and there is limited variability in the marketing exposure children experience within any given country or culture. ¹¹

The evidence is almost exclusively from higher-income countries, with only 6 studies conducted in lower-income to middle-income countries. ^{64,69,95,100,103,124} The representativeness of the data for those populations may be limited and there was no opportunity to examine potential differences by income. Although we could explore the association of BMI and age with some outcomes through metaregression, we could not conduct formal subgroup analyses by age (eg, child vs adolescent), socioeconomic status, gender, or rural/urban residential status owing to inadequate reporting (ie, insufficient studies with data segregated by these characteristics) and a lack of studies of adolescents. Future research should address this.

Conclusions

This review provides a comprehensive update and quantitative synthesis of evidence of food marketing associations with critical behavioral outcomes and demonstrates the evidential value of these studies. WHO has previously recommended that member states enact policies to restrict children's exposure to unhealthy food marketing¹⁴ and the review findings support this position.

ARTICLE INFORMATION

Accepted for Publication: January 19, 2022.

Published Online: May 2, 2022. doi:10.1001/jamapediatrics.2022.1037

Open Access: This is an open access article distributed under the terms of the CC-BY License. © 2022 Boyland E et al. *JAMA Pediatrics*.

Author Contributions: Drs Boyland and Jones had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Boyland, Maden, Hounsome, Boland, Angus.

Acquisition, analysis, or interpretation of data: Boyland, McGale, Maden, Hounsome, Angus, Jones.

Drafting of the manuscript: Boyland, McGale, Maden, Hounsome, Jones.

Critical revision of the manuscript for important intellectual content: Boyland, McGale, Maden, Hounsome, Boland, Angus.

Statistical analysis: Jones.

Obtained funding: Boyland.

Administrative, technical, or material support: Boyland, McGale, Maden, Hounsome, Boland, Jones.

Supervision: Boyland, Hounsome.

Conflict of Interest Disclosures: None reported.

Funding/Support: This article was supported by funding from the World Health Organization.

Role of the Funder/Sponsor: The World Health Organization Nutrition Guidance Expert Advisory Group Subgroup on Policy Actions specified the PICO criteria (including exposure and outcome measures) and confirmed or modified the certainty judgments but otherwise had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We thank Niamh Maloney, PhD, and Katherine Edwards (Liverpool, United Kingdom) for contributions to the screening and data extraction processes for this review. Niamh Maloney was briefly employed as a researcher on the World Health Organization grant that funded this review. Katherine Edwards was not financially compensated.

REFERENCES

- 1. Abarca-Gómez L, Abdeen ZA, Hamid ZA, et al; NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128-9 million children, adolescents, and adults. *Lancet*. 2017;390(10113): 2627-2642. doi:10.1016/S0140-6736(17)32129-3
- 2. Evensen E, Wilsgaard T, Furberg A-S, Skeie G. Tracking of overweight and obesity from early childhood to adolescence in a population-based cohort—the Tromsø Study, Fit Futures. *BMC Pediatr*. 2016;16(1):64. doi:10.1186/s12887-016-0599-5
- **3**. GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*.

- 2020;396(10258):1223-1249. doi:10.1016/S0140-6736(20)30752-2
- 4. Swinburn BA, Sacks G, Hall KD, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet*. 2011;378(9793): 804-814. doi:10.1016/S0140-6736(11)60813-1
- 5. Kelly B, Vandevijvere S, Ng S, et al. Global benchmarking of children's exposure to television advertising of unhealthy foods and beverages across 22 countries. *Obes Rev.* 2019;20(suppl 2): 116-128. doi:10.1111/obr.12840
- **6**. Potvin Kent M, Pauzé E, Roy EA, de Billy N, Czoli C. Children and adolescents' exposure to food and beverage marketing in social media apps. *Pediatr Obes.* 2019;14(6):e12508. doi:10.1111/jjpo. 12508
- 7. Pasch KE, Poulos NS. Outdoor food and beverage advertising: a saturated environment. In: Williams JD, Pasch KE, Collins CA, eds. Advances in Communication Research to Reduce Childhood Obesity. 2013:303-315. doi:10.1007/978-1-4614-5511-0_14
- **8**. Dixon H, Lee A, Scully M. Sports sponsorship as a cause of obesity. *Curr Obes Rep.* 2019;8(4):480-494. doi:10.1007/s13679-019-00363-z
- 9. Rozendaal E, Buijzen M, Valkenburg P. Comparing children's and adults' cognitive advertising competences in the Netherlands. *J Child Media*. 2010;4(1):77-89. doi:10.1080/17482790903407333
- **10**. Truman E, Elliott C. Identifying food marketing to teenagers: a scoping review. *Int J Behav Nutr Phys Act*. 2019;16(1):67. doi:10.1186/s12966-019-0833-2
- 11. Kelly B, King MPsy L, Chapman Mnd K, Boyland E, Bauman AE, Baur LA. A hierarchy of unhealthy food promotion effects: identifying methodological approaches and knowledge gaps. *Am J Public Health*. 2015;105(4):e86-e95. doi:10. 2105/AJPH.2014.302476
- 12. Norman J, Kelly B, Boyland E, McMahon A-T. The impact of marketing and advertising on food behaviours: evaluating the evidence for a causal relationship. *Curr Nutr Rep.* 2016;5(3):139-149. doi:10.1007/s13668-016-0166-6
- 13. WHO Regional Office for Europe. Evaluating implementation of the WHO set of recommendations on the marketing of foods and non-alcoholic beverages to children: progress, challenges and guidance for next steps in the WHO European Region. 2018. Accessed March 31, 2022. http://www.euro.who.int/_data/assets/pdf_file/0003/384015/food-marketing-kids-eng.pdf
- 14. World Health Organization. Set of recommendations for the marketing of food and non-alcoholic beverages to children. 2010. Accessed March 31, 2022. http://whqlibdoc.who.int/publications/2010/9789241500210_eng.pdf
- **15.** Cairns G, Angus K, Hastings G. The extent, nature and effects of food promotion to children: a review of the evidence to December 2008; prepared for the World Health Organization. 2009. Accessed March 31, 2022. https://www.who.int/dietphysicalactivity/Evidence_Update_2009.pdf
- **16.** World Health Organization. Tackling food marketing to children in a digital world: trans-disciplinary perspectives. children's rights,

- evidence of impact, methodological challenges, regulatory options and policy implications for the WHO European Region. 2016. March 31, 2022. https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/publications/2016/tackling-food-marketing-to-children-in-a-digital-world-trans-disciplinary-perspectives.-childrens-rights,-evidence-of-impact,-methodological-challenges,-regulatory-options-and-policy-implications-for-the-who-european-region-2016
- 17. Boyland EJ, Nolan S, Kelly B, et al. Advertising as a cue to consume: a systematic review and meta-analysis of the effects of acute exposure to unhealthy food and nonalcholic beverage advertising on intake in children and adults. *Am J Clin Nutr.* 2016;103(2):519-533. doi:10.3945/ajcn. 115.120022
- 18. Sadeghirad B, Duhaney T, Motaghipisheh S, Campbell NRC, Johnston BC. Influence of unhealthy food and beverage marketing on children's dietary intake and preference: a systematic review and meta-analysis of randomized trials. *Obes Rev.* 2016; 17(10):945-959. doi:10.1111/obr.12445
- **19**. Folkvord F, van 't Riet J. The persuasive effect of advergames promoting unhealthy foods among children: a meta-analysis. *Appetite*. 2018;129:245-251. doi:10.1016/j.appet.2018.07.020
- **20**. Russell SJ, Croker H, Viner RM. The effect of screen advertising on children's dietary intake: a systematic review and meta-analysis. *Obes Rev.* 2019;20(4):554-568. doi:10.1111/obr.12812
- 21. Smith R, Kelly B, Yeatman H, Boyland E. Food marketing influences children's attitudes, preferences and consumption: a systematic critical review. *Nutrients*. 2019;11(4):875. doi:10.3390/nu11040875
- 22. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097
- **23.** World Health Organization. *A Framework for Implementing the Set of Recommendations on the Marketing of Foods and Non-alcoholic Beverages to Children*. World Health Organization; 2012.
- 24. World Health Organization. Call for data on the impact of marketing for foods and non-alcoholic beverages on children's eating behaviours, body weight and health. 2019. https://www.who.int/nutrition/topics/datacall-marketingtochildren-impact-2019/en/
- **25.** Rohatgi A. WebPlotDigitizer, version 3.9. 2015. Accessed March 31, 2022. https://silo.tips/download/webplotdigitizer-user-manual-version-39
- **26**. McKenzie JE, Brennan SE. Chapter 12: Synthesizing and presenting findings using other methods. In: Higgins JPT, Thomas J, Chandler J, et al. *Cochrane Handbook for Systematic Reviews of Interventions, version 62*. Cochhane. 2021
- **27**. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw*. 2010;36(3): 1-48. doi:10.18637/jss.v036.i03
- **28**. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*. 2000;56(2):455-463. doi:10.1111/j.0006-341X.2000. 00455.x

- **29**. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-634. doi:10.1136/bmj.315.7109.629
- **30**. Viechtbauer W, Cheung MW-L. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods*. 2010;1(2):112-125. doi:10.1002/jrsm.11
- **31**. Harrer M, Cuijpers P, Furukawa T, Ebert DD. Companion R package for the guide 'doing meta-analysis in R'. 2019. Accessed March 31, 2022. https://dmetar.protectlab.org
- **32.** Guyatt GH, Oxman AD, Vist GE, et al; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924-926. doi:10.1136/bmj.39489.470347.AD
- **33.** Anderson GH, Khodabandeh S, Patel B, Luhovyy BL, Bellissimo N, Mollard RC. Mealtime exposure to food advertisements while watching television increases food intake in overweight and obese girls but has a paradoxical effect in boys. *Appl Physiol Nutr Metab*. 2015;40(2):162-167. doi:10. 1139/apmr-2014-0249
- **34.** Anschutz DJ, Engels RCME, Van Strien T. Side effects of television food commercials on concurrent nonadvertised sweet snack food intakes in young children. *Am J Clin Nutr*. 2009;89(5): 1328-1333. doi:10.3945/ajcn.2008.27075
- **35.** Anschutz DJ, Engels RCME, Van Strien T. Maternal encouragement to be thin moderates the effect of commercials on children's snack food intake. *Appetite*. 2010;55(1):117-123. doi:10.1016/j. appet.2010.03.014
- **36.** Coates AE, Hardman CA, Halford JCG, Christiansen P, Boyland EJ. Social media influencer marketing and children's food intake: a randomized trial. *Pediatrics*. 2019;143(4):e20182554. doi:10. 1542/peds.2018-2554
- **37**. Coates AE, Hardman CA, Halford JCG, Christiansen P, Boyland EJ. The effect of influencer marketing of food and a "protective" advertising disclosure on children's food intake. *Pediatr Obes*. 2019;14(10):e12540. doi:10.1111/ijpo.12540
- **38**. Emond JA, Lansigan RK, Ramanujam A, Gilbert-Diamond D. Randomized exposure to food advertisements and eating in the absence of hunger among preschoolers. *Pediatrics*. 2016;138(6): e20162361. doi:10.1542/peds.2016-2361
- **39**. Folkvord F, Anschütz DJ, Buijzen M, Valkenburg PM. The effect of playing advergames that promote energy-dense snacks or fruit on actual food intake among children. *Am J Clin Nutr*. 2013;97 (2):239-245. doi:10.3945/ajcn.112.047126
- **40**. Folkvord F, Lupiáñez-Villanueva F, Codagnone C, Bogliacino F, Veltri G, Gaskell G. Does a 'protective' message reduce the impact of an advergame promoting unhealthy foods to children? an experimental study in Spain and the Netherlands. *Appetite*. 2017;112:117-123. doi:10. 1016/j.appet.2017.01.026
- **41**. Folkvord F, Veling H, Hoeken H. Targeting implicit approach reactions to snack food in children: effects on intake. *Health Psychol*. 2016;35 (8):919-922. doi:10.1037/hea0000365
- **42**. Folkvord F, Anschütz DJ, Nederkoorn C, Westerik H, Buijzen M. Impulsivity, "advergames," and food intake. *Pediatrics*. 2014;133(6):1007-1012. doi:10.1542/peds.2013-3384

- **43**. Folkvord F, Anschütz DJ, Wiers RW, Buijzen M. The role of attentional bias in the effect of food advertising on actual food intake among children. *Appetite*. 2015;84:251-258. doi:10.1016/j.appet.2014. 10.016
- **44.** Forman J, Halford JCG, Summe H, MacDougall M, Keller KL. Food branding influences ad libitum intake differently in children depending on weight status. results of a pilot study. *Appetite*. 2009;53(1):76-83. doi:10.1016/j.appet.2009.05.015
- **45**. Gilbert-Diamond D, Emond J, Lansigan RK, et al. Television food advertisement exposure and FTO genotype in relation to excess consumption in children. *Int J Obes (Lond)*. 2016;41(1):23-39. doi:10.1038/ijo.2016.163
- **46**. Gregori D, Ballali S, Gafare CE, et al. Investigating the obesogenic effects of marketing snacks with toys: an experimental study in Latin America. *Nutr J.* 2013;12(1):95. doi:10.1186/1475-2891-12-95
- **47**. Gregori D, Gulati A, Hochdorn A, et al. Food packaged with toys: an investigation on potential obesogenic effects in Indian children. *Indian J Pediatr*. 2014;81(1)(suppl 1):30-38. doi:10. 1007/s12098-014-1448-x
- **48**. Gregori D, Lorenzoni G, Ballali S, Vecchio MG, Verduci E, Berchialla P. Is brand visibility on snacks packages affecting their consumption in children? results from an experimental ad-libitum study. *Arch Latinoam Nutr.* 2017;67:36-49.
- **49**. Harris JL, Bargh JA, Brownell KD. Priming effects of television food advertising on eating behavior. *Health Psychol*. 2009;28(4):404-413. doi:10.1037/a0014399
- **50**. Harris JL, Haraghey KS, Lodolce M, Semenza NL. Teaching children about good health? halo effects in child-directed advertisements for unhealthy food. *Pediatr Obes*. 2018;13(4):256-264. doi:10.1111/ijpo.12257
- **51.** Harris JL, Speers SE, Schwartz MB, Brownell KD. US food co branded advergames on the internet: children's exposure and effects on snack consumption. *J Child Media*. 2012;6(1):51-68. doi:10.1080/17482798.2011.633405
- **52**. Keller KL, Kuilema LG, Lee N, et al. The impact of food branding on children's eating behavior and obesity. *Physiol Behav*. 2012;106(3):379-386. doi:10.1016/j.physbeh.2012.03.011
- **53**. Kotler JA, Schiffman JM, Hanson KG. The influence of media characters on children's food choices. *J Health Commun*. 2012;17(8):886-898. doi:10.1080/10810730.2011.650822
- **54.** Lorenzoni G, Rtskhladze I, Vecchio MG, et al. Effect of TV advertising on energy intake of Georgian children: results of an experimental study. *Med J Nutrition Metab*. 2017;10(3):183-192. doi:10. 3233/MNM-17153
- **55.** Lorenzoni G, Zec S, Farias LF, et al. Does food advertising influence snacks consumption in Chilean children? results from an experimental ad libitum study. *Arch Latinoam Nutr.* 2017;67:24-35.
- **56.** Masterson TD, Bermudez MA, Austen M, et al. Food commercials do not affect energy intake in a laboratory meal but do alter brain responses to visual food cues in children. *Appetite*. 2019;132: 154-165. doi:10.1016/j.appet.2018.10.010
- **57**. McGale LS, Smits T, Halford JCG, Harrold JA, Boyland EJ. The influence of front-of-pack portion size images on children's serving and intake of

- cereal. *Pediatr Obes*. 2020;15(2):e12583. doi:10.1111/ijpo.12583
- **58.** Norman J, Kelly B, McMahon AT, et al. Sustained impact of energy-dense TV and online food advertising on children's dietary intake: a within-subject, randomised, crossover, counter-balanced trial. *Int J Behav Nutr Phys Act.* 2018;15(37). doi:10.1186/s12966-018-0672-6
- **59.** Smith R, Kelly B, Yeatman H, et al. Advertising placement in digital game design influences children's choices of advertised snacks: a randomized trial. *J Acad Nutr Diet*. 2020;120(3): 404-413. doi:10.1016/j.jand.2019.07.017
- **60**. Smits T, Vandebosch H. Endorsing children's appetite for healthy foods: celebrity versus non-celebrity spokes-characters. *Communications*. 2012;37(4):371-391. doi:10.1515/commun-2012-0021
- **61**. Vecchio MG, Nikolakis A, Galasso F, Baldas S, Gregori D. Even a very intense exposure to TV advertising promoting fruit consumption is not enough to make children eat more fruit: results from an experimental study in Italy. *Med J Nutrition Metab*. 2019;12:1-11. doi:10.3233/MNM-180254
- **62.** Dalton MA, Longacre MR, Drake KM, et al. Child-targeted fast-food television advertising exposure is linked with fast-food intake among pre-school children. *Public Health Nutr.* 2017;20(9): 1548-1556. doi:10.1017/S1368980017000520
- **63**. Emond JA, Longacre MR, Drake KM, et al. Exposure to child-directed TV advertising and preschoolers' intake of advertised cereals. *Am J Prev Med*. 2019;56(2):e35-e43. doi:10.1016/j. amepre.2018.09.015
- **64.** Fernandez MMY, Februhartanty J, Bardosono S. Association between food marketing exposure and consumption of confectioneries among pre-school children in Jakarta. *Malays J Nutr.* 2019:25 (supplement):563-573.
- **65**. Kelly B, Freeman B, King L, Chapman K, Baur LA, Gill T. Television advertising, not viewing, is associated with negative dietary patterns in children. *Pediatr Obes*. 2016;11(2):158-160. doi:10. 1111/ijpo.12057
- **66.** Minaker LM, Storey KE, Raine KD, et al. Associations between the perceived presence of vending machines and food and beverage logos in schools and adolescents' diet and weight status. *Public Health Nutr*. 2011;14(8):1350-1356. doi:10. 1017/S13689800110000449
- **67.** Olafsdottir S, Eiben G, Prell H, et al. Young children's screen habits are associated with consumption of sweetened beverages independently of parental norms. *Int J Public Health*. 2014;59(1):67-75. doi:10.1007/s00038-013-0473-2
- **68.** Scully M, Wakefield M, Niven P, et al; NaSSDA Study Team. Association between food marketing exposure and adolescents' food choices and eating behaviors. *Appetite*. 2012;58(1):1-5. doi:10.1016/j. appet.2011.09.020
- **69**. Silva RNMT, Duarte DA, de Oliveira AMG. The influence of television on the food habits of schoolchildren and its association with dental caries. *Clin Exp Dent Res*. 2020;6(1):24-32. doi:10.1002/cre2.244
- **70**. Aerts G, Smits T. Do depicted suggestions of portion size on-pack impact how much (un)healthy food children consume. *Int J Consum Stud.* 2019; 43(3):237-244. doi:10.1111/ijcs.12503

- 71. Boyland EJ, Harrold JA, Dovey TM, et al. Food choice and overconsumption: effect of a premium sports celebrity endorser. *J Pediatr*. 2013;163(2): 339-343. doi:10.1016/j.jpeds.2013.01.059
- **72.** Brown CL, Matherne CE, Bulik CM, et al. Influence of product placement in children's movies on children's snack choices. *Appetite*. 2017;114: 118-124. doi:10.1016/j.appet.2017.03.022
- **73.** Dovey TM, Taylor L, Stow R, Boyland EJ, Halford JCG. Responsiveness to healthy television (TV) food advertisements/commercials is only evident in children under the age of seven with low food neophobia. *Appetite*. 2011;56(2):440-446. doi:10.1016/j.appet.2011.01.017
- **74.** Leonard B, Campbell MC, Manning KC. Kids, caregivers, and cartoons: the impact of licensed characters on food choices and consumption. *J Public Policy Mark.* 2019;38(2):214-231. doi:10. 1177/0743915619827919
- **75.** Masserot C, Brée J. Publicité et obésité enfantine. limpact des annonces publicitaires télévisées sur les choix alimentaires des enfants. *Management & Avenir*. 2010;37(7):97-119. doi:10. 3917/mav.037.0097
- **76.** Arendt F, Naderer B, Abdollahi M, Mittelberger A, Surzhyk O, Zhou L. Television commercials and fading behavioral brand choice effects in Austrian children. *J Child Media*. 2015;9 (4):435-452. doi:10.1080/17482798.2015.1089299
- **77.** Boyland EJ, Kavanagh-Safran M, Halford JCG. Exposure to 'healthy' fast food meal bundles in television advertisements promotes liking for fast food but not healthier choices in children. *Br J Nutr*. 2015;113(6):1012-1018. doi:10.1017/S0007114515000082
- **78**. Bruce AS, Pruitt SW, Ha O-R, et al. The influence of televised food commercials on children's food choices: evidence from ventromedial prefrontal cortex activations. *J Pediatr*. 2016;177:27-32.e1. doi:10.1016/j.jpeds.2016.06.067
- **79.** Dias M, Agante L. Can advergames boost children's healthier eating habits? a comparison between healthy and non-healthy food. *J Consum Behav*. 2011;10(3):152-160. doi:10.1002/cb.359
- **80**. Dixon H, Niven P, Scully M, Wakefield M. Food marketing with movie character toys: effects on young children's preferences for unhealthy and healthier fast food meals. *Appetite*. 2017;117:342-350. doi:10.1016/j.appet.2017.07.014
- **81.** Dixon H, Scully M, Niven P, et al. Effects of nutrient content claims, sports celebrity endorsements and premium offers on pre-adolescent children's food preferences: experimental research. *Pediatr Obes*. 2014;9(2): e47-e57. doi:10.1111/j.2047-6310.2013.00169.x
- **82.** Ferguson CJ, Contreras S, Kilburn M. Advertising and fictional media effects on healthy eating choices in early and later childhood. *Psychol Pop Media Cult*. 2014;3(3):164-173. doi:10.1037/ppm0000016
- **83**. Ferguson CJ, Muñoz ME, Medrano MR. Advertising influences on young children's food choices and parental influence. *J Pediatr*. 2012;160 (3):452-455. doi:10.1016/j.jpeds.2011.08.023
- **84.** Gatou T, Mamai-Homata E, Koletsi-Kounari H, Polychronopoulou A. The short-term effects of television advertisements of cariogenic foods on children's dietary choices. *Int Dent J.* 2016;66(5): 287-294. doi:10.1111/idj.12229

- **85**. Hobin EP, Hammond DG, Daniel S, Hanning RM, Manske S. The Happy Meal effect: the impact of toy premiums on healthy eating among children in Ontario, Canada. *Can J Public Health*. 2012;103(4):e244-e248. doi:10.1007/BF03404228
- **86.** Matthes J, Naderer B. Children's consumption behavior in response to food product placements in movies. *J Consum Behav*. 2015;14(2):127-136. doi:10.1002/cb.1507
- 87. McDarby F, O'Hora D, O'Shea D, Byrne M. Taking the sweetness out of the 'Share a Coke' marketing campaign: the influence of personalized labelling on elementary school children's bottled drink choices. *Pediatr Obes*. 2018;13(1):63-69. doi:10.1111/jipo.12193
- **88.** Naderer B, Matthes J, Binder A, et al. Shaping children's healthy eating habits with food placements? food placements of high and low nutritional value in cartoons, children's BMI, food-related parental mediation strategies, and food choice. *Appetite*. 2018;120:644-653. doi:10. 1016/j.appet.2017.10.023
- **89.** Naderer B, Matthes J, Marquart F, Mayrhofer M. Children's attitudinal and behavioral reactions to product placements: investigating the role of placement frequency, placement integration, and parental mediation. *Int J Advert*. 2018;37(2):236-255. doi:10.1080/02650487.2016.1218672
- **90.** Naderer B, Matthes J, Zeller P. Placing snacks in children's movies: cognitive, evaluative, and conative effects of product placements with character product interaction. *Int J Advert*. 2018;37 (6):852-870. doi:10.1080/02650487.2017.1348034
- **91.** Putnam MM, Cotto CE, Calvert SL. Character apps for children's snacks: effects of character awareness on snack selection and consumption patterns. *Games Health J.* 2018;7(2):116-120. doi:10. 1089/g4h.2017.0097
- **92.** Royne MB, Kowalczyk CM, Levy M, Fox AK. Milk, juice, or cola? exploring the effect of product placement on children's attitudes and behavior. *Health Mark Q.* 2017;34(2):128-141. doi:10.1080/07359683.2016.1275241
- **93**. Tarabashkina L, Quester P, Crouch R. Food advertising, children's food choices and obesity: interplay of cognitive defences and product evaluation: an experimental study. *Int J Obes (Lond)*. 2016;40(4):581-586. doi:10.1038/ijo.2015.234
- **94.** Toomey DA, Francis AL. Branded product placement and pre-teenaged consumers: influence on brand preference and choice. *Young Consumers*. 2013;14(2):180-192. doi:10.1108/17473611311325573
- 95. Ülger B. Packages with cartoon trade characters versus advertising: an empirical examination of preschoolers' food preferences. *J Food Prod Mark*. 2008;15(1):104-117. doi:10.1080/10454440802470649
- **96.** Uribe R, Fuentes-García A. The effects of TV unhealthy food brand placement on children. its separate and joint effect with advertising. *Appetite*. 2015;91:165-172. doi:10.1016/j.appet.2015.03.030
- **97.** Agante L, Pascoal A. How much is "too much" for a brand to use an advergame with children? *J Prod Brand Manage*. 2019;28(2):287-299. doi:10.1108/JPBM-08-2017-1554
- **98**. Hudson S, Elliott C. Measuring the impact of product placement on children using digital brand integration. *J Food Prod Mark*. 2013;19(3):176-200. doi:10.1080/10454446.2013.724370

- **99.** Pempek TA, Calvert SL. Tipping the balance: use of advergames to promote consumption of nutritious foods and beverages by low-income African American children. *Arch Pediatr Adolesc Med.* 2009;163(7):633-637. doi:10.1001/archpediatrics. 2009.71
- **100**. Ueda MH, Porto RB, Vasconcelos LA. Publicidade de alimentos e escolhas alimentares de crianças. *Psic Teor Pesq.* 2014;30:53-61. doi:10. 1590/S0102-37722014000100007
- **101**. McAlister AR, Cornwell TB. Collectible toys as marketing tools: understanding preschool children's responses to foods paired with premiums. *J Public Policy Mark*. 2012;31(2):195-205. doi:10.1509/jppm.10.067
- **102.** Jones SC, Kervin L. An experimental study on the effects of exposure to magazine advertising on children's food choices. *Public Health Nutr.* 2011;14 (8):1337-1344. doi:10.1017/S1368980010002983
- **103**. Letona P, Chacon V, Roberto C, Barnoya J. Effects of licensed characters on children's taste and snack preferences in Guatemala, a low/middle income country. *Int J Obes (Lond)*. 2014;38(11): 1466-1469. doi:10.1038/ijo.2014.38
- **104.** McGale LS, Halford JCG, Harrold JA, Boyland EJ. The influence of brand equity characters on children's food preferences and choices. *J Pediatr*. 2016;177:33-38. doi:10.1016/j. jpeds.2016.06.025
- **105.** Reimann M, Lane K. Can a toy encourage lower calorie meal bundle selection in children? a field experiment on the reinforcing effects of toys on food choice. *PLoS One*. 2017;12(1):e0169638. doi:10.1371/journal.pone.0169638
- **106**. Roberto CA, Baik J, Harris JL, Brownell KD. Influence of licensed characters on children's taste and snack preferences. *Pediatrics*. 2010;126(1):88-93. doi:10.1542/peds.2009-3433
- **107.** Ogle AD, Graham DJ, Lucas-Thompson RG, Roberto CA. Influence of cartoon media characters on children's attention to and preference for food and beverage products. *J Acad Nutr Diet*. 2017;117 (2):265-270.e2. doi:10.1016/j.jand.2016.08.012
- **108**. Ares G, Arrúa A, Antúnez L, et al. Influence of label design on children's perception of two snack foods: comparison of rating and choice-based conjoint analysis. *Food Qual Prefer*. 2016;53:1-8. doi:10.1016/j.foodqual.2016.05.006
- **109**. Castonguay J. Sugar and sports: age differences in children's responses to a high sugar cereal advertisement portraying physical activities. *Communic Res.* 2019;46(5):579-596. doi:10.1177/0093650215587357
- **110.** de Droog SM, Valkenburg PM, Buijzen M. Using brand characters to promote young children's liking of and purchase requests for fruit. *J Health Commun*. 2011;16(1):79-89. doi:10.1080/10810730. 2010.529487
- 111. Dixon H, Scully M, Wakefield M, Kelly B, Pettigrew S. Community junior sport sponsorship: an online experiment assessing children's responses to unhealthy food v. pro-health sponsorship options. *Public Health Nutr.* 2018;21 (6):1176-1185. doi:10.1017/S1368980017003561
- **112.** Gunnarsdottir I, Thorsdottir I. Should we use popular brands to promote healthy eating among children? *Public Health Nutr.* 2010;13(12):2064-2067. doi:10.1017/S1368980010000893

- 113. Lapierre MA, Vaala SE, Linebarger DL. Influence of licensed spokescharacters and health cues on children's ratings of cereal taste. *Arch Pediatr Adolesc Med.* 2011;165(3):229-234. doi:10. 1001/archpediatrics.2010.300
- **114.** Neyens E, Smits T, Boyland E. Transferring game attitudes to the brand: persuasion from age 6 to 14. *Int J Advert*. 2017;36(5):724-742. doi:10. 1080/02650487.2017.1349029
- **115.** Arrúa A, Vidal L, Antúnez L, et al. Influence of label design on children's perception of 2 snack foods. *J Nutr Educ Behav*. 2017;49(3):211-217.e1. doi:10.1016/j.jneb.2016.10.021
- **116.** Boyland EJ, Harrold JA, Kirkham TC, et al. Food commercials increase preference for energy-dense foods, particularly in children who watch more television. *Pediatrics*. 2011;128(1):e93-e100. doi:10. 1542/peds.2010-1859
- 117. Hartmann M, Cash SB, Yeh C-H, Landwehr SC, McAlister AR. Children's purchase behavior in the snack market: can branding or lower prices

- motivate healthier choices? *Appetite*. 2017;117: 247-254. doi:10.1016/j.appet.2017.06.014
- 118. Levin AM, Levin IP. Packaging of healthy and unhealthy food products for children and parents: the relative influence of licensed characters and brand names. *J Consum Behav*. 2010;9(5):393-402. doi:10.1002/cb.326
- 119. Pettigrew S, Tarabashkina L, Roberts M, Quester P, Chapman K, Miller C. The effects of television and internet food advertising on parents and children. *Public Health Nutr.* 2013;16(12): 2205-2212. doi:10.1017/S1368980013001067
- **120.** Heard AM, Harris JL, Liu S, Schwartz MB, Li X. Piloting an online grocery store simulation to assess children's food choices. *Appetite*. 2016;96:260-267. doi:10.1016/j.appet.2015.09.020
- **121.** Simões I, Agante L. The impact of event sponsorship on Portuguese children's brand image and purchase intentions. *Int J Advert.* 2014;33(3): 533-556. doi:10.2501/IJA-33-3-533-556
- **122**. Castetbon K, Harris JL, Schwartz MB. Purchases of ready-to-eat cereals vary across US

- household sociodemographic categories according to nutritional value and advertising targets. *Public Health Nutr.* 2012;15(8):1456-1465. doi:10.1017/
- **123**. Drewnowski A, Michels S, Leroy D. The impact of crunchy Wednesdays on Happy Meal fruit orders: analysis of sales data in France, 2009-2013. *J Nutr Educ Behav*. 2017;49(3):236-240.e1. doi:10.1016/j. ineb.2016.10.005
- **124.** Ghimire N, Rao A. Comparative evaluation of the influence of television advertisements on children and caries prevalence. *Glob Health Action*. 2013;6(1):20066. doi:10.3402/gha.v6i0.20066
- **125**. Reynolds A, Mann J, Cummings J, Winter N, Mete E, Te Morenga L. Carbohydrate quality and human health: a series of systematic reviews and meta-analyses. *Lancet*. 2019;393(10170):434-445. doi:10.1016/S0140-6736(18)31809-9
- **126.** Polanin JR, Snilstveit B. Converting between effect sizes. *Campbell Syst Rev.* 2016;12(1):1-13. doi:10.4073/cmpn.2016.3