

## Photoantimicrobials in agriculture

Gilberto Ú. L. Braga<sup>a,\*</sup>, Geraldo J. Silva-Junior<sup>b</sup>, Guilherme T. P. Brancini<sup>a,\*</sup>, John E. Hallsworth<sup>c,\*</sup>,  
and Mark Wainwright<sup>d,\*</sup>

<sup>a</sup>Departamento de Análises Clínicas, Toxicológicas e Bromatológicas, Faculdade de Ciências  
Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, 14040-903, Brazil.

<sup>b</sup>Fundecitrus, Fundo de Defesa da Citricultura, Araraquara, 14807-000, Brazil.

<sup>c</sup>Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, 19  
Chlorine Gardens, Belfast, BT9 5DL, Northern Ireland, United Kingdom.

<sup>d</sup>School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool,  
L3 3AF, United Kingdom.

\*Corresponding authors. Tel.: +55-16-33154425. E-mail address: gbraga@fcfrp.usp.br (G.U.L.  
Braga); Tel: +55-16-33154723. Email address: [guilherme.brancini@gmail.com](mailto:guilherme.brancini@gmail.com) (G. T. P.  
Brancini); Tel: +44-7522259221. E-mail address: j.hallsworth@qub.ac.uk (J. E. Hallsworth); Tel:  
+44-01512312039. E-mail address: mark\_wainwright@hotmail.com (M. Wainwright).

## 22 ABSTRACT


23 Classical approaches for controlling plant pathogens may be impaired by the development of  
24 pathogen resistance to chemical pesticides and by limited availability of effective antimicrobial  
25 agents. Recent increases in consumer awareness of and/or legislation regarding environmental  
26 and human health, and the urgent need to improve food security, are driving increased demand  
27 for safer antimicrobial strategies. Therefore, there is a need for a step change in the approaches  
28 used for controlling pre- and post-harvest diseases and foodborne human pathogens. The use of  
29 light-activated antimicrobial substances for the so-called antimicrobial photodynamic treatment is  
30 known to be effective not only in a clinical context, but also for use in agriculture to control plant-  
31 pathogenic fungi and bacteria, and to eliminate foodborne human pathogens from seeds,  
32 sprouted seeds, fruits, and vegetables. Here, we take a holistic approach to review and re-  
33 evaluate recent findings on: (i) the ecology of naturally-occurring photoantimicrobials, (ii)  
34 photodynamic processes including the light-activated antimicrobial activities of some plant  
35 metabolites, and (iii) fungus-induced photosensitization of plants. The inhibitory mechanisms of  
36 both natural and synthetic light-activated substances, known as photosensitizers, are discussed  
37 in the contexts of microbial stress biology and agricultural biotechnology. Their modes-of-  
38 antimicrobial action make them neither stressors nor toxins/toxicants (with specific modes of  
39 poisonous activity), but a hybrid/combination of both. We highlight the use of photoantimicrobials  
40 for the control of plant-pathogenic fungi and quantify their potential contribution to global food  
41 security.

42 *Keywords:* antimicrobial photodynamic treatment (APDT); cellular toxicity versus stress;  
43 foodborne pathogens; global food security; photoantimicrobials; plant-pathogenic fungi

## 1. Introduction

The progressive increase in the numbers of fungi and bacteria that are tolerant to currently used antimicrobials is a major threat to human health (Fisher *et al.*, 2018; Revie *et al.*, 2018; Sabino *et al.*, 2020). Indeed, the intensive antimicrobial use raises concerns regarding both human and environmental health. Furthermore, there is an acute need to improve food security on a global scale (Kettles and Luna, 2019). Against this backdrop, it is imperative to develop new and effective strategies for the control of plant-pathogenic microorganisms. Antimicrobial photodynamic treatment (APDT) is a promising alternative to conventional antifungal and antibiotic agents which can be used for the treatment of localized infections in animal and human hosts or to kill plant- or human/animal pathogens in the environment (Calzavara-Pinton *et al.*, 2012; Dai *et al.*, 2011; de Menezes *et al.*, 2014b; de Menezes *et al.*, 2014a; de Menezes *et al.*, 2016; Gonzales *et al.*, 2017; Hamblin, 2016; Rodrigues *et al.*, 2012a; Rodrigues *et al.*, 2013; Smijs and Pavel, 2011; Vera *et al.*, 2012; Wainwright *et al.*, 2017).

To achieve microbial killing, APDT uses three primary components, namely a photosensitizer, light, and molecular oxygen. The accumulation of a photosensitizer in the cell (either inside or at the surface) of the target microbe is followed by exposure to light that, at an appropriate wavelength, excites the photosensitizer. This causes the production of reactive oxygen species (ROS), such as singlet oxygen ( $^1\text{O}_2$ ) and hydroxyl radicals ( $\bullet\text{OH}$ ), which cause biomolecular damage to the cell, effectively killing it with little to no side effects on the host (Fig. 1) (Brancini *et al.*, 2016; Calzavara-Pinton *et al.*, 2012; de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; Gonzales *et al.*, 2010; Gonzales *et al.*, 2017; St. Denis *et al.*, 2011; Wainwright *et al.*, 2017).



68

**Figure 1** – The principle of antimicrobial photodynamic treatment. A photosensitizer (PS), upon exposure to light, is initially excited to a higher-energy electronic state. Then, via intersystem crossing (transitioning between different electronic states), the excited PS transitions to a triplet state, which reacts with molecular oxygen via either an electron transfer or energy transfer reaction; Type I or Type II reactions, respectively. The former produces reactive oxygen species such as superoxide anion radical ( $\text{O}_2^{\bullet-}$ ), hydroxyl radical ( $\bullet\text{OH}$ ), and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ); the latter generates singlet oxygen ( $^1\text{O}_2$ ). These reactive oxygen species cause biomacromolecular damage and pathogen cell death.

76

Several types of photosensitizers have been used successfully to perform APDT. These include plant-produced, microbe-produced, and synthetic or semi-synthetic photoantimicrobials: chlorins, curcumins, flavins, furocoumarins, hypericins, indocyanines, phenothiazines, phthalocyanines, porphyrins, xanthenes, and others (Hamblin, 2016; Hasenleitner and Plaetzer, 2020; Temba *et al.*, 2016; Wainwright *et al.*, 2017). When reactive species such as  $^1\text{O}_2$  are

81

produced at plant surfaces, either via natural plant-produced photosensitizer or via agriculturally applied photosensitizer, they damage fungal spores and mycelia, yeasts, bacteria, as well as the ovipositors of insects that are embedded in the plant tissue (Berenbaum and Larson, 1988; Flors and Nonell, 2006; Gonzales *et al.*, 2017).

The chronic and inevitable drawback of conventional (chemical) antifungals, which, like antibiotics, have target-specific modes-of-action, is the development of microbial resistance (Wainwright *et al.*, 2017). By contrast, there is little evidence of the development of resistance to biophysical stressors (e.g. chaotropic, hydrophobic, and oxidative stressors) that act as antimicrobials at multiple target sites and/or via sites-of-action within the cell (Ball and Hallsworth, 2015; Bhaganna *et al.*, 2010; Cray *et al.*, 2013b; Cray *et al.*, 2013a; Cray *et al.*, 2014; Cray *et al.*, 2015b; Hallsworth, Heim and Timmis, 2003). Furthermore, most of the photosensitizers used in APDT exhibit low mammalian toxicity and are environmentally-friendly relative to conventional pesticides (Andrade *et al.*, 2022; Hamblin, 2016; Wainwright *et al.*, 2017). The APDT has the additional advantage of, unlike most conventional fungicides and antibiotics, being able to kill both metabolically-active and -inactive cells, including bacterial and fungal spores (de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; de Menezes *et al.*, 2016; Eichner *et al.*, 2015; Fracarolli *et al.*, 2016; Gomes *et al.*, 2011; Gonzales *et al.*, 2010; Gonzales *et al.*, 2017; Luksiene, Buchovec and Paskeviciute, 2009; Luksiene, Buchovec and Paskeviciute, 2010a; Rodrigues *et al.*, 2012a).

Additionally, APDT is not only able to control plant pathogens pre- and post-harvest (Ambrosini *et al.*, 2020; de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; Fracarolli *et al.*, 2016; Gonzales *et al.*, 2017; Luksiene and Paskeviciute, 2011; Tang *et al.*, 2021) but can kill foodborne pathogens and inactivate microbial toxins (Huang *et al.*, 2021; Jančula *et al.*, 2010). This said, the identification of effective photosensitizers, and evaluation of potential side-effects on plant- and environmental health, are imperative to the further development of APDT for use in agriculture (de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; Luksiene and Paskeviciute, 2011; Tang *et al.*, 2021; Vol'pin *et al.*, 2000).

Here, we take a compound-oriented approach, but one based on diverse lines of evidence. We evaluate the natural ecology of photosensitizer-driven antimicrobial processes in plants, including the importance of photosensitizers for phytopathogens and for plant defenses. Additionally, we examine the use of photoantimicrobials in an agricultural context to determine the potential to improve global food security. We discuss inhibitory mechanisms of photosensitizers, in relation to microbial stress biology and agricultural biotechnology, with emphasis on their use for the control of plant-pathogenic fungi, preventing spoilage of foods and feeds, and for controlling mycotoxin-producing fungi and foodborne pathogens, and global food security.

## 2. Photodynamic inactivation of plant-pathogens

As opposed to topical applications in a clinical setting, the use of APDT to control agricultural plant-pathogens would require bulk application of photosensitizers over considerable areas of land and at reasonable prices, and environmental safety is paramount. Some photosensitizers can be obtained directly from plants, algae, and cyanobacteria or from by-products of processing of fruits such as Tahiti acid lime (*Citrus aurantifolia*) and grapefruit (*Citrus x paradisi*) (Asthana *et al.*, 1993; de Menezes *et al.*, 2014a; Fracarolli *et al.*, 2016; Hudson and Towers, 1991; Temba *et al.*, 2016). Use of APDT in the field can take advantage of solar radiation, so does not need artificial light for photosensitizer activation. The high irradiances and broad emission spectrum of solar radiation can activate diverse types of photosensitizers, whether they are excited by visible light or by UV radiation (de Menezes *et al.*, 2014a; Hudson and Towers, 1991). Unlike controlled lighting provided by lasers, LEDs, or other artificial sources, incident solar radiation in the field fluctuates. This is due to factors such as climate and weather, time of year, distance from the equator, altitude, atmospheric humidity, dust, and pollution. Furthermore, the periods of illumination in the field are lengthy and the light cycles follow a diurnal pattern (Braga *et al.*, 2015).

In clinical settings, APDT can be designed to target the pathogen rather than the host by applying the photosensitizer topically on a localized area of infection and by restricting delivery of light to that area of infection (Hamblin, 2016; Wainwright *et al.*, 2017). On agricultural crops, however, such a protocol would not be feasible as the photosensitizer is applied indiscriminately on pathogen and plant, which are both exposed to solar radiation. Preventing damage to the crop plant, therefore, must be achieved by other means. Nonetheless, fruits and grains can be readily processed post-harvest, using APDT, to reduce the populations of spoilage microbes and foodborne pathogens under controlled conditions and using artificial light sources (Buchovec *et al.*, 2016; Luksiene and Paskeviciute, 2011).

## 2.1 Photodynamic inactivation of plant-pathogenic fungi

Widespread application of synthetic fungicides which have modes-of-action based on site-specific targets within the pathogen cell has been the treatment-of-choice for pre- and post-harvest control of most plant-pathogenic fungi (Ishii and Holloman, 2015; Kretschmer *et al.*, 2009; Oliver and Hewitt, 2014). However, fungicide resistance has been reported for decades in commercially important pathogens of agricultural crops, including *Alternaria*, *Aspergillus*, *Colletotrichum*, *Erysiphe*, *Fusarium*, *Mycosphaerella*, *Plasmopara*, and *Pythium* (Andrivon *et al.*, 1997; Bartlett *et al.*, 2002; Chitolina *et al.*, 2021; Deising, Reimann and Pascholati, 2008; Ishii and Holloman, 2015; Jensen *et al.*, 2016; Peres *et al.*, 2005; Ribas e Ribas *et al.*, 2016; Wong and Midland, 2007; Wong *et al.*, 2008). Current concerns about environmental and human health have given rise to recent legislation restricting the use of many of the more dangerous agrochemicals in some regions of the world. Combined with microbial resistance, this has been accompanied by decreasing numbers of commercial fungicides that are approved for agricultural use. For instance, top agricultural producing countries around the world have banned the use of or limited the access to a series of harmful pesticides (Ding *et al.*, 2019; Donley, 2019; Gunnell *et al.*, 2017). This included restrictions on the concentrations and overall quantity of approved pesticides that can


be applied (Jess *et al.*, 2014). The need for novel and/or integrated strategies to control fungi both pre- and post-harvest is now, therefore, more urgent than ever.

Taxonomically diverse fungi have been effectively killed by APDT, including human pathogens of the genera *Candida* (Dai *et al.*, 2011; Gonzales and Maisch, 2012; Rodrigues *et al.*, 2013; Rodrigues *et al.*, 2020a; Rodrigues *et al.*, 2020b) and *Trichophyton* (Rodrigues *et al.*, 2012a; Romagnoli *et al.*, 1998; Smijs *et al.*, 2014; Trigos and Ortega-Regules, 2002); entomopathogens used for biological control of insects, such as *Beauveria* (Martin, Mischke and Schroder, 1998) and *Metarhizium* (Gonzales *et al.*, 2010); saprophytic fungi that also act as opportunistic pathogens of humans, such as *Acremonium* (Lukšiene *et al.*, 2005), *Aspergillus* (DiCosmo, Towers and Lam, 1982; Friedberg *et al.*, 2001; Gilaberte *et al.*, 2011; Gonzales *et al.*, 2010; Temba *et al.*, 2016), *Cryptococcus* (Bourque *et al.*, 1985; Rodrigues *et al.*, 2012b), *Emericella* (Trigos and Ortega-Regules, 2002), *Exophiala* (Gao *et al.*, 2016), *Neurospora* (Blanc, Tuveson and Sargent, 1976; Shimizu, Egashira and Takahama, 1979), *Penicillium* (Asthana *et al.*, 1993; Gomes *et al.*, 2011), and *Rhizopus* (Liu *et al.*, 2019; Luksiene, Peciulyte and Lugauskas, 2004); endophytes, such as *Papulaspora* (Trigos and Ortega-Regules, 2002); and plant-pathogens, such as *Alternaria*, *Cladosporium* (DiCosmo, Towers and Lam, 1982; Luksiene, Peciulyte and Lugauskas, 2004; Lukšiene *et al.*, 2005; Teegne, Pretorius and Swart, 2008), *Botrytis* (Ambrosini *et al.*, 2020; Hamminger *et al.*, 2022; Kairyte, Kadys and Luksiene, 2013; Luksiene and Buchovec, 2019; Mares *et al.*, 2004; Tang *et al.*, 2021; Teegne, Pretorius and Swart, 2008), *Botryosphaeria*, *Mycosphaerella*, *Rhizoctonia*, and *Sclerotium* (Tang *et al.*, 2021; Teegne, Pretorius and Swart, 2008), *Colletotrichum* (de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; DiCosmo, Towers and Lam, 1982; Fracaroli *et al.*, 2016; Gonzales *et al.*, 2017), *Fusarium* (Asthana *et al.*, 1993; Bourque *et al.*, 1985; de Menezes *et al.*, 2016; Gao *et al.*, 2016; Kashiwabuchi *et al.*, 2013; Lazzaro *et al.*, 2004; Lukseviciute and Luksiene, 2020; Luksiene, Peciulyte and Lugauskas, 2004; Lukšiene *et al.*, 2005; Mares *et al.*, 2002; Mares *et al.*, 2004; Teegne, Pretorius and Swart, 2008; Vorobey and Pinchuk, 2008), *Magnaporthe* (Vol'pin *et al.*,



2000), *Trichothecium* (Luksiene, Peciulyte and Lugauskas, 2004), as well as the oomycetes *Pythium* and *Saprolegnia* (DiCosmo, Towers and Lam, 1982; Mares *et al.*, 2004; Tang *et al.*, 2021; Tegegne, Pretorius and Swart, 2008). The majority of studies for plant-pathogens have been performed *in vitro*; only a handful of assays have been conducted on a plant host, few experiments have emulated field conditions, and even fewer trials have assessed efficacy in the field. The small number of field trials carried out to test APDT may be explained by the need for wide-scale application of photosensitizers across large areas (where environmental safety is paramount) as opposed to topical applications in a clinical setting.

As explained above, effective APDT of plant-pathogenic fungi relies on the presence of the photosensitizer, simultaneous exposure to solar radiation, and the lifestyle of the fungal species. Some pathogens develop distinct and specialized structures such as asexual spores (e.g., conidia), sexual spores (e.g., ascospores, basidiospores) and other structures (appressoria, fruiting bodies, hyphae/mycelium, sclerotia, biofilms, etc). Invasion and colonization of plant tissue is carried out by hyphae of pathogenic fungi, but spores are usually produced on host-plant surface (Agrios, 2005; Lucas, Dyer and Murray, 2000; Mukherjee *et al.*, 2021; Peres *et al.*, 2005). Thus, these spores are usually exposed to sunlight, so are a vulnerable structure, among others, that can be targeted by APDT (Fig. 2) (de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; de Menezes *et al.*, 2016; Fracarolli *et al.*, 2016).



**Figure 2** – Stages of the pathogen lifecycle at which antimicrobial photodynamic treatment (APDT; red arrows) can prevent conidial germination and formation of appressorium (i), production of secondary conidia (ii) penetration/infection of the host-plant (iii and iv), and pathogen reproduction/dispersal (v). This schematic is based on *Citrus* infection by *Colletotrichum abscissum*.

Therefore, it is fortunate that studies into APDT of fungi have focused on conidia rather than hyphae. The antifungal assays with conidial suspensions can be readily standardized, since conidia are produced by most filamentous fungi, and the inoculum suspension is easy to prepare (Arnason *et al.*, 1986; Aver'yanov *et al.*, 2011; Braga *et al.*, 2015; Clisi, 2017; de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; DiCosmo, Towers and Lam, 1982; Gonzales *et al.*, 2010; Kairyte, Kadys and Luksiene, 2013; Luksiene, Peciulyte and Lugauskas, 2004; Mares *et al.*, 2004; Vorobey and Pinchuk, 2008). Conidia act as agents of dispersal for the majority of plant-pathogens, and can exhibit robust stress biology upon germination (Araújo *et al.*, 2020; Dijksterhuis *et al.*, 2018; Dijksterhuis, 2019; Stevenson *et al.*, 2017b; Stevenson *et al.*, 2017c). In pathogenic species, conidia are also involved in recognition and infection of the host (Barros *et al.*, 2010; Braga, Destéfano and Messias, 1999; Braga *et al.*, 2015; Nascimento *et al.*, 2010;

Peres *et al.*, 2005). The biophysical properties (electrostatic charge, hydrophobicity, etc) and chemical composition of the conidia surface differ greatly from those of hyphae (or the vegetative cells of yeast species) (Barros *et al.*, 2010; Gonzales *et al.*, 2010). Furthermore, conidial properties can vary between developmental stages, nutritional history and physiological status, fungal species and strain, and even within populations of the same strain (Rangel *et al.*, 2005; Wyatt *et al.*, 2015a; Wyatt *et al.*, 2015b). Such factors, particularly surface structure/chemistry, can influence or determine the outcomes of exposure to photosensitizers (de Menezes *et al.*, 2014b; Gonzales *et al.*, 2010; Rodrigues *et al.*, 2012a). Phototreatment of conidia-producing lesions on plant surfaces can cause a marked reduction in the viability of the fungal propagules present, thereby reducing disease transmission (Fig. 2) (Agrios, 2005; Timer and Zitko, 1991; Zulfiqar, Brlansky and Timmer, 1996).

A genus that has been the subject of APDT studies is *Colletotrichum* (de Menezes *et al.*, 2014a; de Menezes *et al.*, 2014b; Fracarolli *et al.*, 2016; Gonzales *et al.*, 2017), an ascomycete genus of common plant pathogens of both wild- and crop-plant species (Ciampi-Guillardi *et al.*, 2022; Gama *et al.*, 2022; Gonçalves *et al.*, 2021; Wharton and Diéguez-Urbeondo, 2004). *Colletotrichum* species are potent pathogens, responsible for major economic losses, especially on temperate, subtropical, and tropical fruits (Wharton and Diéguez-Urbeondo, 2004). During the asexual stage, *Colletotrichum* species produce acervuli on plant surfaces, which release mucilage containing vast numbers of unicellular conidia (Ben *et al.*, 2021; Dowling *et al.*, 2020; Zulfiqar, Brlansky and Timmer, 1996). This mucilage is readily dissolved by water, so conidia spread via rain-splash to other plants, albeit only short distances from the source (Fig. 2) (Madden, Yang and Wilson, 1996; Ntahimpera, Madden and Wilson, 1997). Strategies to minimize *Colletotrichum* epidemics are based on preventive conventional fungicide sprays during the blooming period, particularly on rainy seasons (Gama *et al.*, 2020; Silva-Junior *et al.*, 2014). However, fungicide-resistant *Colletotrichum* isolates have been reported (Deising, Reimann and Pascholati, 2008;


Dowling *et al.*, 2020; Forcelini *et al.*, 2016; Peres, Seijo and Turechek, 2010; Wong and Midland, 2007; Wong *et al.*, 2008).

Chemically diverse photosensitizers have been used in APDT to kill conidia of plant-pathogens *in vitro*, including: (i) plant metabolites, such as coumarins and furocoumarins (de Menezes *et al.*, 2014a; Fracarolli *et al.*, 2016), curcumin (Al-Asmari, Mereddy and Sultanbawa, 2017; Temba *et al.*, 2016), phenylheptatriyne (Bourque *et al.*, 1985), phenylphenalenone (Lazzaro *et al.*, 2004), polyacetylenes (Christensen and Brandt, 2006), and thiophenes (DiCosmo, Towers and Lam, 1982); (ii) semi-synthetic compounds, such as chlorophyllins (Hamming *et al.*, 2022; Luksiene and Paskeviciute, 2011) and porphyrins (Tang *et al.*, 2021; Vandresen *et al.*, 2016; Vorobey and Pinchuk, 2008); phthalocyanine metal complexes (Vol'pin *et al.*, 2000); and (iii) synthetic compounds, such as phenothiazinium dyes (e.g., methylene blue, new methylene blue N, and toluidine blue O) (de Menezes *et al.*, 2014b; de Menezes *et al.*, 2016; Gao *et al.*, 2016; Gonzales *et al.*, 2017; Liu *et al.*, 2019; Paziani *et al.*, 2019; Tonani *et al.*, 2018) and xanthenes [e.g., rose bengal (RB)] (Arboleda *et al.*, 2014). For each of these classes, we closely examine photodynamic inactivation of plant-pathogenic fungi.

### 2.1.1. Photodynamic inactivation of fungi by plant metabolites

Plants employ various strategies to protect themselves against pathogens, including the constitutive and inductive production of secondary metabolites. Some of these compounds exhibit antimicrobial activities upon photoactivation (de Menezes *et al.*, 2014a; Fracarolli *et al.*, 2016; Hudson and Towers, 1991; Larson and Berenbaum, 1988). Some plants, even those not generally considered to be phototoxic, can rapidly synthesize photosensitizers upon infection by a pathogen (Flors and Nonell, 2006; Kourany, Arnason and Schneider, 1988). Photosensitizers of plant origin include alkaloids with a structure that can be based on tryptamine (e.g., hermane), phenylalanine and tyrosine (e.g., berberine, sanguinarine) or anthranilic acid (e.g., skimmianine and other

furanoquinolines); cinnamate derivatives (e.g., coumarins and furocoumarins); polyketides (e.g., polyenes, thiophenes, quinines, and chromenes); and porphyrins that are precursors and degradation products of chlorophylls (Fig. 3) (de Menezes *et al.*, 2014a; Flors and Nonell, 2006; Fracarolli *et al.*, 2016; Hudson and Towers, 1991).



**Figure 3** – The chemical structures of common plant-produced photosensitizers. Chemical classes are shown in brackets. Structures were drawn with Marvin JS (ChemAxon).

Coumarins and furocoumarins (e.g., psoralens, angelicins) are found in the oil ducts and cuticles of species within the Apiaceae (e.g., carrots), Fabaceae (e.g., beans and lentils), Moraceae (e.g., figs), Rutaceae (e.g., *Citrus* species), among others (Asthana *et al.*, 1993; de Menezes *et al.*, 2014a; Hudson and Towers, 1991; Manderfeld *et al.*, 1997; Nigg *et al.*, 1993). They exhibit antimicrobial or insecticidal activities, via either light-independent or light-dependent

mechanisms (Bintsis, Litopoulou-Tzanetaki and Robinson, 2000; Bogucka-Kocka and Krzaczek, 2003). Both coumarins and furocoumarins are typically synthesized continuously (albeit at low levels), so are constitutive. However, their synthesis is upregulated when plants experience bacterial and fungal infection or abiotic stresses (Asthana *et al.*, 1993; Desjardins, Spencer and Plattner, 1989; Manderfeld *et al.*, 1997). In general, the highest concentrations of furocoumarins within the leaf occur at the surface, in the epidermal layer (Zobel and Brown, 1989) and in oil glands within the peel of *Citrus* fruits (Fisher and Trama, 1979).

Treatment of cells using psoralens and UV-A radiation induces pyrimidine monoadducts and interstrand crosslinks within DNA (Bordin *et al.*, 1976; Dardalhon *et al.*, 2007; Grant, Von Borstel and Ashwood-Smith, 1979). The phototoxicity of psoralens was initially thought to be a consequence of light-activated conjugation with DNA. However, Joshi and Pathak (1983) revealed that, whether linear or angular in their molecular configuration, furocoumarins can trigger production of reactive oxygen species upon exposure to light (Joshi and Pathak, 1983). It is likely that both  $^1\text{O}_2$  and superoxide radicals contribute to the phototoxicity of these compounds, especially via their adverse effects on the plasma membrane (Joshi and Pathak, 1983; Llano, Raber and Eriksson, 2003). The photoactivation of furocoumarins, and associated damage to membrane systems have been reviewed previously (Dall'Acqua and Martelli, 1991). It is widely recognized that the damage by furocoumarins occurs via multiple mechanisms (Llano, Raber and Eriksson, 2003; Potapenko, 1991; Sumorek-Wiadro *et al.*, 2020). However, in the context of fungal photosensitization, the relative importance of each of these processes has yet to be determined.

Carotenoids and other (blue-green) pigments of *Fusarium oxysporum* and *Fusarium solani* which cause root-rot and wilt in *Citrus* trees, respectively,) and *Penicillium digitatum* and *Penicillium italicum* (which cause fruit rot, as agents of green mold and blue mold, respectively) were evaluated as protectants against APDT using the plant-derived photosynthesizers 8-

methoxypsoralen (8-MOP) and  $\alpha$ -terthienyl ( $\alpha$ -T; a thiophene). For each of these fungal species, mutants in conidial pigmentation and wild-type strains were treated with each photosensitizer (at 10  $\mu\text{g mL}^{-1}$ ) and exposed to UV radiation (broad-spectrum source; emission from 300 nm to 425 nm; irradiance of 40-43  $\text{W m}^{-2}$ ). Phototreatment of conidia using  $\alpha$ -T was effective, killing most of them, regardless of fungal species. Mutants of *F. oxysporum* and *F. solani* that cannot accumulate carotenoids in their conidia were highly vulnerable to APDT. Likewise, conidial-pigment mutants of *P. digitatum* and *P. italicum* were more sensitive than the wild-type to APDT with  $\alpha$ -T. Comparisons of *Fusarium* wild-type conidia and the carotenoid-deficient mutants showed that carotenoids are less effective at protecting against APDT with 8-MOP than APDT with  $\alpha$ -T. A different result was observed in the study of *Penicillium*. The heavily pigmented blue-and-green wild-type conidia of *P. digitatum* and *P. italicum*, and a rust-colored mutant of *P. digitatum* were more tolerant to APDT with 8-MOP than their (white) mutant counterparts (Asthana and Tuveson, 1992). The authors hypothesized that carotenoids in wild-type *Fusarium* conidia protect against damage by UV-A-activated  $\alpha$ -T by quenching  $^1\text{O}_2$ , while the blue-green pigment(s) of wild-type *P. italicum* conidia (located in the cell wall) prevent DNA damage caused by 8-MOP by filtering out UV wavelengths that would otherwise activate the photosensitizer.

In a similar study, *Citrus jambhiri* leaf extracts, and the pure furocoumarins bergapten (5-methoxypsoralen; 5-MOP) and psoralen, were evaluated for phototoxicity against wild-type conidia of *F. oxysporum*, *F. solani*, *P. digitatum*, and *P. italicum* and their color-mutant strains (Asthana *et al.*, 1993). The wild-type strains of both of these *Penicillium* species were less vulnerable than their mutant strains to APDT using furanocoumarins plus UV-A radiation. A 5- $\log_{10}$  reduction in conidia of *F. oxysporum* viability was observed both in the wild-type strain and the pigmentation-mutant strains. However, wild-type conidia of *F. solani* were at least two orders of magnitude less susceptible than the white mutant conidia. Additionally, Asthana *et al.* (1993) compared ADPT treatment with bergapten of wild-type strains and mutant strains of each

*Penicillium* species, and observed different outcomes. In *P. italicum*, conidia of the wild-type and the mutant with altered brown coloration survived with minimal inactivation, whereas mutant white conidia were extremely susceptible. In *P. digitatum*, killing of wild-type and rust-mutant conidia reached 5 log<sub>10</sub>. For both *Penicillium* species, the white mutant was highly susceptible to phototreatment (with survival decreasing by as much as six orders of magnitude) (Asthana *et al.*, 1993). Similar results were observed with psoralen activated by UV-A radiation (Asthana *et al.*, 1993). Phototreatment using bergapten was one order of magnitude less effective than treatment using psoralen.

A study of APDT using either 8-MOP + isopimpinellin (both furocoumarins) or a mixture of citropten + 7-methoxy coumarin (both coumarins) compared efficacies against conidia of *Colletotrichum abscissum* (former *C. acutatum sensu lato*) and *Aspergillus nidulans* (de Menezes *et al.*, 2014a). Isopimpinellin and the mixture of coumarins were obtained from liquid residues after the industrial processing of *C. aurantifolia*. Upon treatment with the photosensitizers, conidia were exposed to solar radiation. Phototreatment with 8-MOP (50 µM) reduced survival by approximately 2 and 4 log<sub>10</sub> for *C. acutatum* after 1 and 2 h of exposure, respectively; and by approximately 4 log<sub>10</sub> for *A. nidulans*, regardless of the duration of light exposure. Also, APDT using the mixture of coumarins reduced survival by approximately 1 and 3 log<sub>10</sub> for *C. acutatum* after 1 and 2 h of light exposure, respectively. As observed for 8-MOP, phototreatment with the coumarin mixture was more effective for *A. nidulans* conidia, for which the reduction in survival was approximately 4 log<sub>10</sub>, regardless of the duration of light exposure. For *C. acutatum* conidia, isopimpinellin was the least effective treatment, reducing survival by less than 2 log<sub>10</sub> after a 2-h light exposure. Nonetheless, isopimpinellin was effective against *A. nidulans* conidia, reducing survival by approximately 4 log<sub>10</sub>. This study also reported that 8-MOP penetrates conidia and accumulates within cytoplasmic vesicles (de Menezes *et al.*, 2014a). Furthermore, APDT using crude extracts from *C. aurantifolia*, red grapefruit, and white grapefruit at 12.5 mg L<sup>-1</sup> were



performed and killed from 20% to 70% of the conidia. The *C. aurantifolia* extract was the most effective (Fracarolli *et al.*, 2016).

Given that APDT with furocoumarins and coumarins was effective against *C. abscisum*, an important question is whether the host plant would tolerate such treatment. To address this issue, the effects of phototreatment on the leaves of plant hosts *Citrus sinensis* (sweet orange), *Citrus reticulata* × *C. sinensis* hybrid (Murcott tangerine), and *Fragaria ananassa* (strawberry) were evaluated using furocoumarins and coumarins combined with solar radiation (de Menezes *et al.*, 2014a; Fracarolli *et al.*, 2016). Phototreatment with 8-MOP, isopimpinellin, and coumarins did not damage the leaves of *C. sinensis* or Murcott tangerine. However, successive daily applications of phototreatment (for 2 weeks) using the individual furocoumarins and the coumarin mixture caused considerable damage to the leaves of strawberry, with the death of epidermis- and parenchyma cells and oxidation of leaf pigments (de Menezes *et al.*, 2014a). This result is interesting because the photosensitizers used were isolated from *Citrus* spp., so these plants might be expected to have some tolerance to the photosensitizers in order to avoid self-induced damage. Conversely, strawberry plants do not produce these photosensitizers, so self-induced damage is not an issue and mechanisms to avoid it are not necessary. Nonetheless, these results show that host damage can occur, depending on plant species, so that the safety of phototreatment must be assessed on a case-by-case basis.

Polyacetylenes (polyenes) are a highly effective class of photosensitizers that occur in flowers, leaves, stems, and roots of species in the plant families Apiaceae, Asteraceae, and Campanulaceae (Binns *et al.*, 2000; Christensen and Brandt, 2006; Hudson and Towers, 1991; Mares *et al.*, 2004). Several plant species are known to produce and accumulate acetylenes, polyacetylenes, and thiophenes in response to infection by microbial pathogens (Arnason *et al.*, 1986; Bourque *et al.*, 1985; Kourany, Arnason and Schneider, 1988). They are synthesized in plant cells via the desaturation and chain shortening of fatty acids. Derivatives of polyacetylenes include the sulfur-containing thiophenes (Hudson and Towers, 1991). Many polyacetylenes

exhibit antifungal activity, and these are greatly enhanced by solar radiation or near-UV radiation (Arnason *et al.*, 1986; Bourque *et al.*, 1985; DiCosmo, Towers and Lam, 1982; Mares *et al.*, 2002; Mares *et al.*, 2004). For diverse biological systems, studies show that the phototoxicity of polyacetylenes depends on oxygen availability (Gong *et al.*, 1988). Acetylenes, especially polyacetylenes, are linear, rigid (inflexible), and lipophilic molecules that accumulate in cellular membranes. *In vitro* experiments suggested that the fungal plasma membrane is the primary site-of-action for photoactivated acetylenes, and that they are not genotoxic (Arnason *et al.*, 1986). The biological activities of four thiophene photosensitizers on *Saccharomyces cerevisiae* cells (potential genotoxicity and kill rates) were evaluated in the dark or combined with exposure to UV-A radiation (irradiance of 5 W m<sup>-2</sup> and emission peak at 350 nm) (Muzzoli and Sacchetti, 2001). None of these four thiophenes were found to be genotoxic:  $\alpha$ -terthienyl ( $\alpha$ -T); 5-(4-hydroxy-1-butenyl)-2,2'-bithienyl (BBT-OH); 5-(3-buten-1-ynyl)-2,2'-bithienyl (BBT); and 5-(4-acetoxy-1-butenyl)-2,2'-bithienyl (BBT-OAc).

Phototreatments were performed *in vitro* using three naturally-occurring thiophene derivatives as photosensitizers: 5-(3-buten-1-ynyl)-2,2'-bithienyl (Compound I, BBT); 2,2':5',2''-terthienyl (Compound II); and 2-chloro-4-[5-(penta-1,3-diynyl)-2-thienyl]but-3-ynyl acetate (Compound III), combined with exposure to UV-A (320-380 nm) against some plant-pathogenic ascomycetes, oomycetes, and zigomycetes (DiCosmo, Towers and Lam, 1982). Compounds I and II were obtained from *Echinops sphaerocephalus* and compound III from *Tagetes erecta* (both members of the Asteraceae). Conidia of *Alternaria alternata*, *Aspergillus niger*, *Cladosporium variable*, and *Colletotrichum* spp., as well as sporangiospores of *Rhizopus nigricans* were placed on media containing 0.01, 0.1, 1, and 10  $\mu$ g mL<sup>-1</sup> of the photosensitizers and exposed to UV-A radiation either immediately or after incubations of 17 and 24 h. In all cases, APDT reduced mycelial growth by 50-100% regardless of the photosensitizer or fungal/oomycete species. The oomycetes were the most susceptible, irrespective of the photosensitizer used. Phototreatment using Compound II repressed conidiogenesis in *A. niger* and sporangiogenesis in *R. nigricans*.

Germlings were generally more susceptible to APDT than non-germinated propagules. However, the viability of ungerminated conidia of *A. niger* and *R. nigricans* was unaffected by APDT with Compound II. A previous study reported that the UV-mediated cytotoxicity of Compound II occurs in *Escherichia coli* and *S. cerevisiae* only in the presence of available oxygen, which is consistent with the photodynamic basis for its mode-of-action (Arnason *et al.*, 1986).

The APDT was conducted on conidia and mycelia of the cereal pathogen *Fusarium culmorum* using phenylheptatriyne combined with near-UV radiation (300-400 nm, 5 W m<sup>-2</sup>), which was extracted from the plant *Bidens pilosa* (Asteraceae). The treatment strongly inhibited both germination of macroconidia and growth of mycelia (Bourque *et al.*, 1985). Phenylheptatriyne disrupts membrane function in *F. culmorum* via both light-dependent and light-independent mechanisms (Arnason *et al.*, 1986). Phototreatment of mycelia or macroconidia with phenylheptatriyne (10 ppm) led to increasing granulation of the cytoplasm as exposure to near-UV radiation (300-400 nm, 5 W m<sup>-2</sup>) was increased (indicating cellular damage), inhibited <sup>14</sup>C-phenylalanine uptake and respiration, and enhanced K<sup>+</sup> leakage, confirming that the plasma membrane is the primary target site of phenylheptatriyne (Arnason *et al.*, 1986).

Furthermore, the accumulation of phototoxic thiophenes was studied in *T. erecta* that was infected with *F. oxysporum* (Kourany, Arnason and Schneider, 1988). The naturally occurring thiophenes BBT-OH, BBT-OAc,  $\alpha$ -T, BBT, and 5-(3,4-diacetoxy-1-butynyl)-2,2'-bithienyl (BBT-20Ac) completely inhibited spore germination at 5  $\mu$ g mL<sup>-1</sup> (*in vitro*) in the presence of near-UV radiation (300-400 nm, 4 W m<sup>-2</sup>). Also,  $\alpha$ -T was strongly phototoxic against mycelia (Kourany, Arnason and Schneider, 1988).

The plant *Tagetes patula* (French marigold; Asteraceae) also accumulates the thiophenes  $\alpha$ -T and BBT-OH (Romagnoli *et al.*, 1998). The APDT was carried out against the plant-pathogenic fungi *Botrytis cinerea*, *Fusarium moniliforme*, and *Pythium ultimum* using the pure thiophenes  $\alpha$ -T and BBT-OH, and a methanol extract of *T. patula* (Mares *et al.*, 2002; Mares *et al.*, 2004). Mycelia were placed on media containing 5, 10, and 50  $\mu$ g mL<sup>-1</sup> of each pure thiophene,

or *T. patula* extract at a range of dilutions, and then exposed to UV-A radiation (peak at 350 nm, 5 W m<sup>-2</sup>) or simulated solar radiation. Each of these treatments inhibited growth in a concentration-dependent manner and regardless of pathogen species. In terms of reduction of growth-rate, *P. ultimum* was the most susceptible species, and *F. moniliforme* was the least susceptible (Mares *et al.*, 2002). The use of scanning electron microscopy and transmission electron microscopy revealed structural alterations to the plasma membrane of *P. ultimum*, disorganization of the cytoplasm, destruction of the nuclear envelope, and damage to the cell wall (Mares *et al.*, 2004). Comparable damage was observed in the dermatophyte fungus *Nannizzia cajetani* following APDT using BBT-OH (Romagnoli *et al.*, 1998).

Phenylphenalenones are phototoxic polycyclic aromatic compounds found mainly in Haemodoraceae and Musaceae families (Hidalgo, Kai and Schneider, 2015). They protect against pathogens, and their accumulation is upregulated in response to several fungal species (Flors and Nonell, 2006; Luis *et al.*, 1994). Light-induced <sup>1</sup>O<sub>2</sub> production and antifungal activity was reported for phenylphenalenones extracted and purified from pathogen-infected *Musa acuminata* (dwarf banana) (Lazzaro *et al.*, 2004). Conidia of *F. oxysporum* were spread onto potato dextrose agar supplemented with each of the purified phenylphenalenones and then either kept in the dark or exposed to visible light. For each photosensitizer obtained this way, antifungal activity was observed in both light and darkness; but was highest with exposure to light for the majority of the photosensitizers. Furthermore, antifungal activity was proportional to the amount of <sup>1</sup>O<sub>2</sub> produced by phenalenones. Experiments conducted in D<sub>2</sub>O-based culture media confirmed the participation of <sup>1</sup>O<sub>2</sub> in phenylphenalenone phototoxicity (Lazzaro *et al.*, 2004). Interestingly, the synthesis of 4-phenylphenalenone, which exhibited both the highest <sup>1</sup>O<sub>2</sub> yield and greatest antifungal activity of the phenylphenalenones assayed, occurs only in infected plants. By contrast, the less potent 9-phenylphenalenones occur in both healthy and infected plants from other families. Given the adverse effects of <sup>1</sup>O<sub>2</sub> on cellular systems, plants could have evolved to minimize <sup>1</sup>O<sub>2</sub> generation whenever possible (Lazzaro *et al.*, 2004). Also, some fungi have evolved

partial resistance to photosensitizers produced by plants for self-defense: for instance, *Mycosphaerella fijiensis*, the causative agent of the black sigatoka leaf-spot disease of bananas, can convert phenylphenalenones to sulfate conjugates that are inactive (Hidalgo *et al.*, 2016).

Decontaminating fungi-infected grain, maize, peanuts, or other seeds (whether used for sowing or consumption) mitigates against dispersal and mycotoxin contamination of the food supply chain. *Aspergillus flavus* is a commonly-occurring seed-borne pathogen that produces mycotoxins, including aflatoxin that is a potent carcinogen (Temba *et al.*, 2016). Thus far, chemical antifungals have been used to kill seed-borne fungal pathogens, but with varying levels of success (Dweba *et al.*, 2017). Furthermore, seed-decontamination treatments must kill fungi and inactivate mycotoxins without reducing seed viability or vigour (Lukšienė *et al.*, 2007).

Phototreatment of *A. flavus* conidia was evaluated both *in vivo* and *in vitro* using curcumin as the photosensitizer (Temba *et al.*, 2016). Curcumin, a yellow polyphenol, is obtained from the tubers of the plant *Curcuma longa*. Conidia were treated with different photosensitizer concentrations (from 5 to 100  $\mu\text{M}$ ) and exposed to light at 420 nm, both in phosphate buffered saline (PBS) solution and on maize kernels. Fluences used ranged from 12 to 84  $\text{J cm}^{-2}$  and were obtained using a xenon arc lamp with adjustable wavelength selection as the light source. Reductions of conidial viability were up to 3  $\log_{10}$  in suspensions and 2  $\log_{10}$  in maize kernels when optimal combinations of photosensitizer concentration and light fluence were used (Temba *et al.*, 2016). Also, APDT using curcumin (100 to 1000  $\mu\text{M}$ ) combined with white light (24 to 96  $\text{J cm}^{-2}$ ) were evaluated on conidia of *A. flavus*, *A. niger*, *F. oxysporum*, *Penicillium crysogenum*, and *Penicillium griseofulvum* (Al-Asmari, Mereddy and Sultanbawa, 2017). Conidia were killed by curcumin whether in spore suspensions or on the surface of agar plates.


Curcumin was also shown to be effective against *B. cinerea*: phototreatment of spores with a concentration of 800  $\mu\text{M}$  and a light fluence of 120  $\text{J cm}^{-2}$  (430 nm wavelength) completely killed the conidia. Furthermore, the toxins botrydial and dihydrobotrydial, which accumulate in spores under normal conditions, could not be detected in treated conidia (Huang *et al.*, 2021). These

results further emphasize the application of APDT not only for pathogen killing, but also for toxin inactivation or destruction.

## 2.2 Photodynamic inactivation of plant-pathogenic fungi using synthetic and semi-synthetic photosensitizers

### 2.2.1 Phenothiazines

Phenothiazines are tricyclic organic compounds derived from a thiazine heterocyclic (Fig. 4). For control of fungi in both agriculture and medicine, phenothiazinium dyes are among the most-commonly used photosensitizers. Generally, they are not toxic to mammals (Wainwright *et al.*, 2017) and are environmentally safer than widely-used commercial fungicides (Andrade *et al.*, 2022).



**Figure 4** – Chemical structure of phenothiazinium dyes commonly used for photodynamic treatment. Structures were drawn with Marvin JS (ChemAxon).

Methylene blue and other phenothiazinium photosensitizers are used for tracing cell malignancy and to treat methemoglobinemia (a blood disorder arising from the oxidation of Fe<sup>2+</sup>

to Fe<sup>3+</sup> within hemoglobin) at concentrations that are orders of magnitude higher than the minimum concentrations required to kill microbes (Shatila, Verma and Adam, 2017; Wainwright, 2010). Novel phenothiaziniums, such as the pentacyclic S137, new methylene blue N, and derivatives with basic side-chains, exhibit higher levels of antifungal activity compared to methylene blue (Dai *et al.*, 2011; de Menezes *et al.*, 2016; Rodrigues *et al.*, 2012a; Wainwright, Meegan and Loughran, 2011; Wainwright *et al.*, 2015). Recent studies have shown that the antimicrobial activity of phenothiaziniums can be enhanced by urea or inorganic salts such as potassium iodide (Nuñez *et al.*, 2015; Vecchio *et al.*, 2015), which is also observable for other photosensitizers (Bispo, Suhani and van Dijk, 2021; Castro *et al.*, 2020).

In terms of interaction between photosensitizers and target pathogen, the negatively-charged surfaces of both the fungal plasma membrane and fungal cell wall promote interactions with cationic phenothiaziniums, and several lines of evidence suggest that the plasma membrane is the primary site of damage following light-activation of these molecules (de Menezes *et al.*, 2016; Ito, 1978; Paardekooper *et al.*, 1992; Paardekooper *et al.*, 1995).

A study of *Saccharomyces fragilis* revealed that the photosensitizer toluidine blue O, which is commonly used for APDT, interacts with polyphosphates localized outside the plasma membrane without entering the cells (Tussen, Beekes and Van Steveninck, 1981). Also, it was reported that toluidine blue O does not enter cells of *S. cerevisiae* and that the photodynamic activity a consequence of its action on the extracellular medium and/or on the outer surface of the plasma membrane (Ito, 1977). A study of APDT on *S. cerevisiae* showed that toluidine blue O causes rapid oxidation of ergosterol and the subsequent accumulation of oxidized ergosterol within the plasma membrane (Bocking *et al.*, 2000). The damaged plasma membrane facilitates entry of the photosensitizer into the cytosol, which further damages intracellular membranes and biomolecules. This, in turn, leads to impaired mitochondrial function and, ultimately, cell death (Bocking *et al.*, 2000).

Indeed, APDT with methylene blue, toluidine blue O, new methylene blue N, or S137 increased the plasma membrane permeability of *F. moniliforme*, *F. oxysporum*, and *F. solani* conidia. However, only the most-lipophilic photosensitizers, new methylene blue N and S137, caused peroxidation of membrane lipids (de Menezes *et al.*, 2016), which could indicate that photosensitizer localization is heavily dependent on fungal species and/or developmental stage. Furthermore, a recent study compared the mechanism of *C. albicans* photoinactivation with new methylene blue N and S137 (Rodrigues *et al.*, 2020b). Whereas new methylene blue N targets mitochondria and reduce their membrane potential, S137 partitions into the cell membrane due to its high log *P* (6.26), where it causes destabilization and increased permeability (Rodrigues *et al.*, 2020b). Upon light exposure, S137 already present within the cell membrane increases photokilling, especially at lower light fluences (Rodrigues *et al.*, 2020b).

Gonzales and coworkers evaluated the effects of APDT with methylene blue or toluidine blue O on conidia of the saprophyte *A. nidulans* and the entomopathogen *Metarhizium robertsii* (formerly *M. anisopliae lato sensu*) (Gonzales *et al.*, 2010). Conidia of *Metarhizium* species have long been used as inoculum for control of agricultural insect pests, and are still one of the most effective fungal species for this purpose (Braga *et al.*, 2015; Brancini *et al.*, 2022; Fernandes *et al.*, 2015; Rangel *et al.*, 2015). In the study, concentrations of methylene blue and toluidine blue O ranged from 1 to 400  $\mu\text{M}$  and conidia were exposed to broad-spectrum visible light (irradiance of  $50 \text{ W m}^{-2}$ ) for 30 or 60 minutes. Mortality rates of up to 99.7% were achieved according to CFU counts, and germination of conidia which remained viable was delayed, suggesting considerable stress or damage to surviving conidia (Hamill *et al.*, 2020). Washing conidia prior to light exposure slightly reduced the effect of APDT on *M. robertsii* but strongly reduced the effect on *A. nidulans*. These findings suggest that methylene blue and toluidine blue O are taken up by each type of conidia at different rates or interact with conidia via different mechanisms (Gonzales *et al.*, 2010).

Additionally, when APDT of *A. nidulans* and *M. robertsii* was attempted for conidia in potato dextrose broth, no loss of viability occurred, indicating that some constituents of the medium may



act as antioxidants and scavengers of ROS (Gonzales *et al.*, 2010). Furthermore, conidial pigments conferred some protection against APDT with phenothiazinium photosensitizers. The conidia of the *M. robertsii* dark green wild-type and yellow-colored mutants were more resistant to APDT compared with white (albino)- or violet-colored mutants (Gonzales *et al.*, 2010). Similarly, dark green wild-type conidia of *A. nidulans* were more tolerant to APDT using methylene blue or toluidine blue O than mutants with diverse conidial pigmentation (Al-Rubeai and El-Hassi, 1986).

The efficacy of APDT of *C. abscisum*, *Colletotrichum gloeosporioides*, and *A. nidulans* conidia was determined *in vitro* using the using phenothiazinium photosensitizers methylene blue, toluidine blue O, new methylene blue N, and S137 (de Menezes *et al.*, 2014b). Minimum inhibitory concentration (MIC) was determined for each photosensitizer at different light fluences and S137 was found to be the most effective. At fluences  $\geq 20 \text{ J cm}^{-2}$ , for example, an S137 concentration of only 10  $\mu\text{M}$  was sufficient to prevent fungal growth, regardless of species. The dark toxicity of S137 was also greater than that of the other photosensitizers assayed, regardless of the species. Superior activity was also reported for S137 and new methylene blue N relative to methylene blue against *Candida* (Dai *et al.*, 2011; Rodrigues *et al.*, 2013) and *Trichophyton* spp. (Rodrigues *et al.*, 2012a). Furthermore, APDT of conidia with new methylene blue N or S137 and solar radiation or red light (emitted by LEDs) was highly effective, regardless of the photosensitizer, light source or species. When conidia were washed prior to light exposure, APDT using new methylene blue N or S137 was about as effective as APDT without washing, indicating that these photosensitizers were taken in by conidia.

Consistent with this, microscopic examination of *C. abscisum* conidia revealed that new methylene blue N and S137 accumulated in cytoplasmic vesicles (de Menezes *et al.*, 2014b). Photosensitizer entry and accumulation begins upon contact with conidia, and is independent of light exposure. This study also compared localization of these photosensitizers with the localization of the dyes Sudan III and FM4-64<sup>®</sup>, which have affinity for lipid bodies and vacuolar membranes, respectively. Both new methylene blue N and S137 accumulated in lipid bodies and

small vacuoles. Conidial treatment in the dark with S137 at concentrations  $\geq 50 \mu\text{M}$  modified the structures of the cytoplasmic organelles and caused the formation of large vesicles (de Menezes *et al.*, 2014b).

One concern about using these photosensitizers on crop plants in the field is their loss of photosensitizing potential after extended light exposure. If new methylene blue N or S137 are exposed to solar radiation prior to application to conidia, their APDT potency is reduced (de Menezes *et al.*, 2014b), a phenomenon known as photobleaching (Nassar, Wills and Harriman, 2019). For instance, when new methylene blue N and S137 were exposed to solar radiation for 3 h and then used for APDT, conidial survival was reduced only about  $3 \log_{10}$  relative to controls, compared with a reduction of  $5 \log_{10}$  if the photosensitizers were not exposed to solar radiation prior to APDT. After exposure to solar radiation for 12 h, S137 was only weakly active against *C. abscissum* conidia, whereas new methylene blue N retained more of its activity, with an ability to kill 90% ( $1 \log_{10}$ ) of the conidia (de Menezes *et al.*, 2014b). However, it is important to note that these experiments were performed under harsh conditions: photosensitizer solutions were exposed continuously to solar radiation at a tropical site ( $21.2^\circ$  latitude S) during cloudless, early-autumn days. Thus, the longevity of phenothiazinium photosensitizers is likely to be greater for most agricultural scenarios, especially if geographical location or climatic conditions involve less-intense solar exposure (de Menezes *et al.*, 2014b).

The effects of photodynamic treatment on the leaves of *C. sinensis* were evaluated using methylene blue, new methylene blue N, toluidine blue O, and S137 (each at  $50 \mu\text{M}$ ) and solar radiation (de Menezes *et al.*, 2014b). There was no apparent damage to the plant (regardless of the photosensitizer used), presumably because the photosensitizer could not penetrate the  $4\text{-}\mu\text{m}$ -thick leaf cuticle (de Menezes *et al.*, 2014b). As ROS generated during APDT have relatively short half-life, their diffusion can be very limited, thereby restricting damage to the immediate vicinity of the photosensitizer (Skovsen *et al.*, 2005). Therefore, APDT of plant-pathogenic microbes located on the host-plant surfaces proceeds without compromising the integrity of the latter.

In the last decades, human mycoses caused by species considered to be plant pathogens or fungal saprophytes (rather than human pathogens) increased dramatically (Guarro, 2013). Among the causative agents are species of *Aspergillus*, *Exophiala*, *Fusarium*, and *Rhizopus* (Gao *et al.*, 2016; Guarro, 2013; Liu *et al.*, 2019; Woo *et al.*, 2013). Invasive human infections by these fungi are usually refractory to treatment with conventional antifungals (Guarro, 2013; Liu *et al.*, 2019; Paulussen *et al.*, 2017), so APDT of these fungi may have clinical potential. In this sense, a detailed study to evaluate APDT using methylene blue, new methylene blue N, toluidine blue O, and S137 on both ungerminated and germinated microconidia of *F. moliniforme*, *F. oxysporum*, and *F. solani* were evaluated (de Menezes *et al.*, 2016). The intracellular localization of the photosensitizers as well as potential consequences of APDT were determined, including lipid peroxidation, plasma-membrane permeability, and conidial survival. Regardless of the photosensitizer used, APDT killed both ungerminated and germinating microconidia efficiently for all three *Fusarium* species (de Menezes *et al.*, 2016).


Another strategy to control and treat fungi-caused human diseases is the combination of APDT and antifungals. In this sense, the effects of APDT with methylene blue at concentrations of 8 to 32  $\mu\text{g mL}^{-1}$  either alone or in combination with standard antifungal compounds were evaluated (Gao *et al.*, 2016). The treatment was carried out for both planktonic cells and biofilms of clinical isolates of *Exophiala dermatitidis*, *F. oxysporum*, and *F. solani*. Phototreatment with methylene blue reduced survival by up to 3.8  $\log_{10}$  and 6.4  $\log_{10}$  of planktonic *Exophiala* spp. and *Fusarium* spp., respectively. The reductions for biofilms were 4.2  $\log_{10}$  and 5.6  $\log_{10}$ , respectively. However, light fluence used had to be two-fold higher than that used against planktonic cells. Application of APDT prior to the use of standard antifungals resulted in dramatic reduction of MICs when compared to antifungal treatment alone irrespective of fungal species (Gao *et al.*, 2016).

The mucoromycete *Rhizopus oryzae* causes post-harvest fruit rot and is also a common cause of mucormycosis, an aggressive and frequently fatal opportunistic fungal infection in immunocompromised individuals (Uyar and Uyar, 2018; Walther, Wagner and Kurzai, 2020).

Phototreatment with methylene blue ( $32 \mu\text{g mL}^{-1}$ ) and red light (LED, 635 nm,  $12 \text{ J cm}^{-2}$ ) completely inhibited growth and the reduction in CFU counts was up to  $4.3 \log_{10}$ . Also, APDT reduced the MIC for the antifungals itraconazole, posaconazole, and amphotericin B (Liu *et al.*, 2019).

## 2.2.2 Xanthenes

Xanthene is a tricyclic dibenzopyran organic compound that, while not possessing useful photodynamic properties, has many derivatives that are used in APDT, such as eosin Y and rose bengal (Fig. 5). The *in vitro* effects of APDT with eosin Y on the endophyte *Papulaspora immersa* and the plant-pathogen *Emericella rugulosa* were evaluated (Trigos and Ortega-Regules, 2002). Ergosterol oxidation and survival were assessed after APDT. Mycelia of these fungi were no longer viable after the treatment and cell death correlated with ergosterol photooxidation, indicating that ergosterol damage may be an effective way of achieving photoinactivation.



**Figure 5** – Chemical structure of the two best known and most used xanthene dyes. Structures were drawn with Marvin JS (ChemAxon).

In integrated pest management programs, photoactive insecticides, such as xanthene derivatives, may be combined with microbial biocontrol agents (Kim, Je and Choi, 2010; Mischke, Martin and Schroder, 1998). However, only a few studies have addressed the potential issue of APDT causing harm to biological control agents. Such biocontrol agents include viruses (e.g. baculoviruses, entomopoxviruses), bacteria (e.g. *Bacillus thuringiensis*), and entomopathogenic fungi (e.g. *Beauveria bassiana*, *Isaria fumosorosea*, and *Metarhizium* spp.).


One study evaluated APDT using the xanthene dyes phloxine B (an insecticidal photoactive compound), rose bengal, fluorescein, or eosin Y combined with white light and observed inhibited growth of the biocontrol agents *B. thuringiensis* and *B. bassiana* (Martin, Mischke and Schroder, 1998). Also, phloxine B was evaluated for compatibility with selected biocontrol fungi to determine its potential for use in integrated pest management programs (Mischke, Martin and Schroder, 1998). These studies found evidence that phloxine B, like other photosensitizers (see below), inhibit some biocontrol fungi. Phloxine B at 0.01% (w/v) inhibited the growth of *B. bassiana*, *Coniothyrium minitans*, and *Verticillium lecanii* in the presence of light. Growth of the fungus *Trichoderma virens* was inhibited by phloxine B both with and without light exposure, while growth of *Stilbella erythrocephala* was not affected by phloxine B, irrespective of illumination (Mischke, Martin and Schroder, 1998). Treatment with phloxine B at 0.005, 0.01, and 0.02 g L<sup>-1</sup> combined with visible light did not affect conidia germination of the entomopathogen *I. fumosorosea* and, interestingly, a complementary effect of phloxine B on the insecticidal efficacy of the fungus against the greenhouse whitefly, *Trialeurodes vaporariorum*, was observed (Kim, Je and Choi, 2010). Collectively, these data suggest that chemically diverse photosensitizers can inhibit phylogenetically diverse biocontrol agents such that each photosensitizer/biocontrol organism combination must be evaluated prior to consideration for use in pest-management programs.

Fungal infections of the human cornea, frequently caused by species of the genera *Aspergillus* and *Fusarium*, are termed fungal keratitis, a disease potentially leading to blindness

(Thomas, 2003). Fungal keratitis is more common in agricultural communities, and its management is restricted by the availability of effective antifungal agents, which must be able to penetrate corneal tissue (Thomas, 2003). The effects of APDT on clinical isolates of *A. fumigatus* and *F. solani* using rose bengal and riboflavin as photosensitizers were compared (Arboleda *et al.*, 2014). Rose bengal and other xanthene dyes have been routinely used in ophthalmology clinics to visualize degeneration of, or other defects in, the surface epithelium of the eye (Feenstra and Tseng, 1992; Pelloso *et al.*, 2012). Conidia suspensions were treated with rose bengal or riboflavin, both at 0.1% (w/v), and were then exposed to green or UV-A light (375 nm, 29.1 W m<sup>-2</sup>). Phototreatment with rose bengal prevented the growth of both *A. fumigatus* and *F. solani*, but no photoinactivation was observed with riboflavin (Arboleda *et al.*, 2014). The success of APDT with rose bengal at 0.1 or 0.2% and green light (15 min at 5.4 J cm<sup>-2</sup>) were reported in a pilot clinical study with patients with progressive keratitis caused by *Fusarium spp.* and *Curvularia spp.* (Naranjo *et al.*, 2019).

### 2.2.3 Porphyrins

Porphyrins are heterocyclic macrocycles composed by four pyrrole subunits interconnected via methane bridges (Fig. 6) and have been widely evaluated for APDT of phytopathogenic organisms.



**Figure 6** – Chemical structure of porphin, the simplest porphyrin, and the hematoporphyrin dimethyl ether (HPde) derivative. Structures were drawn with Marvin JS (ChemAxon).

*Fusarium* is a genus of filamentous fungi that contains many agriculturally important plant pathogens, mycotoxin producers, and opportunistic human pathogens (Dong *et al.*, 2020; Lysøe *et al.*, 2014; Ma *et al.*, 2013; Stenglein, 2009). The effects of APDT with hematoporphyrin dimethyl ether (HPde) on spores of plant pathogens (*Fusarium avenaceum* and *Trichothecium roseum*) and saprotrophic opportunistic human pathogens (*A. flavus* and *R. oryzae*) were evaluated (Luksiene, Peciulyte and Lugauskas, 2004). Sporangiospores of *R. oryzae* and conidia of the other fungal species were treated with HPde (0.25 to 71  $\mu$ M) and exposed to visible light at 300 W m<sup>-2</sup> for 15 minutes. HPde accumulated within the spores and exhibited dark toxicity regardless of fungal species, though its potency as an inhibitor of germination varied with fungal species, with *A. flavus* being more susceptible than the other species tested. The APDT using HPde and

visible-light exposure inhibited spore germination for all species, but *A. flavus* and *R. oryzae* were more susceptible than the other fungi. *In-vitro* APDT using HPde and visible-light was also effective at killing conidia of the plant-pathogen *A. alternata* and saprotrophic/human pathogen *Acremonium strictum* (Lukšiene *et al.*, 2005).

APDT of *F. culmorum* and *Fusarium poae* conia was evaluated *in vitro* using protoporphyrin IX, which is a hydrophobic dye that localizes to cell membranes (Vorobey and Pinchuk, 2008). Conidia were treated with protoporphyrin IX (1 to 4  $\mu\text{M}$ ) and were then exposed to visible light at 150  $\text{W m}^{-2}$ . Phototreatment resulted in protein and lipid oxidation, increased plasma-membrane permeability, and reduced conidial viability. At 4  $\mu\text{M}$  protoporphyrin IX and a fluence of 20  $\text{J cm}^{-2}$ , germination decreased by 55 and 96% for *F. culmorum* and *F. poae*, respectively.

Many studies have attempted to improve the efficiency of APDT with porphyrins by modifying their structure and producing a series of cationic derivatives. For instance, APDT of conidia from the saprotrophic fungus *Penicillium chrysogenum* was carried out using five cationic porphyrins (each at 50  $\mu\text{M}$ ) and white light (irradiance 2,000  $\text{W m}^{-2}$ , 20 min) (Gomes *et al.*, 2011). The most effective porphyrin, 5,10,15,20-Tetrakis(*N*-methylpyridinium-4-yl)porphyrin tetraiodide, caused a 4.1  $\log_{10}$  reduction in conidial viability. The size of the *N*-alkyl chain was shown to correlate with photoinactivation efficiency, mainly by affecting the solubility of the photosensitizer and its binding to conidia. In this sense, the best photosensitizer was the molecule with the shortest carbon chain, suggesting that the increase of the *N*-alkyl length of all four alkyl chains does not improve the photodynamic efficiency. The amount of photosensitizer incorporated by conidia was a determinant for photoinactivation efficiency and varied among the different porphyrins. Accordingly, examination using light microscopy revealed that all of the porphyrins penetrated conidia, but some showed a more uniform distribution within cells whereas others localized to the plasma membrane (Gomes *et al.*, 2011).

*Colletotrichum graminicola* is a destructive pathogen of maize causing both stalk rot and leaf blight (Damm *et al.*, 2010). Treatment of *C. graminicola* conidia was carried out using five




cationic *meso*-(1-methyl-4-pyridinio)porphyrins which have phenyl or 1-methyl-4-pyridinio group at the macrocycle *meso* position (Vandresen *et al.*, 2016). This was performed using porphyrin concentrations from 1 to 25  $\mu\text{M}$  and fluences ranging from 30 to 120  $\text{J cm}^{-2}$  (emitted from a 250-W halogen lamp). Considering the lowest photosensitizer concentration and the lowest light fluence that enabled photoinactivation, the porphyrins efficiencies were ranked as triple-charged (1  $\mu\text{M}$  with a fluence of 30  $\text{J cm}^{-2}$ ) > double-charged-*trans* (1  $\mu\text{M}$  with a fluence of 60  $\text{J cm}^{-2}$ ) > tetra-charged (15  $\mu\text{M}$  with a fluence of 90  $\text{J cm}^{-2}$ ) > mono-charged (25  $\mu\text{M}$  with a fluence of 120  $\text{J cm}^{-2}$ ). The APDT using the triple-charged porphyrin at 1  $\mu\text{M}$  and 30  $\text{J cm}^{-2}$  killed all conidia. Double-charged-cys-porphyrin killed conidia in the dark, i.e without light-activation. The porphyrins that presented high  $^1\text{O}_2$  quantum yields and accumulated to a high degree in conidia were the best photosensitizer (Vandresen *et al.*, 2016).

APDT with a porphyrin (TMPyP) metal-organic framework (PS@MOF) was evaluated against the plant-pathogenic fungi *Sclerotinia sclerotiorum*, *Pythium aphanidermatum* and *B. cinerea* both *in vitro* and *in planta* (Tang *et al.*, 2021). *In vitro* APDT with PS@MOF strongly inhibited mycelia growth of the three fungal species at photosensitizer concentrations of 6, 12, and 24  $\text{mg L}^{-1}$ . Also, APDT was able to control *S. sclerotiorum* on cucumber with efficiency equal to the dicarboximide fungicide dimethachlon without causing damage to the host plant (Tang *et al.*, 2021). Also, APDT with the anionic porphyrin tetra-4-sulfonatophenyl porphyrin tetra-ammonium (TPPS) was tested against *B. cinerea* both *in vitro* and *in planta* (Ambrosini *et al.*, 2020). *In vitro* APDT with TPPS at 1.5  $\mu\text{M}$  combined with white light caused potent inhibition of mycelium growth. Also, mycelium pre-treated with TPPS was unable to infect detached leaves of any of the three grapevine clones from Chadornnay, Merlot, and Sauvignon. Importantly, treatment with the photosensitizer at 12.5  $\text{M}$  did not damage the plants (Ambrosini *et al.*, 2020).

#### 2.2.4 Chlorins, bacteriochlorins, chlorophyllins, and chitosan

A chlorin, the core chromophore of a chlorophyll, is a dihydroporphyrin macrocycle that contains three pyrrole rings and one pyrroline ring (Fig. 7) (Taniguchi and Lindsey, 2017). Several of the clinically important photosensitizers are chlorins, including m-tetrahydroxyphenylchlorin, benzoporphyrin derivative, radachlorin, and chlorin e6 (Abrahamse and Hamblin, 2016). Structurally, chlorins have a double bond in one pyrrole ring reduced (Fig. 7) whereas bacteriochlorins have two pyrrole rings with reduced double bonds (Fig. 7) (Martinez De Pinillos Bayona *et al.*, 2017). The bacteriochlorin group also includes important clinical photosensitizers (Abrahamse and Hamblin, 2016) and both chlorins and bacteriochlorins have been evaluated as photosensitizers for use in APDT against plant pathogens and foodborne human pathogens (Lopez-Carballo *et al.*, 2008; Luksiene and Paskeviciute, 2011; Uliana *et al.*, 2014).



**Figure 7** – Chemical structures depicting the differences between chlorin (20  $\pi$  electrons) and bacteriochlorin (18  $\pi$  electrons). The structure of a chlorin derivative, sodium chlorophyllin, is also shown. Structures were drawn with Marvin JS (ChemAxon).

As mentioned earlier, positively-charged photosensitizers tend to be more efficient as antimicrobials because of their affinity for the negatively-charged cell surfaces of bacteria and fungi (Hamblin, 2016). Indeed, the attachment of polycationic polymers such as poly-L-lysine and polyethylenimine to chlorins (that otherwise do not possess any intrinsic positive charge) enhanced their efficiency in APDT (Hamblin *et al.*, 2002; Tegos *et al.*, 2006). Interestingly, it was reported that an asymmetric dicationic bacteriochlorin was significantly more active against Gram-positive bacteria and fungi than a symmetrically-substituted tetracationic bacteriochlorin (Huang *et al.*, 2014). Recently, thiopyridinium and methoxypyridinium chlorin derivatives were tested

against *F. oxysporum* (Sierra-Garcia, Cunha and Lourenço, 2022). Among these, a free-base thiopyridinium chlorin was shown to be the most effective compound, achieving complete conidial killing after 15 min ( $45 \text{ J cm}^{-2}$ ) of white-light exposure at a concentration of  $15 \mu\text{M}$ . The other compounds required either higher concentrations and/or longer exposure to light (Sierra-Garcia, Cunha and Lourenço, 2022). These results show that chlorin derivatives can be potent photosensitizers against plant-pathogenic fungi.

Chlorophyllins are semi-synthetic, water-soluble salts derived from chlorophyll and are also approved as food colorants in both the EU and the USA (Fig. 7) (Wrolstad and Culver, 2012). Chlorophyllins have been used as photosensitizers for photodynamic treatment of several types of cancers as well as for APDT (Afrasiabi *et al.*, 2020; Luksiene and Paskeviciute, 2011; Luksiene and Buchovec, 2019; Lukseviciute and Luksiene, 2020). Treatment using sodium salts of chlorophyllin (Na-Chl) and visible light was evaluated for post-harvest control of spoilage microbes on strawberries (Luksiene and Paskeviciute, 2011). Naturally-contaminated strawberry fruits were soaked in Na-Chl at  $1 \text{ mM}$  for 5 min and illuminated for 20 min with visible light ( $400 \text{ nm}$  and irradiance of  $120 \text{ W m}^{-2}$ ). The growth of fungi and total aerobic mesophiles was reduced by 86 and 97%, respectively. Consequently, shelf life of treated fruits was extended by two days. Also, APDT increased total antioxidant activity of the fruit extracts by almost 20% but did not impact the amounts of either anthocyanins or phenols, nor caused changes to fruit color (Luksiene and Paskeviciute, 2011).

Chitosan is a cationic linear polysaccharide produced commercially by deacetylation of chitin. Additionally, chitosan can form films and exhibits antimicrobial activity against a wide range of microorganisms (Dutta, Tripathi and Dutta, 2012; Ke *et al.*, 2021). A chlorophyllin-chitosan complex (Chl-CHS) has been used in APDT to kill microorganisms on fruit and grains surfaces. The APDT using Chl-CHS was assessed for the microbiota of strawberries (Luksiene and Paskeviciute, 2011). Naturally-contaminated strawberries were soaked for 30 min in 0.1% (w/v) chitosan,  $1.5 \cdot 10^{-5} \text{ M}$  chlorophyllin/0.1% chitosan or  $1.5 \cdot 10^{-5} \text{ M}$  chlorophyllin and were exposed to

405-nm radiation for 60 min (fluence of  $38 \text{ J cm}^{-2}$ ). Chitosan combined with light exposure reduced colony forming units of fungi by  $0.4 \log_{10}$ ; chlorophyllin-based APDT reduced colony forming units by as much as  $0.9 \log_{10}$ ; and APDT using Chl-CHS reduced colony forming units by  $1.4 \log_{10}$  (Luksiene and Buchovec, 2019), showing the superior performance of the complex. No additional photosensitization-induced free radical was found in the strawberry matrix and no changes were caused to color, texture, and nutritional or visual quality of the fruits (Luksiene and Buchovec, 2019).


Furthermore, APDT using Chl-CHS was evaluated against fungi present on the surface of wheat grains (Buchovec and Lukšienė, 2015). Wheat grains were soaked in 0.1% Chl-0.001% chitosan solution and were then exposed to 405-nm radiation for 30 min (at a fluence of  $30 \text{ J cm}^{-2}$ ). This treatment reduced the number of colony forming units of fungi by  $0.68 \log_{10}$  (mortality ~80%). APDT with Chl-CHS was also evaluated to inactivate *Fusarium graminearum* mycelia *in vitro* and conidia on artificially-contaminated wheat grains. *In vitro* APDT with 0.005%/Chl-0.5% chitosan combined with exposure to 405-nm radiation inhibited mycelium growth but did not kill the fungus. The results of the study did not make it clear what percentage of conidia was inactivated by APDT on the grain surface, but the treatment did not affect the vigor and viability of the grains (Buchovec and Lukšienė, 2015).

Chitosan has also been chemically combined with other photosensitizers, such as protoporphyrin XI and riboflavin, yielding the conjugates PPIX-CHS and RF-CHS, respectively (Dibona-Villanueva and Fuentealba, 2021; Dibona-Villanueva and Fuentealba, 2022). Both conjugates were used for the APDT of *P. digitatum*. The PPIX-CHS compound inhibited fungal growth by 100% at 0.005% (w/v) after one hour of white-light exposure. Interestingly, using a mixture of unconjugated protoporphyrin IX and chitosan did not result in fungal killing, showing the improved properties of the conjugate (Dibona-Villanueva and Fuentealba, 2022). The RF-CHS conjugate was also effective against the fungus, albeit only at higher concentrations (0.5-0.9%) compared to PPIX-CHS (Dibona-Villanueva and Fuentealba, 2021). Nonetheless, the

conjugation of riboflavin and chitosan greatly improved the photodynamic properties of the former: the RF-CHS conjugate had higher  $^1\text{O}_2$  yield and improved interaction with fungal cells compared to riboflavin alone (Dibona-Villanueva and Fuentealba, 2021).

### 2.2.5 Phthalocyanines

Phthalocyanines are two-dimensional, 18  $\pi$ -electron aromatic porphyrin analogues consisting of four isoindole subunits linked together via nitrogen atoms (Fig. 8) (Claessens, Hahn and Torres, 2008). Phototreatment using different types of phthalocyanines can kill various fungi, including plant-pathogenic species (Prandini *et al.*, 2022; Rodrigues *et al.*, 2020a; Rodrigues *et al.*, 2012b; Vol'pin *et al.*, 2000).



Phthalocyanine

**Figure 8** – Basic chemical structure of a phthalocyanine. The structures were drawn with Marvin JS (ChemAxon).

The filamentous ascomycete *Magnaporthe oryzae* (formerly *Magnaporthe grisea*) causes rice blast, the most economically devastating disease of cultivated rice (Wilson and Talbot, 2009). Rice blast occurs throughout crop areas and is typically responsible for a 10 to 30% reduction of rice yield (Martin-Urdiroz *et al.*, 2016; Yan and Talbot, 2016). Use of conventional antifungals to

control *M. oryzae* has proven expensive and of limited efficacy (Yan and Talbot, 2016). Therefore, the effects of 20 phthalocyanine-metal complexes on conidia germination of *M. grisea* (= *M. oryzae*) both with and without light exposure were evaluated (Vol'pin *et al.*, 2000). Photosensitizer concentrations tested ranged from 0.5 to 100  $\mu\text{g mL}^{-1}$  and light was provided by a xenon lamp with water filter. Inhibition of conidia germination varied between zero and 78% depending on the type of phthalocyanine-metal complex. The authors also evaluated the phytotoxicity of these complexes using rice-plant leaves. Most of the phthalocyanine-metal complexes did not damage the leaves, but some did cause chlorotic or necrotic lesions that were however local and negligible at the low photosensitizer concentrations required to control the pathogen. Interestingly, some of these complexes also protected rice plants from blast disease. The authors hypothesize that, *in planta*, the phthalocyanine redox activity in the dark along with its photosensitizing ability promote the generation of ROS, which damage the fungus and, consequently, favor disease resistance (Vol'pin *et al.*, 2000).

The chemical derivatization of phthalocyanine-metal complexes is an important process to obtain better photosensitizers. For instance, a thiopyridinium derivative of Zn(II) phthalocyanine was shown to be very effective against *C. abscissum*, achieving complete conidial killing at only 5  $\mu\text{M}$  and a fluence of 37.5  $\text{J cm}^{-2}$  (640-nm red light) (Prandini *et al.*, 2022). Furthermore, ammonium derivatives of Si(IV) phthalocyanines were shown to be effective against *E. coli* and *S. aureus*, also at low concentrations (3 and 6  $\mu\text{M}$ ) and a white-light fluence of 540  $\text{J cm}^{-2}$  (Gamelas *et al.*, 2022).

## 2.2.6 5-aminolevulinic acid

Microbial contamination of seed- and bean sprouts that are produced for human consumption is a chronic problem in the food supply chain (Mir *et al.*, 2021; Symes, Goldsmith and Haines, 2015). Sprouts are produced from plants such as legumes (e.g. beans, chickpeas, lentils, peas, and soybean), cereals (e.g. rye, wheat, barley, and oats), and vegetables (including

alfalfa, radish, mustard, and other *Brassica* species). The methods currently employed to decontaminate seeds and beans prior to sprouting (e.g. washing with chlorine, chlorine dioxide, sodium and calcium hypochlorite, and hydrogen peroxide) have a number of drawbacks (Lukšienė *et al.*, 2007; Mir *et al.*, 2021).

Studies have been carried out to evaluate APDT of seeds (prior to sprouting) using 5-aminolevulinic acid (5-ALA) (Lukšienė *et al.*, 2007; Lukšienė and Zukauskas, 2009). 5-ALA is a naturally-occurring precursor to heme synthesis in eukaryotic and prokaryotic cells, which induces the production of the endogenous photosensitizers protoporphyrin IX, uroporphyrin, and coproporphyrin (Kamp *et al.*, 2005). When present in the extracellular milieu, 5-ALA is taken up by cells of bacteria, yeast, and filamentous fungi. This can in turn stimulate synthesis of porphyrin-type photosensitizers which can be light-activated to enable control of the microbes (Harris and Pierpoint, 2012; Kamp *et al.*, 2005; Lukšienė and Zukauskas, 2009; Polmickaitė-Smirnova *et al.*, 2022).

The APDT using 5-ALA has proved effective to control fungal contaminants of wheat grains (Lukšienė *et al.*, 2007). Wheat grains naturally contaminated with fungi were soaked for 4 h in a solution of NaCl (5%) with 5-ALA (6 mM) at 26 °C and were then exposed to light (522 nm, emitted by an incandescent lamp equipped with optical filters). Twelve hours after exposure, grains were examined for the presence of viable fungi, which revealed that *Apergillus* spp., *Fusarium* spp., *Mucor* spp., and *Rhizomucor* spp. were susceptible to APDT with 5-ALA while *Acremonium* was not. Treatment with 5-ALA not only reduced fungal contamination but also stimulated the growth of wheat seedlings and roots during the subsequent sprouting procedure, all without impairing grain germination and viability (Lukšienė *et al.*, 2007). A similar molecule, methyl aminolevulinate, is the methyl ester of 5-ALA and has also been used as photosensitizer in APDT. Methyl aminolevulinate-based APDT was used against finger nail infections of *F. oxysporum* and *Aspergillus terreus* (Gilaberte *et al.*, 2011). This treatment cured refractory onychomycosis caused by these fungi.



### 2.2.7 Riboflavin and riboflavin derivatives

Riboflavin, widely known as vitamin B<sub>2</sub>, is a water-soluble compound which can be synthesized by plants and microorganisms, but is essential for animals as they lack an endogenous biosynthetic pathway (Schwechheimer *et al.*, 2016). Riboflavin acts as cofactors for oxidoreductases as well as prosthetic groups for enzymes in the  $\beta$ -oxidation pathway (Massey, 2000). The vitamin is synthesized biotechnologically using microorganisms, and is mainly used as feed and food additive as well as for pharmaceutical applications (Schwechheimer *et al.*, 2016).

When exposed to visible light, riboflavin acts as a potent photosensitizer by producing  $^1\text{O}_2$  (Bäumler *et al.*, 2012; Cardoso, Libardi and Skibsted, 2012; Fuentealba *et al.*, 2015). However, due to the lack of positive charge, riboflavin is not a very effective photosensitizer for use in APDT (Nielsen *et al.*, 2015). For instance, riboflavin combined with UV-A could not inhibit the growth of either hyphae or conidia of *A. fumigatus*, *F. solani*, and other *Fusarium* spp. (Arboleda *et al.*, 2014; Kashiwabuchi *et al.*, 2013; Sauer *et al.*, 2010). However, the addition of riboflavin at 250  $\mu\text{M}$  significantly enhanced the efficacy of simulated solar disinfection at 150  $\text{W m}^{-2}$  against a variety of microorganisms, including *F. solani*, with mortality rates of 100% being achieved after a 6-h exposure (Heaselgrave and Kilvington, 2010). Pretreatment of fungi with amphotericin B can increase the effectiveness of APDT using riboflavin and UV-A (365 nm, 30  $\text{W m}^{-2}$ ), according to an *in vitro* study of *A. fumigatus*, *C. albicans*, and *Fusarium* spp. (Sauer *et al.*, 2010). Also, synthetic riboflavin derivatives that are positively charged have been successfully used in APDT against *Bacillus* endospores (Eichner *et al.*, 2015). In this sense, and as mentioned earlier, the conjugation of riboflavin with chitosan improved the interaction of the former with *P. digitatum* cells (Dibona-Villanueva and Fuentealba, 2021).

## 3. Post-harvest photoinactivation of foodborne pathogens and microbial contaminants

Plant surfaces are typically subject to extreme fluctuations in water activity (Stevenson *et al.*, 2015b), solar radiation, temperature, and other parameters. As such, they tend to be nutrient-poor (Lievens *et al.*, 2015) and can be inhospitable for human pathogens such as enteric bacteria. Damage to plant tissues, such as that caused by plant pathogens or food processing, can create a habitable substrate for, and so promote the growth of, enteric pathogens of humans (Heaton and Jones, 2008; Weiman, 2014).

Among the main foodborne bacterial pathogens are *Bacillus cereus*, *E. coli*, *Listeria monocytogenes*, and *Salmonella enterica*, none of which is capable of growth on low water-activity surfaces of  $< 0.850$  (do Prado-Silva *et al.*, 2022; Santos *et al.*, 2015; Stevenson *et al.*, 2015a). Microbial contamination of fruits and vegetables and other types of food/feeds can also greatly shorten their shelf life, especially by fungal psychrophiles and xerophiles, some of which are capable of growth even at subzero temperatures and at  $\leq 0.585$  water activity (Chin *et al.*, 2010; Collins and Buick, 1989; Stevenson *et al.*, 2017c; Stevenson *et al.*, 2017b; Stevenson *et al.*, 2017a). Several studies, carried out *in vitro* using different photosensitizers, have established that APDT can efficiently kill diverse foodborne pathogens and spoilage microbes, including cells and spores of bacteria (Aponiene *et al.*, 2015; Buchovec *et al.*, 2017; Eichner *et al.*, 2015; Gulías *et al.*, 2020; Luksiene and Buchovec, 2019; Luksiene and Brovko, 2013; do Prado-Silva *et al.*, 2021; do Prado-Silva *et al.*, 2022; Silva *et al.*, 2018; Sobotta *et al.*, 2019).

Natural and semi-synthetic photosensitizers such as 5-ALA (Buchovec, Vaitonis and Luksiene, 2009; Luksiene, Buchovec and Paskeviciute, 2009), sodium magnesium and sodium copper chlorophyllin (approved as food additives E-140 and E-141, respectively) (Buchovec *et al.*, 2016; Buchovec *et al.*, 2017; Hasenleitner and Plaetzer, 2020; Luksiene and Buchovec, 2019; Luksiene and Paskeviciute, 2011; Luksiene, Buchovec and Paskeviciute, 2010a; Luksiene, Buchovec and Paskeviciute, 2010b), curcumin (approved as the food additive E-100) (Gong *et al.*, 2020; Glueck *et al.*, 2017; Hu *et al.*, 2018; Temba *et al.*, 2016; Tortik, Spaeth and Plaetzer, 2014), furocoumarins (de Menezes *et al.*, 2014a; Fracarolli *et al.*, 2016; Ulate-Rodríguez *et al.*,

1997), hypericin (Aponiene *et al.*, 2015; Kairyte *et al.*, 2012), and riboflavin derivatives (Eichner *et al.*, 2015) are among the most studied in relation the food microbiology.

APDT of the Gram-negative *S. enterica* using 5-ALA efficiently kills the bacterium, according to *in vitro* studies (Buchovec, Vaitonis and Luksiene, 2009). Bacterial cells were taken from an exponential-phase culture and incubated with 7.5 mM 5-ALA for up to 1 h in the dark, and then exposed to visible light (LED, with emission at 400 nm and irradiance of 200 W m<sup>-2</sup>). Bacterial photodynamic inactivation was dependent on the incubation time with 5-ALA and fluence. Viable cell number was reduced by up to 6 log<sub>10</sub>. *In vitro* APDT with 5-ALA was also able to kill the Gram-positive *B. cereus*, regardless of whether vegetative cells or spores were tested (Luksiene, Buchovec and Paskeviciute, 2009). Photoinactivation of *B. cereus* on the surfaces of grains and packaging materials was also observed (Luksiene, Buchovec and Paskeviciute, 2009). Cells and spores of *B. cereus* suspended in PBS or on the surface of food-packaging material and wheat grains were treated with 5-ALA (3 to 7.5 mM) and then exposed to visible light. Survival of vegetative cells in suspension was reduced by 6.3 log<sub>10</sub> and on the surface of food packing by 4 log<sub>10</sub>. *B. cereus* spores both in suspension and on packaging material were also susceptible to APDT and survival was reduced by 3.7 and 2.7 log<sub>10</sub>, respectively. Similar results were observed for the highly pathogenic *L. monocytogenes* which was killed by 5-ALA-based APDT both in suspension (survival reduction up to 4 log<sub>10</sub>) and as biofilm on the surface of packaging material (survival reduction up to 3.1 log<sub>10</sub>) (Buchovec, Paskeviciute and Luksiene, 2010). *B. cereus* spores were also efficiently inactivated by APDT (> 3 log<sub>10</sub> reduction in survival) when a tricationic porphyrin was used as the photosensitizer. However, the susceptibility of spores to porphyrin-based APDT was highly variable among different species of *Bacillus* (da Silva *et al.*, 2012).

Sodium magnesium chlorophyllin (E-140) and sodium copper chlorophyllin (E-141) were incorporated into gelatin films, and their potential to inhibit *Staphylococcus aureus* and *L. monocytogenes* was evaluated (Lopez-Carballo *et al.*, 2008). Bacterial cell suspensions were spread on the surface of tryptone soy agar. Control gelatin films (no photosensitizer) and gelatin

films supplemented with E-140 or E-141 were placed on the surface of the inoculated agar, and then irradiated for 5 or 15 min. Following these treatments, plates were incubated for 24 h and bacterial colonies were then counted. Results showed that the E-140- and E-141-based APDT reduced the number of colony forming units of *S. aureus* and *L. monocytogenes* by 5 and 4 log<sub>10</sub>, respectively. *In vitro* APDT with sodium magnesium chlorophyllin at 5 µM combined with blue light (433 nm, 6.6 J cm<sup>-2</sup>) reduced the survival of *S. aureus* by more than 7 log<sub>10</sub> (Hasenleitner and Plaetzer, 2020). However, APDT with the two chlorophyllins had no effect on the viability of the Gram-negative bacteria *E. coli* and *Salmonella* spp. (Lopez-Carballo *et al.*, 2008).

APDT using Na-Chl on survival of cells and spores of *B. cereus* (Luksiene, Buchovec and Paskeviciute, 2010a) and *L. monocytogenes* (Luksiene, Buchovec and Paskeviciute, 2010b) was evaluated *in vitro*, both in suspension and on the surface of yellow packaging trays (polyolefin-mixture of polyethylene and polypropylene). Cells and spores of *B. cereus* suspended in PBS or on packing material were incubated with Na-Chl ( $7.5 \times 10^{-8}$  to  $7.5 \times 10^{-5}$  M) and then exposed to visible light (peak emission at 400 nm and irradiance of 200 W m<sup>-2</sup>). Treatment with  $7.5 \times 10^{-7}$  M of Na-Chl reduced the survival up to 7 log<sub>10</sub> of the cells of *B. cereus* suspended in PBS and those on the surface of packaging trays. *B. cereus* spores were more tolerant to APDT than cells, but were also killed depending on photosensitizer concentration and light-exposure time. Treatment with  $7.5 \times 10^{-5}$  M of Na-Chl reduced the survival of the spores suspended in PBS up to 4 log<sub>10</sub>. Reduction of the survival of the spores on packaging tray was 5 log<sub>10</sub> at  $7.5 \times 10^{-5}$  M of Na-Chl.

Also, APDT using Na-Chl was tested on the survival of thermosensitive and thermotolerant strains of *L. monocytogenes* both in suspension and on the surface of yellow packaging trays (Luksiene, Buchovec and Paskeviciute, 2010b). Phototreatment reduced the survival of both strains up to 7 log<sub>10</sub> when cells were suspended in PBS and killed all the cells when they were on packaging trays. The APDT of *B. cereus* and *L. monocytogenes* using Na-Chl was considerably more effective than washing with 200 ppm sodium hypochlorite (Luksiene, Buchovec and Paskeviciute, 2010a; Luksiene, Buchovec and Paskeviciute, 2010b). Mechanistic studies

conducted with Gram-negative bacteria has shown that during Na-Chl-based APDT, chlorin binds to the surface of the bacterial cell causing  $^1\text{O}_2$ -mediated membrane damage and cell wall disruptions, increased release of intracellular components, and cell death (Žudytė *et al.*, 2020).

As presented earlier, APDT using Na-Chl and visible light was tested in strawberries that had been inoculated with *L. monocytogenes* (Luksiene and Paskeviciute, 2011). Strawberries were inoculated with *L. monocytogenes*, soaked in 1 mM Na-Chl solution for 5 min, and then exposed to visible light (400 nm, irradiance of  $120 \text{ W m}^{-2}$ ) for 30 min. The treatment reduced the viability of the cells by  $1.8 \log_{10}$  compared to control samples.

The effect APDT with the Chl-CHS chlorophyllin-chitosan complex on survival of *L. monocytogenes* on the surface of wheat grains was evaluated (Buchovec and Lukšienė, 2015). Also, coating of strawberries with Chl-CHS and illumination with visible light at  $76 \text{ J cm}^{-2}$  inactivated yeast/microfungi on the fruits by 1.4 log and prolonged the shelf life by 3 days without any negative effect on the fruits (Luksiene and Buchovec, 2019). Details for these two studies were discussed above and can be found on section 2.2.4. Furthermore, the effects of APDT using Na-Chl alone and combined with chitosan or high-power pulsed UV (200-1000 nm, peak at 260 nm) on the survival of *S. enterica* were determined *in vitro* (Buchovec *et al.*, 2017). The APDT alone reduced the survival of the bacteria by  $2.05 \log_{10}$  while APDT combined with chitosan or pulsed UV reduced the viability by 7.28 and  $7.5 \log_{10}$ , respectively. Interestingly, Na-Chl-based APDT induced the transcription of genes responsible for ROS inactivation in *S. enterica* (Buchovec *et al.*, 2017).

Hypericin-based APDT and hypericin-based APDT combined with high power pulsed light (HPPL) were evaluated for *L. monocytogenes* and *S. enterica* (Kairyte *et al.*, 2012). Cells were incubated with hypericin ( $10^{-5}$  or  $10^{-7} \text{ M}$ ) in PBS and exposed to visible light (peak emission at 585 nm and irradiance of  $38.4 \text{ W m}^{-2}$ ). For the combined treatment, after APDT, bacteria were exposed to 350 pulses of HPPL (UV fluence of  $0.023 \text{ J cm}^{-2}$ ). Hypericin interacted with the cells of both species and APDT reduced the survival of *Listeria* and *Salmonella* by 7 and  $1 \log_{10}$ ,

respectively. Electron microscopy studies showed that APDT induced total collapse of the *Listeria* cell wall, but not that of *Salmonella*. Combined treatment of APDT and pulsed light reduced the survival of *Listeria* and *Salmonella* by 6.7 to 7 log<sub>10</sub>, respectively. The effect of APDT with hypericin ( $1.5 \times 10^{-5}$  to  $1 \times 10^{-8}$  M) and visible light (585 nm, irradiance of 38.4 W m<sup>-2</sup>, and fluences up to 9.2 J cm<sup>-2</sup>) on the survival of *B. cereus* both *in vitro* and inoculated on the surface of fruits (apricots and plumes) and vegetables (cauliflowers) were also investigated (Aponiene *et al.*, 2015). Hypericin-based APDT reduced the survival of the bacteria up to 4.4 log<sub>10</sub> *in vitro*. Inactivation of mesophilic bacteria on the surface of fruits and vegetables reached up to 1.3 log<sub>10</sub>.

The use of APDT employing curcumin bound to polyvinylpyrrolidone (PVP-C) and NovaSol®-curcumin for the decontamination of *S. aureus* from cucumber, pepper, and chicken meat was evaluated (Tortik, Spaeth and Plaetzer, 2014). Both curcumin and PVP-C have been approved as food additives. Vegetables and meat were contaminated with the bacteria, sprinkled with PVP-C and NovaSol®-curcumin at concentrations of 50 and 100 µM, respectively, and illuminated immediately using visible light (emission peak at 435 nm, irradiance 94 W m<sup>-2</sup> and fluence 33.8 J cm<sup>-2</sup>). Photodynamic inactivation of *S. aureus* caused a mean reduction of 2.6 log<sub>10</sub> on cucumbers, 2.5 log<sub>10</sub> on pepper, and 1.7 log<sub>10</sub> on chicken meat relative to controls. Also, no visible changes of the exterior appearance of the foodstuff after APDT were observed (Tortik, Spaeth and Plaetzer, 2014).

APDT using phenothiazinium photosensitisers, porphyrins, and xanthenes have also been tested against foodborne pathogens and microbial contaminants post-harvest. The spore-forming bacterium *Alicyclobacillus acidoterrestris* can cause great losses to fruit juice industries due to its thermal and chemical resistance and spoilage potential. Phototreatment with new methylene blue or tetracationic porphyrin combined with white light inactivated the spores both in suspension (PBS and orange juice) and on orange peels. Reductions in viability reached up to 7.3 log<sub>10</sub> in suspensions and 2.8 log<sub>10</sub> on peels. The presence of potassium iodide increased the effect of APDT (do Prado-Silva *et al.*, 2020).

APDT with cationic porphyrins were also used to photoinactivate the Gram-negative phytopathogenic bacterium *Pseudomonas syringae* pv. *actinidiae* both *in vitro* and in kiwifruit plants under solar radiation. Photoinactivation reached up to 7.4 log<sub>10</sub> *in vitro* and 6.2 log<sub>10</sub> on leaves (Martins *et al.*, 2018). Also, APDT with eosin Y combined with green light was evaluated against the pathogenic bacteria *S. enterica*, *E. coli*, *P. aeruginosa*, *S. aureus*, and *B. cereus*. *P. aeruginosa* was completely inactivated at 10 µmol L<sup>-1</sup>, and reductions in viability reached 2.7 and 1.7 for *B. cereus* and *S. enterica*, respectively. *E. coli* viability was slightly reduced while *S. aureus* was the most susceptible, being completely inactivated by eosin at 5 µmol L<sup>-1</sup> and 5 min of light exposure (Bonin *et al.*, 2018).

#### 4. Fungal tolerance to photoantimicrobials

Plants that produce furocoumarins and other potent photosensitizers can still experience severe microbial infections, and this is due to the development of tolerance or resistance to photosensitization in some specialized pathogens. *Fusarium sambucinum*, as well as some other plant pathogens, can metabolize, and thereby detoxify, xenobiotics such as furocoumarins. *F. sambucinum* is cosmopolitan in terms of habitat and lifestyle (both soil saprophyte and plant-pathogen), but is not generally regarded as any more stress tolerant than comparable species of fungi (Cray *et al.*, 2016). Circumstantial evidence from ecophysiological/toxin-resistance studies suggests that individual strains may preferentially inhabit either soils or the plant host (Desjardins, Spencer and Plattner, 1989). *F. sambucinum* tolerance to the furocoumarin xanthotoxin has been tested *in vitro* for 62 strains obtained from soils and diseased plants. As all the experiments were conducted in the dark, only direct inhibition by compounds was evaluated. Twenty-one out of 24 *F. sambucinum* strains isolated from plants and only two out of 38 strains isolated from soil were found to be highly tolerant to xanthotoxin. Of 16 *F. sambucinum* strains tested against 16 furocoumarin precursors and furocoumarins, all those that had been isolated from plants were highly tolerant and, in most cases, completely able to metabolize all of the compounds assayed.


Conversely, most of the soil-derived strains tested were tolerant to furanocoumarin precursors but sensitive to certain furocoumarins (Desjardins, Spencer and Plattner, 1989).

Metabolic inactivation of phototoxic plant metabolites is not the only mechanism that plant pathogens use to protect themselves against host-induced photosensitization. For instance, conidia of some fungi contain high concentration of pigments, such as carotenoids, that scavenge reactive species and can mitigate stresses such as those induced by solar UV radiation and photodynamic processes (Blanc, Tuveson and Sargent, 1976; Braga *et al.*, 2015; Thomas, Sargent and Tuveson, 1981; Shimizu, Egashira and Takahama, 1979). Pigments such as carotenoids are usually located within the plasma membrane and are able to quench both singlet oxygen and other types of reactive species produced by photosensitizers (Thomas, Sargent and Tuveson, 1981; Shimizu, Egashira and Takahama, 1979; Will, Newland and Reppe, 1984). Other pigments, which are structural components of the conidial cell wall (including melanins and melanin-like compounds), can selectively absorb solar radiation at the wavelengths required to activate the plant's photosensitizers (Asthana and Tuveson, 1992). The wild-type conidia of several *Colletotrichum*, *Fusarium*, and *Neurospora* species that are yellowish accumulate carotenoids, while *Alternaria*, *Aspergillus*, *Metarhizium*, and *Penicillium* conidia that are dark colored (brown-, gray- green- or bluish) are characterized by high levels of melanins or melanin-like pigments within the cell wall (Asthana and Tuveson, 1992; Gonzales *et al.*, 2010).

The above discussion contains examples of plants which produce photosensitizers that act as photoantimicrobials, protecting them against infections by pathogenic microbes. However, in plant-microbe interactions, there are also cases in which the opposite occurs. Plant-pathogenic fungi of the genera *Alternaria*, *Cladosporium*, *Elsinoë* and *Mycosphaerella* produce perylenequinone pigments during host infection, such as cercosporin, elsinochromes, hypocrellins, calphostin, and rubellin which are potent photosensitizers and damage the plant (Chung, 2011; Daub, Herrero and Chung, 2005; Daub, Herrero and Chung, 2013; Heiser, Sachs and Liebermann, 2003; Świdarska-Burek *et al.*, 2020; Thomas *et al.*, 2020). Most of the



characterized perylenequinones produced by fungi share a common 4,9-dihydroxy-3,10-perylenequinone core and differ in side chain composition (Daub, Herrero and Chung, 2013). These pigments are very potent  $^1\text{O}_2$ -generating photosensitizers that have a crucial role in the establishment of pathogenic association between fungi and their plant host (Chung, 2011; Daub, Herrero and Chung, 2013). The production of these photosensitizers during infection causes lipid peroxidation and damage to the plasma membrane of the host cells, leading to leakage of nutrients into the intercellular spaces colonized by the pathogen (Fig. 9) (Daub, 1982; Daub and Briggs, 1983; Daub, Herrero and Chung, 2013).




**Figure 9** – Schematic mechanism depicting the mode-of-action of phytopathogenic fungi that use photosensitizers for pathogenesis. After penetration through stomata, the fungus releases a photosensitizing molecule in the intercellular space. Activation of this molecule by light results in the production of reactive oxygen species that damage lipids at the cell membrane, releasing plant nutrients

into the medium and allowing sustained fungal growth. The depicted mechanism is based on that of *Cercospora* fungi.

Interestingly, the producing fungi are resistant to these photoactivated toxins (Daub, Herrero and Chung, 2013). Among these fungi, those of the genus *Cercospora* are the best studied (Daub, Herrero and Chung, 2013; Świdarska-Burek *et al.*, 2020). *Cercospora* species cause devastating leaf-blighting disease on a wide range of important plant host species worldwide (Beseli *et al.*, 2015). Several species of the genus *Cercospora* produce large quantities of the phototoxin cercosporin, a lipid-soluble 4,9-hydroxyperylene-3,10-quinone derivative, which absorbs light and reacts with oxygen, generating reactive oxygen species, mostly  $^1\text{O}_2$  (Daub and Hangarter, 1983). Unlike free-radical forms of ROS against which resistance mechanism are well characterized, the cellular bases of  $^1\text{O}_2$  resistance are still being elucidated (Beseli, Noar and Daub, 2015; Daub, Herrero and Chung, 2013; Thomas *et al.*, 2020). Light is required, not only for cercosporin activation, but also for cercosporin production (Ehrenshaft and Upchurch, 1991). *Cercospora* species can, under light, produce and thrive in concentrations of cercosporin up to 1000-fold higher than that which is lethal to other organisms (Ehrenshaft *et al.*, 1998). These fungi are highly tolerant not only to cercosporin but also to a broad range of structurally unrelated  $^1\text{O}_2$ -generating photosensitizers, including porphyrins, xanthenes, and phenothiazinium dyes (Ehrenshaft, Jenns and Daub, 1995). Some fungi other than *Cercospora* species, such as *Alternaria solani*, *Cladosporium cucumerinum*, *Cladosporium fulvum*, *Colletotrichum lagenarium*, *Verticillium* sp., *S. cerevisiae* and *Sporobolomyces* sp. are also highly resistant to cercosporin and other  $^1\text{O}_2$ -generating photosensitizers (Daub, 1987). Due to their high intrinsic tolerance, these fungi are excellent models for the elucidation of molecular and genetic bases of resistance to  $^1\text{O}_2$ -generating photosensitizers.

Much of what is known about the tolerance of *Cercospora* to cercosporin and to other  $^1\text{O}_2$ -generating photosensitizers came from Daub's group and some of their reviews are

recommended for an in-depth approach on this subject (Daub and Ehrenshaft, 2000; Daub, Herrero and Chung, 2005; Daub, Herrero and Chung, 2013; Świdarska-Burek *et al.*, 2020). *Cercospora* auto-resistance to light-activated cercosporin is a complex and yet not completely understood characteristic mediated by multiple mechanisms including the reversible reduction and detoxification of the cercosporin inside the fungal cells, the production of  $^1\text{O}_2$  quenchers, and the transport of the toxin out of the cells (Fig. 10) (Beseli *et al.*, 2015; Daub *et al.*, 1992; Daub, Herrero and Chung, 2013).



1158

**Figure 10** – Examples of mechanisms via which cells may be either tolerant or resistant to photodynamic inactivation. These mechanisms include: (I) inactivation of photosensitizers via redox reactions; (II) scavenging of reactive oxygen species by specialized molecules (e.g., carotenoids, vitamin B6); (III) detoxification of reactive oxygen species by antioxidant enzymes (e.g., superoxide dismutase); and (IV) efflux of photosensitizers from the intracellular medium by transporters. These mechanisms are

based on current knowledge on tolerance/resistance to APDT on *Cercospora* fungi, but they are potentially present in all microorganisms.

Carotenoids are the most efficient quenchers of  $^1\text{O}_2$  identified in biological systems, and are able to quench not only  $^1\text{O}_2$ , but also the activated triplet state of  $^1\text{O}_2$ -generating photosensitizers (Daub and Payne, 1989; Daub, Herrero and Chung, 2005; Kruk and Szymańska, 2021). Carotenoids were also the first endogenous compounds identified that can protect fungi against cercosporin-based APDT (Daub and Payne, 1989). Carotenoid-deficient mutants of *Neurospora crassa* are significantly more sensitive to cercosporin than the carotenoid-producing wild-type isolates (Blanc, Tuveson and Sargent, 1976). Carotenoids also protect fungi against other  $^1\text{O}_2$ -generating photosensitizers that damage the plasma membrane. Conidia of *N. crassa* and mycelia of *Ustilago violaceae* mutants lacking carotenoids are less tolerant to methylene blue- and toluidine blue O-based APDT than the wild-type strains (Thomas, Sargent and Tuveson, 1981; Will, Newland and Reppe, 1984). However, experiments performed with *Cercospora nicotianae* showed a different and unexpected result (Ehrenshaft, Jenns and Daub, 1995). Target gene disruption was used to create carotenoid-minus mutants of a wild-type and two cercosporin-sensitive *C. nicotianae* mutants. These carotenoids-deficient mutants had similar sensitivity to either photoactivated cercosporin or five others  $^1\text{O}_2$ -generating photosensitizers (rose bengal, hematoporphyrin, methylene blue, toluidine blue O, eosin Y) compared to the parent strains. Together, these data suggested that carotenoids are important to fungal tolerance to photoactivated cercosporin but they are not involved or have only a minor effect on *Cercospora* resistance to  $^1\text{O}_2$ -generating photosensitizers (Daub and Payne, 1989).

The term vitamin B<sub>6</sub> is used to describe all biologically interconvertible forms of pyridoxine (Bilski *et al.*, 2000). Vitamin B<sub>6</sub> and its derivatives are good  $^1\text{O}_2$  quenchers; also, the pyridoxine moiety can function as a redox quencher for excited cercosporin by forming the cercosporin radical anion (Bilski *et al.*, 2000). Mutants of *C. nicotianae* deficient in biosynthesis of vitamin B<sub>6</sub> are highly sensitive to cercosporin and other  $^1\text{O}_2$ -generating photosensitizers, such as

hematoporphyrin, rose bengal, eosin Y, methylene blue, and toluidine blue O (Ehrenshaft *et al.*, 1998; Ehrenshaft *et al.*, 1999b; Ehrenshaft *et al.*, 1999a). Analysis of cellular levels of vitamin B<sub>6</sub> also showed that *C. nicotianae* has 2- to 3-fold higher levels of B<sub>6</sub> vitamers than the cercosporin-sensitive fungi *A. flavus* and *N. crassa* (Herrero and Daub, 2007).

In contrast to <sup>1</sup>O<sub>2</sub> quenchers, there is little evidence for a role of antioxidant enzymes in cercosporin resistance in *Cercospora*. *C. nicotianae* does not have higher levels of superoxide dismutase (SOD), catalase, or peroxidase activities as compared to cercosporin-based APDT sensitive fungi and there is also no difference in overall antioxidant activity (Daub, 1987).

The ability to transport cercosporin and other photosensitizers out of the cell also contributes to fungal resistance to these compounds (Beseli *et al.*, 2015; Daub, Herrero and Chung, 2013). Both the Major Facilitator Superfamily (MFS) and ATP-binding cassette (ABC) family of transporters are able to transport cercosporin out of *Cercospora* cells and provide partial resistance against cercosporin-based APDT (Beseli *et al.*, 2015). Targeted disruption of the gene for CFP (*Cercosporin Facilitator Protein*), an MFS transporter, in the soybean pathogen *Cercospora kikuchii* drastically reduced the production of cercosporin, greatly impaired virulence of the fungus, and increased sensitivity to exogenous cercosporin in comparison to the wild-type strain (Callahan *et al.*, 1999). Also, the transgenic expression of *CFP* gene in the cercosporin-sensitive fungus *Cochiobolus heterostrophus* resulted in increased tolerance to cercosporin due to its export out of the fungus (Upchurch *et al.*, 2002).

The importance of MFS transporters to the resistance to <sup>1</sup>O<sub>2</sub>-generating photosensitizers was also observed in other fungal pathogens. *Bcmfs1*, an MFS transporter from *B. cinerea*, provides tolerance to cercosporin-based APDT (Hayashi, Schoonbeek and De Waard, 2002). Deletion mutants showed increased sensitivity to photoactivated cercosporin, while overexpression mutants displayed decreased sensitivity (Hayashi, Schoonbeek and De Waard, 2002). Mutants of the citrus pathogen *A. alternata* lacking the *AaMFS19* gene, which encodes an

MFS transporter, display profound sensitivity to the  $^1\text{O}_2$ -generating photosensitizers eosin Y, rose bengal, hematoporphyrin, methylene blue, and cercosporin (Chen *et al.*, 2017).

The importance of ABC transporters to cercosporin resistance was demonstrated in *C. nicotianae*. Mutants with disruption of *ATR1*, which is an ABC transporter gene, had dramatic reductions in cercosporin production and also showed moderately higher sensitivity to cercosporin indicating that *ATR1* acts as a cercosporin efflux pump and has a partial role in cercosporin-based APDT resistance (Amnuaykanjanasin and Daub, 2009). *CnATR2*, another ABC transporter involved in partial resistance to cercosporin, was recently characterized. Transformation and expression of *CnATR2* in the cercosporin-sensitive fungus *N. crassa* significantly increased cercosporin resistance. However, target gene disruption of *CnATR2* in the wild type *C. nicotianae* did not decrease resistance. The overexpression of the gene that codes for Snq2p, a well-characterized multidrug, ABC-type, efflux protein conferred resistance to cercosporin and to other  $^1\text{O}_2$ -generating photosensitizers such as methylene blue and toluidine blue O to an otherwise sensitive *S. cerevisiae* strain (Ververidis *et al.*, 2001). In contrast, the *snq2* null mutant was not more sensitive to methylene blue and toluidine blue O than a wild-type control strain (Ververidis *et al.*, 2001).

Studies with redox-sensitive dyes, reducing agents, and with detection of reduced and oxidized forms of cercosporin indicated that the most relevant mechanism responsible for *Cercospora*'s resistance to cercosporin is the ability of these fungi to maintain cercosporin within the hyphae in a reduced form (Fig. 10) (Daub *et al.*, 1992; Daub and Ehrenshaft, 2000; Daub, Herrero and Chung, 2005; Daub, Herrero and Chung, 2013; Jenns and Daub, 1995; Świdarska-Burek *et al.*, 2020). Reduced cercosporin is a poor generator of  $^1\text{O}_2$ , particularly in aqueous solution (Leisman and Daub, 1992). As the reduced form is labile, and readily reoxidizes on aeration or removal from the reducing agents, cercosporin that diffuses away from the fungal cell spontaneously reoxidizes to the photoactive form needed for the infection of the host plant (Daub *et al.*, 1992; Leisman and Daub, 1992).

Other fungal species were also tested for cercosporin-reducing ability (Daub *et al.*, 1992). *A. alternata*, which is cercosporin-resistant, was able to reduce cercosporin but *A. flavus* and *N. crassa*, which are cercosporin-sensitive, had only limited ability. Evidence suggested that the reduction of the photosensitizer may be a generalized mechanism of resistance for *Cercospora* to other  $^1\text{O}_2$ -generating photosensitizers (Daub *et al.*, 1992). *Cercospora* species were also resistant to eosin Y- but not to rose bengal-based APDT. Microscopic observation showed that *Cercospora* species were not capable of reducing rose bengal but were capable of reducing eosin Y. The reduction of the photosensitizer as a protective mechanism was also observed in other fungal species. The over-expression of the gene *CPD1* (Cercosporin Photosensitizer Detoxification) that codes for a putative plasma membrane-associated reductase conferred resistance to cercosporin, methylene blue and toluidine blue O in *S. cerevisiae* (Ververidis *et al.*, 2001).

It is often stated that, due to its multiple-target mode of action, the selection of fungi displaying resistance to APDT is unlikely. However, the existence of several fungal species that are intrinsically resistant to APDT with  $^1\text{O}_2$ -generating photosensitizer, indicates that the possibility of the emergence of tolerance to APDT in fungal species of medical or agricultural importance deserves more attention. A recent study performed by da Cruz and coworkers has showed that *C. abscisum* submitted to successive cycles of APDT can become more tolerant to the treatment, although the decrease in susceptibility was small. Also, the study revealed that the more-tolerant strain also accumulated higher amounts of carotenoids (da Cruz *et al.*, in preparation). Unfortunately, other studies performing long-term experimentation with filamentous fungi to determine whether or not it is possible to select resistant strains during successive cycles of APDT are still lacking. There is no doubt that the understanding of the mechanisms responsible for the intrinsic resistance to APDT of some fungal species, such as of the genus *Cercospora*, will be important to understand and eventually anticipate a possible emergence of resistance to APDT in species of medical and agricultural importance, which may occur with the expansion of its use.

The discussion above may seem heavily based on Type II reactions, i.e., those producing  $^1\text{O}_2$ . However, some of the mechanisms presented (Fig. 10) can also operate to avoid cell damage arising from Type I reactions. For instance, the redox processes that render photosensitizers not reactive to light would also prevent Type I reactions from occurring. Similarly, efflux pumps that prevent photosensitizers from being light activated inside cells do not discriminate between Type I and Type II photosensitizers.

The same cannot be said about the other two mechanisms, i.e., detoxification and scavenging. Cells have known lines of defense against Type I ROS –such as superoxide anion radical and hydrogen peroxide– in the form of the enzymes superoxide dismutase, catalase, and glutathione reductase, as well as specific scavengers for these reactive species. Both the expression of the enzymes and the production/accumulation of scavengers can be modulated to achieve increased tolerance to photosensitizers operating via Type I reactions. On the other hand, there is no known first line of defense against  $^1\text{O}_2$ , so an effective antioxidant system may prevent exclusively against Type I photosensitizers. Evidence of this can be found on extremophilic microorganisms. The bacterium *Deinococcus radiodurans* has a remarkable antioxidant system that effectively protects the proteome from ionizing radiation, desiccation, and oxidative stresses at high levels (Qi *et al.*, 2020), making the microbe very tolerant to Type I ROS. Nonetheless, *D. radiodurans* cannot sustain the damages imposed by  $^1\text{O}_2$ -producing photosensitizers, with at least one report showing that its tolerance falls below that of *E. coli* (Nitzan and Ashkenazi, 1999; Schafer, Schmitz and Horneck, 1998).

At present, it is not possible to say whether the tolerance mechanisms to Type I and to Type II photosensitizers would emerge and occur concomitantly (if at all). However, if tolerance to APDT is a multifactorial process operating at many levels, then alternating chemically-diverse photosensitizers that operate via different reactions will provide some protection against the emergence of tolerance.



## 5. Conclusions and unresolved questions

Plants, as do microbes, produce a mechanistically (and chemically) diverse array of antimicrobials which can vary with species, habitat, and environmental conditions (Cray *et al.*, 2015a; Lievens *et al.*, 2015; Oren and Hallsworth, 2014; Wecke and Mascher, 2011; Suryawanshi *et al.*, 2015). Of these, photosensitizers can have elegant mechanisms and are some of the most potent antimicrobials; yet, they are relatively undervalued in relation to their biotechnological potential.

The above discussion covers the ecologies of naturally-occurring, photodynamic processes including the light-activated antimicrobial activities of some plant metabolites, and the intriguing use of the photodynamic process by some plant-pathogenic fungi as an important virulence factor. The use of natural and synthetic photosensitizers to kill plant-pathogenic fungi and foodborne pathogens were also reviewed and discussed. The inhibitory mechanisms of both natural and synthetic light-activated substances were covered in the contexts of microbial stress biology and agricultural biotechnology. Implications were also made in relation to treatment of clinical infections caused by opportunistic fungi pathogens, once considered only plant pathogens and/or saprotrophic.

The development of conventional pesticides is a complex, costly, and time-consuming process that can be divided into three main steps: (i) research on the synthesis and screening of molecules, (ii) product development; and (iii) registration. The research evaluates the biological, chemical, toxicological, environmental, and commercial characteristics of candidate molecules to be registered. The development includes several processes, such as optimization of formulation, assessment of products in field trials against different biological targets in a variety of crops, and evaluation of toxicological and environmental impacts. Finally, product data are submitted to different regulatory agencies, which may agree or disagree with the registration and commercialization (McDougall, 2016). Therefore, as in the development of conventional

pesticides, photosensitizers may be submitted to similar processes before large-scale use in agriculture. In addition, the average time spent by a company to develop a conventional pesticide is approximately 11 years and the cost is about US\$ 286 million (McDougall, 2016). Currently, the cost for obtaining a photosensitizer-based product is unknown, and future research on economic feasibility is needed. Furthermore, there is still a considerable knowledge gap due to numerous unresolved questions. For example, although it is well established that APDT with most of the photosensitizers tested is able to kill, to a greater or lesser extent, most of the different species of fungi *in vitro*, little is known about the efficacy of the treatment in the field on different crops. An important issue that needs attention is the negative effect of shadowing, which may be caused by the plant canopy and/or by an extensive cloud cover lasting many days. Similarly, little is known about the side effects of the different photosensitizers on the host plants and in the environment. Also, formulations containing photosensitizers will have to be developed and approved for use in the field.

Despite its great potential, it seems that the development and use of APDT in agriculture has been delayed by the fact that this antimicrobial approach is unknown to the majority of agricultural professionals and by the apparent lack of interest by the chemical and pharmaceutical industries in the development of photosensitizer-based products for agricultural use. This parallels a similar lack of interest from these industries in the development of clinical anti-infectives based on this approach for both human and veterinary application. Here we showed that the use of photoantimicrobials is a viable and needed alternative to control plant- and foodborne pathogens, and has the potential to contribute to improving global food security.

## Acknowledgments

This work was supported by grants 2012/15204-8 and 2016/11386-5 from the State of São Paulo Research Foundation (FAPESP) to GULB. We also thank the National Council for Scientific

and Technological Development for research grants to GULB (425998/2018-5 and 307738/2018-3) and to GJSJ (308202/2017-1 and 316910/2021-0), and for a post-doctoral fellowship (165191/2020-1) to GTPB.

## REFERENCES

- Abrahamse, H. and Hamblin, M. R. (2016) 'New photosensitizers for photodynamic therapy', *Biochemical Journal*, 473(4), pp. 347-364.
- Afrasiabi, S., Pourhajibagher, M., Chiniforush, N., Aminian, M. and Bahador, A. (2020) 'Anti-biofilm and anti-metabolic effects of antimicrobial photodynamic therapy using chlorophyllin-phycoerythrin mixture against *Streptococcus mutans* in experimental biofilm caries model on enamel slabs', *Photodiagnosis and photodynamic therapy*, 29, pp. 101620.
- Agrios, G. N. (2005) *Plant pathology*. Elsevier.
- Al-Asmari, F., Mereddy, R. and Sultanbawa, Y. (2017) 'A novel photosensitization treatment for the inactivation of fungal spores and cells mediated by curcumin', *Journal of Photochemistry and Photobiology B: Biology*, 173, pp. 301-306.
- Al-Rubeai, M. A. and El-Hassi, M. (1986) 'Inactivation of wild type and mutant *Aspergillus nidulans* conidia by far-UV, near-UV, visible and sun lights', *Environmental and experimental botany*, 26(3), pp. 243-252.
- Ambrosini, V., Issawi, M., Sol, V. and Riou, C. (2020) 'Photodynamic inactivation of *Botrytis cinerea* by an anionic porphyrin: an alternative pest management of grapevine', *Scientific Reports*, 10(1), pp. 1-12.
- Amnuaykanjanasin, A. and Daub, M. E. (2009) 'The ABC transporter ATR1 is necessary for efflux of the toxin cercosporin in the fungus *Cercospora nicotianae*', *Fungal genetics and biology*, 46(2), pp. 146-158.
- Andrade, G. C., Brancini, G. T. P., Abe, F. R., de Oliveira, D. P., Nicolella, H. D., Tavares, D. C., Micas, A. F. D., Savazzi, E. A., Silva-Junior, G. J. and Wainwright, M. (2022) 'Phenothiazinium dyes for photodynamic treatment present lower environmental risk compared to a formulation of trifloxystrobin and tebuconazole', *Journal of Photochemistry and Photobiology B: Biology*, 226, pp. 112365.
- Andrivon, D., Ramage, K., Guerin, C., Lucas, J. and Jouan, B. (1997) 'Distribution and fungicide sensitivity of *Colletotrichum coccodes* in French potato-producing areas', *Plant Pathology*, 46(5), pp. 722-728.
- Aponiene, K., Paskeviciute, E., Reklaitis, I. and Luksiene, Z. (2015) 'Reduction of microbial contamination of fruits and vegetables by hypericin-based photosensitization: Comparison with other emerging antimicrobial treatments', *Journal of Food Engineering*, 144, pp. 29-35.

- 1377 Araújo, C. A. S., Ferreira, P. C., Pupin, B., Dias, L. P., Avalos, J., Edwards, J., Hallsworth, J. E.  
1378 and Rangel, D. E. N. (2020) 'Osmotolerance as a determinant of microbial ecology: A study of  
1379 phylogenetically diverse fungi', *Fungal Biology*, 124(5), pp. 273-288.
- 1380 Arboleda, A., Miller, D., Cabot, F., Taneja, M., Aguilar, M. C., Alawa, K., Amescua, G., Yoo, S. H.  
1381 and Parel, J.-M. (2014) 'Assessment of rose bengal versus riboflavin photodynamic therapy for inhibition  
1382 of fungal keratitis isolates', *American journal of ophthalmology*, 158(1), pp. 64-70. e2.
- 1383 Arnason, J., Bourque, G., Madhosilngh, C. and Orr, W. (1986) 'Disruption of membrane functions  
1384 in *Fusarium culmorum* by an acetylenic allelochemical', *Biochemical systematics and ecology*, 14(6), pp.  
1385 569-574.
- 1386 Asthana, A., McCloud, E. S., Berenbaum, M. R. and Tuveson, R. W. (1993) 'Phototoxicity of Citrus  
1387 jambhiri to fungi under enhanced UV-B radiation: Role of furanocoumarins', *Journal of chemical ecology*,  
1388 19(12), pp. 2813-2830.
- 1389 Asthana, A. and Tuveson, R. (1992) 'Effects of UV and phototoxins on selected fungal pathogens  
1390 of citrus', *International journal of plant sciences*, 153(3, Part 1), pp. 442-452.
- 1391 Aver'yanov, A., Lapikova, V., Pasechnik, T., Zakharenkova, T., Pogosyan, S. and Baker, C. (2011)  
1392 'Suppression of cucurbit scab on cucumber leaves by photodynamic dyes', *Crop Protection*, 30(7), pp.  
1393 925-930.
- 1394 Ball, P. and Hallsworth, J. E. (2015) 'Water structure and chaotropicity: their uses, abuses and  
1395 biological implications', *Physical Chemistry Chemical Physics*, 17(13), pp. 8297-8305.
- 1396 Barros, B. H. R., da Silva, S. H., Marques, E. d. R., Rosa, J. C., Yatsuda, A. P., Roberts, D. W. and  
1397 Braga, G. U. L. (2010) 'A proteomic approach to identifying proteins differentially expressed in conidia and  
1398 mycelium of the entomopathogenic fungus *Metarhizium acridum*', *Fungal Biology*, 114(7), pp. 572-579.
- 1399 Bartlett, D. W., Clough, J. M., Godwin, J. R., Hall, A. A., Hamer, M. and Parr-Dobrzanski, B. (2002)  
1400 'The strobilurin fungicides', *Pest Management Science: formerly Pesticide Science*, 58(7), pp. 649-662.
- 1401 Ben, H., Huo, J., Yao, Y., Gao, W., Wang, W., Hao, Y. and Zhang, X. (2021) 'First report of  
1402 *Colletotrichum capsici* causing Anthracnose on *Alocasia macrorrhizos* in China', *Plant Disease*, 105(4),  
1403 pp. 1203-1203.
- 1404 Berenbaum, M. and Larson, R. (1988) 'Flux of singlet oxygen from leaves of phototoxic plants',  
1405 *Experientia*, 44(11), pp. 1030-1032.
- 1406 Beseli, A., Amnuaykanjanasin, A., Herrero, S., Thomas, E. and Daub, M. E. (2015) 'Membrane  
1407 transporters in self resistance of *Cercospora nicotianae* to the photoactivated toxin cercosporin', *Current*  
1408 *Genetics*, 61(4), pp. 601-620.
- 1409 Beseli, A., Noar, R. and Daub, M. E. (2015) 'Characterization of *Cercospora nicotianae*  
1410 Hypothetical Proteins in Cercosporin Resistance', *PLOS ONE*, 10(10), pp. e0140676.
- 1411 Bhaganna, P., Volkers, R. J., Bell, A. N., Kluge, K., Timson, D. J., McGrath, J. W., Ruijsenaars,  
1412 H. J. and Hallsworth, J. E. (2010) 'Hydrophobic substances induce water stress in microbial cells', *Microbial*  
1413 *Biotechnology*, 3(6), pp. 701-716.

- 1414 Bilski, P., Li, M., Ehrenshaft, M., Daub, M. and Chignell, C. (2000) 'Vitamin B6 (pyridoxine) and its  
1415 derivatives are efficient singlet oxygen quenchers and potential fungal antioxidants', *Photochemistry and*  
1416 *photobiology*, 71(2), pp. 129-134.
- 1417 Binns, S., Purgina, B., Bergeron, C., Smith, M., Ball, L., Baum, B. and Arnason, J. (2000) 'Light-  
1418 mediated antifungal activity of Echinacea extracts', *Planta medica*, 66(03), pp. 241-244.
- 1419 Bintsis, T., Litopoulou-Tzanetaki, E. and Robinson, R. K. (2000) 'Existing and potential applications  
1420 of ultraviolet light in the food industry—a critical review', *Journal of the Science of Food and Agriculture*,  
1421 80(6), pp. 637-645.
- 1422 Bispo, M., Suhani, S. and van Dijk, J. M. (2021) 'Empowering antimicrobial photodynamic therapy  
1423 of Staphylococcus aureus infections with potassium iodide', *Journal of Photochemistry and Photobiology*  
1424 *B: Biology*, 225, pp. 112334.
- 1425 Blanc, P., Tuveson, R. and Sargent, M. (1976) 'Inactivation of carotenoid-producing and albino  
1426 strains of Neurospora crassa by visible light, blacklight, and ultraviolet radiation', *Journal of bacteriology*,  
1427 125(2), pp. 616-625.
- 1428 Bocking, T., Barrow, K. D., Netting, A. G., Chilcott, T. C., Coster, H. G. and Hofer, M. (2000) 'Effects  
1429 of singlet oxygen on membrane sterols in the yeast Saccharomyces cerevisiae', *Eur J Biochem*, 267(6),  
1430 pp. 1607-18.
- 1431 Bogucka-Kocka, A. and Krzaczek, T. (2003) 'The furanocoumarins in the roots of Heracleum  
1432 sibiricum L', *Acta Poloniae Pharmaceutica*, 60(5), pp. 391-393.
- 1433 Bonin, E., dos Santos, A. R., Fiori da Silva, A., Ribeiro, L. H., Favero, M. E., Campanerut-Sá, P. A.  
1434 Z., de Freitas, C. F., Caetano, W., Hioka, N. and Mikcha, J. M. G. (2018) 'Photodynamic inactivation of  
1435 foodborne bacteria by eosin Y', *Journal of Applied Microbiology*, 124(6), pp. 1617-1628.
- 1436 Bordin, F., Carlassare, F., Baccichetti, F. and Anselmo, L. (1976) 'DNA repair and recovery in  
1437 Escherichia coli after psoralen and angelicin photosensitization', *Biochimica et Biophysica Acta (BBA)-*  
1438 *Nucleic Acids and Protein Synthesis*, 447(3), pp. 249-259.
- 1439 Bourque, G., Arnason, J., Madhosingh, C. and Orr, W. (1985) 'The photosensitization of the plant  
1440 pathogen Fusarium culmorum by phenylheptatriyne from Bidens pilosa', *Canadian journal of botany*, 63(5),  
1441 pp. 899-902.
- 1442 Braga, G. U., Destéfano, R. H. and Messias, C. L. (1999) 'Oxygen consumption by Metarhizium  
1443 anisopliae during germination and growth on different carbon sources', *Journal of invertebrate pathology*,  
1444 74(2), pp. 112-119.
- 1445 Braga, G. U., Rangel, D. E., Fernandes, E. K., Flint, S. D. and Roberts, D. W. (2015) 'Molecular  
1446 and physiological effects of environmental UV radiation on fungal conidia', *Curr Genet*, 61(3), pp. 405-25.
- 1447 Brancini, G. T., Rodrigues, G. B., Rambaldi, M. S., Izumi, C., Yatsuda, A. P., Wainwright, M., Rosa,  
1448 J. C. and Braga, G. U. (2016) 'The effects of photodynamic treatment with new methylene blue N on the  
1449 Candida albicans proteome', *Photochem Photobiol Sci*, 15(12), pp. 1503-1513.

- 1450 Brancini, G. T. P., Hallsworth, J. E., Corrochano, L. M. and Braga, G. Ú. L. (2022) 'Photobiology of  
1451 the keystone genus *Metarhizium*', *Journal of Photochemistry and Photobiology B: Biology*, 226, pp.  
1452 112374.
- 1453 Buchovec, I., Lukseviciute, V., Marsalka, A., Reklaitis, I. and Luksiene, Z. (2016) 'Effective  
1454 photosensitization-based inactivation of Gram (–) food pathogens and molds using the chlorophyllin—  
1455 chitosan complex: towards photoactive edible coatings to preserve strawberries', *Photochemical &  
1456 Photobiological Sciences*, 15(4), pp. 506-516.
- 1457 Buchovec, I., Lukseviciūtė, V., Kokštaite, R., Labeikyte, D., Kaziukonyte, L. and Luksiene, Z. (2017)  
1458 'Inactivation of Gram (–) bacteria *Salmonella enterica* by chlorophyllin-based photosensitization:  
1459 Mechanism of action and new strategies to enhance the inactivation efficiency', *Journal of Photochemistry  
1460 and Photobiology B: Biology*, 172, pp. 1-10.
- 1461 Buchovec, I. and Lukšienė, Ž. (2015) 'Novel approach to control microbial contamination of  
1462 germinated wheat sprouts: photoactivated chlorophyllin-chitosan complex', *International journal of food  
1463 processing technology*, 2, pp. 26-30.
- 1464 Buchovec, I., Paskeviciute, E. and Luksiene, Z. (2010) 'Photosensitization-based inactivation of  
1465 food pathogen *Listeria monocytogenes* in vitro and on the surface of packaging material', *Journal of  
1466 Photochemistry and Photobiology B: Biology*, 99(1), pp. 9-14.
- 1467 Buchovec, I., Vaitonis, Z. and Luksiene, Z. (2009) 'Novel approach to control *Salmonella enterica*  
1468 by modern biophotonic technology: photosensitization', *Journal of applied microbiology*, 106(3), pp. 748-  
1469 754.
- 1470 Bäumlér, W., Regensburger, J., Knak, A., Felgentraeger, A. and Maisch, T. (2012) 'UVA and  
1471 endogenous photosensitizers—the detection of singlet oxygen by its luminescence', *Photochemical &  
1472 Photobiological Sciences*, 11(1), pp. 107-117.
- 1473 Callahan, T. M., Rose, M. S., Meade, M. J., Ehrenshaft, M. and Upchurch, R. G. (1999) 'CFP, the  
1474 putative cercosporin transporter of *Cercospora kikuchii*, is required for wild type cercosporin production,  
1475 resistance, and virulence on soybean', *Molecular plant-microbe interactions*, 12(10), pp. 901-910.
- 1476 Calzavara-Pinton, P., Rossi, M. T., Sala, R. and Venturini, M. (2012) 'Photodynamic antifungal  
1477 chemotherapy', *Photochemistry and photobiology*, 88(3), pp. 512-522.
- 1478 Cardoso, D. R., Libardi, S. H. and Skibsted, L. H. (2012) 'Riboflavin as a photosensitizer. Effects  
1479 on human health and food quality', *Food & Function*, 3(5), pp. 487-502.
- 1480 Castro, K. A. D. F., Brancini, G. T. P., Costa, L. D., Biazotto, J. C., Faustino, M. A. F., Tomé, A.  
1481 C., Neves, M. G. P. M. S., Almeida, A., Hamblin, M. R., da Silva, R. S. and Braga, G. Ú. L. (2020) 'Efficient  
1482 photodynamic inactivation of *Candida albicans* by porphyrin and potassium iodide co-encapsulation in  
1483 micelles', *Photochemical & Photobiological Sciences*, 19(8), pp. 1063-1071.
- 1484 Chen, L.-H., Tsai, H.-C., Yu, P.-L. and Chung, K.-R. (2017) 'A major facilitator superfamily  
1485 transporter-mediated resistance to oxidative stress and fungicides requires Yap1, Skn7, and MAP kinases  
1486 in the citrus fungal pathogen *Alternaria alternata*', *PLoS One*, 12(1), pp. e0169103.

- Chin, J. P., Megaw, J., Magill, C. L., Nowotarski, K., Williams, J. P., Bhaganna, P., Linton, M., Patterson, M. F., Underwood, G. J. and Mswaka, A. Y. (2010) 'Solutes determine the temperature windows for microbial survival and growth', *Proceedings of the National Academy of Sciences*, 107(17), pp. 7835-7840.
- Chitolina, G. M., Silva-Junior, G. J., Feichtenberger, E., Pereira, R. G. and Amorim, L. (2021) 'Distribution of *Alternaria alternata* isolates with resistance to quinone outside inhibitor (QoI) fungicides in Brazilian orchards of tangerines and their hybrids', *Crop Protection*, 141, pp. 105493.
- Christensen, L. P. and Brandt, K. (2006) 'Bioactive polyacetylenes in food plants of the Apiaceae family: occurrence, bioactivity and analysis', *Journal of pharmaceutical and biomedical analysis*, 41(3), pp. 683-693.
- Chung, K. R. (2011) 'Elsinoë fawcettii and Elsinoë australis: the fungal pathogens causing citrus scab', *Molecular plant pathology*, 12(2), pp. 123-135.
- Ciampi-Guillardi, M., Muñoz, V. N. V., Silva-Junior, G. J. and Massola Júnior, N. S. (2022) 'Molecular detection and quantification of *Colletotrichum abscissum* in sweet orange propagative material', *Plant Pathology*, 71(3), pp. 634-643.
- Claessens, C. G., Hahn, U. and Torres, T. (2008) 'Phthalocyanines: From outstanding electronic properties to emerging applications', *The Chemical Record*, 8(2), pp. 75-97.
- Clsi (2017) 'Reference Method for Broth Dilution Antifungal Susceptibility Testing of Filamentous Fungi', *Clinical and Laboratory Standards Institute*, 3rd Edition.
- Collins, M. and Buick, R. (1989) 'Effect of temperature on the spoilage of stored peas by *Rhodotorula glutinis*', *Food Microbiology*, 6(3), pp. 135-141.
- Cray, J., Bhaganna, P., Singhal, R., Patil, S., Saha, D., Chakraborty, R., Iwaguchi, S., Timson, D. and Hallsworth, J. 'Chaotropic and hydrophobic stress mechanisms of antifungal substances'. *Modern fungicides and antifungal compounds VII. Proceedings of the 17th International Reinhardsbrunn Symposium, April 21-25 2013, Friedrichroda, Germany: Deutsche Phytomedizinische Gesellschaft eV Verlag*, 73-80.
- Cray, J. A., Bell, A. N., Bhaganna, P., Mswaka, A. Y., Timson, D. J. and Hallsworth, J. E. (2013a) 'The biology of habitat dominance; can microbes behave as weeds?', *Microbial biotechnology*, 6(5), pp. 453-492.
- Cray, J. A., Connor, M. C., Stevenson, A., Houghton, J. D., Rangel, D. E., Cooke, L. R. and Hallsworth, J. E. (2016) 'Biocontrol agents promote growth of potato pathogens, depending on environmental conditions', *Microbial biotechnology*, 9(3), pp. 330-354.
- Cray, J. A., Russell, J. T., Timson, D. J., Singhal, R. S. and Hallsworth, J. E. (2013b) 'A universal measure of chaotropicity and kosmotropicity', *Environmental Microbiology*, 15(1), pp. 287-296.
- Cray, J. A., Stevenson, A., Ball, P., Bankar, S. B., Eleutherio, E. C., Ezeji, T. C., Singhal, R. S., Thevelein, J. M., Timson, D. J. and Hallsworth, J. E. (2015a) 'Chaotropicity: a key factor in product tolerance of biofuel-producing microorganisms', *Current opinion in biotechnology*, 33, pp. 228-259.

- 1524 Cray, J. A., Stevenson, A., Ball, P., Bankar, S. B., Eleutherio, E. C. A., Ezeji, T. C., Singhal, R. S.,  
 1525 Thevelein, J. M., Timson, D. J. and Hallsworth, J. E. (2015b) 'Chaotropicity: a key factor in product  
 1526 tolerance of biofuel-producing microorganisms', *Current Opinion in Biotechnology*, 33, pp. 228-259.
- 1527 da Silva, R. N., Tome, A. C., Tome, J. P., Neves, M. G., Faustino, M. A., Cavaleiro, J. A., Oliveira,  
 1528 A., Almeida, A. and Cunha, A. (2012) 'Photo-inactivation of *Bacillus* endospores: Inter-specific variability  
 1529 of inactivation efficiency', *Microbiology and immunology*, 56(10), pp. 692-699.
- 1530 Dai, T., Bil de Arce, V. J., Tegos, G. P. and Hamblin, M. R. (2011) 'Blue dye and red light, a dynamic  
 1531 combination for prophylaxis and treatment of cutaneous *Candida albicans* infections in mice', *Antimicrobial  
 1532 agents and chemotherapy*, 55(12), pp. 5710-5717.
- 1533 Dall'Acqua, F. and Martelli, P. (1991) 'Photosensitizing action of furocoumarins on membrane  
 1534 components and consequent intracellular events', *Journal of Photochemistry and Photobiology B: Biology*,  
 1535 8(3), pp. 235-254.
- 1536 Damm, U., Baroncelli, R., Cai, L., Kubo, Y., O'Connell, R., Weir, B., Yoshino, K. and Cannon, P. F.  
 1537 (2010) 'Colletotrichum: species, ecology and interactions', *IMA fungus*, 1(2), pp. 161-165.
- 1538 Dardalhon, M., Lin, W., Nicolas, A. and Averbeck, D. (2007) 'Specific transcriptional responses  
 1539 induced by 8-methoxypsoralen and UVA in yeast', *FEMS yeast research*, 7(6), pp. 866-878.
- 1540 Daub, M. and Payne, G. (1989) 'The role of carotenoids in resistance of fungi to cercosporin',  
 1541 *Phytopathology (USA)*.
- 1542 Daub, M. E. (1982) 'Peroxidation of tobacco membrane lipids by the photosensitizing toxin,  
 1543 cercosporin', *Plant Physiology*, 69(6), pp. 1361-1364.
- 1544 Daub, M. E. (1987) 'Resistance of fungi to the photosensitizing toxin, cercosporin', *Phytopathology*,  
 1545 77(11), pp. 1515-1520.
- 1546 Daub, M. E. and Briggs, S. P. (1983) 'Changes in tobacco cell membrane composition and structure  
 1547 caused by cercosporin', *Plant Physiology*, 71(4), pp. 763-766.
- 1548 Daub, M. E. and Ehrenshaft, M. (2000) 'The photoactivated *Cercospora* toxin cercosporin:  
 1549 contributions to plant disease and fundamental biology', *Annual review of phytopathology*, 38(1), pp. 461-  
 1550 490.
- 1551 Daub, M. E. and Hangarter, R. P. (1983) 'Light-induced production of singlet oxygen and  
 1552 superoxide by the fungal toxin, cercosporin', *Plant Physiology*, 73(3), pp. 855-857.
- 1553 Daub, M. E., Herrero, S. and Chung, K.-R. (2005) 'Photoactivated perylenequinone toxins in fungal  
 1554 pathogenesis of plants', *FEMS Microbiology Letters*, 252(2), pp. 197-206.
- 1555 Daub, M. E., Herrero, S. and Chung, K.-R. (2013) 'Reactive oxygen species in plant pathogenesis:  
 1556 the role of perylenequinone photosensitizers', *Antioxidants & Redox Signaling*, 19(9), pp. 970-989.
- 1557 Daub, M. E., Leisman, G. B., Clark, R. A. and Bowden, E. F. (1992) 'Reductive detoxification as a  
 1558 mechanism of fungal resistance to singlet oxygen-generating photosensitizers', *Proceedings of the  
 1559 National Academy of Sciences*, 89(20), pp. 9588-9592.



- de Menezes, H. D., Pereira, A. C., Brancini, G. T. P., de Leao, H. C., Massola Junior, N. S., Bachmann, L., Wainwright, M., Bastos, J. K. and Braga, G. U. L. (2014a) 'Furocoumarins and coumarins photoinactivate *Colletotrichum acutatum* and *Aspergillus nidulans* fungi under solar radiation', *Journal of Photochemistry and Photobiology B-Biology*, 131, pp. 74-83.
- de Menezes, H. D., Rodrigues, G. B., Teixeira Sde, P., Massola, N. S., Jr., Bachmann, L., Wainwright, M. and Braga, G. U. (2014b) 'In vitro photodynamic inactivation of plant-pathogenic fungi *Colletotrichum acutatum* and *Colletotrichum gloeosporioides* with Novel Phenothiazinium photosensitizers', *Appl Environ Microbiol*, 80(5), pp. 1623-32.
- de Menezes, H. D., Tonani, L., Bachmann, L., Wainwright, M., Braga, G. U. and von Zeska Kress, M. R. (2016) 'Photodynamic treatment with phenothiazinium photosensitizers kills both ungerminated and germinated microconidia of the pathogenic fungi *Fusarium oxysporum*, *Fusarium moniliforme* and *Fusarium solani*', *J Photochem Photobiol B*, 164, pp. 1-12.
- Deising, H. B., Reimann, S. and Pascholati, S. F. (2008) 'Mechanisms and significance of fungicide resistance', *Brazilian Journal of Microbiology*, 39(2), pp. 286-295.
- Desjardins, A. E., Spencer, G. F. and Plattner, R. D. (1989) 'Tolerance and metabolism of furanocoumarins by the phytopathogenic fungus *Gibberella pulicaris* (*Fusarium sambucinum*)', *Phytochemistry*, 28(11), pp. 2963-2969.
- Dibona-Villanueva, L. and Fuentealba, D. (2021) 'Novel Chitosan–Riboflavin Conjugate with Visible Light-Enhanced Antifungal Properties against *Penicillium digitatum*', *Journal of Agricultural and Food Chemistry*, 69(3), pp. 945-954.
- Dibona-Villanueva, L. and Fuentealba, D. (2022) 'Protoporphyrin IX–Chitosan Oligosaccharide Conjugate with Potent Antifungal Photodynamic Activity', *Journal of Agricultural and Food Chemistry*.
- DiCosmo, F., Towers, G. N. and Lam, J. (1982) 'Photo-induced fungicidal activity elicited by naturally occurring thiophene derivatives', *Pesticide Science*, 13(6), pp. 589-594.
- Dijksterhuis, J. (2019) 'Fungal spores: Highly variable and stress-resistant vehicles for distribution and spoilage', *Food microbiology*, 81, pp. 2-11.
- Dijksterhuis, J., Meijer, M., van Doorn, T., Samson, R. and Rico-Munoz, E. (2018) 'Inactivation of stress-resistant ascospores of Eurotiales by industrial sanitizers', *International journal of food microbiology*, 285, pp. 27-33.
- Ding, Y., Chen, H., Yang, Q., Feng, L., Hua, X. and Wang, M. (2019) 'A fluorescence polarization immunoassay for detection of thiachloprid in environmental and agricultural samples', *RSC Advances*, 9(63), pp. 36825-36830.
- do Prado-Silva, L., Alvarenga, V. O., Braga, G. Ú. and Sant'Ana, A. S. (2021) 'Inactivation kinetics of *Bacillus cereus* vegetative cells and spores from different sources by antimicrobial photodynamic treatment (aPDT)', *LWT*, 142, pp. 111037.

- 1595 do Prado-Silva, L., Brancini, G. T. P., Braga, G. Ú. L., Liao, X., Ding, T. and Sant'Ana, A. S. (2022)  
 1596 'Antimicrobial photodynamic treatment (aPDT) as an innovative technology to control spoilage and  
 1597 pathogenic microorganisms in agri-food products: An updated review', *Food Control*, 132, pp. 108527.
- 1598 do Prado-Silva, L., Gomes, A. T., Mesquita, M. Q., Neri-Numa, I. A., Pastore, G. M., Neves, M. G.,  
 1599 Faustino, M. A., Almeida, A., Braga, G. U. and Sant'Ana, A. S. (2020) 'Antimicrobial photodynamic  
 1600 treatment as an alternative approach for *Alicyclobacillus acidoterrestris* inactivation', *International Journal*  
 1601 *of Food Microbiology*, 333, pp. 108803.
- 1602 Dong, F., Zhang, X., Xu, J. H., Shi, J. R., Lee, Y.-W., Chen, X. Y., Li, Y. P., Mokoena, M. P. and  
 1603 Olaniran, A. O. (2020) 'Analysis of *Fusarium graminearum* species complex from freshly harvested rice in  
 1604 Jiangsu province (China)', *Plant Disease*, 104(8), pp. 2138-2143.
- 1605 Donley, N. (2019) 'The USA lags behind other agricultural nations in banning harmful pesticides',  
 1606 *Environmental Health*, 18(1), pp. 44.
- 1607 Dowling, M., Peres, N., Villani, S. and Schnabel, G. (2020) 'Managing *Colletotrichum* on Fruit  
 1608 Crops: A "Complex" Challenge', *Plant Disease*, 104(9), pp. 2301-2316.
- 1609 Dutta, J., Tripathi, S. and Dutta, P. K. (2012) 'Progress in antimicrobial activities of chitin, chitosan  
 1610 and its oligosaccharides: a systematic study needs for food applications', *Food Science and Technology*  
 1611 *International*, 18(1), pp. 3-34.
- 1612 Dweba, C., Figlan, S., Shimelis, H., Motaung, T., Sydenham, S., Mwadzingeni, L. and Tsilo, T.  
 1613 (2017) 'Fusarium head blight of wheat: Pathogenesis and control strategies', *Crop Protection*, 91, pp. 114-  
 1614 122.
- 1615 Ehrenshaft, M., Bilski, P., Li, M. Y., Chignell, C. F. and Daub, M. E. (1999a) 'A highly conserved  
 1616 sequence is a novel gene involved in de novo vitamin B6 biosynthesis', *Proceedings of the National*  
 1617 *Academy of Sciences*, 96(16), pp. 9374-9378.
- 1618 Ehrenshaft, M., Chung, K.-R., Jenns, A. E. and Daub, M. E. (1999b) 'Functional characterization of  
 1619 SOR1, a gene required for resistance to photosensitizing toxins in the fungus *Cercospora nicotianae*',  
 1620 *Current genetics*, 34(6), pp. 478-485.
- 1621 Ehrenshaft, M., Jenns, A. and Daub, M. (1995) 'Targeted gene disruption of carotenoid  
 1622 biosynthesis in *Cercospora nicotianae* reveals no role for carotenoids in photosensitizer resistance',  
 1623 *Molecular plant-microbe interactions: MPMI (USA)*.
- 1624 Ehrenshaft, M., Jenns, A. E., Chung, K. R. and Daub, M. E. (1998) 'SOR1, a Gene Required for  
 1625 Photosensitizer and Singlet Oxygen Resistance in *Cercospora* Fungi, Is Highly Conserved in Divergent  
 1626 Organisms', *Molecular Cell*, 1(4), pp. 603-609.
- 1627 Ehrenshaft, M. and Upchurch, R. G. (1991) 'Isolation of light-enhanced cDNAs of *Cercospora*  
 1628 *kikuchii*', *Applied and environmental microbiology*, 57(9), pp. 2671-2676.
- 1629 Eichner, A., Gollmer, A., Späth, A., Bäuml, W., Regensburger, J., Koenig, B. and Maisch, T.  
 1630 (2015) 'Fast and effective inactivation of *Bacillus atrophaeus* endospores using light-activated derivatives  
 1631 of vitamin B2', *Photochemical & Photobiological Sciences*, 14(2), pp. 387-396.

- 1632 Feenstra, R. P. and Tseng, S. C. (1992) 'Comparison of fluorescein and rose bengal staining',  
1633 *Ophthalmology*, 99(4), pp. 605-617.
- 1634 Fernandes, E. K., Rangel, D. E., Braga, G. U. and Roberts, D. W. (2015) 'Tolerance of  
1635 entomopathogenic fungi to ultraviolet radiation: a review on screening of strains and their formulation',  
1636 *Curr Genet*, 61(3), pp. 427-40.
- 1637 Fisher, J. F. and Trama, L. A. (1979) 'High-performance liquid chromatographic determination of  
1638 some coumarins and psoralens found in citrus peel oils', *Journal of Agricultural and Food Chemistry*, 27(6),  
1639 pp. 1334-1337.
- 1640 Fisher, M. C., Hawkins, N. J., Sanglard, D. and Gurr, S. J. (2018) 'Worldwide emergence of  
1641 resistance to antifungal drugs challenges human health and food security', *Science*, 360(6390), pp. 739-  
1642 742.
- 1643 Flors, C. and Nonell, S. (2006) 'Light and singlet oxygen in plant defense against pathogens:  
1644 phototoxic phenalenone phytoalexins', *Accounts of chemical research*, 39(5), pp. 293-300.
- 1645 Forcelini, B. B., Seijo, T. E., Amiri, A. and Peres, N. A. (2016) 'Resistance in strawberry isolates of  
1646 *Colletotrichum acutatum* from Florida to quinone-outside inhibitor fungicides', *Plant Disease*, 100(10), pp.  
1647 2050-2056.
- 1648 Fracarolli, L., Rodrigues, G. B., Pereira, A. C., Massola Junior, N. S., Silva-Junior, G. J., Bachmann,  
1649 L., Wainwright, M., Bastos, J. K. and Braga, G. U. (2016) 'Inactivation of plant-pathogenic fungus  
1650 *Colletotrichum acutatum* with natural plant-produced photosensitizers under solar radiation', *J Photochem*  
1651 *Photobiol B*, 162, pp. 402-11.
- 1652 Friedberg, J. S., Skema, C., Baum, E. D., Burdick, J., Vinogradov, S. A., Wilson, D. F., Horan, A.  
1653 D. and Nachamkin, I. (2001) 'In vitro effects of photodynamic therapy on *Aspergillus fumigatus*', *Journal of*  
1654 *Antimicrobial Chemotherapy*, 48(1), pp. 105-107.
- 1655 Fuentealba, D., López, J. J., Palominos, M., Salas, C. O. and Soto-Arriaza, M. A. (2015) 'Gramicidin  
1656 conformational changes during riboflavin photosensitized oxidation in solution and the effect of N-  
1657 methylation of tryptophan residues', *Photochemical & Photobiological Sciences*, 14(4), pp. 748-756.
- 1658 Gama, A. B., Baggio, J. S., Rebello, C. S., Lourenço, S. d. A., Gasparoto, M. C. d. G., da Silva  
1659 Junior, G. J., Peres, N. A. and Amorim, L. (2020) 'Sensitivity of *Colletotrichum acutatum* Isolates from  
1660 Citrus to Carbendazim, Difenoconazole, Tebuconazole, and Trifloxystrobin', *Plant Disease*, 104(6), pp.  
1661 1621-1628.
- 1662 Gama, A. B., Peres, N. A., Singerman, A. and Dewdney, M. M. (2022) 'Evaluation of disease alert  
1663 systems for postbloom fruit drop of citrus in Florida and economic impact of adopting the Citrus Advisory  
1664 System', *Crop Protection*, 155, pp. 105906.
- 1665 Gamelas, S. R. D., Vieira, C., Bartolomeu, M., Faustino, M. A. F., Tomé, J. P. C., Tomé, A. C.,  
1666 Almeida, A. and Lourenço, L. M. O. (2022) 'Photodynamic inactivation of pathogenic Gram-negative and  
1667 Gram-positive bacteria mediated by Si(IV) phthalocyanines bearing axial ammonium units', *Journal of*  
1668 *Photochemistry and Photobiology B: Biology*, 233, pp. 112502.

- 1669 Gao, L., Jiang, S., Sun, Y., Deng, M., Wu, Q., Li, M. and Zeng, T. (2016) 'Evaluation of the effects  
1670 of photodynamic therapy alone and combined with standard antifungal therapy on planktonic cells and  
1671 biofilms of *Fusarium* spp. and *Exophiala* spp', *Frontiers in microbiology*, 7, pp. 617.
- 1672 Gilaberte, Y., Aspiroz, C., Martes, M. P., Alcalde, V., Espinel-Ingroff, A. and Rezusta, A. (2011)  
1673 'Treatment of refractory fingernail onychomycosis caused by nondermatophyte molds with  
1674 methylaminolevulinate photodynamic therapy', *Journal of the American Academy of Dermatology*, 65(3),  
1675 pp. 669-671.
- 1676 Glueck, M., Schamberger, B., Eckl, P. and Plaetzer, K. (2017) 'New horizons in microbiological  
1677 food safety: Photodynamic Decontamination based on a curcumin derivative', *Photochemical &  
1678 Photobiological Sciences*, 16(12), pp. 1784-1791.
- 1679 Gomes, M. C., Woranovicz-Barreira, S. M., Faustino, M. A., Fernandes, R., Neves, M. G., Tomé,  
1680 A. C., Gomes, N., Almeida, A., Cavaleiro, J. A. and Cunha, A. (2011) 'Photodynamic inactivation of  
1681 *Penicillium chrysogenum* conidia by cationic porphyrins', *Photochemical & Photobiological Sciences*,  
1682 10(11), pp. 1735-1743.
- 1683 Gong, C., Li, Y., Gao, R., Xiao, F., Zhou, X., Wang, H., Xu, H., Wang, R., Huang, P. and Zhao, Y.  
1684 (2020) 'Inactivation of specific spoilage organism (*Pseudomonas*) of sturgeon by curcumin-mediated  
1685 photodynamic inactivation', *Photodiagnosis and Photodynamic Therapy*, 31, pp. 101827.
- 1686 Gong, H. H., Kagan, J., Seitz, R., Stokes, A. B., Meyer, F. A. and Tuveson, R. (1988) 'The  
1687 phototoxicity of phenylheptatriyne: oxygen-dependent hemolysis of human erythrocytes and inactivation  
1688 of *Escherichia coli*', *Photochemistry and photobiology*, 47(1), pp. 55-63.
- 1689 Gonzales, F. P., da Silva, S. H., Roberts, D. W. and Braga, G. U. (2010) 'Photodynamic inactivation  
1690 of conidia of the fungi *Metarhizium anisopliae* and *Aspergillus nidulans* with methylene blue and toluidine  
1691 blue', *Photochem Photobiol*, 86(3), pp. 653-61.
- 1692 Gonzales, F. P. and Maisch, T. (2012) 'Photodynamic inactivation for controlling *Candida albicans*  
1693 infections', *Fungal biology*, 116(1), pp. 1-10.
- 1694 Gonzales, J. C., Brancini, G. T. P., Rodrigues, G. B., Silva-Junior, G. J., Bachmann, L., Wainwright,  
1695 M. and Braga, G. U. L. (2017) 'Photodynamic inactivation of conidia of the fungus *Colletotrichum*  
1696 *abscissum* on *Citrus sinensis* plants with methylene blue under solar radiation', *J Photochem Photobiol B*,  
1697 176, pp. 54-61.
- 1698 Gonçalves, F. P., Nogueira Júnior, A. F., Silva-Junior, G. J., Ciampi-Guillardi, M. and Amorim, L.  
1699 (2021) 'Environmental requirements for infection of *Colletotrichum acutatum* and *C. gloeosporioides* sensu  
1700 lato in citrus flowers and prevalence of these pathogens in Brazil', *European Journal of Plant Pathology*,  
1701 160(1), pp. 27-37.
- 1702 Grant, E., Von Borstel, R. and Ashwood-Smith, M. (1979) 'Mutagenicity of cross-links and  
1703 monoadducts of furocoumarins (psoralen and angelicin) induced by 360-nm radiation in excision-repair-  
1704 defective and radiation-insensitive strains of *saccharomyces cerevisiae*', *Environmental Mutagenesis*,  
1705 1(1), pp. 55-63.

- Guarro, J. (2013) 'Fusariosis, a complex infection caused by a high diversity of fungal species refractory to treatment', *European Journal of Clinical Microbiology & Infectious Diseases*, 32(12), pp. 1491-1500.
- Gulías, Ò., McKenzie, G., Bayó, M., Agut, M. and Nonell, S. (2020) 'Effective photodynamic inactivation of 26 Escherichia coli strains with different antibiotic susceptibility profiles: A planktonic and biofilm study', *Antibiotics*, 9(3), pp. 98.
- Gunnell, D., Knipe, D., Chang, S.-S., Pearson, M., Konradsen, F., Lee, W. J. and Eddleston, M. (2017) 'Prevention of suicide with regulations aimed at restricting access to highly hazardous pesticides: a systematic review of the international evidence', *The Lancet Global Health*, 5(10), pp. e1026-e1037.
- Hallsworth, J. E., Heim, S. and Timmis, K. N. (2003) 'Chaotropic solutes cause water stress in Pseudomonas putida', *Environmental Microbiology*, 5(12), pp. 1270-1280.
- Hamblin, M. R. (2016) 'Antimicrobial photodynamic inactivation: a bright new technique to kill resistant microbes', *Current Opinion in Microbiology*, 33, pp. 67-73.
- Hamblin, M. R., O'Donnell, D. A., Murthy, N., Rajagopalan, K., Michaud, N., Sherwood, M. E. and Hasan, T. (2002) 'Polycationic photosensitizer conjugates: effects of chain length and Gram classification on the photodynamic inactivation of bacteria', *Journal of Antimicrobial Chemotherapy*, 49(6), pp. 941-951.
- Hamill, P. G., Stevenson, A., McMullan, P. E., Williams, J. P., Lewis, A. D. R., S, S., Stevenson, K. E., Farnsworth, K. D., Khroustalyova, G., Takemoto, J. Y., Quinn, J. P., Rapoport, A. and Hallsworth, J. E. (2020) 'Microbial lag phase can be indicative of, or independent from, cellular stress', *Scientific Reports*, 10(1), pp. 5948.
- Hamminger, C., Glueck, M., Fefer, M., Ckurshumova, W., Liu, J., Tenhaken, R. and Plaetzer, K. (2022) 'Photodynamic Inactivation of plant pathogens part II: fungi', *Photochemical & Photobiological Sciences*, 21(2), pp. 195-207.
- Harris, F. and Pierpoint, L. (2012) 'Photodynamic therapy based on 5-aminolevulinic acid and its use as an antimicrobial Agent', *Medicinal research reviews*, 32(6), pp. 1292-1327.
- Hasenleitner, M. and Plaetzer, K. (2020) 'In the Right Light: Photodynamic Inactivation of Microorganisms using a LED-based illumination device tailored for the Antimicrobial Application', *Antibiotics*, 9(1), pp. 13.
- Hayashi, K., Schoonbeek, H.-j. and De Waard, M. A. (2002) 'Bcmfs1, a novel major facilitator superfamily transporter from Botrytis cinerea, provides tolerance towards the natural toxic compounds camptothecin and cercosporin and towards fungicides', *Applied and environmental microbiology*, 68(10), pp. 4996-5004.
- Heaselgrave, W. and Kilvington, S. (2010) 'Antimicrobial activity of simulated solar disinfection against bacterial, fungal, and protozoan pathogens and its enhancement by riboflavin', *Applied and Environmental Microbiology*, 76(17), pp. 6010-6012.

- Heaton, J. C. and Jones, K. (2008) 'Microbial contamination of fruit and vegetables and the behaviour of enteropathogens in the phyllosphere: a review', *Journal of applied microbiology*, 104(3), pp. 613-626.
- Heiser, I., Sachs, E. and Liebermann, B. (2003) 'Photodynamic oxygen activation by rubellin D, a phytotoxin produced by *Ramularia collo-cygni* (Sutton et Waller)', *Physiological and Molecular Plant Pathology*, 62(1), pp. 29-36.
- Herrero, S. and Daub, M. E. (2007) 'Genetic manipulation of vitamin B-6 biosynthesis in tobacco and fungi uncovers limitations to up-regulation of the pathway', *Plant science*, 172(3), pp. 609-620.
- Hidalgo, W., Chandran, J. N., Menezes, R. C., Otálvaro, F. and Schneider, B. (2016) 'Phenylphenalenones protect banana plants from infection by *Mycosphaerella fijiensis* and are deactivated by metabolic conversion', *Plant, cell & environment*, 39(3), pp. 492-513.
- Hidalgo, W., Kai, M. and Schneider, B. (2015) '4-Methoxycinnamic acid—An unusual phenylpropanoid involved in phenylphenalenone biosynthesis in *Anigozanthos preissii*', *Phytochemistry*, 117, pp. 476-481.
- Hu, J., Lin, S., Tan, B. K., Hamzah, S. S., Lin, Y., Kong, Z., Zhang, Y., Zheng, B. and Zeng, S. (2018) 'Photodynamic inactivation of *Burkholderia cepacia* by curcumin in combination with EDTA', *Food Research International*, 111, pp. 265-271.
- Huang, L., Krayner, M., Roubil, J. G., Huang, Y.-Y., Holten, D., Lindsey, J. S. and Hamblin, M. R. (2014) 'Stable synthetic mono-substituted cationic bacteriochlorins mediate selective broad-spectrum photoinactivation of drug-resistant pathogens at nanomolar concentrations', *Journal of Photochemistry and Photobiology B: Biology*, 141, pp. 119-127.
- Huang, L., Yong, K. W. L., Fernando, W. C., Carpinelli de Jesus, M., De Voss, J. J., Sultanbawa, Y. and Fletcher, M. T. (2021) 'The Inactivation by Curcumin-Mediated Photosensitization of *Botrytis cinerea* Spores Isolated from Strawberry Fruits', *Toxins*, 13(3).
- Hudson, J. and Towers, G. (1991) 'Therapeutic potential of plant photosensitizers', *Pharmacology & therapeutics*, 49(3), pp. 181-222.
- Ishii, H. and Holloman, D. (2015) 'Fungicide resistance in plant pathogens', *Tokyo: Springer, doi*, 10, pp. 978-4.
- Ito, T. (1977) 'Toluidine blue: the mode of photodynamic action in yeast cells', *Photochemistry and photobiology*, 25(1), pp. 47-53.
- Ito, T. (1978) 'Cellular and subcellular mechanisms of photodynamic action: the  $^1O_2$  hypothesis as a driving force in recent research', *Photochemistry and Photobiology*, 28(4-5), pp. 493-506.
- Jančula, D., Bláhová, L., Karásková, M. and Maršálek, B. (2010) 'Degradation of natural toxins by phthalocyanines—example of cyanobacterial toxin, microcystin', *Water Science and Technology*, 62(2), pp. 273-278.
- Jenns, A. and Daub, M. (1995) 'Characterization of mutants of *Cercospora nicotianae* sensitive to the toxin cercosporin', *Phytopathology*.

- Jensen, R., Hagen, F., Astvad, K., Tyron, A., Meis, J. and Arendrup, M. (2016) 'Azole-resistant *Aspergillus fumigatus* in Denmark: a laboratory-based study on resistance mechanisms and genotypes', *Clinical Microbiology and Infection*, 22(6), pp. 570. e1-570. e9.
- Jess, S., Kildea, S., Moody, A., Rennick, G., Murchie, A. K. and Cooke, L. R. (2014) 'European Union policy on pesticides: implications for agriculture in Ireland', *Pest management science*, 70(11), pp. 1646-1654.
- Joshi, P. C. and Pathak, M. A. (1983) 'Production of singlet oxygen and superoxide radicals by psoralens and their biological significance', *Biochemical and biophysical research communications*, 112(2), pp. 638-646.
- Kairyte, K., Kadys, A. and Luksiene, Z. (2013) 'Antibacterial and antifungal activity of photoactivated ZnO nanoparticles in suspension', *Journal of Photochemistry and Photobiology B: Biology*, 128, pp. 78-84.
- Kairyte, K., Lapinskas, S., Gudelis, V. and Luksiene, Z. (2012) 'Effective inactivation of food pathogens *Listeria monocytogenes* and *Salmonella enterica* by combined treatment of hypericin-based photosensitization and high power pulsed light', *Journal of applied microbiology*, 112(6), pp. 1144-1151.
- Kamp, H., Tietz, H. J., Lutz, M., Piazena, H., Sowyrda, P., Lademann, J. and Blume-Peytavi, U. (2005) 'Antifungal effect of 5-aminolevulinic acid PDT in *Trichophyton rubrum*', *Mycoses*, 48(2), pp. 101-107.
- Kashiwabuchi, R. T., Carvalho, F. R., Khan, Y. A., Hirai, F., Campos, M. S. and McDonnell, P. J. (2013) 'Assessment of fungal viability after long-wave ultraviolet light irradiation combined with riboflavin administration', *Graefes Archive for Clinical and Experimental Ophthalmology*, 251(2), pp. 521-527.
- Ke, C.-L., Deng, F.-S., Chuang, C.-Y. and Lin, C.-H. (2021) 'Antimicrobial actions and applications of chitosan', *Polymers*, 13(6), pp. 904.
- Kettles, G. J. and Luna, E. (2019) 'Food security in 2044: How do we control the fungal threat?', *Fungal biology*, 123(8), pp. 558-564.
- Kim, J. S., Je, Y. H. and Choi, J. Y. (2010) 'Complementary effect of Phloxine B on the insecticidal efficacy of *Isaria fumosorosea* SFP-198 wettable powder against greenhouse whitefly, *Trialeurodes vaporariorum* West', *Pest management science*, 66(12), pp. 1337-1343.
- Kourany, E., Arnason, J. T. and Schneider, E. (1988) 'Accumulation of phototoxic thiophenes in *Tagetes erecta* (Asteraceae) elicited by *Fusarium oxysporum*', *Physiological and molecular plant pathology*, 33(2), pp. 287-297.
- Kretschmer, M., Leroch, M., Mosbach, A., Walker, A.-S., Fillinger, S., Mernke, D., Schoonbeek, H.-J., Pradier, J.-M., Leroux, P. and De Waard, M. A. (2009) 'Fungicide-driven evolution and molecular basis of multidrug resistance in field populations of the grey mould fungus *Botrytis cinerea*', *PLoS pathogens*, 5(12), pp. e1000696.
- Kruk, J. and Szymańska, R. (2021) 'Singlet oxygen oxidation products of carotenoids, fatty acids and phenolic prennylipids', *Journal of Photochemistry and Photobiology B: Biology*, 216, pp. 112148.

- Larson, R. A. and Berenbaum, M. R. (1988) 'Environmental phototoxicity', *Environmental Science & Technology*, 22(4), pp. 354-360.
- Lazzaro, A., Corominas, M., Martí, C., Flors, C., Izquierdo, L. R., Grillo, T. A., Luis, J. G. and Nonell, S. (2004) 'Light-and singlet oxygen-mediated antifungal activity of phenylphenalenone phytoalexins', *Photochemical & Photobiological Sciences*, 3(7), pp. 706-710.
- Leisman, G. B. and Daub, M. E. (1992) 'Singlet oxygen yields, optical properties, and phototoxicity of reduced derivatives of the photosensitizer cercosporin', *Photochemistry and photobiology*, 55(3), pp. 373-379.
- Lievens, B., Hallsworth, J. E., Pozo, M. I., Belgacem, Z. B., Stevenson, A., Willems, K. A. and Jacquemyn, H. (2015) 'Microbiology of sugar-rich environments: diversity, ecology and system constraints', *Environmental Microbiology*, 17(2), pp. 278-298.
- Liu, Z., Tang, J., Sun, Y. and Gao, L. (2019) 'Effects of Photodynamic Inactivation on the Growth and Antifungal Susceptibility of *Rhizopus Oryzae*', *Mycopathologia*, 184(2), pp. 315-319.
- Llano, J., Raber, J. and Eriksson, L. A. (2003) 'Theoretical study of phototoxic reactions of psoralens', *Journal of Photochemistry and Photobiology A: Chemistry*, 154(2-3), pp. 235-243.
- Lopez-Carballo, G., Hernández-Muñoz, P., Gavara, R. and Ocio, M. (2008) 'Photoactivated chlorophyllin-based gelatin films and coatings to prevent microbial contamination of food products', *International journal of food microbiology*, 126(1-2), pp. 65-70.
- Lucas, J. A., Dyer, P. S. and Murray, T. D. (2000) 'Pathogenicity, host-specificity, and population biology of *tapesia* spp., causal agents of eyespot disease of cereals', *Advances in Botanical Research: Academic Press*, pp. 225-258.
- Luis, J. G., Fletcher, W. Q., Echeverri, F. and Grillo, T. A. (1994) 'Phenalenone-type phytoalexins from *Musa acuminata* synthesis of 4-phenyl-phenalenones', *Tetrahedron*, 50(37), pp. 10963-10970.
- Luksevičiute, V. and Luksiene, Z. (2020) 'Inactivation of molds on the surface of wheat sprouts by chlorophyllin-chitosan coating in the presence of visible LED-based light', *Journal of Photochemistry and Photobiology B: Biology*, 202, pp. 111721.
- Luksiene, Z. and Brovko, L. (2013) 'Antibacterial photosensitization-based treatment for food safety', *Food Engineering Reviews*, 5(4), pp. 185-199.
- Luksiene, Z. and Buchovec, I. (2019) 'Impact of chlorophyllin-chitosan coating and visible light on the microbial contamination, shelf life, nutritional and visual quality of strawberries', *Innovative Food Science & Emerging Technologies*, 52, pp. 463-472.
- Luksiene, Z., Buchovec, I. and Paskeviciute, E. (2009) 'Inactivation of food pathogen *Bacillus cereus* by photosensitization in vitro and on the surface of packaging material', *Journal of applied microbiology*, 107(6), pp. 2037-2046.
- Luksiene, Z., Buchovec, I. and Paskeviciute, E. (2010a) 'Inactivation of *Bacillus cereus* by Na-chlorophyllin-based photosensitization on the surface of packaging', *Journal of applied microbiology*, 109(5), pp. 1540-1548.



- 1852        Luksiene, Z., Buchovec, I. and Paskeviciute, E. (2010b) 'Inactivation of several strains of *Listeria*  
1853        *monocytogenes* attached to the surface of packaging material by Na-chlorophyllin-based  
1854        photosensitization', *Journal of Photochemistry and Photobiology B: Biology*, 101(3), pp. 326-331.
- 1855        Luksiene, Z. and Paskeviciute, E. (2011) 'Novel approach to the microbial decontamination of  
1856        strawberries: chlorophyllin-based photosensitization', *Journal of Applied Microbiology*, 110(5), pp. 1274-  
1857        1283.
- 1858        Luksiene, Z., Peculyte, D. and Lugauskas, A. (2004) 'Inactivation of fungi in vitro by  
1859        photosensitization: preliminary results', *Annals of Agricultural and Environmental Medicine*, 11(2).
- 1860        Lukšienė, Z. and Zukauskas, A. (2009) 'Prospects of photosensitization in control of pathogenic  
1861        and harmful micro-organisms', *Journal of Applied Microbiology*, 107(5), pp. 1415-1424.
- 1862        Lukšienė, Ž., Pečiulyte, D., Jurkonienė, S. and Puras, R. (2005) 'Inactivation of possible fungal food  
1863        contaminants by photosensitization', *Food Technology and Biotechnology*, 43(4), pp. 335-341.
- 1864        Lukšienė, Ž., Danilčenko, H., Tarasevičienė, Ž., Anusevičius, Ž., Marozienė, A. and Nivinskas, H.  
1865        (2007) 'New approach to the fungal decontamination of wheat used for wheat sprouts: effects of  
1866        aminolevulinic acid', *International journal of food microbiology*, 116(1), pp. 153-158.
- 1867        Lysøe, E., Harris, L. J., Walkowiak, S., Subramaniam, R., Divon, H. H., Riiser, E. S., Llorens, C.,  
1868        Gabaldón, T., Kistler, H. C. and Jonkers, W. (2014) 'The genome of the generalist plant pathogen *Fusarium*  
1869        *avenaceum* is enriched with genes involved in redox, signaling and secondary metabolism', *PLoS One*,  
1870        9(11), pp. e112703.
- 1871        Ma, L.-J., Geiser, D. M., Proctor, R. H., Rooney, A. P., O'Donnell, K., Trail, F., Gardiner, D. M.,  
1872        Manners, J. M. and Kazan, K. (2013) 'Fusarium pathogenomics', *Annual review of microbiology*, 67, pp.  
1873        399-416.
- 1874        Madden, L., Yang, X. and Wilson, L. (1996) 'Effects of rain intensity on splash dispersal of  
1875        *Colletotrichum acutatum*', *Phytopathology*, 86(8), pp. 864-874.
- 1876        Manderfeld, M. M., Schafer, H. W., Davidson, P. M. and Zottola, E. A. (1997) 'Isolation and  
1877        identification of antimicrobial furocoumarins from parsley', *Journal of food protection*, 60(1), pp. 72-77.
- 1878        Mares, D., Tosi, B., Poli, F., Andreotti, E. and Romagnoli, C. (2004) 'Antifungal activity of *Tagetes*  
1879        *patula* extracts on some phytopathogenic fungi: ultrastructural evidence on *Pythium ultimum*',  
1880        *Microbiological Research*, 159(3), pp. 295-304.
- 1881        Mares, D., Tosi, B., Romagnoli, C. and Poli, F. (2002) 'Antifungal activity of *Tagetes patula* extracts',  
1882        *Pharmaceutical biology*, 40(5), pp. 400-404.
- 1883        Martin, P. A., Mischke, S. and Schroder, R. F. (1998) 'Compatibility of photoactive dyes with insect  
1884        biocontrol agents', *Biocontrol Science and Technology*, 8(4), pp. 501-508.
- 1885        Martin-Urdiroz, M., Osés-Ruiz, M., Ryder, L. S. and Talbot, N. J. (2016) 'Investigating the biology  
1886        of plant infection by the rice blast fungus *Magnaporthe oryzae*', *Fungal Genetics and Biology*, 90, pp. 61-  
1887        68.

- 1888 Martinez De Pinillos Bayona, A., Mroz, P., Thunshelle, C. and Hamblin, M. R. (2017) 'Design  
1889 features for optimization of tetrapyrrole macrocycles as antimicrobial and anticancer photosensitizers',  
1890 *Chemical biology & drug design*, 89(2), pp. 192-206.
- 1891 Martins, D., Mesquita, M. Q., Neves, M. G., Faustino, M. A., Reis, L., Figueira, E. and Almeida, A.  
1892 (2018) 'Photoinactivation of *Pseudomonas syringae* pv. *actinidiae* in kiwifruit plants by cationic porphyrins',  
1893 *Planta*, 248(2), pp. 409-421.
- 1894 Massey, V. (2000) 'The chemical and biological versatility of riboflavin', *Biochemical Society*  
1895 *Transactions*, 28(4), pp. 283-296.
- 1896 Mir, S. A., Farooq, S., Shah, M. A., Sofi, S. A., Dar, B., Hamdani, A. M. and Khaneghah, A. M.  
1897 (2021) 'An overview of sprouts nutritional properties, pathogens and decontamination technologies', *LWT*,  
1898 141, pp. 110900.
- 1899 Mischke, S., Martin, P. A. and Schroder, R. F. (1998) 'Compatibility of phloxine B, an insecticidal  
1900 photoactive dye, with selected biocontrol fungi', *Biocontrol Science and Technology*, 8(4), pp. 509-515.
- 1901 Mukherjee, R., Gruszeński Hope, A., Bilyeu Landon, T., Schmale David, G. and Boreyko Jonathan,  
1902 B. (2021) 'Synergistic dispersal of plant pathogen spores by jumping-droplet condensation and wind',  
1903 *Proceedings of the National Academy of Sciences*, 118(34), pp. e2106938118.
- 1904 Muzzoli, M. and Sacchetti, G. (2001) 'Biological activity of four thiophene compounds in resting  
1905 *Saccharomyces cerevisiae* cells', *Pharmaceutical biology*, 39(1), pp. 40-42.
- 1906 Naranjo, A., Arboleda, A., Martinez, J. D., Durkee, H., Aguilar, M. C., Relhan, N., Nikpoor, N., Galor,  
1907 A., Dubovy, S. R. and Leblanc, R. (2019) 'Rose Bengal photodynamic antimicrobial therapy for patients  
1908 with progressive infectious keratitis: a pilot clinical study', *American journal of ophthalmology*, 208, pp.  
1909 387-396.
- 1910 Nascimento, E., da Silva, S. H., Marques Edos, R., Roberts, D. W. and Braga, G. U. (2010)  
1911 'Quantification of cyclobutane pyrimidine dimers induced by UVB radiation in conidia of the fungi  
1912 *Aspergillus fumigatus*, *Aspergillus nidulans*, *Metarhizium acridum* and *Metarhizium robertsii*', *Photochem*  
1913 *Photobiol*, 86(6), pp. 1259-66.
- 1914 Nassar, S. J. M., Wills, C. and Harriman, A. (2019) 'Inhibition of the Photobleaching of Methylene  
1915 Blue by Association with Urea', *ChemPhotoChem*, 3(10), pp. 1042-1049.
- 1916 Nielsen, H. K., Garcia, J., Væth, M. and Schlafer, S. (2015) 'Comparison of riboflavin and toluidine  
1917 blue O as photosensitizers for photoactivated disinfection on endodontic and periodontal pathogens in  
1918 vitro', *PLoS One*, 10(10), pp. e0140720.
- 1919 Nigg, H., Nordby, H., Beier, R., Dillman, A., Macias, C. and Hansen, R. (1993) 'Phototoxic  
1920 coumarins in limes', *Food and chemical toxicology*, 31(5), pp. 331-335.
- 1921 Nitzan, Y. and Ashkenazi, H. (1999) 'Photoinactivation of *Deinococcus radiodurans*: An Unusual  
1922 Gram-Positive Microorganism', *Photochemistry and Photobiology*, 69(4), pp. 505-510.
- 1923 Ntahimpera, N., Madden, L. and Wilson, L. (1997) 'Effect of rain distribution alteration on splash  
1924 dispersal of *Colletotrichum acutatum*', *Phytopathology*, 87(6), pp. 649-655.

- 1925 Nuñez, S. C., Yoshimura, T. M., Ribeiro, M. S., Junqueira, H. C., Maciel, C., Coutinho-Neto, M. D.  
1926 and Baptista, M. S. (2015) 'Urea enhances the photodynamic efficiency of methylene blue', *Journal of*  
1927 *Photochemistry and Photobiology B: Biology*, 150, pp. 31-37.
- 1928 Oliver, R. P. and Hewitt, H. G. (2014) *Fungicides in crop protection*. Cabi.
- 1929 Oren, A. and Hallsworth, J. E. (2014) 'Microbial weeds in hypersaline habitats: the enigma of the  
1930 weed-like *Haloferax mediterranei*', *FEMS Microbiology Letters*, 359(2), pp. 134-142.
- 1931 Paardekooper, M., Bruune, A. W. D., Steveninck, J. V. and Broek, P. J. V. d. (1995) 'Intracellular  
1932 damage in yeast cells caused by photodynamic treatment with toluidine blue', *Photochemistry and*  
1933 *photobiology*, 61(1), pp. 84-89.
- 1934 Paardekooper, M., Van den Broek, P. J., De Bruijne, A. W., Elferink, J. G., Dubbelman, T. M. and  
1935 Van Steveninck, J. (1992) 'Photodynamic treatment of yeast cells with the dye toluidine blue: all-or-none  
1936 loss of plasma membrane barrier properties', *Biochimica et Biophysica Acta (BBA)-Biomembranes*,  
1937 1108(1), pp. 86-90.
- 1938 Paulussen, C., Hallsworth, J. E., Álvarez-Pérez, S., Nierman, W. C., Hamill, P. G., Blain, D.,  
1939 Rediers, H. and Lievens, B. (2017) 'Ecology of aspergillosis: insights into the pathogenic potency of  
1940 *Aspergillus fumigatus* and some other *Aspergillus* species', *Microbial biotechnology*, 10(2), pp. 296-322.
- 1941 Paziani, M. H., Tonani, L., de Menezes, H. D., Bachmann, L., Wainwright, M., Braga, G. Ú. L. and  
1942 von Zeska Kress, M. R. (2019) 'Antimicrobial photodynamic therapy with phenothiazinium photosensitizers  
1943 in non-vertebrate model *Galleria mellonella* infected with *Fusarium keratoplasticum* and *Fusarium*  
1944 *moniliforme*', *Photodiagnosis and Photodynamic Therapy*, 25, pp. 197-203.
- 1945 Pellosi, D. S., Estevão, B. M., Semensato, J., Severino, D., Baptista, M. S., Politi, M. J., Hioka, N.  
1946 and Caetano, W. (2012) 'Photophysical properties and interactions of xanthene dyes in aqueous micelles',  
1947 *Journal of Photochemistry and Photobiology A: Chemistry*, 247, pp. 8-15.
- 1948 Peres, N. A., Seijo, T. E. and Turechek, W. W. (2010) 'Pre-and post-inoculation activity of a  
1949 protectant and a systemic fungicide for control of anthracnose fruit rot of strawberry under different wetness  
1950 durations', *Crop Protection*, 29(10), pp. 1105-1110.
- 1951 Peres, N. A., Timmer, L. W., Adaskaveg, J. E. and Correll, J. C. (2005) 'Lifestyles of *Colletotrichum*  
1952 *acutatum*', *Plant Disease*, 89(8), pp. 784-796.
- 1953 Polmickaitė-Smirnova, E., Buchovec, I., Bagdonas, S., Sužiedėlienė, E., Ramanavičius, A. and  
1954 Anusevičius, Ž. (2022) 'Photoinactivation of *Salmonella enterica* exposed to 5-aminolevulinic acid: Impact  
1955 of sensitization conditions and irradiation time', *Journal of Photochemistry and Photobiology B: Biology*,  
1956 231, pp. 112446.
- 1957 Potapenko, A. Y. (1991) 'New trends in photobiology: mechanisms of photodynamic effects of  
1958 furocoumarins', *Journal of Photochemistry and Photobiology B: Biology*, 9(1), pp. 1-33.
- 1959 Prandini, J. A., Castro, K. A. D. F., Biazotto, J. C., Brancini, G. T. P., Tomé, J. P. C., Lourenço, L.  
1960 M. O., Braga, G. Ú. L. and da Silva, R. S. (2022) 'Thiopyridinium phthalocyanine for improved

- photodynamic efficiency against pathogenic fungi', *Journal of Photochemistry and Photobiology B: Biology*, 231, pp. 112459.
- Qi, H.-z., Wang, W.-z., He, J.-y., Ma, Y., Xiao, F.-z. and He, S.-y. (2020) 'Antioxidative system of *Deinococcus radiodurans*', *Research in Microbiology*, 171(2), pp. 45-54.
- Rangel, D. E., Braga, G. U., Anderson, A. J. and Roberts, D. W. (2005) 'Variability in conidial thermotolerance of *Metarhizium anisopliae* isolates from different geographic origins', *J Invertebr Pathol*, 88(2), pp. 116-25.
- Rangel, D. E., Braga, G. U., Fernandes, E. K., Keyser, C. A., Hallsworth, J. E. and Roberts, D. W. (2015) 'Stress tolerance and virulence of insect-pathogenic fungi are determined by environmental conditions during conidial formation', *Curr Genet*, 61(3), pp. 383-404.
- Revie, N. M., Iyer, K. R., Robbins, N. and Cowen, L. E. (2018) 'Antifungal drug resistance: evolution, mechanisms and impact', *Current opinion in microbiology*, 45, pp. 70-76.
- Ribas e Ribas, A. D., Spolti, P., Del Ponte, E. M., Donato, K. Z., Schrekker, H. and Fuentefria, A. M. (2016) 'Is the emergence of fungal resistance to medical triazoles related to their use in the agroecosystems? A mini review', *brazilian journal of microbiology*, 47, pp. 793-799.
- Rodrigues, G. B., Brancini, G. T. P., Pinto, M. R., Primo, F. L., Wainwright, M., Tedesco, A. C. and Braga, G. U. L. (2020a) 'Photodynamic inactivation of *Candida albicans* and *Candida tropicalis* with aluminum phthalocyanine chloride nanoemulsion', *Fungal Biology*, 124(5), pp. 297-303.
- Rodrigues, G. B., Brancini, G. T. P., Uyemura, S. A., Bachmann, L., Wainwright, M. and Braga, G. U. L. (2020b) 'Chemical features of the photosensitizers new methylene blue N and S137 influence their subcellular localization and photoinactivation efficiency in *Candida albicans*', *Journal of Photochemistry and Photobiology B: Biology*, 209, pp. 111942.
- Rodrigues, G. B., Dias-Baruffi, M., Holman, N., Wainwright, M. and Braga, G. U. (2013) 'In vitro photodynamic inactivation of *Candida* species and mouse fibroblasts with phenothiazinium photosensitisers and red light', *Photodiagnosis Photodyn Ther*, 10(2), pp. 141-9.
- Rodrigues, G. B., Ferreira, L. K., Wainwright, M. and Braga, G. U. (2012a) 'Susceptibilities of the dermatophytes *Trichophyton mentagrophytes* and *T. rubrum* microconidia to photodynamic antimicrobial chemotherapy with novel phenothiazinium photosensitizers and red light', *Journal of Photochemistry and Photobiology B: Biology*, 116, pp. 89-94.
- Rodrigues, G. B., Primo, F. L., Tedesco, A. C. and Braga, G. U. (2012b) 'In vitro photodynamic inactivation of *Cryptococcus neoformans* melanized cells with chloroaluminum phthalocyanine nanoemulsion', *Photochemistry and Photobiology*, 88(2), pp. 440-447.
- Romagnoli, C., Mares, D., Sacchetti, G. and Bruni, A. (1998) 'The photodynamic effect of 5-(4-hydroxy-1-butinyl)-2, 2'-bithienyl on dermatophytes', *Mycological research*, 102(12), pp. 1519-1524.
- Sabino, C. P., Wainwright, M., Ribeiro, M. S., Sellera, F. P., dos Anjos, C., Baptista, M. d. S. and Lincopan, N. (2020) 'Global priority multidrug-resistant pathogens do not resist photodynamic therapy', *Journal of Photochemistry and Photobiology B: Biology*, 208, pp. 111893.

- 1998 Santos, R., de Carvalho, C. C., Stevenson, A., Grant, I. R. and Hallsworth, J. E. (2015)  
 1999 'Extraordinary solute-stress tolerance contributes to the environmental tenacity of mycobacteria',  
 2000 *Environmental microbiology reports*, 7(5), pp. 746-764.
- 2001 Sauer, A., Letscher-Bru, V., Speeg-Schatz, C., Touboul, D., Colin, J., Candolfi, E. and Bourcier, T.  
 2002 (2010) 'In vitro efficacy of antifungal treatment using riboflavin/UV-A (365 nm) combination and  
 2003 amphotericin B', *Investigative ophthalmology & visual science*, 51(8), pp. 3950-3953.
- 2004 Schafer, M., Schmitz, C. and Horneck, G. (1998) 'High sensitivity of *Deinococcus radiodurans* to  
 2005 photodynamically-produced singlet oxygen', *International Journal of Radiation Biology*, 74(2), pp. 249-253.
- 2006 Schwechheimer, S. K., Park, E. Y., Revuelta, J. L., Becker, J. and Wittmann, C. (2016)  
 2007 'Biotechnology of riboflavin', *Applied microbiology and biotechnology*, 100(5), pp. 2107-2119.
- 2008 Shatila, W., Verma, A. and Adam, S. (2017) 'Plasmapheresis in severe methemoglobinemia  
 2009 following occupational exposure', *Transfusion and Apheresis Science*, 56(3), pp. 341-344.
- 2010 Shimizu, M., Egashira, T. and Takahama, U. (1979) 'Inactivation of *Neurospora crassa* conidia by  
 2011 singlet molecular oxygen generated by a photosensitized reaction', *Journal of bacteriology*, 138(2), pp.  
 2012 293-296.
- 2013 Sierra-Garcia, I. N., Cunha, Â. and Lourenço, L. M. O. (2022) 'In vitro photodynamic treatment of  
 2014 *Fusarium oxysporum* conidia through the action of thiopyridinium and methoxypyridinium chlorins', *Journal*  
 2015 *of Photochemistry and Photobiology A: Chemistry*, 432, pp. 114081.
- 2016 Silva, A. F., Borges, A., Giaouris, E., Graton Mikcha, J. M. and Simoes, M. (2018) 'Photodynamic  
 2017 inactivation as an emergent strategy against foodborne pathogenic bacteria in planktonic and sessile  
 2018 states', *Critical reviews in microbiology*, 44(6), pp. 667-684.
- 2019 Silva-Junior, G. J., Spósito, M. B., Marin, D. R. and Amorim, L. (2014) 'Efficacy and timing of  
 2020 application of fungicides for control of citrus postbloom fruit drop', *Crop Protection*, 59, pp. 51-56.
- 2021 Skovsen, E., Snyder, J. W., Lambert, J. D. and Ogilby, P. R. (2005) 'Lifetime and diffusion of singlet  
 2022 oxygen in a cell', *The Journal of Physical Chemistry B*, 109(18), pp. 8570-8573.
- 2023 Smijs, T., Dame, Z., de Haas, E., Aans, J.-B., Pavel, S. and Sterenborg, H. (2014) 'Photodynamic  
 2024 and Nail Penetration Enhancing Effects of Novel Multifunctional Photosensitizers Designed for The  
 2025 Treatment of Onychomycosis', *Photochemistry and Photobiology*, 90(1), pp. 189-200.
- 2026 Smijs, T. G. and Pavel, S. (2011) 'The susceptibility of dermatophytes to photodynamic treatment  
 2027 with special focus on *Trichophyton rubrum*', *Photochemistry and photobiology*, 87(1), pp. 2-13.
- 2028 Sobotta, L., Skupin-Mrugalska, P., Piskorz, J. and Mielcarek, J. (2019) 'Porphyrinoid  
 2029 photosensitizers mediated photodynamic inactivation against bacteria', *European journal of medicinal*  
 2030 *chemistry*, 175, pp. 72-106.
- 2031 St. Denis, T. G., Dai, T., Izikson, L., Astrakas, C., Anderson, R. R., Hamblin, M. R. and Tegos, G.  
 2032 P. (2011) 'All you need is light: antimicrobial photoinactivation as an evolving and emerging discovery  
 2033 strategy against infectious disease', *Virulence*, 2(6), pp. 509-520.

- 2034 Stenglein, S. (2009) 'Fusarium poae: a pathogen that needs more attention', *Journal of Plant*  
 2035 *Pathology*, pp. 25-36.
- 2036 Stevenson, A., Burkhardt, J., Cockell, C. S., Cray, J. A., Dijksterhuis, J., Fox-Powell, M., Kee, T.  
 2037 P., Kminek, G., McGenity, T. J. and Timmis, K. N. (2015a) 'Multiplication of microbes below 0.690 water  
 2038 activity: implications for terrestrial and extraterrestrial life', *Environmental microbiology*, 17(2), pp. 257-277.
- 2039 Stevenson, A., Cray, J. A., Williams, J. P., Santos, R., Sahay, R., Neuenkirchen, N., McClure, C.  
 2040 D., Grant, I. R., Houghton, J. D. and Quinn, J. P. (2015b) 'Is there a common water-activity limit for the  
 2041 three domains of life?', *The ISME journal*, 9(6), pp. 1333-1351.
- 2042 Stevenson, A., Hamill, P. G., Dijksterhuis, J. and Hallsworth, J. E. (2017a) 'Water-, pH-and  
 2043 temperature relations of germination for the extreme xerophiles *Xeromyces bisporus* (FRR 0025),  
 2044 *Aspergillus penicillioides* (JH 06 THJ) and *Eurotium halophilicum* (FRR 2471)', *Microbial biotechnology*,  
 2045 10(2), pp. 330-340.
- 2046 Stevenson, A., Hamill, P. G., Medina, Á., Kminek, G., Rummel, J. D., Dijksterhuis, J., Timson, D.  
 2047 J., Magan, N., Leong, S.-L. L. and Hallsworth, J. E. (2017b) 'Glycerol enhances fungal germination at the  
 2048 water-activity limit for life', *Environmental Microbiology*, 19(3), pp. 947-967.
- 2049 Stevenson, A., Hamill, P. G., O'Kane, C. J., Kminek, G., Rummel, J. D., Voytek, M. A., Dijksterhuis,  
 2050 J. and Hallsworth, J. E. (2017c) '*Aspergillus penicillioides* differentiation and cell division at 0.585 water  
 2051 activity', *Environmental Microbiology*, 19(2), pp. 687-697.
- 2052 Sumorek-Wiadro, J., Zając, A., Maciejczyk, A. and Jakubowicz-Gil, J. (2020) 'Furanocoumarins in  
 2053 anticancer therapy—For and against', *Fitoterapia*, 142, pp. 104492.
- 2054 Suryawanshi, R., Patil, C., Borase, H., Narkhede, C., Stevenson, A., Hallsworth, J. and Patil, S.  
 2055 (2015) 'Towards an understanding of bacterial metabolites prodigiosin and violacein and their potential for  
 2056 use in commercial sunscreens', *International journal of cosmetic science*, 37(1), pp. 98-107.
- 2057 Symes, S., Goldsmith, P. and Haines, H. (2015) 'Microbiological safety and food handling practices  
 2058 of seed sprout products in the Australian State of Victoria', *Journal of food protection*, 78(7), pp. 1387-  
 2059 1391.
- 2060 Tang, J., Tang, G., Niu, J., Yang, J., Zhou, Z., Gao, Y., Chen, X., Tian, Y., Li, Y. and Li, J. (2021)  
 2061 'Preparation of a porphyrin metal–organic framework with desirable photodynamic antimicrobial activity for  
 2062 sustainable plant disease management', *Journal of Agricultural and Food Chemistry*, 69(8), pp. 2382-  
 2063 2391.
- 2064 Taniguchi, M. and Lindsey, J. S. (2017) 'Synthetic chlorins, possible surrogates for chlorophylls,  
 2065 prepared by derivatization of porphyrins', *Chemical reviews*, 117(2), pp. 344-535.
- 2066 Tegegne, G., Pretorius, J. and Swart, W. (2008) 'Antifungal properties of *Agapanthus africanus* L.  
 2067 extracts against plant pathogens', *Crop Protection*, 27(7), pp. 1052-1060.
- 2068 Tegos, G. P., Anbe, M., Yang, C., Demidova, T. N., Satti, M., Mroz, P., Janjua, S., Gad, F. and  
 2069 Hamblin, M. R. (2006) 'Protease-stable polycationic photosensitizer conjugates between

- polyethyleneimine and chlorin (e6) for broad-spectrum antimicrobial photoinactivation', *Antimicrobial agents and chemotherapy*, 50(4), pp. 1402-1410.
- Temba, B. A., Fletcher, M. T., Fox, G. P., Harvey, J. J. and Sultanbawa, Y. (2016) 'Inactivation of *Aspergillus flavus* spores by curcumin-mediated photosensitization', *Food control*, 59, pp. 708-713.
- Thomas, E., Herrero, S., Eng, H., Gomaa, N., Gillikin, J., Noar, R., Beseli, A. and Daub, M. E. (2020) 'Engineering *Cercospora* disease resistance via expression of *Cercospora nicotianae* cercosporin-resistance genes and silencing of cercosporin production in tobacco', *Plos one*, 15(3), pp. e0230362.
- Thomas, P. (2003) 'Fungal infections of the cornea', *Eye*, 17(8), pp. 852-862.
- Thomas, S. A., Sargent, M. and Tuveson, R. (1981) 'Inactivation of normal and mutant *Neurospora crassa* conidia by visible light and near-UV: role of  $^1O_2$ , carotenoid composition and sensitizer location', *Photochemistry and Photobiology*, 33(3), pp. 349-354.
- Timer, L. and Zitko, S. (1991) 'Evaluation of fungicides and application frequency for control of postbloom fruit drop of citrus caused by *Colletotrichum gloeosporioides*', 31. *Reunión Anual de la Sociedad Americana de Fitopatología. División Caribe, San José (Costa Rica), 20-25 May 1991*.
- Tonani, L., Morosini, N. S., Dantas de Menezes, H., Nadaletto Bonifacio da Silva, M. E., Wainwright, M., Leite Braga, G. U. and Regina von Zeska Kress, M. (2018) 'In vitro susceptibilities of *Neoscytalidium* spp. sequence types to antifungal agents and antimicrobial photodynamic treatment with phenothiazinium photosensitizers', *Fungal Biol*, 122(6), pp. 436-448.
- Tortik, N., Spaeth, A. and Plaetzer, K. (2014) 'Photodynamic decontamination of foodstuff from *Staphylococcus aureus* based on novel formulations of curcumin', *Photochemical & Photobiological Sciences*, 13(10), pp. 1402-1409.
- Trigos, A. and Ortega-Regules, A. (2002) 'Selective destruction of microscopic fungi through photo-oxidation of ergosterol', *Mycologia*, 94(4), pp. 563-568.
- Tussen, J., Beekes, H. and Van Steveninck, J. (1981) 'Localization of polyphosphates at the outside of the yeast cell plasma membrane', *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 649(3), pp. 529-532.
- Ulate-Rodríguez, J., Schafer, H. W., Zottola, E. A. and Davidson, P. M. (1997) 'Inhibition of *Listeria monocytogenes*, *Escherichia coli* O157: H7, and *Micrococcus luteus* by linear furanocoumarins in a model food system', *Journal of food protection*, 60(9), pp. 1050-1054.
- Uliana, M. P., Pires, L., Pratavieira, S., Brocksom, T. J., de Oliveira, K. T., Bagnato, V. S. and Kurachi, C. (2014) 'Photobiological characteristics of chlorophyll a derivatives as microbial PDT agents', *Photochemical & Photobiological Sciences*, 13(8), pp. 1137-1145.
- Upchurch, R. G., Rose, M. S., Eweida, M. and Callahan, T. M. (2002) 'Transgenic assessment of CFP-mediated cercosporin export and resistance in a cercosporin-sensitive fungus', *Current genetics*, 41(1), pp. 25-30.
- Uyar, G. E. Ö. and Uyar, B. (2018) 'Effects of ethanol and ultraviolet-C treatments on inactivation of *Rhizopus oryzae* spores which cause postharvest rot', *Food Science and Technology*, 39, pp. 691-695.

- 2107 Vandresen, C. C., Gonçalves, A. G., Ducatti, D. R. B., Murakami, F. S., Nosedá, M. D., Duarte, M.  
2108 E. R. and Barreira, S. M. W. (2016) 'In vitro photodynamic inactivation of conidia of the phytopathogenic  
2109 fungus *Colletotrichum graminicola* with cationic porphyrins', *Photochemical & Photobiological Sciences*,  
2110 15(5), pp. 673-681.
- 2111 Vecchio, D., Gupta, A., Huang, L., Landi, G., Avci, P., Rodas, A. and Hamblin, M. R. (2015)  
2112 'Bacterial photodynamic inactivation mediated by methylene blue and red light is enhanced by synergistic  
2113 effect of potassium iodide', *Antimicrobial agents and chemotherapy*, 59(9), pp. 5203-5212.
- 2114 Vera, D. M. A., Haynes, M. H., Ball, A. R., Dai, T., Astrakas, C., Kelso, M. J., Hamblin, M. R. and  
2115 Tegos, G. P. (2012) 'Strategies to potentiate antimicrobial photoinactivation by overcoming resistant  
2116 phenotypes', *Photochemistry and photobiology*, 88(3), pp. 499-511.
- 2117 Ververidis, P., Davrazou, F., Dailianas, G., Georgakopoulos, D., Kanellis, A. and Panopoulos, N.  
2118 (2001) 'A novel putative reductase (Cpd1p) and the multidrug exporter Snq2p are involved in resistance  
2119 to cercosporin and other singlet oxygen-generating photosensitizers in *Saccharomyces cerevisiae*',  
2120 *Current genetics*, 39(3).
- 2121 Vol'pin, M. E., Novodárova, G. N., Krainova, N. Y., Lapikova, V. P. and Aver'yanov, A. A. (2000)  
2122 'Redox and fungicidal properties of phthalocyanine metal complexes as related to active oxygen', *Journal*  
2123 *of Inorganic Biochemistry*, 81(4), pp. 285-292.
- 2124 Vorobey, A. and Pinchuk, S. (2008) 'Photodamage to spores of *Fusarium* fungi sensitized by  
2125 protoporphyrin IX', *Biophysics*, 53(5), pp. 386-389.
- 2126 Wainwright, M. (2010) 'Safe photoantimicrobials for skin and soft-tissue infections', *International*  
2127 *journal of antimicrobial agents*, 36(1), pp. 14-18.
- 2128 Wainwright, M., Antczak, J., Baca, M., Loughran, C. and Meegan, K. (2015) 'Phenothiazinium  
2129 photoantimicrobials with basic side chains', *Journal of Photochemistry and Photobiology B: Biology*, 150,  
2130 pp. 38-43.
- 2131 Wainwright, M., Maisch, T., Nonell, S., Plaetzer, K., Almeida, A., Tegos, G. P. and Hamblin, M. R.  
2132 (2017) 'Photoantimicrobials-are we afraid of the light?', *Lancet Infect Dis*, 17(2), pp. e49-e55.
- 2133 Wainwright, M., Meegan, K. and Loughran, C. (2011) 'Phenothiazinium photosensitisers IX. Tetra-  
2134 and pentacyclic derivatives as photoantimicrobial agents', *Dyes and Pigments*, 91(1), pp. 1-5.
- 2135 Walther, G., Wagner, L. and Kurzai, O. (2020) 'Outbreaks of Mucorales and the species involved',  
2136 *Mycopathologia*, 185(5), pp. 765-781.
- 2137 Wecke, T. and Mascher, T. (2011) 'Antibiotic research in the age of omics: from expression profiles  
2138 to interspecies communication', *Journal of Antimicrobial Chemotherapy*, 66(12), pp. 2689-2704.
- 2139 Weiman, S. (2014) 'Farm to Table: Predicting, Preventing Foodborne Outbreaks', *Microbe*  
2140 *Magazine*, 9, pp. 357-358.
- 2141 Wharton, P. S. and Diéguez-Urbeondo, J. 'The biology of *Colletotrichum acutatum*'. *Anales del*  
2142 *jardín botánico de Madrid*, 3-22.



- Will, O. H., Newland, N. A. and Reppe, C. R. (1984) 'Photosensitivity of pigmented and nonpigmented strains of *Ustilago violacea*', *Current Microbiology*, 10(5), pp. 295-301.
- Wilson, R. A. and Talbot, N. J. (2009) 'Under pressure: investigating the biology of plant infection by *Magnaporthe oryzae*', *Nature Reviews Microbiology*, 7(3), pp. 185-195.
- Wong, F. P., De la Cerda, K. A., Hernandez-Martinez, R. and Midland, S. L. (2008) 'Detection and characterization of benzimidazole resistance in California populations of *Colletotrichum cereale*', *Plant disease*, 92(2), pp. 239-246.
- Wong, F. P. and Midland, S. L. (2007) 'Sensitivity distributions of California populations of *Colletotrichum cereale* to the DMI fungicides propiconazole, myclobutanil, tebuconazole, and triadimefon', *Plant Disease*, 91(12), pp. 1547-1555.
- Woo, P. C., Ngan, A. H., Tsang, C. C., Ling, I. W., Chan, J. F., Leung, S.-Y., Yuen, K.-Y. and Lau, S. K. (2013) 'Clinical spectrum of *Exophiala* infections and a novel *Exophiala* species, *Exophiala hongkongensis*', *Journal of Clinical Microbiology*, 51(1), pp. 260-267.
- Wrolstad, R. E. and Culver, C. A. (2012) 'Alternatives to those artificial FD&C food colorants', *Annual review of food science and technology*, 3, pp. 59-77.
- Wyatt, T. T., Golovina, E. A., van Leeuwen, R., Hallsworth, J. E., Wösten, H. A. and Dijksterhuis, J. (2015a) 'A decrease in bulk water and mannitol and accumulation of trehalose and trehalose-based oligosaccharides define a two-stage maturation process towards extreme stress resistance in ascospores of *N. eosartorya fischeri* (A *spargillus fischeri*)', *Environmental Microbiology*, 17(2), pp. 383-394.
- Wyatt, T. T., Van Leeuwen, M. R., Golovina, E. A., Hoekstra, F. A., Kuenstner, E. J., Palumbo, E. A., Snyder, N. L., Visagie, C., Verkennis, A. and Hallsworth, J. E. (2015b) 'Functionality and prevalence of trehalose-based oligosaccharides as novel compatible solutes in ascospores of *N. eosartorya fischeri* (A *spargillus fischeri*) and other fungi', *Environmental Microbiology*, 17(2), pp. 395-411.
- Yan, X. and Talbot, N. J. (2016) 'Investigating the cell biology of plant infection by the rice blast fungus *Magnaporthe oryzae*', *Current opinion in microbiology*, 34, pp. 147-153.
- Zobel, A. M. and Brown, S. A. (1989) 'Histological localization of furanocoumarins in *Ruta graveolens* shoots', *Canadian journal of botany*, 67(3), pp. 915-921.
- Zulfiqar, M., Brlansky, R. and Timmer, L. (1996) 'Infection of flower and vegetative tissues of citrus by *Colletotrichum acutatum* and *C. gloeosporioides*', *Mycologia*, 88(1), pp. 121-128.
- Świdarska-Burek, U., Daub, M. E., Thomas, E., Jaszek, M., Pawlik, A. and Janusz, G. (2020) 'Phytopathogenic cercosporoid fungi—from taxonomy to modern biochemistry and molecular biology', *International Journal of Molecular Sciences*, 21(22), pp. 8555.
- Žudytė, B., Velička, M., Šablinskas, V. and Lukšienė, Ž. (2020) 'Understanding *Escherichia coli* damages after chlorophyllin-based photosensitization', *Journal of Biophotonics*, 13(11), pp. e202000144.