

HUMAN EXPOSURE TO ORGANOCHLORINE PESTICIDES IN VEGETABLES FROM MAJOR CITIES IN SOUTH-SOUTH NIGERIA

Godswill Okeoghene **Tesi**^{1*}, Grace **Obi-Iyeke**², Jude Chinedu **Ossai**³, Anthony A. **Ogbuta**¹, Evieva F. **Ogbara**⁴, Daniel I. **Olorunfemi**^{4,5} and Iwekumo E. **Agbozu**^{1,6}

¹Department of Chemical Sciences, University of Africa, Toru-Orua, Nigeria

²Department of Botany, Delta State University Abraka, Nigeria

³Department of Chemistry, Delta State University Abraka, Nigeria

⁴Department of Biological Sciences, University of Africa, Toru-Orua, Nigeria

⁵Department of Environmental Management and Toxicology, University of Benin, Benin-City

⁶Department of Environmental Management and Toxicology, Federal University of Petroleum Resources, Effurun, Nigeria

*Correspondence author: godswillinfodesk@yahoo.com; godswill.tesi@uat.edu.ng;
+234(0)7035312293

HIGHLIGHTS

- The vegetables were contaminated with OCPs
- The OCPs in the vegetables were lower than their respective MRLs
- The daily intakes were below their respective FAO/WHO acceptable limits
- The TCR values indicated low to moderate carcinogenic risks
- OCPs in the vegetables originated from both historical and recent usage

RESPONSES TO REVIEWERS COMMENTS

Reviewer #1: Overview and general recommendation:

This work contains useful information on the human exposure of OCPs in vegetables from major cities in Nigeria. The paper needs some revisions as there are a few things to be checked and/or corrected. They are given below:

Specific comments:

1. **Comment:** In the introduction, please more elaborate the reason why the authors only selected four vegetables as the samples in the study, e.g. if most consumed, how many consumption per year. Why these vegetables are so important to be analyzed?
Response: More elaborate reasons why the authors only selected four vegetables as the samples in the study have been given.
2. **Comment:** Why the samples only collected from four major cities in Nigeria, are they most populated or vegetables producer? Please provide the profiles of sampling area.
Response: Samples were collected from four major cities in the south-south region of Nigeria and not Nigeria in general. They are the most populated cities in the region. The profiles of sampling area have been included.
3. **Comment:** Are the vegetables locally produced? Or are cultivated from other cities.
Responses: The vegetables are locally produced.
4. **Comment:** Page 3. Please provide the sampling period in the method.
Response: The sampling period has been added in the method section
5. **Comment:** Page 10. The authors have tried to identify potential sources of contamination and suggested that several OCPs are from recent usage. Please provide data or reference of OCPs usage in Nigeria, e.g. from other study or exports and imports of OCPs
Response: Data or reference of OCPs usage in Nigeria, e.g. from other study or exports and imports of OCPs has been included in the introduction and source identification section of discussion.
6. **Comment:** In the results section, a PCA was performed. Please provide the method of PCA in the method section.
Response: This has been included

Reviewer #3:

The paper presents results on vegetables as a source of persistent pollutant exposure to the consumers in the study area. Although similar studies have been carried out on leafy vegetables in the southwest and Northern Nigeria, the study adds to the information available on presence of pesticide residues in leafy vegetables sampled from Nigeria.

Abstract

Comment: The abstract was well written

Response: Thank you

Comment: In the first few mentions of ng g⁻¹, the Author should indicate that concentrations are fresh weights; ng g⁻¹fresh wt or (fw) or dry weight (dwt).

Response: This has been included

Comment: Other editorial corrections are indicated in the annotated manuscript

Responses: These has been effected

Introduction

Comment: Syntax errors are indicated in the annotated manuscript

Response: These has been effected

Materials & Methods

Sampling and sample pre-treatment

Comment: How significant are the markets that the samples were obtained from? There is no information on this in the study.

Response: The significance of the markets has been added

Comment: In Figure 1, is it possible to add a sample collection map showing the 4 markets used as sample locations in the 4 cities?

Response: This has been added

Sample Preparation and Extraction.

Comment: Line 99-100, The type of extractor used and volume of solvent was not stated. I am surprised that the extraction was carried out for 10 hours.

Response: These were not stated because the detailed procedure have been previously reported.

Comment: QUECHERS (Quick Easy Cheap Effective Rugged Safe) method are highly beneficial analytical approach which simplifies the analysis of multiple pesticide residues in fruit, vegetables and cereals within a shorter time. This is will be good for future work of this nature.

Response: This has been noted by authors. Thank you

Comment: What was the percentage purity of the stream of Nitrogen used for extraction?

Response: This has been included

Quantification of the extract samples

Comment: The GC-MS mode adopted for the qualitative and quantitative analysis was not stated.

Response: This has been added

Quality Assurance

Comment: Recovery results look good and appear to be well carried out. However, were any certified plant-based reference materials extracted to guarantee the analytical precision. If not, the addition of a certified/standard reference material in a plant tissue would be a good addition to future reports of this nature.

Response: This has been noted by authors. Thank you

Risk assessment

Comment: Page 5, Line 134-135, "In this study, the body weight of 80 Kg for adults and 15 Kg for children were used and ingestion rate = 165 g day⁻¹ was adopted (Tesi et al., 2021)." The reference cited in SM2 is different from what was cited in the manuscript.

Response: This section has been rephrase and revised accordingly

Comment: For the ingestion rate, the author needs to rely on/use sub-national or national survey data or world health organization/FAO data such as cluster diet as used by numerous authors across the world. In reality, children and adults do not consume the same quantity of vegetables daily!

Response: WHO/FAO value has been used to recompute the health risk.

Comment: Bodyweight of 80kg is on the high side for adults, the author needs to check up on other works carried out in your country not necessary from your research group. The average weight for adults in Africa should be around 60-70

(https://stats.areppim.com/stats/stats_weightboysng.htm; <https://doi.org/10.1186/1471-2458-12-439>)

Response: Body weight of 60 kg has been used to recomputed the health risk

Comment: In the risk assessment, what were the health endpoints for which the authors computed the hazard risk index? Where are these for Cardiovascular? Respiratory? Etc.... The specific health endpoint is highly important in risk assessment and characterization. The authors should search for health endpoints and the target organs of the oral Reference Dose and cancer slope factor used for the risk estimation.

Response: The health endpoints has been included in the manuscript while the target organs of the oral Reference Dose and cancer slope factor used for the risk estimation has been included in the supplemental material SM3.

Comment: The limitation or shortcoming of the assumed values used for the study needs to be stated. For example; the absorption and bioavailability rates of pesticide residues in the children and adults, the age of adults and children, and the concentration used for the analysis (mean concentration, 10 percentile, 90th percentile.....)

Response: The limitation or shortcoming has been added

Data Analysis

Comment: For analysis of variance, were the data tested for normality and equal variance? Were any transformations necessary? Almost all contaminant or toxicology data is left-censored with a skewed distribution best treated by log-transformation (Log (n+1)).

Response: Data were tested for normality using the Shapiro Wilks and Kolmogorov Smirnov tests and no transformation was done.

Comment: ANOVA can only be carried out when the assumptions are satisfied and if failed, a non-parametric test can be carried out. None of the Table is showing the result from the ANOVA or the means separation. Can the author utilise statistics (either parametric or non-parametric test) to check if differences exist among/between concentrations across the 4 locations as well as the 4 vegetables?

Response: The non-parametric Kruskal Wallis H test was applied to determine significant variation in the concentrations of OCPs in the vegetables since the data were not normal based on the Shapiro Wilks and Kolmogorov Smirnov tests. The results of the Kruskal Wallis H test are shown in supplemental material.

Results Presentation

Comment: In Table 1, DDT, DDD, and DDE should be written as p,p'-DDT, p,p'-DDD, and p,p'-DDE respectively. This correction should be made on all mentions (Figures and Tables inclusive).

Response: This has been effected.

Comment: The classification of the OCP as shown in Tables 1-2 is wrong. Methoxychlor also known as Methoxy-DDT or Dimethoxy-DDT or p,p'-Dimethoxydiphenyltrichloroethane should belong to the DDT group.

Response: This has been effected

Comment: Line 163, the range presented for green leaves is different from the range 7.63 - 54.2 ng g⁻¹ reported in Table 2.

Response: This has been corrected

Comment: Line 168, maximum residue limits (MRLs) stipulated by the European Union for the respective congeners were not presented in any of the Tables.

Response: This has been corrected

Comment: 178-180, the statement "The occurrence pattern of OCPs in the vegetables followed the order Σ HCHs > Σ Chlordanes > Σ Drins > Σ DDTs > Σ Endosulfans for all the sampled cities in southern Nigeria" is not true for all cases. The author should cross-check. For example; in Table 1, the statement is not true for all the cities, Chlordane was highest for Warri (Bitter leaves), Yenagoa (Bitter leaves) and so on.

Response: This has been revised

Comment: In Figure 1, is it possible to add a sample collection map showing the 4 markets used as sample locations in the 4 cities?

Response: This has been added

Comment: In Fig 2, italicise the scientific names and use another design such as a pie chart where the percentage contribution of each class/group can be shown

Response: Pie chart has been used here

Comment: In Fig 3 C-E, the pie chart should be bigger or use other designs to make it easier for the reader to grasp or maintain the current size, remove the caption and include a colour identification code for each of the congeners.

Response: This has been done

Comment: In Fig 4, The authors should reference the benchmark values on the graphs with a horizontal delimitation which will show clearly what is "above" and "below" those thresholds. It would be much nicer to read and easier to see. For example, a line showing the benchmark values (HRI=1) behind the bars.

Response: Benchmark values has been referenced on the graphs with a horizontal delimitation in yellow colour

Comment: There is a need for authors to cross-check the data presented in Tables and Figures with data presented in the result section.

Response: This has been done

DISCUSSION

Comment: Re-writing presenting this study in a longitudinal format, reflecting on other research on vegetables and food products across Nigeria and other countries of the world might be good and show clearly that it's an ongoing problem.

Response: This has been done (See Table 3).

Comment: Page 6, Line 170-176, there is a need to state the mean concentration or range of the OCP reported in the studies. Were their concentrations in wet weight or dry weight?

Response: This has been added. However, no information was available whether the concentrations were in wet or dry weight

Comment: Page 9, Line 235-247, The possible health effects (both non-carcinogenic or systemic and carcinogenic) due to exposure to the OCP congeners were not stated. How does the risk estimate in this study differs or in similarity to other studies in your country?

Response: This has been added

Comment: Can the author present a longitudinal review to show current trends or levels of the pesticides in the country's vegetables in comparison with detected levels in this study?

Response: This has been done (See Table 3)

Conclusion

Comment: See annotated manuscript


Response: This has been corrected

References

Comment: The references cited in supplementary materials were not listed in the supplementary file.

Response: The references has been listed

GRAPHICAL ABSTRACT



HUMAN EXPOSURE TO ORGANOCHLORINE PESTICIDES IN VEGETABLES FROM MAJOR CITIES IN SOUTH-SOUTH NIGERIA

ABSTRACT

Contamination of vegetables with organochlorine pesticides (OCPs) during cultivation could affect their nutritional value and also results in adverse health effects to consumers. Thus, this study evaluates human exposure to OCPs in vegetables from major cities in south-south Nigeria. A total of eighty vegetables consisting of *Vernonia amygdalina*, *Telfairia occidentalis*, *Desmodium intortum* cv and *Talinum triangulare* obtained from four major cities (Warri, Benin, Yenagoa and Port-Harcourt) in south-south Nigeria were analysed for OCPs using a gas chromatograph coupled with a mass selective detector. The results showed that the mean concentration of $\sum 20$ OCPs in the vegetables ranged from 11.6 to 37.7 ng g⁻¹ fresh wt for *Vernonia amygdalina*, 2.48 to 37.0 ng g⁻¹ fresh wt for *Telfairia occidentalis*, 7.63-54.2 ng g⁻¹ fresh wt for *Desmodium intortum* cv and 27.3 to 57.3 ng g⁻¹ fresh wt for *Talinum triangulare*. The concentrations of OCPs were generally lower than their respective Maximum Residue Limits (MRLs). The EDI values of the $\sum 20$ OCPs in the vegetables ranged from 141 to 464 ng kg⁻¹ bw day⁻¹ for children and 26.5 to 87.0 ng kg⁻¹ bw day⁻¹ for adults. The EDI of \sum Drins, \sum Chlordanes and \sum DDTs for the vegetables were below their respective acceptable EDI values set by FAO/WHO Joint Meeting on Pesticide Residue. The hazard index values for children and adults were generally < 1 suggesting that there is no potential non-carcinogenic risk for children and adults consuming the vegetables. The total cancer risk values were above 1×10^{-6} and indicated that children and adults have moderate and low carcinogenic risks respectively from ingestion of these vegetables. The isomeric ratios and principal component analysis result showed that OCPs in these vegetables originated from both historical and recent usage in vegetable cultivation.

Keywords: Vegetables, OCPs, Daily intake, Risks, Nigeria

1.1 Introduction

Vegetables constituted the second most consumed food category in West Africa after cereals (Stadlmayr *et al.*, 2013). Vegetables are bought from farmers or traders and consumed raw or cooked. They provide extra vitamins and minerals to the body and play essential nutritional, social and economic roles for consumers (Bolor *et al.*, 2018; Adeleye *et al.*, 2019; Tesi *et al.*, 2021). During cultivation, several pests are attracted to vegetables. To control and manage these pests as well as increase yield, farmers applied pesticides. Approximately, between 125,000 and 130,000 metric tonnes of pesticides is used annually in Nigeria (Asogwa and Dongo, 2009). The importation of pesticides rose progressively from \$13 million in 2001 to

\$28 million in 2003 (FAO, 2005) and this trend has continue till date (PAN, 2007; Oyekunle *et al.*, 2011). For instance, a total of one hundred and forty-seven thousand, four hundred and forty-six (147, 446) tonnes of pesticides comprising five hundred and eighty-four (584) tonnes of toxic pesticides were imported into Nigeria for crops production in 2018 (FAO Statistics, 2020). In 2019, Nigeria import \$306 million worth of pesticides while in 2020, Nigeria import \$321 million worth of pesticides resulting in 4.87% import growth (<https://oec.world/en/profile/hs/pesticides>). Organochlorine pesticides are classic persistent organic pollutants extensively applied to prevent and eradicate fungi, insects, weeds, and bacterial effects in agriculture (Gereslassie *et al.*, 2019; Tesi *et al.*, 2020). OCPs are of global concern because of their toxicity, persistency, bioaccumulation tendency, ecological and human health effects via the food chain (Yang *et al.*, 2012; Bai *et al.*, 2015). Some health effects of OCPs include endocrine disruption, development of cancer cells, immunological and neurological disorders, foetal and reproductive, neuro-toxicological and immunotoxicological disorders; reproductive and foetal defects, enzyme inhibition; cryptorchidism, and low sperm concentration (Botella *et al.*, 2004; Barlow, 2005; Yucra *et al.*, 2006; Qin *et al.*, 2019; Tesi *et al.*, 2020; Emoyan *et al.*, 2021).

There is currently an increase in the rate of vegetables consumption worldwide (Thompson and Agbugba, 2013; Adetunde *et al.*, 2018; Tesi *et al.*, 2021). Thus, studies assessing the healthy state of vegetables with reference to their OCPs contents are needed as residues of OCPs may contaminate and accumulate in vegetables during cultivation. In Nigeria, some studies on OCPs in vegetables have been documented (Benson and Arowajoye, 2011; Akan *et al.*, 2014; Njoku *et al.*, 2017; Adefemi *et al.*, 2018; Ibrahim *et al.*, 2018; Adeleye *et al.*, 2019). However, none of these studies have evaluated OCPs content in *Vernonia amygdalina*, *Desmodium intortum* cv and *Talinum triangulare* and only two studies have assessed OCPs in *Telfairia occidentalis*. Also, these previous studies were done in northern and south western

Nigeria. *Vernonia amygdalina*, *Telfairia occidentalis*, *Desmodium intortum* cv and *Talinum triangulare* are very popular, readily available, cheap and mostly consumed leafy vegetables by majority of the populace in south-south Nigeria. They are generally cultivated in any open land space near homes, in roadsides, farmlands and gardens. Data on OCPs in the above mentioned vegetables are however lacking in south-south region of Nigeria. Thus, the aim of the present study is to assess OCPs concentrations in vegetables, health risks of OCPs associated with consumption of vegetables and sources of OCPs in the vegetables from four major cities in south-south Nigeria. Such data is essential for quality control of food and moreover helps consumers to make an informed decision about the choice of foods in order to reduce risk.

1.2 Materials and Methods

1.2.1 Study area

The study area comprised four major cities of Warri, Benin, Yenagoa and Port-Harcourt in south-south Nigeria (Figure 1). These four major cities are the commercial capitals of Delta, Edo, Bayelsa and Rivers States respectively. These cities are densely populated with population of 943,000 for Warri, 1,841,000 for Benin, 352,285 for Yenagoa and 3,325,000 for Port Harcourt in 2022 based on the United Nations – World Population Prospect (<https://www.macrotrends.net/cities>). Four markets from each of the cities were used (Figure 1). These markets include Main market (WM1), Okere market (WM2), Igbudu market (WM3) and Polokor market (WM4) from Warri City; Oba market (BM1), Ekiosa market (BM2), New Benin market (BM3) and Ikpoba Hill market BM4) from Benin City; Opolo market (YM1), Swali market (YM2), Tombia market (YM3) and Kpasia market (YM4) from Yenagoa City; Oil Mill market (PM1), Mile 1 market (PM2), Abuloma market (PM3) and Rumuigbo market (PM4) from Port Harcourt City. These markets are significant because

they are the biggest, most visited and located at the centre of these cities. Thus, they were selected based on their sizes, location and their popularity.

1.2.2 Sample collection

Four vegetables were sampled and they include; *Vernonia amygdalina*, *Telfairia occidentalis*, *Desmodium intortum* cv and *Talinum triangulare* popularly called bitter leaves, pumpkin leaves, green leaves and water leaves respectively. Sample collection was done between October and November, 2021. Five composites of each vegetable were purchased from four markets in each of the four major cities giving eighty samples in all. Vegetables were locally cultivated in these cities and also from neighbouring towns. Vegetable samples were wrapped with aluminium foil, labeled and transported to the laboratory. Thereafter, the vegetables were rinsed with distilled water, sliced into pieces and kept in refrigerator at 4 °C prior to analysis.

1.2.3 Reagents

The reagents used in this study were of spectra grade and include OCPs standards from Accu Standard, USA, dichloromethane from GFS Chemicals, Columbus, n-hexane from Ultrafine Limited, London, Florisil from Labtech Chemicals, Italy and anhydrous sodium sulfate from Merck, Germany.

1.2.4 Sample extraction and clean up

The USEPA method 3550C as earlier described by Tesi *et al.* (2021) was followed for OCPs extraction and cleans up in the vegetables. Briefly, 10 g of the homogenous vegetable sample was weighed into an extraction thimble, and extraction was carried out for 10 hours using

dichloromethane/hexane mixture. The extract was concentrated using a rotary evaporator, cleaned up was carried out on a column packed with acidified silica gel, Florisil, anhydrous Na₂SO₄ and copper powder. Elution was carried out with 40 mL of DCM/hexane. The eluent was collected and evaporated to near dryness with under a stream of nitrogen gas (99.99 % pure) and kept in a vial for analysis.

1.2.5 Instrumental analysis

The determination of the OCPs in the vegetables was performed using a gas chromatograph (Agilent 6890N) coupled with a mass selective detector (GC–MSD). The injection temperature of 250 °C, detector temperature was 290 °C and column was DB-17 (30m × 250 µm × 0.25 µm). High purity helium gas with a steady flow rate of 0.8 mL/min was used as the carrier gas. A 1 µL sample was injected into the GC–MS in splitless mode. Initial oven temperature was 150 °C, increase to 280 °C at 6 °C/min and final temperature was 300 °C. The GC-MSD mode adopted for the analysis was the selective ion mode (SIM).

1.2.6 Quality control and assurance

Spiked recovery technique and blank analysis were used for quality control in this study. Already analysed sample was spiked with standard solution of the OCPs and the spiked sample was analysed. Then the percent recovery was computed. The percent recovery of the OCPs ranged from 88.6%–102 %. The relative standard deviation for replicate analyses (*n* = 3) was < 6 %. Analysis of blank samples was done to obtain the limits of detection (LOD) and quantification (LOQ). The LOD was obtained from the concentration of the OCPs that formed a signal/noise ratio of 3, while the LOQ is the concentration of the OCPs that formed a signal/noise ratio of 10. The values of LODs, LOQs, R² and % recoveries of OCPs are shown in Supplemental Material SM1.

1.2.7 Statistical Analysis

Data analyses were completed using the IBM Statistical Product and Service Solutions (IBM SPSS version 25). Descriptive statistics such as mean, standard deviation and range were used. Also, inferential statistics such as the non-parametric Kruskal Wallis H test was applied to determine significant variation in the concentrations of OCPs in the vegetables after the Shapiro Wilks and Kolmogorov Smirnov tests were used to determine the normality of the data. Principal component analysis and isomeric ratios were used to determine the sources of OCPs in the vegetables.

1.3 Risk assessment of OCPs in the vegetables

1.3.1 Estimation Daily Intake (EDI)

The EDI of OCPs from consumption of these vegetables was computed from the equation:

$$EDI (ng\ kg^{-1}\ bwday^{-1}) = \frac{OCPs\ Concentration \times Ingestion\ Rate}{Body\ Weight} \quad (1)$$

In this study, the body weight of 60 Kg for adults and 15 Kg for children were used (Tesi *et al.*, 2021) and ingestion rate = 46.4 g day⁻¹ was adopted (WHO, 2012a, b).

1.3.3 Non-carcinogenic and carcinogenic risks

The OCPs non-carcinogenic risk from ingestion of the vegetables was assessed as hazard index (HI) while the carcinogenic risk was assessed as total cancer risk (TCR) using the equations below (USEPA, 2009):

$$HI = \left[\frac{C \times IngR \times EF \times ED}{BW \times AT_{nc}} \times 10^{-6} \right] / RfD \quad (2)$$

$$TCR = \frac{C \times IngR \times EF \times ED \times SFO}{BW \times AT_{ca}} \times 10^{-6} \quad (3)$$

The definition of terms and values for each of the variables in the equations (1) and (2) are given in supplemental material SM2 and SM3 and Tesi *et al.* (2021). The HI value > 1 suggests that non-cancer risk is present and vice versa while TCR value above 1×10^{-6} suggests that presence of cancer risk is present and vice versa (USEPA, 2010). However, New York State Department of Health (2007) classified TCR values into: very low risk (TCR value $< 10^{-6}$), low risk (TCR value $< 10^{-6}$ but $< 10^{-4}$), moderate risk (TCR value $> 10^{-4}$ but $< 10^{-3}$), high risk (TCR value $> 10^{-3}$ but 10^{-1}) and very high risk (TCR value $\geq 10^{-1}$). In this study, the health endpoints for which the risks were computed include neurotoxicity, systemic toxicity, immunotoxicity, reproductive toxicity and developmental toxicity.

1.4 Results and Discussions

1.4.1 Concentration of OCPs in vegetables

The concentrations of OCPs in the vegetables are shown in Tables 1 and 2. There was no significant difference ($p > 0.05$) in the OCPs concentrations in each vegetable type across the four locations (Tables SM4-SM7) as well as among the different vegetables (Table SM8). However, the mean concentrations of $\sum 20$ OCPs in *Vernonia amygdalina* followed the order: Benin $>$ Port-Harcourt $>$ Warri $>$ Yenagoa while the mean concentrations of $\sum 20$ OCPs in *Telfairia occidentalis* followed the order: Port-Harcourt $>$ Benin $>$ Yenagoa $>$ Warri. For *Desmodium intortum cv* and *Talinum triangulare*, the order of $\sum 20$ OCPs concentrations were: Port-Harcourt $>$ Benin $>$ Yenagoa $>$ Warri and Benin $>$ Yenagoa $>$ Port-Harcourt $>$ Warri respectively (Table 1). The $\sum 20$ OCPs concentrations found in the vegetables varied from 11.6 to 37.7 ng g⁻¹ for *Vernonia amygdalina* 2.48 to 37.0 ng g⁻¹ for *Telfairia occidentalis*, 7.63-54.2 ng g⁻¹ for *Desmodium intortum cv* and 27.3 to 57.3 ng g⁻¹ for *Talinum triangulare*. On the average, the concentrations of OCPs in the vegetables obtained in this study followed the trend: *Talinum triangulare* $>$ *Desmodium intortum cv* $>$ *Vernonia*

amygdalina > *Telfairia occidentalis* (Table 2). The concentrations of OCPs obtained in this study were generally lower than their respective maximum residue limits (MRLs) stipulated by the European Union (2005; 2012) and World Health Organization (2010). The concentrations of OCPs obtained in our study with others reported in literature are shown in Table 3. The OCPs concentrations in vegetables from south-south, Nigeria were comparable to OCPs concentrations reported in some vegetables from Lagos, Nigeria (Oyeyiola *et al.*, 2017), Ghana (Bolor *et al.*, 2018), Togo (Kolani *et al.*, 2016) and Pakistan (Majeed *et al.*, 2020). However, the concentrations of OCPs obtained in this study were higher than those reported by Shoiful *et al.* (2013) for vegetables in Indonesia but lower than others (Adefemi *et al.*, 2018; Agnandji *et al.*, 2018; Adeleye *et al.*, 2019; Odewale *et al.*, 2021; Olutona *et al.*, 2021; Omeje *et al.*, 2021; Suleiman *et al.*, 2021).

1.4.2 Profiles of OCPs homologues in vegetables

The profiles of OCPs homologues in the vegetables are given in Figures 2 and 3 (a-e). On the average, the occurrence pattern of OCPs in the vegetables followed the order $\sum\text{HCHs} > \sum\text{Chlordanes} > \sum\text{DDTs} > \sum\text{Drins} > \sum\text{Endosulfans}$ for *Vernonia amygdalina*; $\sum\text{Drins} > \sum\text{HCHs} > \sum\text{Chlordanes} > \sum\text{Endosulfans} > \sum\text{DDTs}$ for *Telfairia occidentalis*, $\sum\text{Drins} > \sum\text{Chlordanes} > \sum\text{HCHs} > \sum\text{Endosulfans} > \sum\text{DDTs}$ for *Desmodium intortum* cv and $\sum\text{HCHs} > \sum\text{DDTs} > \sum\text{Chlordanes} > \sum\text{Drins} > \sum\text{Endosulfans}$ for *Talinum triangulare* (Figure 2). The mean $\sum\text{HCHs}$ concentrations ranged from 1.18 to 30.5 ng g⁻¹. *T. triangulare* from Port-Harcourt and *T. occidentalis* from Warri have the maximum and minimum $\sum\text{HCHs}$ concentrations respectively. The concentrations of $\sum\text{HCHs}$ in these vegetables followed the order *Talinum triangulare* > *Desmodium intortum* cv > *Vernonia amygdalina* > *Telfairia occidentalis*. Among the HCHs γ -HCH was the dominant congener in *V. amygdalina* and *D. intortum* cv accounting for 41 % and 42 % of the total $\sum\text{HCHs}$ respectively while α -HCH was the dominant HCH congener in *T. occidentalis* and *T. triangulare* constituting 65 % and

213 32 % respectively of the Σ HCHs (Figure 2a). The mean concentrations of Σ DDTs ranged
 214 from not detected to 19.5 ng g⁻¹. The maximum Σ DDTs concentration was found in *T.*
 215 *triangulare* from Benin whereas Σ DDTs was not found in *T. occidentalis* and *T. triangulare*
 216 from Yenagoa and *D. intortum cv* from Warri. The concentrations of Σ DDTs in these
 217 vegetables followed the order: *Talinum triangulare* > *Vernonia amygdalina* > *Desmodium*
 218 *intortum cv* > *Telfairia occidentalis*. Among the DDTs, DDE is the dominant congener. DDE
 219 constituted 86 %, 64 %, 100% and 63 % of the Σ DDTs for *V. amygdalina*, *T. occidentalis*, *D.*
 220 *intortum cv* and *T. triangulare* respectively (Figure 2b). Only DDE was detected in *D.*
 221 *intortum cv*. However, DDD was not detected in *V. amygdalina* and DDT was not detected in
 222 *T. occidentalis*. The mean concentrations of Σ Chlordanes in the vegetables from southern
 223 Nigeria ranged from 0.08 ng g⁻¹ for *Telfairia occidentalis* from Warri to 16.2 ng g⁻¹ for
 224 *Desmodium intortum cv* from Port-Harcourt. The concentrations of Σ Chlordanes in these
 225 vegetables followed the order: *Desmodium intortum cv* > *Talinum triangulare* > *Vernonia*
 226 *amygdalina* > *Telfairia occidentalis*. γ -chlordane (45 %), heptachlor epoxide (69 %),
 227 heptachlor (52 %) and α -chlordane were the predominant chlordane congeners in *Vernonia*
 228 *amygdalina*, *Telfairia occidentalis*, *Desmodium intortum cv* and *Talinum triangulare*
 229 respectively (Figure 2c). The mean Σ Endosulfan concentrations ranged from not detected to
 230 6.77 ng g⁻¹. The Σ endosulfan was not detected in *Vernonia amygdalina* from Warri and Port-
 231 Harcourt and *Desmodium intortum cv* from Yenagoa. The concentrations of Σ Endosulfans in
 232 these vegetables followed the order: *Talinum triangulare* > *Vernonia amygdalina* >
 233 *Desmodium intortum* > *Telfairia occidentalis*. Endosulfan II was not detected in any of the
 234 vegetables in this study while endosulfan sulphate was found only in *Telfairia occidentalis*
 235 from Warri and *Desmodium intortum cv* from Port-Harcourt. Endosulfan I was the
 236 predominant congener among the endosulfans constituting 71 to 100 % of the Σ endosulfans
 237 in these vegetables. The mean concentrations of Σ Drins varied between not detected in

Vernonia amygdalina from Yenagoa to 19.5 ng g⁻¹ in *Desmodium intortum* cv from Port-Harcourt. The concentrations of Σ Drins in these vegetables followed the order: *Desmodium intortum* cv > *Talinum triangulare* > *Telfairia occidentalis* > *Vernonia amygdalina*. Aldrin was the dominant congener among the Σ Drins in *Vernonia amygdalina*, *Telfairia occidentalis* constituting 94 % and 49 % respectively of the Σ Drins. Dieldrin was dominant in *Desmodium intortum* cv accounting for 49 % of the Σ Drins while endrin constituted 44 % of the Σ Drins in *Talinum triangulare* to be the dominant congener. Endrin aldehyde was found only in *Telfairia occidentalis* from Benin, *Talinum triangulare* from Benin and Port-Harcourt while endrin ketone was found only in *Telfairia occidentalis* from Warri.

1.4.3 Estimated Dietary Intake

The estimated EDI values of OCPs are displayed in supplemental material SM9. The EDI (in ng kg⁻¹ bw day⁻¹) from ingestion of the vegetables varied between 42.4 and 179, 4.6 to 107, 35.9 to 91.4, 8.2 to 45.7, 30.8 to 95.9 and 141 to 464 for Σ HCHs, Σ DDTs, Σ Chlordanes, Σ Endosulfans Σ Drins and Σ 20 OCPs respectively for children ingestion and 8.0 to 33.5, 0.9 to 20.0, 6.7 to 17.1, 1.5 to 8.6, 5.8 to 18.0 and 26.5 to 87.0 for Σ HCHs, Σ DDTs, Σ Chlordanes, Σ Endosulfans, Σ Drins and Σ 20 OCPs respectively for adults ingestion. The estimated daily intake obtained in this study were lower than the acceptable daily intake of 100, 500 and 10,000 ng kg⁻¹ bw day⁻¹ for Σ Drins, Σ Chlordanes and Σ DDTs respectively established by FAO/WHO Joint Meeting on Pesticide Residue (WHO, 2010).

1.4.4 Estimated non-carcinogenic and carcinogenic risks

The HI and TCR values of OCPs in the vegetables are displayed in Figure 4 and supplemental materials SM10 and SM11. The HI values for children and adults consuming the vegetables varied between 0.06 and 2.75 and from 0.01 to 0.69 respectively. The HI

values for children and adults were generally < 1 suggesting that non-cancer risk is absent from consuming these vegetables except for *Vernonia amygdalina*, *Telfairia occidentalis*, and *Desmodium intortum* cv from Port Harcourt for child exposure. The HI values obtained in this study were far lower than the HI values previously reported for OCPs in vegetables (Adefemi *et al.*, 2018; Adeleye *et al.*, 2019). Adefemi *et al.* (2018) and Adeleye *et al.* (2019) reported HI values of 15.8 to 54.1 and 68.916 to 116 respectively for children; and 7.91 to 27.2 and 19.182 to 32.35 for adults respectively.

The TCR values from children and adults consuming the vegetables varied between 5.96×10^{-5} and 1.08×10^{-3} and from 5.42×10^{-6} to 9.77×10^{-5} respectively. The TCR values were $> 1.0 \times 10^{-6}$ indicating that there is cancer risk for children and adults consuming the vegetables. Based on the NYSDOH classification, the TCR values for children fall into the moderate risk while those of adults fall into the low risk category. Heptachlor epoxide and aldrin contributed significantly to the HI and TCR values. The possible long term carcinogenic effects of OCPs exposure from consumption of the vegetables include skin, lung, liver, pancreas, prostate, breast, kidney and brain cancers. In this study, the health risk assessment was done following the assumptions that the concentrations of the OCPs remain unchanged during the whole exposure duration. It was also assumed that absorption and bioavailability rates are 100%. Thus, the results of the health risk assessment of OCPs in the vegetables should be considered as indicative and not to be interpreted as accurate prediction of observed health outcomes.

1.5 Source Identification of OCPs in the vegetables

1.5.1 Isomeric ratios

The isomeric ratios of OCPs used in source identification in the vegetables are shown in supplemental material SM12. For *Vernonia amygdalina*, the ratios of heptachlor

epoxide/heptachlor and DDT/ Σ DDTs indicated that the OCPs in the vegetables were from historical usage whereas other isomeric ratios indicated that the OCPs were from recent usage. For *Telfairia occidentalis*, the ratios of γ -HCH/ Σ HCHs and β -HCH/ γ -HCH indicated that the OCPs are from recent usage whereas the other isomeric ratios indicated that the OCPs were from historical usage. For *Desmodium intortum* cv, all the isomeric ratios suggested that the OCPs were from recent usage. For *Talinum triangulare*, ratios of α -HCH/ γ -HCH and γ -HCH/ Σ HCHs suggested that the OCPs were from recent usage whereas the other isomeric ratios suggested that the OCPs were from historical usage. This result implies and also confirmed that OCPs are still being imported and used in farming in Nigeria despite their banned.

1.5.2 Principal Component Analysis (PCA)

The PCA result of the OCPs in these vegetables is displayed in supplemental material SM13 and Figure SM1. Two factors were extracted and make up for 73.439 % of the OCPs data set. Factor 1 explained 39.743 % of the total variance with positive high loading on HCHs (.847) and endosulfans (.886) and moderate loading on DDTs. The association of these OCP homologues with component 1 indicates that they are from similar source and have related physicochemical properties (Kim and Smith, 2001; Devi *et al.*, 2013). Factor 2 explained 33.697 % of total variance in the OCPs dataset with high positive loading on chlordanes (.920) and drins (.887). The association of the chlordanes and drins in this component suggested that they originated from similar source(s) in these vegetables.

1.6 Conclusion

The results of this study have shown that the vegetables from south-south Nigeria were contaminated with OCPs but the concentrations of OCPs in the vegetables were below their respective MRLs. The HI values suggested a potential non-carcinogenic health risk for

children consumers and no potential risks for adult consumers of the vegetables. The TCR values indicated that children have moderate carcinogenic risk while adults have low carcinogenic risk from consuming the vegetables. The isomeric ratios and PCA results showed that OCPs in these vegetables from southern Nigeria originated from both historical and recent usage in the vegetable cultivation.

Acknowledgements

The authors hereby appreciate Dickson Okiti Sarah, Akio Serina, Chinedu Esther and Maxwell Osaye and Disi-Thompson Diseye for their assistance during the collection of samples.

Conflict of Interest

There is no conflicting interest among authors.

References

- Adefemi, S.O., Asaolu, S.S., Ibigbami, O.A., Orege, J.I., Azeez, M.A., Akinsola, A.F., 2018. Multi-residue levels of persistent organochlorine pesticides in edible vegetables: a human health risk assessment. *J. Agric. Food Chem.* 7, 143–152.
- Adeleye, A.O., Sosan, M.B., Oyekunle, A.O., 2019. Dietary exposure assessment of organochlorine pesticides in two commonly grown leafy vegetables in South-western Nigeria. *Heliyon*, 5, e01895 <https://doi.org/10.1016/j.heliyon.2019.e01895>

342 Adetunde, O.T., Mills, G.A., Oluseyi, T.O., Oyeyiola, A.O., Olayinka, K.O., Alo, B.I., 2018.
 343 Polycyclic aromatic hydrocarbon in vegetables grown on contaminated soils in a sub-
 344 saharan tropical environment – Lagos, Nigeria. *Polycycl. Aromat. Compd.*, DOI:
 345 10.1080/10406638.2018.1517807

346 Agnandji, P., Ayi-Fanou, L., Gbaguidi, M.A.N., Cachon, B.F., Hounha, M., Dikpo, M.T.,
 347 Cazier, F., Sanni, A., 2018. Distribution of organochlorine pesticides residues in
 348 *Solanum macrocarpum* and *Lactuca sativa* cultivated in south of Benin (Cotonou and
 349 Seme-Kpodji). *Am. J. Food Sci. Technol.*, 6, 19-25.

350 Akan, J.C., Jafiya, L., Chellube, Z.M., Mohammed, Z., Abdulrahman, F.I., 2014.
 351 Determination of some organochlorine pesticide residues in vegetable and soil
 352 samples from Alau dam and Gongulong agricultural sites, Borno State, North Eastern
 353 Nigeria. *Int. J. Environ. & Eco. Engr.* 8, 325–332.

354 Asogwa, E.U., Dongo, L.N., 2009. Problems associated with pesticides usage and
 355 applications in Nigeriacocoa production: A review. *Afr. J. Agric. Res.*, 4, 675-683.

356 Bai, J., Lu, Q., Zhao, Q., Wang, J., Gao, Z., Zhang, G., 2015. Organochlorine pesticides
 357 (OCPs) in wetland soils under different land uses along a 100-year chronosequence of
 358 reclamation in a Chinese estuary. *Sci. Rep.*, 5, 17624

359 Barlow, S.M., 2005. Agricultural chemicals and endocrine-mediated chronic toxicity or
 360 carcinogenicity. *Scand J Work Environ Health*, 31, 119–122.

361 Benson, N.U., Aruwajoye, O.I., 2011. Assessment of contamination by organochlorine
 362 pesticides in *Solanum lycopersicum* L. and *Capsicum annum* L: a market survey in
 363 Nigeria. *Afr. J. Environ. Sci. Technol.* 5, 437–442.

364 Bolor, V.K., Boadi, N.O., Borquaye, L.S., Afful, S., 2018. Human risk assessment of
 365 organochlorine pesticide residues in vegetables from Kumasi, Ghana. *J. Chem.*,
 366 <https://doi.org/10.1155/2018/3269065>.

367 Botella, B., Crespo J., Rivas, A., Cerrillo I., Olea-Serrano, M.F., Olea, N., 2004. Exposure of
 368 women to organochlorine pesticides in Southern Spain. *Environ Res.*, 96, 34-40.

369 Devi, N.L., Chakraborty, P., Shihua, Q., Zhang, G., 2013. Selected organochlorine pesticides
 370 (OCPs) in surface soils from three major states from the northeastern part of India.
 371 *Environ. Monit. Assess.* 185, 6667–6676

372 Emoyan, O.O., Peretiemo-Clarke, B.O., Tesi, G.O., Ohwo, E., 2021. Occurrence, origin,
 373 ecological and human health risks of organochlorine pesticides in soils from selected
 374 urban, suburban and rural storm water reservoirs. *Soil Sediment Contam.*,
 375 <https://doi.org/10.1080/15320383.2021.1913993>

376 European Union (EU) 2012. Pesticide Database, Pesticide Residues MRLs, Directorate
 377 General for Health & Consumers, European Union, City of Brussels, Belgium, 2012.

378 European Union (EU). 2005. European Union Commission Regulation (2005) No 396/2005
 379 of the European Parliament and of the Council of 23 February 2005 on Maximum
 380 Residue Levels of Pesticides in or on Food and Feed of Plant and Animal Origin and
 381 Amending Council Directive 91/414/E-Text with EEA Relevance.

382 FAO. 2005. *AOSTAT statistical database*. Rome, Food and Agricultural Organization

FAOSTAT 2020. Statistics of production crops and pesticide trade for all area of the world. Available at: <http://www.fao.org/0e3121c5-b1bf-4689-9efb-c431428f61a8> (accessed 15 May 2022).

Gereslassie, T., Workineh, A., Atieno, O.J., Wang, J., 2019. Determination of occurrences, distribution, health impacts of organochlorine pesticides in soils of Central China. *Int. J. Environ. Res. Public Health*, 16, 146

Ibrahim, E.G., Yakubu, N., Nnamonu, L., Yakubu, J.M., 2018. Determination of organochlorine pesticide residues in pumpkin, spinach and sorrel leaves grown in Akwanga, Nasarawa State, Nigeria. *J. Environ. Prot.* 9, 508–515.

Kim, J. H., Smith, A., 2001. Distribution of organochlorine pesticides in soils from South Korea. *Chemosphere*, 43, 137–140

Kolani, L., Mawussi, G., Sanda, K., 2016. Assessment of organochlorine pesticide residues in vegetable samples from some agricultural areas in Togo. *Am. J. Anal. Chem.* 7, 32–341.

Majeed, R., Khan, M.A., Fatima, S.U., Khan, M.A., Shaukat, S.S., 2020. Occurrence of organochlorine pesticide (OCPs) residues in farm vegetables cultivated through untreated wastewater in Karachi. *Int. J. Biol. Biotech.*, 17, 513-516

New York States Department of Health (NYS DOH) 2007. Hopewell precision area contamination: Appendix C-NYS DOH, In: Procedure for evaluating potential health risks for contaminants of concern.

Njoku, K.L., Ezech, C.V., Obidi, F.O., Akinola, M.O., 2017. Assessment of pesticide residue levels in vegetables sold in some markets in Lagos State, Nigeria. *Nig. J. Biotech.* 32, 53–60.

Odewale, G.O., Sosan, M.B., Oyekunle, J.A.O., Adeleye, A.O., 2021. Assessment of systemic and carcinogenic health risks of persistent organochlorine pesticide residues in four fruit vegetables in southwestern Nigeria. *Br. Food J*, DOI 10.1108/BFJ-05-2021-0573

Olutona, G.O., Fakunle, I.A., Adegbola, R.A., 2021. Detection of organochlorine pesticides residue and trace metals in vegetables obtained from Iwo market, Iwo, Nigeria. *Int. J. Environ. Sci. Technol.*, <https://doi.org/10.1007/s13762-021-03431-x>

Omeje, K.O., Ezema, B.O., Okonkwo, F., Onyishi, N.C., Ozioko, J., Rasag, W.A., Sardo, G., Okpala, C.O.R. 2021. Quantification of heavy metals and pesticide residues in widely consumed Nigerian food crops using atomic absorption spectroscopy (AAS) and gas chromatography (GC). *Toxins*, 13, 870. <https://doi.org/10.3390/toxins13120870>

Oyekunle, J.A.O., Ogunfowokan, A.O., Torto, N., Akanni, M.S. 2011. Determination of organochlorine pesticides in the agricultural soil of Oke-Osun farm settlement, Osogbo, Nigeria. *Environ. Monit. Assess.*, 177, 51–61

Oyeyiola, A.O., Fatunsin, O.T., Akanbi, L.M., Fadahunsi, D.E., Moshood, M.O., 2017. Human health risk of organochlorine pesticides in foods grown in Nigeria. *J. Health Pollut.* 7, 63–70.

PAN. 2007. Pesticides Action Network Strategic Assessment of the Status of POPs Pesticides Trading in South Western Nigeria (p. 61). Africa Stockpile Programme. www.africastockpiles.net

Qin, F., Yang, Y., Wang, S.T., Dong, Y.N., Xu, M.X., Wang, Z.W., Zhao, J.X., 2019. Exercise and air pollutants exposure: A systematic review and meta-analysis. *Life Sci.*, 218, 153–164.

- Shoiful, A., Fujita, H., Watanabe, I., Honda, K., 2013. Concentrations of organochlorine pesticides (OCPs) residues in foodstuffs collected from traditional markets in Indonesia. *Chemosphere*, 90, 1742–1750
- Stadlmayr, B., Charrondiere, U.R. and Burlingame, B., 2013. Development of a regional food composition table for West Africa. *Food Chem.*, 140, 443–446.
- Sulaiman, M., Maigari, A., Ihedioha, J., Lawal, R., Gimba, A., Shuaibu, A., 2021. Levels and health risk assessment of organochlorine pesticide residues in vegetables from Yamaltu area in Gombe, Nigeria. *Fr. -Ukr. j. chem.*, 9, 19–31
- Tesi, G.O., Iniaghe, P.O., Lari, B. Obi-Iyeke, G., Ossai, J.C., 2021. Polycyclic aromatic hydrocarbons (PAHs) in leafy vegetables consumed in southern Nigeria: concentration, risk assessment and source apportionment. *Environ. Monit. Assess.* 193, 443 <https://doi.org/10.1007/s10661-021-09217-5>
- Tesi, J.N., Tesi, G.O., Ossai, J.C., Agbozu, I.E. 2020. Organochlorine pesticides (OCPs) in agricultural soils of southern Nigeria: Spatial distribution, source identification, ecotoxicological, and human health risks assessment. *Environ. Forensics* <https://doi.org/10.1080/15275922.2020.1850570>
- Thompson, D., Agbugba, I.K., 2013. Marketing of tropical vegetable in Aba area of Abia State Nigeria. *J Agric Econ. Dev.*, 2, 272–279
- USEPA (United States Environmental Protection Agency) 2009. Risk assessment guidance for superfund. Volume 1: Human Health Evaluation Manual (F, supplemental guidance for Inhalation Risk Assessment) EPA/540/R/070/002, Office of Superfund Remediation and Technology Innovation, Washington, DC.
- USEPA (United States Environmental Protection Agency) 2010. Regional screening levels (RSL) summary tables. <http://www.epa.gov/risk/risk-based-screening-tablegeneric-tables>. (Accessed 21 April, 2021).
- WHO (World Health Organization) 2010. Pesticide residue in food and feed. <<http://www.codexalimentarius.net/pestres/data/pesticides/index.html>>.
- WHO (World Health Organization) (2012a). Country assignments to the 17 proposed GEMS/food consumption cluster diets of World Health Organization. Available at: http://www.who.int/foodsafety/chem/cluster_diets_2012.pdf.
- WHO (World Health Organization) (2012b). GEMS/Food consumption database for all 17 GEMS/Cluster diets of World Health Organization 8pp. Available at: http://www.who.int/nutrition/landscape_analysis/nlis_gem_food/en
- Yang, D., Qi, S. H., Zhang, J. Q., Tan, L. Z., Zhang, J. P., Zhang, Y., Xu, F., Xing, X. L., Hu, Y., Chen, W., Yang, J. H., Xu, M.H., 2012. Residues of organochlorine pesticides (OCPs) in agricultural soils of Zhangzhou City, China. *Pedosphere* 22, 178–179.
- Yucra, S., Rubio, J., Gasco, M., Gonzales, C., Steenland, K. & Gonzales, G. F. 2006. Semen quality and reproductive sex hormones levels in Peruvian pesticide sprayers. *Int J Occup Environ Health*, 12, 355–361

Table 1: Mean OCPs concentrations (ng g⁻¹) in vegetables with respect to locations

<i>Vernonia amygdalina</i>					<i>Telfairia occidentalis</i>				<i>Desmodium intortum cv</i>				<i>Talinum triangulare</i>				EU/WHO MRLs
	WR	BN	YG	PH	WR	BN	YG	PH	WR	BN	YG	PH	WR	BN	YG	PH	
α-HCH	ND	0.89± 0.5	0.94±0.5	ND± 0.04	0.07±0.05	0.80±0.08	0.73±0.03	8.37±0.05	0.59±0.07	ND	3.14±0.07	2.02±0.05	3.57±0.02	8.85±0.03	8.25±0.04	ND	10
β-HCH	1.70±0.5	7.12± 0.09	ND	ND± 0.01	0.56±0.08	0.24±0.38	ND	ND	0.69±0.05	2.39±0.01	1.94±0.03	6.27±0.06	1.94±0.06	2.41±0.09	6.37±0.03	8.25±1.05	10
γ-HCH	ND	9.67± 1.00	2.24± 0.09	ND± 0.09	ND	1.36±0.80	0.15±0.01	ND	1.76±0.02	6.68±0.06	2.68±0.08	1.96±0.02	ND	4.44±0.06	8.45±0.09	0.03±1.01	10
σ-HCH	1.60±0.09	ND	ND	6.19± 0.5	0.55±0.6	1.07±0.09	0.95±0.07	0.57±0.5	0.26±0.09	ND	ND	0.71±0.04	2.37±0.09	1.50±0.08	7.45±0.06	0.03±0.06	10
ΣHCHs	2.30±1.20	17.68±1.00	3.18± 0.90	6.19± 1.00	1.18±0.80	3.47±1.05	1.83±0.80	8.94±0.40	3.30±0.50	9.07±0.66	7.76±0.80	10.96±0.60	8.06±0.90	17.20±0.93	30.52±1.50	9.15±0.78	
p,p'-DDE	2.30±0.6	7.09± 0.05	ND	0.18± 0.05	ND	0.08±0.03	ND	0.98±0.60	ND	1.00±0.50	0.27±0.03	0.90±0.04	8.34±0.06	8.02±0.06	ND	8.14±0.08	50
p,p'-DDD	ND	ND	ND	ND	0.07±0.01	ND	ND	0.53±0.01	ND	ND	ND	ND	2.72±0.09	8.25±0.07	ND	ND	50
p,p'-DDT	ND	ND	1.70± 0.50	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	3.27±0.08	ND	ND	50
Methoxychlor	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
ΣDDTs	2.80±0.55	7.09± 0.9	1.70± 1.01	0.18±0.70	0.07±0.90	0.08±0.60	ND	1.51±0.05	ND	1.00±0.11	0.27±1.05	0.90±1.35	11.06±1.04	19.25±1.05	ND	8.14±1.05	
α-Chlordane	ND	ND	1.54± 0.02	5.24±0.08	0.08±0.05	ND	ND	2.08±0.04	ND	3.06±0.04	ND	ND	2.00±1.05	ND	1.15±0.06	10.24±0.08	
γ-Chlordane	4.37± 1.10	1.66± 0.9	ND	6.16±0.05	ND	ND	ND	0.96±0.02	0.65±0.2	4.22±0.1	ND	ND	3.71±0.03	0.75±0.06	ND	0.61±0.01	
Heptachlor	ND	0.12± 0.5	2.70± 0.90	0.93±0.02	ND	0.16±0.90	0.78±0.06	ND	0.72±0.05	7.60±0.06	ND	8.94±0.07	1.56±0.08	ND	1.93±0.09	0.53±0.05	10
Heptachlor epoxide	ND	2.50± 1.00	ND	1.88±0.04	ND	1.55±0.03	0.54±0.08	6.91±0.05	0.26±0.06	ND	0.51±0.50	7.27±1.05	0.22±0.04	1.71±0.05	2.72±0.04	2.18±0.04	10
ΣChordanes	4.37±0.9	4.28± 1.50	4.24± 0.70	14.21±0.50	0.08±0.25	1.71±1.02	1.32±0.55	9.95±0.55	1.63±0.90	14.88±0.89	0.51±0.68	16.21±0.94	7.49±0.88	2.46±1.45	5.80±0.83	13.56±0.92	
Endosulfan I	ND	6.77± 0.77	2.48± 0.6	ND	ND	0.91±0.04	1.54±0.04	0.51±0.08	0.87±0.09	0.25±0.05	ND	1.13±0.07	0.68±0.06	5.78±0.03	6.02±0.08	4.15±0.07	50
Endosulfan II	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	50
Endosulfan sulfate	ND	ND	ND	ND	0.02±0.60	ND	ND	ND	ND	ND	ND	5.49±0.08	ND	ND	ND	ND	50
ΣEndosulfans	0± 0.5	6.77± 0.9	2.48± 0.2	0±0.5	0.02±1.05	0.91±0.5	1.54±0.70	0.51±0.60	0.87±1.05	0.25±1.05	0±0.50	6.62±0.99	0.68±1.05	5.78±1.50	6.02±1.04	4.15±1.00	
Aldrin	3.47±1.50	1.92± 0.3	ND	5.17±0.07	ND	ND	ND	8.00±0.03	1.83±0.50	ND	ND	4.07±0.08	0.01±0.02	ND	ND	3.43±0.06	10
Dieldrin	ND	ND	ND	ND	1.10±1.00	ND	ND	1.50±0.08	ND	7.85±0.04	ND	9.32±0.50	ND	ND	ND	ND	10
Endrin	ND	ND	ND	0.65±0.4	ND	ND	ND	6.60±0.01	ND	4.07±0.01	1.61±0.09	6.13±0.02	ND	7.49±0.01	0.94±1.05	ND	10
Endrin aldehyde	ND	ND	ND	ND	ND	0.20±0.07	ND	ND	ND	ND	ND	ND	ND	4.77±0.08	ND	2.50±0.08	10
Endrin Ketone	ND	ND	ND	ND	0.03±0.01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
ΣDrins	3.47±1.40	1.92± 0.80	0± 0.3	5.82±1.10	1.13±0.55	0.2±1.40	0.82±1.03	16.1±1.50	1.83±1.22	11.92±1.30	1.61±1.06	19.52±1.50	0.01±0.03	12.26±0.63	0.94±1.50	5.93±1.36	
Σ20 OCPs	12.94±1.05	37.7± 1.50	11.6± 0.80	26.40±0.90	2.48±1.05	6.37±1.11	5.51±0.90	37.01±0.99	7.63±1.05	37.12±1.00	10.15±1.50	54.21±0.99	27.30±0.68	57.24±1.22	43.28±1.04	40.93±1.50	

WR = Warri; BN = Benin; YG = Yenagoa; PH = Port-Harcourt

Table 2: Summary statistics of OCPs concentrations (ng g⁻¹) in the vegetables

	<i>Vernonia amygdalina</i> (n=20)		<i>Telfairia occidentalis</i> (n=20)		<i>Desmodium intortum cv</i> (n=20)		<i>Talinum triangulare</i> (n=20)	
	Mean± SD	RANGE	Mean± SD	RANGE	Mean± SD	RANGE	Mean± SD	RANGE
α-HCH	0.46±0.53	ND-0.94	2.49±3.93	0.07-8.37	1.44± 1.42	ND-3.14	5.21± 4.15	ND-8.85
β-HCH	1.96±3.46	ND-7.12	0.2±0.27	ND-0.56	2.82± 2.41	0.69-6.27	4.74± 3.07	1.94-8.25
γ-HCH	2.98±4.58	ND-9.67	0.38±0.66	ND-1.36	3.27± 2.31	1.76-6.68	3.23± 3.05	ND-8.45
σ-HCH	1.95±2.93	ND-6.19	0.79±0.26	0.55-1.07	0.24± 0.33	ND-0.71	3.05± 3.00	0.87-7.45
ΣHCHs	7.34±7.09	2.3-17.68	3.86±3.52	1.18-8.94	7.77± 3.26	3.3-11.0	16.2± 10.4	8.06-30.5
p,p'-DDE	2.52±3.31	ND-7.09	0.27±0.48	ND-0.98	0.54± 0.49	ND-1.00	6.13± 4.09	ND-8.34
p,p'-DDD	ND	ND	0.15±0.26	ND-0.53	ND	ND	2.74± 3.89	ND-8.25
p,p'-DDT	0.43±0.85	ND-1.70	ND	ND	ND	ND	0.82± 1.64	ND-8.27
Methoxychlor	ND	ND	ND	ND	ND	ND	ND	ND
ΣDDTs	2.94±2.97	0.18-7.08	0.42±0.73	ND-1.51	0.54± 0.49	ND-1.00	9.69± 8.07	ND-3.27
α-Chlordane	1.7±2.47	ND-5.24	0.54±1.03	ND-2.08	0.77± 1.53	ND-3.06	3.35± 4.67	ND-19.5
γ-Chlordane	3.05±2.75	ND-6.16	0.24±0.48	ND-0.96	1.22± 2.02	ND-4.22	1.27± 1.66	ND-10.2
Heptachlor	0.94±1.25	ND-2.70	0.24±0.37	ND-0.78	4.32± 4.61	ND-8.94	1.01± 0.89	ND-3.71
Heptachlor epoxide	1.1±1.29	ND-2.50	2.25±3.17	ND-6.91	2.01± 3.51	ND-7.27	1.71± 1.07	0.22-2.72
ΣChordanes	6.78±4.96	4.24-14.2	3.27±4.51	0.08-9.95	8.31± 8.39	0.51-16.2	7.33± 4.65	2.46-13.6
Endosulfan I	2.31±3.19	ND-6.77	0.74±0.65	ND-1.54	0.56± 0.53	ND-1.13	4.16± 2.46	0.68-6.02
Endosulfan II	ND	ND	ND	ND	ND	ND	ND	ND
Endosulfan sulfate	ND	ND	0.01±0.01	ND-0.02	1.37± 2.75	ND-5.49	ND	ND
ΣEndosulfans	2.31±3.19	ND-6.77	0.75±0.64	0.02-1.54	1.94± 3.14	ND-6.62	4.16± 2.46	0.68-6.02
Aldrin	2.64±2.20	ND-5.17	2.21±3.88	ND-8.00	1.48± 1.93	ND-4.07	0.86± 1.71	ND-3.43
Dieldrin	ND	ND	0.65±0.77	ND-1.50	4.29± 4.99	ND-9.32	ND	ND
Endrin	0.16±0.33	ND-0.65	1.65±3.30	ND-6.60	2.95± 2.70	ND-6.13	2.11± 3.62	ND-7.49
Endrin aldehyde	ND	ND	0.05±0.10	ND-0.20	ND	ND	1.82± 2.29	ND-4.77
Endrin Ketone	ND	ND	0.01±0.02	ND-0.03	ND	ND	ND	ND
ΣDrins	2.8±2.46	ND-5.82	4.56±7.70	0.2-16.10	8.72± 8.66	1.61-19.52	4.79± 5.62	0.01-12.3
Σ20 OCPs	22.2±12.3	11.6-37.7	12.8±16.2	2.48-37.0	27.3± 22.4	7.63-54.2	42.2± 12.3	27.3-57.2

Table 3: Comparison of OCPs concentrations (ng g⁻¹) in vegetables from south-south Nigeria with others reported in literature

Location	Vegetable types	No. of samples	No. of OCPs compounds analysed	ΣHCHs	ΣDDTs	ΣChlordanes	ΣEndosulfans	ΣDrins	ΣOCPs	References
South-south, Nigeria	<i>Vernonia amygdalina</i> , <i>Telfairia occidentalis</i> , <i>Desmodium intortum</i> cv and <i>Talinum triangulare</i>	80	20	1.18-30.5	ND-7.08	0.08-16.2	ND-6.77	ND-19.5	2.48-57.3	This Study
Gombe, Northern Nigeria	Tomato, onion, pepper, and chili pepper	36	10	242-329		98.0-124	214-286	257-373	257-492	Suleiman <i>et al.</i> (2021)
Nsukka and Enugu, South-east, Nigeria	Fluted pumpkin, <i>Amaranthus</i> leaf, waterleaf, and scent leaf	NA	9	<DL-504	<DL-221	<DL-733	<DL-54.8	<DL-771	461-1255	Omeje <i>et al.</i> (2021)
Nsukka and Enugu, South-east, Nigeria	Okro, cucumber, carrot, and watermelon	NA	9	<DL-517	<DL-80.8	<DL-733	<DL-73.3	<DL-242	144-1255	Omeje <i>et al.</i> (2021)
Iwo, Southwest, Nigeria	Carrot, Onion, Cabbage, Garlic and Ginger	50	17	<DL-950	<DL-4898	<DL-2700	<DL-4320	<DL-44200	<DL-14150	Olutona <i>et al.</i> (2021)
Southwest, Nigeria	Carrot, Cucumber, Tomato and watermelon	144	9	-	-	903-1368	1438-4256	1986-4182	4329-9508	Odewale <i>et al.</i> (2021)
Lagos, Southwest, Nigeria	Cabbage, cameroun, green and chilli peppers, carrot, lettuce, tomato and scotch honnet	NA	13	1.03-2.88	10.75	0.73-1.57	0.09-0.29	0.4-2.06	14.8-131	Oyeyiola <i>et al.</i> (2017)
Southwest, Nigeria	Fluted pumpkin and <i>Amaranthus</i>	32	9	-	-	103-349	788-3449	3999-5698	5366-9496	Adeleye <i>et al.</i> (2019)
Ekiti Southwest, Nigeria	<i>Senecio bialfrae</i> (Wet season)	8	17	ND-4.0	41.0-166	ND-68.0	ND-245	ND-299	136-932	Adefemi <i>et al.</i> (2018)
Ekiti Southwest, Nigeria	<i>Senecio bialfrae</i> (Dry season)	8	17	ND-91.0	63.0-379	29.0-299	ND-286	20.0-246	189-908	Adefemi <i>et al.</i> (2018)
Cotonou and Seme-Kpodji, Benin Republic	<i>Lactuca sativa</i> L and <i>Solanum macrocarpum</i> L	31	10	66-415	44-127	-	310-2313	5.0-66.0	569-2780	Agnandji <i>et al.</i> (2018)
Kumasi,	Cabbage, Lettuce and	15	8	0.78-9.54	175-190	2.29	0.60-28.47	25.8	-	Bolor <i>et al.</i> (2018)

Ghana	Onion									
Togo	Tomato, cabbage and lettuce	130	19	<0.001-93.8	<0.001-1.52	<0.001-1.59	-	<0.001-0.70	<0.001-97.9	Kolani <i>et al.</i> (2016)
Indonesia	Carrot, potato, cucumber, corn and onions	21	21	<0.05	<0.03	<0.03	-	<0.03	<0.03	Shoiful <i>et al.</i> (2013)
Pakistan	Tomato, brinjal, gourd, okra and spinash	45	-	0.06-0.72	0.09-2.53	-	0.04-1.17	-	-	Majeed <i>et al.</i> (2020)




Figure 1: Map of Study area showing the locations of the markets




Figure 2: Occurrence pattern of OCPs homologues in the vegetables (a) *Vernonia amygdalina* (b) *Telfairia occidentalis* (c) *Desmodium intortum* cv (d) *Talinum triangulare*




Figure 3a: Percentage composition of HCHs congeners in the vegetables




Figure 3b: Percentage composition of DDTs congeners in the vegetables




Figure 3c: Percentage composition of Chlordane congeners in the vegetables




Figure 3d: Percentage composition of Endosulfans congeners in the vegetables




Figure 3e: Percentage composition of Drins congeners in the vegetables




Figure 4a: Hazard index values of OCPs in the vegetables (The yellow line behind the bars indicates the benchmark value)




Figure 4b: Total cancer risk values of OCPs in the vegetables (The yellow line behind the bars indicates the acceptable risk value of 1×10^{-6})