

**Energy expenditure of female international standard soccer players: a doubly labelled
water investigation**

James C. Morehen^{1,2}, Christopher Rosimus², Bryce P. Cavanagh², Catherine Hambly³, John
R. Speakman³, Kirsty J. Elliot-Sale⁴, Marcus P. Hannon¹ and James P. Morton¹

¹Research Institute for Sport and Exercise Sciences (RISES)
Liverpool John Moores University
Byrom Street
Liverpool
L3 3AF
UK

²The Football Association,
London
UK

³Institute of Biological and Environmental Sciences
University of Aberdeen
Aberdeen
UK

⁴Musculoskeletal Physiology Research Group,
Sport Health and Performance Enhancement Research Centre,
Nottingham Trent University,
Nottingham, UK

Address for correspondence:

Dr James C Morehen
Research Institute for Sport and Exercise Sciences (RISES)
Liverpool John Moores University
Byrom Street
Liverpool
L3 3AF
Email: j.c.morehen@ljmu.ac.uk

Abstract

Purpose: To quantify total daily energy expenditure (TEE) of international adult female soccer players. **Methods:** Twenty-four professional players were studied during a twelve-day period where they participated in an international training camp (also inclusive of two competitive games) representing the English national team. TEE was assessed via the doubly labelled water (DLW) method during the full 12 days as well as the initial 4-day period prior to game one. Energy intake (EI) was also assessed (via weighed food analysis) during the initial 4-day period to permit estimation of energy availability (EA). **Results:** Mean TEE did not differ ($P=0.31$) between the 12-day (2693 ± 432 kcal.day⁻¹; range: 2105-3507; 54 ± 6 kcal.kg⁻¹ fat free mass, FFM) versus the 4-day assessment period (2753 ± 359 kcal.day⁻¹; range: 1942-3280; 56 ± 8 kcal.kg⁻¹ FFM). Mean four-day EI was 1923 ± 357 kcal.day⁻¹ (range: 1639-2172) and mean activity energy expenditure was 1069 ± 278 kcal.day⁻¹ (range: 155-1549 kcal.day⁻¹). When assessed for estimated EA, 88% of players were categorised with low EA status according to the threshold of <30 kcal.kg⁻¹ FFM. Mean daily carbohydrate intake equated to 3.3 ± 0.7 g.kg⁻¹ body mass. **Conclusion:** When compared with previously published data from adult male players, we demonstrate that the relative daily energetic requirements of engaging in professional soccer training and match play is comparable between sexes. From a practical perspective, data suggest that practitioners should likely focus education and behaviour change strategies on “fuelling” for match play and training to optimise both player health and performance.

Keywords: carbohydrate, energy availability, RED-S

Introduction

In adult male professional soccer players, the physical demands of both match play (1–3) and training (4–6) are well documented. Such data typically demonstrate that the absolute loads completed in training are lower than those experienced in match play, as is the case for total distance (<7 km vs. ~10-13 km), high-speed running distance (<300 m vs. >900 m), sprint distance (<150 m vs. >200 m), and average speed (<80 m/min vs. ~100-120 m.min⁻¹) (7–9). When assessed during a typical in-season weekly micro-cycle comprising one or two games, outfield professional players typically expend 3000-4000 kcal.d⁻¹ (40-60 kcal.kg⁻¹ fat free mass, FFM), as quantified using the gold standard doubly labelled water method (9–11). Accordingly, evidence-based guidelines for the recommended energy and macronutrient intake to support both daily training and match play have recently been published (12). In this regard, it is suggested that daily carbohydrate (CHO) intake should equate to 3-8 g.kg⁻¹ body mass to allow for flexibility between rest days, training days and match days.

In contrast to adult male players, the energetic requirements and external training loads completed by elite female players are not as well understood (13–18). This is of specific interest given recent reports documenting the prevalence of low energy availability (LEA, defined as <30 kcal.kg⁻¹ FFM per day) in female professional players from the English Women's Super League (13). Indeed, these researchers observed that between 50-70% of players were classified with LEA status on both match day and "heavy" training days where daily activity energy expenditure was >700 kcal.d⁻¹, as estimated by global positioning systems (GPS). Analysis of self-reported energy intakes (EI) also demonstrated that these players consumed a consistent daily CHO intake of 3-3.5 g.kg⁻¹ body mass, thereby failing to adjust daily CHO intake in accordance with alterations to training load or in preparation for match play. Such data build on previous observations that female players apparently "under-fuel" in relation to

daily CHO intake (14–17). Given that 80% and 69% of type 1 and II muscle fibres from elite female players are classified as empty or almost empty of muscle glycogen immediately post-match play (18), such relative CHO intakes are likely sub-optimal in relation to promoting physical performance.

The reported prevalence of LEA is of particular concern given the potential for players to develop negative symptoms associated with the Female Athlete Triad (19,20) or Relative Energy Deficiency in Sport (RED-S) models (21). Nonetheless, despite previous assessments of activity energy expenditure and energy availability (EA) in such populations (13,15,16,22), it remains difficult to prescribe evidence based nutritional guidelines owing to the indirect methodologies employed to quantify daily *total* energy expenditure (TEE) (e.g., activity diaries and accelerometry which may under- or over-estimate non-exercise activity). In this regard, the doubly labelled water (DLW) method is the gold standard method of assessing total daily energy expenditure in free-living conditions *in vivo* (23). Importantly, this non-invasive method allows for an assessment of energy expenditure over a 7-14 day period (i.e. a typical in-season micro-cycle) without interfering in day-to-day activities such as soccer training or match play (23).

Accordingly, the primary aim of the present study was to therefore assess TEE of female soccer players via the gold standard DLW method. To this end, we studied 24 English female soccer players during a twelve-day period where players participated in an international training camp (also inclusive of two competitive games) representing the English national team. As a secondary measure, we also assessed energy intake (via weighed food analysis) during the initial four days of the assessment period to allow for an estimation of energy availability (EA). Given that this cohort represent players of the highest standard, it is hoped that these data may

provide a platform for which to develop evidence based nutritional guidelines that optimise the health and performance of female players.

Methods

Participants

Twenty-four female professional international soccer players volunteered to take part in the study. Cohort participant characteristics (also categorised according to playing position) are presented in Table 1. All players remained injury free for the duration of the study. All experimental procedures and associated risks were explained to players and written informed consent was obtained. The study was conducted according to the Declaration of Helsinki and was approved by the University Ethics Committee of Liverpool John Moores University.

Overview of study design

An overview of the experimental protocol is shown in Figure 1. All players completed a 9-day international training camp in November 2019 comprising 4 training days, 1 rest day, 2 travel days and 2 match days. Players completed the training prescribed by the national team's coaching staff and were available for team selection to play in 2 competitive international matches on days 5 (home game) and 8 (away game) during the study period. Three players did not play in either match and where appropriate, these players' data are not reported (indicated accordingly). TEE was assessed during a 12-day (9-day camp followed by 3-days at home) and 4-day assessment period using the DLW method whilst energy intake was also assessed during the 4 days prior to match one. TEE was assessed over 12 days (as opposed to 9-days) due to logistical challenges of urine collection on days 9 to 11 of the study. Players completed the second international football match abroad in Croatia on day 8. On day 9, players travelled back to the UK and were then driven from the airport to their homes. This resulted in no

opportunity to collect urine samples on this day. It was decided between international staff and domestic club staff that players were to rest at home on day 10 and 11 without any interruptions. On day 12, players arrived back at their respective clubs for duty, allowing a final urine sample to be collected. External loading was quantified from all pitch-based training sessions and games. To compare data across time, days are expressed in proximity to the match e.g., one day before the game is referred to as match day (MD) minus one (i.e., MD-1) whereas the day after the game is referred to as MD+1 etc.

Baseline measures

Due to logistical issues associated with player availability, body composition was assessed for 18 players only, occurring 2-4 weeks prior to the training camp via whole-body dual-energy X-ray absorptiometry (DXA) (Hologic QDR Series, Discovery A, Bedford, MA, USA), where the effective radiation dose was 0.01 mSv per person. All scans were performed and analysed by the same trained operator in accordance with best practice procedures (24). Resting metabolic rate (RMR) was estimated for each player using a recent female athlete specific predictive equation (25). This equation ($RMR = 120.81 + (4.88 \times \text{Stature}[\text{cm}]) + 8.24 \times \text{FFM}[\text{kg}] + (5.71 \times \text{Age}[\text{years}])$) was selected as it was developed using healthy female athletes of a similar age-range and FFM to those in the present study. On the morning of day 1 of the training camp, all players (i.e. n=24) were assessed for body mass and stature. Under standardised conditions (>8 hours overnight fast), measurement of stature (SECA, model-217, Hamburg Germany) and body mass (SECA, model-875, Hamburg, Germany) were measured to the nearest 0.1 cm and 0.1 kg, respectively according to the International Society for the Advancement of Kinanthropometry (ISAK) guidelines (26) by an ISAK Level-1 practitioner.

Quantification of external training and match load

The decision to wear GPS units during training was left to the players (goalkeepers do not wear these units). As such, thirteen outfield players who completed all training sessions and matches wore the same portable global GPS units (Apex, STATSports, Newry, Northern Ireland) for all pitch-based training sessions and both matches. Pitch-based sessions were monitored using the GPS units as previously described in professional soccer players (4,27,28). The GPS unit was placed inside a custom-made manufacturer provided vest (Apex, STATSports, Newry, Northern Ireland) that held the unit on the upper back between both scapulae, allowing clear exposure of the GPS antennae to acquire a clear satellite connection. External load variables selected for analysis from the training and match data were duration of activity (min), total distance covered (km) and high-speed running (defined as >5.30 to 6.30 m.s^{-1} , >19.08 to 22.68 km.h^{-1}).

Measurement of energy expenditure using the DLW method

Twenty-four players were available for assessment of TEE. Energy expenditure was determined via the DLW method (the gold standard method of measuring energy expenditure in free-living conditions) which we have previously used in professional team sport athletes (9,11,29). During the evening of day zero, between the hours of 18:00-20:00, players provided a background urine sample. Players then consumed a single bolus oral dose weighed to four d.p. of deuterium (^2H) and oxygen (^{18}O) stable isotopes in the form of water ($^2\text{H}_2^{18}\text{O}$), with a desired enrichment of 10% ^{18}O and 5% $^2\text{H}_2$ using the calculation:

$$\text{Dose (mL)} = 0.65(\text{body mass, g}) \times \text{DIE} / \text{IE},$$

Where 0.65 is the approximate proportion of the body comprised of water, DIE is the desired initial enrichment ($\text{DIE} = 618.923 \times \text{body mass (kg)}^{-0.305}$) and IE is the initial enrichment (10%)

100,000 parts per million (30) dosed according to body weight two-to-three weeks prior to the national camp. To ensure the whole dose was administered, participants were observed consuming each bolus dose and each glass vial was refilled with additional water which players were asked to consume. Time of dosing was recorded. Isotopes were purchased from Sercon (Cheshire, UK).

During the morning of day one (07:00-10:00), body mass was assessed (SECA, model-875, Hamburg, Germany), and participants were asked to provide a urine sample, collected in a 50 ml tube. This allowed initial isotope enrichment to be determined following total body water equilibrium (30). Thereafter, body mass was collected during the morning of day two, three, four, five, six and 12 and urine samples (second pass of the day) were collected on day two, three, four, five, six, seven, eleven and twelve (in line with logistical constraints), to determine elimination rates of both isotopes via the multi-point method (23).

For the DLW analysis, urine was encapsulated into capillaries, which were then vacuum distilled (31), and water from the resulting distillate was used. This water was analysed using a liquid water analyser (Los Gatos Research; (32)). Samples were run alongside three laboratory standards for each isotope and three International standards (Standard Light Artic Precipitate, Standard Mean Ocean Water and Greenland Ice Sheet Precipitation; (30,33)) to account for machine day to day variation and correct delta values to parts per million. Isotope elimination rates were converted to EE using an updated two-pool model equation (34) and a mean calculated food quotient of 0.85 ± 0.2 . The results from the energy expenditure data are expressed as a daily average from the 12-day data collection period and also the initial 4-day collection period. Physical activity level (PAL) was also calculated for each player by dividing TEE by RMR. PAL data is provided for 18 players only, given that 6 players were not available for DXA assessment (hence predicted RMR was not calculated for these players).

Assessment of energy and macronutrient intake

All twenty-four players on camp completed assessment of dietary and energy intake. Dietary intake was assessed for the first four days of the study via weighed food inventory. A four-day assessment period was chosen due to logistical issues with overseas travel for the rest of the study. This method of energy intake assessment has previously been used alongside DLW with athletes (35). All main meals were consumed (i.e., breakfast, lunch, and dinner) in the presence of the research team. Any snacks consumed outside of these meals was reported to the research team via the remote food photography method, as described previously (9,11,36). All players were free to self-select food choices and had received no prior education on nutrition strategies for training days. As such, players were asked to continue with their habitual nutritional practices through the study period. The information gained from this study was then used to produce individualised education and behaviour change strategies. Weighed food intake was assessed using an identified weighing station for main meals only, which included four separate calibrated weighing scales (Salter 1160 BKDR, Tonbridge, Kent, UK) placed on top of four separate A3 1cm cubed template place mats. The members of research team operating the stations during breakfast, lunch and dinner included three Sport and Exercise Register (SENr) registered performance nutritionists. Once participants had selected their first item of food, they arrived at the weighing station, placed their plate on the scale and informed the registered nutritionist the weight of the plate. This number was then populated into a pre-designed spreadsheet with a description of the food item underneath their name. For example, the participant would tell the member of staff the weight of their food item i.e., 762 g of white pasta, to inform both the weight and item of food. The participant would then place their second chosen item of food on the plate, for example chicken, and would return to the weighing station to re-weigh their plate, by calling out the weight and food item to the member of staff. Participants would follow the same process of calling out the new total weight and food item

to one of the three nutritionists who again would populate the spreadsheet. The spreadsheet was pre-designed to subtract the weight of the plate from the initial food item to allow quantification of food item number 1. Subsequently, as each food item was then added to the participant plate, the spreadsheet would automatically subtract the previous food item away from the measured food item so quantification of each food item could be calculated independently. This process was repeated until all participants had completed their total meal choice, at which point a photographic picture was captured of the complete final meal and weight and stored for later analysis. If players had finished eating and still had food left on their plate, they were asked to return to the weighing station to see a member of the research team who would subtract any food items left off the original completed meal total via the spreadsheet. In addition to weighing food, the remote food photographic method (RFPM) was used (11), to understand and retrieve information on what players consumed away from the three main mealtimes. This included EI consumed during “snack windows” provided on camp and EI consumed in hotel rooms. Players were asked to provide a photograph of the food or drink that they consumed and were sent to the research team on a smart phone via WhatsApp messaging service, as described previously (36). Thirdly, to further enhance reliability and ensure that participants missed no food or drink consumption, six random 24-hr food recalls were also performed by two members of the research team to cross check methods one and two. To obtain energy and macronutrient composition, professional dietary analysis software (Nutritics Ltd, Ireland) was used by a Sport and Exercise Nutrition register accredited practitioner with experience working with Nutritics Ltd. All energy intake is reported in kilocalories (kcal) and kilocalories per kilogram of total body mass ($\text{kcal}\cdot\text{kg}^{-1}$). Macronutrient intakes were also analysed and reported in grams (g) and grams per kilogram of body mass ($\text{g}\cdot\text{kg}^{-1}$).

Menu construction and the preparation of meals and snacks were undertaken by the national team's professional chef and performance nutrition team and developed in line with the demands of the training camp and consideration of proximity to each game. Throughout the duration of energy intake assessment, meals were consumed at the base camp hotel for the squad with menus provided on a buffet style basis. Breakfast options available daily included: eggs, beans, toast, porridge, muesli, fruits and yoghurts. Lunch and dinner had different options that included one red meat option, one poultry option, one fish option, three-to-four carbohydrate options (e.g., pasta, rice, potatoes, quinoa), three vegetable options alongside a salad bar and snacks such as yoghurts, nuts, cereal bars and condiments. During training sessions, players were provided with low calorie isotonic sports drinks (Lucozade Lite), water and upon request, isotonic energy gels (Science in Sport, GO Isotonic Gels, UK). Protein drinks (Science in Sport, Whey Protein, UK) were provided after training sessions. All carbohydrate provided during training were optional and consumed *ad libitum* as opposed to individualised prescription to players.

Estimation of energy availability

Given that FFM was known for 18 players only (due to completion of DXA assessment), EA was initially estimated for this cohort. However, due to a sample error with the urine sample provided by one player on day 4, this player's 4-day analysis of TEE was not completed, hence EA is estimated for 17 players. The thermic effect of food (TEF) was assumed to be 10% of EI for all individuals (37), subsequently enabling estimations of activity energy expenditure ($AEE = TEE - [RMR + TEF]$) and energy availability ($EA = EI - [AEE/FFM]$) (38) during the initial four days of the training camp. Energy availability was defined using the following thresholds: optimal ($> 45 \text{ kcal.kg FFM}^{-1}.\text{day}^{-1}$), reduced ($30\text{-}45 \text{ kcal.kg FFM}^{-1}.\text{day}^{-1}$) and low ($<30 \text{ kcal.kg FFM}^{-1}.\text{day}^{-1}$) (20).

Statistical Analysis

All data were initially assessed for normality of distribution using Shapiro-Wilk's test. Differences in training load, match load and energy intake across days were analysed using a one-way repeated measures ANOVA. Where significant main effects were present, Tukey post-hoc analysis was conducted to locate specific differences. Comparisons between energy intake and expenditure were analysed using a paired t-test. Ninety-five percent confidence intervals (95% CI) for the differences are also presented. Relationships between TEE and body mass, fat-free mass, stature, RMR and four-day AEE were assessed using Pearson's correlation. All statistical analysis were completed using SPSS (version 27, SPSS, Chicago, IL) where $P < 0.05$ is indicative of statistical significance. Data are presented as mean \pm SD.

Results

Baseline characteristics

Player characteristics including stature, body mass, fat-free mass, fat mass, percent body fat, bone mineral content and bone mineral density are presented in Table 1. Data are presented for the full cohort as well as mean data from positional groups.

Training and match load

External loading variables are presented for $n=13$ in accordance with those players who wore GPS monitors across all training sessions and games. Training duration (Figure 2A) was longer on MD-4 (89 ± 4 min) compared to MD-1 for match one (61 ± 2 min; 95% CI = 22 to 32 min; $P < 0.01$) and MD-1 for match two (63 ± 7 min; 95% CI = 17 to 34 min; $P < 0.01$). Similarly, MD-3 training duration (89 ± 5 min) was also longer than MD-1 training duration for match

one (95% CI = 21 to 33 min; $P < 0.01$) and match two (95% CI = 18 to 33 min; $P < 0.01$). In contrast, no difference was apparent for the duration of match one (64 ± 33 min) and match two (73 ± 31 min) compared to the remaining training days ($P > 0.05$).

In accordance with exercise duration, more distance (Figure 2B) was covered on MD-4 (6020 ± 620 m) compared to MD-1 for match one (2927 ± 862 m; 95% CI = 2090 to 4095 m; $P < 0.01$) and MD-1 for match two (4063 ± 540 m; 95% CI = 1177 to 2736 m; $P < 0.01$). Similarly, MD-3 distance covered (6340 ± 537 m) was greater than MD-1 distance covered for match one (95% CI = 2264 to 4562 m; $P < 0.01$) and match two (95% CI = 1721 to 2833 m; $P < 0.01$). The distance covered on MD-1 for match one was significantly lower than both the distance covered on MD-1 for match two ($P = 0.012$) and the distance covered in match two (7430 ± 3237 m; 95% CI = -7734 to -1272 m; $P = 0.004$). There was no significant difference in distance covered between match day one (6243 ± 340 m) and all other days ($P > 0.05$).

High-speed running distance (Figure 2C) was significantly greater during match one (361 ± 183 m) compared to MD-4 (126 ± 85 m; 95% CI = 73 to 395 m; $P < 0.01$), MD-1 for match one (85 ± 79 m; 95% CI = 102 to 450 m; $P < 0.01$) and MD-1 for match two (77 ± 41 m; 95% CI = 107 to 460 m; $P < 0.01$). High-speed running distance was significantly greater during match two (337 ± 197 m) when compared to MD-1 for both match one ($P < 0.01$) and match two ($P = 0.013$), although no significant difference was apparent with other training days or match one ($P > 0.05$). There was no significant difference in high-speed running distance between other training days ($P > 0.05$).

Energy expenditure

Mean TEE for the whole cohort ($n = 24$) across the full 12-day period was 2693 ± 432 kcal.day⁻¹ (range: 2105-3507 kcal.day⁻¹), 43 ± 6 kcal.kg⁻¹ (range 33-55 kcal.kg⁻¹) and 54 ± 6 kcal.kg⁻¹ FFM (range: 45-68 kcal.kg⁻¹ FFM). Mean four-day TEE ($n = 23$) was 2753 ± 359 kcal.day⁻¹

(range: 1942-3280 kcal.day⁻¹), 44 ± 7 kcal.kg⁻¹ (range 29-55 kcal.kg⁻¹) and 56 ± 8 kcal.kg⁻¹ FFM (range: 37-68 kcal.kg⁻¹ FFM). There was no significant difference between 12-day TEE and 4-day absolute TEE (P=0.307). Mean four-day AEE (n=23) was 1058 ± 352 kcal.day⁻¹ (range: 155-1549 kcal.day⁻¹) and mean PAL values (n=18) was 1.79 ± 0.24 (range: 1.4-2.2). For illustrative purposes, individual data points (where players are represented within their positional groups) are displayed in Figure 3 A-D.

Energy intake and macronutrient intake

Mean energy intake (n=24) during the 4-day assessment period was 1923 ± 232 kcal.day⁻¹ (range: 1639-2172 kcal.day⁻¹). Both absolute (P<0.01) and relative (P<0.01) mean energy intake (Figure 4A and B) was significantly different between training days. In absolute terms, players consumed less energy on MD-3 (1639 ± 285 kcal.day⁻¹) compared to MD-4 (2172 ± 373 kcal.day⁻¹, 95% CI -807 to -259 kcal.day⁻¹, P<0.01), MD-2 (1919 ± 319 kcal.day⁻¹, 95% CI -554 to -5 kcal.day⁻¹, P=0.04) and MD-1 (1962 ± 452 kcal.day⁻¹, 95% CI -597 to -48 kcal.day⁻¹, P=0.01). In contrast, there was no difference between the MD-4 and MD-2 (P=0.80) or MD-1 (P=0.19) and between MD-2 and MD-1 (P=0.97). In relative terms, players consumed less energy on MD-3 (26 ± 5 kcal.kg⁻¹.day⁻¹) compared with MD-4 (34 ± 6 kcal.kg⁻¹.day⁻¹, 95% CI 34 to 13 kcal.kg⁻¹.day⁻¹, P<0.01) and MD-1 (31 ± 8 kcal.kg⁻¹.day⁻¹, 95% CI -10 to 1 kcal.kg⁻¹.day⁻¹, P=0.02). In contrast, no difference was apparent between MD-3 and MD-2 (30 ± 6 kcal.kg⁻¹.day⁻¹, P=0.07), MD-4 and MD-2 (P=0.11) or MD-1 (P=0.25) and between MD-2 and MD-1 (P=0.97).

Mean absolute CHO intake (Figure 4C) was similar (P=0.37) between MD-4 (218 ± 56 g.day⁻¹), MD-3 (203 ± 57 g.day⁻¹), MD-2 (192 ± 45 g.day⁻¹) and MD-1 (203 ± 71 g.day⁻¹). Similarly, mean relative CHO intake (Figure 4D) was similar (P=0.38) between MD-4 ($3.5 \pm$

0.9 g.kg⁻¹.day⁻¹), MD-3 (3.2 ± 1.0 g.kg⁻¹.day⁻¹), MD-2 (3.0 ± 0.7 g.kg⁻¹.day⁻¹) and MD-1 (3.2 ± 1.1 g.kg⁻¹.day⁻¹).

Mean absolute protein intake was significantly different (P<0.01; Figure 4E) between training days such that on MD-4 (123 ± 21 g.day⁻¹), MD-3 (120 ± 33 g.day⁻¹) and MD-1 (135 ± 24 g.day⁻¹) more protein was consumed than on MD-2 (100 ± 23 g.day⁻¹; 95% CI = 5 to 41 g.day⁻¹; P<0.01, 95% CI = 2 to 39 g.day⁻¹; P=0.02 and 95% CI = 18 to 52 g.day⁻¹; P<0.01, respectively). No difference was observed between MD-4, MD-3, and MD-1 (P>0.05). Mean relative protein intake was significantly different (P<0.01; Figure 4F) between training days such that on MD-4 (1.9 ± 0.2 g.kg⁻¹.day⁻¹), MD-3 (1.9 ± 0.4 g.kg⁻¹.day⁻¹) and MD-1 (2.1 ± 0.4 g.kg⁻¹.day⁻¹) more protein was consumed than on MD-2 (1.6 ± 0.4 g.kg⁻¹.day⁻¹; 95% CI = 0.0 to 0.6 g.kg⁻¹.day⁻¹; P<0.01, 95% CI = 0.0 to 0.5 g.kg⁻¹.day⁻¹; P=0.03 and 95% CI = 0.3 to 0.8 g.kg⁻¹.day⁻¹; P<0.01, respectively).

Mean absolute fat intake was significantly different (P<0.01; Figure 4G) between training days such that on MD-4 (90 ± 21 g.day⁻¹), more fat was consumed than on MD-3 (38 ± 14 g.day⁻¹; 95% CI = 37 to 66 g.day⁻¹; P<0.01) and MD-1 (67 ± 24 g.day⁻¹; 95% CI = 3 to 42 g.day⁻¹; P<0.01). Similarly, more fat was consumed on MD-2 (87 ± 33 g.day⁻¹; 95% CI = 28 to 69 g.day⁻¹; P<0.01) than MD-3 and MD-1 (67 ± 24 g.day⁻¹; 95% CI = 15 to 43 g.day⁻¹; P<0.01) compared to MD-3. Mean relative fat intake was significantly different (P<0.01; Figure 4H) between training days such that on MD-4 (1.4 ± 0.3 g.kg⁻¹.day⁻¹), more fat was consumed compared to MD-3 (0.6 ± 0.2 g.kg⁻¹.day⁻¹; 95% CI = 0.5 to 1.0 g.kg⁻¹.day⁻¹; P<0.01) and MD-1 (1.0 ± 0.4 g.kg⁻¹.day⁻¹; 95% CI = 0.0 to 0.6 g.kg⁻¹.day⁻¹; P<0.01). Similarly, more fat was consumed on MD-2 (1.3 ± 0.5 g.kg⁻¹.day⁻¹) when compared to MD-3 (95% CI = 0.4 to 1.1 g.kg⁻¹.day⁻¹; P<0.01) and on MD-1 when compared to MD-3 (95% CI = 0.2 to 0.6 g.kg⁻¹.day⁻¹; P<0.01).

Energy intake versus energy expenditure (n = 24) and energy availability (n = 17)

In relation to the initial 4-day assessment period, there was a significant difference between EI and TEE ($-825 \pm 419 \text{ kcal.day}^{-1}$; 95% CI -1006 to $-643 \text{ kcal.day}^{-1}$; $P < 0.01$) (see Figure 5A). However, despite significant differences in EI and TEE, body mass did not change across this time period (see Figure 5B) ($0.01 \pm 1.16 \text{ kg}$; 95% CI -0.48 to 0.51 kg ; $P = 0.95$). Mean daily (n = 17) estimated energy availability was $18 \pm 9 \text{ kcal.kg FFM}^{-1}.\text{day}^{-1}$ (range: 2 - $36 \text{ kcal.kg FFM}^{-1}.\text{day}^{-1}$). Overall, 88% of players assessed for EA represented with $< 30 \text{ kcal.kg FFM}^{-1}.\text{day}^{-1}$ (see Figure 5C).

Factors affecting TEE and AEE

There was a significant positive relationship between 12-day TEE and body mass ($r^2 = 0.56$; $P < 0.01$), fat-free mass ($r^2 = 0.65$; $P < 0.01$) and predicted RMR ($r^2 = 0.51$; $P < 0.01$). There was also a significant positive relationship between four-day TEE and four-day AEE ($r^2 = 0.97$; $P < 0.01$). There was no significant relationship between TEE and stature ($r^2 = 0.15$; $P > 0.05$). Data are presented in Figure 6.

Discussion

In using the DLW method, we provide the first direct assessment of total daily energy expenditure of adult female professional soccer players. Our measurements were obtained from players of the highest standard and were collected over a 12-day period when players were representing their national team. When compared with previously published data from adult male players, we demonstrate that the relative daily energetic requirements of engaging in professional soccer training and match play is comparable between sexes. As such, these data now provide a platform for which to develop evidence based nutritional guidelines for this population. From a practical perspective, our data suggest that practitioners should likely focus education and behaviour change strategies (at least for the present cohort) on “fuelling” for match play and training to optimise both player health and performance.

Previous assessments of daily TEE and AEE in female soccer players have been quantified using a combination of indirect methods such as accelerometers, heart rate monitors, activity logs and prediction equations (19, 24, 25, 44, 46). In absolute terms, such studies report that the TEE of female soccer players ranges from ~2400-2700 kcal.day⁻¹ (22,39,40). In using the DLW method, we observed comparable mean four-day (three training days, one rest day) TEE of 2753 ± 359 kcal.day⁻¹ (range: 1942-3280 kcal.day⁻¹) whilst mean TEE from the full 12-day assessment period was 2693 ± 423 kcal.day⁻¹ (range: 2105-3507 kcal.day⁻¹). In absolute terms, our data demonstrate a lower TEE to that previously observed in adult male professional players where mean expenditure was approximately 3500 kcal.d⁻¹ (9–11). Nonetheless, when expressed in relative terms (alongside comparable PAL values of 1.4-2.2), it is therefore apparent that the daily energetic requirements of both males and females engaging in professional soccer training and match play typically equates to 40-60 kcal.kg⁻¹ FFM.

Notwithstanding the limitations of comparing indirect and direct assessment methods, the present data also suggest that the energy requirements of competing and training at an “international” level may be higher than that associated with the players’ respective domestic level competition. For example, when compared with players from the English Women’s Super League (WSL), assessments of the AEE of the goalkeepers ($924 \pm 133 \text{ kcal.day}^{-1}$), defenders ($964 \pm 436 \text{ kcal.day}^{-1}$), midfielders ($1318 \pm 195 \text{ kcal.day}^{-1}$) and attackers ($1073 \pm 348 \text{ kcal.day}^{-1}$) studied here is greater than the mean AEE ($418 \text{ kcal.day}^{-1}$) of those players training within the domestic WSL (13). It is noteworthy, however, that the DLW derived assessment of AEE documented here is inclusive of all activity “outside” of pitch-based training such as strength-based sessions undertaken in the gym, recovery swimming pool sessions, as well as non-training related activity such as walking to and from the training centre and hotel and walking up and down stairs etc. In contrast, the AEE quantified by Moss et al. (19) is derived from a combination of metabolic equivalents and/or accelerometers worn during training, matches and strength and conditioning sessions only. Additionally, the training loads completed by Moss et al. (19) was completed in the final month of the season (May), a time when training loads are typically reduced in comparison to other phases of the season.

The external training and match loads observed here are lower than the respective loads associated with other international and domestic level soccer match play (41–43). For example, total distance and high speed running distance covered by outfield players is lower in our study ($8.8 \pm 1.4 \text{ km}$ and $0.35 \pm 0.18 \text{ km}$ respectively) compared with other international ($9.9 \pm 1.8 \text{ km}$ and $1.5 \pm 0.1 \text{ km}$ respectively) and domestic ($9.7 \pm 1.4 \text{ km}$ and 1.3 ± 0.9 respectively) soccer matches (42). Difference between studies are most likely due to variation in methods used to collect match load data, where in previous studies, distance covered and high speed running was estimated from time motion analysis as opposed to GPS adopted here.

457 Additionally, the thresholds used for high-speed running in previous studies ($>18 \text{ km.h}^{-1}$) is
458 lower than this study ($>19 \text{ km.h}^{-1}$) and makes it difficult to compare between studies. Such
459 challenges in the lack of a definitive approach to identify high-intensity actions and the
460 subsequent ambiguity in this area has recently been documented (44).

461
462 In relation to energy intake, previous studies in female soccer players have reported estimated
463 energy intakes of $2124 \pm 444 \text{ kcal.day}^{-1}$ (13), $2226 \pm 368 \text{ kcal.day}^{-1}$ (39) and 2387 ± 177
464 kcal.day^{-1} (16). In contrast, we report estimated energy intakes that are approximately 200-300
465 kcal.d^{-1} lower (mean of four-days: $1923 \pm 357 \text{ kcal.day}^{-1}$), a finding that may be due, in part,
466 to the differing methods employed (e.g., self-reported food diaries versus researcher supervised
467 weighed food intakes, the latter which may have influenced player food choices towards under-
468 consumption of foods). In agreement with recent observations from players from the English
469 Women's Super League (13), we also observed minimal CHO periodisation with players
470 reporting comparable and consistent daily CHO intakes of 3.0 to 3.5 g.kg^{-1} . Notably, only one
471 player consumed the recommended range of 6-8 g.kg^{-1} on the day before the match (12), thus
472 it is likely that players commenced the first game with sub-optimal muscle glycogen stores
473 (18). In contrast, mean protein intake across all training days ($1.8 \pm 0.4 \text{ g.kg}^{-1}.\text{day}^{-1}$; range 1.6
474 to 2.1 $\text{g.kg}^{-1}.\text{day}^{-1}$) was aligned to supporting training adaptations (45) and in accordance with
475 recommendations for professional soccer players (12). When taken together, it therefore
476 appears that female soccer players may not consume (or periodise) sufficient CHO intake to
477 meet the demands of training and competition, a factor that could lead to chronically low
478 energy availability and symptoms associated with the female athlete triad (21) or RED-S
479 models (19). Unfortunately, we are limited in that we do not currently provide any data
480 assessing the impacts of the energy intake reported here on health and performance outcomes.
481 Nonetheless, from a practical perspective, our data suggest that practitioners should likely

target education and behaviour change strategies on “fuelling” for match play and training to optimise both player health and performance. Based on our assessment of TEE, it is suggested that relative intakes of CHO, fat and protein corresponding to 4-8 (to account for rest-days, training days, match day minus 1, match day etc), 1.5-2 and 1.6-2 g·kg⁻¹·day⁻¹ body mass would provide a reasonable starting point for which to meet the daily energy requirements of female soccer players of professional standard.

Although we readily acknowledge the difficulties in assessing energy availability (46) as well as the limitation of our four-day assessment period via weighed food inventory (i.e. players may alter food intake because of researcher presence), it is noteworthy that the estimated prevalence of LEA observed here (i.e., 88%, 15 out of 17, players presented with LEA <30 kcal.kg⁻¹ FFM) is greater than previous reports where 70, 24 and 65% of players presented with LEA in English (13), American (14) and Polish national leagues (40) respectively. The lower absolute energy intakes reported here coupled with the potentially increased physical demands associated with competing at international level (when compared to domestic level competition) may be a contributing factor. Whilst we also acknowledge the limitations (35,36) associated with dietary assessment and potential under-reporting (as evidenced by the lack of statistical change in body mass), further work is required to ascertain whether players’ chosen dietary choices were an unconscious or conscious decision that is based upon beliefs surrounding optimal nutritional practices. We also acknowledge that the classification of LEA status as <30 kcal.kg⁻¹ FFM is based upon laboratory studies that typically adopt short-term periods of “consistent” daily EI, EE and therefore EA. For example, studies which established EA concepts did so over short (four-to-seven days) periods where careful but artificial control of diet and exercise was prescribed (20). The application of such a threshold to real world situations is likely limited by the fact that daily energy expenditure fluctuates day-to-day in

507 accordance with alterations to eating schedules, training load, and competitive demands.
508 Accordingly, the prevalence of LEA status in the present study (and associated long term
509 physiological implications) may be over-estimated. Further studies are required to evaluate the
510 prevalence of LEA using longer assessment timeframes. Furthermore, assessment of within-
511 day and between-day EA combined with screening tools (21,47,48) and clinical markers would
512 help gain greater accuracy with current assessments of EA in female athletes in the applied
513 field.

514
515 In summary, we provide the first report to directly assess total daily energy expenditure in a
516 cohort of adult female professional soccer players of international standard. Our data suggest
517 that the relative daily energetic requirements of engaging in professional soccer training and
518 match play is comparable in males and females. From a practical perspective, our data suggest
519 that individualised education and behaviour change strategies should focus on “fuelling” (i.e.
520 increasing daily CHO intake) for match play and training to optimise health and performance.

523 **Acknowledgements**

524 This research was funded by The Football Association. The authors would like to thank all
525 of the players for their commitment during this study.

527 **Conflict of interest**

528 The authors report no potential conflict of interest. The results of the present study do not
529 constitute endorsement by ACSM. All results presented here are done so clearly, honestly,
530 and without fabrication, falsification, or inappropriate data manipulation.

531

532 **References**

- 533 1. Bloomfield J, Polman R, O'Donoghue P. Physical demands of different positions in
534 FA Premier League soccer. *J Sport Sci Med*. 2007; 6(1):63-32.
- 535 2. Bush M, Barnes C, Archer DT, Hogg B, Bradley PS. Evolution of match performance
536 parameters for various playing positions in the English Premier League. *Hum Mov Sci*
537 [Internet]. 2015 Feb;39:1–11. Available from:
538 <https://linkinghub.elsevier.com/retrieve/pii/S0167945714001808>
- 539 3. Carling C, Bloomfield J, Nelsen L, Reilly T. The role of motion analysis in elite
540 soccer: Contemporary performance measurement techniques and work rate data.
541 *Sports Medicine*. 2008; 38(10):839-862.
- 542 4. Anderson L, Orme P, Di Michele R, Close GL, Morgans R, Drust B, et al.
543 Quantification of training load during one-, two- and three-game week schedules in
544 professional soccer players from the English Premier League: implications for
545 carbohydrate periodisation. *J Sports Sci*. 2016; 34(13):1250-1259.
- 546 5. Malone JJ, Di Michele R, Morgans R, Burgess D, Morton JP, Drust B. Seasonal
547 training-load quantification in elite English Premier League soccer players. *Int J Sports*
548 *Physiol Perform*. 2015; 10(4):489-497.
- 549 6. Akenhead R, Harley JA, Tweddle SP. Examining the External Training Load of an
550 English Premier League Football Team With Special Reference to Acceleration. *J*
551 *Strength Cond Res* [Internet]. 2016 Sep;30(9):2424–32. Available from:
552 <http://journals.lww.com/00124278-201609000-00008>
- 553 7. Bangsbo J, Mohr M, Krstrup P. Physical and metabolic demands of training and
554 match-play in the elite football player. *J Sport Sci* [Internet]. 2006;24(7):665–74.
555 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16766496>

- 556 8. Bradley PS, Sheldon W, Wooster B, Olsen P, Boanas P, Krstrup P. High-intensity
557 running in English FA Premier League soccer matches. *J Sports Sci.* 2009; 27(2):159-
558 168.
- 559 9. Anderson L, Orme P, Naughton RJ, Close GL, Milsom J, Rydings D, et al. Energy
560 intake and expenditure of professional soccer players of the English Premier league:
561 Evidence of carbohydrate periodization. *Int J Sport Nutr Exerc Metab.* 2017;
562 27(3):228-238.
- 563 10. Brinkmans NYJ, Iedema N, Plasqui G, Wouters L, Saris WHM, van Loon LJC, et al.
564 Energy expenditure and dietary intake in professional football players in the Dutch
565 Premier League: Implications for nutritional counselling. *J Sports Sci.* 2019;
566 37(24):2759-2767.
- 567 11. Hannon MP, Parker LJF, Carney DJ, McKeown J, Speakman JR, Hambly C, et al.
568 Energy Requirements of Male Academy Soccer Players from the English Premier
569 League. *Med Sci Sport Exerc* [Internet]. 2020 Jul 21; 53(1):200-210. Available from:
570 <https://journals.lww.com/10.1249/MSS.00000000000002443>
- 571 12. Collins J, Maughan RJ, Gleeson M, Bilsborough J, Jeukendrup A, Morton JP, et al.
572 UEFA expert group statement on nutrition in elite football. Current evidence to inform
573 practical recommendations and guide future research. *Br J Sports Med* [Internet]. 2021
574 Apr;55(8):416–416. Available from:
575 <https://bjsm.bmj.com/lookup/doi/10.1136/bjsports-2019-101961>
- 576 13. Moss SL, Randell RK, Burgess D, Ridley S, ÓCairealláin C, Allison R, et al.
577 Assessment of energy availability and associated risk factors in professional female
578 soccer players. *Eur J Sport Sci* [Internet]. 2020 Aug 6;1–10. Available from:
579 <https://www.tandfonline.com/doi/full/10.1080/17461391.2020.1788647>
- 580 14. Reed JL, De Souza MJ, Kindler JM, Williams NI. Nutritional practices associated with

- low energy availability in Division I female soccer players. *J Sports Sci*. 2014;
32(16):1499-509
15. Martin L, Lambeth A, Scott D. Nutritional practices of national female soccer players: analysis and recommendations. *J Sports Sci Med [Internet]*. 2006;5(1):130–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24198690>
16. Reed JL, de Souza MJ, Williams NI. Changes in energy availability across the season in Division I female soccer players. *J Sports Sci*. 2013; 31(3):314-24.
17. Dobrowolski H, Włodarek D. Dietary Intake of Polish Female Soccer Players. *Int J Environ Res Public Health [Internet]*. 2019 Mar 29 [cited 2021 Nov 3];16(7):1134. Available from: <https://pubmed.ncbi.nlm.nih.gov/30934885/>
18. Krstrup P, Mohr M, Nybo L, Draganidis D, Randers MB, Ermidis G, et al. Muscle metabolism and impaired sprint performance in an elite women's football game. *Scand J Med Sci Sports [Internet]*. 2021 Jun 25 [cited 2021 Jul 3];sms.13970. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/sms.13970>
19. Areta JL, Taylor HL, Koehler K. Low energy availability: history, definition and evidence of its endocrine, metabolic and physiological effects in prospective studies in females and males. *Eur J Appl Physiol [Internet]*. 2020 Oct 23; 121(1):1-21. Available from: <http://link.springer.com/10.1007/s00421-020-04516-0>
20. Loucks AB, Kiens B, Wright HH. Energy availability in athletes. *J Sports Sci [Internet]*. 2011 Jan 28;29(sup1):S7–15. Available from: <https://www.tandfonline.com/doi/full/10.1080/02640414.2011.588958>
21. Mountjoy M, Sundgot-Borgen J, Burke L, Carter S, Constantini N, Lebrun C, et al. The IOC consensus statement: Beyond the Female Athlete Triad-Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med*. 2014; 48(7):491-7.
22. Mara JK, Thompson KG, Pumpa KL. Assessing the energy expenditure of elite female

- 606 soccer players: A preliminary study. *J Strength Cond Res.* 2015; 29(10):2780-6.
- 607 23. Westerterp KR. Doubly labelled water assessment of energy expenditure: principle,
608 practice, and promise. *Eur J Appl Physiol* [Internet]. 2017 Jul;117(7):1277–85.
609 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28508113>
- 610 24. Nana A, Slater GJ, Hopkins WG, Halson SL, Martin DT, West NP, et al. Importance
611 of standardized DXA protocol for assessing physique changes in athletes. *Int J Sport*
612 *Nutr Exerc Metab.* 2016; 26(3):259-67.
- 613 25. Watson AD, Zabriskie HA, Witherbee KE, Sulavik A, Gieske BT, Kerksick CM.
614 Determining a Resting Metabolic Rate Prediction Equation for Collegiate Female
615 Athletes. *J strength Cond Res.* 2019; 2426-2434.
- 616 26. Marfell-jones M, Olds T, Stewart A, Carter L. International standards for
617 anthropometric assessment. International Society for the Advancement of
618 Kinanthropometry. *Int Soc Adv Kinanthropometry.* 2006;
- 619 27. Hannon MP, Coleman NM, Parker LJF, McKeown J, Unnithan VB, Close GL, et al.
620 Seasonal training and match load and micro-cycle periodization in male Premier
621 League academy soccer players. *J Sports Sci* [Internet]. 2021 Mar 24;1–12. Available
622 from: <https://www.tandfonline.com/doi/full/10.1080/02640414.2021.1899610>
- 623 28. Anderson L, Orme P, Michele R Di, Close GL, Milsom J, Morgans R, et al.
624 Quantification of Seasonal-Long Physical Load in Soccer Players With Different
625 Starting Status From the English Premier League: Implications for Maintaining Squad
626 Physical Fitness. *Int J Sports Physiol Perform* [Internet]. 2016 Nov;11(8):1038–46.
627 Available from: [https://journals.humankinetics.com/view/journals/ijsp/11/8/article-](https://journals.humankinetics.com/view/journals/ijsp/11/8/article-p1038.xml)
628 [p1038.xml](https://journals.humankinetics.com/view/journals/ijsp/11/8/article-p1038.xml)
- 629 29. Morehen JC, Bradley WJ, Clarke J, Twist C, Hambly C, Speakman JR, et al. The
630 Assessment of Total Energy Expenditure During a 14-Day In-Season Period of

631 Professional Rugby League Players Using the Doubly Labelled Water Method. *Int J*
632 *Sport Nutr Exerc Metab* [Internet]. 2016;26(5):464–72. Available from:
633 <https://www.ncbi.nlm.nih.gov/pubmed/27096279>

634 30. Speakman JR. Doubly Labelled Water; Theory and Practice. London: Chapman &
635 Hall; 1997.

636 31. Nagy K. The Doubly Labelled Water (3HH18O) Method: a guide to its use. Los
637 Angeles, CA: UCLA Publication; 1983. 12–1417.

638 32. Berman ESF, Fortson SL, Snaith SP, Gupta M, Baer DS, Chery I, et al. Direct analysis
639 of $\delta^2\text{H}$ and $\delta^{18}\text{O}$ in natural and enriched human urine using laser-based, off-axis
640 integrated cavity output spectroscopy. *Anal Chem* [Internet]. 2012 Nov
641 20;84(22):9768–73. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23075099>

642 33. Craig H. Standard for Reporting Concentrations of Deuterium and Oxygen-18 in
643 Natural Waters. *Science* (80-) [Internet]. 1961;133(3467):1833–4. Available from:
644 <https://www.ncbi.nlm.nih.gov/pubmed/17819002>

645 34. Speakman JR, Yamada Y, Sagayama H, Berman ESF, Ainslie PN, Andersen LF, et al.
646 A standard calculation methodology for human doubly labeled water studies. *Cell*
647 *Reports Med* [Internet]. 2021 Feb 16 [cited 2021 Jun 30];2(2):100203. Available from:
648 <https://pubmed.ncbi.nlm.nih.gov/33665639/>

649 35. Capling L, Beck K, Gifford J, Slater G, Flood V, O'Connor H. Validity of Dietary
650 Assessment in Athletes: A Systematic Review. *Nutrients*. 2017; 9(12):1313.

651 36. Stables RG, Kasper AM, Sparks SA, Morton JP, Close GL. An Assessment of the
652 Validity of the Remote Food Photography Method (Termed Snap-N-Send) in
653 Experienced and Inexperienced Sport Nutritionists. *Int J Sport Nutr Exerc Metab*.
654 2021; 31(2):125-134.

655 37. Westerterp KR. Diet induced thermogenesis [Internet]. Vol. 1, Nutrition and

Metabolism. BioMed Central; 2004 [cited 2021 Jun 25]. p. 5. Available from:
/pmc/articles/PMC524030/

38. Loucks AB. Energy balance and body composition in sports and exercise. *J Sports Sci*; 2004;22(1):1–14. Available from: <https://pubmed.ncbi.nlm.nih.gov/14974441/>

39. Braun H, Von Andrian-Werburg J, Schänzer W, Thevis M. Nutrition status of young elite female German football players. *Pediatr Exerc Sci*. 2018; 30(1):157-167.

40. Dobrowolski H, Włodarek D. Low energy availability in group of Polish female soccer players. *Rocz Państwowego Zakładu Hig* [Internet]. 2020; 71(1):89-96. Available from: http://wydawnictwa.pzh.gov.pl/roczniki_pzh/pobierz-artykul?id=1323

41. Datson N, Hulton A, Andersson H, Lewis T, Weston M, Drust B, et al. Applied Physiology of Female Soccer: An Update. *Sport Med* [Internet]. 2014 Sep 7;44(9):1225–40. Available from: <http://link.springer.com/10.1007/s40279-014-0199-1>

42. Andersson HÅ, Randers MB, Heiner-Møller A, Krstrup P, Mohr M. Elite Female Soccer Players Perform More High-Intensity Running When Playing in International Games Compared With Domestic League Games. *J Strength Cond Res* [Internet]. 2010 Apr;24(4):912–9. Available from: <https://insights.ovid.com/crossref?an=00124278-201004000-00004>

43. Strauss A, Sparks M, Pienaar C. The Use of GPS Analysis to Quantify the Internal and External Match Demands of Semi-Elite Level Female Soccer Players during a Tournament. *J Sports Sci Med* [Internet]. 2019 Mar;18(1):73–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30787654>

44. Randell RK, Clifford T, Drust B, Moss SL, Unnithan VB, De Ste Croix MBA, et al. Physiological Characteristics of Female Soccer Players and Health and Performance Considerations: A Narrative Review [Internet]. Vol. 51, *Sports Medicine*. Springer

Science and Business Media Deutschland GmbH; 2021 [cited 2021 Jul 4]. p. 1377–99.
Available from: <https://doi.org/10.1007/s40279-021-01458-1>

45. Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med*. 2018; 52(6):376-384.
46. Burke LM, Lundy B, Fahrenholtz IL, Melin AK. Pitfalls of Conducting and Interpreting Estimates of Energy Availability in Free-Living Athletes. *Int J Sport Nutr Exerc Metab* [Internet]. 2018 Jul 1;28(4):350–63. Available from: <https://journals.humankinetics.com/view/journals/ijsnem/28/4/article-p350.xml>
47. Melin A, Tornberg ÅB, Skouby S, Faber J, Ritz C, Sjödén A, et al. The LEAF questionnaire: a screening tool for the identification of female athletes at risk for the female athlete triad. *Br J Sport Med*. 2014;48(7):540–5.
48. Souza MJ De, Nattiv A, Joy E, Misra M, Williams NI, Mallinson RJ, et al. 2014 Female Athlete Triad Coalition Consensus Statement on Treatment and Return to Play of the Female Athlete Triad: 1st International Conference held in San Francisco, California, May 2012 and 2nd International Conference held in Indianapolis, Indiana, May 2013. *Br J Sports Med* [Internet]. 2014 Feb 1 [cited 2021 Jul 11];48(4):289–289. Available from: <https://bjsm.bmj.com/content/48/4/289>

Table 1. Baseline player characteristics of elite English female soccer players competing at international level. Stature, body mass, fat-free mass, fat mass and percent body fat values are presented according to playing position. Stature and body mass n=24. Fat-free mass, fat mass, percent body fat, bone mineral content, bone mineral density, pelvis bone mineral density, Z-score derived from DXA n=18. Predicted resting metabolic rate (RMR) n=18. Predicted RMR = $120.81 + (4.88 \times \text{Stature}[\text{cm}]) + 8.24 \times \text{FFM}[\text{kg}] + (5.71 \times \text{Age}[\text{years}])$ (25).

Position	Goalkeepers	Defenders	Midfielders	Attackers	Squad
Stature (cm)	174.3 ± 0.5 (n=3)	169.7 ± 2.4 (n=9)	168.2 ± 9.2 (n=4)	163.0 ± 3.5 (n=8)	168.1 ± 5.9 (n=24)
Body Mass (kg)	67.0 ± 8.7 (n=3)	62.4 ± 3.2 (n=9)	60.4 ± 5.0 (n=4)	60.1 ± 1.1 (n=8)	62.1 ± 4.7 (n=24)
Fat-Free Mass (kg)	45.5 ± 3.5 (n=3)	44.1 ± 3.6 (n=6)	42.8 ± 3.9 (n=4)	41.6 ± 2.1 (n=5)	43.2 ± 3.4 (n=18)
Fat Mass (kg)	14.4 ± 5.1 (n=3)	11.1 ± 1.3 (n=6)	10.3 ± 3.0 (n=4)	12.2 ± 1.4 (n=5)	11.8 ± 2.7 (n=18)
Percent Body Fat (%)	22.9 ± 5.2 (n=3)	19.5 ± 2.6 (n=6)	18.6 ± 4.6 (n=4)	20 ± 2.7 (n=5)	20.6 ± 3.7 (n=18)
Whole Body Bone Mineral Content (g)	2808 ± 361 (n=3)	2837 ± 158 (n=6)	2803 ± 236 (n=4)	2637 ± 165 (n=5)	2766 ± 213 (n=18)
Whole Body Bone Mineral Density (g/cm²)	1.26 ± 0.12 (n=3)	1.33 ± 0.06 (n=6)	1.35 ± 0.11 (n=4)	1.26 ± 0.10 (n=5)	1.31 ± 0.10 (n=18)
Pelvis Bone Mineral Density (g/cm²)	1.37 ± 0.19 (n=3)	1.28 ± 0.11 (n=6)	1.35 ± 0.19 (n=4)	1.42 ± 0.11 (n=5)	1.38 ± 0.13 (n=18)
Whole Body Z-score	2.7 ± 1.0 (n=3)	2.4 ± 0.5 (n=6)	2.7 ± 1.2 (n=4)	2.1 ± 0.5 (n=5)	2.4 ± 0.7 (n=18)
Predicted RMR (kcal.day⁻¹)	1549 ± 56 (n=3)	1515 ± 71 (n=6)	1494 ± 95 (n=4)	1449 ± 46 (n=5)	1486 ± 66 (n=18)

Figure 1. Schematic overview of the 12-day study period including the 9-day national training camp. TEE was assessed over 12 days and 4-days (as opposed to 9-days) due to logistical challenges of urine collection on day 9 to 11 of the study. Day 6 and days 9-12 represented rest days during which no scheduled training took place.

Figure 2. (A) Training and match-play duration, (B) total distance, and (C) high speed running distance in during an international training camp from female soccer players. White bars represent training days, denoted as days away from match day (MD), i.e., MD-5, etc., and grey bars represent match day. No training was completed on days with no data bars. ^a denotes significant difference from MD-4, $P<0.05$. ^b denotes significant difference from MD-3, $P<0.05$. ^c denotes significant difference from MD-1 prior to match 1, $P<0.05$. ^d denotes significant difference from MD one, $P<0.05$. ^f denotes significant difference from MD two, $P<0.05$. Black circles represent individual players. All data are representative of $n=13$ in accordance with players who wore GPS monitors.

Figure 3. (A) Mean twelve daily total energy expenditure ($n=24$), (B) mean four-day total energy expenditure ($n=23$), (C) mean four-day activity energy expenditure ($n=23$), (D) physical activity level ($n=18$) within each positional group. Black circles represent individual players.

Figure 4. (A) Absolute and (B) relative energy intake, (C) absolute and (D) relative carbohydrate intake, (E) absolute and (F) relative protein intake and (G) absolute and (H) relative fat intake across the initial 4-day assessment period ($n=24$ for all variables). Black circles represent individual players. ^a denotes significant difference from MD-4, ^b denotes significant difference from MD-3, ^c denotes significant difference from MD-2, ^d denotes significance difference from MD-1

739

740 **Figure 5.** (A) Difference between TEE and EI (n=23), (B) changes in body mass (n=24) and
741 (C) mean estimated daily energy availability (n=17) when assessed across the initial 4-day
742 assessment period. Black circles represent individual players.

743

744 **Figure 6.** The relationship between mean 12-day total energy expenditure (TEE) and (A) body
745 mass ($P<0.01$), (B) fat free-mass ($P<0.01$), (C) stature ($P>0.05$), predicted resting metabolic
746 rate (RMR; $P<0.01$) and (E) 4-day TEE versus 4-day activity energy expenditure (AEE;
747 $P<0.01$). Black circles represent individual players.

748

