



LJMU Research Online

Watkins, JD, Carter, S, Atkinson, G, Koumanov, F, Betts, JA, Holst, JJ and Gonzalez, JT

Glucagon-like peptide-1 secretion in people with versus without type 2 diabetes: A systematic review and meta-analysis of cross-sectional studies

<http://researchonline.ljmu.ac.uk/id/eprint/18345/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)


Watkins, JD, Carter, S, Atkinson, G, Koumanov, F, Betts, JA, Holst, JJ and Gonzalez, JT (2022) Glucagon-like peptide-1 secretion in people with versus without type 2 diabetes: A systematic review and meta-analysis of cross-sectional studies. *Metabolism*. 140. ISSN 0026-0495

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



Glucagon-like peptide-1 secretion in people with *versus* without type 2 diabetes: a systematic review and meta-analysis of cross-sectional studies

J.D. Watkins^{a,*}, S. Carter^a, G. Atkinson^b, F. Koumanov^a, J.A. Betts^a, J.J. Holst^c, J.T. Gonzalez^{a,*}

^a Centre for Nutrition, Exercise and Metabolism, Department for Health, University of Bath, UK

^b Liverpool John Moores University, Liverpool, UK

^c Biomedical Sciences, Endocrinology Research Section, University of Copenhagen, Denmark

ARTICLE INFO

Keywords:

Endocrinology
Glucagon-like peptide-1
Meta-analysis
Type 2 diabetes

ABSTRACT

Aims/Hypothesis: The aim of this systematic review was to synthesise the study findings on whether GLP-1 secretion in response to a meal tolerance test is affected by the presence of type 2 diabetes (T2D). The influence of putative moderators such as age, sex, meal type, meal form, and assay type were also explored.

Methods: A literature search identified 32 relevant studies. The sample mean and SD for fasting GLP-1_{TOTAL} and GLP-1_{TOTAL} iAUC were extracted and used to calculate between-group standardised mean differences (SMD), which were meta-analysed using a random-effects model to derive pooled estimates of Hedges' *g* and 95 % prediction intervals (PI).

Results: Pooled across 18 studies, the overall SMD in GLP-1_{TOTAL} iAUC between individuals with T2D ($n = 270$, 1047 ± 930 pmol·L⁻¹·min) and individuals without T2D ($n = 402$, 1204 ± 937 pmol·L⁻¹·min) was very small, not statistically significant and heterogenous across studies ($g = -0.15$, $p = 0.43$, PI: -1.53 , 1.23). Subgroup analyses demonstrated an effect of assay type whereby Hedges' *g* for GLP-1 iAUC was greater in individuals with, *versus* those without T2D when using ELISA or Mesoscale ($g = 0.67$ [moderate], $p = 0.009$), but not when using RIA ($g = -0.30$ [small], $p = 0.10$). Pooled across 30 studies, the SMD in fasting GLP-1_{TOTAL} between individuals with T2D ($n = 580$, 16.2 ± 6.9 pmol·L⁻¹) *versus* individuals without T2D ($n = 1363$, 12.4 ± 5.7 pmol·L⁻¹) was small and heterogenous between studies ($g = 0.24$, $p = 0.21$, PI: -1.55 , 2.02).

Conclusions: Differences in fasting GLP-1_{TOTAL} and GLP-1_{TOTAL} iAUC between individuals with, *versus* those without T2D were generally small and inconsistent between studies. Factors influencing study heterogeneity such as small sample sizes and poor matching of groups may help to explain the wide prediction intervals observed. Considerations to improve comparisons of GLP-1 secretion in T2D and potential mediating factors more important than T2D diagnosis *per se* are outlined. PROSPERO ID: CRD42020195612.

1. Introduction

The 'incretin effect' is the amplification of insulin secretion observed following oral ingestion *versus* intravenous (isoglycaemic) glucose administration [1]. This effect is mainly mediated by the incretin hormones, GIP and GLP-1. GIP and GLP-1 are peptide hormones secreted by intestinal K- and L-cells, respectively [2]. These hormones both

stimulate glucose-dependent insulin secretion, while GLP-1 also reduces the rate of gastric emptying and suppresses food intake [3,4]. In individuals with type 2 diabetes (T2D), the incretin effect is, in spite of a preservation of GIP secretion, reduced in comparison to individuals without T2D, which contributes to reduced glucose tolerance [5]. This reduction could be due to either 1) lower GLP-1 release following meals, and/or 2) reduced beta-cell sensitivity to GIP and/or GLP-1. Unlike GIP,

Abbreviations: BMI, Body mass index; CI, Confidence interval; ELISA, Enzyme-linked immunosorbent assay; GIP, Glucose-dependent insulinotropic peptide; GLP-1, Glucagon-like peptide-1; iAUC, Incremental area under the curve; MMTT, Mixed meal tolerance test; MPGF, Major proglucagon fragment; OGTT, Oral glucose tolerance test; PI, Prediction interval; RIA, Radioimmunoassay; SD, Standard deviation; SGLT-1, sodium-glucose transporter protein-1; SMD, Standardised Mean Difference; T, Tau; T2D, Type 2 diabetes; tAUC, Total area under the curve.

* Corresponding authors at: Centre for Nutrition, Exercise and Metabolism, University of Bath, UK.

E-mail addresses: watkins47@sky.com (J.D. Watkins), j.t.gonzalez@bath.ac.uk (J.T. Gonzalez).

<https://doi.org/10.1016/j.metabol.2022.155375>

Received 4 July 2022; Accepted 4 December 2022

Available online 9 December 2022

0026-0495/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

the insulinotropic action of GLP-1 is partially preserved in individuals with T2D [6,7] although its potency is clearly impaired [8]. Nevertheless, reduced GLP-1 secretion could account for some of the reduced incretin effect observed amongst individuals with T2D.

Despite the plausibility of the above notion, there have been conflicting findings suggesting that GLP-1 secretion is reduced [9,10], similar [11], and even increased [12] in individuals with T2D compared to individuals without T2D. The authors of two previous meta-analyses concluded that there was no evidence of reduced GLP-1 secretion in individuals with T2D *versus* matched individuals without T2D [13,14]. Despite these conclusions, there is still general uncertainty in the literature, especially after further relevant data have been published since these meta-analyses were undertaken. There have also been recent calls for meta-analyses to incorporate considerations of prediction intervals [15] and double-counting of control groups from multiple treatment groups being present in some studies [16]. The following sections will outline the new aspects of this meta-analysis including fasting GLP-1, meal type and form, and assay type.

In contrast to postprandial GLP-1, fasting GLP-1 concentrations have yet to be analysed in a systematic review. Fasting GLP-1 concentrations have been demonstrated to be greater in adolescent and adult men and women with overweight/obesity compared to lean individuals [17,18], with fasting GLP-1 concentrations positively associated with BMI, fasting insulinemia, and HOMA-IR [17]. If individuals with T2D also display greater fasting GLP-1 concentrations, this may lead to de-sensitisation of GLP-1 releasing L-cells, leading to lower meal-induced GLP-1 release [19]. This could also have important implications for the use of total area under the curve (tAUC) to measure meal-induced GLP-1 release, since this may be influenced to a major extent by variations in the fasting concentrations.

Such variability in the literature regarding GLP-1 secretion in individuals with T2D may also suggest the presence of moderator variables. One such variable may be related to meal type, given the wide array of signalling pathways involved in GLP-1 secretion. Glucose-mediated GLP-1 secretion is not known to be dependent on receptor activation, but appears to depend on absorption *via* SGLT-1 [20–22], whereas fat- and protein- mediated GLP-1 secretion relies primarily on the activation of a number of nutrient sensors and transporters expressed in enteroendocrine cells [23–25].

Another important factor is the measurement of GLP-1 which is not straightforward due to its rapid and extensive degradation in the circulation, the low concentrations, and the specificity required of assays to measure the relevant GLP-1 isoforms [2]. GLP-1_{TOTAL}, comprising of the intact form and the primary metabolite, is therefore a more appropriate measure compared to the rapidly degraded intact GLP-1 moiety for estimations of secretion. GLP-1_{TOTAL} can be measured using single antibody assays with antibodies specific for the amidated C-terminus [26]. Because of the importance of antisera directed against specific GLP-1 epitopes, the majority of studies employ a well-defined radioimmunoassay [27]. Commercially available antisera and assay kits are often poorly characterised which limits the certainty of which GLP-1 isoforms are detected [28]. As a result, commercially available assays tend to exhibit varying specificity [29], which suggests the type of assay may influence the quantified difference in GLP-1 secretion between individuals with and without T2D. A particular problem applies to the use of the sandwich ELISA which, despite its potentially superior sensitivity, depends on two binding epitopes of the molecule. Given the extensive degradation of GLP-1 from both termini, this almost invariably affects the binding of the antibodies, which is why the assays exhibit serious variability in specificity for the various isoforms [2]. Another potential problem is sample pre-treatment. Some assays [9,30] use plasma extraction to remove potentially interfering substances. This pre-treatment approach may markedly influence both fasting levels and meal responses [2].

The aim of the present review was to synthesise the published literature in order to determine whether GLP-1 secretion is reduced in

individuals with *versus* without T2D in response to oral glucose and/or mixed meal ingestion, and whether there are any differences in fasting GLP-1 concentrations between study samples. The putative roles of potential mediating factors (e.g. meal composition and form, and assay type) were also explored.

2. Methods

This systematic review and meta-analysis was undertaken in alignment with the PRISMA (Preferred Reporting Items for Systematic Review and Meta-analyses) guidelines [31], and was prospectively registered with PROSPERO (CRD42020195612).

2.1. Literature search

Eligible studies were identified by electronic and manual searches in literature references. Electronic searches were conducted *via* the Web of Science, PubMed, Cochrane Library, and Embase from July 2020 through November 2022. The search terms included (“glucagon-like peptide-1” OR “GLP-1”) AND (“secretion” OR “release”) AND (“type 2 diabetes”) AND (“human*” OR “men” OR “man” OR “women”) AND (“OGTT” OR “MMTT” OR “protein” OR “fat” OR “carbohydrate” OR “meal” OR “postprandial”). Eligible studies were assessed independently by two authors according to inclusion criteria. There were no restrictions of the publication date of studies, however studies were required to be in English or able to be translated.

2.2. Inclusion and exclusion criteria

In general, clinical trials measuring GLP-1_{TOTAL} responses to a test meal in individuals with T2D *versus* individuals without T2D were required. For inclusion, studies had to meet the following requirements:

- participants were over the age of 18
- participants did not have >1 comorbidity unrelated to metabolic syndrome
- Individuals with T2D were clinically diagnosed with fasting glucose and/or 2-h glucose >7.0 and >11.1 mmol·L⁻¹, respectively (American Diabetes Association, 2015)

Furthermore, studies were excluded if they:

- used non-specific assays which cross-react with the major proglucagon fragment
- used intact GLP-1 measurements only
- were not peer reviewed
- did not have enough data

All studies were required to involve measurements of either fasting plasma GLP-1 concentrations, or plasma GLP-1_{TOTAL} iAUC in response to a meal challenge in individuals clinically diagnosed with T2D and individuals without T2D. Studies that employed an intravenous infusion in parallel with a test meal were considered eligible if they included a placebo infusion condition and studies involving weight-loss surgery were included if they employed a test meal for a group with T2D and a matched control group prior to surgery. Two researchers (JW and SC) independently assessed studies for inclusion and any disagreements were resolved by a third reviewer (JG). Potential studies that could be included based on their title or abstract were retrieved in full-text and reviewed against the inclusion/exclusion criteria independently by JW and SC. Similarly, JG was consulted in the event of any disputes. In total, 32 studies met criteria for inclusion and were included in this systematic review and meta-analysis (Fig. 1).




Fig. 1. PRISMA flow diagram of the study screening and selection process.

2.3. Data extraction

From eligible published studies, data (mean and SD) were extracted independently by two researchers (JW and SC). Data were extracted into a standardised spreadsheet, which included characteristics of articles, outcome data, and conditions for the Cochrane collaboration's tool for assessing risk of bias [32]. The participant characteristics, assay employed, meal characteristics, fasting plasma glucose and GLP-1, and plasma GLP-1_{TOTAL} iAUC were collected. Where it was not possible to extract this information, authors were contacted for clarification.

Where GLP-1_{TOTAL} values were reported in $\text{pg}\cdot\text{ml}^{-1}$, values were converted to $\text{pmol}\cdot\text{L}^{-1}$ by dividing by 4. When values were only presented in figure form, the figure was digitised and the means and SD/SE were measured manually to the scale provided on the figure. SEs were converted to SDs using $SD = SE \sqrt{n}$.

2.4. Assessment of risk of bias in included studies

The Cochrane risk of bias tool was implemented to assess internal validity of eligible studies [32] independently by two researchers (JW and SC). Each study was assessed in the following 6 domains: incomplete outcome data, selective reporting, group similarity at baseline, intention to treat analysis, timing of outcome assessments, and other bias e.g. conflicts of interest, sample size. Each domain was judged to be either high, low or unclear (if insufficient detail was provided) risk. Disagreements were resolved by a third reviewer (JG). The following domains were not deemed appropriate for the research designs of included studies (because the outcome measure is not subject to conscious bias): random sequence generation, allocation concealment, blinding of

participants and personnel, blinding of outcome assessment, and co-intervention and compliance bias and were omitted from the risk of bias (Fig. 2).

2.5. Statistical analysis

Missing standard deviations were estimated from reported standard errors or confidence intervals according to Cochrane Handbook guidelines. [33] Reported mean intervention effects (quantified using the Hedges g standardised mean difference) were pooled in a random effects meta-analysis using Stata 16 software (StataCorp LLC, TX, USA). Random effects meta-analysis models, rather than a fixed effects models, were selected *a priori* based on the philosophical absence of a consistent “true effect” across the studies [34]. A random effects model was selected as it was more realistic and appropriate to assume that the true effect could vary from study to study. Nevertheless, fixed effects sensitivity analyses were undertaken as these can help interpret the seriousness of any small study effects (publication bias) that are detected [33].

Data entry was carried out for all results by JW & SC and peer-reviewed by GA. Study effect sizes were weighted using the inverse variance approach. The restricted maximum likelihood approach was adopted with the Knapp-Hartung modification to standard errors applied [35]. Ninety-five percent confidence intervals (CIs) and prediction intervals (PI) were calculated for group differences in GLP-1 iAUC and fasting GLP-1. The PI is better aligned to the selection of a random effects model than a CI. A PI quantifies the likely range in which a new study's effect size will fall into, assuming that this future study is of a similar nature to those meta-analysed [15]. A meta-analysis was




Fig. 2. Judgement about each risk of bias domain presented as a percentage across all included studies.

undertaken only where there were at least five studies to ensure minimally acceptable statistical power and relatively precise inferences [36]. I-squared and tau statistics were used to quantify between-study

heterogeneity in the meta-analyses. Tau (T) is the estimated standard deviation of underlying true effects across studies [34]. “Double counting” is an important consideration in any meta-

Table 1 Study characteristics.

| Reference | Outcomes | Sample size | | Meal test | Meal dose (kcal) | Duration of test (min) | Assay |
|------------------------------|---------------|---------------|--------------|----------------------------|------------------|------------------------|------------------------|
| | | T2D (n = 628) | C (n = 1409) | | | | |
| Plourde et al. 2014 [39] | Fasting, iAUC | 9 | 9 | MMTT ^b | 295 | 180 | ELISA |
| Alsalm et al. 2018 [40] | Fasting | 12 | 12 | – | – | 400 | ELISA |
| Nauck et al. 2016 [41] | iAUC | 32 | 29 | MMTT ^b | 541 | 240 | RIA |
| Faerch et al. 2015 [10] | Fasting | 163 | 774 | – | – | 120 | RIA |
| Kozawa et al. 2010 [42] | Fasting | 4 | 5 | – | – | 180 | RIA |
| Muscelli et al. 2008 [43] | Fasting | 10 | 24 | – | – | 180 | RIA |
| Bagger et al. 2011 [44] | Fasting | 8 | 8 | – | – | 240 | RIA |
| Alsalm et al. 2016 [45] | Fasting | 18 | 18 | – | – | 300 | RIA |
| Jørgensen et al. 2012 [46] | Fasting, iAUC | 13 | 12 | MMTT ^a | 300 | 240 | RIA |
| Knop et al. 2007a [47] | Fasting, iAUC | 8 | 8 | OGTT ^a | 200 | 240 | RIA |
| Vilsbøll et al., 2001 [48] | Fasting | 12 | 12 | – | – | 180 | RIA |
| Sonne et al. 2013 [49] | Fasting, iAUC | 15 | 15 | OGTT ^a | 300 | 240 | RIA |
| | | | | Low Fat MMTT ^a | 500 | | |
| | | | | Mod Fat MMTT ^a | 500 | | |
| | | | | High Fat MMTT ^a | 500 | | |
| Toft-Nielsen et al. 2001 [9] | Fasting, iAUC | 54 | 33 | MMTT ^b | 538 | 240 | RIA |
| Theodorakis et al. 2006 [50] | Fasting | 17 | 36 | – | – | 120 | RIA |
| Højberg et al. 2008 [51] | Fasting, iAUC | 9 | 9 | MMTT ^b | 538 | 240 | RIA |
| Yabe et al. 2015 [52] | Fasting | 28 | 54 | – | – | 120 | RIA |
| Ryskjær et al. 2006 [53] | Fasting, iAUC | 8 | 8 | MMTT ^b | 566 | 180 | RIA |
| Astiarraga et al. 2018 [54] | Fasting | 13 | 10 | – | – | 180 | ELISA |
| Martinussen et al. 2015 [55] | Fasting, iAUC | 10 | 12 | MMTT ^a | 300 | 180 | RIA |
| Tricò et al. 2015 [56] | Fasting, iAUC | 10 | 12 | OGTT ^a | 300 | 120 | Mesoscale ^c |
| Alssema et al. 2013 [57] | Fasting, iAUC | 20 | 163 | OGTT ^a | 300 | 120 | RIA |
| | | | | MMTT ^b | 833 | | |
| Lund et al. 2015 [58] | Fasting, iAUC | 15 | 18 | OGTT ^a | 100 | 120 | RIA |
| Ruetten et al. 2018 [59] | Fasting, iAUC | 22 | 23 | MMTT ^b | 470 | 120 | RIA |
| Knop et al., 2007b [60] | Fasting, iAUC | 10 | 10 | OGTT ^a | 200 | 240 | RIA |
| Rohde et al. 2017 [61] | iAUC | 9 | 10 | MMTT ^a | 525 | 240 | RIA |
| Vaag et al., 1996 [62] | Fasting, iAUC | 12 | 13 | OGTT ^a | 300 | 180 | RIA |
| Ahrén et al., 2000 [63] | Fasting | 6 | 6 | – | – | 120 | RIA |
| Kuwata et al. 2016 [64] | Fasting | 12 | 10 | – | – | 240 | Mesoscale ^c |
| St-Jean et al., 2017 [65] | Fasting | 8 | 9 | – | – | 360 | ELISA |
| Yan et al. 2014 [66] | Fasting, iAUC | 10 | 10 | MMTT ^a | 75 | 180 | ELISA |
| | | | | MMTT ^a | 150 | | |
| | | | | MMTT ^a | 300 | | |
| Purnell et al. 2018 [67] | Fasting | 40 | 22 | – | – | 240 | Mesoscale ^c |
| Greenfield et al. 2009 [68] | Fasting, iAUC | 4 | 8 | OGTT ^a | 300 | 120 | RIA |

T2D, Type 2 Diabetes, C, Control group, OGTT, Oral glucose tolerance test, MMTT, Mixed meal tolerance test, RIA, Radioimmunoassay, ELISA, enzyme-linked immunosorbent assay, iAUC, incremental area under the curve.

^a Liquid meal.

^b Mixed meal.

^c Mesoscale provides sandwich immunoassays that measure GLP-1 by electrochemiluminescence.

analysis and refers to problems due to including the same study, or groups from the same study, more than once in the meta-analysis, leading to spurious statistical precision [16]. We addressed this problem by pooling multiple study groups from the same study using the approach reported in the Cochrane Handbook [33].

In the absence of definitive information about the clinical relevance of a unit change in GLP-1 secretion on morbidity and mortality [37] we used thresholds of standardised mean difference to interpret the magnitude of any pooled effects. These thresholds were interpreted as follows; very small <0.2, small 0.2–0.3, moderate 0.4–0.8, and large >0.8 [38]. A two-tailed *p* value of ≤0.05 was deemed statistically significant.

Sub-group analyses were undertaken to explore whether meal type, meal form and assay type influenced study effect sizes. A random effects multivariable meta-regression was undertaken to explore the influence of fasting glucose, meal dose, age and BMI on study effect sizes.

3. Results

3.1. Study selection

A summary of data searching and extraction is provided in Fig. 1. In total, 6011 records were identified, of which 3285 records remained following the removal of duplicates. Following title and abstract screening, 72 full texts were retrieved and evaluated against the inclusion criteria. Consequently, a further 40 studies were excluded and, therefore, 32 studies remained for analysis in our review. The various study characteristics of included studies are presented in Table 1.

3.2. Methodological quality

The risk of bias for all studies was generally low across studies (Fig. 2). However, the areas of greatest risk of bias appeared to be in the

domains for group matching and other biases (low sample sizes). The most common bias appeared to be inequalities in sex, age, and BMI between study groups. Notably, a large number of studies (~33 % of whole sample of studies) recruited <10 participants for each group which decreases the precision for estimate of effect size in each of these studies. While the small samples sizes and poor matching are problematic for the individual studies, these are precisely the limitations that can be addressed by collating conceptually similar studies in a meta-analysis such as the present one.

3.3. GLP-1 iAUC

Data from 270 individuals with T2D and 402 individuals without T2D were pooled from 18 studies with 25 datasets (pooled to avoid double counting of studies). Following the random-effects meta-analysis, we found that the pooled standardised mean difference in GLP-1 iAUC between individuals with and without T2D was not statistically significant (*p* = 0.43) and very small in magnitude (according to our selected thresholds), and the prediction interval for this pooled mean difference was very wide (Fig. 3; *g* = -0.15 [very small], PI [-1.53, 1.23]). The mean absolute values of GLP-1 iAUC were 1047 ± 930 and 1204 ± 938 pmol·L⁻¹·min for individuals with and without T2D, respectively. A regression-based Egger test did not show any clear evidence of small study effects (Fig. 4; *p* = 0.33). Subgroup analyses (See supplementary file) demonstrated no meaningful or statistically significant sub-group effects in GLP-1 iAUC when data were separated by meal type (*p* = 0.39) or meal form (*p* = 0.72). However, there was a large effect of assay type, whereby GLP-1 iAUC was greater in individuals with T2D compared to individuals without T2D when using ELISA or Meso-scale (*g* = 0.67 [moderate], *p* = 0.009, 95 % CI [0.41, 0.93]), but not when using RIA (*g* = -0.30 [small], *p* = 0.10, 95 % CI [-0.72, 0.12]). The difference between these two effect sizes was statistically significant (*p* = 0.002). A random-effects multivariable meta-regression of GLP-1




Fig. 3. Forest plot of GLP-1 iAUC (pmol·L⁻¹·min) for individuals with T2D (treatment) and individuals without T2D (control). A negative standardised mean difference is indicative of mean GLP-1 iAUC being lower in T2D than non-T2D.




Fig. 4. Funnel plot for studies that presented GLP-1 iAUC. A negative Hedges' g is indicative of mean GLP-1 iAUC being higher in T2D than non-T2D.

iAUC suggested that BMI, age, fasting glucose, and meal dose did not influence the GLP-1 iAUC effect sizes (all $p > 0.05$, See Supplementary File). Heterogeneity between studies was high ($I^2 = 74.4\%$; $T^2 = 0.39$). We undertook a sensitivity analysis whereby a fixed effects model was selected. The pooled effect size from this model was a similar -0.17 (95%CI: -0.33 to 0.002 , $p = 0.052$).

3.4. Fasting GLP-1

Data from 580 individuals with T2D and 1363 individuals without T2D were pooled from 30 studies. This random-effects meta-analysis demonstrated that the standardised mean difference in fasting GLP-1 between individuals with T2D versus individuals without T2D was not statistically significant ($p = 0.21$) and small in magnitude (according to our selected thresholds). Again, the prediction interval was wide indicating that a future study of the same nature could result in large effect sizes in either direction (Fig. 5; $g = 0.24$ [small], PI [-1.55 , 2.02]). Mean fasting GLP-1 was 16.2 ± 6.9 and 12.4 ± 5.7 pmol·L⁻¹ for individuals with and without T2D, respectively. A regression-based Egger test did show some evidence of publication bias due to asymmetry of small studies where one very large negative effect size ($g = -3.9$) was reported (Fig. 6; $p = 0.08$). Subgroup analyses demonstrated no differences in fasting GLP-1 between groups when accounting for assay type ($p > 0.05$). Random-effects meta-regression of fasting GLP-1 suggested that BMI, age, and fasting glucose did not affect the fasting GLP-1 outcome. Heterogeneity between studies was high ($I^2 = 87.7\%$; $T^2 = 0.73$).

Although Muscelli et al. (2008) reported that their data were presented as mean and standard deviation (SD), in one of their Figure legends they reported that mean and standard error (SE) were presented. It was, therefore, not wholly certain whether the SDs we extracted for the fasting GLP-1 data were, in fact, standard errors (SE). Therefore, a sensitivity analysis was conducted on the basis that error bars in the relevant Figure represented SEs rather than SDs. We then calculated the SD using the equation $=SE \sqrt{n}$. After this was applied, the meta-analysis pooled SMD was found to be 0.25 (PI: -0.28 , 0.78), which was similar to the original estimate. We also undertook a sensitivity analysis whereby a fixed effects model was selected. The effect size from this model was a similar 0.26 (95%CI: 0.15 to 0.37 , $P < 0.0001$). This similarity indicates that the small study effects we detected for fasting

GLP-1 are not serious [33].

4. Discussion

The findings from this systematic review demonstrate that the standardised mean differences in postprandial GLP-1 secretion and fasting GLP-1 between individuals with versus without T2D are very small to small when compared with traditional thresholds for standardised effects. Furthermore, this analysis revealed that studies utilising ELISA or sandwich immunoassays tended to report higher GLP-1 iAUC in individuals with T2D compared to individuals without T2D, which was not apparent with studies utilising RIAs. Importantly, it is difficult to make judgement about the biological relevance of the pooled standardised effect sizes partly because of the absence of robust “anchors” to morbidity risk and partly because of the wide prediction intervals obtained for the effect sizes.

Similar to previous meta-analyses, in the present study, there was no clear difference in plasma GLP-1 iAUC between individuals with and without T2D [13,14]. Despite this, some of the largest single studies to date have reported lower GLP-1 secretion in individuals with versus without T2D [9,10]. Large heterogeneity between studies resulting in positive, neutral, and negative effect sizes may partly explain why differences in SMDs between groups were very small to small. Additionally, important moderating variables such as BMI and visceral fat content may vary largely within groups. These differences within and between studies present a challenge for determining the most important factors influencing GLP-1 secretion. The present analysis suggests that diagnosis of type 2 diabetes is not associated with large differences in GLP-1 secretion, and thus these other moderating variables may be more important than diabetes diagnosis *per se*.

There were no clear differentiating effects of meal type *i.e.*, OGTT or MMTT, or meal form *i.e.*, liquid or mixed, on GLP-1 iAUC. Given the diversity of intestinal receptors/transporters that are specific to different nutrients it was hypothesised that a potential reduction in GLP-1 secretion in individuals with T2D may occur as a result of receptor/transporter resistance to a specific macronutrient(s), thereby manifesting in different responses based on whether an OGTT or MMTT was administered. While these results do not support this hypothesis, there have not been enough studies to isolate the specific effects of different




Fig. 5. Forest plot of fasting GLP-1 (pmol·L⁻¹) for individuals with T2D (treatment) and individuals without T2D (control). A negative standardised mean difference is indicative of mean fasting GLP-1 being lower in T2D than non-T2D.

macronutrients on GLP-1 release in T2D and it is likely that MMTTs incorporating a combination of macronutrients may have masked any specific nutrient resistance that may have influenced GLP-1 secretion. In 33 participants of normal weight, overweight and obesity, GLP-1, oxyntomodulin and glicentin were increased to a greater extent following lipid *versus* glucose ingestion, with variation in size of response between these peptides following the ingestion of different macronutrients [69]. Moreover, It was recently demonstrated that the exaggerated GLP-1 response to meal intake after gastric bypass operations is mainly due to the effect of carbohydrates [70]. While rate of gastric emptying in the instance of gastric bypass may account for an exaggerated GLP-1 response, a recent paper suggested high levels of inter-individual variation for GLP-1 secretion following the ingestion/administration of different macronutrients, independent of gastric emptying [71]. Thus, individual food items may affect GLP-1 (and other peptides) secretion differentially and this may be impacted by certain metabolic conditions including T2D.

One factor that did appear to influence the estimates of GLP-1 secretion was the assay type employed. As mentioned in the introduction, reliable measurement of GLP-1 is difficult due to the existence of a number of additional proglucagon (GCG) -derived fragments and GLP-1 moieties which can cross-react with the measurement of the intended form of GLP-1 [2,26]. A well-defined RIA previously described

by [27] was utilised by the majority of studies, while only 5 out of 25 studies utilised ELISA or sandwich immunoassay (Mesoscale) to measure GLP-1 iAUC. Interestingly, the studies using ELISA/sandwich immunoassay demonstrated greater GLP-1 iAUC in individuals with T2D *versus* those without T2D, whereas no such difference was apparent in studies that measured GLP-1 by RIA. Previous research comparing the specificity of many commercially available assays suggests the ability of the mesoscale immunoassay to detect non-active forms is incomplete and the recovery of non-amidated forms when using the Merck Millipore ELISA is variable [29]. One limitation of the assay type sub-analyses is that it was only possible to group assays into RIA's or ELISA/mesoscale. The findings from [29] suggest that the reliability and accuracy of ELISAs and RIAs can vary by manufacturer/antibody selection. These findings suggest that choice of assay is an important consideration and should be based on the research question and conditions of the study, but unless there is greater consistency of assay use between studies it will remain difficult to compare findings between studies. It is possible that new, more accurate, methods for measuring GLP-1, including highly specific ELISAs and methods based on mass spectrometry, will become more widely available in the not too distant future. A recent sandwich ELISA recognising both GLP-1_{9-36NH2} and GLP-1₉₋₃₇ has been developed and validated [72], and a new ELISA assay (from Ansh) has been reported to show specificity to a number of proglucagon-derived




Fig. 6. Funnel plot for studies that presented fasting GLP-1. A negative Hedges' g is indicative of mean fasting GLP-1 being lower in T2D than non-T2D.

peptides including oxyntomodulin, glicentin, glucagon and MPGF [73].

To our knowledge this is the first systematic review to investigate fasting GLP-1 in studies involving individuals with and without T2D. The findings of the present study suggest that differences in fasting GLP-1 between these two groups are small (Hedges' g). Unlike for GLP-1 iAUC, the assay type had no clear influence on fasting GLP-1. This suggests that assay type may make more of a difference at higher concentrations of GLP-1, and/or that assays are not sensitive or accurate enough to determine the low circulating fasting concentrations regardless of the presence of T2D or not. The prediction interval generated suggests that any future study comparing fasting GLP-1 between individuals with and without T2D could find greater, equal, or lower fasting GLP-1 in individuals with *versus* without T2D. Nevertheless, the Egger's test demonstrated evidence of publication bias where there was an asymmetry of effect sizes reported by included studies. This asymmetry related to the existence of a few large negative effect sizes. The majority of studies reported positive effect sizes indicating that fasting GLP-1 was greater in individuals with T2D compared to individuals without T2D (16.0 ± 6.8 and 12.3 ± 5.7 pmol·L⁻¹, respectively) which is in agreement with the mean fasting glucose for individuals with and without T2D. However, the positive effect sizes were mostly small and large prediction intervals suggest uncertainty in future studies comparing GLP-1 concentrations in individuals with T2D compared to without T2D.

The studies included in the present systematic review and meta-analyses were highly heterogenous according to the I^2 and T^2 statistic, and the wide prediction intervals. Despite all included studies being conceptually similar, large between-study differences in specific methodologies are a limitation for the present systematic review and may explain why there is such conflicting evidence in the literature. Interestingly, similarly high study heterogeneity was reported in a recent meta-analysis which showed a variable impairment in the incretin effect in T2D [74]. Taken together, this highlights the importance of acknowledging possible sources of study heterogeneity in the GLP-1 related literature, and these are addressed below. Firstly, a number of included studies were at high risk of bias due to small sample sizes and/or differences between groups at baseline. It could be possible that these studies were underpowered to detect a genuine effect and that subsequent variability in these studies contributed to large PIs. Additional

inconsistency may come from meal ingestion. Most commonly, an OGTT or MMTT is ingested to measure GLP-1 responses, however within each type of test meal there is large variability in terms of caloric dose and/or macronutrient composition of the meal consumed. Therefore, grouping studies into those that fed an OGTT and/or MMTT may not be a truly matched comparison. Another confounding factor is the varying relationship between the meal size and the body mass of the participants. It cannot be excluded that identical meals will variably affect GLP-1 responses in individuals with large differences in body mass. In general, GLP-1 responses have been smaller in people with overweight [10], and weight loss has been reported to restore responses [74]. Thus, overweight may be an important factor for reduced GLP-1 responses in patients with T2DM, most of whom are also overweight/obese [2].

Within the literature, there is also poor consistency for reporting GLP-1 secretion. In the present meta-analyses, a number of studies were excluded as the units provided for AUC could not be interpreted. Units may differ based on whether AUCs are time-averaged and also by the length of the postprandial period. To avoid uncertainty and improve the comparability of studies it would be helpful for authors to report their data clearly and thoroughly. For example, authors could report AUC for each segment of the postprandial period separately to allow greater comparison and insight between studies. Tools such as the Time Series Response Analyser [75] would streamline this process. Further issues surround the grouping of individuals with impaired glucose tolerance and T2D. This makes it difficult to separate the physiology of GLP-1 secretion in individuals with a clinical diagnosis of T2D and those with deteriorating glucose control. Indeed, a study with a comparatively large group of people with T2D and controls also included a group with impaired glucose tolerance, and their GLP-1 responses were positioned between those of the controls and individuals with T2D [9]. Lastly, studies are generally poor at describing whether diabetes medication is withheld during data collection, and if so, for how long. This is particularly important considering some common medication, such as metformin, has been shown to increase GLP-1 secretion [76,77], and could therefore influence some of the effect sizes reported. Based on the thorough review of studies reporting GLP-1 secretion in T2D, considerations for the comparability of research studies and recommended solutions are provided in Table 2.

In conclusion, differences in fasting GLP-1 and GLP-1 iAUC between

Table 2
Considerations for GLP-1 literature and recommended solutions.

| Issue within literature | Consequence of this issue | Recommended solution |
|--|---|---|
| Unclear/inconsistent use of AUC and appropriate units | Difficulty comparing across studies, introduces risk of misinterpretation | Comprehensive reporting of method and units |
| Use of tAUC only, to measure meal-induced GLP-1 release | Total area under the curve is not the most suitable measure for postprandial change in GLP-1 secretion as it does not consider the change relative to baseline. This also introduces the possibility of overestimating GLP-1 release in groups with higher fasting GLP-1 relative to the comparator group(s). | Both measures add value. Consider the use of both total and incremental area under the curve when reporting postprandial responses |
| Inconsistent postprandial period | There is evidence of early and/or late phase GLP-1 secretion differences in T2D and inappropriate postprandial sampling protocols (i.e. too short, too infrequent) may limit full characterisation of postprandial GLP-1 responses. | While the length of the postprandial period may differ dependent on context, a standard approach could be to sample at least every 30 min, and report AUC data for each segment. |
| Selection of assay | Circulating concentrations of GLP-1 are in the low picomolar range. The existence of a number of preproglucagon fragments which can cross-react with the intended measure of GLP-1 | Carefully consider research aims, choose a well-defined assay that includes key information on the use of antisera and the forms of GLP-1 captured. |
| Measure of active/inactive GLP-1 | Rapid degradation in the circulation. Concentrations potentially below limit of detection of assays | Measure total GLP-1, which incorporates the intact form and metabolite. |
| Grouping of T2D with individuals with impaired glucose tolerance | If GLP-1 secretion is influenced by degree of glucose control, it is likely that data will be variable. Difficult to identify a true effect in individuals with T2D. | Group by T2D classification or if possible, treat glucose tolerance as a continuous variable. |
| Poor reporting of medication | Information regarding the use of medication and whether medication is withheld during data collection is inconsistent. Certain medication e.g. Metformin, can increase GLP-1 secretion, which can confound the true effect of meal-induced GLP-1 release in T2D. | Authors should clearly report the type of medication individuals are using, the dose/frequency, and also whether or not they withhold medication during data collection, and if so, for how long. |
| Poor reporting of meal characteristics | Level of detail (such as type, form and composition) provided for ingested meals is often inconsistent. Missing information can introduce risk of misinterpretation. | Authors should clearly report the type of meal fed i.e. OGTT, MMTT, the type and form of food ingested, as well as the macronutrient composition and calorie content. |

individuals with T2D and individuals without T2D were small and inconsistent. This suggests that reduced beta-cell responsiveness to GIP and/or GLP-1 as opposed to reduced GLP-1 secretion may provide a better explanation for the reduced incretin effect in T2D. When studies employed either ELISA or sandwich immunoassay (Mesoscale), GLP-1 iAUC was higher in individuals with T2D compared to individuals without T2D. This demonstrates the difficulty surrounding GLP-1 measurement and the importance of using a well-defined assay. Evidence of high heterogeneity between studies, small sample sizes, and poor matching of groups may help to explain the wide prediction intervals

and conflicting evidence in the literature surrounding GLP-1 secretion in T2D. Recommendations for improving comparisons for GLP-1 secretion between individuals with and without T2D were outlined, as well as potential mediators more important than T2D diagnosis *per se*.

Declaration of competing interest

JW has received research funding as part of their PhD studentship from Arla Foods Ingredients (DK). SC, and GA have no conflicts of interest. JJH is a member of the advisory board to Novo Nordisk. FK is supported by MRC (UK) funding (MR/P002927/1). JAB is an investigator on research grants funded by BBSRC, MRC, British Heart Foundation, Rare Disease Foundation, EU Hydration Institute, GlaxoSmithKline, Nestlé, Lucozade Ribena Suntory, ARLA foods, Kennis Centrum Suiker and Salus Optima (L3M Technologies Ltd); has completed paid consultancy for PepsiCo, Kellogg's, SVGC and Salus Optima (L3M Technologies Ltd); is Company Director of Metabolic Solutions Ltd.; receives an annual honorarium as a member of the academic advisory board for the International Olympic Committee Diploma in Sports Nutrition; and receives an annual stipend as Editor-in Chief of *International Journal of Sport Nutrition & Exercise Metabolism*. J.T.G. is an investigator on research grants funded by BBSRC, MRC, British Heart Foundation, The Rank Prize Funds, The European Society for Clinical Nutrition and Metabolism (ESPEN), Lucozade Ribena Suntory, ARLA Foods Ingredients, Kenniscentrum Suiker and Voeding and Clasado Biosciences; and has completed paid consultancy for PepsiCo and SVGC.

References

- Holst JJ. The physiology of glucagon-like peptide 1. *Physiol Rev* Oct 2007;87(4): 1409–39. <https://doi.org/10.1152/physrev.00034.2006> (in eng).
- Holst JJ, Albrechtsen NJW, Rosenkilde MM, Deacon CF. Physiology of the incretin hormones, GIP and GLP-1-regulation of release and posttranslational modifications (in eng). *Compr Physiol* 2019;9(4):1339–81. <https://doi.org/10.1002/cphy.c180013.09>.
- Flint A, Raben A, Astrup A, Holst JJ. Glucagon-like peptide 1 promotes satiety and suppresses energy intake in humans. *J Clin Invest* Feb 1998;101(3):515–20. <https://doi.org/10.1172/JCI990> (in eng).
- Wettergren A, Schjoldager B, Mortensen PE, Myhre J, Christiansen J, Holst JJ. Truncated GLP-1 (proglucagon 78–107-amide) inhibits gastric and pancreatic functions in man. *Dig Dis Sci* Apr 1993;38(4):665–73. <https://doi.org/10.1007/bf01316798> (in eng).
- Nauck MA, Stöckmann F, Ebert R, Creutzfeldt W. Reduced incretin effect in type 2 (non-insulin-dependent) diabetes. *Diabetologia* 1986;29(1):46–52. <https://doi.org/10.1007/BF02427280>.
- Nauck MA, Heimesaat MM, Orskov C, Holst JJ, Ebert R, Creutzfeldt W. Preserved incretin activity of glucagon-like peptide 1 [7–36 amide] but not of synthetic human gastric inhibitory polypeptide in patients with type-2 diabetes mellitus. *J Clin Invest* Jan 1993;91(1):301–7. <https://doi.org/10.1172/JCI116186> (in eng).
- Nauck MA, Klein N, Orskov C, Holst JJ, Willms B, Creutzfeldt W. Normalization of fasting hyperglycaemia by exogenous glucagon-like peptide 1 (7–36 amide) in type 2 (non-insulin-dependent) diabetic patients. *Diabetologia* Aug 1993;36(8):741–4. <https://doi.org/10.1007/bf00401145> (in eng).
- Kjems LL, Holst JJ, Vølund A, Madsbad S. The influence of GLP-1 on glucose-stimulated insulin secretion: effects on beta-cell sensitivity in type 2 and nondiabetic subjects. *Diabetes* Feb 2003;52(2):380–6. <https://doi.org/10.2337/diabetes.52.2.380> (in eng).
- Toft-Nielsen MB, et al. Determinants of the impaired secretion of glucagon-like peptide-1 in type 2 diabetic patients. *J Clin Endocrinol Metab* Aug 2001;86(8): 3717–23. <https://doi.org/10.1210/jcem.86.8.7750> (in eng).
- Faerch K, et al. GLP-1 response to oral glucose is reduced in prediabetes, screen-detected type 2 diabetes, and obesity and influenced by sex: the ADDITION-PRO study. *Diabetes* Jul 2015;64(7):2513–25. <https://doi.org/10.2337/db14-1751> (in eng).
- Vollmer K, et al. Predictors of incretin concentrations in subjects with normal, impaired, and diabetic glucose tolerance. *Diabetes* 2008;57(3):678. <https://doi.org/10.2337/db07-1124>.
- Romero F, et al. Comparable early changes in gastrointestinal hormones after sleeve gastrectomy and Roux-En-Y gastric bypass surgery for morbidly obese type 2 diabetic subjects. *Surg Endosc* Aug 2012;26(8):2231–9. <https://doi.org/10.1007/s00464-012-2166-y> (in eng).
- Nauck MA, Vardarli I, Deacon C, Holst J, Meier J. Secretion of glucagon-like peptide-1 (GLP-1) in type 2 diabetes: what is up, what is down? *Diabetologia* 2011; 54(1):10–8. <https://doi.org/10.1007/s00125-010-1896-4>.
- Calanna S, et al. Secretion of glucagon-like peptide-1 in patients with type 2 diabetes mellitus: systematic review and meta-analyses of clinical studies. *Diabetologia* 2013;56(5):965–72. <https://doi.org/10.1007/s00125-013-2841-0>.

- [15] IntHout J, Ioannidis JP, Rovers MM, Goeman JJ. Plea for routinely presenting prediction intervals in meta-analysis (in eng) *BMJ Open* 2016;6(7):e010247. <https://doi.org/10.1136/bmjopen-2015-010247>. 07 12.
- [16] Senn SJ. Overstating the evidence: double counting in meta-analysis and related problems (in eng) *BMC Med Res Methodol* Feb 13 2009;9(10). <https://doi.org/10.1186/1471-2288-9-10>.
- [17] Stinson SE, et al. Fasting plasma GLP-1 is associated with overweight/obesity and cardiometabolic risk factors in children and adolescents. *J Clin Endocrinol Metab* 2021;106(6):1718–27. <https://doi.org/10.1210/clinem/dgab098>.
- [18] Muñoz JSG, Rodríguez DJ, Morante JH. Diurnal rhythms of plasma GLP-1 levels in normal and overweight/obese subjects: lack of effect of weight loss. *J Physiol Biochem* 2015;71(1):17–28. <https://doi.org/10.1007/s13105-014-0375-7>. 2015/03/01.
- [19] Muscelli E, et al. Impact of incretin hormones on β -cell function in subjects with normal or impaired glucose tolerance. *Am J Physiol Endocrinol Metab* 2006;291(6):E1144–50. <https://doi.org/10.1152/ajpendo.00571.2005>. 2006/12/01.
- [20] Moriya R, Shirakura T, Ito J, Mashiko S, Seo T. Activation of sodium-glucose cotransporter 1 ameliorates hyperglycemia by mediating incretin secretion in mice. *Am J Physiol Endocrinol Metab* Dec 2009;297(6):E1358–65. <https://doi.org/10.1152/ajpendo.00412.2009> (in eng).
- [21] Reimann F, Gribble FM. Glucose-sensing in glucagon-like peptide-1-secreting cells. *Diabetes Sep* 2002;51(9):2757–63 (in eng).
- [22] Kuhre RE, Frost CR, Svendsen B, Holst JJ. Molecular mechanisms of glucose-stimulated GLP-1 secretion from perfused rat small intestine. *Diabetes Feb* 2015;64(2):370–82. <https://doi.org/10.2337/db14-0807> (in eng).
- [23] Spreckley E, Murphy KG. The I-cell in nutritional sensing and the regulation of appetite. *Front Nutr* 2015;2:23. <https://doi.org/10.3389/fnut.2015.00023> (in eng).
- [24] Bodnaruc AM, Prud'homme D, Blanchet R, Giroux I. Nutritional modulation of endogenous glucagon-like peptide-1 secretion: a review. *Nutr Metab December* 09 2016;13(1):92. <https://doi.org/10.1186/s12986-016-0153-3> (Lond.), journal article.
- [25] Modvig IM, Kuhre RE, Holst JJ. Peptide-mediated glucagon-like peptide-1 secretion depends on intestinal absorption and activation of basolaterally located calcium-sensing receptors. *Physiol Rep Apr* 2019;7(8):e14056. <https://doi.org/10.14814/phy2.14056> (in eng).
- [26] Kuhre RE, Wewer Albrechtsen NJ, Hartmann B, Deacon CF, Holst JJ. Measurement of the incretin hormones: glucagon-like peptide-1 and glucose-dependent insulinotropic peptide (in eng) *J Diabetes Complications Apr* 2015;29(3):445–50. <https://doi.org/10.1016/j.jdiacomp.2014.12.006>.
- [27] Orskov C, Rabenhøj L, Wettergren A, Kofod H, Holst JJ. Tissue and plasma concentrations of amidated and glycine-extended glucagon-like peptide I in humans. *Diabetes Apr* 1994;43(4):535–9 (in eng).
- [28] Deacon CF, Holst JJ. Immunoassays for the incretin hormones GIP and GLP-1. *Best Pract Res Clin Endocrinol Metab* 2009;23(4):425–32. <https://doi.org/10.1016/j.beem.2009.03.006>. 2009/08/01/.
- [29] Bak MJ, et al. Specificity and sensitivity of commercially available assays for glucagon-like peptide-1 (GLP-1): implications for GLP-1 measurements in clinical studies. *Diabetes Obes Metab Nov* 2014;16(11):1155–64. <https://doi.org/10.1111/dom.12352> (in eng).
- [30] Toft-Nielsen MB, Madsbad S, Holst JJ. Determinants of the effectiveness of glucagon-like peptide-1 in type 2 diabetes. *J Clin Endocrinol Metab Aug* 2001;86(8):3853–60. <https://doi.org/10.1210/jcem.86.8.7743> (in eng).
- [31] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535. <https://doi.org/10.1136/bmj.b2535>.
- [32] Higgins JP, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ Oct* 18 2011;343:d5928. <https://doi.org/10.1136/bmj.d5928>.
- [33] Higgins JPT, et al., editors. *Cochrane handbook for systematic reviews of interventions version 6.3 (updated February 2022); 2022. www.training.cochrane.org/handbook*.
- [34] M. Borenstein, L. Hedges, and H. Rothstein, "Meta-analysis: Fixed effect vs. random effects."
- [35] Röver C, Knapp G, Friede T. Hartung-Knapp-Sidik-Jonkman approach and its modification for random-effects meta-analysis with few studies (in eng) *BMC Med Res Methodol* Nov 14 2015;15:99. <https://doi.org/10.1186/s12874-015-0091-1>.
- [36] Jackson D, Turner R. Power analysis for random-effects meta-analysis. *Res Synth Methods Sep* 2017;8(3):290–302. <https://doi.org/10.1002/jrsm.1240> (in eng).
- [37] Cook JA, et al. DELTA2 guidance on choosing the target difference and undertaking and reporting the sample size calculation for a randomised controlled trial. *Trials* 2018;19(1):1–8. <https://doi.org/10.1186/s13063-018-2884-0>.
- [38] Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc Jan* 2009;41(1):3–13. <https://doi.org/10.1249/MSS.0b013e31818cb278> (in eng).
- [39] Plourde C, et al. Biliopancreatic diversion with duodenal switch improves insulin sensitivity and secretion through caloric restriction. *Obesity (Silver Spring) Aug* 2014;22(8):1838–46. <https://doi.org/10.1002/oby.20771> (in eng).
- [40] Alsalam W, et al. Effect of single-dose DPP-4 inhibitor sitagliptin on β -cell function and incretin hormone secretion after meal ingestion in healthy volunteers and drug-naïve, well-controlled type 2 diabetes subjects. *Diabetes Obes. Metab. Apr* 2018;20(4):1080–5. <https://doi.org/10.1111/dom.13192> (in eng).
- [41] Nauck MA, et al. Quantification of the contribution of GLP-1 to mediating insulinotropic effects of DPP-4 inhibition with vilaglitin in healthy subjects and patients with type 2 diabetes using exendin [9-39] as a GLP-1 receptor antagonist. *Diabetes Aug* 2016;65(8):2440–7. <https://doi.org/10.2337/db16-0107> (in eng).
- [42] Kozawa J, et al. Similar incretin secretion in obese and non-obese Japanese subjects with type 2 diabetes. *Biochem Biophys Res Commun Mar* 2010;393(3):410–3. <https://doi.org/10.1016/j.bbrc.2010.01.134> (in eng).
- [43] Muscelli E, et al. Separate impact of obesity and glucose tolerance on the incretin effect in normal subjects and type 2 diabetic patients. *Diabetes May* 2008;57(5):1340–8. <https://doi.org/10.2337/db07-1315> (in eng).
- [44] Bagger JI, Knop FK, Lund A, Vestergaard H, Holst JJ, Vilsbøll T. Impaired regulation of the incretin effect in patients with type 2 diabetes. *J Clin Endocrinol Metab Mar* 2011;96(3):737–45. <https://doi.org/10.1210/jc.2010-2435> (in eng).
- [45] Alsalam W, et al. Mixed meal ingestion diminishes glucose excursion in comparison with glucose ingestion via several adaptive mechanisms in people with and without type 2 diabetes. *Diabetes Obes Metab Jan* 2016;18(1):24–33. <https://doi.org/10.1111/dom.12570> (in eng).
- [46] Jørgensen NB, et al. Acute and long-term effects of Roux-en-Y gastric bypass on glucose metabolism in subjects with type 2 diabetes and normal glucose tolerance. *Am J Physiol Endocrinol Metab Jul* 2012;303(1):E122–31. <https://doi.org/10.1152/ajpendo.00073.2012> (in eng).
- [47] Knop FK, et al. Reduced incretin effect in type 2 diabetes: cause or consequence of the diabetic state? *Diabetes Aug* 2007;56(8):1951–9. <https://doi.org/10.2337/db07-0100> (in eng).
- [48] Vilsbøll T, Krarup T, Deacon CF, Madsbad S, Holst JJ. Reduced postprandial concentrations of intact biologically active glucagon-like peptide 1 in type 2 diabetic patients. *Diabetes Mar* 2001;50(3):609–13. <https://doi.org/10.2337/diabetes.50.3.609> (in eng).
- [49] Sonne DP, et al. Postprandial gut hormone responses and glucose metabolism in cholecystectomized patients. *Am J Physiol Gastrointest Liver Physiol* 2013;304(4):G413–9. <https://doi.org/10.1152/ajpgi.00435.2012> (in eng).
- [50] Theodorakis MJ, et al. Human duodenal enteroendocrine cells: source of both incretin peptides, GLP-1 and GIP. *Am J Physiol Endocrinol Metab Mar* 2006;290(3):E550–9. <https://doi.org/10.1152/ajpendo.00326.2004> (in eng).
- [51] Højberg PV, et al. Four weeks of near-normalization of blood glucose has no effect on postprandial GLP-1 and GIP secretion, but augments pancreatic B-cell responsiveness to a meal in patients with type 2 diabetes. *Diabet Med Nov* 2008;25(11):1268–75. <https://doi.org/10.1111/j.1464-5491.2008.02579.x> (in eng).
- [52] Yabe D, et al. Early phase glucagon and insulin secretory abnormalities, but not incretin secretion, are similarly responsible for hyperglycemia after ingestion of nutrients. *J Diabetes Complications Apr* 2015;29(3):413–21. <https://doi.org/10.1016/j.jdiacomp.2014.12.010> (in eng).
- [53] Ryskjaer J, et al. Plasma dipeptidyl peptidase-IV activity in patients with type-2 diabetes mellitus correlates positively with HbA1c levels, but is not acutely affected by food intake. *Eur J Endocrinol Sep* 2006;155(3):485–93. <https://doi.org/10.1530/eje.1.02221> (in eng).
- [54] Astiarraga B, et al. Effects of acute NEFA manipulation on incretin-induced insulin secretion in participants with and without type 2 diabetes. *Diabetologia Aug* 2018; 61(8):1829–37. <https://doi.org/10.1007/s00125-018-4633-z> (in eng).
- [55] Martinussen C, et al. Immediate enhancement of first-phase insulin secretion and unchanged glucose effectiveness in patients with type 2 diabetes after Roux-en-Y gastric bypass. *Am J Physiol Endocrinol Metab* 2015;308(6):E535–44. <https://doi.org/10.1152/ajpendo.00506.2014> (in eng).
- [56] Tricò D, et al. Mechanisms through which a small protein and lipid preload improves glucose tolerance. *Diabetologia Nov* 2015;58(11):2503–12. <https://doi.org/10.1007/s00125-015-3710-9> (in eng).
- [57] Alsama M, et al. Preserved GLP-1 and exaggerated GIP secretion in type 2 diabetes and relationships with triglycerides and ALT. *Eur J Endocrinol Oct* 2013;169(4):421–30. <https://doi.org/10.1530/eje-13-0487> (in eng).
- [58] Lund MT, et al. Preoperative β -cell function in patients with type 2 diabetes is important for the outcome of Roux-en-Y gastric bypass surgery. *J Physiol* 2015;593(14):3123–33. <https://doi.org/10.1113/jp270264> (in eng).
- [59] Ruetten H, et al. Mixed meal and intravenous L-arginine tests both stimulate incretin release across glucose tolerance in man: lack of correlation with β cell function. *Metab Syndr Relat Disord Oct* 2018;16(8):406–15. <https://doi.org/10.1089/met.2018.0022> (in eng).
- [60] Knop FK, Vilsbøll T, Madsbad S, Holst JJ, Krarup T. Inappropriate suppression of glucagon during OGTT but not during isoglycaemic i.v. glucose infusion contributes to the reduced incretin effect in type 2 diabetes mellitus. *Diabetologia Apr* 2007;50(4):797–805. <https://doi.org/10.1007/s00125-006-0566-z> (in eng).
- [61] Rohde U, et al. The impact of EndoBarrier gastrointestinal liner in obese patients with normal glucose tolerance and in patients with type 2 diabetes. *Diabetes Obes Metab Feb* 2017;19(2):189–99. <https://doi.org/10.1111/dom.12800> (in eng).
- [62] Vaag AA, Holst JJ, Vølund A, Beck-Nielsen HB. Gut incretin hormones in identical twins discordant for non-insulin-dependent diabetes mellitus (NIDDM)—evidence for decreased glucagon-like peptide 1 secretion during oral glucose ingestion in NIDDM twins. *Eur J Endocrinol Oct* 1996;135(4):425–32. <https://doi.org/10.1530/eje.0.1350425> (in eng).
- [63] Åhrén B, Holst JJ, Efendic S. Antidiabetogenic action of cholecystokinin-8 in type 2 diabetes. *J Clin Endocrinol Metab Mar* 2000;85(3):1043–8. <https://doi.org/10.1210/jcem.85.3.6431> (in eng).
- [64] Kuwata H, et al. Meal sequence and glucose excursion, gastric emptying and incretin secretion in type 2 diabetes: a randomised, controlled crossover, exploratory trial. *Diabetologia Mar* 2016;59(3):453–61. <https://doi.org/10.1007/s00125-015-3841-z> (in eng).
- [65] St-Jean M, Boudreau F, Carpentier AC, Hivert M-F. HNF1 α defect influences post-prandial lipid regulation. *PLoS One* 2017;12(5). <https://doi.org/10.1371/journal.pone.0177110>. e0177110-e0177110.

- [66] Yan W, et al. Effects of meal size on the release of GLP-1 and PYY after Roux-en-Y gastric bypass surgery in obese subjects with or without type 2 diabetes. *Obes Surg* 2014;24(11):1969–74. <https://doi.org/10.1007/s11695-014-1316-9> (in eng).
- [67] Purnell JQ, et al. Prospective evaluation of insulin and incretin dynamics in obese adults with and without diabetes for 2 years after Roux-en-Y gastric bypass. *Diabetologia* May 2018;61(5):1142–54. <https://doi.org/10.1007/s00125-018-4553-y> (in eng).
- [68] Greenfield JR, et al. Oral glutamine increases circulating glucagon-like peptide 1, glucagon, and insulin concentrations in lean, obese, and type 2 diabetic subjects. *Am J Clin Nutr* Jan 2009;89(1):106–13. <https://doi.org/10.3945/ajcn.2008.26362> (in eng).
- [69] Perakakis N, et al. Circulating levels of five proglucagon-derived peptides in response to intravenous or oral glucose or lipids and to a mixed-meal in subjects with normal weight, overweight, and obesity. *Clin Nutr* 2022;41(9):1969–76. <https://doi.org/10.1016/j.clnu.2022.07.001>. 09.
- [70] Jensen CZ, et al. Responses of gut and pancreatic hormones, bile acids, and fibroblast growth factor-21 differ to glucose, protein, and fat ingestion after gastric bypass surgery. *Am J Physiol Gastrointest Liver Physiol* 2020;318(4):G661–72. <https://doi.org/10.1152/ajpgi.00265.2019>.
- [71] Xie C, et al. Plasma GLP-1 response to oral and intraduodenal nutrients in health and type 2 diabetes-impact on gastric emptying. *J Clin Endocrinol Metab* 2022;107(4), 03(24):e1643–52. <https://doi.org/10.1210/clinem/dgab828> (in eng).
- [72] Albrechtsen NJWewer, et al. A sandwich ELISA for measurement of the primary glucagon-like peptide-1 metabolite (in eng) *Am J Physiol Endocrinol Metab* 2017; 313(3):E284–91. <https://doi.org/10.1152/ajpendo.00005.2017>. 09 01.
- [73] Perakakis N, et al. Methods paper: performance characteristics of novel assays for circulating levels of proglucagon-derived peptides and validation in a placebo controlled cross-over randomized clinical trial. *Metabolism* 2022;129:155157. <https://doi.org/10.1016/j.metabol.2022.155157>. 04.
- [74] Verdich C, Lysgård Madsen J, Toubro S, Buemann B, Holst JJ, Astrup A. Effect of obesity and major weight reduction on gastric emptying. *Int J Obes (Lond)* 2000;24(7):899–905. <https://doi.org/10.1038/sj.ijo.0801250>. 2000/07/01.
- [75] Narang BJ, Atkinson G, Gonzalez JT, Betts JA. A tool to explore discrete-time data: the time series response analyser. *Int J Sport Nutr Exerc Metab* 2020;30(5):374–81. <https://doi.org/10.1123/ijsem.2020-0150> (in English).
- [76] Bahne E, et al. Metformin-induced glucagon-like peptide-1 secretion contributes to the actions of metformin in type 2 diabetes. *JCI Insight* 2018;3(23):e93936. <https://doi.org/10.1172/jci.insight.93936> (in eng).
- [77] Mannucci E, et al. Effects of metformin on glucagon-like peptide-1 levels in obese patients with and without type 2 diabetes. *Diabetes Nutr Metab* Dec 2004;17(6): 336–42 (in eng).