



LJMU Research Online

Kellis, E, Sahinis, C and Baltzopoulos, V

Is hamstrings-to-quadriceps torque ratio useful for predicting anterior cruciate ligament and hamstring injuries? A systematic and critical review

<http://researchonline.ljmu.ac.uk/id/eprint/19006/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Kellis, E, Sahinis, C and Baltzopoulos, V (2022) Is hamstrings-to-quadriceps torque ratio useful for predicting anterior cruciate ligament and hamstring injuries? A systematic and critical review. Journal of Sport and Health Science. 12 (3). pp. 343-358. ISSN 2095-2546

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Review

Is hamstrings-to-quadriceps torque ratio useful for predicting anterior cruciate ligament and hamstring injuries? A systematic and critical review

Eleftherios Kellis^{a,*}, Chrysostomos Sahinis^a, Vasilios Baltzopoulos^b

^a Laboratory of Neuromechanics, Department of Physical Education and Sport Sciences at Serres, Aristotle University of Thessaloniki, Serres 62100, Greece

^b Research Institute for Sport and Exercise Sciences (RISES), Faculty of Science, Liverpool John Moores University, Liverpool L3 5UX, UK

Received 2 August 2021; revised 11 November 2021; accepted 8 December 2021

Available online 19 January 2022

2095-2546/© 2023 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Background: For the past 30 years, the hamstring (H)-to-quadriceps (Q) (H:Q) torque ratio has been considered an important index of muscle strength imbalance around the knee joint. The purpose of this systematic review was to examine the value of H:Q torque ratio as an independent risk factor for hamstring and anterior cruciate ligament (ACL) injuries.

Methods: Database searches were performed to identify all relevant articles in PubMed, MEDLINE, Cochrane Library, and Scopus. Prospective studies evaluating the conventional (concentric H:Q), functional (eccentric H: concentric Q), and mixed (eccentric H at 30°/s: concentric Q at 240°/s) H:Q ratios as risk factors for occurrence of hamstring muscle strain or ACL injury were considered. Risk of bias was assessed using the Quality In Prognosis Studies tool.

Results: Eighteen included studies reported 585 hamstrings injuries in 2945 participants, and 5 studies documented 128 ACL injuries in 2772 participants. Best evidence synthesis analysis indicated that there is very limited evidence that H:Q strength ratio is an independent risk factor for hamstring and ACL injury, and this was not different between various ratio types. Methodological limitations and limited evidence for ACL injuries and some ratio types might have influenced these results.

Conclusion: The H:Q ratio has limited value for the prediction of ACL and hamstring injuries. Monitoring strength imbalances along with other modifiable factors during the entire competitive season may provide a better understanding of the association between H:Q ratio and injury.

Keywords: ACL; Isokinetic; Prospective; Strains; Strength imbalance

1. Introduction

Muscle strength evaluation is an integral part of the fitness assessment of athletes or rehabilitation of individuals with various pathologies or injuries. One of the key parameters in these applications is the hamstrings (H)-to-quadriceps (Q) (H:Q) torque ratio as an index of a strength imbalance between antagonistic muscle groups around the knee joint.^{1,2} Identification of relative weakness in one muscle relative to the other can serve 2 main purposes: first, to identify athletes with strength imbalance in the pre-season period and then use this information to design appropriate training programs to restore H:Q ratio to “normal” levels³ and improve their performance;^{4,5} second,

to identify imbalances after injury or other pathological conditions and use this information to design appropriate rehabilitation programs.^{6–8}

The H:Q ratio can be calculated using various combinations of contraction types (concentric, eccentric, and isometric), angular velocities, and type of torque (peak or angle-specific torque).⁹ The conventional ratio is calculated using torque values of the same contraction type for both antagonistic muscle groups. These include the concentric H:concentric Q (Q_{CON}); the eccentric H (H_{ECC}):eccentric Q; and the H:Q torque ratio.⁹ It has been proposed that an isometric H:Q ratio of approximately 0.66 represents the average value in a typical population sample.¹⁰ This is partly confirmed by a recent systematic review of evidence showing that the isokinetic conventional ratio, when tested at slow (12°/s) to intermediate (180°/s) angular velocities, is in the range of 0.52–0.67 and that it increases further at faster angular velocities.⁸

Peer review under responsibility of Shanghai University of Sport.

* Corresponding author.

E-mail address: ekellis@phed-sr.auth.gr (E. Kellis).

<https://doi.org/10.1016/j.jshs.2022.01.002>

Cite this article: Kellis E, Sahinis C, Baltzopoulos V. Is hamstrings-to-quadriceps torque ratio useful for predicting anterior cruciate ligament and hamstring injuries? A systematic and critical review. *J Sport Health Sci* 2023;12:343–58.

When the quadriceps produce force by shortening (concentric contraction), the hamstrings are activated as antagonists while lengthening (eccentric contraction).⁹ For this reason, several authors have proposed that a more functional approach should be taken in the calculation of H:Q ratio^{1,11,12} by considering the typical muscle actions during a knee extension. Therefore, the dynamic control¹² or functional¹¹ torque ratio has been established as the ratio of eccentric hamstrings torque over concentric quadriceps torque (H_{ECC}/Q_{CON}). Normative values for the functional ratio also vary depending on test angular velocity, being around 0.79 at 60°/s and exceeding 1 at angular velocities greater than 240°/s.⁸ Another modification of the functional ratio is the mixed ratio, which is calculated by dividing the H_{ECC} torque at an angular velocity of 30°/s by the Q_{CON} torque at 240°/s.¹³ Hence, by taking the H_{ECC} torque at a slow speed and the Q_{CON} at a faster speed, the resultant H:Q ratio for the typical sport population would be centered around a value of 1.¹³ According to the authors who proposed the mixed ratio,^{13,14} the main advantage of this ratio is its greater experimental validity (i.e., it is less affected by inertial artifacts). This is because the recorded torque at fast angular velocity tests shows better experimental validity in isokinetic fast angular velocity concentric contraction tests (Q_{CON} torque at 240°/s) than in eccentric contraction tests, while eccentric torque values at a slow angular velocity (H_{ECC} torque at 30°/s) have better validity than fast velocity tests. A recent review has examined the associations between different types of H:Q ratios,¹⁵ but they did not systematically review the association of H:Q ratio with injury.

One important application of the H:Q ratio has been to monitor strength imbalances in individuals with anterior cruciate ligament (ACL) deficiency^{12,16} or reconstruction.^{17,18} This is based on the observation that the hamstrings co-contract near full extension and generate an opposing force to the anterior component of the quadriceps force.^{19,20} This reduces anterior shear loading and contributes to a physiological range of motion and safer relative position of the femur and tibia bones in the knee joint, thus maintaining knee joint stability and reducing ACL load.^{19,20} Consequently, it has been proposed that a low force generating capacity of the hamstrings relative to the quadriceps may be related to ACL injury and re-injury, especially in female athletes.²¹ To our knowledge, published evidence on the association between H:Q ratio and ACL injury has not been reviewed.

An altered H:Q ratio is also frequently considered a risk factor for hamstring strains.²² This is often attributed to a reduced relative ability of the hamstrings to control combined hip flexion and knee extension during dynamic movements such as sprinting or stretching.^{23,24} Systematic reviews have shown that there is strong evidence that strength variables, including H:Q ratio, are not independent risk factors for hamstring injury.^{7,25} However, these studies have examined H:Q ratio generically together with various other potential risk factors and, hence, specific issues that may influence H:Q ratio measurement and its relation to injury have not been considered. These include torque ratio measurement issues (e.g., reliability, gravity correction, validity), calculation method and the type of ratio (e.g., conventional, dynamic, mixed), and interpretation (e.g., the determination of cut-off values). These

considerations are very important because previous studies have identified methodological limitations and issues with the experimental protocol that influence the calculation of H:Q ratio.^{1,2} These limitations are often underestimated, which may lead to erroneous interpretation of the obtained H:Q ratio values. While studies have advocated the use of one type of ratio over another for different applications, there is still confusion about which type of H:Q ratio is better for predicting or preventing injury.⁸ The use of various types of H:Q ratios in combination with limitations in injury survey data collection and analysis may also influence conclusions regarding the usefulness of H:Q ratio for injury prevention. Three recent reviews^{6–8} have shown that it is unclear whether H:Q ratio is sensitive enough to detect differences between injured and non-injured legs or between injured and non-injured individuals suffering from hamstring strains; these studies, however, did not examine the association between H:Q ratio and future injury.

The purpose of this systematic review was to examine the ability of different types of H:Q strength ratios to prospectively predict hamstrings and ACL injuries in athletes. Two specific questions were addressed in this review: (a) Is the H:Q ratio a significant risk factor for ACL and hamstring strains and, if so, which type of H:Q ratio shows a greater association with injury? and (b) do methodological issues and limitations influence the association between H:Q ratio and ACL or hamstring injuries?

2. Methods

2.1. Search strategy and selection of studies

This review was not registered. A comprehensive, multi-step search strategy using Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines was performed in June 2021 using the (Population, Intervention, Comparison, Outcomes of interest; PICO) framework for prognostic studies to identify all relevant articles in 4 electronic databases:²⁶ PubMed, MEDLINE, Cochrane Library, and Scopus. Population: sport athletes; problem: hamstrings or ACL injury; prognostic factor: ipsilateral strength ratios; comparison: prospective comparison of injured and uninjured groups; and outcome: hamstrings or ACL injury and associated risk estimates. The electronic databases were searched independently by 2 authors (EK and CS), to reduce the probability of study selection bias, with no limit for year of publication. The following combination of terms with the proper Boolean (“AND”, “OR”, and “NOT”) operators were used: “isokinetic strength” or “strength ratio” or “strength balance” or “hamstrings to quadriceps ratio” or “knee flexors to knee extensors ratio” or “hamstrings injury” or “posterior thigh injury” or “hamstring strain” or “hamstring tear” or “anterior cruciate ligament injury” or “anterior cruciate ligament rupture” or “risk factors” or “predictor” or “prediction” or “prevention”. Additionally, secondary searches were conducted by screening the reference list of the selected studies and by evaluating the reference list of previous related reviews. After duplicates were removed, 2 independent reviewers (EK and CS) assessed the potential studies by comparing the lists of included and excluded studies. Any discrepancies between the 2 authors

(EK and CS) with regard to the included and excluded studies were resolved through discussion and consensus.

2.2. Eligibility criteria

Studies that satisfied the following criteria were included in the review: (a) publications in international English-language peer-reviewed journals, (b) studies that examined healthy individuals of any age and playing level (ex-athletes or athletic populations) with subsequent occurrence of hamstring muscle or ACL injury during either a sport and competition-related activity or training period, and (c) studies that employed any type of H:Q strength ratio (conventional, functional, and mixed) as a measurement outcome to investigate the risk of injury. Reviews, case and brief reports, letters to editors, thesis, and conference abstracts were excluded.

2.3. Risk of bias assessment

The methodological quality of the included studies was independently examined by 2 reviewers (CS and EK) using a modified version of the Quality In Prognosis Studies tool described by Hayden et al.²⁷ This risk of bias assessment tool examines 6 potential bias domains (study participation, study attrition, prognostic factor measurement, outcome measurement, study confounding, and statistical analysis and reporting) to evaluate different aspects of the potential risk of bias in prognostic studies.⁷ Each of the 6 study design elements contain 5 – 7 items of potential bias that were evaluated according to specific criteria used to examine the degree of risk with a score of “yes” or “no”. If fewer than 75% of the items within a single category were scored with a “yes”, this domain was determined to have a “High risk” of bias, while “Low risk” of bias was considered if the number of items scored with “yes” was greater than or equal to 75%. To obtain an overall low risk of bias, a study should have a low bias on at least 5 of the 6 domains and the outcome measurement. The domain “outcome measurement” (Domain 4) refers to 3 criteria: first, whether the study provides a clear definition of the outcome measurement; second, whether information about the validity and reliability of the measurements is provided; and, finally, whether the same methodology was applied for all participants.

2.4. Data extraction

Two authors (EK and VB) independently extracted the following data from each study to an Excel spreadsheet (Microsoft Office Excel 2016; Microsoft, Redmond, WA, USA): (a) authors and year of publication, (b) details regarding the characteristics of participants who underwent isokinetic evaluation (such as sample size, sport, and sex), (c) methodological characteristics (test angular velocity, types of ratios, gravity correction, participant position, and reliability measures), (d) study duration, (e) strength ratio data, and (f) key findings related to injury prediction.

2.5. Data synthesis and analysis

Preliminary evaluation of the identified literature indicated relatively few studies examining different versions of the H:Q ratio. Further, the data from various studies originated from different sports, and this might have influenced the heterogeneity of the sample for each specific ratio type. For this reason, the final number of studies for each type of H:Q ratio was considered insufficient for meta-analysis. Therefore, a best evidence synthesis⁷ was used to facilitate the interpretation of data derived from included studies. It consisted of a 5 levels of evidence-based qualitative analysis:

1. Strong evidence: consistent results in 2 or more low risk of bias studies (high quality), with generally consistent findings in $\geq 75\%$ of studies.
2. Moderate evidence: provided by 1 low risk of bias study (high quality) and/or 2 or more studies with high risk of bias (low quality) and by generally consistent results across all studies ($\geq 75\%$ of the studies reported consistent findings).
3. Limited evidence: provided by single-study findings from high risk of bias study (low quality).
4. Conflicting evidence: inconsistent findings in multiple studies ($< 75\%$ of the studies reported consistent findings).
5. No evidence: no studies (randomized controlled trials or non-randomized controlled trials) available for assessment.

Best evidence synthesis of the included studies was applied twice: first, to examine the association of each H:Q ratio type and injury across all sports; and second, to examine the association of H:Q ratio and injury in each sport.

3. Results

3.1. Search results

A total of 10,154 potentially relevant published studies were identified through the electronic search (Fig. 1). Of these, 4032 duplicate articles were removed, while after title and abstract screening, 6083 documents were excluded. Of the remaining 41 articles, 18 additional studies were excluded on the basis of full-text screening. Therefore, 23 articles were retained.

3.2. Description of the included studies

Of the 23 studies included in this review, 18 studies examined hamstring injuries and 5 studies examined ACL injuries. These studies documented 585 hamstrings injuries in 2945 participants across football (soccer),^{3,13,28–36} Australian rules football,^{37–39} sprinting,^{40,41} American football,⁴² and Rugby.⁴³ Additionally, 128 ACL injuries in 2772 participants across football (soccer),^{44,45} basketball,^{44,46,47} handball,⁴⁵ floorball,⁴⁶ and in military academy cadets.⁴⁸

From the included studies, demographics, strength measurement methodology, and main data treatment techniques were tabulated. Further, H:Q torque ratio values (mean and SD) for injured and control groups were visualized according to type of H:Q ratio as well as type of injury.




Fig. 1. Search flow diagram. ACL = anterior cruciate ligament; H:Q = hamstrings-to-quadriceps torque ratio.

3.3. Methodological characteristics of the included studies

The details of prospective studies are presented in Table 1. Some studies on hamstring injury have examined more than 1 type of ratio. Hence, 14 studies^{3,13,28,31,33–39,41–43} measured the conventional H:Q ratio, 6 studies^{3,29,32,37,40,41} used the functional H:Q ratio, and 5 studies^{3,13,33–35} examined the mixed ratio, while there were some studies that calculated a different version of the strength imbalance between the 2 muscle groups.^{30,40} Of the 5 included studies on ACL injury, all of them examined the conventional ratio,^{44,45–48} only 1 examined the functional ratio,⁴⁸ and none of them assessed the mixed ratio (Table 1). All studies except for 2^{32,44} used a slow angular velocity (60°/s), 7 studies^{3,31,32,37,38,41,43} used a 180°/s test, 7 studies^{3,13,33–36,41} used a 240°/s test, 4 studies used a 300°/s test,^{29–31,38} while only one³ used a 120°/s angular velocity test.

Twelve of the included studies^{3,13,28–30,37–40,44,45,47} evaluated strength from a seated position (hip angle range 80°–105°), while the remaining 11 studies^{31–36,41–43,46,48} did not specify the participant testing position (Table 1). Additionally, only 9 studies^{3,13,31,33,34,37,38,40,48} included information

regarding gravity correction procedure, and only 4 studies^{3,41,45,46} reported the reliability of their H:Q ratio measurements. Twenty-one studies calculated H:Q ratio using peak torque values, and only 2 studies^{29,41} used both peak and angle-specific H:Q ratios.

3.4. Risk of bias assessment

The scores on the potential risk of bias domains of the included studies are presented in Fig. 2. Particularly, 11 studies on hamstring injury were scored as low risk, while the remaining 7 studies had a high risk (Fig. 2). Of the 5 studies that examined ACL injury risk, 3 were rated as low risk of bias and 2 were rated as high risk of bias (Fig. 2).

3.5. Main findings

Fig. 3 presents the reported conventional H:Q ratios for groups with and without hamstring and ACL injury, while the corresponding data for the functional, mixed, and other ratio types are presented in Fig. 4. Of the 18 studies, non-significant

Table 1
Methodological details of the identified prospective studies examining the association between the H:Q torque ratio and hamstring strain or ACL injury.

Study	Sample	Hip angle	GC	Type	Rel	Legs tested	Ratio type	Angular velocity (°/s)	Seasons tracked	Number of injuries; data treatment
Hamstring injuries										
Bennell et al. (1998) ³⁷	102 M (I: 12, CG: 90) Australian rules football players	90°	Yes	PT	No	Both	CR, FR	60,180	1–2	12 injuries; comparison between players with and without CR and FR H:Q < 0.60, group comparison
Cameron et al. (2003) ³⁹	20M (I: 6, CG: 14) Pro Australian rules football players	Seated	N/R	PT	No	Both	CR	60	2	6 injuries; ROC curve, comparison between players with and without H:Q < 0.60 and > 0.66, group comparison
Croisier et al. (2008) ¹³	462 (I: 35, CG: 427) Pro soccer players	105°	Yes	PT	No	Both	CR, MX	60,240	1	35 injuries; logistic regression in groups with and without 2 of the following criteria: CR H:Q ratios > 0.55 or 0.57, MX H:Q ratio > 0.98 or 1.05 and 15% bilateral differences in strength
Dauty et al. (2016) ³	136 M (I: 57, CG: 79) Pro soccer players	85°	Yes	PT	Yes	Both	CR, FR, MX	60,180,240	10	57 injuries; likelihood ratio for positive and negative tests for peak H:Q ratios and cutoffs for CR < 0.47 and FR and MX < 0.80, group comparison
Dauty et al. (2018) ³³	194 M (I: 36, CG: 158) Pro soccer players	N/R	Yes	PT	No	Both	CR, MX	60,240	15	36 injuries; odds ratio in groups with and without cutoffs CR ≤ 0.47 or < 0.60 and MX < 0.8 or < 1.0, group comparison
Dauty et al. (2020) ³⁴	91 M (I: 31, CG: 60) Pro soccer players	N/R	Yes	PT	No	Both	CR, MX	60,240		31 injuries; logistic regression, ROC curve, odds ratio in groups with and without cutoffs < 0.80 or < 1.00, group comparison
Fousekis et al (2011) ³²	99 M (I: 16, CG: 83) Pro soccer players	N/R	N/R	PT	No	Both	FR	180	1	16 injuries; logistic regression for players with FR H:Q < 1
Henderson et al. (2010) ³¹	143 M (I: 10, CG: 26) elite soccer players	N/R	Yes	PT	No	Both	CR	60,180,300	1	10 injuries; logistic regression, group comparisons, comparison between players with and without CR H:Q < 0.60
Lee et al. (2018) ³⁵	146 (I: 41, CG: 105) Pro soccer players	N/R	N/R	PT	No	Both	CR, MX	60,240		41 injuries; odds ratio for “low” and “high” H:Q value groups, based on a cut-off value, determined using ROC curve analysis, group and leg comparisons, comparison between players with and without CR H:Q < 0.505
Orchard et al. (1997) ³⁸	37 (I: 6, CG: 31) Australian rules football players	Seated	Yes	PT	No	Both	CR	60,180,300	1	6 injuries; discriminant analysis, comparison between players with and without CR H:Q < 0.61, group comparison
Scippers and Leach (2020) ²⁸	89 M (I: 4, CG: 85) amateur soccer players	Seated	N/R	PT	No	Both	CR	60		4 injuries; logistic regression, group comparison
Shalaj et al. (2020) ³⁶	143 M (I: 40, CG: 103) elite soccer players	N/R	N/R	PT	No	Both	CR	60,240	1	40 injuries; logistic regression, group comparison
Sugiura et al. (2008) ⁴⁰	30 M (I: 6, CG: 24) elite sprinters	80°	Yes	PT	No	Both	FR, OT	60	1 year	6 injuries; group and leg comparisons
van Dyk et al. (2016) ³⁰	563 M (I: 167, CG: 396) elite soccer players	90°	N/R	PT	No	Both	OT	60,300	4	167 injuries; logistic regression, ROC curve, group comparisons
van Dyk et al. (2017) ²⁹	413 M (I: 66, CG: 347) Pro football players	90°	N/R	PT, AST	No	Both	FR	60,300	2	66 injuries; Cox regression, odds ratios for the groups with lowest (<1 SD below the mean) and the highest (>1 SD above the mean) value, group and leg comparisons
Yamada and Mastumoto (2009) ⁴³	21 (I: 6, CG: 15) rugby players	N/R	N/R	PT	No	Both	CR	60,180	6 months	6 injuries; discriminant analysis, ROC curve, group comparison

(continued on next page)

Table 1 (Continued)

Study	Sample	Hip angle	GC	Type	Rel	Legs tested	Ratio type	Angular velocity (°/s)	Seasons tracked	Number of injuries; data treatment
Yeung et al. (2009) ⁴¹	44 M and F (I: 8, CG: 36) sprinters	N/R	N/R	PT, AST	Yes	Both	CR, FR	60, 180, 240	1	8 injuries; Cox regression, group comparison, comparison between players with and without CR H:Q < 0.60
Zvijac et al. (2013) ⁴²	212 (I: 38, CG: 174) Pro American football players	N/R	N/R	PT	No	Both	CR	60	5	38 injuries; group comparisons, sensitivity analysis, comparison between players with and without CR H:Q < 0.60
Anterior cruciate ligament injuries										
Steffen et al. (2016) ⁴⁵	420 F elite handball and 447 F elite football players (I: 57, CG: 810)	90°	N/R	PT	Yes	Both	CR	60	8	57 injuries; odds ratio, logistic regression, group comparison
Shimozaki et al. (2018) ⁴⁷	168 F (I: 12, CG: 156) high school basketball players	90°	N/R	PT	No	Both	CR	60	3	12 injuries; logistic regression, group comparison
Hietamo et al. (2020) ⁴⁶	96 F basketball and 78 F football athletes (I: 14, CG: 160)	N/R	N/R	PT	Yes	Both	CR	60	3	14 injuries; Cox regression analysis, Cox hazard ratios, ROC curve analysis, group comparison
Uhorchak et al. (2003) ⁴⁸	755 M and F (I: 23, CG: 732) military academy cadets	N/R	Yes	PT	No	Both	CR, FR	60	4	23 injuries; logistic regression, risk ratio calculation, group comparison
Myer et al. (2009) ⁴⁴	110 F collegiate soccer and basketball players (I: 22, CG: 88)	90°	N/R	PT	No	Both	CR	300	5	22 injuries; group comparison

Abbreviations: ACL = anterior cruciate ligament; AST = angle-specific torque; CG = control group; CR = conventional ratio; F = female; FR = functional ratio; GC = gravity correction; H:Q = hamstrings-to-quadriceps; I = injured; M = male; MX = mixed ratio; N/R = not reported; OT = other strength ratio; Pro = professional; PT = peak torque; ROC = receiver operating characteristics.

differences between injured and uninjured groups or legs were reported by 14 studies on hamstring injury, while none of the included studies reported differences between ACL injured and non-injured athletes.

The main findings of the included studies are presented in Table 2. Of the 14 studies, 11 reported no association between conventional H:Q ratio measurements at 60°/s and hamstring injury,^{3,13,28,31,33,34,36,37,41–43} while 4 studies^{31,37,38,43} reached a similar conclusion using an H:Q ratio at 180°/s. Similarly, non-significant associations with hamstring injury were reported by 5 (out of 6) studies^{3,29,32,37,41} for the functional ratio and by 4 (out of 5) studies for the mixed ratio.^{13,33–35} Of the studies listed above, only 4 showed a high risk of bias.^{3,32,37,43} In contrast, of the 18 total studies that examined the H:Q ratio following hamstring injury, only 5^{3,35,38–40} reported a significant association, of which, 3 were rated as having a high risk of bias.^{35,38,39} None of the prospective studies reported a significant association between H:Q ratio and ACL injury. Of the 5 studies^{44,45–48} that looked at this potential association, 2 studies^{44,45} had a high risk of bias (Table 2).

Summarizing the results for each individual sport indicates that 11 studies^{3,13,28–36} examined soccer players, of which, 9 reported a non-significant association.^{13,28–34,36} There was also no association reported for American football⁴² and rugby⁴³ players. Out of the 3 studies^{37–39} that examined Australian rules football athletes, 2 reported a significant association between a different type of H:Q ratio and hamstring injury.^{38,39} In addition, 2 studies in sprinters^{40,41} reported significant associations between injury and a specific H:Q ratio type. As for ACL injuries, no association with injury was reported by all included studies.

3.6. Best evidence synthesis

Best evidence synthesis strongly indicated that neither the conventional ratio at 60°/s and 240°/s, the functional ratio at 60°/s, nor other strength ratios were associated with hamstring injury risk (Table 3). Moderate evidence for no association was found for the conventional H:Q ratio at 300°/s and the functional H:Q ratio at 180°/s. There was also limited evidence for no association with injury for the conventional H:Q ratio at 120°/s, the functional H:Q ratio at 60°/s, 120°/s, 240°/s, and 300°/s. Finally, evidence for the conventional H:Q ratio at 180°/s and the mixed ratio was conflicting. For ACL injuries, there is strong (conventional ratio) or limited (conventional ratio at 300°/s and functional ratio) evidence that the H:Q ratio is not associated with injury (Table 3).

Table 4 presents the results of best evidence synthesis analysis for each sport. For Australian footballers, there was limited to moderate evidence that H:Q ratios are not associated with hamstring injury; exception was the conventional H:Q ratio at 60°/s, where the evidence was conflicting. Similar results were also found for soccer players. For sprinters, there was conflicting evidence for one H:Q ratio type, moderate evidence for no association with injury for 2 ratio types, and







Fig. 2. Risk of bias assessment.

moderate evidence for an association between injury and the conventional H:Q ratio at 180°/s. As for ACL injuries, there was limited to strong evidence of no association with H:Q ratio for all sports.

4. Discussion

The results of this review indicate that there is moderate-to-strong evidence that the H:Q strength ratio is not an independent risk factor for either hamstring or ACL injury. This result

applies for both the conventional and functional H:Q ratio, while evidence for the mixed H:Q ratio is conflicting. Several methodological limitations and differences between studies might have influenced these results.

4.1. Association of H:Q ratio with injury

With respect to the first question raised in this study, best-evidence-synthesis analysis has shown that the H:Q ratio is not associated with hamstring and ACL injury (Table 3). Overall, of




Fig. 3. Mean conventional H:Q torque ratio values of Inj and Uinj groups reported by the studies included in this review. For all studies but that of Sugiura et al.,⁴⁰ 95%CI were estimated based on the reported means, SDs, and study sample sizes. Solid circle means injured and circle means uninjured. * $p < 0.05$, significant differences between groups. 95%CI = 95% confidence interval; ACL = anterior cruciate ligament; conc = concentric; ecc = eccentric; H:Q = hamstrings-to-quadriceps; Inj = injured; Uinj = uninjured.

the 15 studies with low risk of bias, only 3^{3,13,41} reported that the H:Q ratio is a significant risk factor for hamstring strain, while all studies found that there is no association of H:Q ratio and ACL injury (Table 3). The mean group values reported by various

studies show some similarity. In particular, the conventional ratio values at 60°/s are centered around 0.60 (Fig. 2), the functional ratios at 60°/s around 0.80, and the mixed ratios around 1.30 (Fig. 3). Nevertheless, the variability in each measurement in both




Fig. 4. Mean (95%CI) functional, mixed, and other type of H:Q torque ratio values of groups with (Inj) and without (Uinj) hamstring injury reported by the studies included in this review. For all studies but that of Sugiura et al.,⁴⁰ 95%CI were estimated based on the reported means, SDs, and study sample sizes. Solid circle means injured and circle means uninjured. **p* < 0.05, significant differences between groups. 95%CI = 95% confidence interval; H:Q = hamstrings-to-quadriceps; Inj = injured; L = left; R = right; Uinj = uninjured.

injured and non-injured groups prevents users from deciding which value should be considered as typical and which ones might be indicative of injury risk. Therefore, it is not certain that an individual who has an H:Q ratio that is almost 2 SDs less than the average group value is at increased risk for injury. On the other

hand, it is also not certain that an individual who displays an H:Q ratio that is similar to the average group value has a lower risk for sustaining an injury. This substantial overlap between players with high and low risk of injury is seen in almost all existing screening tests.⁴⁹

Table 2
Results of the included studies on the association between the H:Q ratio and hamstring or ACL injury.

Study	Type of ratio			
	Conventional	Functional	Mixed	Other
Hamstring injury				
<i>60°/s</i>				
Bennell et al. (1998) ³⁷	NS for H:Q < 0.60	NS for H:Q < 0.60		
Cameron et al. (2003) ³⁹	S for H:Q < 0.60, SP = 50% S for H:Q > 0.66 SP = 100%			
Croisier et al. (2008) ¹³	Players with H:Q > 0.55–0.57 and/or 15% bilateral strength differences had 4.66 greater relative risk for injury			
Croisier et al. (2008) ¹³	NS		NS	
Dauty et al. (2016) ³	NS for H:Q < 0.47	NS for H:Q < 0.80	S for H:Q < 0.80, SE = 2.5%, SP 99%	
Dauty et al. (2018) ³³	NS for H:Q < 0.47 or < 0.60		NS	
Dauty et al. (2020) ³⁴			NS for H:Q < 0.80 or < 1.00	
Lee et al. (2018) ³⁵	S for H:Q < 0.505, OR = 3.14 (95% CI:1.37–7.22)		NS	NS
Orchard et al. (1997) ³⁸	S for H:Q < 0.61, positive predictive value = 77.4%			
Sugiura et al. (2008) ⁴⁰	N/R	S		N/R
Yeung et al. (2009) ⁴¹	NS	NS		
Zvijac et al. (2013) ⁴²	NS for H:Q < 0.60			
van Dyk et al. (2016) ³⁰				NS
van Dyk et al. (2017) ²⁹		NS		
Others ^{28,31,36,43}	NS			
<i>120°/s</i>				
Dauty et al. (2016) ³	NS for H:Q < 0.47	NS for H:Q < 0.80		
<i>180°/s</i>				
Bennell et al. (1998) ³⁷	NS for H:Q < 0.60	NS for H:Q < 0.60		
Dauty et al. (2016) ³	S for H:Q < 0.47, SE = 7% SP = 97%			
Yeung et al. (2009) ⁴¹	S for H:Q < 0.60, (Hazard ratio = 17.4, 95%CI: 1.3–231.4)	NS		
Fousekis et al. (2011) ³²		NS for H:Q < 1.00		
Others ^{31,38,43}	NS			
<i>240°/s</i>				
Lee et al. (2018) ³⁵	S for H:Q < 0.505, OR = 0.93 (95%CI: 0.89–0.98)			
Dauty et al. (2016) ³	NS H:Q < 0.47			
Yeung et al. (2009) ⁴¹	NS	NS		
Others ^{31,36}	NS			
<i>300°/s</i>				
van Dyk et al. (2016) ³⁰				NS
van Dyk et al. (2017) ²⁹		NS		
Others ^{31,38}	NS			
ACL injury				
<i>60°/s</i>				
Uhorchak et al. (2003) ⁴⁸	NS	NS		
Others ^{45–47}	NS			
<i>300°/s</i>				
Myer et al. (2009) ⁴⁴	NS			

Abbreviations: 95%CI = 95% confidence interval; ACL = anterior cruciate ligament; H:Q = hamstring-to-quadriceps; NS = non-significant; N/R = not reported; OR = odds ratio; S = significant; SE = sensitivity; SP = specificity.

This study also explored whether different types of H:Q ratio have greater association with injury. However, current evidence indicates that the functional ratio does not offer a greater injury predictive capacity than the conventional one, while evidence on the value of the mixed ratio is conflicting (Table 3). It should be mentioned, however, that of the total 15 studies with lower risk bias, only 5 examined the functional ratio and 2 examined the mixed ratio (Table 1). This indicates that there is much more evidence available for the conventional ratio than for the other ratio types. Furthermore, no studies have examined the association of the mixed H:Q ratio and ACL injury (Table 1). Collectively, these results indicate that the limitations of the conventional ratio as an injury screening tool are not overcome if

a more “functional” approach in the H:Q ratio calculation is followed.

There are various factors that may have influenced these results. In particular, studies used different H:Q cutoff values to discriminate the sample (Table 2). The cutoff values varied from 0.47 to 0.66 for the conventional ratio at 60°/s, from 0.78 to 1.05 for the functional H:Q ratio (across speeds), and from 0.80 to 1 for the mixed H:Q ratio. This could be attributed mainly to differences in the method used to determine these cutoff values as well as to the size of the sample examined in each study. Some studies tested the idea that the optimum value of the conventional H:Q should be 0.60;^{37,42} others determined cutoff values by calculating the area under the

Table 3
Results of best evidence synthesis of studies for each H:Q ratio.

	Low risk of bias	High risk of bias	Association with injury (↑/=)	Association with injury (Best synthesis result)	Level of evidence
Hamstring injury study					
<i>Conventional ratio</i>					
60°/s	Dauty et al. ³ Dauty et al., ³⁴ Croisier et al. ¹³ Zvjajac et al. ⁴² Shalaj et al. ³⁶ Henderson et al. ³¹ Schipper and Leach ²⁸ Yeung et al. ⁴¹	Dauty et al. ³³ Orchard et al. ³⁸ Cameron et al. ³⁹ Yamada and Mastumoto ⁴³ Bennell et al. ³⁷ Lee et al. ³⁵	↑ Orchard et al., ³⁸ Cameron et al., ³⁹ Lee et al. ³⁵ = Dauty et al. ³³ Yamada and Mastumoto, ⁴³ Bennell et al., ³⁷ Dauty et al., ³ Dauty et al., ³⁴ Croisier et al., ¹³ Zvjajac et al., ⁴² Shalaj et al., ³⁶ Henderson et al., ³¹ Schipper and Leach, ²⁸ Yeung et al. ⁴¹	No	Strong
120°/s 180°/s	Dauty et al. ³ Dauty et al. ³ Henderson et al. ³¹ Yeung et al. ⁴¹	Orchard et al. ³⁸ Yamada and Mastumoto ⁴³ Bennell et al. ³⁷	= Dauty et al. ³ ↑ Dauty et al., ³ Yeung et al. ⁴¹ = Henderson et al., ³¹ Orchard et al. ⁸ Yamada and Mastumoto, ⁴³ Bennell et al., ³⁷	No Unknown	Moderate Conflicting
240°/s	Dauty et al. ³ Croisier et al. ¹³ Shalaj et al. ³⁶ Yeung et al. ⁴¹	Lee et al. ³⁵	↑ Lee et al. ³⁵ = Dauty et al., ³ Croisier et al., ¹³ Shalaj et al., ³⁶ Yeung et al. ⁴¹	No	Strong
300°/s	Henderson et al. ³¹	Orchard et al. ³⁸	= Orchard et al. ³⁸ Henderson et al. ³¹	No	Moderate
<i>Functional ratio</i>					
60°/s	Dauty et al. ³ Yeung et al. ⁴¹ Sugiura et al. ⁴⁰	Bennell et al. ³⁷	↑ Sugiura et al. ⁴⁰ = Dauty et al., ³ Yeung et al., ⁴¹ Bennell et al. ³⁷	No	Strong
60°/s–300°/s 120°/s 180°/s	van Dyk et al. ²⁹ Dauty et al. ³ Yeung et al. ⁴¹	Bennell et al. ³⁷ Fousekis et al. ³²	= van Dyk et al. ²⁹ = Dauty et al. ³ = Bennell et al. ³⁷ Fousekis et al., ³² Yeung et al., ⁴¹ = Yeung et al. ⁴¹	No No No	Moderate Moderate Moderate
240°/s	Yeung et al. ⁴¹			No	Moderate
<i>Mixed ratio</i>					
	Dauty et al. ³ Dauty et al., ³⁴ Croisier et al. ¹³	Dauty et al. ³³ Lee et al. ³⁵	↑ Dauty et al. ³ = Dauty et al., ³⁴ Croisier et al., ¹³ Dauty et al., ³³ Lee et al., ³⁵	No	Strong
<i>Other</i>	Sugiura et al., ⁴⁰ van Dyk et al. ³⁰		= Sugiura et al., ⁴⁰ van Dyk et al. ³⁰	No	Strong
ACL injury study					
<i>Conventional ratio</i>					
60°/s	Shimozaki et al. ⁴⁷ Hietamo et al., ⁴⁶ Uhorchak et al. ⁴⁸	Steffen et al. ⁴⁵	= Shimozaki et al., ⁴⁷ Hietamo et al., ⁴⁶ Uhorchak et al., ⁴⁸ Steffen et al., ⁴⁵	No	Strong
300°/s		Myer et al. ⁴⁴	= Myer et al. ⁴⁴	No	Limited
<i>Functional ratio</i>					
60°/s	Uhorchak et al. ⁴⁸		= Uhorchak et al. ⁴⁸	No	Moderate

Notes: ↑ means significant association with future injury; = means no significant association with future injury.

Abbreviations: ACL = anterior cruciate ligament; H:Q = hamstrings-to-quadriceps.

Table 4
Results of best evidence synthesis analysis of studies that examined the association of injury and H:Q ratio for each sport.

	Low risk of bias	High risk of bias	Association with injury (↑/=)	Association with injury (Best synthesis result)	Level of evidence
Hamstring injury study					
<i>Australian rules football</i>					
CR 60°/s		Orchard et al. ³⁸ Cameron et al. ³⁹ Bennell et al. ³⁷	↑ Orchard et al., ³⁸ Cameron et al. ³⁹ = Bennell et al. ³⁷	Unknown	Conflicting
CR 180°/s		Orchard et al. ³⁸ Bennell et al. ³⁷	= Orchard et al., ³⁸ Bennell et al. ³⁷	No	Moderate
CR 300°/s		Orchard et al. ³⁸	= Orchard et al. ³⁸	No	Limited
FR 60°/s, 180°/s		Bennell et al. ³⁷	= Bennell et al. ³⁷	No	Limited
<i>Soccer/football</i>					
CR 60°/s	Dauty et al. ³ Dauty et al., ³⁴ Croisier et al. ¹³ Shalaj et al. ³⁶ Henderson et al. ³¹ Schippers and Leach ²⁸	Dauty et al. ³³ Lee et al. ³⁵	↑ Lee et al. ³⁵ = Croisier et al. ¹³ Dauty et al., ³³ Dauty et al., ³ Dauty et al., ³⁴ Shalaj et al., ³⁶ Henderson et al., ³¹ Schippers and Leach, ²⁸	No	Strong
CR 120°/s	Dauty et al. ³		= Dauty et al. ³	No	Moderate
CR 180°/s	Dauty et al. ³ Henderson et al. ³¹		↑ Dauty et al. ³ = Henderson et al. ³¹	Unknown	Conflicting
CR 240°/s	Dauty et al. ³ Croisier et al. ¹³ Shalaj et al. ³⁶	Lee et al. ³⁵	= Dauty et al. ³ Croisier et al., ¹³ Shalaj et al., ³⁶ Lee et al. ³⁵	No	Strong
CR 300°/s	Henderson et al. ³¹	Orchard et al. ³⁸	= Orchard et al. ³⁸ Henderson et al. ³¹	No	Moderate
FR 60°/s, 120°/s	Dauty et al. ³		= Dauty et al. ³	No	Moderate
FR 60°/s–300°/s	van Dyk et al. ²⁹		= van Dyk et al. ²⁹	No	Moderate
FR 180°/s		Fousekis et al. ³²	= Fousekis et al. ³²	No	Limited
MR	Dauty et al. ³ Dauty et al., ³⁴ Croisier et al. ¹³	Dauty et al. ³³ Lee et al. ³⁵	↑ Dauty et al. ³ = Dauty et al. ³⁴ Croisier et al., ¹³ Dauty et al., ³³ Lee et al. ³⁵	No	Strong
Other	van Dyk et al. ³⁰		= van Dyk et al. ³⁰	No	Moderate
<i>Sprinting</i>					
CR 60°/s, 240°/s	Yeung et al. ⁴¹		= Yeung et al. ⁴¹	No	Moderate
CR 180°/s	Yeung et al. ⁴¹		↑ Yeung et al. ⁴¹	Yes	Moderate
FR 60°/s	Yeung et al. ⁴¹ Sugiura et al. ⁴⁰		↑ Sugiura et al. ⁴⁰ = Yeung et al. ⁴¹	Unknown	Conflicting
FR 180°/s, 240°/s	Yeung et al. ⁴¹		= Yeung et al. ⁴¹	No	Moderate
<i>American football</i>					
CR 60°/s	Zvjic et al. ⁴²		= Zvjic et al. ⁴²	No	Moderate
<i>Rugby</i>					
CR 60°/s		Yamada and Mastumoto ⁴³	= Yamada and Mastumoto ⁴³	No	Limited
ACL injury study					
<i>Soccer/football</i>					
CR 60°/s		Steffen et al. ⁴⁵	= Steffen et al. ⁴⁵	No	Limited
CR 300°/s		Myer et al. ⁴⁴	= Myer et al. ⁴⁴	No	Limited
<i>Basketball</i>					
CR 60°/s	Shimozaki et al. ⁴⁷ Hietamo et al. ⁴⁶		= Shimozaki et al., ⁴⁷ Hietamo et al. ⁴⁶	No	Strong
CR 300°/s		Myer et al. ⁴⁴	= Myer et al. ⁴⁴	No	Limited
<i>Floorball</i>					
CR 60°/s	Hietamo et al. ⁴⁶		= Hietamo et al. ⁴⁶	No	Moderate
<i>Handball</i>					
CR 60°/s		Steffen et al. ⁴⁵	= Steffen et al. ⁴⁵	No	Limited
<i>Military cadets</i>					
CR and FR 60°/s	Uhorchak et al. ⁴⁸		= Uhorchak et al. ⁴⁸	No	Moderate

Notes: ↑ means significant association with future injury; = means no significant association with future injury.

Abbreviations: ACL = anterior cruciate ligament; CR = conventional ratio; FR = functional ratio; H:Q = hamstring-to-quadriceps; MR = mixed ratio.

curve of receiver operating characteristics based on the raw data^{29,30,34,35,39,41,43} (Table 1); still others tested the cutoff values reported by previous experiments.^{3,32,33} Perhaps the most influential study is that done by Croisier et al.,¹³ which examined an almost 4 times greater number of players than all other studies.¹³ These authors did not find a significant association between the conventional H:Q ratio at a specific angular velocity and injury (Table 2). They observed, however, that injured athletes showed a much greater deficit in H_{ECC} torque at 30°/s than that observed for Q_{CON} torque at 240°/s. Hence, they were able to show that soccer players had a 4.66 times greater risk of sustain a hamstring injury if they met 2 of the following criteria: a conventional ratio less than 0.55, a mixed ratio greater than 0.98, and/or 15% bilateral differences in strength when compared with players showing no imbalance in the preseason. Based on the suggestions made by these authors, the H:Q ratio should be combined with bilateral differences in strength to detect players who are at greater injury risk. The use of these specific cutoff values, however, was based on previous experiments in a very small number of players,¹⁴ and hence, it has been criticized.^{50,51} In addition, the cutoff values proposed by Croisier et al.¹³ were tested by other studies,^{3,32,33} which reported a non-significant association between H:Q ratio and hamstring injury. In the most recent of those prospective studies, Dauty et al.³³ failed to find a significant prediction of hamstring injury for a wide range of the conventional and mixed H:Q ratio cutoff values.

Another factor that contributed to the present findings is that the conditions at which the H:Q ratio is measured have limited association with muscle function at the instant of injury. Hamstring injuries are frequently attributed to different mechanisms, one occurring during high-speed sprinting and the other occurring during excessive stretching.^{23,24} ACL injuries occur during the deceleration phase of dynamic movements, such as landing from jumping or change of direction, while the knee is near full extension, frequently in combination with knee valgus and internal/external tibial rotation.⁵² In these movements, the hamstrings work at relatively long muscle lengths. Evidence indicates that isokinetic strength has moderate correlation with kicking⁵³ or sprinting,⁵⁴ but this relationship is lower between H:Q ratio and performance,⁵⁵ obviously, because H:Q ratio is an expression of the relative maximum strength when each muscle works as agonist and not the torque that is exerted simultaneously by the 2 antagonist muscles.⁵⁶ Related to this is the criticism that the position and speed of testing do not resemble actual conditions, such as sprinting or change in direction.⁷ In the included studies, isokinetic testing angular velocities ranged from 60°/s to 300°/s (Table 1), which are lower than those developed during fast sport movements, such as fast kicking (~550°/s–1720°/s,⁵⁷ sprinting (~500°/s–800°/s),⁵⁸ or landing (~78°/s–570°/s).^{59,60} This may also explain the observation that angular velocity does not influence the association of H:Q ratio with injury (Table 2). For this reason, evaluation of the conventional or functional H:Q at a slow angular velocity, such as 60°/s, is recommended.

An important contributor to the low predictive capacity of the H:Q ratio is the multifactorial origin of both hamstring²²

and ACL^{61,62} injuries. Although various factors are often considered to increase the risk for sustaining an injury, very few of them have been identified as independent risk factors for hamstring²² or ACL^{61,62} injury. Hence, preseason evaluation of strength imbalances between antagonistic muscle groups may not identify all players who are at a greater risk for sustaining an injury unless it is part of a wider screening program with a broader range of risk factors examined. Future studies that use multifactorial designs that incorporate measurements of various modifiable factors may assist in this direction.⁶³

Another observation is that all included studies have calculated H:Q ratio using peak torque during the test (Table 1). The H:Q ratio changes throughout the range of motion due to differences in torque–joint angular position curves between the quadriceps and the hamstrings. Hamstrings torque is generally greater near full extension, reaching peak values at 40° of knee flexion.⁵⁶ In contrast, peak quadriceps torque is smaller near full extension, reaching its peak at 60° of knee flexion.⁵⁶ Hence, starting from 90° of knee flexion, the H:Q torque ratio is very low, and it increases almost linearly, reaching its peak at approximately 30°–20° of knee flexion with values around 1.^{56,64} Taking the peak torque provides an index of the maximum strength generation capacity of each muscle, but it may not reflect the instantaneous relationship between hamstrings and quadriceps torque that occurs at a specific joint position. In theory, the use of angle-specific H:Q torque is ecologically more valid as it is closer to the injury mechanism of the ACL^{64,65} as well as the hamstrings muscles.^{66,67} Nevertheless, only 2 low-risk studies^{29,41} examined angle-specific strength values, and they reported that the resulting H:Q ratio has no association with hamstrings injury.

4.2. Methodological issues

The second aim of this study was to identify methodological aspects of the included studies that may have had an influence on the results (Table 1). First, it was observed that only 3 low-risk studies^{3,41,46} examined the reliability of measurements, which appeared to be moderate or high (intraclass correlation coefficients: 0.80–0.98 for peak torque values). Studies have shown a moderate reliability of H:Q ratios in repeated sessions.^{68–70} Impellizzeri et al.⁷⁰ noticed lower H:Q values for the first session as compared with the second and third sessions, suggesting a learning effect. Neuromuscular inhibition, which often takes place during eccentric testing, has also been considered as an important threat to the reliability and validity of the functional or mixed H:Q ratio.³³ In fact, these studies have shown that percentage changes in H:Q should be greater than 18%⁶⁸ or even 28%⁶⁹ to detect real changes. None of the included studies have identified such large differences in any type of ratio (Figs. 3–4) between injured and non-injured athletes. This represents an important limitation of most studies, and it should be taken into consideration when interpreting the results of this review. In addition, threats to the validity of the H:Q ratio, such as gravitational torque¹ and inertial force,⁷¹ are often not considered. Using the peak torque value during the non-isokinetic phase of the torque–angle curve, for example, may lead to an error of up to

16% at an angular velocity of 300°/s.⁷¹ This may influence the association of H:Q ratio with injury in a specific cohort, and it also provides erroneous conclusions regarding the extent of muscle imbalances in these particular athletes.

Differences in injury recording methodology between studies might also have had an influence on the results of this review. First, several studies examined a relatively small sample of players reporting a low number of injuries. For example, 5 out of 15 studies with low risk of bias reported less than 16 injuries each (Table 1). In this case, comparisons between injured and uninjured players is based on unequal sample sizes, and this can dramatically influence the statistical power of the results.⁷² More important, with a small sample size, the validity of using specific cutoff values to discriminate the sample is questionable. Second, concerns have been raised regarding the definition, type, and severity of injury in various studies.¹³ For example, hamstring muscle strains can be mild, moderate, or severe,⁷³ or they may be sprint-type or stretch-type.^{23,24} Studies included in this systematic review did not identify the mechanism that led to the registered injury. One may hypothesize that sprint-type and stretch-type injuries exhibit differences in muscle function and that, hence, a single preseason measurement of H:Q ratio is inadequate for predicting both injury types. Third, inclusion of athletes with a history of injury may also influence the results as previous injury history may increase injury risk.³³ All but 2 studies^{31,44} included in this review excluded individuals with a previous injury incident from their sample. Finally, the reviewed literature provided evidence for the association of H:Q ratio with injury in a specific sample. Based on Bahr,⁴⁹ this is only the first step for an effective screening strategy. The next step is validating the cutoff values using a new sample and the final step is to test whether a prevention program can reduce risk of injury in those athletes who are classified as high risk based on their screening results.

An additional factor related to the heterogeneity of the samples' characteristics is that the participants were athletes from various sports, each with different performance demands (Table 1). Re-examination of our data collected on ACL injuries indicates it is unlikely that the type of sport influenced the results, as all studies failed to show an association between ACL injury and H:Q ratio (Table 4). As for the hamstring injuries, studies in American football, rugby, and soccer athletes indicated no association with H:Q ratio (Table 4). Interestingly, for Australian rules football athletes^{37–39} and sprinters,^{38,39} there was evidence that supports a significant association, but this originates from studies with high risk of bias.^{38,39} For this reason, there was moderate evidence for a significant association with injury in only 1 case (Table 4). Overall, based on the collected evidence, it appears that the absence of association between H:Q ratio and hamstring injury is common in all sports. Nevertheless, subject to more evidence, the possibility that a different association exists for sports such as sprinting or Australian rules football cannot be excluded.

4.3. Implications

There are important implications to be gleaned from the findings of this review. First, due to the low association

between H:Q ratio and injury, one may suggest that preseason evaluation of muscle strength imbalance is not useful for identifying players with a greater risk potential. This does not indicate that preseason evaluation of strength should not be performed. Previous studies have shown that strength training may prevent hamstring injuries and, therefore, monitoring of strength is necessary.⁷⁴ There have been suggestions that other tests such as Nordics, hand-held dynamometers, or force plates may substitute or complement isokinetic testing for better injury prediction results.^{7,75} However, it should be stressed that these tests have no⁷⁶ or only moderate correlation with performance,⁷⁷ and there is no evidence that they are associated with injury.²⁹ Isokinetic dynamometers offer maximum resistance throughout the movement at both concentric and eccentric conditions, and they can be used to monitor strength progress of players effectively, safely, and correctly. Obviously, any test that is performed under controlled laboratory conditions is not expected to have a direct association with muscle function during movements that lead to injury. This observation aligns with the latest research by Bahr,⁴⁹ who commented that “any existing screening test is unlikely to be able to predict injury with sufficient accuracy”.

Strength levels are greater in the preseason period and tend to decrease over the in-season period.⁷⁸ It is possible that some players display greater strength losses than others and that this may influence their performance and injury risk. Further, while preseason evaluation levels provide an evaluation of strength imbalances, it is unclear how in-season training influences strength imbalances. For example, it has been shown that preseason strength was reduced after a 12-week in-season soccer training period without any strength training.⁷⁸ Performance of one strength training session per week during the in-season period maintained strength,⁷⁹ while strength training twice a week may decrease strength due to overtraining.⁸⁰ This indicates that injury epidemiology studies should monitor strength sessions adopted by players during the in-season period as they may affect strength imbalances relative to the preseason measurements. In addition, monitoring strength imbalances at selected time instances during both the preseason and in-season periods will allow for a closer examination of the association between injury and strength imbalance by using strength values obtained during the specific period when the injury occurs.

Limitations of this review are acknowledged. First, we restricted eligibility to studies in English only while we searched 2 of the 4 databases. Second, we did not contact study authors to request clarifications where needed; instead, we extracted information from the published data and cross-checked our results with those reported by previous meta-analysis studies.

5. Conclusion

Preseason evaluation of H:Q ratio has limited value for the prediction of hamstring and ACL injuries. The type of H:Q ratio does not influence the association with injury. Monitoring strength imbalances along with other modifiable factors during the entire competitive season may provide a better understanding of the association between H:Q ratio and injury.

Authors' contributions

EK conceived the idea for the manuscript and wrote the first versions of the manuscript; CS and VB provided intellectual contribution as part of the redrafting process within their areas of expertise. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

Data availability

<https://data.mendeley.com/datasets/vpp87gykvh/draft? a=9f05cca6-878c-4aa0-9808-7cbcb1171246>

References

- Kellis E, Baltzopoulos V. Isokinetic eccentric exercise. *Sports Med* 1995;**19**:202–22.
- Baltzopoulos V, Brodie DA. Isokinetic dynamometry: Applications and limitations. *Sports Med* 1989;**8**:111–6.
- Dauty M, Menu P, Fouasson-Chailloux A, Ferréol S, Dubois C. Prediction of hamstring injury in professional soccer players by isokinetic measurements. *MLTJ* 2016;**6**:116–23.
- O'Donnell SR, Eitan DN, Roper JL. A comparison of quadriceps-to-hamstrings ratios during isokinetic testing, cutting, and drop landings in male soccer players. *Int J Exerc Sci* 2020;**13**:157–66.
- Delextrat A, Gregory J, Cohen D. The use of the functional H:Q ratio to assess fatigue in soccer. *Int J Sport Med* 2010;**31**:192–7.
- Coombs R, Garbutt G. Developments in the use of the hamstring/quadriceps ratio for the assessment of muscle balance. *J Sport Sci Med* 2002;**1**:56–62.
- Green B, Bourne MN, Pizzari T. Isokinetic strength assessment offers limited predictive validity for detecting risk of future hamstring strain in sport: A systematic review and meta-analysis. *Br J Sports Med* 2018;**52**:329–36.
- Baroni BM, Ruas CV, Ribeiro-Alvares JB, Pinto RS. Hamstring-to-quadriceps torque ratios of professional male soccer players: A systematic review. *J Strength Cond Res* 2020;**34**:281–93.
- Kellis E. Quantification of quadriceps and hamstring antagonist activity. *Sports Med* 1998;**25**:37–62.
- Steindler A. *Kinesiology of the human body under normal and pathological conditions*. Springfield, IL: Charles C. Thomas, Publisher; 1955.
- Aagaard P, Simonsen EB, Magnusson SP, Larsson B, Dyhre-Poulsen P. A new concept for isokinetic hamstring: Quadriceps muscle strength ratio. *Am J Sports Med* 1998;**26**:231–7.
- Dvir Z, Eger G, Halperin N, Shklar A. Thigh muscle activity and anterior cruciate ligament insufficiency. *Clin Biomech* 1989;**4**:87–91.
- Croisier JLL, Ganteaume S, Binet J, Genty M, Ferret JMM. Strength imbalances and prevention of hamstring injury in professional soccer players: A prospective study. *Am J Sports Med* 2008;**36**:1469–75.
- Croisier JL, Forthomme B, Namurois MH, Vanderthommen M, Crielaard JM. Hamstring muscle strain recurrence and strength performance disorders. *Am J Sports Med* 2002;**30**:199–203.
- Ruas CV, Pinto RS, Haff GG, Lima CD, Pinto MD, Brown LE. Alternative methods of determining hamstrings-to-quadriceps ratios: A comprehensive review. *Sports Med Open* 2019;**5**:11. doi:10.1186/s40798-019-0185-0.
- Kannus P, Latvala K, Jarvinen M. High muscle strengths in the anterior cruciate ligament deficient knee. Isokinetic and isometric long-term results. *J Orthop Sports Phys Ther* 1987;**9**:223–7.
- Fischer F, Fink C, Herbst E, et al. Higher hamstring-to-quadriceps isokinetic strength ratio during the first post-operative months in patients with quadriceps tendon compared to hamstring tendon graft following ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2018;**26**:418–25.
- Kannus P. Isokinetic evaluation of muscular performance: Implications for muscle testing and rehabilitation. *Int J Sport Med* 1994;**15**(Suppl. 1): S11–8.
- Baratta R, Solomonow M, Zhou BH, Letson D, Chuinard R, D'Ambrosia R. Muscular coactivation. The role of the antagonist musculature in maintaining knee stability. *Am J Sports Med* 1988;**16**:113–22.
- Kellis E, Baltzopoulos V. The effects of the antagonist muscle force on intersegmental loading during isokinetic efforts of the knee extensors. *J Biomech* 1999;**32**:19–25.
- Hewett TE, Myer GD, Zazulak BT. Hamstrings to quadriceps peak torque ratios diverge between sexes with increasing isokinetic angular velocity. *J Sci Med Sport* 2008;**11**:452–9.
- Opar DA, Williams MD, Shield AJ. Hamstring strain injuries: Factors that lead to injury and re-injury. *Sports Med* 2012;**42**:209–26.
- Askling CM, Tengvar M, Saartok T, Thorstensson A. Acute first-time hamstring strains during high-speed running: A longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med* 2007;**35**:197–206.
- Askling CM, Tengvar M, Saartok T, Thorstensson A. Acute first-time hamstring strains during slow-speed stretching: Clinical, magnetic resonance imaging, and recovery characteristics. *Am J Sports Med* 2007;**35**:1716–24.
- Freckleton G, Pizzari T. Risk factors for hamstring muscle strain injury in sport: A systematic review and meta-analysis. *Br J Sports Med* 2013;**47**:351–8.
- Santos CMD, Pimenta CADM, Nobre MRC. The PICO strategy for the research question construction and evidence search. *Rev Lat Am Enfermagem* 2007;**15**:508–11.
- Hayden JA, van Tulder MW, Tomlinson G. Systematic review: Strategies for using exercise therapy to improve outcomes in chronic low back pain. *Ann Intern Med* 2005;**142**:776–85.
- Schippers RC, Leach LL. Relationship between selected physiological characteristics and hamstring injuries in amateur male soccer players. *Afri J Phys Act Health Sci* 2020;**26**. doi:10.37597/ajphes.2020.26.2.5.
- van Dyk N, Witvrouw E, Bahr R. Interseason variability in isokinetic strength and poor correlation with Nordic hamstring eccentric strength in football players. *Scand J Med Sci Sports* 2018;**28**:1878–87.
- van Dyk N, Bahr R, Whiteley R, et al. Hamstring and quadriceps isokinetic strength deficits are weak risk factors for hamstring strain injuries. *Am J Sports Med* 2016;**44**:1789–95.
- Henderson G, Barnes CA, Portas MD. Factors associated with increased propensity for hamstring injury in English Premier League soccer players. *J Sci Med Sport* 2010;**13**:397–402.
- Fousekis K, Tsepis E, Poulmedis P, Athanasopoulos S, Vagenas G. Intrinsic risk factors of non-contact quadriceps and hamstring strains in soccer: A prospective study of 100 professional players. *Br J Sports Med* 2011;**45**:709–14.
- Dauty M, Menu P, Fouasson-Chailloux A. Cutoffs of isokinetic strength ratio and hamstring strain prediction in professional soccer players. *Scand J Med Sci Sports* 2018;**28**:276–81.
- Dauty M, Menu P, Fouasson-Chailloux A. Hamstring muscle injury prediction by isokinetic ratios depends on the method used. *Clin J Sport Med* 2020;**30**:40–5.
- Lee JWY, Mok KMM, Chan HCK, Yung PSH, Chan KMM. Eccentric hamstring strength deficit and poor hamstring-to-quadriceps ratio are risk factors for hamstring strain injury in football: A prospective study of 146 professional players. *J Sci Med Sport* 2018;**21**:789–93.
- Shalaj I, Gjaka M, Bachl N, Wessner B, Tschann H, Tishukaj F. Potential prognostic factors for hamstring muscle injury in elite male soccer players: A prospective study. *PLoS One* 2020;**15**:e0241127. doi:10.1371/journal.pone.0241127.
- Bennell K, Wajswelner H, Lew P, et al. Isokinetic strength testing does not predict hamstring injury in Australian Rules footballers. *Br J Sports Med* 1998;**32**:309–14.
- Orchard J, Marsden J, Lord S, Garlick D. Preseason hamstring muscle weakness associated with hamstring muscle injury in Australian footballers. *Am J Sports Med* 1997;**25**:81–5.

39. Cameron M, Adams R, Maher C. Motor control and strength as predictors of hamstring injury in elite players of Australian football. *Phys Ther Sport* 2003;**4**:159–66.
40. Sugiura Y, Saito T, Sakuraba K, Sakuma K, Suzuki E. Strength deficits identified with concentric action of the hip extensors and eccentric action of the hamstrings predispose to hamstring injury in elite sprinters. *J Orthop Sports Phys Ther* 2008;**38**:457–64.
41. Yeung SS, Suen AMY, Yeung EW. A prospective cohort study of hamstring injuries in competitive sprinters: Preseason muscle imbalance as a possible risk factor. *Br J Sports Med* 2009;**43**:589–94.
42. Zvijac JE, Toriscelli TA, Merrick S, Kiebzak GM. Isokinetic concentric quadriceps and hamstring strength variables from the NFL scouting combine are not predictive of hamstring injury in first-year professional football players. *Am J Sports Med* 2013;**41**:1511–8.
43. Yamada M, Mastumoto D. The reaction time of mental rotation predicts strain in rugby players. *J Phys Ther Sci* 2009;**21**:177–81.
44. Myer GD, Ford KR, Barber Foss KD, Liu C, Nick TG, Hewett TE. The relationship of hamstrings and quadriceps strength to anterior cruciate ligament injury in female athletes. *Clin J Sport Med* 2009;**19**:3–8.
45. Steffen K, Nilstad A, Kristianslund EK, Myklebust G, Bahr R, Krosshaug T. Association between lower extremity muscle strength and noncontact ACL injuries. *Med Sci Sport Exerc* 2016;**48**:2082–9.
46. Hietamo J, Parkkari J, Leppänen M, et al. Association between lower extremity muscular strength and acute knee injuries in young team-sport athletes. *Transl Sports Med* 2020;**3**:626–37.
47. Shimozaki K, Nakase J, Takata Y, Shima Y, Kitaoka K, Tsuchiya H. Greater body mass index and hip abduction muscle strength predict noncontact anterior cruciate ligament injury in female Japanese high school basketball players. *Knee Surg Sports Traumatol Arthrosc* 2018;**26**:3004–11.
48. Uhorchak JM, Scoville CR, Williams GN, Arciero RA, St Pierre P, Taylor DC. Risk factors associated with noncontact injury of the anterior cruciate ligament: A prospective four-year evaluation of 859 West Point cadets. *Am J Sports Med* 2003;**31**:831–42.
49. Bahr R. Why screening tests to predict injury do not work-and probably never will. . . : A critical review. *Br J Sports Med* 2016;**50**:776–80.
50. Grygorowicz M, Michałowska M, Walczak T, et al. Discussion about different cut-off values of conventional hamstring-to-quadriceps ratio used in hamstring injury prediction among professional male football players. *PLoS One* 2017;**12**:e0188974. doi:10.1371/journal.pone.0188974.
51. Houweling TAW, Head A, Hamzeh M. Validity of isokinetic testing for previous hamstring injury detection in soccer players. *Isokinet Exerc Sci* 2009;**17**:213–20.
52. Shimokochi Y, Shultz SJ. Mechanisms of noncontact anterior cruciate ligament injury. *J Athl Train* 2008;**43**:396–408.
53. Kellis E, Katis A. The relationship between isokinetic knee extension and flexion strength with soccer kick kinematics: An electromyographic evaluation. *J Sports Med Phys Fitness* 2007;**47**:385–94.
54. Dowson MN, Nevill ME, Lakomy HKA, Nevill AM, Hazeldine RJ. Modelling the relationship between isokinetic muscle strength and sprint running performance. *J Sports Sci* 1998;**16**:257–65.
55. Struzik A, Pietraszewski B. Relationships between hamstrings-to-quadriceps ratio and variables describing countermovement and drop jumps. *Appl Bionics Biomech* 2019;**2019**: 4505481. doi:10.1155/2019/4505481.
56. Kellis E, Katis A. Quantification of functional knee flexor to extensor moment ratio using isokinetics and electromyography. *J Athl Train* 2007;**42**:477–85.
57. Kellis E, Katis A. Biomechanical characteristics and determinants of instep soccer kick. *J Sport Sci Med* 2007;**6**:154–65.
58. Kellis E, Liassou C. The effect of selective muscle fatigue on sagittal lower limb kinematics and muscle activity during level running. *J Orthop Sports Phys Ther* 2009;**39**:210–20.
59. Wang L-I. The lower extremity biomechanics of single- and double-leg stop-jump tasks. *J Sport Sci Med* 2011;**10**:151–6.
60. Yin L, Sun D, Mei Q, Gu Y, Baker J, Feng N. The kinematics and kinetics analysis of the lower extremity in the landing phase of a stop-jump task. *Open Biomed Eng J* 2015;**9**:103–7.
61. Alentorn-Geli E, Myer GD, Silvers HJ, et al. Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc* 2009;**17**:705–29.
62. Shultz SJ, Schmitz RJ, Benjaminse A, Collins M, Ford K, Kulas AS. ACL research retreat VII: An update on anterior cruciate ligament injury risk factor identification, screening, and prevention. *J Athl Train* 2015;**50**:1076–93.
63. Bittencourt NFN, Meeuwisse WH, Mendonça LD, Nettel-Aguirre A, Ocarino JM, Fonseca ST. Complex systems approach for sports injuries: Moving from risk factor identification to injury pattern recognition - Narrative review and new concept. *Br J Sports Med* 2016;**5**:1309–14.
64. Baumgart C, Welling W, Hoppe MW, Freiwald J, Gokeler A. Angle-specific analysis of isokinetic quadriceps and hamstring torques and ratios in patients after ACL-reconstruction. *BMC Sports Sci Med Rehabil* 2018;**10**:23. doi:10.1186/s13102-018-0112-6.
65. Bates NA, Hewett TE. Motion analysis and the anterior cruciate ligament: Classification of injury risk. *J Knee Surg* 2016;**29**:117–25.
66. Liu H, Garrett WE, Moorman CT, Yu B. Injury rate, mechanism, and risk factors of hamstring strain injuries in sports: A review of the literature. *J Sport Health Sci* 2012;**1**:92–101.
67. Ruas C V, Pinto MD, Brown LE, Minozzo F, Mil-Homens P, Pinto RS. The association between conventional and dynamic control knee strength ratios in elite soccer players. *Isokinet Exerc Sci* 2015;**23**:1–12.
68. Ayala F, De Ste Croix M, Sainz De Baranda P, Santonja F. Absolute reliability of hamstring to quadriceps strength imbalance ratios calculated using peak torque, joint angle-specific torque and joint ROM-specific torque values. *Int J Sports Med* 2012;**33**:909–16.
69. Sole G, Hamrén J, Milosavljevic S, Nicholson H, Sullivan SJ. Test-retest reliability of isokinetic knee extension and flexion. *Arch Phys Med Rehabil* 2007;**88**:626–31.
70. Impellizzeri FM, Bizzini M, Rampinini E, Cereda F, Maffiuletti NA. Reliability of isokinetic strength imbalance ratios measured using the Cybex NORM dynamometer. *Clin Physiol Funct Imaging* 2008;**28**:113–9.
71. Iossifidou AN, Baltzopoulos V. Inertial effects on the assessment of performance in isokinetic dynamometry. *Int J Sport Med* 1998;**19**:567–73.
72. Rusticus SA, Lovato CY. Impact of sample size and variability on the power and type I error rates of equivalence tests: A simulation study. *Pract Assess Res Eval* 2014;**19**:11. doi:10.7275/4s9m-4e81.
73. Mueller-Wohlfahrt HW, Haensel L, Mithoefer K, et al. Terminology and classification of muscle injuries in sport: The Munich consensus statement. *Br J Sports Med* 2013;**47**:342–50.
74. Hibbert O, Cheong K, Grant A, Beers A, Moizumi T. A systematic review of the effectiveness of eccentric strength training in the prevention of hamstring muscle strains in otherwise healthy individuals. *N Am J Sports Phys Ther* 2008;**3**:67–81.
75. McCall A, Carling C, Davison M, et al. Injury risk factors, screening tests and preventative strategies: A systematic review of the evidence that underpins the perceptions and practices of 44 football (soccer) teams from various premier leagues. *Br J Sports Med* 2015;**49**:583–9.
76. Suarez-Arrones L, Lara-Lopez P, Rodriguez-Sanchez P, et al. Dissociation between changes in sprinting performance and Nordic hamstring strength in professional male football players. *PLoS One* 2019;**14**: e0213375. doi:10.1371/journal.pone.0213375.
77. Prince C, Morin JB, Mendiguchia J, et al. Sprint specificity of isolated hamstring-strengthening exercises in terms of muscle activity and force production. *Front Sports Act Living* 2021;**2**:609636. doi:10.3389/fspor.2020.609636.
78. DeRenne C, Hetzler RK, Buxton BP, Ho KW. Effects of training frequency on strength maintenance in pubescent baseball players. *J Strength Cond Res* 1996;**10**:8–14.
79. Rønnestad BR, Nymark BS, Raastad T. Effects of inseason strength maintenance training frequency in professional soccer players. *J Strength Cond Res* 2011;**25**:2653–60.
80. Kraemer WJ, French DN, Paxton NJ, et al. Changes in exercise performance and hormonal concentrations over a big ten soccer season in starters and nonstarters. *J Strength Cond Res* 2004;**18**:121–8.