

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Phytoplankton communities exhibit a stronger response to environmental changes than bacterioplankton in three subtropical reservoirs

Journal:	<i>Environmental Science & Technology</i>
Manuscript ID:	es-2015-02637j.R1
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Liu, Lemian; Institute of Urban Environment, Chinese Academy of Sciences, Yang, Jun; Institute of Urban Environment, Chinese Academy of Sciences, Lv, Hong; Institute of Urban Environment, Chinese Academy of Sciences, Yu, Xiaoqing; Institute of Urban Environment, Chinese Academy of Sciences, Wilkinson, David; Liverpool John Moores University, School of Natural Science and Psychology Yang, Jun; Institute of Urban Environment, Chinese Academy of Sciences,

SCHOLARONE™
Manuscripts

1 **Phytoplankton communities exhibit a stronger response to**
2 **environmental changes than bacterioplankton in three**
3 **subtropical reservoirs**

4
5 Lemian Liu,^{†, a} Jun Yang,^{†, ‡, a} Hong Lv,^{†, a} Xiaoqing Yu,[†] David M. Wilkinson[§] and
6 Jun Yang^{*, †}

7
8 [†]Aquatic EcoHealth Group, Key Laboratory of Urban Environment and Health, Institute of Urban
9 Environment, Chinese Academy of Sciences, Xiamen 361021, People's Republic of China

10 [‡]University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

11 [§]School of Natural Science and Psychology, Liverpool John Moores University, Byrom Street,
12 Liverpool L3 3AF, UK

13

14

15 Key words: algae, bacteria, spatiotemporal pattern, time-lag analysis, community ecology

16

17 ^aThese authors contributed equally to this work.

18 ^{*}For correspondence. E-mail: jyang@iue.ac.cn; Tel. (+86)5926190775; Fax (+86)5926190775.

ABSTRACT

The simultaneous analysis of multiple components of ecosystems is crucial for comprehensive studies of environmental changes in aquatic ecosystems - but such studies are rare. In this study, we analyzed simultaneously the bacterioplankton and phytoplankton communities in three Chinese subtropical reservoirs, and compared the response of these two components to seasonal environmental changes. Time-lag analysis indicated that the temporal community dynamics of both bacterioplankton and phytoplankton showed significant directional changes, and variance partitioning suggested that the major reason was the gradual improvement of reservoir water quality from middle eutrophic to oligo-mesotrophic levels during the course of our study. In addition, we found a higher level of temporal stability or stochasticity in the bacterioplankton community than in the phytoplankton community. Potential explanations are that traits associated with bacteria such as high abundances, widespread dispersal, potential for rapid growth rates and rapid evolutionary adaptation may underlie the different stability or stochasticity of bacterioplankton and phytoplankton communities to the environmental changes. In addition, the indirect response to nitrogen and phosphorus of bacterioplankton may result in the fact that environmental deterministic selection was stronger for the phytoplankton than for the bacterioplankton communities.

INTRODUCTION

Bacterioplankton and phytoplankton are critical components of aquatic microbial food webs, and play essential roles in the structure and function of aquatic ecosystems.^{1,2} Understanding the processes and mechanisms that underlie the abundance and/or biovolume variations of bacterioplankton and phytoplankton communities, are major goals in both pure and applied microbial community ecology. Many previous studies of bacterioplankton and phytoplankton variation have focused on traditional biogeographical concepts, such as distance decay relationships,^{3, 4} species-area relationships,^{5,6} and the niche vs. neutral debate.⁷⁻⁹ Data on the temporal variation in aquatic microbial community composition is more mixed and limited. There is a long history of monitoring phytoplankton in lakes in some parts of the world – for example, such studies have a history of over 100 years in the English Lake District, with extensive data sets existing from the mid-twentieth century onwards.¹⁰ However, bacteria (excluding cyanobacteria) were much more challenging to study until the rise of molecular methods in the late 1980's, so even in well studied regions such as the English Lake District there is much less known about the patterns in bacterial abundance, compared with the data on phytoplankton.¹¹ Macan and Worthington (p83) summarized the mid-twentieth century position in writing that 'The bacteria... play an important, though as yet little understood role in the economy of all fresh waters.'¹² Even today the global distribution of such studies is very patchy with the vast majority of molecular studies of freshwater bacterial communities being confined to Europe, and to a lesser extent North America. The rest of the world is covered by a very small number of studies¹³ with the tropics and sub-tropics being particularly poorly sampled - but see Dalu *et al.* for a rare African example.¹⁴ Arguably, this lack of understanding of the temporal behavior of both bacterioplankton and phytoplankton communities hinders the development of theories about how the stability of microbial community structure and function is maintained across time.¹⁴

In aquatic ecosystems, it has become increasingly clear that the temporal variation in composition of bacterioplankton and phytoplankton communities primarily depends on environmental changes through space and time.¹⁴⁻²⁰ However, the response of these microbial communities to such changes are also mediated by their properties, including their history, metabolic flexibility, physiological tolerance, dispersal capacity, and taxonomic and functional diversity.^{3, 21, 22} Therefore, we hypothesize that bacterioplankton and phytoplankton will have different sensitivities to environmental changes. To date, most aquatic microbial diversity studies have focused on just bacterioplankton or phytoplankton; comprehensive studies simultaneously analyzing bacterioplankton and phytoplankton components of microbial communities across time are much rarer – especially in regions other than Europe.^{13, 23} Analyzing simultaneously the dynamics of different microbial groups is likely to be crucial for understanding the dynamics and responses of the ecosystems to environmental changes.

Additionally, to date, studies estimating temporal variation of microbial communities have mainly used multivariate statistics to illustrate the lack of variation

81 in rates and patterns of community change.²⁴ Few studies have used metrics to
82 specifically quantify temporal variation of microbial communities.^{25, 26} Time-lag
83 analysis (TLA) has proved a useful diagnostic tool to quantify the temporal variation
84 of ecological communities, and can be considered as an extension of autocorrelation
85 analysis for short time series (fewer than 20 time points) of community data.²⁷ A
86 significant and positive regression slope denotes a community undergoing directional
87 change; while a significant and negative regression slope indicates a community with
88 convergent dynamics (e.g., the species composition is becoming more similar to a
89 community type characteristic of the earlier samples in the series). Moreover, a
90 non-significant slope for the regression implies that there is either stochastic variation
91 or high stability over time. Furthermore, the slope of the regression and the regression
92 R^2 can be used as a measure of rate of change across sampling intervals. TLA has
93 been successfully used for estimating anthropogenically perturbed fish,²⁸ plant, and
94 freshwater zooplankton assemblages over a time scale of one to a few decades.²⁷
95 However studies on freshwater bacterioplankton and/or phytoplankton communities
96 using this approach are rare. Although questions have been raised about the power,
97 and so usefulness of time lag analyses,²⁹ it has been widely applied and has the
98 advantages of 'computational ease, its easy comprehensibility... and the possibility of
99 characterizing and comparing the temporal dynamics of large numbers of
100 communities with a single measure'.³⁰

101 In this study, we used classical denaturing gradient gel electrophoresis (DGGE) to
102 analyze the abundant bacterioplankton community and used microscopy to investigate
103 the abundant phytoplankton community in three subtropical reservoirs, southeast
104 China. Although DGGE is starting to be superseded by sequencing techniques, it has
105 advantages of lower cost and is a reasonable approach for characterizing the more
106 dominant taxa. These three drinking water reservoirs provide interesting systems for
107 investigating the response of microbial communities to environmental changes,
108 because, these reservoirs were exhibiting an early stage of eutrophication - their
109 trophic state was unstable, and it declined gradually from middle eutrophic to
110 oligo-mesotrophic levels during the study period. We further quantified the temporal
111 variation of both bacterioplankton and phytoplankton communities using TLA, and
112 compared the response of both communities to the changed environment.
113 Cyanobacteria have in the past been considered 'algae' (blue green algae) and
114 considered alongside the photosynthetic eukaryotic plankton, indeed because they can
115 be identified and counted in water samples using microscopy they are usually
116 considered part of the phytoplankton – while in freshwater ecology 'bacteria' often
117 mainly refers to heterotrophic bacteria.³¹ In our analyses we treat cyanobacteria
118 mainly as phytoplankton (a functional/ecological classification that makes sense
119 because of their photosynthetic nature) but also explore the implications of classing
120 them as bacteria (the correct phylogenetic classification). The aims of this study were
121 (1) to quantify and compare the temporal patterns between bacterioplankton and
122 phytoplankton communities in three subtropical reservoirs; (2) to reveal the response
123 mechanisms of bacterioplankton and phytoplankton communities to the reservoir
124 environmental changes.

125 MATERIAL AND METHODS

126 **Sample collection.** This study was carried out in three reservoirs near Xiamen city,
127 southeast China (Shidou Reservoir - SD, 118°00'E, 24°42'N; Bantou Reservoir - BT,
128 118°01'E, 24°40'N, Tingxi Reservoir - TX, 118°08'E, 24°48'N); full details of study
129 reservoirs information were showed in **our previous study.**³² The main purposes of
130 these reservoirs are flood control, hydroelectric power, irrigation, and water supply
131 for the city of Xiamen. This area has a subtropical humid monsoon climate with an
132 annual mean precipitation of 1350 mm and an annual mean temperature of 20 °C. The
133 rainfall is concentrated in warm months (April to September), while in cold months
134 (October to March) rainfall is much lower.³³

135 Three sampling stations were selected at each reservoir in the riverine zone,
136 transitional zone, and lacustrine zone, respectively. Surface water samples (upper 50
137 cm) were collected bi-monthly in each station from May 2010 to March 2011,
138 **therefore 18 samples were collected in total from each reservoir.** Water samples were
139 subsequently divided into three subsamples: one for water chemistry analyses, the
140 others for bacterioplankton and phytoplankton analyses, respectively. **All water**
141 **samples were stored in the dark at 4 °C and returned to the laboratory within two**
142 **hours for further processing.** For the phytoplankton analysis, a total of 2.5 L surface
143 water samples were fixed in situ with 1% Lugol's iodine solution and were
144 concentrated to a final volume of 50 mL.¹⁹ For the bacterioplankton analysis, 400 ml
145 water was filtered through a 0.22-μm pore size polycarbonate filter (47 mm diameter,
146 Millipore, Billerica, MA, USA). The filters were stored at -80 °C until further use.

147 **Environmental analysis.** Water temperature (WT), pH, dissolved oxygen (DO),
148 electrical conductivity (EC) and chlorophyll *a* (Chl *a*) were measured in situ with a
149 Hydrolab DS5 multi-parameter water quality analyzer (Hach, Loveland, CO, USA).
150 Water transparency was determined with a 30 cm Secchi disk. Total nitrogen (TN),
151 ammonium nitrogen (NH₄-N), nitrite and nitrate nitrogen (NO_x-N), total phosphorus
152 (TP), and phosphate phosphorus (PO₄-P) were measured following methods used in
153 our previous study.²³

154 The comprehensive trophic state index was calculated according to classical
155 Carlson TSI based on three limnological parameters namely chlorophyll *a*, Secchi
156 disk transparency, and total phosphorus.^{32, 34} Where $0 < \text{TSIc} \leq 30$ = oligotrophic, 30
157 $< \text{TSIc} \leq 40$ = oligo-mesotrophic, $40 < \text{TSIc} \leq 50$ = mesotrophic, $50 < \text{TSIc} \leq 60$ =
158 light eutrophic, $60 < \text{TSIc} \leq 70$ = middle eutrophic, and $70 < \text{TSIc} \leq 100$ =
159 hypereutrophic.

160 **Phytoplankton analysis.** Phytoplankton were identified and counted using an
161 inverted microscope (Motic, Xiamen, China) according to Shen *et al.*, Zhang and
162 Huang, and Hu and Wei.³⁵⁻³⁷ Three subsamples were investigated for each sample and
163 at least 500 individuals were counted for each subsample. To compare the
164 bacterioplankton and phytoplankton communities, we transformed the phytoplankton
165 abundance to biovolume according to Paver *et al.*³⁸ Biovolume was estimated from

cell numbers and cell size measurements.^{19, 39} In using denaturing gradient gel electrophoresis it is difficult to detect microbes with abundances of < 1% of the total community.^{40, 41} Therefore, to improve the comparability between bacterioplankton and phytoplankton, we performed the statistical analyses using only abundant phytoplankton species (> 1% biovolume in a sample).

DNA extraction and PCR amplification. Total DNA was extracted directly from the filter using an E.Z.N.A. DNA Kit (Omega Bio-Tek, Norcross, GA, USA) according to the manufacturer's instructions. The extracted DNA was dissolved in 50 µl TE buffer, quantified by spectrophotometer and stored at -20 °C until further use.

The 16S rRNA gene fragments were amplified with the primers 341F-GC (5'-CGC CCG CCG CGC CCC GCG CCC GTC CCG CCG CCC CCG CCC GCC TAC GGG AGG CAG CAG-3') and 907R (5'-CCG TCA ATT CMT TTG AGT TT-3')⁴² under the following PCR conditions: 5 min denaturation at 94°C and 10 touchdown cycles at 94°C for 0.5 min, 67°C (with the temperature decreasing 1°C each cycle) for 0.5 min, 72°C for 1 min; followed by 20 cycles at 94°C for 0.5 min, 57°C for 0.5 min, 72°C for 1 min and a final extension at 72°C for 10 min. Each 50 µl PCR reaction contained 0.3 µM each primer, 2.5 U *Taq* DNA polymerase (TaKaRa, Otsu, Shiga, Japan), 1.5 mM MgCl₂, 200 µM each deoxynucleoside triphosphate, and approximately 40 ng of template DNA in 1× PCR buffer.

Denaturing gradient gel electrophoresis and sequencing. Denaturing gradient gel electrophoresis (DGGE) was performed using a DCode mutation detection system (Bio-Rad, Hercules, CA, USA). Samples containing equal amounts of PCR amplicons were loaded onto 6% (w/v) polyacrylamide gels (37.5 : 1 acrylamide : bisacrylamide) in 1 × Tris-acetate-EDTA (TAE) buffer. The denaturing gradient of 30%–60% was applied for separation of the 16S rRNA genes and 100% denaturant is defined as 7 M urea and 40% (v/v) deionized formamide, respectively. Electrophoresis was performed at 60 °C with a constant voltage of 100 V for 16 h. The DGGE gels were stained with SYBR Green I nucleic acid stain for 30 min in 1× TAE buffer, rinsed in distilled water, and then visualized with UV radiation by using Gel Doc EQ imager (Bio-Rad, Hercules, CA, USA). DGGE patterns were analyzed using the Quantity One software (Bio-Rad, Hercules, CA, USA)⁴³, and were carefully checked and corrected manually. The bands occupying the same position in the different lanes of the gel were identified. The relative abundance matrix was constructed for all lanes, taking into account the relative intensity of individual bands in each lane.

Dominant DGGE bands were excised from the gels and eluted overnight in autoclaved Milli-Q water at 4 °C. The eluted DNA was reamplified with the original primer set (without GC clamp). PCR products were purified with the TaKaRa Agarose Gel DNA Purification Kit (Takara, Otsu, Shiga, Japan), then cloned into a pMD18-vector (Takara, Otsu, Shiga, Japan) and transformed into *Escherichia coli* DH5α competent cells (Takara, Otsu, Shiga, Japan). The successfully inserted plasmids were sequenced unidirectionally using an automated sequencer (ABI 3730 Genetic Analyzer, Applied Biosystems, Foster City, CA, USA). All bacterial 16S

208 rRNA sequences were manually checked and modified with BIOEDIT v7.0.9⁴⁴ and
209 then compared with the GenBank database using BLASTN⁴⁵. The bacterial 16S rRNA
210 sequences were classified by the Ribosomal Database Project Classifier with 80%
211 confidence.⁴⁶

212 **Data analysis.** We used principal component analysis (PCA) to display the overall
213 trends in environmental variables.

214 Two Bray-Curtis dissimilarity matrices were constructed using the bacterioplankton
215 relative abundance data and phytoplankton relative biovolume data generated from
216 each sample, respectively. The non-metric multidimensional scaling (NMDS)
217 ordination was used to investigate differences in microbial communities among
218 samples based on Bray-Curtis dissimilarity.⁴⁷

219 The coefficient of variation (CV) was calculated to compare the temporal
220 variability between the relative abundance of bacterioplankton OTUs and relative
221 biovolume of phytoplankton taxa in the reservoirs. Median absolute deviation (MAD)
222 was used to compare the temporal variability in Bray-Curtis dissimilarity between
223 bacterioplankton and phytoplankton communities.⁴⁸

224 To explore the temporal patterns of environment and community dynamics, we
225 performed linear regressions on Bray-Curtis dissimilarity of community composition
226 (dependent variables) versus the square root of the time lags (independent variables)
227 and Euclidean distance of all environmental variables (dependent variables) versus the
228 square root of the time lags (independent variables) through TLA.²⁷

229 We divided the twelve environmental variables into three groups: the first group
230 which is related to eutrophication includes TN, NH₄-N, NO_x-N, TP, PO₄-P,
231 transparency, Chl *a* and TSIC; the second group constitutes of three variables (EC, pH,
232 DO) which are related to physico-chemical factors; the third group comprised only
233 water temperature. We then used a forward-selection procedure with Monte Carlo
234 permutation tests to select the environmental variables which explained a significant
235 ($P < 0.05$) variation of the bacterial and phytoplankton data in each group.⁴⁹ To
236 eliminate collinearity among variables within each group, explanatory variables with
237 the highest variance inflation factor (VIF) were sequentially removed until all VIFs
238 were less than 20.⁴⁹ Finally, significant variables in each group were selected to
239 perform variance partitioning using varpart function with adjusted R² (vegan package
240 in R software). Before the forward-selection procedure, the microbial data were
241 Hellinger transformed. We used principal component analysis (PCA) to show main
242 gradients in explained variance.

243 For all the statistical analyses, the cyanobacteria (bands 7 and 28) and Chloroplast
244 bands (bands 32-34) from the DGGE profile were removed from the bacterioplankton
245 data sets (Supporting Information Figure S1). Before the PCA, forward-selection, and
246 TLA, the environmental variables were log($x+1$) transformed, with the exception of
247 pH, to improve normality and homoscedasticity. All the statistical analyses were
248 performed in CANOCO 4.5, PRIMER 5.0 and R language environment.⁵⁰⁻⁵²

249 **Accession number.** The 16S rRNA gene sequences from this study were deposited
250 in the GenBank under the accession numbers KP721939 to KP721985.

RESULTS

Variations in environmental variables. All of the 12 physico-chemical and biological parameters generally showed clear temporal variations and represented a wide range of environmental conditions (Figure 1 and Supporting Information Figure S2). $\text{NH}_4\text{-N}$ showed seasonal cycle patterns. However, water temperature, EC, pH, TN, TP, and Chl *a* decreased, while DO, transparency, and $\text{NO}_x\text{-N}$ increased gradually during the study period. The trend of $\text{PO}_4\text{-P}$ was irregular in three reservoirs. In addition, the comprehensive trophic state index (TSIc) decreased from middle eutrophic to oligo-mesotrophic levels in the three reservoirs. The highest TSIc (62.3) appeared in the BT riverine zone in May 2010, and the lowest TSIc (34.3) appeared in the SD transitional zone in March 2011 (Supporting Information Figure S2).

Microbial diversity and taxonomic composition. There were 49, 36, and 48 distinct bacterial DGGE bands in the Shidou Reservoir (SD), the Bantou Reservoir (BT), and the Tingxi Reservoir (TX), respectively. The average band number per month was 26 in all three reservoirs (the mean band numbers of SD, BT, and TX were 28, 21, and 30, respectively) (Supporting Information Figure S1). For the phytoplankton, 221 taxa were detected in all three reservoirs, and 50 of them were abundant ($> 1\%$ biovolume in a sample). Further, 35, 28, and 33 abundant phytoplankton taxa were found in the SD, BT, and TX, respectively. The mean richness for abundant taxa was 19 in all three reservoirs. The mean abundant taxa richness of BT (21) was the highest, followed by TX (20) and SD (16).

There were 18, 13, and 16 prominent DGGE bands that were successfully sequenced in SD, BT, and TX, respectively. They were affiliated with the divisions Acidobacteria, Actinobacteria, Bacteroidetes, Chloroflexi, Chloroplast, Cyanobacteria, Proteobacteria, and Verrucomicrobia. In general, Actinobacteria and Proteobacteria were the dominant groups, but they showed no pronounced seasonal variation. In contrast, the phytoplankton community showed a distinct seasonal shift at phylum level (Supporting Information Figure S3). Cyanophyta dominated the phytoplankton communities in the warmest months, but were subsequently replaced by Bacillariophyta and Chlorophyta. In March, Euglenophyta and Cryptophyta were dominant taxa in the SD and TX reservoirs, respectively. The most dominant Cyanophyta species in the SD and BT reservoirs was *Cylindrospermopsis raciborskii*. However, in the TX reservoir, *Raphidiopsis* sp. and *Microcystis flosaquae* were the dominant Cyanophyta species. Interestingly, one sequenced band was affiliated with *Cylindrospermopsis raciborskii* in the SD reservoir (band 7, Supporting Information Figure S1).

Temporal variability of community composition. We obtained NMDS ordination plots for bacterioplankton and phytoplankton separately (Supporting Information Figure S4). Both bacterioplankton and phytoplankton communities showed a seasonal succession from May 2010 to March 2011 in all three reservoirs. Further, the variation in phytoplankton community between months was larger than in the bacterioplankton

community. Also, both the coefficient of variation (CV) of abundant phytoplankton taxa and median absolute deviation (MAD) of phytoplankton Bray-Cutis dissimilarity were pronounced higher than those of bacterioplankton community (Figure 2).

Quantified temporal change in microbial communities and environmental factors. In general, the TLA regressions had significant positive slopes, indicating the microbial communities and environmental conditions were undergoing a directional change (Figure 3). In addition, a steeper regression slope (faster rate of change) and higher R^2 (lower stochasticity or stability) for the phytoplankton communities were detected compared to the bacterioplankton communities (Figure 3). Interestingly, the regression R^2 were slightly lower (All bacterioplankton: 0.111, SD: 0.076, BT: 0.112, TX: 0.157) if the Cyanobacteria and Chloroplast bands were included in the bacterioplankton data sets.

Relationships between microbial communities and environmental factors. The variables which were significantly related to bacterioplankton or phytoplankton communities in each group are shown in Supporting Information Table S1. Results of the variance partitioning showed that the environmental variables explained 72-90% variation of phytoplankton community, but only explained 36-67% variation of the bacterioplankton community. Moreover, the phytoplankton community had a greater explained variance than the bacterioplankton community for eutrophic related factors (pure eutrophic factors + eutro-physico covariation + eutro-temp covariation + covariation of all variables) (Figure 4 and Supporting Information Figure S5). The first two axes of PCA explained 94.8 % of the total variability and effectively captured the main patterns of variation in the original variables.

DISCUSSION

In general, the TN, TP, $\text{NO}_x\text{-N}$, Chl a , and TS_{lc}, which are closely related to eutrophication,⁵³ decreased gradually during the study period. In particular, the TS_{lc} values decreased from middle eutrophic to oligo-mesotrophic levels. In addition, the EC and pH decreased (note that the pH gradually declined to close to 7), while the transparency and DO increased. These results indicated that reservoir water quality gradually improved during the study (Supporting Information Figure S2). There are two potential explanations for this. First, the monsoon climate brought large precipitation in the warmer months, which would have resulted in large nutrient runoff from the reservoir watershed. Following the monsoon, the nutrients concentration decreased due to reduced precipitation. Second, the Cyanophyta dominated the phytoplankton communities in the warm months due to their higher optimum growth temperature.⁵⁴ These Cyanophyta blooms resulted in the high phytoplankton biovolume in warm months. However, the phytoplankton biovolume decreased with the decreasing of Cyanophyta in the SD and BT reservoirs associated with the decline of water temperature (Supporting Information Figure S3). These two explanations are not entirely independent as it is well known the high water

temperature and nutrient level can combine to increase the likelihood of cyanobacterial blooms – indeed modifying nutrient levels has been suggested as a potentially tractable approach to reducing the incidence of such blooms in a warming world.⁵⁵

Microbial communities are considered one of the most promising indicators of environmental changes and aquatic ecosystem states due to their rapid response to environmental changes compared with larger animals and plants.^{56, 57} However, microbial communities with different properties may have different and diverse responses to the environmental changes. Therefore, quantifying the response patterns of bacterioplankton and phytoplankton communities to environmental changes is essential for quantifying and understanding the ecosystem recovery process from water pollution (e.g., eutrophication). In this study, we demonstrated that TLA is a useful diagnostic tool to evaluate the direction, rates, and patterns of community change that were not obvious from our more conventional multivariate methods.

We found that environmental conditions changed gradually over time - however, it is interesting to note that the direction and variation of community dynamics were stronger in the phytoplankton than in the bacterioplankton (Figure 3). Clearly, the environmental effects measured in this study were stronger for the phytoplankton than for the bacterioplankton (Figure 4). In other words, the temporal stability (or stochasticity) of the community was stronger for the bacterioplankton than for the phytoplankton in these reservoirs. A possible explanation is that the dispersal probability of bacteria is greater than that of phytoplankton. Jones *et al.* investigated the spatial and temporal dynamics of bacterioplankton beta diversity based on the decay of similarity across time and space, and identified equivalent temporal (1 day) and spatial (10 m) scales of variation in bacterial community composition.⁵⁸ The equivalence of a day and a few meters in their impact on bacterial community similarity suggests similar ecological processes driving community assembly occur over both space and time.^{58, 59} Soininen highlighted factors, such as dispersal rate, as likely drivers of bacterial community turnover in both space and time.⁵⁹ Over time, dispersal may have important effects on the temporal dynamics of microorganisms.^{21, 25} High dispersal ability allows the microorganisms to have a higher probability of colonizing suitable habitats from regional pools, therefore potentially reducing the variation of community composition through time.^{60, 61} Therefore, it is possible that some, or many, of these microbial populations in reservoirs should be thought of as metapopulations – an idea that is now widely applied to the populations of many macroscopic organisms.^{21, 62} In most cases, the cell size of bacteria is smaller than the size of phytoplankton. Due to their small bodies, free-living bacteria are often assumed to be ubiquitous dispersers and they are presumably more likely to become widely dispersal – possibly by mechanisms such as becoming airborne as waves break.⁶³ It has often been suggested that smaller cell size should lead to the wider dispersal probability in microbes.^{60, 64, 65} It follows that one reason the bacterial community may be more stable in the face of environmental change than the phytoplankton community is because of recolonization after extinction and/or the supplementation of populations by individuals dispersing from elsewhere.

In addition to dispersal, other factors such as high abundance, the potential for rapid growth rates, and rapid evolutionary adaptation through mutations and/or horizontal gene transfer could also allow bacteria to quickly adapt to new environmental conditions and maintain the stability of community composition.³ Indeed, the potential for widespread horizontal gene transfer potentially blurs the distinction between individual and community in prokaryotic ecology.^{66, 67} Therefore, the response of bacterial communities to environmental change may be less sensitive than that of the eukaryotic microbial community. Recently, Jones and colleagues compared the seasonality of bacterioplankton and micro-eukaryotic planktonic communities in a freshwater lake, and found that the eukaryotic species richness at both sampling locations exhibited strong fluctuations with algal blooms and other environmental changes, whereas the annual fluctuations in the numbers of bacterial OTUs were relative stable.⁶⁸ Furthermore, these authors suggested that the bacterial taxonomic richness was less sensitive to seasonal forcing factors (i.e. temperature, salinity, *Prymnesium parvum* cell abundances, and large spring rain event) than the micro-eukaryotic diversity.⁶⁸ In another similar study, Lear and colleagues compared the epilithic bacterial and benthic macroinvertebrate communities as indicator of ecological health in New Zealand streams.⁶⁹ Although these authors considered that the relationship between localized influences and sessile bacteria may be closer than that between localized influences and bacterioplankton, they found that macroinvertebrate community composition showed a clear gradient with the increasing localized human impact, while epilithic bacterial communities were only different at the most impacted sites.⁶⁹ It appears that bacterial communities provided a less sensitive indicator of the prevailing environmental conditions, than did macroinvertebrates at community level.⁶⁹

Last but not least, the phytoplankton communities had larger variance explained by eutrophic related factors (pure eutrophic factors + eutro-physico covariation + eutro-temp covariation + covariation of all variables) than the bacterioplankton communities in our study (Figure 4 and Supporting Information Figure S5). The eutrophication factors such as nitrogen, phosphorus, and transparency can be directly related to the phytoplankton.^{70, 71} On the other hand, the interaction between the temperature and eutrophication factors or between the physico-chemical and eutrophication factors was also highly and directly related to the phytoplankton. As described above, water temperature has a positive effect increasing eutrophication. For example, Cyanophyta may benefit from high temperature, since they have high optimum growth temperatures.⁵⁴ Additionally, algal growth and bloom have dramatic effects on or closely relationship with EC, DO, and pH.⁷²⁻⁷⁴ In contrast, the eutrophication factors have both direct effects on the bacterial community and also indirect effects through changes in phytoplankton community.^{38, 75} In addition, the bacterioplankton community had larger unexplained variance than the phytoplankton. It is possible that unmeasured carbon flux is an important factor that influences the succession of bacterial community.⁷⁶ However, we can safely conclude that the bacterioplankton were less sensitive to the environmental changes of eutrophication variables comparing to phytoplankton in the subtropical reservoir ecosystems.

There are, however, potential limitations in our approach that merit further discussion. We should note the different detectability between DGGE and microscopy. The DGGE is a well-established method and most easily detects microorganisms with abundances in the ecosystem of 1% of the total community. Interestingly, the different in detectability seems to have an obvious influence on bacterial α diversity but not on β diversity, thus DGGE has been most useful to compare community structural changes across time and space.⁴⁰ Galand *et al.* defined the abundant bacterioplankton phylotypes as the phylotypes with a relative abundance >1% within a sample. These authors indicated that the composition of abundant bacterial communities was similar to that of the entire bacterial communities.⁷⁷ Similarly, we previously investigated Chinese lakes and reservoirs using high-throughput sequencing, and we also found very similar spatial patterns in both abundant (relative abundance >1% in a sample, and mean relative abundance of > 0.1% in all samples) and entire bacterioplankton communities (RELATE $\rho = 0.934$, $P < 0.01$).⁷⁸ Another limitation of DGGE is that with bacterial 16S rRNA genes it can detect some phytoplankton including prokaryotic Cyanophyta but also a few Chloroplasts (from eukaryotes). However, the relative abundance of such phytoplankton is low in our bacterial data (Supporting Information Figure S3). Further, Niu *et al.* explored the relationship between phytoplankton blooms and temporal variation of bacterioplankton community composition.⁷⁹ These authors found a serious *Microcystis* bloom and high biomass of Bacillariophyta and Cryptophyta using microscopy, whereas only three of seventy eight DGGE bands were found to affiliate with Cyanobacteria using universal bacterial 16S rRNA gene primer.⁷⁹

In conclusion, our results demonstrated that the temporal community dynamics of both abundant bacterioplankton and phytoplankton showed a significant directional change, corresponding to the environmental changes in the reservoirs. Due to high levels of dispersal, growth rates, evolutionary adaptation, and indirect response to the nitrogen and phosphorus, the temporal stability or stochasticity of abundant bacterioplankton community was greater than that of phytoplankton community. These indicated that the phytoplankton community was more sensitive to environmental changes (i.e. improvements in water quality from a human perspective) than the bacterioplankton community in these reservoir ecosystems. This is important, because analyzing multi-components of ecosystem (e.g. primary producers and decomposers) simultaneously can provide a more comprehensive picture when identifying an aquatic ecosystem experiencing or recovering from an environmental change or stress. Additional investigations of more microbial components at different time scales and environmental gradients are needed in order to better understand the relative roles of eutrophication and global warming in affecting aquatic ecosystems.

ASSOCIATED CONTENT

Supporting Information

Supplementary Figures S1–S5 and Supplementary Table S1 showing additional study details. The Supporting Information is available free of charge on the ACS

462 Publications website.

463 **AUTHOR INFORMATION**

464 **Corresponding Author**

465 *Phone: (+86)-592-6190-775; fax (+86)-592-6190-775; e-mail: jyang@iue.ac.cn

466 **Notes**

467 The authors declare no competing financial interest.

468

469 **ACKNOWLEDGEMENTS**

470 This work was supported by the National Basic Research Program of China
471 (2012CB956103), the National Natural Science Foundation of China (31172114 and
472 31370471), the Knowledge Innovation Program of the Chinese Academy of Sciences
473 (IUEQN201305), and the Natural Science Foundation of Fujian Province
474 (2015J05075).

475 **AUTHOR CONTRIBUTIONS**

476 J.Y.* and L.L. designed research; L.L., J.Y., H.L., and X.Y. performed the
477 experiments; L.L., D.M.W., and J.Y.* analyzed data and wrote the paper.

478

REFERENCES

- (1) Newton, R. J.; Jones, S. E.; Eiler, A.; McMahon, K. D.; Bertilsson, S. A guide to the natural history of freshwater lake bacteria. *Microbiol. Mol. Biol. Rev.* **2011**, *75*, 14–49.
- (2) Reynolds, C. S. *The Ecology of Freshwater Phytoplankton*; Cambridge University Press: Cambridge, UK, 2006.
- (3) Hanson, C. A.; Fuhrman, J. A.; Horner-Devine, M. C.; Martiny, J. B. H. Beyond biogeographic patterns: processes shaping the microbial landscape. *Nat. Rev. Microbiol.* **2012**, *10*, 497–506.
- (4) Zhang, Y.; Zhao, Z. H.; Dai, M. H.; Jiao, N. Z.; Herndl, G. J. Drivers shaping the diversity and biogeography of total and active bacterial communities in the South China Sea. *Mol. Ecol.* **2014**, *23*, 2260–2274.
- (5) Reche, I.; Pulido-Villena, E.; Morales-Baquero, R.; Casamayor, E. O. Does ecosystem size determine aquatic bacterial richness? *Ecology* **2005**, *86*, 1715–1722.
- (6) Smith, V. H.; Foster, B. L.; Grover, J. P.; Holt, R. D.; Leibold, M. A.; deNoyelles, F. Phytoplankton species richness scales consistently from laboratory microcosms to the world's oceans. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 4393–4396.
- (7) Downing, A. S.; Hajdu, S.; Hjerne, O.; Otto, S. A.; Blenckner, T.; Larsson, U.; Winder, M. Zooming in on size distribution patterns underlying species coexistence in Baltic Sea phytoplankton. *Ecol. Lett.* **2014**, *17*, 1219–1227.
- (8) Jones, S. E.; Newton, R. J.; McMahon, K. D. Evidence for structuring of bacterial community composition by organic carbon source in temperate lakes. *Environ. Microbiol.* **2009**, *11*, 2463–2472.
- (9) Wang, J. J.; Shen, J.; Wu, Y. C.; Tu, C.; Soininen, J.; Stegen, J. C.; He, J. Z.; Liu, X. Q.; Zhang, L.; Zhang, E. L. Phylogenetic beta diversity in bacterial assemblages across ecosystems: deterministic versus stochastic processes. *ISME J.* **2013**, *7*, 1310 – 1321.
- (10) Macan, T. T. *Biological Studies of The English Lakes*; Longman: London, UK, 1970.
- (11) Rhodes, G.; Porter, J.; Pickup, R. W. The bacteriology of Windermere and its catchment: insights from 70 years of study. *Freshwater Biol.* **2012**, *57*, 305–320.
- (12) Macan, T. T.; Worthington, E. B. *Life in Lakes and Rivers*; Collins: London, UK, 1951.
- (13) Logue, J. B.; Lindström, E. S. Biogeography of bacterioplankton in inland waters. *Freshwater Rev.* **2008**, *1*, 99–114.
- (14) Dalu, T.; Clegg, B.; Nihwatiwa, T. Temporal variation of the plankton communities in a small tropical reservoir (Malilangwe, Zimbabwe). *T. Roy. Soc. S. Afr.* **2013**, *68*, 85–96.
- (15) Berdjeb, L.; Ghiglione, J. F.; Jacquet, S. Bottom-up versus top-down control of hypo- and epilimnion free-living bacterial community structures in two neighboring freshwater lakes. *Appl. Environ. Microbiol.* **2011**, *77*, 3591–3599.
- (16) Hambright, K. D.; Beyer, J. E.; Easton, J. D.; Zamor, R. M.; Easton, A. C.; Hallidayschult, T. C. The niche of an invasive marine microbe in a subtropical freshwater impoundment. *ISME J.* **2015**, *9*, 256–264.
- (17) Hatosy, S. M.; Martiny, J. B. H.; Sachdeva, R.; Steele, J.; Fuhrman, J. A.; Martiny, A. C. Beta diversity of marine bacteria depends on temporal scale. *Ecology* **2013**, *94*, 1898–1904.
- (18) Jochimsen, M. C.; Kummerlin, R.; Straile, D. Compensatory dynamics and the stability of phytoplankton biomass during four decades of eutrophication and oligotrophication. *Ecol. Lett.* **2013**, *16*, 81–89.
- (19) Lv, H.; Yang, J.; Liu, L. M.; Yu, X. Q.; Yu, Z.; Chiang, P. C. Temperature and nutrients are

- significant drivers of seasonal shift in phytoplankton community from a drinking water reservoir, subtropical China. *Environ. Sci. Pollut. Res.* **2014**, *21*, 5917–5928.
- (20) Reynolds, C. S.; Maberly, S. C.; Parker, J. E.; De Ville, M. M. Forty years of monitoring water quality in Grasmere (English Lake District): separating the effects of enrichment by treated sewage and hydraulic flushing on phytoplankton ecology. *Freshwater Biol.* **2012**, *57*, 384–399.
- (21) Heino, J.; Melo, A. S.; Siqueira, T.; Soininen, J.; Valanko, S.; Bini, L. M. Metacommunity organisation, spatial extent and dispersal in aquatic systems: patterns, processes and prospects. *Freshwater Biol.* **2015**, *60*, 845–869.
- (22) Logue, J. B.; Langenheder, S.; Andersson, A. F.; Bertilsson, S.; Drakare, S.; Lanzen, A.; Lindstrom, E. S. Freshwater bacterioplankton richness in oligotrophic lakes depends on nutrient availability rather than on species-area relationships. *ISME J.* **2012**, *6*, 1127–1136.
- (23) Liu, L. M.; Yang, J.; Yu, X. Q.; Chen, G. J.; Yu, Z. Patterns in the composition of microbial communities from a subtropical river: effects of environmental, spatial and temporal Factors. *PLoS ONE*. **2013**, *8*, e81232.
- (24) Dimitriu, P. A.; Boyce, G.; Samarakoon, A.; Hartmann, M.; Johnson, P.; Mohn, W. W. Temporal stability of the mouse gut microbiota in relation to innate and adaptive immunity. *Environ. Microbiol. Rep.* **2013**, *5*, 200–210.
- (25) Dini-Andreote, F.; Stegen, J. C.; van Elsas, J. D.; Salles, J. F. Disentangling mechanisms that mediate the balance between stochastic and deterministic processes in microbial succession. *Proc. Natl. Acad. Sci. U.S.A.* **2015**, *112*, E1326–E1332.
- (26) Shade, A.; Read, J. S.; Welkie, D. G.; Kratz, T. K.; Wu, C. H.; McMahon, K. D. Resistance, resilience and recovery: aquatic bacterial dynamics after water column disturbance. *Environ. Microbiol.* **2011**, *13*, 2752–2767.
- (27) Collins, S. L.; Micheli, F.; Hartt, L. A method to determine rates and patterns of variability in ecological communities. *Oikos* **2000**, *91*, 285–293.
- (28) Pyron, M.; Lauer, T. E.; Gammon, J. R. Stability of the Wabash River fish assemblages from 1974 to 1998. *Freshwater Biol.* **2006**, *51*, 1789–1797.
- (29) Angeler, D. G.; Viedma, O.; Moreno, J. M. Statistical performance and information content of time lag analysis and redundancy analysis in time series modeling. *Ecology* **2009**, *90*, 3245–3257.
- (30) Kampichler, C.; van der Jeugd, H. P. Determining patterns of variability in ecological communities: time lag analysis revisited. *Environ. Ecol. Stat.* **2013**, *20*, 271–284.
- (31) Moss, B. *Ecology of Freshwaters*; Wiley-Blackwell: Chichester, UK, 2010.
- (32) Yang, J.; Yu, X. Q.; Liu, L. M.; Zhang, W. J.; Guo, P. Y. Algae community and trophic state of subtropical reservoirs in southeast Fujian, China. *Environ. Sci. Pollut. Res.* **2012**, *19*, 1432–1442.
- (33) Wikipedia Website; <http://en.wikipedia.org/wiki/Xiamen>.
- (34) Carlson, R. E. A trophic state index for lakes. *Limnol. Oceanogr.* **1977**, *22*, 361–369.
- (35) Hu, H. J.; Wei, Y. X. *The Freshwater Algae of China: Systematics, Taxonomy and Ecology*; Science Press: Beijing, China, 2006 (in Chinese).
- (36) Shen, Y. F.; Zhang, Z. S.; Gong, X. J.; Gu, M. R.; Shi, Z. X.; Wei, Y. X. *Modern Biomonitoring Techniques Using Freshwater Microbiota*; China Architecture & Building Press: Beijing, China, 1990 (in Chinese).
- (37) Zhang, Z. S.; Huang, X. F. *Method for Study on Freshwater Plankton*; Science Press: Beijing, China, 1991 (in Chinese).
- (38) Paver, S. F.; Hayek, K. R.; Gano, K. A.; Fagen, J. R.; Brown, C. T.; Davis-Richardson, A. G.;

- Crabb, D. B.; Rosario-Passapera, R.; Giongo, A.; Triplett, E. W.; Kent, A. D. Interactions between specific phytoplankton and bacteria affect lake bacterial community succession. *Environ. Microbiol.* **2013**, *15*, 2489–2504.
- (39) Hillebrand, H.; Durselen, C. D.; Kirschtel, D.; Pollinger, U.; Zohary, T. Biovolume calculation for pelagic and benthic microalgae. *J. Phycol.* **1999**, *35*, 403–424.
- (40) Nocker, A.; Burr, M.; Camper, A. K. Genotypic microbial community profiling: a critical technical review. *Microb. Ecol.* **2007**, *54*, 276–289.
- (41) Pedrós-Alió, C. The rare bacterial biosphere. *Annu. Rev. Mar. Sci.* **2012**, *4*, 449–466.
- (42) Schäfer, H.; Muyzer, G. Denaturing gradient gel electrophoresis in marine microbial ecology. In *Methods in Microbiology*; Paul, J. H., Ed.; Academic: London, UK, 2001; Vol. 30, pp 425–468.
- (43) Schauer, M.; Massana, R.; Pedrós-Alió, C. Spatial differences in bacterioplankton composition along the Catalan coast (NW Mediterranean) assessed by molecular fingerprinting. *FEMS Microbiol. Ecol.* **2000**, *33*, 51–59.
- (44) Hall, T. *Biological Sequence Alignment Editor for Win95/98/NT/2K/XP*; Ibis Biosciences: Carlsbad, CA, USA, 1999.
- (45) Altschul, S. F.; Gish, W.; Miller, W.; Myers, E. W.; Lipman, D. J. Basic local alignment search tool. *J. Mol. Biol.* **1990**, *215*, 403–410.
- (46) Wang, Q.; Garrity, G. M.; Tiedje, J. M.; Cole, J. R. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* **2007**, *73*, 5261–5267.
- (47) Clarke, K. R.; Gorley, R. N. *PRIMER v5: User Manual/Tutorial*; PRIMER-E Ltd: Plymouth, UK, 2001.
- (48) Shade, A.; Caporaso, J. G.; Handelsman, J.; Knight, R.; Fierer, N. A meta-analysis of changes in bacterial and archaeal communities with time. *ISME J.* **2013**, *7*, 1493–1506.
- (49) Ter Braak, C. J.; Smilauer, P. *CANOCO Reference Manual and CanoDraw for Windows User's Guide: Software for Canonical Community Ordination (version 4.5)*; Microcomputer Power: Ithaca, New York, 2002.
- (50) Lepš, J.; Šmilauer, P. *Multivariate Analysis of Ecological Data Using CANOCO*; Cambridge University Press: Cambridge, UK, 2003.
- (51) Scheiner, S. *Design and Analysis of Ecological Experiments*; Chapman & Hall: New York, USA, 1998.
- (52) R Development Core Team. R: A language and environment for statistical computing (Vienna: R Foundation for Statistical Computing). 2008.
- (53) Lewis Jr, W. M.; Wurtsbaugh, W. A.; Paerl, H. W. Rationale for control of anthropogenic nitrogen and phosphorus to reduce eutrophication of inland waters. *Environ. Sci. Technol.* **2011**, *45*, 10300–10305.
- (54) Kosten, S.; Huszar, V. L. M.; Becares, E.; Costa, L. S.; van Donk, E.; Hansson, L. A.; Jeppesen, E.; Kruk, C.; Lacerot, G.; Mazzeo, N.; De Meester, L.; Moss, B.; Lurling, M.; Nöges, T.; Romo, S.; Scheffer, M. Warmer climates boost cyanobacterial dominance in shallow lakes. *Global Change Biol.* **2012**, *18*, 118–126.
- (55) Scheffer, M.; Barrett, S.; Carpenter, S. R.; Folke, C.; Green, A. J.; Holmgren, M.; Hughes, T. P.; Kosten, S.; van de Leemput, I. A.; Nepstad, D. C.; van Nes, E. H.; Peeters, E. T. H. M.; Walker, B. Creating a safe operating space for iconic ecosystems. *Science* **2015**, *347*, 1317–1319.
- (56) McCormick, P. V.; Cairns, J. Algae as Indicators of Environmental-Change. *J. Appl. Phycol.* **1994**, *6*, 509–526.

- (57) Paerl, H. W.; Dyble, J.; Moisander, P. H.; Noble, R. T.; Piehler, M. F.; Pinckney, J. L.; Steppe, T. F.; Twomey, L.; Valdes, L. M. Microbial indicators of aquatic ecosystem change: current applications to eutrophication studies. *FEMS Microbiol. Ecol.* **2003**, *46*, 233–246.
- (58) Jones, S. E.; Cadkin, T. A.; Newton, R. J.; McMahon, K. D. Spatial and temporal scales of aquatic bacterial beta diversity. *Front. Microbiol.* **2012**, *3*, 318.
- (59) Soininen, J. Species turnover along abiotic and biotic gradients: patterns in space equal patterns in time? *Bioscience* **2010**, *60*, 433–439.
- (60) Wilkinson, D. M.; Koumoutsaris, S.; Mitchell, E. A.; Bey, I. Modelling the effect of size on the aerial dispersal of microorganisms. *J. Biogeogr.* **2012**, *39*, 89–97.
- (61) Yang, J.; Smith, H. G.; Sherratt, T. N.; Wilkinson, D. M. Is there a size limit for cosmopolitan distribution in free-living microorganisms? A biogeographical analysis of testate amoebae from polar areas. *Microb. Ecol.* **2010**, *59*, 635–645.
- (62) Nee, S. Metapopulations and their spatial dynamics. In *Theoretical Ecology*; May, R. M.; McLean, A. R., Eds.; Oxford University Press: Oxford, UK, 2007; pp 35–45.
- (63) Hamilton, W. D.; Lenton, T. M. Spora and Gaia: How microbes fly with their clouds. *Etho. Ecol. Evol.* **1998**, *10*, 1–16.
- (64) De Bie, T.; De Meester, L.; Brendonck, L.; Martens, K.; Goddeeris, B.; Ercken, D.; Hampel, H.; Denys, L.; Vanhecke, L.; Van der Gucht, K.; Van Wichelen, J.; Vyverman, W.; Declerck, S. A. J. Body size and dispersal mode as key traits determining metacommunity structure of aquatic organisms. *Ecol. Lett.* **2012**, *15*, 740–747.
- (65) Ragon, M.; Fontaine, M. C.; Moreira, D.; López-García, P. Different biogeographic patterns of prokaryotes and microbial eukaryotes in epilithic biofilms. *Mol. Ecol.* **2012**, *21*, 3852–3868.
- (66) Margulis, L.; Sagan, D. *Microcosmos*; Simon and Schuster: New York, USA, 1986.
- (67) O'Malley, M. A. *Philosophy of Microbiology*; Cambridge University Press: Cambridge, UK, 2014.
- (68) Jones, A. C.; Liao, T. S. V.; Najar, F. Z.; Roe, B. A.; Hambright, K. D.; Caron, D. A. Seasonality and disturbance: annual pattern and response of the bacterial and microbial eukaryotic assemblages in a freshwater ecosystem. *Environ. Microbiol.* **2013**, *15*, 2557–2572.
- (69) Lear, G.; Boothroyd, I.; Turner, S.; Roberts, K.; Lewis, G. A comparison of bacteria and benthic invertebrates as indicators of ecological health in streams. *Freshwater Biol.* **2009**, *54*, 1532–1543.
- (70) Calijuri, M. C.; dos Santos, A. C. A.; Jati, S. Temporal changes in the phytoplankton community structure in a tropical and eutrophic reservoir (Barra Bonita, SP-Brazil). *J. Plankton Res.* **2002**, *24*, 617–634.
- (71) Elser, J. J.; Marzolf, E. R.; Goldman, C. R. Phosphorus and nitrogen limitation of phytoplankton growth in the freshwaters of North America: a review and critique of experimental enrichments. *Can. J. Fish. Aquat. Sci.* **1990**, *47*, 1468–1477.
- (72) Sotero-Santos, R. B.; Carvalho, E. G.; Dellamano-Oliveira, M. J.; Rocha, O. Occurrence and toxicity of an *Anabaena* bloom in a tropical reservoir (Southeast Brazil). *Harmful Algae* **2008**, *7*, 590–598.
- (73) Bere, T.; Tundisi, J. G. Influence of ionic strength and conductivity on benthic diatom communities in a tropical river (Monjolinho), São Carlos-SP, Brazil. *Hydrobiologia* **2011**, *661*, 261–276.
- (74) Hallegraeff, G. M. A review of harmful algal blooms and their apparent global increase. *Phycologia* **1993**, *32*, 79–99.

- 654 (75) Chrzanowski, T. H.; Sterner, R. W.; Elser, J. J. Nutrient enrichment and nutrient regeneration
655 stimulate bacterioplankton growth. *Microb. Ecol.* **1995**, *29*, 221–230.
- 656 (76) Vrede, K.; Vrede, T.; Isaksson, A.; Karlsson, A. Effects of nutrients (phosphorous, nitrogen, and
657 carbon) and zooplankton on bacterioplankton and phytoplankton—a seasonal study. *Limnol. Oceanogr.*
658 **1999**, *44*, 1616–1624.
- 659 (77) Galand, P. E.; Casamayor, E. O.; Kirchman, D. L.; Lovejoy, C. Ecology of the rare microbial
660 biosphere of the Arctic Ocean. *Proc. Natl. Acad. Sci. U.S.A.* **2009**, *106*, 22427–22432.
- 661 (78) Liu, L.; Yang, J.; Yu, Z.; Wilkinson, D. M. The biogeography of abundant and rare
662 bacterioplankton in the lakes and reservoirs of China. *ISME J.* **2015**, doi:10.1038/ismej.2015.29.
- 663 (79) Niu, Y.; Shen, H.; Chen, J.; Xie, P.; Yang, X.; Tao, M.; Ma, Z. M.; Qi, M. Phytoplankton
664 community succession shaping bacterioplankton community composition in Lake Taihu, China. *Water*
665 *Res.* **2011**, *45*, 4169–4182.

666

667 **Figure Legends**

668 **Figure 1.** PCA plots showing the overall seasonal trends of environmental variables in Shidou
669 (SD), Bantou (BT) and Tingxi (TX) reservoirs. Temp – water temperature, Trans - transparency,
670 EC – electrical conductivity, TSIC – comprehensive trophic state index. The heavy monsoon rains
671 begin from April to September in the study area (see Material and Methods).

672


673 **Figure 2.** Temporal variability in abundant bacterial OTU abundance and abundant phytoplankton
674 biovolume, and median absolute deviation (MAD) in Bray-Curtis dissimilarity between samples.
675 Data are means \pm standard error (error bars); $N=18$. Statistical analysis is t test (** $P < 0.01$; * $P <$
676 0.05).

677

678 **Figure 3.** Time-lag regression analysis of changes in abundant bacterioplankton, abundant
679 phytoplankton communities, and environmental variables.

680

681 **Figure 4.** Results of abundant bacterioplankton and phytoplankton variance partitioning for each
682 reservoir. Eutro (eutrophic factors), Physico (physico-chemical factors), Temp (water
683 temperature).



Abstract Graphic
45x32mm (300 x 300 DPI)




Figure 1. PCA plots showing the overall seasonal trends of environmental variables in Shidou (SD), Bantou (BT) and Tingxi (TX) reservoirs. Temp – water temperature, Trans - transparency, EC – electrical conductivity, TSic – comprehensive trophic state index. The heavy monsoon rains begin from April to September in the study area (see Material and Methods).
54x16mm (300 x 300 DPI)




Figure 2. Temporal variability in abundant bacterial OTU abundance and abundant phytoplankton biovolume, and median absolute deviation (MAD) in Bray-Curtis dissimilarity between samples. Data are means \pm standard error (error bars); N = 18. Statistical analysis is t test (**P < 0.01; *P < 0.05).
65x24mm (300 x 300 DPI)




Figure 3. Time-lag regression analysis of changes in abundant bacterioplankton, abundant phytoplankton communities, and environmental variables.
180x183mm (300 x 300 DPI)




Figure 4. Results of abundant bacterioplankton and phytoplankton variance partitioning for each reservoir. Eutro (eutrophic factors), Physico (physico-chemical factors), Temp (water temperature). 58x39mm (300 x 300 DPI)