



The effects of exposure to appetitive cues on inhibitory control: A meta-analytic investigation

Andrew Jones^{a,b,*}, Eric Robinson^{a,b}, Jay Duckworth^{a,b}, Inge Kersbergen^{a,b}, Natasha Clarke^{a,b}, Matt Field^{a,b}

^a Department of Psychological Sciences, University of Liverpool, UK

^b UK Centre for Tobacco and Alcohol Studies, University of Liverpool, UK

ARTICLE INFO

Keywords:

Appetitive
Alcohol
Cues
Food
Inhibitory control
Overweight/obese

ABSTRACT

Inhibitory control refers to the ability to stop, change or delay a response, and is often used in order to protect higher order goals. Theoretical models suggest that appetitive cues such as pictures of alcoholic drinks or food evoke strong automatic appetitive responses which lead to transient impairments in inhibitory control, and that these effects of cues may be related to individual differences (e.g. in body mass index, or alcohol consumption). In order to investigate these claims we conducted a random effects meta-analysis of 66 effect sizes (35 alcohol, 31 food) from 37 articles that tested the effect of exposure to appetitive (alcohol/food) cues on indices of inhibitory control. The overall effect of cue exposure was small, but robust (SMD = -0.12 [95% CI -0.23 , -0.02]; $Z = 2.34$, $p = .02$, $I^2 = 84\%$). Exposure to alcohol-related cues significantly impaired inhibitory control (SMD = -0.21 [95% CI = -0.32 , -0.11]; $Z = 4.17$, $p < .001$), however exposure to food-related cues did not lead to impairments (SMD = -0.03 [95% CI = -0.21 , 0.15]; $Z = 0.36$, $p = .720$). There was no evidence that drinking or weight status significantly moderated the effects of cues on inhibitory control. Similarly, cue modality (words, pictures, or smells) did not significantly moderate the effects. Trim and Fill analysis suggested bias in the literature, which when corrected, made the overall effect of cues non-significant. Overall, these findings provide some tentative support for theoretical claims that exposure to appetitive cues prompts transient impairments in inhibitory control. Further research is required to determine the clinical significance of these observations. However, care should be taken when drawing conclusions from a potentially biased evidence base.

1. Introduction

Inhibitory control refers to the ability to stop, change or delay a response that is inappropriate given current environment demands (Logan, Cowan, & Davis, 1984; Logan, Schachar, & Tannock, 1997). This (in)ability is a key component of impulsivity and executive functioning (Bickel, Jarmolowicz, Mueller, Gatchalian, & McClure, 2012), and it overlaps considerably with broader constructs such as self-control and disinhibition (Baumeister, 2014; Tarter, Kirisci, Reynolds, & Mezzich, 2004). Effective inhibition of behaviour permits the suppression of automatic appetitive responses evoked by cues related to unhealthy foods or alcohol. This may allow people the opportunity to make controlled decisions (Wiers, Gladwin, Hofmann, Salemink, & Ridderinkhof, 2013) and maintain their higher order goals such as abstinence from alcohol or weight-loss, even when tempted by environmental cues such as the sight or smell of appetising foods or alcoholic drinks (Jones, Hardman, Lawrence, & Field, 2017; Verbruggen,

Best, Bowditch, Stevens, & McLaren, 2014).

Theoretical models of both addiction and obesity posit that deficits in inhibitory control are an important contributor to the development and maintenance of these conditions (Goldstein & Volkow, 2011; Volkow, Wang, Tomasi, & Baler, 2013). Laboratory research using established measures of inhibitory control such as the Stop Signal and Go/No-Go tasks (Verbruggen & Logan, 2008) generally supports the predictions made by these models. For example, there are robust cross-sectional associations between impairments in inhibitory control and hazardous drinking (Christiansen, Cole, Goudie, & Field, 2012; Smith, Mattick, Jamadar, & Iredale, 2014), and also Body Mass Index (BMI) and indicators of unhealthy eating (Nederkoorn, Smulders, Havermans, Roefs, & Jansen, 2006; Vainik, Dagher, Dubé, & Fellows, 2013). Longitudinal studies demonstrate that deficits in inhibitory control predict subsequent alcohol use (Fernie et al., 2013), transition to dependence (Rubio et al., 2008) and treatment outcomes (Petit et al., 2014). Similar studies also show poorer inhibitory control predicts weight gain over

* Corresponding author. Department of Psychological Sciences, University of Liverpool, Bedford Street South, L69 7ZA. UK.
E-mail address: ajj@liv.ac.uk (A. Jones).

one year (Nederkoorn, Houben, Hofmann, Roefs, & Jansen, 2010).

Findings such as these are generally interpreted as indicating that inhibitory control is a stable trait characteristic that differs between individuals but remains fairly constant within individuals, which is why it reliably predicts between-subject variability in behaviour months or years later. However, more recent theoretical models have suggested that inhibitory control functions as a *transient state* which can fluctuate in response to environmental or internal ‘events’ (De Wit, 2009), and these short term impairments in inhibitory control may increase the immediate risk of temptation and subsequent (re) lapse. In a recent narrative review we (Jones, Christiansen, Nederkoorn, Houben, & Field, 2013a) suggested that individuals have a general stable capacity for inhibitory control, however this capacity can fluctuate (both improve and worsen) in response to environmental and internal events. Exposure to appetitive cues is one potential environmental event that may negatively influence inhibitory control, because those cues evoke automatic appetitive tendencies (Brockmeyer, Hahn, Reetz, Schmidt, & Friederich, 2015; Field, Kiernan, Eastwood, & Child, 2008; Kemps & Tiggemann, 2015; Kemps, Tiggemann, Martin, & Elliott, 2013), and these responses should conflict with inhibition of behaviour. Indeed, exposure to both alcohol (Czapla et al., 2015a,b) and food-related cues (Phelan et al., 2011) results in short-lived impairments in inhibitory control. Furthermore, these ‘cue-specific’ deficits in inhibition may predict greater variance in individual differences in health-related behaviours than general inhibition deficits (Houben, Nederkoorn, & Jansen, 2013; Petit, Kornreich, Noël, Verbanck, & Campanella, 2012), including *ad-libitum* food and alcohol consumption in the laboratory (Field & Jones, 2017; Price, Lee, & Higgs, 2016). However, as with many research questions that are studied intensively, there are some null or equivocal findings in the literature, in which cue-exposure has not impaired inhibitory control (Mainz et al., 2012; Nederkoorn, Baltus, Guerrieri, & Wiers, 2009).

The aim of the present meta-analytic investigation was to quantify the extent to which exposure to appetitive-cues (alcohol and food-related pictures) causes transient impairments in the ability to inhibit behaviour, and to identify procedural variables or participant characteristics that may moderate this effect. Following initial scoping searches, we limited our investigation to alcohol and food-related cues because the vast majority of studies in the field were limited to these domains (however, the disinhibiting effects of smoking- (Luijten, Littel, & Franken, 2011) and drug-related cues (Pike, Stoops, Fillmore, & Rush, 2013), have been investigated). We sought to identify potential moderators of the effects of appetitive cues on inhibitory control, including: drinking/weight status, the modality of cue-exposure (pictorial, lexical, olfactory), type of task used to measure inhibitory control, and to determine if cue-specific inhibitory deficits are associated with unhealthy behaviours or outcomes, such as alcohol consumption or BMI. We hypothesised that inhibitory control would be worse during or after exposure to appetitive cues compared to neutral cues, or the absence of cues. For our moderator analyses we predicted that this effect would be larger in heavier drinkers, individuals with alcohol use disorder and individuals with overweight/obesity because, theoretically, automatic appetitive responses to appetitive cues should be stronger in people who consume them more frequently (Volkow et al., 2013; Wiers et al., 2007). We had no *a priori* predictions regarding cue-modality, but we investigated this on the basis of findings from some individual studies which demonstrated differential effects of cues presented in different modalities on both inhibitory control and subjective craving (Boswell & Kober, 2016; Monk, Sunley, Qureshi, & Heim, 2016). Similarly, we examined the effects of different inhibitory control tasks as each have differing inhibitory pressures and may measure a different type of inhibitory control, e.g. action cancellation versus action inhibition (Eagle, Bari, & Robbins, 2008). However, we made no specific hypothesis as to which may be most affected by cue-exposure.

2. Methods

2.1. Information sources and search strategy

We conducted scoping searches using three commonly used electronic databases (Scopus, PubMed and PsycInfo) in November and December 2015. We pre-registered our protocol and analysis strategy on Open Science Framework (<https://osf.io/c9jf8/>). Full searches were carried out in October 2017. Our literature search was guided by the Preferred Reporting Items for Systematic Review (PRISMA) guidelines. See supplementary materials for full search strategy and terms. Following identification of full text articles we conducted manual searches on reference lists, and identified further articles based on authors' knowledge. In total we identified 35 effect sizes for alcohol and 31 for food.

2.2. Eligibility criteria

All studies had to meet the following criteria in order to be included in the meta-analysis; (i) include human participants aged 18+, (ii) include alcohol or food-related (appetitive) cue exposure, i.e. olfactory or visual cues, prior to or during an inhibitory control task, (iii) a control comparison, for example exposure to neutral cues during, or the absence of cue exposure (baseline) prior to or during, an inhibitory control task. Cue exposure involved food/alcohol and neutral images/words that were embedded into an inhibitory control task (Houben et al., 2013; Jones et al., 2015), or the holding and sniffing of food/alcohol prior to completing an inhibitory control task (Gauggel et al., 2010; Lattimore & Mead, 2015).

2.3. Outcome measure(s)

Studies were required to have an outcome measure of inhibitory control during/following appetitive cue exposure, and either a measure of inhibitory control at baseline (prior to cue-exposure) or during exposure to non-appetitive (neutral) cues. Proposed measures of inhibitory control were cross-checked against previous literature and review papers to ensure that they were validated measures (e.g. Diamond, 2013). All authors agreed on the tasks for inclusion.

2.4. Data extraction and coding

Three independent coders (JD, IK, NC) performed the searches and identified the relevant articles. After removal of duplicates, 4151 unique articles were identified. These articles were screened via title and abstract, which resulted in exclusion of 3819 articles with agreement from all coders. Data were extracted by the coders and cross-checked by the first author. In cases where insufficient data was available the authors were contacted to provide this data. If the authors did not respond to the data request and it was possible, we used Web Plot Digitizer (Version 3.10, Rohatgi, 2016) to estimate means and variances from figures presented in publications, as recommended (Jelicic Kadic, Vucic, Dosenovic, Sapunar, & Puljak, 2016; Vucic, Jelicic Kadic, & Puljak, 2015).

To code moderator variables such as drinking and weight status we first examined if any participants were described in the article in a specific way (e.g. alcohol dependent, overweight). If no explicit claims were made we made group level inferences on alcohol use using established cut-offs for ‘heavy’ or ‘hazardous’ drinking via scores on the AUDIT (score > 8 indicative of hazardous drinking (see Saunders, Aasland, Babor, De La Fuente, & Grant, 1993) or estimates of units of alcohol consumed per week (> 14 units per week indicative of heavy drinking¹). For weight status we examined if group mean BMI > 25 kg/

¹ Note that these cut offs differ by country. Fourteen units per week is the UK

m^2 (for overweight/obese). Three studies compared alcohol-dependent participants to a control group (Noel et al., 2005; Noel et al., 2007; Sion, Jurado-Barba, Alonso, & Rubio-Valladolid, 2017), but provided no information as to whether the control group drank any alcohol. Therefore, the control groups in these studies were not included in any analysis.

2.5. Variables of interest

The indices of inhibition used for each task are stated in Table 1. The most common tasks were the Stop Signal, Go/No-Go and Go/No-Go shifting tasks. The Stop Signal and Go/No-Go tasks require motor inhibition of a pre-potent response following a visual or auditory ‘stop signal’ or ‘No-Go cue’. In the Stop Signal task this cue is presented following a variable delay after initial stimulus onset and therefore motor behaviour has to be *cancelled*, whereas in the No-Go task the No-Go cue is presented concurrently with the target stimulus, and therefore behaviour must be *restrained* rather than cancelled (Eagle et al., 2008). In the shifting version of the task the cues for ‘Go’ and ‘No-Go’ are switched on a block-by-block basis (Meule, 2017). In the anti-saccade task participants have to inhibit an involuntary oculomotor response (saccade) to a visual stimulus that appears in the periphery of a visual display (Hallett, 1978). In the Stroop task (Stroop, 1935) participants have to name the colour of target words whilst ignoring the semantic content of the word (e.g., the word ‘red’ printed in blue ink). Finally, in the flanker task participants have to categorise a target stimulus whilst ignoring distractor stimuli that appear alongside it (Eriksen & Eriksen, 1974). Stop Signal Reaction Time and Commission errors were the most common outcomes from these tasks. We also extracted and coded a number of variables for our main and supplementary analyses, including; type of task used, modality of cue exposure, drinking and weight status, and any correlations with BMI, typical alcohol use or AUDIT scores (see Table 1). We selected these variables as they were the most commonly measured across all studies.

2.6. Statistical analyses

Our main statistical analyses were carried out using Review Manager 5.3 (Cochrane Informatics & Knowledge Management Department, UK, 2014), with supplementary analyses conducted using JASP (JASP team, Version 0.8.4). All outcomes were continuous, therefore we computed the Standardised Mean Difference (SMD) effect size using the equation $SMD = (M_a - M_c)/S_p$, where M_a is the mean inhibitory control measure following exposure to appetitive cues, M_c is the mean following neutral cues (or no cues) and S_p is the pooled standard deviation. We also computed the Standard Error of the SMD; most studies used within-subject designs so we adjusted the SE based on the within-subject correlation (Elbourne et al., 2002), which was requested from authors when not presented in the article. When this correlation was not available it was estimated based on the mean correlation across all effect sizes ($r \sim .60$). The SMD is interpreted as; 0.2 = small effect, 0.5 = moderate effect, and 0.8 = large effect. In this case a negative SMD is indicative of a reduction in inhibitory control following appetitive cue exposure. In order to ensure consistency, effect sizes from outcome measures in which a larger number was indicative of poorer inhibition (e.g. Stop Signal Reaction times, Number/proportion of commission errors) were reversed.

To assess between-study heterogeneity we used the I^2 statistic, calculated as $I^2 = (Q - df/Q) \times 100\%$, where Q is the chi-squared statistic and df is the accompanying degrees of freedom. We used random-effects models due to substantial heterogeneity between studies (Riley, Higgins, & Deeks, 2011). In the case of substantial

heterogeneity we provided estimates of subgroup effects to aid interpretation of the data. Finally, to remove outlying effect sizes we calculated z-scores and identified any effect size which was an extreme value at .001 alpha level (i.e. $Z > 3.30$). As a result one effect size (Petit et al., 2012): Light drinkers: SMD > 6) was excluded from all subsequent analyses. This decision was made a-priori, as evidenced in our pre-registration.

2.7. Characteristics of studies

The majority of studies employed within-subject (repeated measures) designs, in which participants either (a) completed similar inhibitory control tasks during or after exposure to appetitive cues and neutral cues, or (b) completed one inhibitory control task with embedded appetitive and neutral cues that permitted separate indices of inhibitory control to be computed for each type of cue (e.g. (Jones & Field, 2015; Nederkooij et al., 2009; Petit et al., 2012)). We also identified some studies that employed between-subject designs in which participants were randomized to exposure to either appetitive or neutral cues (e.g. (Jones, Rose, Cole, & Field, 2013b; Lattimore & Mead, 2015; Muraven & Shmueli, 2006)). Field and Jones (2017) employed a mixed design in which inhibitory control was measured at baseline in all participants, before a between-subjects cue exposure manipulation (appetitive, neutral), followed by a second measure of inhibitory control. In this case we took the difference between the two groups after cue exposure.

Some studies also contrasted the effects of appetitive cues on inhibitory control in different groups of participants using mixed designs. For example, heavy vs. light drinkers (Nederkooij et al., 2009), people with alcohol dependence vs. controls (Czapla et al., 2015a,b), obese/overweight vs. normal weight (Loeber et al., 2012). In these studies we computed within-subject comparisons based on these groups where possible to allow for individual comparisons to be included in different moderator analyses (see Table 1). Finally, some studies used multiple inhibitory control tasks or parameters (e.g., Adams, Ataya, Attwood, & Munafò, 2013) and in these cases we adjusted the sample sizes in the control conditions ($N^{\text{Control}}/\text{number of tasks}$) accordingly to ensure each comparison could be included in our pooled analyses, as recommended (Higgins & Green, 2011).

3. Results

The article selection process and flow is shown in Fig. 1. Following exclusion of irrelevant articles by title and abstract scanning we identified 37 full-text articles. See Table 1 for full details.

3.1. Pre-registered analyses

3.1.1. Primary hypothesis: the overall effect of appetitive cues on inhibitory control

Our main analysis considered 66 effect sizes. We included a subgroup of appetitive cue-type: alcohol-related ($k = 35$) or food-related ($k = 31$). The overall effect of appetitive cues was small but statistically significant (SMD = -0.12 [95% CI $-0.23, -0.02$]; $Z = 2.34$, $p = .02$, $I^2 = 84\%$). Exposure to alcohol-related cues significantly impaired inhibitory control (SMD = -0.21 [95% CI $-0.32, -0.11$]; $Z = 4.17$, $p < .001$). There was no evidence that food-related cues led to observable deficits in inhibitory control (SMD = -0.03 [95% CI $-0.21, 0.15$]; $Z = 0.36$, $p = .720$). There was weak statistical evidence for subgroup differences ($\chi^2 (1) = 2.97$, $p = .090$). These results suggest that exposure to appetitive-cues impairs inhibitory control compared to neutral cue exposure/no cue exposure, and overall the impairing effects of alcohol-related cues are more robust (see Supplementary Fig. 1).

(footnote continued)

Department of Health Guidelines for low risk drinking.

Table 1
Details of studies included in the meta-analysis.

Authors and Study	Participants and Design	Cue exposure and Control	Outcome	Notes	Included subgroup analyses
Adams et al. (2013)	N = 96 Mean age: 21.0 Inclusion criteria: Social drinkers Exclusion criteria: None stated	Pictorial and Lexical cue exposure Appetitive: Alcohol-related cues and words were embedded into the task Control: Soft-drink images and musical instrument words were embedded into the task	Commission errors to alcohol cues (pictorial) and words (lexical)	Participants completed the task under placebo and alcohol intoxication sessions. Data from the placebo session only was analysed. Participants were also split into heavy and light drinkers based on units consumed	Drinking status: Heavy drinkers, Light drinkers Cue modality: Pictorial, Lexical Task: Go/No-Go Switching
(Czapla et al., 2015a,b)	N = 165 Mean age: 47.0 Inclusion criteria: Alcohol dependent patients or health matched controls Exclusion criteria: Current drug abuse or dependence (except alcohol/nicotine); severe somatic, neurological or psychiatric diseases; pregnancy	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the inhibition task Control: Neutral (geometric shapes) were embedded into the task.	Commission errors	We considered the control group to be light drinkers based on their drinking characteristics (number of drinking days and cumulative alcohol consumption) reported in the article.	Drinking status: Alcoholics, Light drinkers Cue modality: Pictorial Task: Go/No-Go
Czapla et al. (2015b)	N = 16 Mean age: 23.8 Inclusion criteria: Good physical health Exclusion criteria: None stated	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the inhibition task Control: Neutral (geometric shapes) were embedded into the task.	Commission errors	Participants were split into binge and non-binge drinkers. We considered non-binge drinkers to be light drinkers based on their drinking characteristics reported in the article.	Drinking status: Heavy drinkers, Light drinkers Cue modality: Pictorial Task: Go/No-Go
Field & Jones (2017)	N = 81 Mean age = 20.0 Inclusion criteria: Heavy drinking Exclusion criteria: No history of substance use disorder or ADHD	Pictorial and olfactory cue exposure Appetitive: Participants sniffed beer during a task with alcohol cues embedded Control: Participants sniffed water during a task with neutral (water) cues embedded	Stop Signal Reaction time		Drinking status: Heavy drinkers Cue modality: Pictorial, olfactory Task: Stop Signal
Gauget et al. (2010)	N = 20 Mean age: 44.9 Inclusion criteria: Detoxified alcohol dependent patients Exclusion criteria: None stated	Olfactory cue exposure Appetitive: Participants sniffed beer prior to the inhibition task Control: Participants sniffed water prior to the inhibition task	Stop Signal Reaction Time		Drinking Status: Alcoholics Cue exposure type: Olfactory Task: Stop Signal
Jones et al. (2013a,b)	N = 60 Mean age: 21.15 years Inclusion criteria: Heavy drinking Exclusion criteria: No history of alcohol-related problems	Olfactory cue exposure Appetitive: Participants sniffed beer prior to the inhibition task Control: Participants sniffed water prior to the inhibition task	Stop Signal Reaction Time		Drinking Status: Heavy drinkers Cue modality: Pictorial, olfactory Task: Stop Signal
Jones and Field (2015) study 1	N = 64 Mean age: 22.34 years Inclusion criteria: Social drinking Exclusion criteria: No history of alcohol-related problems or ADHD	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the inhibition task Control: Neutral (scenery) cues were embedded into the inhibition task	Stop Signal Reaction Time	Alcohol vs Neutral cues comparison was used. Positively and negatively valenced cues were also included in the Stop Signal task	Drinking status: Heavy drinkers, Light drinkers Cue exposure type: Pictorial Task: Stop Signal
Jones and Field (2015) study 2	N = 117 Mean age: 24.8 Inclusion criteria: Social drinking Exclusion criteria: No history of alcohol-related problems or ADHD	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: Neutral (scenery) cues were embedded into the task	Proportion of correct responses	Alcohol vs Neutral cues comparison was used. Positively and negatively valenced cues were also included in the Stop Signal task	Drinking status: Heavy drinkers, Light drinkers Cue modality: Pictorial Task: Anti-saccade
Kreusch, Vilenne, and Quertemont (2013) study 1	N = 71 Mean age: 21.1 Inclusion criteria: Social drinker Exclusion criteria: Drug consumption; psychiatric symptoms	Pictorial cue exposure Appetitive: Alcohol related cues were embedded into the task Control: Neutral (object images) were embedded into the task	Proportion of commission errors	Sample was split by problem and non-problematic drinkers based on AUDIT scores.	Drinking status: Heavy drinkers and Light drinkers Cue modality: Pictorial Task: Go/No-Go switching

(continued on next page)

Table 1 (continued)

Alcohol-cue exposure articles				
Authors and Study	Participants and Design	Cue exposure and Control	Outcome	Notes
Kreusch et al. (2013) study 2	N = 81 Mean age: 20.0 Inclusion criteria: Social drinking Exclusion criteria: Drug consumption; psychiatric symptoms	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: Neutral (object images) were embedded into the task	Proportion of commission errors	Drinking status: no information Cue modality: Pictorial Task: Go/No-Go Switching
	N = 14 Mean age: 43.5 Inclusion criteria: Detoxified alcohol dependent patients Exclusion criteria: No history of other substance dependence of schizophrenia/schizophreniform disorders	Lexical cue exposure Appetitive: Alcohol-related words were embedded into the task Control: Neutral-related words were embedded into the task	Proportion of commission errors on Stop trials	Drinking Status: Alcoholics Cue modality: Lexical, Olfactory Task: Stop Signal
Mainz et al. (2012)	N = 11 Mean age: 44.0 Inclusion criteria: Detoxified alcohol dependent patients; males Exclusion criteria: None stated	Olfactory cue exposure and imagery Appetitive: Participants imagined situations in which they usually drank along with presentation of the smell of alcohol Control: Participants imagined a place in which they never drink along with presentation of the smell of oranges	Stop Signal Reaction Time	Drinking status: Alcoholics Cue modality: Olfactory Task: Stop Signal
Monk et al. (2016)	N = 40 Mean age: 23.7 Inclusion criteria: None stated Exclusion criteria: None stated	Olfactory and pictorial cue exposure Appetitive: Alcohol-related images were embedded into the task and the smell of vodka was administered via an olfactory mask Control: Letters were embedded into the task and the smell of citrus was administered via an olfactory mask	Commission errors	Drinking Status: Heavy drinkers Cue modality: Pictorial, Olfactory Task: Go/No task
Muraven & Schmeuli (2006)	N = 160 Mean age: 27.03 Inclusion criteria: Social drinking Exclusion criteria: Score > 2 on Short Michigan Alcohol Screening test; planning for pregnancy; taking anti-craving medicine; serious mental illness	Olfactory cue Appetitive: Participants sniffed beer prior to the inhibition task Control: Participants sniffed water prior to the inhibition task	Correct inhibitory responses	Drinking status: no information provided Cue modality: Olfactory Task: Stop Signal
Nederkoorn et al. (2009)	N = 64 Mean age: 21.05 Inclusion criteria: Social drinking Exclusion criteria: None stated	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: Neutral (shades of grey) cues were embedded into the task	Stop Signal Reaction Time	Drinking status: Heavy drinkers, Light drinkers Cue modality: Pictorial Task: Stop Signal
Nikolaou, Field, and Duka (2013)	N = 14 Mean age: 23.9 Inclusion criteria: Individuals who drank > 2 units per week Exclusion criteria: History of psychiatric, neurological or physical conditions; treatment for dependence; use of medication	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: (object images) were embedded into the task	Accuracy rate on incongruent flanker trials	Drinking status: Heavy drinkers Cue modality: Pictorial Task: Flanker
Noel et al. (2007)		Lexical cue exposure Appetitive: Alcohol-related words were embedded into the task Control: Neutral-related words were embedded into the task	Commission errors to alcohol words	Drinking status: Alcoholics, no information on controls Cue modality: Lexical Task: Go/No-Go Switching

(continued on next page)

Table 1 (continued)

Alcohol-cue exposure articles				
Authors and Study	Participants and Design	Cue exposure and Control	Outcome	Notes
Noel et al. (2009)	N = 40 Mean age: 44.1 Inclusion criteria: Alcoholics had to meet DSM criteria; controls aged matched Exclusion criteria: Current DSM-IV Axis I diagnoses, a history of significant medical illness, head injury; use of other psychotropic drugs or substances that influence cognition; overt cognitive dysfunction.	Lexical cue exposure Appetitive: Alcohol-related words were embedded into the task Control: Neutral-related words were embedded into the task	Decision bias (C)	No information was given for control group alcohol consumption. Therefore we were unable to include this group in our analyses.
	N = 30 Mean age: 33.6 Inclusion criteria: Meeting DSM-IV criteria for substance dependence Exclusion criteria: Psychosis, documented head injury or seizure disorder	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: Neutral (non-alcohol related) cues were embedded into the task	Commission errors	Drinking status: Alcoholics, no information on controls Cue modality: Lexical Task: Go/No-Go Switching
	N = 35 Mean age: 21.3 Inclusion criteria: Regular alcohol consumption Exclusion criteria: Alcohol abstainers; major medical problems; history of alcohol-related problems	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: Neutral (non-alcohol related) cues were embedded into the task	Commission errors	Drinking status: Heavy drinkers; Light Drinkers Cue modality: Pictorial Task: Go/No-Go task
Petit et al. (2012)	N = 54 Mean age: 45.0 Inclusion criteria: Patients with alcoholism undergoing treatment or healthy age/sex matched controls Exclusion criteria: Diagnosis of axis I disorders (DSM-IV) or significant CNS or visual impairment	Pictorial cue exposure Appetitive: Alcohol-related cues were embedded into the task Control: Neutral (non-alcohol related) cues were embedded into the task	Commission errors	Drinking status: Alcoholics, Light drinkers Cue modality: Pictorial Task: Go/No-Go task
Slon et al. (2017)	N = 85 Mean age: Not reported Inclusion criteria: Alcohol dependent patients attending detoxification and recovery Exclusion criteria: Psychiatric morbidities	Lexical cue exposure Appetitive: Alcohol-related words were embedded into the task Control: Neutral (non-alcohol related) words were embedded into the task	Stop Signal Reaction time	Drinking status: Alcoholics Cue modality: Lexical Task: Stop Signal
Weafer and Fillmore (2012)	N = 50 Mean age: 23.9 Inclusion criteria: Adult beer drinkers Exclusion criteria: Head trauma, psychiatric disorder, or substance abuse disorder	Pictorial cue exposure Appetitive: Alcohol-related cues (Beer only) were embedded into the task Control: Neutral (non-alcohol related) cues were embedded into the task	Proportion of Inhibition errors	Drinking status: Heavy drinkers Cue modality: Pictorial Task: Go/No-Go task
Weafer and Fillmore (2014)	N = 40 Mean age: 23.3 Inclusion criteria: Adult beer drinkers Exclusion criteria: Head trauma, psychiatric disorder, or substance abuse disorder	Pictorial cue exposure Appetitive: Alcohol-related cues (Beer only) were embedded into the task Control: Neutral (non-alcohol related) cues were embedded into the task	Proportion of Inhibition errors	Drinking status: Heavy drinkers Cue modality: Pictorial Task: Go/No-Go task
Food-cue exposure articles				
He et al. (2014)	N = 30 Mean age: 19.7 Inclusion criteria: None stated Exclusion criteria: Individuals with neuropsychiatric disorders, medication or health issues which interfered with neuroimaging data Between Subjects	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Low calorie food-related cues were embedded into the task	Commission errors	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go Switching

(continued on next page)

Table 1 (continued)

Alcohol-cue exposure articles					
Authors and Study	Participants and Design	Cue exposure and Control	Outcome	Notes	Included subgroup analyses
Houben et al. (2013)	N = 87 Mean age: 26.2 Inclusion criteria: Female Exclusion criteria: Pregnancy or current dieters	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Arbitrary letters were embedded into the task	Stop Signal Reaction time		Weight Status: Normal weight Cue modality: Pictorial Task: Stop Signal
Hume, Howells, Rauch, Kroff, and Lambert (2015)	N = 81 Mean age: 31.7 Inclusion criteria: Female Exclusion criteria: Known metabolic disease, pregnancy or lactation in the last three months	Pictorial cue exposure Appetitive: Palatable food-related cues were embedded into the task Control: Neutral (office equipment) cues were embedded into the task	Errors	Participants were split into normal weight, overweight and obese based on their BMI	Weight Status: Normal weight; Overweight/Obese Cue modality: Lexical Task: Stroop
Lattimore and Mead (2015)	N = 51 Mean age: 25.0 Inclusion criteria: Female Exclusion criteria: BMI < 18.5 or > 39.5; pregnant; diabetes diagnosis; sought medical help for an eating disorder; using medication	Olfactory cue exposure Appetitive: Participants had to smell food items and touch them against their lips without eating them. Control: no information given	Stop Signal Reaction Time	Participants were split into low-impulsive and high impulsive based on scores on the Barratt Impulsivity Scale	Weight Status: Normal weight, Overweight/Obese Cue modality: Olfactory Task: Stop Signal
Loeber et al. (2012)	N = 40 Mean age: 46.4 Inclusion criteria: Obese or Normal weight Exclusion criteria: Severe psychiatric, neurological, or somatic diseases; untreated endocrine illnesses; history of surgical interventions in the gastro-intestinal system; medications	Lexical cue exposure Appetitive: Food-related words were embedded into the task Control: Neutral (object) words were embedded into the task	Commission errors	Participants were recruited specifically as obese (BMI > 30) or normal weight (BMI 18.5–25).	Weight Status: Normal weight, Overweight/Obese Cue modality: Lexical Task: Go/No-Go Switching
Loeber, Grosshans, Herpertz, Kiefer, and Herpertz (2013)	N = 48 Mean age: 24.5 Inclusion criteria: 18 and 65; BMI between 18.5 and 25 Exclusion criteria: Severe psychiatric, neurological, or somatic diseases; untreated endocrine illnesses as well as psychoactive medication; pregnancy	Lexical cue exposure Appetitive: Food-related words were embedded into the task Control: Neutral (clothing) words were embedded into the task	Commission errors	Participants were split into hungry or satiated based on scores on a Grand Hunger Scales	Weight Status: Normal weight Cue modality: Lexical Task: Go/No-Go Switching
Lyu, Zheng, Chen, and Jackson (2017)	N = 62 Mean age: 21.5 Inclusion criteria: None stated Exclusion criteria: None stated	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (household items) cues were embedded into the task	Commission errors	Participants were recruited specifically for Binge Eating tendencies or age-matched controls. High calorie vs Neutral comparison was used. Low calorie images were also used in the task	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go Switching
Manasse et al. (2016)	N = 90 Mean age: 50.36 Inclusion criteria: Treatment seeing overweight and obese females. Exclusion criteria: None stated	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (office equipment) cues were embedded into the task	Stop Signal Reaction time	Participants were recruited specifically as overweight binge eaters or overweight with no history of binge eating. Food vs Neutral comparison was used. Positive images were also used in the task.	Weight Status: Overweight/Obese Cue modality: Pictorial Task: Stop Signal
Meule, Lukito, Vögele, and Kubler (2011)	N = 61 Mean age: 22.1 Inclusion criteria: Female Exclusion criteria: Self-reported current diagnoses of psychiatric disorders and psychopharmacological medication	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (office equipment) cues were embedded into the task	Commission errors	Participants were split into restrained or unrestrained eaters based on the Restraint Scale	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go
Meule, Lutz, Vögele, and Kübler (2012)	N = 50 Mean age: 22.3 Inclusion criteria: Female Exclusion criteria: Mental disorders, psychoactive medication, under- or overweight; aged > 40	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (office equipment) cues were embedded into the task	Commission errors	Participants were split into high and low food addiction groups based on the Yale Food Addiction Scale	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go

Continued on next page

(continued on next page)

Table 1 (continued)

Alcohol-cue exposure articles				
Authors and Study	Participants and Design	Cue exposure and Control	Outcome	Notes
Meule, Lutz, Vögele, and Kibler (2014a) study 1	N = 50 Mean age: 22.3 Inclusion criteria: Female Exclusion criteria: Mental disorders, psychoactive medication, under- or overweight; aged > 40	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (office equipment) cues were embedded into the task	Commission errors	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go Switching
Meule et al. (2014a) study 2	N = 102 Mean age: 22.8 Inclusion criteria: Female Exclusion criteria: Mental disorders, psychoactive medication, aged > 40	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (office equipment) cues were embedded into the task	Commission errors	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go Switching
Meule and Kubler (2014)	N = 55 Mean age: 24.4 Inclusion criteria: Female Exclusion criteria: None stated	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Low calorie food-related were embedded into the task	Commission errors	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go Switching
Meule et al. (2014b)	N = 50 Mean age: 22.3 Inclusion criteria: Female Exclusion criteria: Mental disorders, psychoactive medication, aged > 40	Pictorial cue exposure Appetitive: High calorie food-related cues were embedded into the task Control: Neutral (household items) cues were embedded into the task	Stop Signal Reaction Time	Weight Status: Normal weight Cue modality: Pictorial Task: Stop Signal
Phelan et al. (2011)	N = 48 Mean age: 46.8 Inclusion criteria: None stated Exclusion criteria: Binge eating, food allergies, and vegetarianism	Lexical cue exposure Appetitive: High calorie food-related words were embedded into the task Control: Neutral words were embedded into the task	Number of valid words	Weight Status: Normal weight, Overweight/Obese Cue modality: Lexical Task: Stroop
Schag et al. (2013)	N = 76 Mean age: 39.67 Inclusion criteria: Females Exclusion criteria: Impaired vision; somatic diseases; medications; pregnancy or lactation; psychosis or bipolar disorder	Pictorial cue exposure Appetitive: Food-related cues were embedded into the task Control: Neutral (non-food) related cues were presented prior to the task	Anti-saccade errors	Weight Status: Normal weight, Overweight/Obese Cue modality: Pictorial Task: Anti-saccade
Yeomans and Brace (2015)	N = 96 Mean age: 21.4 Inclusion criteria: None stated Exclusion criteria: None stated	Pictorial cue exposure Appetitive: Food-related cues were embedded into the task Control: Neutral related cues were presented prior to the task	Commission errors	Weight Status: Normal weight Cue modality: Pictorial Task: Go/No-Go task




Fig. 1. Meta-analysis search results and flow chart.

3.1.2. Moderation by drinking status

We identified $k = 9$ effect sizes for light drinkers, $k = 15$ effect sizes for heavy drinkers and $k = 8$ effect sizes for alcohol dependent patients². There was no significant effect of alcohol-related cues on inhibitory control in light drinkers ($SMD = -0.15$ [95% CI = $-0.33, 0.03$]; $Z = 1.59$, $p = .110$; $I^2 = 53\%$). There was a significant effect of alcohol-related cues on inhibitory control in heavy drinkers ($SMD = -0.26$ [95% CI = $-0.46, -0.06$]; $Z = 2.59$, $p < .010$; $I^2 = 80\%$), and alcohol-dependent patients ($SMD = -0.22$ [95% CI = $-0.41, -0.04$]; $Z = 2.35$, $p = .020$; $I^2 = 58\%$). However, there was no significant subgroup effect ($X^2(2) = 0.79$, $p = .67$) suggesting that drinking status did not reliably moderate the effects of alcohol-related cues on inhibitory control.

3.1.3. Moderation by weight status

We identified $k = 10$ effect sizes for overweight/obese participants and $k = 21$ effect sizes for normal weight or underweight participants. There was no significant effect of food-related cues on inhibitory control in overweight/obese individuals ($SMD = -0.31$ [95% CI = $-0.73, 0.12$]; $Z = 1.42$, $p = .16$; $I^2 = 91\%$) or normal weight individuals ($SMD = 0.09$ [95% CI = $-0.10, 0.28$]; $Z = 0.95$, $p = .34$; $I^2 = 86\%$). There was weak evidence for a subgroup effect ($X^2(1) = 2.83$, $p = .09$).

² We were unable to categorise individuals in Kreusch et al. (2013, study 2), or Muraven and Schmeuli (2006) as there was no information provided on participants' alcohol consumption.

3.1.4. Moderation by modality of cue-exposure

Across both food and alcohol cues we identified $K = 42$ effect sizes from studies that employed pictorial cues, $k = 16$ effect sizes from studies that employed lexical (word) cues and $k = 10$ effect sizes from studies that employed olfactory/*in vivo* cues. Note that two studies³ used combined cue-exposure paradigms. Therefore they contributed to more than one group in these analyses, but removal of effect sizes from these studies did not significantly alter the results. Exposure to olfactory/*in vivo* cues led to significant impairments in inhibitory control ($SMD = -0.24$ [95% CI = $-0.41, -0.07$]; $Z = 2.83$, $p < .001$, $I^2 = 65\%$). Whereas, pictorial ($SMD = -0.07$ [95% CI = $-0.19, 0.05$]; $Z = 1.07$, $p = .280$, $I^2 = 83\%$) and lexical ($SMD = -0.26$ [95% CI = $-0.57, 0.06$]; $Z = 1.62$, $p = .110$, $I^2 = 90\%$) cues did not significantly impair inhibitory control. However, the test for subgroup differences was not statistically significant ($X^2(2) = 3.47$, $p = .180$).

Due to the differences between alcohol-related and food-related cues we analysed the effects of cue-exposure modality separately. For alcohol, we identified $k = 23$ effect sizes from studies that employed pictorial cues, $k = 6$ effect sizes were from studies that employed lexical cues and $k = 8$ effect sizes were from studies that employed olfactory/*in vivo* cues. Pictorial alcohol-related cues ($SMD = -0.22$ [95% CI = $-0.35, -0.08$]; $Z = 3.11$, $p = .002$; $I^2 = 75\%$) and olfactory alcohol-related cues ($SMD = -0.27$ [95% CI = $-0.46, -0.08$]; $Z = 2.76$, $p < .001$; $I^2 = 69\%$) significantly impaired inhibitory control, however lexical cues did not have a significant effect ($SMD = -0.13$ [95%

³ Field and Jones (2017) combined pictorial and olfactory cues, whereas Kreusch et al. (2017) combined lexical and olfactory cues.

CI = -0.27, 0.02; $Z = 1.74$, $p = .080$; $I^2 = 0\%$). The test for subgroup differences was not significant ($X^2(2) = 1.51$, $p = .470$).

For food, we identified $k = 19$ effect sizes from studies that employed pictorial cues, $k = 10$ effect sizes from studies that employed from lexical cues and $k = 2$ effect sizes were from studies that employed olfactory/*in vivo* cues. Only lexical cues impaired inhibitory control (SMD = -0.30 [95% CI = -0.83, -0.24]; $Z = 1.09$, $p = .28$, $I^2 = 94\%$); pictorial (SMD = 0.10 [95% CI = -0.06, 0.27]; $Z = 1.20$, $p = .23$, $I^2 = 82\%$) and olfactory cues (SMD = -0.12 [95% CI = -0.53, 0.295]; $Z = 0.58$, $p = .56$, $I^2 = 54\%$) did not. However, the test for subgroup differences was not statistically significant ($X^2(2) = 2.69$, $p = .260$).

3.2. Exploratory analyses

3.2.1. Effect of inhibitory control task

We conducted exploratory analyses on the type of task used to operationalize inhibitory control (see Table 1). Following appetitive cue exposure, inhibitory control was impaired on the Stop Signal ($k = 18$; SMD = -0.15 [95% CI = -0.25, -0.05]; $Z = 3.00$, $p < .001$, $I^2 = 47\%$), Stroop ($k = 6$; SMD = -0.66 [95% CI = -1.28, -0.03]; $Z = 2.04$, $p = .040$, $I^2 = 92\%$), and Anti-saccade tasks ($k = 5$; SMD = -0.19 [95% CI = -0.33, -0.04]; $Z = 2.57$, $p = .010$, $I^2 = 0\%$), but there was no reliable effect on Go/No-Go tasks ($k = 16$; SMD = -0.15 [95% CI = -0.35, 0.05]; $Z = 1.46$, $p = .14$, $I^2 = 84\%$) or Go/No-Go shifting tasks ($k = 20$; SMD = 0.10 [95% CI = -0.15, 0.35]; $Z = 0.80$, $p = .430$, $I^2 = 89\%$). However, the subgroup effect was not statistically significant ($X^2(4) = 6.64$, $p = .160$).

Due to methodological considerations regarding the Go/No-Go shifting task (see Meule, 2017 and discussion) we repeated our primary analysis after excluding studies that used this task. In this case both alcohol-related cues ($k = 26$; SMD = -0.23 [95% CI = -0.35, -0.11]; $Z = 3.85$, $p < .001$; $I^2 = 73\%$) and food-related cues ($k = 20$; SMD = -0.19 [95% CI = -0.37, -0.01]; $Z = 2.03$, $p = .04$; $I^2 = 82\%$) impaired inhibitory control. We note that this was not an a-priori analysis. We present all analyses with exclusion of effect sizes from the Go/No-Go shifting task in online supplementary materials.

3.2.2. Examination of bias

Visual inspection of the funnel plot (see Supplementary Fig. 2) for all studies suggested asymmetry, and Trim and Fill analyses suggested 17 effect sizes would need to be added to achieve symmetry (see Supplementary Fig. 3). Adding these effect sizes made the overall point estimate non-significant (SMD = 0.05 [95% CI = -0.08, 0.18]). This suggests some degree of bias was evident across the effect sizes included. We also conducted Egger's test to formally examine asymmetry by regressing the effect size against the precision, however the test was not statistically significant ($Z = -0.56$, $p = .574$).

4. Discussion

The results of this meta-analytic investigation demonstrate that exposure to alcohol-related cues prompts robust, albeit small impairments in inhibitory control, although the evidence for comparable effects of food-related cues was not reliable. We observed substantial heterogeneity across effect sizes, which remained high despite several subgroup analyses that attempted to identify moderating variables. There was limited evidence to suggest that drinking status moderated the effect of alcohol-related cues, or that weight status moderated the effect of food-related cues on inhibitory control. Similarly, the modality of cue exposure did not significantly moderate our findings. Statistical correction for bias made the main effect of appetitive cues on inhibitory control no longer statistically significant.

Our primary hypothesis – exposure to appetitive cues would prompt a deficit in inhibitory control – was not fully supported. Overall, appetitive cues impaired inhibitory control, however this was driven by a

significant effect for alcohol-related cues. These findings provide partial support for theoretical models which suggest inhibitory control is a transient process that is sensitive to environmental and internal events (Jones, Christiansen, Nederkoorn, Houben, & Field, 2013a; Verbruggen et al., 2014), and the transient nature of inhibitory control cues may be one psychological mechanism that underlies the influence of alcohol cues on drinking behaviour (De Wit, 2009; Field & Jones, 2017). We found no evidence that transient impairments in inhibitory control were associated with individual differences in alcohol or food intake over the longer term, such as (self-reported) quantity of alcohol consumed per week, hazardous drinking scores or BMI. Future studies should attempt to clarify the associations between the disinhibiting effects of appetitive cues and food or alcohol intake that is measured immediately afterwards, such as *ad-libitum* intake (Field & Jones, 2017; Jones et al., 2015) or operant choice (Veling, Aarts, & Stroebe, 2013).

Consideration of publication bias suggests that even these small effects of appetitive cues may be inflated, and when we accounted for 'missing' (small and non-significant; $k = 17$) effect sizes using Trim and Fill analyses (Duval & Tweedie, 2000), the overall effect size was no longer statistically significant. This suggests that this literature is characterised by 'small study effects', often of poor methodological quality, and reporting biases that can substantially influence pooled-estimates (Schwarzer, Carpenter, & Rücker, 2015). Future research should conduct well powered studies and aim to publish all data to mitigate these biases and improve our confidence in pooled estimates of effect.

As hypothesised, impairments in inhibitory control following exposure to alcohol-related cues were comparable heavy drinkers and people with alcohol dependence, but these effects were absent in light drinkers. However, the test for subgroup differences was not statistically significant. Therefore, these subgroup differences are merely suggestive, and they should be interpreted with caution because many subgroups were poorly defined and created post-hoc using median split techniques (Jones & Field, 2015; Nederkoorn et al., 2009). Future studies might use established criteria to define heavy drinking or alcohol dependence (Babor, Higgins-Biddle, Saunders, & Monteiro, 2001; Edwards, 1996), in order to determine if these subgroup differences are indeed robust.

The absence of a robust effect of food-related cues on inhibitory control was surprising but is difficult to interpret given methodological features of the original studies. Eleven (35%) of studies used the Go/No-Go switching task for food-related cues. In this task, contingencies between making a motor response and inhibiting to high caloric vs. control cues are regularly switched on a block-by-block basis (Loeber et al., 2012). The repeated shifting of task contingencies between blocks mean that this task is likely to capture the effect of cues on inhibition and shifting, two distinct subcomponents of executive functions, and therefore this task provides an impure measure of inhibitory control (Miyake et al., 2010). Furthermore, variations on this task have also used low-caloric food cues (rather than non-food items) which may still be appetitive (He et al., 2014). Notably, when we removed effect sizes generated from Go/No-Go shifting tasks from our analysis the effect of food cues on inhibitory control was robust.

The modality in which appetitive cues were presented (e.g. visual vs. olfactory) did not moderate the effect of appetitive cues on inhibitory control. The absence of a moderation effect here should be interpreted cautiously until further direct comparisons across modalities are attempted (cf. Boswell & Kober, 2016; Monk et al., 2016). Overall, our moderator analyses suggest that we were unable to reduce the substantial heterogeneity by identifying variables that might moderate the influence of appetitive cues on inhibitory control. It is possible that other variables may influence this relationship, but we did not identify enough studies to examine this. For example, heterogeneity may have been caused by considerable variability in food-related and control images (see Table 1) and individual differences in reactivity to these cues, or differing levels of motivation to restrict unhealthy

behaviours across the samples. Future studies should investigate these potential moderators in more detail.

Finally, the effect of alcohol-related cues on inhibitory control supports the recent development of Inhibitory Control Training (ICT) as a behavioural intervention, to mitigate against cue-specific inhibition deficits. ICT creates an associative link between appetitive cues and inhibition of behaviour, which is thought to extinguish the associative link between appetitive cues and approach behaviour (Stice, Lawrence, Kemps, & Velting, 2016; Verbruggen et al., 2014). Promising effects of ICT have been demonstrated for both *ad-libitum* food and alcohol consumption in the laboratory (Allom, Mullan, & Hagger, 2015; Jones et al., 2016).

Some limitations of our analyses should be taken into consideration. First, we only included studies that examined cue reactivity to alcohol or food-related cues. There is evidence in other domains for cue-specific impairments of inhibitory control, including smoking (Luijten et al., 2011) and illicit drug use (Pike et al., 2013; Verdejo-García et al., 2012) which we did not integrate into our current analysis due to the limited number of available studies. Similarly, although we limited our findings to adult participants (18 + in United Kingdom, the legal drinking age) research also demonstrates inhibitory deficits following alcohol cues and food cues in younger people (Ames et al., 2014; Korucuoglu, Gladwin, & Wiers, 2015). Second, we were unable to directly measure associations between subjective and physiological cue-reactivity (e.g. craving and arousal) and inhibitory control as few studies measured this consistently. Finally, it is unknown whether these deficits in inhibitory control are a capacity deficit, or a motivational deficit (i.e. participants do not evoke effortful inhibition; Fujita, 2011). Future research should aim to overcome these limitations but also identify the mechanisms through which appetitive cues impair inhibitory control, for example through competition with attentional processes (Pessoa, Padmala, Kenner, & Bauer, 2012), reductions in limited self-regulatory resources (Muraven & Shmueli, 2006) or a reduction in their motivations (Pessoa, 2009). Furthermore, the influence of appetitive cues on inhibitory control in real-world settings (outside of the laboratory) should be investigated in order to elucidate the significance of these deficits for health-related behaviour. An interesting way to do this would be to examine real-time cue-exposure and inhibitory control using Ecological Momentary Assessment techniques (Shiffman, 2009). Given that cue-reactivity demonstrates substantial within- and between-subject variability (Serre, Fatseas, Swendsen, & Auriacombe, 2015) the real-world effects of repeated (LaRowe, Saladin, Carpenter, & Upadhyaya, 2007), cumulative and personalised cues (MacKillop et al., 2010) may identify more robust effects of exposure to appetitive cues on inhibitory control.

To conclude, the results from this meta-analytic investigation demonstrate that inhibitory control is sensitive to the presence of alcohol-related cues. The effect of food-related cues was less robust and may be confounded by methodological features of the tasks used. Overall, these findings provide some tentative support for theoretical predictions that inhibitory control is sensitive to exposure to appetitive cues. However whether these effects are robust, and if they play an important role in health-related behaviour, are important questions for future research.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.appet.2018.06.024>.

References

Contributed data to meta-analyses.

- (*)Adams, S., Ataya, A. F., Attwood, A. S., & Munafò, M. R. (2013). Effects of alcohol on disinhibition towards alcohol-related cues. *Drug and Alcohol Dependence*, 127, 137–142.
- Allom, V., Mullan, B., & Hagger, M. S. (2015). Does inhibitory control training improve health behaviour? A meta-analysis. *Health Psychology Review*, 10, 168–186.
- Ames, S. L., Kisbu-Sakarya, Y., Reynolds, K. D., Boyle, S., Cappelli, C., Cox, M. G., et al.

- (2014). Inhibitory control effects in adolescent binge eating and consumption of sugar-sweetened beverages and snacks. *Appetite*, 81, 180–192.
- Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G. (2001). *The alcohol use disorders identification test: Guidelines for use in primary care*. World Health Organisation.
- Baumeister, R. F. (2014). Self-regulation, ego depletion, and inhibition. *Neuropsychologia*, 65, 313–319.
- Bickel, W. K., Jarmolowicz, D. P., Mueller, E. T., Gatchalian, K. M., & McClure, S. M. (2012). Are executive function and impulsivity antipodes? A conceptual reconstruction with special reference to addiction. *Psychopharmacology*, 221, 361–387.
- Boswell, R. G., & Kober, H. (2016). Food cue reactivity and craving predict eating and weight gain: A meta-analytic review. *Obesity Reviews*, 17, 159–177.
- Brockmeyer, T., Hahn, C., Reetz, C., Schmidt, U., & Friederich, H.-C. (2015). Approach bias and cue reactivity towards food in people with high versus low levels of food craving. *Appetite*, 95, 197–202.
- Christiansen, P., Cole, J. C., Goudie, A. J., & Field, M. (2012). Components of behavioural impulsivity and automatic cue approach predict unique variance in hazardous drinking. *Psychopharmacology*, 219, 501–510.
- (*)Czapla, M., Simon, J. J., Friederich, H. C., Herpertz, S. C., Zimmermann, P., & Loeber, S. (2015b). Is binge drinking in young adults associated with an alcohol-specific impairment of response inhibition? *European Addiction Research*, 21, 105–113.
- (*)Czapla, M., Simon, J. J., Richter, B., Kluge, M., Friederich, H. C., Herpertz, S., et al. (2015a). The impact of cognitive impairment and impulsivity on relapse of alcohol-dependent patients: Implications for psychotherapeutic treatment. *Addiction Biology*, 21, 873–884.
- De Wit, H. (2009). Impulsivity as a determinant and consequence of drug use: A review of underlying processes. *Addiction Biology*, 14, 22–31.
- Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135–168.
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455–463.
- Eagle, D. M., Bari, A., & Robbins, T. W. (2008). The neuropsychopharmacology of action inhibition: Cross-species translation of the stop-signal and go/no-go tasks. *Psychopharmacology*, 199, 439–456.
- Edwards, G. (1996). Sensible drinking - doctors should stick with the independent medical advice. *British Medical Journal*, 312, 1.
- Elbourne, D. R., Altman, D. G., Higgins, J. P. T., Curtin, F., Worthington, H. V., & Vail, A. (2002). Meta-analyses involving cross-over trials: Methodological issues. *International Journal of Epidemiology*, 31, 140–149.
- Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & Psychophysics*, 16, 143–149.
- Fernie, G., Peeters, P., Gullo, M. J., Christiansen, P., Cole, J., Sunmull, H., et al. (2013). Multiple components of impulsivity predict prospective alcohol involvement in adolescents. *Addiction*, 108, 1916–1923.
- (*)Field, M., & Jones, A. (2017). Elevated alcohol consumption following alcohol cue exposure is partially mediated by reduced inhibitory control and increased craving. *Psychopharmacology (Berl)*.
- Field, M., Kiernan, A., Eastwood, B., & Child, R. (2008). Rapid approach responses to alcohol cues in heavy drinkers. *Journal of Behavior Therapy and Experimental Psychiatry*, 39, 209–218.
- Fujita, K. (2011). On conceptualizing self-control as more than the effortful inhibition of impulses. *Personality and Social Psychology Review*, 15, 352–366.
- (*)Guggel, S., Heusinger, A., Forkmann, T., Boecker, M., Lindenmeyer, J., Miles Cox, W., et al. (2010). Effects of alcohol cue exposure on response inhibition in detoxified alcohol-dependent patients. *Alcoholism: Clinical and Experimental Research*, 34, 1584–1589.
- Goldstein, R. Z., & Volkow, N. D. (2011). Dysfunction of the prefrontal cortex in addiction: Neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*, 12, 652–669.
- Hallett, P. E. (1978). Primary and secondary saccades to goals defined by instructions. *Vision Research*, 18, 1279–1296.
- (*)He, Q., Xiao, L., Xue, G., Wong, S., Ames, S. L., Schembre, S. M., et al. (2014). Poor ability to resist tempting calorie rich food is linked to altered balance between neural systems involved in urge and self-control. *Nutrition Journal*, 13, 92.
- Higgins, J. P. T., & Green, S. (Eds.). (2011). *Cochrane handbook for systematic reviews of interventions*. The Cochrane Collaboration.
- (*)Houben, K., Nederkoorn, C., & Jansen, A. (2013). Eating on impulse: The relation between overweight and food-specific inhibitory control. *Obesity*.
- (*)Hume, D. J., Howells, F. M., Rauch, H. G., Kroff, J., & Lambert, E. V. (2015). Electrophysiological indices of visual food cue-reactivity: Differences in obese, overweight and normal weight women. *Appetite*, 85, 126–137.
- Jelicic Kadic, A., Vucic, K., Dosenovic, S., Sapunar, D., & Puljak, L. (2016). Extracting data from figures with software was faster, with higher interrater reliability than manual extraction. *Journal of Clinical Epidemiology*, 74, 119–123.
- Jones, A., Button, E., Rose, A. K., Robinson, E., Christiansen, P., Di Lemma, L., et al. (2015). The ad-libitum alcohol 'taste test': Secondary analyses of potential confounds and construct validity. *Psychopharmacology*, 1–8.
- Jones, A., Christiansen, P., Nederkoorn, C., Houben, K., & Field, M. (2013a). Fluctuating Disinhibition: Implications for the understanding and treatment of alcohol and other substance use disorders. *Frontiers in Psychiatry*, 22.
- Jones, A., Di Lemma, L. C. G., Robinson, E., Christiansen, P., Nolan, S., Tudur-Smith, C., et al. (2016). Inhibitory control training for appetitive behaviour change: A meta-analytic investigation of mechanisms of action and moderators of effectiveness. *Appetite*, 97, 16–28.
- (*)Jones, A., & Field, M. (2015). Alcohol-related and negatively-valenced cues increase motor and oculomotor disinhibition in social drinkers. *Experimental and Clinical Psychopharmacology*, 23, 122–129.
- Jones, A., Hardman, C. A., Lawrence, N., & Field, M. (2017). Cognitive training as a potential treatment for overweight and obesity: A critical review of the evidence. *Appetite*.
- (*)Jones, A., Rose, A. K., Cole, J., & Field, M. (2013b). Effects of alcohol cues on craving

- and ad libitum alcohol consumption in social drinkers: The role of disinhibition. *Journal of Experimental Psychopathology*, 4, 239–249.
- Kemps, E., & Tiggemann, M. (2015). Approach bias for food cues in obese individuals. *Psychology and Health*, 30, 370–380.
- Kemps, E., Tiggemann, M., Martin, R., & Elliott, M. (2013). Implicit approach-avoidance associations for craved food cues. *Journal of Experimental Psychology: Applied*, 19, 30–38.
- Korucuoglu, O., Gladwin, T. E., & Wiers, R. W. (2015). Alcohol-induced changes in conflict monitoring and error detection as predictors of alcohol use in late adolescence. *Neuropsychopharmacology*, 40, 614–621.
- (*)Kreusch, F., Billieux, J., & Quertemont, E. (2017). Alcohol-cue exposure decreases response inhibition towards alcohol-related stimuli in detoxified alcohol-dependent patients. *Psychiatry Research*, 249, 232–239.
- (*)Kreusch, F., Vilenne, A., & Quertemont, E. (2013). Response inhibition toward alcohol-related cues using an alcohol go/no-go task in problem and non-problem drinkers. *Addictive Behaviors*, 38, 2520–2528.
- LaRowe, S. D., Saladin, M. E., Carpenter, M. J., & Upadhyaya, H. P. (2007). Reactivity to nicotine cues over repeated cue reactivity sessions. *Addictive Behaviors*, 32, 2888–2899.
- (*)Lattimore, P., & Mead, B. R. (2015). See it, grab it, or STOP! Relationships between trait impulsivity, attentional bias for pictorial food cues and associated response inhibition following in-vivo food cue exposure. *Appetite*, 90, 248–253.
- Loeber, S., Grosshans, M., Herpertz, S., Kiefer, F., & Herpertz, S. C. (2013). Hunger modulates behavioral disinhibition and attention allocation to food-associated cues in normal-weight controls. *Appetite*, 71, 32–39.
- (*)Loeber, S., Grosshans, M., Korucuoglu, O., Vollmert, C., Vollstädt-Klein, S., Schneider, S., et al. (2012). Impairment of inhibitory control in response to food-associated cues and attentional bias of obese participants and normal-weight controls. *International Journal of Obesity* (2005), 36, 1334–1339.
- (*)Logan, G. D., Cowan, W. B., & Davis, K. A. (1984). On the ability to inhibit simple and choice reaction time responses: A model and a method. *Journal of Experimental Psychology: Human Perception and Performance*, 10, 276–291.
- Logan, G. D., Schachar, R. J., & Tannock, R. (1997). Impulsivity and inhibitory control. *Psychological Science*, 8, 60–64.
- Luijten, M., Littell, M., & Franken, I. H. A. (2011). Deficits in inhibitory control in smokers during a Go/Nogo task: An investigation using event-related brain potentials. *PLoS One*, 6, 6.
- (*)Lyu, Z., Zheng, P., Chen, H., & Jackson, T. (2017). Approach and inhibition responses to external food cues among average-weight women who binge eat and weight-matched controls. *Appetite*, 108, 367–374.
- MacKillop, J., O'Hagen, S., Lisman, S. A., Murphy, J. G., Ray, L. A., Tidey, J. W., et al. (2010). Behavioral economic analysis of cue-elicited craving for alcohol. *Addiction*, 105, 1599–1607.
- (*)Mainz, V., Dürke, B., Boecker, M., Kessel, R., Gauggel, S., & Forkmann, T. (2012). Influence of cue exposure on inhibitory control and brain activation in patients with alcohol dependence. *Frontiers in Human Neuroscience*.
- (*)Manasse, S. M., Goldstein, S. P., Wyckoff, E., Forman, E. M., Juarascio, A. S., Butryn, M. L., et al. (2016). Slowing down and taking a second look: Inhibitory deficits associated with binge eating are not food-specific. *Appetite*, 96, 555–559.
- Meule, A. (2017). Reporting and interpreting task performance in go/No-go affective shifting tasks. *Frontiers in Psychology*, 8, 701.
- (*)Meule, A., & Kübler, A. (2014). Double trouble. Trait food craving and impulsivity interactively predict food-cue affected behavioral inhibition. *Appetite*, 79, 174–182.
- (*)Meule, A., Lutz, A. P. C., Krawietz, V., Stützer, J., Vögele, C., & Kübler, A. (2014a). Food-cue affected motor response inhibition and self-reported dieting success: A pictorial affective shifting task. *Frontiers in Psychology*, 5.
- (*)Meule, A., Lutz, A., Vögele, C., & Kübler, A. (2012). Women with elevated food addiction symptoms show accelerated reactions, but no impaired inhibitory control, in response to pictures of high-calorie food-cues. *Eating Behaviors*, 13, 423–428.
- Meule, A., Lukito, S., Vögele, C., & Kübler, A. (2011). Enhanced behavioral inhibition in restrained eaters. *Eating Behaviour*, 12, 152–155.
- (*)Meule, A., Lutz, A. P. C., Vögele, C., & Kübler, A. (2014b). Impulsive reactions to food-cues predict subsequent food craving. *Eating Behaviors*, 15, 99–105.
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: A latent variable analysis. *Cognitive Psychology*, 41, 49–100.
- (*)Monk, R. L., Sunley, J., Qureshi, A. W., & Heim, D. (2016). Smells like inhibition: The effects of olfactory and visual alcohol cues on inhibitory control. *Psychopharmacology*, 233, 1331–1337.
- (*)Muraven, M., & Shmueli, D. (2006). The self-control costs of fighting the temptation to drink. *Psychology of Addictive Behaviors*, 20, 154–160.
- (*)Nederkoorn, C., Baltus, M., Guerrieri, R., & Wiers, R. W. (2009). Heavy drinking is associated with deficient response inhibition in women but not in men. *Pharmacology Biochemistry and Behavior*, 93, 331–336.
- Nederkoorn, C., Houben, K., Hofmann, W., Roefs, A., & Jansen, A. (2010). Control yourself or just eat what you like? weight gain over a year is predicted by an interactive effect of response inhibition and implicit preference for snack foods. *Health Psychology*, 29, 389–393.
- Nederkoorn, C., Smulders, F. T. Y., Havermans, R. C., Roefs, A., & Jansen, A. (2006). Impulsivity in obese women. *Appetite*, 47, 253–256.
- Nikolaou, K., Field, M., & Duka, T. (2013). Alcohol-related cues reduce cognitive control in social drinkers. *Behavioural Pharmacology*, 24, 29–36.
- (*)Noel, X., Van Der Linden, M., D'Acremont, M., Bechara, A., Dan, B., Hanak, C., et al. (2007). Alcohol cues increase cognitive impulsivity in individuals with alcoholism. *Psychopharmacology*, 192, 291–298.
- (*)Noel, X., Van Der Linden, M., D'Acremont, M., Colmant, M., Hanak, C., Pelc, I., et al. (2005). Cognitive biases toward alcohol-related words and executive deficits in polysubstance abusers with alcoholism. *Addiction*, 100, 1302–1309.
- Pessoa, L., Padmala, S., Kenner, A., & Bauer, A. (2012). Interactions between cognition and emotion during response inhibition. *Emotion*, 12, 192–197.
- (*)Petit, G., Cimochovska, A., Kornreich, C., Hanak, C., Verbanck, P., & Campanella, S. (2014). Neurophysiological correlates of response inhibition predict relapse in detoxified alcoholic patients: Some preliminary evidence from event-related potentials. *Neuropsychiatric Disease and Treatment*, 10, 1025–1037.
- (*)Petit, G., Kornreich, C., Noël, X., Verbanck, P., & Campanella, S. (2012). Alcohol-related context modulates performance of social drinkers in a visual go/no-go task: A preliminary assessment of event-related potentials. *PLoS One*, 7, e37466.
- (*)Phelan, S., Hassenstab, J., McCaffery, J. M., Sweet, L., Raynor, H. A., Cohen, R. A., et al. (2011). Cognitive interference from food cues in weight loss maintainers, normal weight, and obese individuals. *Obesity*, 19, 69–73.
- Pike, E., Stoops, W. W., Fillmore, M. T., & Rush, C. R. (2013). Drug-related stimuli impair inhibitory control in cocaine abusers. *Drug and Alcohol Dependence*, 133, 768–771.
- Price, M., Lee, M., & Higgins, S. (2016). Food-specific response inhibition, dietary restraint and snack intake in lean and overweight/obese adults: A moderated-mediation model. *International Journal of Obesity*, 40, 877–882.
- Riley, R. D., Higgins, J. P. T., & Deeks, J. J. (2011). Interpretation of random effects meta-analyses. *BMJ*, 342, 964–967.
- Rubio, G., Jimenez, M., Rodriguez-Jimenez, R., Martinez, I., Avila, C., Ferre, F., et al. (2008). The role of behavioral impulsivity in the development of alcohol dependence: A 4-year follow-up study. *Alcoholism: Clinical and Experimental Research*, 32, 1681–1687.
- Saunders, J. B., Aasland, O. G., Babor, T. F., De La Fuente, J. R., & Grant, M. (1993). Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-ii. *Addiction*, 88, 791–804.
- Schag, K., Teufel, M., Junne, F., Preissl, H., Hautzinger, M., Zipfel, S., et al. (2013). Impulsivity in binge eating disorder: Food cues elicit increased reward responses and disinhibition. *PLoS ONE*, 8.
- Schwarzer, G., Carpenter, J. R., & Rücker, G. (2015). Small-study effects in meta-analysis. In G. Schwarzer, J. R. Carpenter, & G. Rücker (Eds.). *Meta-analysis with R* (pp. 107–141). Cham: Springer International Publishing.
- Serre, F., Fatseas, M., Swendsen, J., & Auriacombe, M. (2015). Ecological momentary assessment in the investigation of craving and substance use in daily life: A systematic review. *Drug and Alcohol Dependence*, 148, 1–20.
- Shiffman, S. (2009). Ecological momentary assessment (EMA) in studies of substance use. *Psychological Assessment*, 21, 486–497.
- (*)Sion, A., Jurado-Barba, R., Alonso, M. J., & Rubio-Valladolid, G. (2017). Inhibitory capacity assessment in alcohol dependent patients: Translation from a modified stop signal task. *Actas Españolas de Psiquiatría*, 45, 21–31.
- Smith, J., Mattick, R., Jamadar, S., & Iredale, J. (2014). Deficits in behavioural inhibition in substance abuse and addiction: A meta-analysis. *Drug and Alcohol Dependence*, 145, 1–33.
- Stice, E., Lawrence, N. S., Kemps, E., & Veling, H. (2016). Training motor responses to food: A novel treatment for obesity targeting implicit processes. *Clinical Psychology Review*, 49, 16–27.
- Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, 18, 643–662.
- Tarter, R. E., Kirisci, L., Reynolds, M., & Mezzich, A. (2004). Neurobehavior disinhibition in childhood predicts suicide potential and substance use disorder by young adulthood. *Drug and Alcohol Dependence*, 76.
- Vainik, U., Dagher, A., Dubé, L., & Fellows, L. K. (2013). Neurobehavioural correlates of body mass index and eating behaviours in adults: A systematic review. *Neuroscience & Biobehavioral Reviews*, 37, 279–299.
- Veling, H., Aarts, H., & Stroebe, W. (2013). Using stop signals to reduce impulsive choices for palatable unhealthy foods. *British Journal of Health Psychology*, 18, 354–368.
- Verbruggen, F., Best, M., Bowditch, W. A., Stevens, T., & McLaren, I. P. L. (2014). The inhibitory control reflex. *Neuropsychologia*, 65, 263–278.
- Verbruggen, F., & Logan, G. D. (2008). Automatic and controlled response inhibition: Associative learning in the go/No-go and stop-signal paradigms. *Journal of Experimental Psychology: General*, 137, 649–672.
- Verdejo-García, A., Lubman, D. I., Schwert, A., Roffel, K., Vilar-López, R., MacKenzie, T., et al. (2012). Effect of craving induction on inhibitory control in opiate dependence. *Psychopharmacology*, 219, 519–526.
- Volkow, N. D., Wang, G. J., Tomasi, D., & Baler, R. D. (2013). Obesity and addiction: Neurobiological overlaps. *Obesity Reviews*, 14, 2–18.
- Vucic, K., Jelacic Kadic, A., & Puljak, L. (2015). Survey of Cochrane protocols found methods for data extraction from figures not mentioned or unclear. *Journal of Clinical Epidemiology*, 68, 1161–1164.
- (*)Weafer, J., & Fillmore, M. T. (2012). Alcohol-related stimuli reduce inhibitory control of behavior in drinkers. *Psychopharmacology*, 222, 489–498.
- (*)Weafer, J., & Fillmore, M. T. (2014). Alcohol-related cues potentiate alcohol impairment of behavioral control in drinkers. *Psychology of Addictive Behaviors*, 29, 290–299.
- Wiers, R. W., Gladwin, T. E., Hofmann, W., Salemink, E., & Ridderinkhof, K. R. (2013). Cognitive bias modification and cognitive control training in addiction and related psychopathology: Mechanisms, clinical perspectives, and ways forward. *Clinical Psychological Science*, 1, 192–212.
- Wiers, R. W., Bartholow, B. D., van den Wildenberg, E., Thush, C., Engels, R. C. M. E., Sher, K. J., et al. (2007). Automatic and controlled processes and the development of addictive behaviors in adolescents: A review and a model. *Pharmacology Biochemistry and Behavior*, 86, 263–283.
- (*)Yeomans, M. R., & Brace, A. (2015). Cued to act on impulse: More impulsive choice and risky decision making by women susceptible to overeating after exposure to food stimuli. *PLoS One*, 10, e0137626.