

## LJMU Research Online

Rooijakkers, MJP, Stens, NA, Van Wely, MH, Van Der Wulp, K, Rodwell, L, Gehlmann, H, Van Garsse, LAFM, Geuzebroek, GSC, Verkroost, MWA, Habets, J, El Messaoudi, S, Thijssen, DHJ, Nijveldt, R and Van Royen, N

Diastolic delta best predicts paravalvular regurgitation after transcatheter aortic valve replacement as assessed by cardiac magnetic resonance: the APPOSE trial

https://researchonline.ljmu.ac.uk/id/eprint/22995/

Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Rooijakkers, MJP ORCID logoORCID: https://orcid.org/0000-0002-6815-7667, Stens, NA, Van Wely, MH, Van Der Wulp, K, Rodwell, L, Gehlmann, H, Van Garsse, LAFM, Geuzebroek, GSC ORCID logoORCID: https://orcid.org/0000-0003-0407-2901. Verkroost. MWA. Habets. J. El Messaoudi. S. Thiissen. DHJ

LJMU has developed LJMU Research Online for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact <a href="mailto:researchonline@ljmu.ac.uk">researchonline@ljmu.ac.uk</a>

http://researchonline.ljmu.ac.uk/



## Diastolic delta best predicts paravalvular regurgitation after transcatheter aortic valve replacement as assessed by cardiac magnetic resonance: the APPOSE trial

Maxim J.P. Rooijakkers (1)<sup>1</sup>, Niels A. Stens<sup>1,2</sup>, Marleen H. van Wely<sup>1</sup>, Kees van der Wulp<sup>1</sup>, Laura Rodwell<sup>3</sup>, Helmut Gehlmann<sup>1</sup>, Leen A.F.M. van Garsse<sup>4</sup>, Guillaume S.C. Geuzebroek (1)<sup>4</sup>, Michel W.A. Verkroost<sup>4</sup>, Jesse Habets<sup>5</sup>, Saloua El Messaoudi<sup>1</sup>, Dick H.J. Thijssen<sup>2,6</sup>, Robin Nijveldt<sup>1</sup>, and Niels van Royen<sup>1</sup>\*

<sup>1</sup>Department of Cardiology, Radboud University Medical Centre, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, The Netherlands; <sup>2</sup>Department of Physiology, Radboud University Medical Centre, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, The Netherlands; <sup>3</sup>Department of Health Sciences, Section Biostatistics, Radboud Institute for Health Sciences, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, The Netherlands; <sup>4</sup>Department of Cardiothoracic Surgery, Radboud University Medical Centre, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, The Netherlands; <sup>5</sup>Department of Medical Imaging, Radboud University Medical Centre, Geert Grooteplein Zuid 10, 6525 GA, The Netherlands; and <sup>6</sup>Research Institute for Sports and Exercise Sciences, Liverpool John Moores University, Liverpool L3 5UX, UK

Received 31 October 2022; revised 19 January 2023; accepted 4 February 2023; online publish-ahead-of-print 14 March 2023

Aims	Paravalvular regurgitation (PVR) is a common complication after transcatheter aortic valve replacement (TAVR) that poses an increased risk of rehospitalization for heart failure and mortality. The aim of this study was to assess the accuracy of haemodynamic indices to predict relevant PVR.
Methods and results	In this prospective single-centre clinical trial, four haemodynamic indices of PVR measured during TAVR were assessed for their correlation with gold standard cardiac magnetic resonance (CMR)-derived regurgitant fraction (CMR-RF) at 1 month follow-up: diastolic delta (DD), heart rate-adjusted diastolic delta (HR-DD), aortic regurgitation index (ARI), and aortic regurgitation index ratio (ARI ratio). These haemodynamic indices were analysed for their ability to predict relevant PVR (defined as CMR-RF > 20%) using receiver operating characteristic (ROC) curves with corresponding area under the ROC curves (AUCs). A total of 77 patients were included and had CMR performed 41 $\pm$ 14 days after TAVR. Mean CMR-RF was 12.4 $\pm$ 9.3%. Fifteen (19.5%) patients had CMR-RF > 20%. DD had the best correlation with CMR-RF and the highest AUC to predict relevant PVR (0.82; 95% CI, 0.72–0.92), followed by HR-DD (AUC 0.78; 95% CI, 0.67–0.89), ARI (AUC 0.78; 95% CI, 0.66–0.89), and ARI ratio (AUC 0.65; 95% CI, 0.49–0.81). The optimal cut-off value for DD was 32 mmHg, with sensitivity of 69% and specificity of 77% in predicting relevant PVR.
Conclusion	DD measured during TAVR best predicts relevant PVR. Correction for heart rate (HR-DD) or systolic blood pressure (ARI, ARI ratio) did not improve this predictive value.

\* Corresponding author: Tel: +31 24 361 11 11, Fax: +31 24 354 05 76, E-mail: niels.vanroyen@radboudumc.nl

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (https://creativecommons.org/licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

#### **Graphical Abstract**

## Diastolic delta best predicts paravalvular regurgitation after TAVR



Schematic overview of the procedures and main results of the study. The blue and red asterisks denote pigtail catheters measuring pressure in the aorta and left ventricle, respectively. Haemodynamic indices were derived from these pressure measurements. AUC, area under the curve; CMR, cardiac magnetic resonance; TAVR, transcatheter aortic valve replacement; TTE, transthoracic echocardiography.

**Keywords** 

aortic stenosis • cardiac magnetic resonance • outcomes • paravalvular regurgitation • transcatheter aortic valve replacement

## Introduction

Transcatheter aortic valve replacement (TAVR) is a well-established minimally invasive treatment option in patients with severe symptomatic aortic stenosis (AS). Initially established for inoperable patients at high or prohibitive surgical risk, TAVR is nowadays also indicated in intermediate-risk patients as an alternative for surgical aortic valve replacement (SAVR). With the non-inferiority or even superiority of TAVR over SAVR in low-risk patients, the indications for TAVR are continuously expanding.<sup>1–3</sup>

One potential shortcoming of TAVR though is paravalvular regurgitation (PVR). The occurrence of PVR is common after TAVR, with incidences of mild PVR up to 40%, and incidences of greater than or equal to moderate PVR up to 10% in contemporary TAVR studies.<sup>4,5</sup> In patients with greater than or equal to moderate PVR, mortality is three times higher compared to patients with none to trace PVR.<sup>6,7</sup> Recently, also mild-moderate PVR was shown to be associated with an increased risk of mortality at 5 years after TAVR.<sup>8</sup> Therefore, it is important to grade PVR during the procedure, both for prognostic and therapeutic purposes. Identification of relevant PVR during the TAVR procedure could guide additional interventions (e.g. post-dilation) to reduce PVR and improve patient outcome.

Adoption of the 'minimalist TAVR' approach has precluded the use of peri-procedural transoesophageal echocardiography (TEE) in procedural detection of PVR. Transthoracic echocardiography (TTE) has several limitations in the assessment of PVR, potentially underestimating the degree of PVR when compared with gold standard cardiac magnetic resonance (CMR).<sup>9,10</sup> Angiographic grading is well feasible during the procedure but lacks diagnostic accuracy.<sup>11</sup> Alternatively, haemodynamic indices to estimate PVR can be derived from transvalvular pressure tracings post-TAVR. Such transvalvular pressure tracings are used to assess remaining pressure gradients over the aortic valve prosthesis.<sup>12</sup> Hence, these pressure tracings do not require additional instrumentation and do not involve additional contrast or radiation exposure. Commonly used haemodynamic indices to estimate PVR are diastolic delta (DD), heart rate-adjusted diastolic delta (HR-DD), aortic regurgitation index (ARI), and aortic regurgitation index ratio (ARI ratio). The latter three are in fact derivatives of DD, which is computed as the difference between end-diastolic blood pressure and left ventricular end-diastolic pressure (LVEDP). All four haemodynamic indices are correlated to mortality after TAVR.<sup>13–16</sup> However, the predictive value of these indices for relevant PVR, in which gold standard CMR is used as the reference modality, has never been determined.

Therefore, the aim of the Assessment of Paravalvular Regurgitation After Transcatheter Aortic Valve Replacement by Hemodynamic Measurements and Cardiac Magnetic Resonance (APPOSE) trial was to assess the predictive value of peri-procedural haemodynamic indices for the occurrence of relevant PVR at 1 month follow-up as quantified by CMR.

## Methods

#### **Population and design**

In this prospective single-centre study, we included consecutive patients who underwent a transfemoral or transaxillary TAVR for severe symptomatic AS at the Radboud University Medical Centre. In all patients, a selfexpanding valve (Portico; Abbott Structural Heart, Minneapolis, MN, USA) was implanted, with valve sizes ranging between 23 and 29 mm. The main exclusion criteria were the presence of a pre-existing prosthetic cardiac device or valve, left ventricular ejection fraction (LVEF) <30%, and a serum creatinine >250 µmol/L or end-stage renal disease. The complete list of inclusion and exclusion criteria is provided in Supplementary data online, Table S1. The study protocol was approved by the local Medical Research Ethics Committee and by the institutional review board of the Radboud University Medical Centre. Written informed consent was obtained from all patients prior to inclusion. The APPOSE trial is registered in the ClinicalTrials.gov database (NCT04281771). This trial was funded by a research grant from Abbott. The sponsor of the trial had no role in the design and conduct of the trial; in the enrolment of participants; in the collection, analysis, and interpretation of the data; in the preparation, review, or approval of the manuscript; or in the decision to submit the manuscript for publication.

## **TAVR** procedure

All patients were discussed in a multi-disciplinary Heart Team and were deemed eligible for a TAVR procedure. TAVR was performed according to routine protocol. Procedures were performed in a hybrid catheterization laboratory with a standard operating team consisting of an interventional cardiologist, cardiothoracic surgeon, and anaesthesiologist. TAVR was performed either under general anaesthesia or under conscious sedation, which was left to the discretion of the operating team.

#### Haemodynamic measurements

We used 5F fluid-filled pigtail catheters (Impulse; Boston Scientific, Marlborough, MA, USA) to simultaneously measure continuous pressure in the aorta and the left ventricle. Heart rate was derived from continuous electrocardiographic monitoring. Haemodynamic data were displayed and stored in Mac-Lab (GE Healthcare, Chicago, IL, USA).

Haemodynamic measurements were performed at baseline (i.e. before pre-dilation), directly after implantation of the bioprosthetic valve, and (when applicable) after post-dilation. Data were captured and stored using Castor (Castor EDC, Amsterdam, the Netherlands) and exported for analysis to SPSS Statistics (version 25.0.0.1, IBM Corporation, Armonk, NY,

USA). Haemodynamic data at baseline and after final TAVR result were used for the calculation of haemodynamic indices. For each haemodynamic index, the averaged pressures of at least three representative cardiac cycles during sinus rhythm (or paced rhythm) were used or at least five representative cycles during atrial fibrillation.

Haemodynamic indices were computed using the following calculations:

- DD: end-diastolic blood pressure LVEDP.
- HR-DD: (DD/heart rate) \* 80.
- ARI: (DD/systolic blood pressure) \* 100.
- ARI ratio: ARI after implantation/ARI prior to implantation.

### Echocardiographic assessment of PVR

TTE to determine the degree of PVR was performed 4–6 weeks after the TAVR procedure. Echocardiographic grading of PVR was based on an integrative multi-parametric approach that mainly included visual assessment of the number of PVR jets, jet width at the origin, and the circumferential extent of PVR. The degree of PVR was classified into none/trace, mild, moderate, or severe, according to Valve Academic Research Consortium-3 (VARC-3) criteria.<sup>17</sup> Two researchers (M.R. and S.E.M.) independently assessed the echocardiographic degree of PVR. If consensus was not reached, a third researcher got involved (N.V.R.).

### **CMR** measurements

All patients were scanned on the same day as the TTE assessment (4-6 weeks after TAVR) on a commercially available 1.5 T CMR scanner (Siemens Avanto; Siemens, Erlangen, Germany). Using a 2D phase-contrast velocity encoded spoiled gradient echo pulse sequence, the slice for the through-plane velocity quantification was placed perpendicular to the direction of the flow, just above the struts of the TAVR bioprosthesis. Flow acquisitions were performed during successive end-expiratory breath holds, using both a high velocity encoding (Venc) of at least 180 cm/s (or higher, depending on the velocity at which no aliasing was observed) and a low Venc of 75 cm/s. One cardiac cycle consisted of 25 phases. The high Venc was used for accurate assessment of the forward volume, the low Venc for determination of the regurgitant volume. CMR-derived regurgitant fraction (CMR-RF) was measured by dividing the regurgitant volume by the forward volume, multiplied by 100. The software package Medis Suite MR (Medis Medical Imaging, Leiden, the Netherlands) was used to analyse the CMR-RF.

#### Endpoints

The primary endpoint was the CMR-RF. Relevant PVR was defined as CMR-RF > 20%, whereas non-relevant PVR was defined as CMR-RF  $\leq$  20%. <sup>18</sup> Secondary endpoints included procedural haemodynamic indices of PVR (DD, HR-DD, ARI, and ARI ratio) and short-term outcomes according to VARC-3 criteria. <sup>17</sup>

## Statistical analysis

Haemodynamic indices with cut-offs previously associated with PVR were expected to be more prevalent in patients with greater than or equal to moderate PVR compared to patients with less than or equal to mild PVR. Based on previous literature, the prevalence of greater than or equal to moderate PVR was expected to be 30% in patients with levels of ARI above the median.<sup>19</sup> Based on the information above, using  $\chi^2$  test, a *P* value of 0.0125, and power of 80%, a total of 76 patients with analysable CMR scans and paired haemodynamic measurements were required. The sample size is comparable to previous studies in which CMR was compared with echocardiography in 71 patients<sup>20</sup> and to aortic root angiography in 69 patients.<sup>11</sup>

Data are presented as mean  $\pm$  standard deviation, median (interquartile range), or number (percentage), as appropriate. Baseline characteristics are displayed according to CMR-RF (CMR-RF > 20% vs. CMR-RF  $\leq$  20%). Categorical variables were compared using the  $\chi^2$  test. Continuous variables



Figure 1 Study flowchart. CMR, cardiac magnetic resonance; SAVR, surgical aortic valve replacement; TAVR, transcatheter aortic valve replacement.

were compared using Student's *t*-test or Mann–Whitney *U* test, depending on normal distribution.

Correlation analysis was performed between the haemodynamic indices (DD, HR-DD, ARI, and ARI ratio) and CMR-RF, using Pearson's correlation coefficient. Receiver operating characteristic (ROC) analyses were performed to assess the predictive value of the haemodynamic indices for CMR-RF > 20%. To allow ROC analysis, the continuous CMR-RF was dichotomized to CMR-RF > 20% and CMR-RF  $\leq 20\%$ .<sup>18</sup> The area under the ROC curve (AUC) was used to quantify diagnostic performance of the haemodynamic indices, which were subsequently compared using DeLong's test for correlated AUCs.<sup>21</sup> Optimal cut-offs for all indices were defined as the cut-off value of the index at which the sensitivity (SE) and specificity (SP) curves intersect. When a higher SP was possible with the SE corresponding to the point of intersection, or vice versa, that cut-off value was chosen. Analyses were performed in SPSS Statistics (version 25.0.0.1, IBM Corporation, Armonk, NY, USA) and R (version 4.1.2, R Foundation for Statistical Computing, Vienna, Austria).

## Results

## **Baseline characteristics**

Between November 2019 and October 2021, 103 consecutive patients who were accepted for TAVR provided written informed consent for the study. After exclusion of patients denoted as screen failure (n = 11) and patients without a complete CMR evaluation (n = 15), we examined a total of 77 patients (*Figure 1*). Baseline characteristics for patients with either CMR-RF > 20% or CMR-RF  $\leq$  20% are shown in *Table 1*. Mean age was 80.4 ± 5.1 years and 46.8% of patients were men. Median New York Heart Association (NYHA) function class was II, with 39.0% of patients being in NYHA class III or IV. Mean Society of Thoracic Surgeons (STS) and European System for Cardiac Operative Risk Evaluation (EuroSCORE) II were 2.38 ± 0.96 and 2.53 ± 1.63, respectively. Mean LVEF was 54.2 ± 8.4%, with a mean aortic valve area (AVA) and aortic valve mean gradient of 0.79 ± 0.18 cm<sup>2</sup>

#### M.J.P. Rooijakkers et al.

	Study population (n = 77)	CMR-RF ≤ 20% ( <i>n</i> = 62)	CMR-RF > 20% ( <i>n</i> = 15)	P value
Demographics	·····	· · · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · · ·	
Age, years	80.4 ± 5.1	80.5 ± 5.0	80.4 ± 5.9	0.972
Male sex, n (%)	36 (46.8)	27 (43.5)	9 (60)	0.252
Body mass index (BMI), kg/m <sup>2</sup>	27.3 ± 4.0	27.1 ± 4.0	28.1 ± 4.1	0.402
Obesity, n (%)	17 (22.1)	12 (19.4)	5 (33.3)	0.300
Smoker, n (%)	5 (6.5)	3 (4.8)	2 (13.3)	0.249
Medical history				
STS score	2.38 ± 0.96	2.38 ± 0.93	2.35 ± 1.11	0.893
NYHA class	2 [2–3]	2 [2–3]	2 [2–3]	0.928
NYHA class III/IV, n (%)	30 (39.0)	24 (38.8)	6 (40.0)	0.927
Diabetes mellitus, n (%)	20 (26.0)	13 (21.0)	7 (46.7)	0.054
Coronary artery disease, n (%)	41 (53.2)	33 (53.2)	8 (53.3)	0.994
COPD, <i>n</i> (%)	10 (13.0)	8 (12.9)	2 (13.3)	1.000
Atrial fibrillation, n (%)	17 (22.1)	14 (22.6)	3 (20.0)	1.000
MDRD-GFR, mL/min	64.8 ± 17.5	66.7 ± 17.0	57.0 <u>+</u> 17.8	0.053
Haemoglobin level, mmol/L	$7.9 \pm 0.9$	8.0 ± 0.9	7.6 ± 1.0	0.122
Pre-procedural echocardiographic				
parameters				
Aortic valve area, cm <sup>2</sup>	0.79 ± 0.18	$0.78 \pm 0.19$	$0.80 \pm 0.11$	0.794
Aortic valve mean gradient, mmHg	45.2 ± 12.4	44.9 ± 11.6	46.5 ± 15.7	0.642
Aortic valve maximum velocity, m/s	4.3 ± 0.6	$4.3 \pm 0.6$	$4.3 \pm 0.6$	0.583
Left ventricular ejection fraction, %	54.2 ± 8.4	54.8 ± 8.3	51.7 ± 8.4	0.203
Moderate or severe aortic regurgitation, n (%)	7 (9.1)	6 (9.7)	1 (6.7)	1.000

#### Table 1 Baseline characteristics

Data are presented as mean  $\pm$  standard deviation, median [interquartile range], or as number (%).

CMR-RF, cardiac magnetic resonance-regurgitant fraction; COPD, chronic obstructive pulmonary disease; MDRD-GFR, Modification of Diet in Renal Disease—glomerular filtration rate; NYHA, New York Heart Association; STS, Society of Thoracic Surgeons.

 $45.2\pm12.4$  mmHg, respectively. No significant differences were found in baseline characteristics between patients with either CMR-RF >20% or CMR-RF  $\leq20\%.$ 

# Procedural characteristics and clinical outcomes

Details of the procedural characteristics and clinical outcomes are provided in *Table 2*. A total of 17 (22.1%) TAVR procedures were performed under general anaesthesia, and 60 (77.9%) patients underwent TAVR under conscious sedation. In 70 (90.9%) patients, the transfemoral approach was used, and the transaxillary approach was used in the remaining seven (9.1%) patients. Pre-dilation was performed in 97.4% of all TAVR procedures, and post-dilation was performed in 19.5% of patients. A second valve was implanted in two (2.6%) patients due to migration of the first TAVR bioprosthesis. Median implantation depth below aortic annulus was 4 mm (IQR 4–6).

Within 30 days of the procedure, stroke/TIA occurred in three (3.9%) patients, permanent pacemaker implantation in eight (10.3%) patients, major vascular complications in two (2.6%) patients, and major bleeding complications in two (2.6%) patients. Acute kidney injury requiring temporary dialysis occurred in one (1.3%) patient. Technical success was 96.1%, and device success was 94.8%.

## Haemodynamic measurements

Procedural haemodynamic measurements are presented in *Table 3*. Pre-implantation, mean heart rate was  $67 \pm 15$  BPM, LVEDP was  $16 \pm 6$  mmHg, and peak-to-peak gradient was  $53 \pm 17$  mmHg. Mean DD was  $38 \pm 11$  mmHg, HR-DD was  $47 \pm 14$  mmHg/BPM, and ARI was  $33 \pm 9$ . Except for DD, all parameters changed significantly after implantation. Mean heart rate increased to  $78 \pm 18$  BPM (P < 0.001), LVEDP increased to  $19 \pm 6$  mmHg (P = 0.010), and peak-to-peak gradient decreased to  $4 \pm 3$  mmHg (P < 0.001). HR-DD decreased to  $42 \pm 13$  mmHg/BPM (P = 0.001), and ARI decreased to  $29 \pm 9$  (P = 0.003). Mean ARI ratio was  $0.95 \pm 0.45$ .

## Echocardiographic assessment of PVR

Mean duration between TAVR and TTE was 41  $\pm$  14 days. TTE assessment showed none/trace PVR in 41 (53.2%) patients, mild PVR in 34 (44.2%) patients, moderate PVR in two (2.6%) patients, and no patients with severe PVR.

# CMR quantification of regurgitant fraction (CMR-RF)

Mean forward volume measured with a high ( $\geq$ 180 cm/s) velocity ending (Venc) was 77.8 ± 19.0 mL. Mean regurgitant volume measured with a low Venc (75 cm/s) was 10.1 ± 8.5 mL, resulting in a mean

Table 2         Procedural characteristics and clinical outcomes		
	Study population (n = 77)	
Procedural characteristics		
Anaesthesia		
General anaesthesia, $n$ (%)	17 (22.1)	
Conscious sedation, n (%)	60 (77.9)	
Approach	· · · ·	
Transfemoral, n (%)	70 (90.9)	
Transaxillary, n (%)	7 (9.1)	
Pre-dilation, n (%)	75 (97.4)	
Post-dilation, n (%)	15 (19.5)	
Implantation depth, millimetre below	4 [4–6]	
aortic annulus		
Second valve implanted, $n$ (%)	2 (2.6)	
Clinical outcomes		
All stroke at 30 days, n (%)	3 (3.9)	
Stroke, n (%)	2 (2.6)	
TIA, n (%)	1 (1.3)	
Acute kidney injury, n (%)	1 (1.3)	
Permanent pacemaker implantation < 30 days, <i>n</i> (%)	8 (10.3)	
Vascular complications $< 30$ days, $n$ (%)	13 (16.9)	
Major, <i>n</i> (%)	2 (2.6)	
Minor, <i>n</i> (%)	11 (14.3)	
Bleeding complications < 30 days, n (%)	11 (14.3)	
Major, <i>n</i> (%)	2 (2.6)	
Minor, <i>n</i> (%)	9 (11.7)	
Technical success (at exit from	74 (96.1)	
procedure room), n (%)		
Device success (at 30 days), n (%)	73 (94.8)	
Early safety (at 30 days), n (%)	66 (85.7)	

Data are presented as mean  $\pm$  standard deviation, median [interquartile range], or as number (%). TIA, transient ischemic attack.

regurgitant fraction of  $12.4 \pm 9.3\%$ . Fifteen (19.5%) patients had CMR-RF > 20% (*Table 4*).

## Relation between haemodynamic indices and CMR-RF

Correlation analysis showed that DD had the best (inverse) correlation with CMR-RF (correlation coefficient: -0.352, P = 0.002), followed by HR-DD (correlation coefficient: -0.344, P = 0.002), ARI (correlation coefficient: -0.251 P = 0.027), and ARI ratio (correlation coefficient: -0.145, P = 0.209). DD also had the highest AUC (0.82; 95% CI, 0.72-0.92), followed by HR-DD (AUC 0.78; 95% CI, 0.67-0.89), ARI (AUC 0.78; 95% CI, 0.66-0.89), and ARI ratio (AUC 0.65; 95% CI, 0.49-0.81) (DD vs. ARI ratio, P = 0.01, other comparisons between AUCs non-significant) (*Figure 2*). Optimization of the cut-offs resulted in the following values: 32 mmHg for DD (SE: 69%, SP: 77%), 37 mmHg/ BPM for HR-DD (SE: 69%, SP: 67%), 25 for ARI (SE: 69%, SP: 69%), and 0.85 for ARI ratio (SE: 62%, SP: 58%). The previously proposed cut-off

of  $\leq$ 18 mmHg for DD had a SE of 0% and SP of 98% for the detection of CMR-RF > 20%. The SE and SP curves for DD to detect CMR-RF > 20% are shown in *Figure 3*.

## Discussion

In this study, we assessed four haemodynamic indices for their predictive value for relevant PVR based on gold standard CMR. This imaging modality served as the reference standard, in order to identify the most accurate haemodynamic index to grade PVR.

The key findings can be summarized as follows. First, in this prospective non-selective cohort of consecutive TAVR patients, we found a mean CMR-RF of 12%, with 19.5% of patients showing a CMR-RF > 20%. Second, the haemodynamic index DD had the best correlation with CMR-RF and the highest predictive value for CMR-RF > 20%, followed by HR-DD, ARI, and ARI ratio. Third, a cut-off value of 32 mmHg for DD provides a SE of 69% and a SP of 77% for the presence of CMR-RF > 20% at 1-month follow-up.

The mean CMR-RF of 12% in our study is comparable to the CMR-RF of other TAVR devices.<sup>9,11</sup> Echocardiographic assessment of PVR at 1-month follow-up also demonstrated that the majority of patients in our study had none/trace PVR, with only two patients with moderate PVR, and no cases of severe PVR. These numbers are in line with previous studies reporting on the echocardiographic performance of the Abbott Portico valve, as well as other TAVR platforms.<sup>22,23</sup>

A variety of CMR classifications have been proposed in literature, using different cut-offs to grade PVR. In a study performed by Ribeiro et  $al_{2}^{24}$  it was shown that a CMR-RF  $\geq$  30% was associated with an increased risk of mortality and the composite of mortality and rehospitalization for heart failure. In the 2022 EACVI and ESC position paper on multi-modality imaging assessment of native valvular regurgitation, the CMR cut-offs for mild, moderate, and severe aortic regurgitation are <30%, 30–49%, and  $\geq$ 50%, respectively.<sup>25</sup> Using this CMR grading scheme in our cohort would categorize 74 (96.1%) patients into less than or equal to mild PVR, three (3.9%) patients into moderate PVR, and zero patients into severe PVR. This closely resembles the degree of PVR as assessed with TTE (see Supplementary data online, Table S2). However, others have shown negative impact on clinical outcome at lower cut-offs.<sup>18</sup> Thus, for the present study, we adopted the more conservative CMR-RF cut-off value of 20%, aiming to diminish the risk of missing patients with potentially clinically relevant PVR. Another argument to use a more conservative cut-off in the present setting is the fact that a very high degree of PVR is generally already observed by angiography, thereby reducing the additional value of haemodynamic indices.

Although the incidences of mild and greater than or equal to moderate PVR as assessed by echocardiography have decreased with newer-generation TAVR devices and advancing techniques, risk of PVR is still considered a major shortcoming of TAVR. Moderate or severe PVR strongly and negatively affects clinical outcome, as illustrated by a three-fold increase in all-cause and cardiovascular mortality after TAVR compared with patients with none/trace PVR.<sup>7,26,27</sup> Recently, also mild–moderate PVR has been associated with an increased risk of mortality at 5 years after TAVR, implicating that also with this degree of PVR, additional interventions should be considered to reduce PVR.<sup>8</sup>

Moreover, we recently identified PVR as risk factor for recurrence of gastrointestinal bleeding after TAVR in patients with Heyde syndrome, further stressing the need for reduction of PVR.<sup>28</sup> Thus, assessment of PVR, including the use of the correct parameters and/or adoption of sensitive cut-offs, is of utmost importance in determining prognosis. Also, it could guide additional therapies aiming to reduce PVR and improve clinical outcome after TAVR.

#### M.J.P. Rooijakkers et al.

	Pre-implantation	Post-implantation	<i>P</i> value
Heart rate, beats per minute (BPM)	67 <u>±</u> 15	78 ± 18	<0.001
Systolic blood pressure (SBP), mmHg	118 ± 22	140 ± 28	<0.001
Diastolic blood pressure (DBP), mmHg	54 <u>±</u> 10	58 <u>+</u> 9	0.006
Left ventricular systolic pressure (LVSP), mmHg	171 ± 27	144 <u>+</u> 27	<0.001
Left ventricular end-diastolic pressure (LVEDP), mmHg	16 ± 6	19 <u>±</u> 6	0.010
Peak–peak gradient, mmHg	53 <u>+</u> 17	4 ± 3	<0.001
Mean gradient, mmHg	47 ± 12	9 <u>±</u> 4	<0.001
Haemodynamic indices			
Diastolic delta (DD)	38 ± 11	39 <u>±</u> 11	0.385
Heart rate-adjusted diastolic delta (HR-DD)	47 <u>+</u> 14	42 ± 13	0.001
Aortic regurgitation index (ARI)	33 ± 9	29 <u>+</u> 9	0.003
Aortic regurgitation index ratio (ARI ratio)	N/A	$0.95 \pm 0.45$	N/A

#### Table 3 Haemodynamic measurements

Data are presented as mean  $\pm$  standard deviation. N/A, not applicable.

 
 Table 4
 Cardiac magnetic resonance (CMR) and echocardiographic assessment of paravalvular regurgitation (PVR)

	Study population ( <i>n</i> = 77)
Days after TAVR	41 <u>±</u> 14
Flow measurements	
Forward volume, mL	77.8 <u>+</u> 19.0
Regurgitant volume, mL	10.1 ± 8.5
Regurgitant fraction, %	12.4 ± 9.3
CMR-RF classification	
CMR-RF > 20%, n (%)	15 (19.5)
CMR-RF ≤ 20%, n (%)	62 (80.5)
TTE classification of PVR <sup>a</sup>	
None/trace, n (%)	41 (53.2)
Mild, n (%)	34 (44.2)
Moderate, n (%)	2 (2.6)
Severe, n (%)	0

Data are presented as mean  $\pm$  standard deviation or as number (%).

CMR-RF, cardiac magnetic resonance-regurgitant fraction; TAVR, transcatheter aortic valve replacement; TTE, transthoracic echocardiography.

<sup>a</sup>TTE was performed on the same day as CMR.

Treatment options for PVR include balloon post-dilation (BPD), transcatheter valve-in-valve (ViV), and percutaneous closure by a vascular plug. BPD is usually considered first since this is easily performed during TAVR procedure and particularly effective in case of inadequate apposition or relative under sizing of the valve. Although BPD is an effective strategy in reducing PVR, the incidence of stroke after BPD is significantly higher,<sup>29</sup> and BPD can cause aortic rupture. Therefore, unnecessary BPD should be avoided. Transcatheter placement of a second valve (ViV) can adequately reduce significant PVR in case of malposition of the prosthesis (either too high or too low compared

with the aortic annulus).<sup>30</sup> Also, transcatheter closure of a paravalvular leak is feasible, although its use is still limited.<sup>31,32</sup>

Echocardiography also allows assessment of PVR, especially TEE. However, with the advent of the 'minimalist TAVR' approach, a commonly used term for performing TAVR under local anaesthesia with minimal sedation and promoting early discharge, the use of peri-procedural TEE as imaging modality is more or less precluded. TTE is less accurate in the assessment of PVR compared with TEE.<sup>33,34</sup> Various patient factors such as airway disease and habitus often limit acoustic windows for TTE. Regurgitant jets are often multiple and eccentric, making traditional echocardiographic regurgitation assessment techniques invalid. As a result, TTE potentially underestimates the degree of PVR when compared with gold standard CMR.<sup>9,10,18</sup>

Angiographic assessment using the visual Sellers' method also has several limitations, including subjectivity and lack of accuracy in quantifying PVR.<sup>34,35</sup> Moreover, aortic root angiography correlates only moderately to CMR in the classification of PVR.<sup>11</sup> This highlights the limitations of conventional modalities and the need for objective methodologies to aid in PVR assessment.

Transvalvular haemodynamic indices are cheap and readily available, can be performed at multiple instances during the procedure, and have very low inter- and intra-observer variability. Therefore, several studies have assessed the prognostic value of different peri-procedural haemodynamic indices. In these studies, it was shown that DD  $\leq$  18 mmHg,<sup>13</sup> HR-DD < 25 mmHg/BPM,<sup>14</sup> ARI < 25,<sup>15</sup> and ARI ratio < 0.6<sup>16</sup> are all independently associated with (1-year) mortality after TAVR. Previously, we showed that ARI ratio < 0.6 is the strongest haemodynamic predictor of 1-year mortality after TAVR.<sup>36</sup> In the present study, interestingly ARI ratio was least predictive for the occurrence of CMR-RF > 20%. Indeed, for clinical outcome not only the post-TAVR occurrence of PVR is important but also the pre-TAVR haemodynamic circumstances like diastolic dysfunction and pre-existing aortic regurgitation.<sup>37</sup> Therefore, ARI ratio and DD provide different and supplemental information for guidance of therapeutic strategy and prognosis.

We found that, regarding DD, the best cut-off to predict CMR-RF > 20% was 32 mmHg. This implies that in patients with a DD  $\leq$  32 mmHg, it could be considered to perform additional post-dilation, especially when conventional imaging modalities like aortic root angiography or echocardiography point in the same direction. Naturally, this



Figure 2 ROC analyses showing correlation between haemodynamic indices and relevant PVR expressed as CMR-RF > 20%. ARI, aortic regurgitation index; ARI ratio, aortic regurgitation index ratio; AUC, area under the curve; CMR-RF, cardiac magnetic resonance-regurgitant fraction; DD, diastolic delta; HR-DD, heart rate-adjusted diastolic delta; PVR, paravalvular regurgitation; Venc, velocity encoding. Asterisk (\*) indicates statistically significant difference between AUCs of DD and ARI ratio (P = 0.01).





should be left to the treating physician, also considering the risks of additional post-dilation. Future studies need to validate such an approach.

HR-DD, ARI, and ARI ratio had lower AUCs than DD, with a significantly lower AUC for ARI ratio than for DD. The aforementioned parameters are derivatives of DD, necessitating an additional calculation compared with the DD. In other words, our findings suggest that the most simple and most readily available haemodynamic index yields the highest accuracy to predict CMR-RF > 20%.

Besides the four haemodynamic indices assessed in this study, additional haemodynamic parameters for assessment of PVR have been addressed in literature. Integration of the systolic and diastolic time components during measurement of the ARI has resulted in the timeintegrated aortic regurgitation index (TIARI), which shows a higher AUC compared with the ARI,<sup>38</sup> and can be useful to guide BPD after valve deployment.<sup>39</sup> Dividing the area between aortic and left ventricular pressure-time curves by the duration of diastole generates the diastolic pressure-time index (DPTI). When DPTI is divided by the systolic blood pressure and multiplied by 100, the DPTI adjusted is obtained, which could be considered to differentiate between relevant and nonrelevant PVR after TAVR.<sup>40</sup> However, since both the TIARI and DPTI are time-integrated measures that are not readily available during TAVR procedure, we did not integrate these in the present study.

### **Future perspectives**

Future studies need to be conducted to prospectively assess the newly proposed cut-offs for the haemodynamic parameters described in our study. The DD in particular can be used to aid peri-procedural decision-making. Furthermore, videodensitometry has emerged as a complementary modality for grading PVR. This technique shows a high correlation with CMR-RF and can be used both offline and online.<sup>41,42</sup>

## Limitations

This trial has some limitations. First, the use of CMR might have led to selection bias whereby patients with an overall impaired health status could tend to decline participation in this study. Second, assessment of the accuracy of haemodynamic indices to predict a CMR-RF  $\geq$  30% was prohibited due to the limited number of patients in this group. Third, this study is solely performed with the self-expanding Abbott Portico bioprosthesis; therefore, the results are not directly applicable to other types of TAVR devices.

## Conclusion

In conclusion, we have demonstrated that the diastolic delta has the highest predictive value for the occurrence of relevant PVR (defined as CMR-RF > 20%) 1 month after TAVR. Routine assessment of this readily available haemodynamic parameter should be encouraged.

## Supplementary data

Supplementary data are available at European Heart Journal— Cardiovascular Imaging online.

## Acknowledgements

None declared.

## Funding

This research was funded by a research grant from Abbott Laboratories (Chicago, Illinois, USA).

**Conflict of interest**: M.J.P.R., N.A.S., K.v.d.W., L.R., H.G., L.A.F.M.v.G., G.S.C.G., M.W.A.V., J.H., S.E.M., and D.H.J.T. do not have potential conflicts

of interest or disclosures to report. M.H.v.W. has been a proctor and consultant for Abbott Vascular. R.N. has received research funding from Philips Volcano and Biotronik. N.v.R. has received research funding from Abbott, Philips, Medtronic, and Biotronik; has served as a consultant for RainMed, Castor, and Medtronic; and received speaker fees from Abbott and Bayer.

## Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

## References

- Popma JJ, Deeb GM, Yakubov SJ, Mumtaz M, Gada H, O'Hair D et al. Transcatheter aortic-valve replacement with a self-expanding valve in low-risk patients. N Engl J Med 2019;380:1706–15.
- Mack MJ, Leon MB, Thourani VH, Makkar R, Kodali SK, Russo M et al. Transcatheter aortic-valve replacement with a balloon-expandable valve in low-risk patients. N Engl J Med 2019;380:1695–705.
- Forrest JK, Deeb GM, Yakubov SJ, Rovin JD, Mumtaz M, Gada H et al. 2-Year outcomes after transcatheter versus surgical aortic valve replacement in low-risk patients. J Am Coll Cardiol 2022;79:882–96.
- Tamburino C, Bleiziffer S, Thiele H, Scholtz S, Hildick-Smith D, Cunnington M et al. Comparison of self-expanding bioprostheses for transcatheter aortic valve replacement in patients with symptomatic severe aortic stenosis: SCOPE 2 randomized clinical trial. *Circulation* 2020;**142**:2431–42.
- Kalogeropoulos AS, Redwood SR, Allen CJ, Hurrell H, Chehab O, Rajani R et al. A 20-year journey in transcatheter aortic valve implantation: evolution to current eminence. Front Cardiovasc Med 2022;9:971762.
- Athappan G, Patvardhan E, Tuzcu EM, Svensson LG, Lemos PA, Fraccaro C et al. Incidence, predictors, and outcomes of aortic regurgitation after transcatheter aortic valve replacement: meta-analysis and systematic review of literature. J Am Coll Cardiol 2013;61:1585–95.
- Chau KH, Chen S, Crowley A, Redfors B, Li D, Hahn RT et al. Paravalvular regurgitation after transcatheter aortic valve replacement in intermediate-risk patients: a pooled PARTNER 2 study. *EuroIntervention* 2022;**17**:1053–60.
- Okuno T, Tomii D, Heg D, Lanz J, Praz F, Stortecky S et al. Five-year outcomes of mild paravalvular regurgitation after transcatheter aortic valve implantation. *EuroIntervention* 2022;**18**:33–42.
- Crouch G, Tully PJ, Bennetts J, Sinhal A, Bradbrook C, Penhall AL et al. Quantitative assessment of paravalvular regurgitation following transcatheter aortic valve replacement. J Cardiovasc Magn Reson 2015;17:32.
- Orwat S, Diller GP, Kaleschke G, Kerckhoff G, Kempny A, Radke RM et al. Aortic regurgitation severity after transcatheter aortic valve implantation is underestimated by echocardiography compared with MRI. *Heart* 2014;**100**:1933–8.
- Frick M, Meyer CG, Kirschfink A, Altiok E, Lehrke M, Brehmer K et al. Evaluation of aortic regurgitation after transcatheter aortic valve implantation: aortic root angiography in comparison to cardiac magnetic resonance. EuroIntervention 2016;11:1419–27.
- Saikrishnan N, Kumar G, Sawaya FJ, Lerakis S, Yoganathan AP. Accurate assessment of aortic stenosis: a review of diagnostic modalities and hemodynamics. *Circulation* 2014; 129:244–53.
- Patsalis PC, Konorza TF, Al-Rashid F, Plicht B, Riebisch M, Wendt D et al. Incidence, outcome and correlates of residual paravalvular aortic regurgitation after transcatheter aortic valve implantation and importance of haemodynamic assessment. *EuroIntervention* 2013;8:1398–406.
- 14. Jilaihawi H, Chakravarty T, Shiota T, Rafique A, Harada K, Shibayama K et al. Heart-rate adjustment of transcatheter haemodynamics improves the prognostic evaluation of paravalvular regurgitation after transcatheter aortic valve implantation. *EuroIntervention* 2015;**11**:456–64.
- Sinning JM, Hammerstingl C, Vasa-Nicotera M, Adenauer V, Lema Cachiguango SJ, Scheer AC et al. Aortic regurgitation index defines severity of peri-prosthetic regurgitation and predicts outcome in patients after transcatheter aortic valve implantation. J Am Coll Cardiol 2012;59:1134–41.
- Sinning JM, Stundl A, Pingel S, Weber M, Sedaghat A, Hammerstingl C et al. Pre-procedural hemodynamic status improves the discriminatory value of the aortic regurgitation index in patients undergoing transcatheter aortic valve replacement. JACC Cardiovasc Interv 2016;9:700–11.
- Généreux P, Piazza N, Alu MC, Nazif T, Hahn RT, Pibarot P et al. Valve Academic Research Consortium 3: updated endpoint definitions for aortic valve clinical research. *Eur Heart J* 2021;42:1825–57.
- Hartlage GR, Babaliaros VC, Thourani VH, Hayek S, Chrysohoou C, Ghasemzadeh N et al. The role of cardiovascular magnetic resonance in stratifying paravalvular leak severity after transcatheter aortic valve replacement: an observational outcome study. J Cardiovasc Magn Reson 2014;16:93.

- Vasa-Nicotera M, Sinning JM, Chin D, Lim TK, Spyt T, Jilaihawi H et al. Impact of paravalvular leakage on outcome in patients after transcatheter aortic valve implantation. JACC Cardiovasc Interv 2012;5:858–65.
- Altiok E, Frick M, Meyer CG, Al Ateah G, Napp A, Kirschfink A et al. Comparison of two- and three-dimensional transthoracic echocardiography to cardiac magnetic resonance imaging for assessment of paravalvular regurgitation after transcatheter aortic valve implantation. Am J Cardiol 2014;**113**:1859–66.
- DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;44:837–45.
- Søndergaard L, Rodés-Cabau J, Hans-Peter Linke A, Fichtlscherer S, Schäfer U, Kuck KH et al. Transcatheter aortic valve replacement with a repositionable self-expanding prosthesis: the PORTICO-I trial 1-year outcomes. J Am Coll Cardiol 2018;72:2859–67.
- Mollmann H, Linke A, Holzhey DM, Walther T, Manoharan G, Schafer U et al. Implantation and 30-day follow-up on all 4 valve sizes within the portico transcatheter aortic bioprosthetic family. JACC Cardiovasc Interv 2017;10:1538–47.
- Ribeiro HB, Orwat S, Hayek SS, Larose É, Babaliaros V, Dahou A et al. Cardiovascular magnetic resonance to evaluate aortic regurgitation after transcatheter aortic valve replacement. J Am Coll Cardiol 2016;68:577–85.
- Lancellotti P, Pibarot P, Chambers J, La Canna G, Pepi M, Dulgheru R et al. Multi-modality imaging assessment of native valvular regurgitation: an EACVI and ESC council of valvular heart disease position paper. *Eur Heart J Cardiovasc Imaging* 2022; 23:e171–232.
- Leon MB, Smith CR, Mack MJ, Makkar RR, Svensson LG, Kodali SK et al. Transcatheter or surgical aortic-valve replacement in intermediate-risk patients. N Engl J Med 2016; 374:1609–20.
- Makkar RR, Thourani VH, Mack MJ, Kodali SK, Kapadia S, Webb JG et al. Five-year outcomes of transcatheter or surgical aortic-valve replacement. N Engl J Med 2020;382: 799–809.
- Goltstein L, Rooijakkers MJP, Görtjes NCC, Akkermans RP, Zegers ES, Pisters R et al. Reduction of gastrointestinal bleeding in patients with Heyde syndrome undergoing transcatheter aortic valve implantation. *Circ Cardiovasc Interv* 2022;**15**:e011848.
- Wang N, Lal S. Post-dilation in transcatheter aortic valve replacement: a systematic review and meta-analysis. J Interv Cardiol 2017;30:204–11.
- Stundl A, Rademacher MC, Descoups C, Weber M, Sedaghat A, Grube M et al. Balloon post-dilation and valve-in-valve implantation for the reduction of paravalvular leakage with use of the self-expanding CoreValve prosthesis. *EuroIntervention* 2016;**11**:1140–7.

- Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J et al. 2021 ESC/ EACTS guidelines for the management of valvular heart disease. Eur Heart J 202;43: 561–632.
- Giblett JP, Rana BS, Shapiro LM, Calvert PA. Percutaneous management of paravalvular leaks. Nat Rev Cardial 2019;16:275–85.
- Kronzon I, Jelnin V, Ruiz CE, Saric M, Williams MR, Kasel AM et al. Optimal imaging for guiding TAVR: transesophageal or transthoracic echocardiography, or just fluoroscopy? JACC Cardiovasc Imaging 2015;8:361–70.
- Abdelghani M, Soliman OI, Schultz C, Vahanian A, Serruys PW. Adjudicating paravalvular leaks of transcatheter aortic valves: a critical appraisal. Eur Heart J 2016;37:2627–44.
- Sellers RD, Levy MJ, Amplatz K, Lillehei CW. Left retrograde cardioangiography in acquired cardiac disease: technic, indications and interpretations in 700 cases. Am J Cardiol 1964;14:437–47.
- 36. van Wely M, van der Wulp K, Rooijakkers M, Vart P, Morshuis W, van Royen N et al. Aortic regurgitation index ratio is a strong predictor of 1-year mortality after transcatheter aortic valve implantation using self-expanding devices. Semin Thorac Cardiovasc Surg 2021;**33**:923–30.
- Halkin A, Steinvil A, Aviram G, Biner S, Banai S, Keren G et al. Aortic regurgitation following transcatheter aortic valve replacement: impact of preprocedural left ventricular diastolic filling patterns on late clinical outcomes. *Catheter Cardiovasc Interv* 2016;87: 1156–63.
- Bugan B, Kapadia S, Svensson L, Krishnaswamy A, Tuzcu EM. Novel hemodynamic index for assessment of aortic regurgitation after transcatheter aortic valve replacement. *Catheter Cardiovasc Interv* 2015;86:E174–9.
- Kumar A, Sato K, Jobanputra Y, Betancor J, Halane M, George R et al. Time-integrated aortic regurgitation index helps guide balloon postdilation during transcatheter aortic valve replacement and predicts survival. J Am Heart Assoc 2019;8:e012430.
- Höllriegel R, Woitek F, Stativa R, Mangner N, Haußig S, Fuernau G et al. Hemodynamic assessment of aortic regurgitation after transcatheter aortic valve replacement: the diastolic pressure-time index. JACC Cardiovasc Interv 2016;9:1061–8.
- Modolo R, van Mourik M, El Bouziani A, Kawashima H, Rosseel L, Abdelghani M et al. Online quantitative aortographic assessment of aortic regurgitation after TAVR: results of the OVAL study. JACC Cardiovasc Interv 2021;14:531–8.
- Abdel-Wahab M, Abdelghani M, Miyazaki Y, Holy EW, Merten C, Zachow D et al. A novel angiographic quantification of aortic regurgitation after TAVR provides an accurate estimation of regurgitation fraction derived from cardiac magnetic resonance imaging. JACC Cardiovasc Interv 2018;11:287–97.