

LJMU Research Online

Dillamore, AM, Belokurov, V, Kravtsov, A and Font, AS

Taking the Milky Way for a spin: disc formation in the ARTEMIS simulations

https://researchonline.ljmu.ac.uk/id/eprint/23121/

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Dillamore, AM ORCID logoORCID: https://orcid.org/0000-0003-0807-5261, Belokurov, V ORCID logoORCID: https://orcid.org/0000-0002-0038-9584, Kravtsov, A ORCID logoORCID: https://orcid.org/0000-0003-4307-634X and Font. AS ORCID logoORCID: https://orcid.org/0000-0001-8405-9883 (2023)

LJMU has developed LJMU Research Online for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

http://researchonline.ljmu.ac.uk/

Taking the Milky Way for a spin: disc formation in the ARTEMIS simulations

Adam M. Dillamore⁽⁰⁾,¹* Vasily Belokurov⁽⁰⁾,¹ Andrey Kravtsov⁽⁰⁾,^{2,3,4} and Andreea S. Font⁽⁰⁾

¹Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

²Department of Astronomy and Astrophysics, The University of Chicago, Chicago, IL 60637, USA

³Kavli Institute for Cosmological Physics, The University of Chicago, Chicago, IL 60637, USA

⁴Enrico Fermi Institute, The University of Chicago, Chicago, IL 60637, USA

⁵Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, UK

Accepted 2023 October 26. Received 2023 October 25; in original form 2023 September 15

ABSTRACT

We investigate the formation (spin-up) of galactic discs in the ARTEMIS simulations of Milky Way (MW)-mass galaxies. In almost all galaxies, discs spin up at higher [Fe/H] than the MW. Those galaxies that contain an analogue of the *Gaia* Sausage-Enceladus (GSE) spin up at a lower average metallicity than those without. We identify six galaxies with spin-up metallicity similar to that of the MW, which formed their discs ~8–11 Gyr ago. Five of these experience a merger similar to the GSE. The spin-up times correlate with the halo masses at early times: galaxies with early spin-up have larger virial masses at a lookback time $t_L = 12$ Gyr. The fraction of stars accreted from outside the host galaxy is smaller in galaxies with earlier spin-ups. Accreted fractions small enough to be comparable to the MW are only found in galaxies with the earliest disc formation and large initial virial masses ($M_{200c} \approx 2 \times 10^{11}$ M_{\odot} at $t_L = 12$ Gyr). We find that discs form when the halo's virial mass reaches a threshold of $M_{200c} \approx (6 \pm 3) \times 10^{11}$ M_{\odot}, independent of the spin-up time. However, the failure to form a disc in other galaxies appears to be instead related to mergers at early times. We also find that discs form when the central potential is not particularly steep. Our results indicate that the MW assembled its mass and formed its disc earlier than the average galaxy of a similar mass.

Key words: Galaxy: disc – Galaxy: evolution – Galaxy: formation – Galaxy: kinematics and dynamics.

1 INTRODUCTION

The Milky Way (MW), like all other spiral galaxies, was not born with a rotating disc, it had to form one. Exactly when and how galaxies acquire stable stellar discs is an open question, currently pursued both with cutting-edge lookback observations and through theoretical and numerical calculations.

The ratio of the rotational velocity to the velocity dispersion (v/σ) is measured to decrease significantly with increasing redshift for galaxies of a given stellar mass (see Kassin et al. 2012; Wisnioski et al. 2015). This disc *settling* is a mass-dependent process: more massive galaxies attain higher v/σ values earlier (Wisnioski et al. 2019). Mixed in with these trends is the information on the disc *emergence*, i.e. the galaxy's transformation from a state characterized by random motions to a configuration dominated by a coherently rotating stellar disc. Currently available samples of high-redshift galaxies with morphological and kinematic measurements are still too small to answer this question through population studies (but see recent studies by Ferreira et al. 2022, 2023; Jacobs et al. 2023; Nelson et al. 2023; Robertson et al. 2023). However, in the MW, archaeological records containing accurate kinematic information provided by *Gaia* (Gaia Collaboration 2016) and detailed chemical

abundances supplied by surveys such as APOGEE (Majewski et al. 2017) and GALAH (De Silva et al. 2015) have recently been used to crack the puzzle of the disc emergence.

For example, Belokurov & Kravtsov (2022) show that the stars born *in situ* in the MW (separated from the accreted stellar debris using the abundance ratio of aluminium to iron [Al/Fe]) exhibit a characteristic trend of the increasing rotational velocity with metallicity [Fe/H]. At low metallicity, i.e. at [Fe/H] < -1.3, the MW stars possess little to no net spin; however, from [Fe/H] \approx -1.3 to [Fe/H] \approx -0.9, the median azimuthal velocity v_{ϕ} rises from \approx 50 to \approx 150 km s⁻¹. Belokurov & Kravtsov (2022) associate this rapid *spin-up* with the emergence of the old Galactic disc, following the turbulent and chaotic state of the Galaxy imprinted in the ancient *in situ* stellar population *Aurora*.¹ While more contaminated in their selection of the *in situ* population, the follow-up studies of Conroy et al. (2022) and Rix et al. (2022) report similar trends of the rotational velocity with metallicity.

Belokurov & Kravtsov (2022) also highlight a discrepancy between the observations and the numerical simulations of galaxy formation. While the spin-up itself is a ubiquitous feature in chemokinematic histories of model MW-sized galaxies, the metal-

¹Named after Aurora – the Latin name of the goddess of dawn Eos in Greek mythology.

© The Author(s) 2023.

Published by Oxford University Press on behalf of Royal Astronomical Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

^{*} E-mail: amd206@cam.ac.uk

licity at which the disc forms in the two simulation suites analysed (Auriga and FIRE) is noticeably higher compared to the observations. Similar results were also produced by McCluskey et al. (2023) using the FIRE-2 simulations, although one of these galaxies (named Romeo) does have a similar time of disc formation to the MW. This question is further explored in Semenov et al. (2023a) using a larger sample of MW-sized galaxies from the Illustris TNG50 simulation to study the statistics of the spin-up metallicity. They conclude that in TNG50 only \approx 10 per cent of MW-sized galaxies form their disc at metallicities similar to those measured by Belokurov & Kravtsov (2022). They also show that the haloes of these galaxies assemble their mass early.

One of the most significant events in the MW's history was the merger with Gaia Sausage-Enceladus (GSE; Belokurov et al. 2018; Helmi et al. 2018), a massive satellite of total mass $\sim 10^{11} M_{\odot}$, that was accreted by the MW 8-11 Gyr ago (e.g. Belokurov et al. 2018; Fattahi et al. 2019). This is observed as a population of stars in the stellar halo with highly eccentric orbits and relatively high metallicity. MW analogues with GSE-like mergers have been studied in various suites of high-resolution zoomed-in cosmological simulations, such as Auriga (Grand et al. 2017, 2020; Fattahi et al. 2019) and ARTEMIS (Font et al. 2020; Dillamore et al. 2022, 2023). These have shown that approximately one-third of MW-like galaxies possess a dominant radially anisotropic feature in their stellar haloes resembling the GSE (Fattahi et al. 2019), and that the associated mergers have transformative effects on their host galaxies (Dillamore et al. 2022). These include creation of an in situ stellar halo (Belokurov et al. 2020; Grand et al. 2020) and tipping of the disc (Dillamore et al. 2022; Orkney et al. 2023).

Following the above early attempts to gauge the conditions necessary for the disc emergence, in this study we focus on the connection between the disc spin-up in model MW-like galaxies and their mass assembly histories using the ARTEMIS simulation suite. This paper is arranged as follows: We briefly describe the ARTEMIS simulations in Section 2 and our methods in Section 3. Our results are presented and discussed in Section 4, and summarized in Section 5. Finally, in Appendix A we show the star formation density of a selection of ARTEMIS galaxies across time to illustrate the changes in morphology involved in disc formation.

2 SIMULATIONS

ARTEMIS (Font et al. 2020) is a zoomed-in hydrodynamical simulation suite, consisting of 45 haloes with MW-like masses. Details can be found in Font et al. (2020) and are outlined below.

The simulations were run using the GADGET-3 code (Springel 2005) in a *WMAP* Lambda cold dark matter cosmology. While the hydrodynamics code and subgrid physics are shared with the EAGLE project (Crain et al. 2015; Schaye et al. 2015), the stellar feedback was recalibrated to better match the stellar mass–halo mass relation (Font et al. 2020).

An initial collisionless simulation was run to redshift zero in a base periodic box of side length 25 Mpc h^{-1} . A set of 45 galaxies was then selected based purely on their virial masses. These lie in the range $0.8 < M_{200c}/10^{12} M_{\odot} < 2.0$ at redshift z = 0, where M_{200c} is the mass enclosed within a volume containing a mean density of 200 times the critical density.

The final high-resolution simulations have a dark matter particle mass of $1.17 \times 10^5 \,\mathrm{M_{\odot}} \,h^{-1}$ and an initial baryon particle mass of $2.23 \times 10^4 \,\mathrm{M_{\odot}} \,h^{-1}$. The Plummer-equivalent softening length is 125 pc h^{-1} . Haloes and subhaloes were identified with the SUBFIND algorithm (Dolag et al. 2009).

The size–stellar mass relation of the ARTEMIS galaxies closely matches the observations by Shen et al. (2003) in the Sloan Digital Sky Survey (SDSS, Font et al. 2020). However, the typical disc masses of the ARTEMIS galaxies are $\approx 10^{10}\,M_{\odot}$, smaller than the estimate for the MW of $\approx (5\pm0.5)\,\times\,10^{10}\,M_{\odot}$ (e.g. McMillan 2017).

3 METHOD

3.1 Selection of disc galaxies

We define a disc galaxy using the corotation parameter κ_{co} (Correa et al. 2017), the fraction of stellar kinetic energy associated with motion in the positive sense about the *z*-axis. Specifically,

$$\kappa_{\rm co} = \frac{K_{\rm rot}}{K} = \frac{1}{K} \sum_{i}^{L_{z,i} > 0,} \frac{1}{2} m_i \left(\frac{L_{z,i}}{m_i R_i}\right)^2, \tag{1}$$

where m_i is the mass of star particle *i*, while $L_{z,i}$ and R_i are its *z*-angular momentum and projected galactocentric radius in the *x*-*y* plane, respectively. The sum is over all particles with r < 30 kpc and positive angular momentum (i.e. circulating in the same sense as the net rotation). *K* is the total kinetic energy of stars with r < 30 kpc.

In Dillamore et al. (2022), we defined a disc galaxy using threshold $\kappa_{co} \geq 0.4$ (Correa et al. 2017) and used this method to identify 27 of the 45 ARTEMIS galaxies with discs at the present day. In this study, we use a stricter threshold of $\kappa_{co} \geq 0.5$, which removes the borderline cases of weak discs and retains only galaxies in which more than half of the stellar kinetic energy is associated with net rotation. This results in a sample of 18 galaxies that we use in this study. In three of these, κ_{co} decreases below 0.5 before recovering to exceed it again after t_{spin} . There are also 17 examples of galaxies where $\kappa_{co} \geq 0.5$ at some earlier snapshot but not at the present day. We include these in Fig. 2.

Since κ_{co} defines the presence of a stellar disc, we also use it to define the lookback time of disc emergence or spin-up time, t_{spin} . We let t_{spin} be the lookback time of the earliest snapshot at which $\kappa_{co} \ge 0.5$. Since κ_{co} takes into account all stars present within r = 30 kpc at a particular snapshot, this should be seen as the time at which the galaxy becomes disc dominated rather than when the disc starts forming. We therefore find that star formation in a disc-like configuration generally begins a few snapshots (or $\approx 1-2$ Gyr) before t_{spin} , as can be seen in Fig. A1. We also note that this figure shows that, generally, stars born before disc formation have very irregular distribution, consistent with results of Belokurov & Kravtsov (2022, see their fig. 12) based on the FIRE-2 simulations. Thus, simulations do not show the existence of 'ancient' low-metallicity discs before the actual disc spin-up at metallicities [Fe/H] $\gtrsim -1.5$.

3.2 Selection of the GSE sample

In Dillamore et al. (2022), we identified ARTEMIS galaxies containing a feature resembling the GSE in their accreted stellar halo. This selection process is briefly described below [see Dillamore et al. (2022) for full details and discussion of the fitting procedure]

Particles in ARTEMIS are classed as *in situ* if they were born bound to the host subhalo, defined as the most massive subhalo identified by the SUBFIND algorithm (Dolag et al. 2009) at each simulation snapshot. They are otherwise classed as accreted.

We calculate the velocity components in spherical coordinates of accreted stars in a solar neighbourhood-like region [5 < R (kpc) < 15, 0 < |z| (kpc) < 9] in each galaxy at the redshift z = 0



Figure 1. Left-hand panel: Median v_{ϕ} as a function of age, with different colours indicating GSE and non-GSE galaxies. The solid lines and bands indicate the medians and 16th–84th percentile ranges of each sample. Middle panel: As above, but as a function of [Fe/H]. Data from the MW are shown with the black dashed line. The dotted lines mark the individual ARTEMIS galaxies selected as 'early spin-up', those with the highest median v_{ϕ} at [Fe/H] = -1. These have similar spin-up profiles to the MW. Right-hand panel: As above, but median [Fe/H] as a function of age. Measurements for the MW's globular clusters are shown with horizontal error bars. This demonstrates that our corrected [Fe/H] values in ARTEMIS align the [Fe/H] versus age tracks with the MW's *in situ* globular clusters (those with higher [Fe/H] at a given age; see Myeong et al. 2018; Massari, Koppelman & Helmi 2019).

snapshot. A two-component Gaussian mixture model is fitted to these velocity distributions using the expectation–maximization algorithm GAUSSIANMIXTURE from the SCIKIT-LEARN library (Pedregosa et al. 2011). All stars within the above cuts are included in these fits. The two Gaussian components represent a radially anisotropic and a more isotropic distribution. The radial anisotropy is characterized by the anisotropy parameter $\beta \equiv 1 - (\sigma_{\theta}^2 + \sigma_{\phi}^2)/(2\sigma_r^2)$ (Binney & Tremaine 2008). For the observed GSE feature in the MW, $\beta \approx 0.86$ (Fattahi et al. 2019).

We include a halo in our GSE sample if the more anisotropic (higher β) Gaussian component has $\beta > 0.8$ and a contribution to the accreted solar neighbourhood population of greater than 40 per cent. This follows a similar criteria used by Fattahi et al. (2019). The GSE sample therefore consists of galaxies with significant, highly radial features in their accreted haloes. This gives us a total of 23 galaxies in the GSE sample, 10 of which also have $\kappa_{co} \ge 0.5$ at the present day.

Our final sample of 18 disc galaxies therefore contains 10 GSE galaxies. We refer to the remaining eight as non-GSE galaxies; these are galaxies with $\kappa_{co} \ge 0.5$ at the present day but no GSE-like feature.

4 RESULTS

4.1 Rotational velocity, age, and [Fe/H]

We follow Belokurov & Kravtsov (2022) and Semenov et al. (2023a) and investigate the relations between the median azimuthal velocity v_{ϕ} , age, and [Fe/H] of stars in the solar neighbourhood. We work in a coordinate system with its *z*-axis aligned with the total angular momentum of stars within r = 30 kpc of the galactic centre. We select stars from the redshift z = 0 snapshot from the region 5 < R (kpc) <11, |z| < 3 kpc that are flagged as *in situ* in origin. To correct for the metallicity differences between different galaxies, we subtract the median [Fe/H] of each sample such that the corrected median [Fe/H] is zero in each galaxy. This means that the median solar neighbourhood metallicities approximately match that of the MW (Haywood 2001).

For each galaxy, we divide the stars into bins of age and [Fe/H], and calculate the median v_{ϕ} and [Fe/H] in each bin. The results are shown

in Fig. 1. The galaxies are divided into GSE (red) and non-GSE (blue) samples, with the medians and 16th–84th percentile ranges of each shown by the coloured lines and bands, respectively. For comparison, we show data for the MW from APOGEE DR17 (middle panel; Majewski et al. 2017; Abdurro'uf et al. 2022; Belokurov & Kravtsov 2022) and the MW's globular clusters (right-hand panel; VandenBerg et al. 2013).

Belokurov & Kravtsov (2022) found that the transition from low to high v_{ϕ} (spin-up) generally occurs at higher metallicities in the FIRE simulations than in the MW (except for the galaxy in the Romeo halo; see McCluskey et al. 2023). The middle panel of Fig. 1 shows that this is also true in the ARTEMIS galaxies, with a median $v_{\phi} = 150 \text{ km s}^{-1}$ occurring at [Fe/H] ≈ -0.5 compared to -1 in the MW. However, there is a difference between the GSE and non-GSE samples. The galaxies that spin-up at lower metallicities and earlier times are more likely to contain a GSE analogue, whereas the median v_{ϕ} of the non-GSE galaxies spins up at later times and higher metallicities.

From the middle panel, we select six galaxies that have the most MW-like v_{ϕ} versus [Fe/H] tracks, henceforth 'early spin-up galaxies'. These are the six galaxies with the highest median v_{ϕ} at [Fe/H] = -1, five of which are in the GSE sample. The tracks of these galaxies are shown with dotted lines in each panel of Fig. 1. The left-hand panel confirms that they do indeed spin-up at earlier than average times, reaching a median $v_{\phi} \sim 150 \text{ km s}^{-1}$ by $t_{\rm L} \sim 10 \text{ Gyr}$.

4.2 Spin-up times and evolution of haloes and their mass concentration

We show the relation between the spin-up time $t_{\rm spin}$ and the galaxies' masses at early times in Fig. 2. The top and middle panels show the virial mass M_{200c} and maximum circular velocity $v_{\rm max}$, respectively, both calculated at a lookback time of $t_{\rm L} = 12$ Gyr. The galaxies with discs ($\kappa_{\rm co} \ge 0.5$) at the z = 0 snapshot are shown by circles, with the early spin-up galaxies selected from Fig. 1 shown by circles with black edges. We also show galaxies where $\kappa_{\rm co}$ exceeds 0.5 at some earlier snapshot, but is less than 0.5 at the present day ('former discs', marked with crosses). There is a moderate correlation between $t_{\rm spin}$ and both the virial mass and $v_{\rm max}$ at $t_{\rm L} = 12$ Gyr, with earlier



Figure 2. Top panel: Virial mass M_{200c} at a lookback time of $t_{\rm L} = 12$ Gyr against spin-up time $t_{\rm spin}$ for all haloes with $\kappa_{\rm co} \ge 0.5$ at the present day (circles). Here, we also show galaxies with $\kappa_{\rm co} \ge 0.5$ at earlier snapshots only, but not at the present day ('former discs'; crosses). Red (blue) points indicate haloes in the (non-)GSE samples, and the early spin-up galaxies selected from Fig. 1 are marked with black rings. **Middle panel:** As above, but showing the maximum circular velocity $v_{\rm max}$ (at $t_{\rm L} = 12$ Gyr) against $t_{\rm spin}$. In both cases there is a correlation with spin-up time, with earlier spin-up soccurring in galaxies with higher masses and circular velocities at $t_{\rm L} = 12$ Gyr. **Bottom panel:** Fraction of accreted stars within r = 30 kpc at the present day against spin-up time. An estimate and its uncertainty for the MW is shown with a black line and grey band. Disc galaxies with earlier spin-ups tend to have lower accreted stellar fractions, while the former discs (crosses) have much larger proportions of accreted stars.

spin-ups occurring in galaxies with larger $M_{200c}(t_{\rm L} = 12 \,{\rm Gyr})$ and $v_{\rm max}(12 \,{\rm Gyr})$. In particular, the early spin-up galaxies have both larger $M_{200c}(12 \,{\rm Gyr})$ and $v_{\rm max}(12 \,{\rm Gyr})$ than almost all other present-day disc galaxies.

There is a large number of non-GSE galaxies with $t_{\rm spin}$ between 6 and 10 Gyr and $M_{200c}(12 \,{\rm Gyr}) \lesssim 10^{11} {\rm M_{\odot}}$, lower than the $M_{200c}(12 \,{\rm Gyr})$ values of early spin-up galaxies' masses. This is likely related to the different accretion histories of the GSE and non-GSE galaxies. We previously showed (Dillamore et al. 2023) that the ARTEMIS galaxies lacking a GSE analogue undergo more massive mergers than those with a GSE-like feature, many of which happen late. Their typical accreted satellites have stellar masses of $\sim 10^{10} \,{\rm M_{\odot}}$, roughly four times larger than those of GSE galaxies. This means that a greater proportion of the galaxies' final mass is assembled after $t_{\rm L} = 12 \,{\rm Gyr}$. Hence, to reach a final halo mass within the ARTEMIS selection criteria, these galaxies typically have lower masses at early times than those with a GSE analogue.

To show how accretion history relates to the spin-up time, we calculate the fraction of stars within r = 30 kpc that are classed as accreted. This is plotted against t_{spin} in the bottom panel of Fig. 2. For comparison, we calculate an estimate of the MW's accreted fraction as follows. We take the measurement of the MW's total stellar mass from McMillan (2017), $M^* = (54.3 \pm 5.7) \times 10^9 \text{ M}_{\odot}$, and the total stellar halo mass from Deason, Belokurov & Sanders (2019), $M^*_{\text{halo}} = (1.4 \pm 0.4) \times 10^9 \text{ M}_{\odot}$. We assume that a fraction $f_{\text{acc,halo}} = 0.81$ of the stellar halo is accreted (Naidu et al. 2020), and calculate our estimated overall accreted fraction $f_{\text{acc}} = f_{\text{halo,acc}} M^*_{\text{halo}} / M^* = 0.021 \pm 0.006$. This estimate and its uncertainty are shown in the bottom panel of Fig. 2 by the black solid line and grey shaded region. Since this calculation assumes that all accreted stars are in the halo, we have checked that our results do not significantly change if we exclude accreted stars on disc-like orbits in the simulations.

For the present-day disc galaxies, the galaxies with earlier spinups tend to have smaller accreted stellar fractions. A large majority of the ARTEMIS galaxies have accreted fractions significantly larger than that of the MW. The exceptions (G19, G38, and to a lesser extent G34) all have early spin-ups ($t_{spin} > 9$ Gyr) with MW-like v_{ϕ} versus [Fe/H] relations (see Fig. 1). The former disc galaxies (mostly lacking a GSE feature) tend to have larger accreted fractions, including those with early spin-ups. This suggests that mergers are responsible for destroying these discs, and are consistent with the conclusions of Dillamore et al. (2023) that non-GSE galaxies on average undergo more major mergers.

In Fig. 3, we show how the total accreted stellar fractions change across lookback time. At each snapshot, we calculate the fraction of all stars within r = 30 kpc that have been accreted before that time. The medians and 16th–84th percentile ranges are plotted versus lookback time t_L for the GSE and non-GSE present-day disc galaxy samples, as well as individually for the six early spin-up galaxies. The GSE galaxies have lower median accreted fractions at all times after $t_L \approx 11$ Gyr. The non-GSE galaxies have much larger contributions from accreted stars, peaking at ≈ 0.3 at $t_L = 10$ Gyr. This is likely a reflection of their more active accretion histories (Dillamore et al. 2023). Five of the six early spin-up galaxies have small accreted fractions across all ages, at ≤ 0.1 at most snapshots. Finally, we note that the median accreted fractions of $\sim 0.1-0.3$ are roughly consistent with the results of Semenov et al. (2023a).

In the top panel of Fig. 4, we show the evolution of the virial mass M_{200c} with lookback time $t_{\rm L}$ for the 18 present-day disc galaxies. We mark the spin-up times and corresponding virial masses with coloured points. The 16th–84th percentile range of the former discs is also shown as a function of time. The bottom panel of Fig. 4 shows the ratio of the circular velocity $v_c \equiv [GM_{tot}(< r)/r]^{1/2}$ at the stellar half-mass radius, $v_c(R_{\rm eff})$, and $v_c(3R_{\rm eff})$, again marking the corresponding values at the disc spin-up. In Hopkins et al. (2023,



Figure 3. Accreted fraction of stars within r = 30 kpc as a function of lookback time. In each snapshot all accreted stars present in the host galaxy at that time (within 30 kpc) are included in the calculations. The lines and bands have the same meanings as in Fig. 1. At $t_L < 11$ Gyr, the non-GSE galaxies have a higher median fraction of accreted stars, with a median of about 30 per cent at $t_L = 10$ Gyr. With one exception the early spin-up galaxies generally have low accreted fractions (≤ 0.1).

see their figs 6 and 7), the galaxies that form gas discs all have $v_c(R_{\rm eff})/v_c(3R_{\rm eff}) \gtrsim 1$ at the present day, and mass concentration (represented by the shape of the circular velocity profile) was used as a measure of the steepness of the central potential.

Fig. 4 shows that the virial masses at spin-up span a rather narrow range of $\sim 3-9 \times 10^{11} \,\mathrm{M}_{\odot}$, considerably smaller than the range of masses $M_{200c}(t_{\rm L} = 12 \,\mathrm{Gyr})$ in Fig. 2. Moreover, there is no apparent trend of $M_{200c}(t_{\rm spin})$ with $t_{\rm spin}$. This indicates that in the ARTEMIS simulations discs in $M_{200c}(z = 0) \approx 10^{12} \,\mathrm{M}_{\odot}$ haloes form when their host halo reaches $M_{200c} \approx (6 \pm 3) \times 10^{11} \,\mathrm{M}_{\odot}$. We see this more clearly in the top-left panel of Fig. 5, where we plot M_{200c} versus κ_{co} for each individual present-day disc galaxy. κ_{co} remains small (<0.3), while M_{200c} increases from $\sim 10^{10} \,\mathrm{to} \sim 10^{11} \,\mathrm{M}_{\odot}$.

$$M_{200c}(t_{\rm L} = 12\,{\rm Gyr})$$
 (2)

Interestingly, this virial mass value is close to the mass threshold at which a hot halo forms and gas accretion transitions from coldmode to hot-mode accretion (Kereš et al. 2005, 2009; Dekel & Birnboim 2006; Dekel et al. 2009). Thus, this result is consistent with the scenario in which a hot halo is essential for disc formation by mediating the accretion of gas (e.g. Dekel et al. 2020; Stern et al. 2021; Hafen et al. 2022). Indeed, fig. 12 in Semenov et al. (2023b) shows that a hot gaseous halo forms over a significant fraction of the virial radius immediately before disc spin-up.

At the same time, the bottom panel of Fig. 4 shows that disc spinup generally occurs when $v_c(R_{eff})/v_c(3R_{eff}) < 1$, i.e. when the mass distribution was not particularly centrally concentrated. This is also seen in the lower panels of Fig. 5, where the circular velocity ratio is plotted as a function of κ_{co} . This result is consistent with findings of Semenov et al. (2023b, see their figs 7–9) that the mass distribution becomes more concentrated after the disc forms and that halo mass at spin-up has a narrower range than their mass concentration indicator.

We note that all discs have $v_c(R_{\text{eff}})/v_c(3R_{\text{eff}}) \gtrsim 1$ by z = 0, in agreement with findings of Hopkins et al. (2023), who found that all simulated galaxies that form a disc have $v_c(R_{\text{eff}})/v_c(3R_{\text{eff}}) \gtrsim 1$ at z = 0 (see e.g. their figs 4–7). Fig. 4, however, shows that the mass



Figure 4. Top panel: Virial mass M_{200c} versus time for the 18 presentday disc galaxies (coloured lines). The spin-up times and masses at the corresponding snapshots are shown by the coloured points, where the colours have the same meanings as in Fig. 2. The population of former discs is also shown with the grey band, which spans between the 16th and 84th percentiles of M_{200c} as a function of time. The panel shows that disc in the ARTEMIS galaxies forms when their haloes reach mass of $M_{200c} \approx (6 \pm 3) \times 10^{11} \text{ M}_{\odot}$. **Bottom panel:** The ratio of the circular velocity $v_c = [GM_{tot}(< r)/r]^{0.5}$ at the stellar half-mass radius R_{eff} to that at $3R_{eff}$, which we use as a measure of mass concentration and steepness of the potential. The panel shows that discs in the ARTEMIS galaxies generally form when the mass is not very centrally concentrated $(v_c(R_{eff})/v_c(3R_{eff}) < 1)$, but the mass distribution becomes centrally concentrated after the disc forms. The former disc galaxies generally behave similarly to the present-day discs.

concentration increases both before *and* after the disc forms and is thus not a cause of disc spin-up.

Hopkins et al. (2023) also showed that a gas disc can form in a halo with z = 0 mass of $M_{200c} = 4 \times 10^{10} \,\mathrm{M}_{\odot}$ (their m11a object), which is well outside the range of $M_{200c}(t_{\rm spin}) = (6 \pm 3) \times 10^{11} \,\mathrm{M}_{\odot}$ found in the ARTEMIS simulations and the virial mass threshold expected for the hot halo formation. Nevertheless, this does not necessarily imply a discrepancy. Hopkins et al. (2023) focus on the formation of *gas* discs that may not necessarily lead to the formation of a stellar disc. Our definition of a disc, on the other hand, is based purely on stellar kinematics. Our findings are therefore not inconsistent with the formation of gas discs in dwarf galaxies. In fact, Hopkins et al. (2023) also find that *at the time of formation of gaseous discs* the circular velocity is decreasing with decreasing radius at $r < 3R_{\rm eff}$ and increases generally *after* disc formation due to the growth of central mass concentration. If stellar discs were also found to be present in



Figure 5. Like Fig. 4, but M_{200c} and the circular velocity ratio are plotted as functions of κ_{co} . The left-hand column shows present-day discs and the right column shows former discs. The spin-up threshold of $\kappa_{co} = 0.5$ is marked with a red dashed line. This confirms that spin-up occurs when the virial mass reaches the range $M_{200c} \approx (6 \pm 3) \times 10^{11}$, whereas the circular velocity ratios span a wide range of values when κ_{co} crosses 0.5. The behaviour is generally more chaotic for the former discs, in which κ_{co} exceeds 0.5 before decreasing again.

lower mass galaxies, this may indicate that a hot gaseous inner halo mediating gas accretion and inducing disc formation may form in different ways. It may form when the host halo has a regular mass concentration and reaches a threshold of $M_{200c}(t_{spin}) = (6 \pm 3) \times 10^{11} \,\mathrm{M_{\odot}}$ or it may form in haloes of lower virial mass if a strong central mass concentration capable of compressing gas to a sufficiently high temperature forms via some physical process.

These possibilities warrant further investigation. Our results, however, indicate that regular haloes with typical concentrations form discs when their central potential is not particularly steep.

4.3 Comparison to galaxies without discs

So far we have focused on the ARTEMIS galaxies that form discs (i.e. reach $\kappa_{co} \geq 0.5$) at some snapshot. We now compare these to the general properties of those that do not reach this threshold at any time. Note that only 10 ARTEMIS galaxies fail to become discs at any time, so the following analysis is based on a reasonably small sample.

In Fig. 6, we show the corotation parameter κ_{co} , virial mass M_{200c} , circular velocity ratio, and accreted stellar fraction versus lookback time $t_{\rm L}$. As in Figs 1 and 3, the coloured lines and bands correspond to the median and 16th–84th percentile ranges of the quantities as a function of $t_{\rm L}$ for the two samples. Galaxies with $\kappa_{co} \ge 0.5$ in any snapshot are shown in green and those without $\kappa_{co} \ge 0.5$ are shown in purple.



Figure 6. Various quantities against lookback time $t_{\rm L}$ for galaxies with discs ($\kappa_{\rm co} \ge 0.5$) in any snapshot (green) and those that never form discs ($\kappa_{\rm co} < 0.5$) in every snapshot, purple). From top to bottom, the panels show the corotation parameter $\kappa_{\rm co}$, virial mass M_{200c} , circular velocity ratio, and total accreted stellar fraction (as shown in Fig. 3). Apart from $\kappa_{\rm co}$ itself, the most striking difference between the two samples is the larger fraction of accreted stars at early times in the non-disc galaxies. This suggests that the failure to form a disc may be due to merger events.

The κ_{co} dependence shows the expected behaviour. The disc galaxies reach $\kappa_{co} \sim 0.5$ by $t_L \sim 8$ Gyr, with some decreasing below 0.5 later (these are the 'former discs'). The non-disc galaxies have significantly lower values of κ_{co} , never exceeding 0.5 by construction. The virial mass behaviour is, however, similar between the two samples. This suggests that while the virial mass at early times determines the epoch of disc formation (see Figs 2, 4, and 5), it does not dictate whether or not a disc forms. The same can be said of the circular velocity ratio, although the non-discs do have slightly higher values on average in most snapshots.

The most significant difference between the two samples can be seen in the accreted stellar fraction. At $t_{\rm L} \approx 10-12$ Gyr, the median accreted fraction in non-disc galaxies increases to more than double that in those that form discs, with the 84th percentile being much higher. This period corresponds to the time at which the discs are forming (top panel and Fig. 1). This is a strong indication that the failure to form a dominant disc may be due to massive mergers at $t_{\rm L} \sim 10$ Gyr.

5 SUMMARY AND CONCLUSIONS

We have used the ARTEMIS simulations of galaxies in MW-mass haloes to investigate the spin-up of discs. We have focused on the relations between the time of disc spin-up and the mass assembly histories of parent haloes, including the presence of a merger and associated kinematic signatures similar to the GSE merger in the MW. Our results and conclusions can be summarized as follows:

(i) Most galaxies with discs at z = 0 form their discs at higher metallicity than the MW. ARTEMIS galaxies with a GSE-like feature form discs earlier and at lower metallicities than those without, and are closer to the track of the median $v_{\phi} - [Fe/H]$ of the MW (see Fig. 1). We select six early spin-up galaxies that are similar to the MW (those with the highest median v_{ϕ} at [Fe/H] = -1), five of which have a GSE-like feature.

(ii) The distribution of stars before the disc forms is irregular without a well-defined disc or flattening (see Fig. A1). There is thus no evidence in simulations for the existence of an ancient metal-poor disc before the main disc spin-up.

(iii) There is a correlation between spin-up time $t_{\rm spin}$ and halo mass at a lookback time of $t_{\rm L} = 12$ Gyr: earlier spin-ups occur in galaxies with larger M_{200c} and $v_{\rm max}$ at $t_{\rm L} = 12$ Gyr (see top and middle panels of Fig. 2). In particular, the six early spin-up galaxies have the largest halo masses and $v_{\rm max}$ at that epoch and experience the earliest spin-ups ($t_{\rm spin} > 8$ Gyr).

(iv) The spin-up time also anticorrelates with the fraction of accreted stars at z = 0. Galaxies with early spin-ups and discs tend to have smaller accreted fractions than those with later spin-ups (bottom panel of Figs 2 and 3). We estimate the accreted stellar mass fraction in the MW, and find that only two ARTEMIS galaxies have comparable fractions. Both of these have MW-like spin-ups (at early times and low metallicities) and are massive at early times. Galaxies with a GSE-like feature also tend to have lower accreted fractions, supporting the conclusions of Dillamore et al. (2023) that these undergo fewer major mergers than those without.

(v) We show that discs in ARTEMIS galaxies form when their haloes reach masses of $M_{200c} \approx (6 \pm 3) \times 10^{11} \,\mathrm{M_{\odot}}$ (top panel of Fig. 4), similar to the mass at which galaxies are expected to form hot gaseous haloes and transition from the cold- to hot-mode accretion regime. We also show that the mass distribution becomes concentrated *after* the disc forms (see the bottom panel of Fig. 4 and associated discussion in Section 4.2), suggesting that it is not the cause of disc formation.

(vi) While the virial mass is related to the time at which the disc forms, it does not exclusively determine whether or not disc formation happens at all. The galaxies that never form discs have similar mass growth profiles to those that do, but a significantly larger proportion of their stellar populations are accreted, particularly at early times ($t_L \sim 10$ Gyr). This suggests that the absence of dominant disc formation may be linked to massive mergers occurring at these times.

It is clear that the MW experienced mass assembly and disc formation earlier than average for haloes of its mass. Our results show that the accretion histories of MW analogues are closely linked to the formation of their discs.

ACKNOWLEDGEMENTS

We thank the anonymous referee for a helpful report that has improved this manuscript. We are grateful to Vadim Semenov for useful comments on the draft of this paper. AMD thanks the Science and Technology Facilities Council (STFC) for a PhD studentship. This work was funded by UKRI grant 2604986. AK was supported by the National Science Foundation grants AST-1714658 and AST-1911111 and NASA ATP grant 80NSSC20K0512. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

DATA AVAILABILITY

Data from the ARTEMIS simulations may be shared on reasonable request to the corresponding author.

REFERENCES

- Abdurro'uf et al., 2022, ApJS, 259, 35
- Belokurov V., Kravtsov A., 2022, MNRAS, 514, 689
- Belokurov V., Erkal D., Evans N. W., Koposov S. E., Deason A. J., 2018, MNRAS, 478, 611
- Belokurov V., Sanders J. L., Fattahi A., Smith M. C., Deason A. J., Evans N. W., Grand R. J. J., 2020, MNRAS, 494, 3880
- Binney J., Tremaine S., 2008, Galactic Dynamics, 2nd edn. Princeton Univ. Press, Princeton, NJ
- Conroy C. et al., 2022, preprint (arXiv:2204.02989)
- Correa C. A., Schaye J., Clauwens B., Bower R. G., Crain R. A., Schaller M., Theuns T., Thob A. C. R., 2017, MNRAS, 472, L45
- Crain R. A. et al., 2015, MNRAS, 450, 1937
- Deason A. J., Belokurov V., Sanders J. L., 2019, MNRAS, 490, 3426
- Dekel A., Birnboim Y., 2006, MNRAS, 368, 2
- Dekel A. et al., 2009, Nature, 457, 451
- Dekel A., Ginzburg O., Jiang F., Freundlich J., Lapiner S., Ceverino D., Primack J., 2020, MNRAS, 493, 4126
- De Silva G. M. et al., 2015, MNRAS, 449, 2604
- Dillamore A. M., Belokurov V., Font A. S., McCarthy I. G., 2022, MNRAS, 513, 1867
- Dillamore A. M., Belokurov V., Evans N. W., Font A. S., 2023, MNRAS, 519, L87
- Dolag K., Borgani S., Murante G., Springel V., 2009, MNRAS, 399, 497
- Fattahi A. et al., 2019, MNRAS, 484, 4471
- Ferreira L. et al., 2022, ApJ, 938, L2
- Ferreira L. et al., 2023, ApJ, 955, 94
- Font A. S. et al., 2020, MNRAS, 498, 1765
- Gaia Collaboration, 2016, A&A, 595, A1
- Grand R. J. J. et al., 2017, MNRAS, 467, 179
- Grand R. J. J. et al., 2020, MNRAS, 497, 1603

- Hafen Z. et al., 2022, MNRAS, 514, 5056
- Haywood M., 2001, MNRAS, 325, 1365
- Helmi A., Babusiaux C., Koppelman H. H., Massari D., Veljanoski J., Brown A. G. A., 2018, Nature, 563, 85
- Hopkins P. F. et al., 2023, MNRAS, 525, 2241
- Jacobs C. et al., 2023, ApJ, 948, L13
- Kassin S. A. et al., 2012, ApJ, 758, 106
- Kereš D., Katz N., Weinberg D. H., Davé R., 2005, MNRAS, 363, 2
- Kereš D., Katz N., Fardal M., Davé R., Weinberg D. H., 2009, MNRAS, 395, 160
- McCluskey F., Wetzel A., Loebman S. R., Moreno J., Faucher-Giguere C.-A., 2023, preprint (arXiv:2303.14210)
- McMillan P. J., 2017, MNRAS, 465, 76
- Majewski S. R. et al., 2017, AJ, 154, 94
- Massari D., Koppelman H. H., Helmi A., 2019, A&A, 630, L4
- Myeong G. C., Evans N. W., Belokurov V., Sanders J. L., Koposov S. E., 2018, ApJ, 863, L28
- Naidu R. P., Conroy C., Bonaca A., Johnson B. D., Ting Y.-S., Caldwell N., Zaritsky D., Cargile P. A., 2020, ApJ, 901, 48
- Nelson E. J. et al., 2023, ApJ, 948, L18
- Orkney M. D. A. et al., 2023, MNRAS, 525, 683
- Pedregosa F. et al., 2011, J. Mach. Learn. Res., 12, 2825
- Rix H.-W. et al., 2022, ApJ, 941, 45
- Robertson B. E. et al., 2023, ApJ, 942, L42
- Schaye J. et al., 2015, MNRAS, 446, 521
- Semenov V. A., Conroy C., Chandra V., Hernquist L., Nelson D., 2023a, preprint (arXiv:2306.09398)
- Semenov V. A., Conroy C., Chandra V., Hernquist L., Nelson D., 2023b, preprint (arXiv:2306.13125)
- Shen S., Mo H. J., White S. D. M., Blanton M. R., Kauffmann G., Voges W., Brinkmann J., Csabai I., 2003, MNRAS, 343, 978
- Springel V., 2005, MNRAS, 364, 1105
- Stern J. et al., 2021, ApJ, 911, 88
- VandenBerg D. A., Brogaard K., Leaman R., Casagrande L., 2013, ApJ, 775, 134
- Wisnioski E. et al., 2015, ApJ, 799, 209
- Wisnioski E. et al., 2019, ApJ, 886, 124

APPENDIX A: STAR FORMATION RATE DENSITY

In Fig. A1, we show the projected star formation rate density in each of the 18 present-day disc galaxies, from $t_{\rm L} = 12.4-0$ Gyr. Each row corresponds to a different galaxy, labelled on the righthand side. The z-axis corresponds to the angular momentum of the gas within r = 10 kpc in each snapshot, with the z-unit vector from the previous snapshot marked with a red arrow. The galaxies are ordered by t_{spin} , with the spin-up (marked with a red border) becoming earlier moving down the rows. The star formation is initially clumpy and irregular, with the angular momentum direction of the gas changing dramatically between subsequent snapshots. The stellar discs (parallel to the x-axis) can then be seen forming in the few snapshots prior to the spin-up time, after which star formation usually takes place only close to this plane. The relation between early-time mass and spin-up time shown in Fig. 2 can also be discerned, with rates of star formation in the first few snapshots being visibly lower in the galaxies with the latest spin-ups (top few rows)

Note that the presence of a disc (i.e. $\kappa_{co} > 0.5$) does not necessarily imply that star formation is occurring in a disc-like configuration. For example, the star formation in G26 and G32 at $t_L = 0$ is more dispersed. Conversely, star formation in a disc does not necessarily imply that $\kappa_{co} > 0.5$ (e.g. G24 and G27 before spin-up). In both scenarios, this may be a reflection of the fact that κ_{co} describes the overall state of all stars, whereas Fig. A1 only shows where star formation is actively ongoing. A disc may therefore be present but not star forming (e.g. G26 and G32), or actively forming stars but not yet dominant (e.g. G24 and G27). We also find in a few cases (including G24 and G27) that κ_{co} approaches close to 0.5 a few Gyr before it exceeds that value. Hence, their late spin-up times are partly due to their discs not becoming dominant enough until sometime after their initial formation.

	12.4	12.2	12.0	11.6	11.1	10.8	10.5	10.0	9.4	8.8	7.9	7.3	6.6	5.9	5.1	4.1	3.2	2.3	1.3	0.0	
den hen hen hen hen hen hen hen hen hen h	5 0 % **		- Sider -	1		•		N		1	1	1	1	1	1	1	1	4		1	G24
	5 0 5	S.	*				X		X				11		1	T			1		G40
	5	2 4	*			*		-	׼.		1	1				2	1	1	1	1	G27
	5 0 5	- dés		-						\		L		T.					1		G15
	505			\$	×.	1	1	Ľ.	\mathbf{X}	1	1		1				<u>.</u>	er i en Kenstern	1	1	G25
	5	-	*	*	de la	1		 	X	1		Ľ.	1	1			$\mathbf{\lambda}$		A		G26
	505	A.				1	1	1			4	T			49					1	G29
			A.				Ň		L		- 1				4					4	G18
	5 0 5				-		1	1		1	X	and the second sec	C.d.			1					G42
	5 0 5	× 1	2	l.)	1	1	1		ana <mark>r</mark> as Nasar	-	-						1	1	G01
	5 05			1		-	-	1			H	N			<u> </u>						G16
	5	A.	*			1	1	Ţ		÷.		-	X		\ 	1	. /	1	1	\ \	G32
	5 0 5		1			1		1	7-12	N.		est b Streft		1							G17
	5		Ĩ		I	1						1	1						25		G34
	5 0 5		1	1		A				<u>ale</u>	X			A.		¥.	-1				G23
	5 0 5				1	1	1	<u> </u>	1			1	1		T		-				G38
	5 0 5	1		-	1		1	L	1			1		-							G44
	505	1	1	4	1			~	Å						en <mark>klas</mark>			1		1	G19
	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5 S	$r \left[\mathrm{kpc} \right]$	-50 5 2]	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	-50 5	

 t_L [Gyr]

Figure A1. Star formation rate density in the inner 10 kpc of all 18 galaxies in our sample (rows) across many snapshots (columns). The *z*-axis in each panel is parallel to the instantaneous angular momentum of gas particles within a radius of r = 10 kpc. For each galaxy, the snapshot at the spin-up time t_{spin} is marked with a red border, with t_{spin} becoming earlier from top to bottom. The red arrows mark the direction of the *z*-axis in the previous snapshot, which indicates how much this direction changes between consecutive snapshots. As in the other figures, the red (blue) colours of the galaxy IDs correspond to (non-)GSE galaxies, and the early spin-up galaxies highlighted in Figs 1 and 2 are underlined.

This paper has been typeset from a TEX/LATEX file prepared by the author.

© The Author(s) 2023. Published by Oxford University Press on behalf of Royal Astronomical Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.