

Effects of endurance exercise training on left ventricular structure in healthy adults: a systematic review and meta-analysis

Barbara N. Morrison ¹, Keith George ², Elizabeth Kreiter³, Duncan Dixon³, Lyndon Rebello¹, Raffaele J. Massarotto¹, and Anita T. Cote ^{1*}

¹School of Human Kinetics, Trinity Western University, 22500 University Drive, Langley, BC, Canada V2Y 1Y1; ²Research and Enterprise, Liverpool John Moores University, Egerton Court, 2 Rodney Street, Liverpool L1 2UA, UK; and ³Reference and Information Literacy, Trinity Western University, 22500 University Drive, Langley, BC, V2Y 1Y1, Canada

Received 15 August 2022; revised 24 January 2023; accepted 26 January 2023; online publish-ahead-of-print 31 January 2023

See the editorial comment for this article ‘Endurance training: what is the expected left ventricle remodelling?’, by F. Bandera, <https://doi.org/10.1093/eurjpc/zwad109>.

Aims

To determine the impact of endurance training (ET) interventions on left ventricular (LV) chamber size, wall thickness, and mass in healthy adults.

Methods and results

Electronic databases including CINAHL, MEDLINE, PsycINFO, SPORTDiscus, Cochrane library, and EBM Reviews were searched up to 4 January 2022. Criteria for inclusion were healthy females and/or males (>18 years), ET intervention for ≥ 2 weeks, and studies reporting pre- and post-training LV structural parameters. A random-effects meta-analysis with heterogeneity, publication bias, and sensitivity analysis was used to determine the effects of ET on LV mass (LVM) and diastolic measures of interventricular septum thickness (IVSd), posterior wall thickness (PWTd), and LV diameter (LVDd). Meta-regression was performed on mediating factors (age, sex, training protocols) to assess their effects on LV structure. Eighty-two studies met inclusion criteria ($n = 1908$; 19–82 years, 33% female). There was a significant increase in LVM, PWTd, IVSd, and LVDd following ET [standardized mean difference (SMD) = 0.444, 95% confidence interval (CI): 0.361, 0.527; $P < 0.001$; SMD = 0.234, 95% CI: 0.159, 0.309; $P < 0.001$; SMD = 0.237, 95% CI: 0.159, 0.316; $P < 0.001$; SMD = 0.249, 95% CI: 0.173, 0.324; $P < 0.001$, respectively]. Trained status, training type, and age were the only mediating factors for change in LVM, where previously trained, mixed-type training, young (18–35 years), and middle-aged (36–55 years) individuals had the greatest change compared with untrained, interval-type training, and older individuals (>55 years). A significant increase in wall thickness was observed in males, with a similar augmentation of LVDd in males and females. Trained individuals elicited an increase in all LV structures and ET involving mixed-type training and rowing and swimming modalities conferred the greatest increase in PWTd and LVDd.

Conclusion

Left ventricular structure is significantly increased following ET. Males, young and trained individuals, and ET interventions involving mixed training regimes elicit the greatest changes in LV structure.

Lay summary

Heart structure significantly increases the following endurance training (ET) ≥ 2 weeks.

- Changes in heart structure were most prominent in males, who are young (18–35 years), already trained, and following concurrent continuous and interval training.
- Changes in heart size were not shown in older individuals (>55 years) compared with young and middle-aged individuals.
- While both males and females similarly increase their cavity size and heart mass, sex differences were revealed for wall thickness where significant increases were seen in males but not females.

Keywords

Endurance exercise training • Left ventricle • Athletes • Sex differences

* Corresponding author. Tel: +1 604 513 2121 ext. 3726. Email: anita.cote@twu.ca

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Introduction

Participation in regular endurance training (ET) is associated with structural and functional cardiac adaptations that, in turn, improve cardiorespiratory fitness needed for higher peak oxygen uptake required for enhancing athletic performance.^{1,2} These structural adaptations include increases in left ventricular (LV) chamber size, wall thickness, and mass (LVM) and require careful consideration when differentiating between normal physiological adaptation and potential pathology.¹ Previous cross-sectional literature has demonstrated that endurance-trained individuals have an increased LVM compared with age- and sex-matched controls,^{3–8} but that these physiological cardiac adaptations can be mediated by gender, training volume, and sport type.^{5,6,8,9} Evidence suggests that male athletes,^{5,8} those with greater training volume,^{5,8} and those that participate in high dynamic–low static sports and high dynamic–high static sports have the largest LVM.^{6,10} Cross-sectional data are, however, limited by selection bias and cannot determine the cause-and-effect relationship between ET and cardiac phenotype. This requires intervention-based, longitudinal training studies. There is a growing evidence-base available suggesting that longitudinal training studies can lead to cardiac adaptation, at least qualitatively similar to that observed in cross-sectional, athlete-control studies. For example, a 1-year intensive ET programme in sedentary individuals resulted in a significant increase in LVM, approaching similar levels to that seen in elite endurance-trained athletes.³

Although research has demonstrated increases in LV chamber size, wall thickness, and mass in response to ET, these changes have been highly variable, study-to-study, and there has been limited systematic evaluation of potential factors that may mediate the cardiac structure ET response. Confounding factors such as age, sex, ET history (untrained or previously trained), modality of ET (i.e. running, cycling, rowing, swimming, or combination of more than one modality), or type of training programme (continuous, interval, or mixture of both types) require further study and the complexity and breadth of potential confounders lends itself to a systematic review and meta-analysis. Therefore, the aim of the current systematic review was to determine the impact of prospective ET interventions on LV structure in healthy adults. The secondary purpose was to assess how variation in potential confounders may mediate change in LVM. We hypothesized that LV structure would be significantly greater following ET, with males, younger, trained individuals, and ET interventions involving high dynamic–high static modalities eliciting greater changes in LVM and other structural variables.

Methods

The review is reported in accordance with the PERSiST (implementing PRISMA in Exercise, Rehabilitation, Sport medicine and Sports science) consensus statement.¹¹ This review is registered with Prospero (CRD42017072090).

Eligibility criteria

Studies included had to meet the following inclusion criteria: (i) ET interventions in healthy men and women ≥ 18 years; (ii) LV structural parameters reported prior to and after an ET intervention; and (iii) ≥ 2 weeks in duration. Cross-sectional studies comparing training methods across sporting disciplines and observational longitudinal studies, including multi-day races, were excluded, as were conference abstracts, editorial, and review papers. Studies that included clinical populations were excluded; however, if there was a healthy control population that underwent ET, the control population results were included. Individually or in any combination, ET that included running, aerobics, cycling, swimming, rowing, soccer, or other ET modality of at least moderate intensity were eligible. Studies that were solely strength-based, or low-intensity ET, were excluded. Interventions that reported an element of strength training within their moderate or higher

intensity ET intervention were included. There were no restrictions on the setting in which the ET were performed (i.e. classes, supervised individual, or group training). Furthermore, exercise could not be combined with pharmacological interventions. In the event the same study had multiple publications, the report that included the greatest sample size of the eligible outcome measures was included. In studies that assessed more than two time-points during the intervention, the pre-training and last training block measurements were included in the analysis.^{3,12–22} In instances where studies only broadly discussed the training phases (i.e. pre-season training, competitive season, maintenance phases), and did not prospectively prescribe the training programme, they were excluded as observational studies.

Search strategy

The systematic search was conducted by a librarian (E.K.) who had expertise in systematic reviews and verified by a senior librarian (D.D.). The databases searched included CINAHL, MEDLINE, PsycINFO, SPORTDiscus, Cochrane (Database of Systematic Reviews, Clinical Answers, Central Register of Controlled Trials, Methodology Register), and EBM Reviews (Health Technology Assessment, NHS Economic Evaluation Database, ACP Journal Club, Database of Abstracts of Reviews of Effects). These databases were last searched on 4 January 2022 and yielded 3481 results. The search strategy combined the two concepts of intervention and outcome, investigating the impact of ET (intervention) on LV structure (outcome). Keyword searches, with the phrase and truncation indications, were performed in the Title and Abstract fields. Subject headings were employed where available. See [Supplementary material online, Table S1](#) for the full list of keywords and subject headings used in each database, according to their interface (EBSCO or Ovid). The English-language filter was the only limiter applied to the search results; results were not limited according to date or publication type.

Selection process

A total of 3481 articles were identified through database searching with an additional 13 articles identified through reference lists and hand searching. The articles were imported into a reference management software (Endnote X9) where duplicates were removed. The remaining 1631 results were imported into a systematic review management software (COVIDENCE, Melbourne, Australia) for screening so that investigators could work independently. Two reviewers (N.S., A.R.) reviewed all study titles and abstracts and excluded studies according to the pre-specified criteria with a unanimous result required to exclude a study ([Figure 1](#)). Two investigators (A.T.C., B.N.M.) reviewed the full-text articles and assessed them according to the search criteria for inclusion. Disagreement in any of the stages was resolved by consensus between the two reviewers and a third reviewer (K.G.) if consensus could not be reached.

Data extraction

One investigator (B.N.M.) extracted data using a standardized data extraction form agreed upon by three reviewers (A.T.C., K.G., B.N.M.). A second investigator (L.R.) reviewed the data for accuracy, and any inconsistencies were resolved by discussion between these investigators, and a third investigator (A.T.C.) solved any disagreements. General study information (authors, publication year, sample size), participant information (age, sex), imaging modality (echocardiography, echo; cardiac magnetic resonance imaging, CMR), LV structural parameters (interventricular septum thickness in end-diastole, IVSd; posterior wall thickness in end-diastole, PWTd; LV end-diastolic diameter, LVDd), and LV functional parameters [($\text{VO}_{2\text{max}}$, end-diastolic volume, EDV; end-systolic volume, ESV; and stroke volume, SV), ET intervention features: trained status, modality (running/aerobics, cycling, swimming, rowing, soccer, or any combination of these), type of training, inclusion of strength training, duration of the study (weeks), frequency of training (days/week), time per session (h); total weekly hours] were extracted from included studies. Participants' trained status were classified as *untrained* (individuals who had not previously exercised regularly, on an organized basis or competed in previous competitions) and *trained* (individuals who followed a regular training programme or previously competed, including those who were at the beginning an ET programme (i.e. college matriculation, training for a marathon) and athletes who went through a period of detraining prior to the ET intervention). The type of training included: *continuous* (CONT) training defined as ET that performed with a




Figure 1 PRISMA flow diagram of the systematic process in article selection.

continuous intensity throughout that did not involve rest periods, *interval (INT)* training that included alternating bouts of high intensity with rest periods, and *mixed (MIX)* training that included a combination of these, either within the same session or during separate training sessions. Studies assessing more than one independent group were extracted as individual interventions for the analyses. Where full-text articles could not be found, or if data were unavailable from tables or the results section, the authors of the study in question were contacted by email. If the authors could not be contacted, the study was excluded. In instances where the training details were not reported, these studies were left out of the respective sub-analysis.

Data synthesis and analysis

Meta-analysis

The raw data extracted from the studies were transformed into effect sizes and were calculated as standardized mean differences (SMD) between pre- and post-training values using sample size and paired P -values with a meta-analysis software, using a random-effects model (Comprehensive Meta-Analysis V3). This model was selected due to different imaging techniques used to assess functional and structural variables as well as differences in the estimation equations to derive LVM across the studies. Absolute values for all variables of interest were the preferred value for the meta-analysis. Indexed values were used if absolute values or body surface area (BSA, pre- and post-measurement) were not reported. Since the model calculates the difference between pre- and post-values, the inclusion of the indexed values in this manner was justified. In cases where exact P -values were not reported, significant levels of $P < 0.05$, $P < 0.01$, and $P < 0.001$ were reported as 0.05, 0.01, and 0.001, respectively, and when reported as non-significant or $P > 0.05$, the values were reported as

0.999. An overall meta-analysis was conducted for each of the LV structural outcomes (LVM, LVDd, IVSd, PWTD) and additional meta-analyses were performed independently for each moderator variable to investigate their effect on LV structures. The planned moderators to be assessed were sex, age group [young (18–35 years), middle-aged (36–55 years), or older (>55 years)], training status, type of training, and mode of training. A meta-regression was conducted to ascertain if any effect moderator variable influenced the SMD in LV structures.

Heterogeneity among studies was assessed using I^2 statistics (the percentage of total variation between studies due to heterogeneity rather than by chance) and classified as low, moderate, and high at <50, 51–75, and >75%, respectively.²³ Relative influence of each study on the SMD was assessed by omitting one study at a time for sensitivity analysis ('one study removed analysis'). Publication bias was investigated by funnel plots for a visual inspection of asymmetry and statistically assessed using Egger's test and the trim-and-fill method.^{24,25} The National Heart, Lung, and Blood Institute (NHLBI) quality assessment tool for pre-post studies with no control group to assess for potential flaws in study methods including sources of bias, sampling, confounding variables, study power, and other relevant factors.²⁶ Each study was judged as 'good', 'fair', or 'poor' quality based on ratings from 11 items included in the tool. The 12th item was not included in this assessment as it pertained to group-level interventions (e.g. a whole hospital, a community) which was not relevant in the current review. Globally, a good study had the least risk of bias and was considered valid. A fair study is prone to some bias, but insufficient to invalidate its findings, varying in its strengths and weaknesses. A poor-quality study has a high risk of bias and is considered invalid. If a study was rated 'poor', it was removed from the analysis to determine its influence on the SMD. If the removal of the poor-rated study did not change the overall outcome of the

SMD when the study was removed, the study remained in the analysis. The quality assessment was performed by two reviewers (B.N.M. and R.J.M.), with a third reviewer (A.T.C.) involved in cases of uncertainty.

Descriptive analysis

The pre- and post-values for structural and functional values were reported as estimated pooled means and were calculated using a weighted sample dependent upon the sample size of each study [pooled average = $((n_1 \times \text{mean}_1) + (n_2 \times \text{mean}_2) \dots (n_k \times \text{mean}_k)) / \text{sum of all samples}$] and pooled standard deviations were calculated as:

$$S_{\text{pooled}} = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2 + \dots + (n_k - 1)s_k^2}{n_1 + n_2 + \dots + n_k - k}}$$

When the SD was not reported, it was estimated from the standard error of the mean, 95% confidence interval (CI), or IQR as suggested in the Cochrane handbook.²⁷ In situations where provided measures were not able to be converted to SD, they were excluded from the pooled means table and the sample size and number of studies were adapted accordingly. In instances where the relative $\text{VO}_{2\text{max}}$ (mL/kg/min) was not provided, it was estimated by dividing the group absolute mean VO_2 (L/min) by the group mean body weight (L/min/kg \times 1000). In studies where the SV was not reported, it was calculated using the following equation $\text{SV} = \text{LVEDV} - \text{LVESV}$. Per cent change was reported in consideration of different measurement techniques and modalities and calculated as $((\text{post} - \text{pre}) / \text{pre}) \times 100$ for all LV structural variables (LVM, IVSd, PWTd, LVDd) and their mediating factors (i.e. sex, age group, training status, type of training, mode of training). Independent samples t-test or one-way ANOVA was used to determine differences between groups with the Tukey *post hoc* analysis. The Pearson correlation was used to establish associations between training status and protocols with changes in LVM. Statistical significance was considered statistically significant at a *P*-value of <0.05 . Statistical analyses were performed using SPSS software Version 27.0 (IBM Corp., Armonk, NY, USA).

Results

Study selection and characteristics

The PRISMA flow diagram (Figure 1) illustrates the process of article selection, removal, and inclusion. Eighty-two studies satisfied the inclusion criteria ($n = 1908$; 19–82 years, 33% female).^{3,12–22,28–97} Baseline characteristics for the included studies are reported in Table 1. There were 19 studies that included more than one population. Specifically, 8 studies reported more than one type^{16,43,44,51,53,62,65} or mode³¹ of ET intervention and 11 studies reported more than one population type (i.e. sex,^{19,60,63} age,⁶⁹ trained status,^{14,21,50,54} menopause status,⁴¹ genotype³⁰) for a total of 105 analyses. Of these studies, 15 were randomized to either the ET or control group.^{28,31,36,37,48,51,53,63,66,67,76,77,89,96} Left ventricular structural characteristics were assessed before and after ET using echo or CMR. Table 2 summarizes the ET interventions. The average duration (range) of the ET interventions was 16.3 weeks (2–52 weeks), 4 sessions per week (1–8), and 4.8 h per week (0.2–32.5), with the majority of ET sessions between 30 and 60 min in duration. There were 11 studies that reported an element of strength training within their ET intervention.^{18,32,39,57,61,72,73,75,78,91,93} Trained individuals engaged in a significantly greater number of hours per week compared with untrained individuals (14.3 ± 8.8 vs. 3.1 ± 2.2 ; $P < 0.001$, respectively), with similar length of ET programs (18.9 ± 11.9 vs. 15.9 ± 12.3 weeks; $P = 0.378$). Table 3 summarizes pooled means and % change for before and after ET intervention for all structural and functional parameters in untrained and trained individuals. Figure 2A–E illustrates the per cent change for all LV structural variables and their mediating factors.

Overall ET had a high tolerability, with three participants reporting the training to be burdensome and 19 reporting musculoskeletal

injuries out of 10 studies. No studies reported major adverse outcomes; however, there were four cardiac abnormalities identified (i.e. three dysrhythmias and one left bundle branch block) in two studies. There were 127 (7%) dropouts amongst all studies, although 48 studies did not report adherence.

Effect of endurance training on left ventricular structure

Meta-analysis

The results of all meta-analyses including the overall SMD for LV structures and their independent moderators, heterogeneity, and publication bias are reported in Supplementary material online, S2–S5. The meta-analyses revealed a significant increase in all LV structures (LVM: SMD = 0.444; $P < 0.001$, IVSd: SMD = 0.237; $P < 0.001$, PWTd: SMD = 0.234; $P < 0.001$, LVDd: SMD = 0.249; $P < 0.001$) (see Supplementary material online, S9–S11), with the greatest increase in LVM (Figure 3). Left ventricular mass and LVDd increased in both males and females (LVM: SMD = 0.498; $P < 0.001$ and SMD = 0.280; $P = 0.004$; LVDd: SMD = 0.216; $P < 0.001$ and SMD = 0.245; $P = 0.003$, respectively), whereas PWTd and IVSd only increased in males. Egger's test revealed publication bias for LVM ($P = 0.001$) and LVDd ($P = 0.013$). Using the Trim and Fill method in those with publication bias, the SMD decreased slightly for LVM (0.368, 95% CI: 0.279, 0.458) and LVDd (0.188, 95% CI: 0.107, 0.269). The heterogeneity was low to moderate in all studies. None of the studies had high ($>75\%$) heterogeneity. Each of the included studies were of good ($n = 33$), fair ($n = 47$), and poor ($n = 2$) quality as indicated by the NHLBI quality assessment and agreed upon by both reviewers. The removal of the two studies with 'poor' quality did not elicit a significant change in the SMD for LV structural parameters; therefore, these studies remained in the analyses. Sensitivity analyses using the one study removed protocol revealed that the omission of any single study showed no change in the overall statistical significance, indicating that the presented results are statistically robust.

Moderator analysis

Several moderator variables influenced the SMD of the LV structures. The meta-regression bubble plots are reported in Supplementary material online S6–S8. LVM: There was a significant association between age group and LVM ($P = 0.0408$), with young and middle-aged individuals ($B = 0.2836$, $P = 0.0185$; $B = 0.3444$, $P = 0.0256$, respectively) conferring the greatest change in LVM compared with older individuals. Trained individuals conferred a greater change in LVM ($B = 0.2778$, $P = 0.0110$) than untrained and mixed training ($B = 0.2976$, $P = 0.0335$) elicited the greatest increase in LVM compared with continuous- and interval-type training. IVSd: Sex influenced the SMD of IVSd, with males demonstrating the greatest change in IVSd compared with females ($B = 0.2981$, $P = 0.0053$). Training status also demonstrated a significant association, with those that were trained ($B = 0.2362$, $P = 0.0261$) conferring a greater change than untrained individuals. PWTd: There was a significant association between sex and SMD of PWTd, with males conferring the greatest change in PWTd ($B = 0.1954$, $P = 0.0443$). There was also a significant association between trained status ($P = 0.0010$), training type ($P = 0.0492$), and mode of exercise ($P = 0.0046$) on SMD of PWTd. Those that were trained ($B = 0.3162$, $P = 0.0010$) had a greater change compared with those that were untrained, mixed, and continuous training regimes elicited the greatest change ($B = 0.2650$, $P = 0.0263$; $B = 0.2404$, $P = 0.0215$, respectively) compared with interval training, and swimming, rowing, and running ($B = 1.4014$, $P = 0.0093$; $B = 0.3084$, $P = 0.0340$; $B = 0.2831$, $P = 0.0022$, respectively) conferred the greatest change compared with cycling. LVDd: Those that were trained ($B = 0.2344$, $P = 0.0328$) conferred a greater increase compared with the untrained. Of the training types, only those that

Table 1 Baseline characteristics of journal articles meeting eligibility criteria

Study (author, year)	n	Control (n)	Female (%)	Age [mean \pm SD or (range)]	Population	VO _{2max} [mL/min/kg] (mean \pm SD or (range))	Imaging modality (dimensions included)
Untrained							
Adams_1981 ²⁸	22	9 ^a	0	22 (18–25)	College students (no endurance training > 3 months in last 5 years)	48.6 \pm 6.0	Echo (IVSd, PWTd, LVDd)
Aksakai_2013 ²⁹	34	n/a	0	22 \pm 2	No history of prior exercise exposure	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Alves_2009_1, 2, 3, 4 ³⁰	83	n/a	0	27 \pm 1	Brazilian policeman	47–50	Echo (LVM, IVSd, PWTd, LVDd)
Andersen_2010_A ³¹	18	10 ^a	100	37 \pm 8 ^b	Sedentary (no physical training for at least 2 years)	35.5 \pm 5.9	Echo (IVSd, PWTd, LVDd)
Andersen_2010_B ⁴	19	10 ^a	100	37 \pm 8 ^b	Sedentary (no physical training for at least 2 years)	32.5 \pm 4.6	Echo (IVSd, PWTd, LVDd)
Arbab-Zadeh_2014 ³	12	n/a	42	29 \pm 6	Untrained (< 30 min/day, < 3 \times /week regularly using either dynamic or static exercise)	40.3 \pm 5.5	CMR (LVM)
Bates_2013 ³³	10	n/a	40	39 \pm 12	Untrained	27.6 \pm 6.3	CMR (LVM)
Boone_2014 ³⁵	9	n/a	0	27 \pm 3	Moderately active (< 3 \times /week of any training)	53	Echo (LVM, IVSd, LVDd)
Camargo_2008 ³⁶	6	7 ^a	0	29 \pm 4	Healthy (no regular physical training in previous year, 35–42 VO _{2peak} (mL/min/kg))	38.1 \pm 2	CMR (LVM)
Cornelissen_2011 ³⁷	16	n/a	44	59 (55–71) ^b	Sedentary	21.8 \pm 0.8	CMR (LVM)
Cox_1986 ³⁸	11	5 ^a	55	23 \pm 1	Sedentary, inactive (no formal training for at least 3 months prior)	40.8 \pm 2.1	Echo (LVM, IVSd, PWTd, LVDd)
Dart_1992 ¹²	10	n/a	50	20–30	Sedentary	38	Echo (LVM, IVSd, PWTd, LVDd)
DeMaria_1978 ⁴⁰	24	n/a	46	26	Sacramento Police Academy	35.5 \pm 1.7	Echo (LVM, IVSd, PWTd, LVDd)
Egelund_2017_1 ⁴¹	36	n/a	100	49 \pm 2	Late pre-menopausal, sedentary (<2 h of physical training/week during previous 2 years, < 40 VO ₂ (mL/min/kg))	30.1 \pm 4.4	Echo (LVM, IVSd, PWTd, LVDd)
Egelund_2017_2 ⁴¹	37	n/a	100	53 \pm 3	Early post-menopausal, sedentary (<2 h of physical training/week during previous 2 years, < 40 VO ₂ (mL/min/kg))	30.4 \pm 3.3	Echo (LVM, IVSd, PWTd, LVDd)
Ehsani_1991 ¹³	10	n/a	0	64 \pm 3	Sedentary	29.6 \pm 4.1	Echo (LVM, PWTd, LVDd)
Esfandiari_2014_A ⁴³	8	n/a	0	25 \pm 3	Untrained (<2 h of <6 METs)	39.5 \pm 7.1	Echo (LVM, IVSd, PWTd, LVDd)
Esfandiari_2014_B ⁴³	8	n/a	0	26 \pm 5	Untrained (<2 h of <6 METs)	39.9 \pm 5.9	Echo (LVM, IVSd, PWTd, LVDd)
Eskelinen_2016_A ⁴⁴	14	n/a	0	47 \pm 9	Untrained (< 2 \times /week, no active training background, < 40 VO ₂ (mL/min/kg))	34.7 \pm 4.1	CMR (LVM)
Eskelinen_2016_B ⁴⁴	14	n/a	0	48 \pm 10	Untrained (< 2 \times /week, no active training background, < 40 VO ₂ (mL/min/kg))	33.7 \pm 3.9	CMR (LVM)
Fujimoto_2010 ⁴⁷	9	n/a	33	71 \pm 3	Sedentary (\leq 30 min, 3 \times /week)	22.8 \pm 3.4	CMR (LVM)
Fujimoto_2013 ⁴⁶	15	13 ^a	71	67 \pm 6	Sedentary (\leq 30 min, 3 \times /week)	23.0 \pm 4.7	CMR (LVM)
Grace_2018_1 ¹⁴	22	n/a	0	62 \pm 5	Sedentary (no participation in formal exercise training)	28.3	Echo (LVM, IVSd, PWTd, LVDd)

Continued

Table 1 Continued

Study (author, year)	n	Control (n)	Female (%)	Age [mean \pm SD or (range)]	Population	VO _{2max} [mL/min/kg] (mean \pm SD or (range))	Imaging modality (dimensions included)
Haykowsky_2005 ⁴⁸	8	8 ^a	100	66 \pm 3	No regular participation in aerobic or strength training	22	Echo (LVM, IVSd, PWTd, LVDd)
Hedman_2017 ⁴⁹	21	21 ^a	100	34 \pm 7	No regular participation in aerobic training within last year	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Hickson_1982_1, 2 ⁵⁰	15	n/a	40	29 \pm 5	Moderately active in recreational sports, but no training in last 6 months	39.6–45.2	Echo (LVM, IVSd, PWTd, LVDd)
Holloway_2018 ¹⁵	12	9 ^a	0	21 \pm 2	Healthy	42.5	Echo (IVSd, PWTd)
Huang_2019_1 ⁵¹	18	9 ^a	0	22 \pm 1	Sedentary (exercise <1/week, < 20 min)	20.4 \pm 1.4	Echo (LVM, IVSd, LVDd)
Huang_2019_2 ⁵¹	18	9 ^a	0	22 \pm 0	Sedentary (exercise <1/week, < 20 min)	21.0 \pm 1.0	Echo (LVM, IVSd, LVDd)
Hulke_2012_1 ⁵²	42	n/a	100	20 \pm 2	Healthy (DNR physical activity criteria)	37.4 \pm 3.3	Echo (LVM, IVSd, PWTd, LVDd)
Hulke_2012_2 ⁵²	43	n/a	0	20 \pm 1	Healthy (DNR physical activity criteria)	45.1 \pm 5.9	Echo (LVM, IVSd, PWTd, LVDd)
Hulke_2012_A ¹⁶	14	12	DNR	20 \pm 1	Healthy (DNR physical activity criteria)	34.1 \pm 6.1	Echo (LVM, IVSd, PWTd, LVDd)
Hulke_2012_B ¹⁶	13	12	DNR	20 \pm 1	Healthy (DNR physical activity criteria)	33.3 \pm 8.5	Echo (LVM, IVSd, PWTd, LVDd)
Hwang_2016_1 ⁵³	15	14 ^a	67	65 \pm 1	Sedentary	23.1 \pm 0.7	Echo (LVM, IVSd, PWTd, LVDd)
Hwang_2016_2 ⁵³	14	14 ^a	50	66 \pm 2	Sedentary	25.9 \pm 1.9	Echo (LVM, IVSd, PWTd, LVDd)
Kanakis_1982_1,2 ⁵⁴	12	n/a	58	23 (19–32)	Moderately active in recreational sports	35.9–42.5	Echo (IVSd, PWTd, LVDd)
Kiflom_2022 ⁵⁵	20	n/a	0	19–23	No regular participation in exercise training	n/a	Echo (IVSd, PWTd, LVDd)
Kivisto_2006 ⁵⁶	14	n/a	71	43 (23–58)	Sedentary	n/a	CMR (LVM)
Krzeminski_1989 ¹⁷	18	n/a	0	21 \pm 8	No active participation in sports	46.3 \pm 4.3	Echo (PWTd, LVDd)
Landry_1985 ⁵⁹	20	n/a	60	25 \pm 4	Sedentary (never trained)	37	Echo (LVM, IVSd, PWTd, LVDd)
Lane_2014_1 ⁶⁰	28	n/a	0	24 \pm 1	Sedentary (<30 min, < 1/week)	38	Echo (LVM)
Lane_2014_2 ⁵⁹	25	n/a	100	24 \pm 1	Sedentary (<30 min, < 1/week)	30	Echo (LVM)
Lusiani_1986 ⁶¹	13	9 ^a	0	20 \pm 7	Prior to beginning training program	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Mahdiabadi_2013_A ⁶²	10	n/a	0	21 \pm 2	Non-athletic	DNR	Echo (IVSd, PWTd, LVDd)
Mahdiabadi_2013_B ⁶²	10	n/a	0	21 \pm 1	Non-athletic	DNR	Echo (IVSd, PWTd, LVDd)
Marsh_1983 ⁶⁴	12	12	100	28 \pm 4	Running on unorganized basis <6 months – 4 years	46.4 \pm 3.6	Echo (LVM, IVSd, PWTd, LVDd)
Marsh_2021_1 ⁶³	46	COC	100	24 \pm 5	Untrained (< 150 min/week of organized exercise)	DNR	CMR (LVM)
Marsh_2021_2 ⁶³	26	COC	0	28 \pm 6	Untrained (< 150 min/week of organized exercise)	DNR	CMR (LVM)
Masoomah_2012 ⁹⁷	10	9	100	25 \pm 4	Non-athletic (no previous regular exercise training)	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Matsuo_2014_A ⁶⁵	14	n/a	0	26 \pm 7	Sedentary (no participation in regular activities for 1 year)	43.9 \pm 6.7	CMR (LVM)

Continued

Table 1 Continued

Study (author, year)	n	Control (n)	Female (%)	Age [mean \pm SD or (range)]	Population	VO _{2max} [mL/min/kg] (mean \pm SD or (range))	Imaging modality (dimensions included)
Matsuo_2014_B ⁶⁵	14	n/a	0	27 \pm 6	Sedentary (no participation in regular activities for 1 year)	41.9 \pm 5.6	CMR (LVM)
Matsuo_2014_C ⁶⁵	14	n/a	0	26 \pm 6	Sedentary (no participation in regular activities for 1 year)	42.0 \pm 6.8	CMR (LVM)
Morrison_1986 ⁶⁶	17	8	100	52 \pm 4	Untrained	27.3 \pm 4.6	Echo (IVSd, PWTd, LVDd)
O'Driscoll_2018 ⁶⁷	40	COC	0	21 \pm 2	Inactive/sedentary (<2.5 MET-h/week, > 8 h/day sitting time)	43.2 \pm 5.2	Echo (LVM, IVSd, PWTd, LVDd)
Park_2003 ⁶⁸	8	n/a	100	63 \pm 2	Sedentary	21.6 \pm 2.6	Echo (LVDd)
Perrault_1982_1 ⁶⁹	11	n/a	0	19 \pm 1	DNR	DNR	Echo (IVSd, PWTd, LVDd)
Perrault_1982_2 ⁶⁹	13	n/a	0	40 \pm 3	DNR	DNR	Echo (IVSd, PWTd, LVDd)
Pickering_1997 ⁷⁰	10	n/a	60	62 \pm 2	Sedentary	25.0 \pm 3.2	Echo (IVSd, PWTd, LVDd)
Rahimi_2018_1 ⁷¹	10	n/a	100	32 \pm 7	Non-athlete (no specific exercise activities)	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Rodrigues_2006 ⁷²	23	n/a	0	31 \pm 4	Sedentary	39 \pm 5	Echo (LVM, IVSd, PWTd, LVDd)
Rojek_2015 ⁷³	21	0	24	33 \pm 6	Starting to prepare for triathlon competition	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Rubal_1987_1 ¹⁹	10	n/a	100	19–31	No participation in regular programme of physical conditioning	38 \pm 8	Echo (LVM, IVSd, PWTd, LVDd)
Rubal_1987_2 ¹⁹	10	n/a	0	19–30	No participation in regular programme of physical conditioning	42 \pm 7	Echo (LVM, IVSd, PWTd, LVDd)
Saadatnia_2016 ⁹⁶	12	10 ^a	0	23 \pm 3	Untrained (currently no regular exercise), non-athlete	37.6	Echo (LVM, IVSd)
Sagiv_1989 ⁷⁴	20	n/a	0	67 \pm 4	Physically active in supervised aerobic programme for > 1 year, 3 \times /week	29.9	Echo (IVSd, PWTd)
Sayevand_2015 ⁷⁶	10	10 ^a	100	23 \pm 1	No regular training in the previous year	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Scharf_2015 ⁷⁷	42	n/a	0	44 \pm 5	Sedentary (<3 h/week of physical training)	DNR	CMR (LVM)
Shapiro_1983 ²⁰	15	15	0	26	No recent participation in sport	48.6 \pm 4.0	Echo (LVM, IVSd, PWTd, LVDd)
Skattebo_2020 ⁷⁹	12	n/a	42	29 \pm 6	Untrained (\leq 1 exercise training session/week during previous year)	44.5	Echo (LVM, IVSd, PWTd, LVDd)
Slordahl_2004 ⁸⁰	12	n/a	100	22 \pm 1	Sedentary (no formal training programme <3 months prior to the study)	42.6 \pm 2.9	Echo (LVM, IVSd, PWTd, LVDd)
Soto_2008 ⁸¹	12	n/a	50	69 \pm 6	Sedentary (\leq 30 min/day, <2 days/week)	23 \pm 3	Echo (LVM)
Spence_2011 ⁸²	10	n/a	0	27 \pm 1	<3 h/week of structured activity	45.8 \pm 1.6	CMR (LVM, IVSd, PWTd, LVDd)
Spina_1992 ⁸⁴	17	n/a	41	27 \pm 16	Sedentary (no regular exercise for 6 months prior to study)	40.9	Echo (LVDd)
Spina_1997 ⁸⁶	8	n/a	0	66 \pm 5	Sedentary (no regular exercise for 6 months prior to study)	28.7	Echo (LVM, IVSd, PWTd, LVDd)
Spina_2000 ⁸⁵	10	n/a	100	54 \pm 3	Sedentary (<2 \times /month of regular physical activity)	22.0 \pm 0.5	Echo (LVM, IVSd, PWTd, LVDd)

Continued

Table 1 Continued

Study (author, year)	n	Control (n)	Female (%)	Age [mean \pm SD or (range)]	Population	VO _{2max} [mL/min/kg] (mean \pm SD or (range))	Imaging modality (dimensions included)
Spina_2004 ⁸³	22	14 ^a	50	82 \pm 4	Sedentary	17 \pm 3	Echo (LVM, IVSd, PWTd, LVDd)
Vance_2014 ⁸⁷	19	22 ^a	0	22–37	Untrained (<2 h/week of aerobic training prior to enrolment, <50 mL/kg/min)	37.1 \pm 7.1	Echo (LVM, IVSd, PWTd, LVDd)
Vanhees_1992 ⁸⁸	27	n/a	0	38 \pm 2	Sedentary (<1 h/week of sport activities)	n/a	Echo (LVM, IVSd, PWTd, LVDd)
Wieling_1981_1 ²¹	9	17 ^d	0	20 \pm 2	Freshmen (active in high school sports, but no daily training programme)	58 \pm 8	Echo (IVSd, PWTd, LVDd)
Windecker_2002 ⁹²	8	n/a	0	36 \pm 5	Healthy (DNR prior physical activity levels)	46 \pm 6	Echo (LVM, IVSd, PWTd, LVDd)
Wolfe_1979 ²²	12	10 ^a	0	37	No engagement in regular exercise for at least 5 years prior	42.4 \pm 2.4	Echo (LVM, IVSd, PWTd, LVDd)
Wolfe_1992 ⁹³	7	4 ^a	100	19 \pm 2	Sedentary (no previous participation in endurance-type sport)	41.3 \pm 3.3	Echo (LVM, IVSd, PWTd, LVDd)
Younis_1987 ⁹⁴	19	n/a	0	19	Non-athlete, active	53 \pm 6	Echo (LVM, IVSd, PWTd, LVDd)
Trained							
Baggish_2009 ^{e32}	38	n/a	50	20 \pm 1	University students participating in official competitive athletics	DNR	Echo (LVM, IVSd, PWTd)
Bonaduce_1998 ³⁴	15	15 ^d	0	21 \pm 4	High level bicyclists competing in Italian amateur national teams for >3 years	62 \pm 4	Echo (LVM, IVSd, PWTd, LVDd)
D'Ascenzi_2015 ³⁹	91	n/a	0	23 \pm 6	Professional athletes (soccer, basketball, volleyball) competing in national or international level	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Ehansi_1978 ⁴²	8	n/a	13	17–19	University swim team (no regular exercise for 2–7 months before the beginning of training)	51.8 \pm 2.0	Echo (LVM, PWTd, LVDd)
Fagard_1983 ⁴⁵	12	12 ^d	0	24 \pm 1	Belgian cyclists (professional and amateur)	60.9 \pm 1.5	Echo (IVSd, PWTd, LVDd)
Grace_2018_2 ¹⁴	17	n/a	0	61 \pm 5	Masters athletes (national competitors, in triathlon, athletics, sprint cycling, racquet sports)	40.4	Echo (LVM, IVSd, PWTd, LVDd)
Kleinnibbelink_2021_1 ⁵⁷	19	n/a	0	26 \pm 4	Elite (Olympic) rowers	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Kleinnibbelink_2021_2 ⁵⁷	8	n/a	100	27 \pm 2	Elite (Olympic) rowers	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Lamont_1980 ⁵⁸	11	n/a	100	21 \pm 4	University swim team	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Naylor_2005 ¹⁸	22	12 ^a	23	20 \pm 3	Elite rowers	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Rahimi_2018_2 ⁷¹	10	n/a	100	31 \pm 9	Athlete (>5 \times /week, 2 h)	DNR	Echo (LVM)
Sareban_2018 ⁷⁵	15	n/a	0	20 \pm 3	National or international level rowers	67 \pm 6	Echo (LVM, LVDd)
Shah_2018 ⁷⁸	9	n/a	0	19 \pm 1	Newly matriculated, recruited members of varsity rowing	39.2	Echo (LVM, IVSd, PWTd, LVDd)
Venckunas_2006 ⁸⁹	12	11 ^a	0	25 \pm 7	Distance runners	DNR	Echo (LVM, IVSd, PWTd, LVDd)
Wasfy_2018 ⁹⁰	8	4 ^d	0	DNR	Newly matriculated, recruited members of varsity rowing	DNR	Echo (LVM)

Continued

Table 1 Continued

Study (author, year)	n	Control (n)	Female (%)	Age [mean \pm SD or (range)]	Population	VO _{2max} [mL/min/kg] (mean \pm SD or (range))	Imaging modality (dimensions included)
Wasfy_2019 ⁹¹	17	n/a	47	19 \pm 0	Collegiate swimmers	DNR	Echo (LVM, LVDd)
Wieling_1981_2 ²¹	14	17 ^d	0	22 \pm 6	Senior oarsmen (completed at least 1 rowing season)	62.1 \pm 7.6	Echo (IVSd, PWTd, LVDd)
Zilinski_2015 ⁹⁵	45	n/a	0	48 \pm 7	Recreational athletes	44.6 \pm 5.2	Echo (LVM, PWTd)

Articles assessing independent training groups were evaluated as individual studies and were distinguished by letters (A, B, C, D); articles assessing independent population groups were evaluated as individual studies and were distinguished by number (1, 2, 3, 4).COC, cross-over control study; CMR, cardiac magnetic resonance imaging; DNR, did not report; Echo, echocardiography.

^aUnderwent pre- and post-measurement.

^bOnly reported age as a group.

^cControl group in a clinical study that underwent intervention.

^dPre-measurement only.

^eOnly included group with parents who did not have hypertension.

participated in a mixed training regime elicited an increase in SMD of LVDd ($B = 0.3459$, $P = 0.0053$). Furthermore, only those that participated in swimming and rowing conferred an increase in SMD of LVDd ($B = 0.6527$, $P = 0.0056$; 0.3839 , $P = 0.0198$, respectively) compared with running.

Impact of training protocol and status on left ventricular structural change

The length of the ET intervention was not associated with an increase in LVM when all studies from 2 to 52 weeks were evaluated together ($r = 0.103$; $P = 0.349$). The dose–response association between the length of the training programme and LVM between sexes are presented in [Figure 4A and B](#). In the males, there was a significant positive association between the length of study and change in LVM ($r = 0.424$; $P < 0.005$), whereas in females, there was not ($r = 0.264$; $P = 0.274$). In the studies that reported VO_{2max}, all reported an increase in VO_{2max} except for one group of elite rowers.⁷⁵ The mean per cent change for VO_{2max} was higher in the untrained ($N = 69$) vs. the trained ($N = 8$) group (14.7 ± 7.6 and 8.6 ± 5.3 ; $P = 0.014$, respectively); however, the mean per cent change for LVM was higher in the trained ($N = 15$) vs. untrained ($N = 71$) group (19.7 ± 11.7 and 9.9 ± 10.1 ; $P = 0.007$, respectively). Additionally, pre-intervention VO_{2max} and LVM ($N = 63$) were positively associated with the change in LVM ($r = 0.285$, $P = 0.024$).

Discussion

This systematic review and meta-analysis synthesized data from 82 studies and 1908 participants examining the impact of ET intervention on LV structure in healthy men and women. The main findings were: (i) all LV structures increased in response to ET; (ii) LVM increased in young and middle-aged individuals but not in older individuals; (iii) a significant increase in wall thickness was observed in males, whereas an increase in LVM and LVDd was similarly augmented in males and females; (iv) trained status was associated with a greater increase in all LV structural parameters compared with untrained; (v) type of training influenced the adaptation of LVM, PWTd, and LVDd with mixed-type training eliciting the greatest change; and (vi) mode of training conferred an increase in PWTd and LVDd with rowing and swimming eliciting the greatest increase ([Figure 5](#)). These findings in the context of influencing factors are discussed below.

Age

Ageing is associated with numerous cardiac changes, such as an increase in wall thickness and a decline in ventricular contractility.^{3,98,99} Additionally, LVM and LVM index increases naturally with age in both males and females.¹⁰⁰ It is accepted that intense and long-lasting ET increases wall thickness beyond age-associated norms with a concomitant increase in cavity size in younger individuals; however, these changes have not been consistently observed in older individuals initiating an ET programme.^{47,98} In a study comparing the impact of 6 months of unsupervised training on cardiac structure in younger (mean age 29 ± 4) and older (mean age 46 ± 7) individuals preparing for a marathon, both groups demonstrated a similar increase in LVM; however, the younger group also saw an increase in LV cavity size.⁹⁸ Another study reported that in sedentary seniors (71 ± 3) compared with Masters athletes (68 ± 3), the sedentary seniors' baseline LVM was significantly less than the Masters athletes'.⁴⁷ After 1 year of progressive training, the sedentary seniors' LVM increased closer to the LVM of the Masters athletes; however, they did not see a change in their mass–volume ratio.⁴⁷ A potential reason for this could be that long and intense programmes required to elicit such changes are cautiously initiated in the previously inactive older population due to the concern of causing a musculoskeletal injury or an adverse cardiac event. However, it would be important to determine if high intensity and longer types of ET could elicit positive remodelling in these older age groups, safely to prevent age-associated cardiac structural and functional changes such as LV hypertrophy and LV stiffening.⁹⁹ In order to prevent the age-related, cross-linked advanced glycation end products in the vascular and LV walls and to elicit change in the number and volume of cardiac myocytes, exercise training needs to start early, and if started later in life, may be too late.¹⁰¹ Secondly, sedentary adults may need a longer period of time to see improvements in cardiac structure and function compared with their younger counterparts.¹⁰¹ In the present study, the average length of study in the older group was 24.1 ± 18.7 (6–52) weeks and the average hours of ET per week was 2.5 ± 1.3 h. Further research examining the age ET needs to be undertaken to attenuate age-related decline in cardiac structure and function and thereby cardiac risk as well as determining age-appropriate training programmes.

Sex differences

Many of the mechanisms that contribute to cardiac sex differences are unclear, including the magnitude and temporal sequence of cardiac

Table 2 Endurance training intervention details

Study (author, year)	Length of study (weeks)	Modality	Intensity	Progressive training	Type of training	Frequency (days/week)	Time per session, min	Total hours (h/week)
Untrained								
Adams_1981 ²⁸	11	RUN ^a	85% HR _{max}	DNR	CONT	5	50	4.2
Aksakal_2013 ²⁹	26	RUN	11 → ≥17 RPE (Borg scale)	↑ volume	CONT	DNR	60–240	DNR
Alves_2009_1,2,3,4 ³⁰	12	RUN	HR between VT → HR slightly above RCP	DNR	CONT	3	60	3
Andersen_2010_A ³¹	16	RUN	82% MHR	DNR	CONT	2–3	60	2.5 ^c
Andersen_2010_B ³¹	16	SOCCER	Football session	DNR	MIX	2–3	60	2.5 ^c
Arbab-Zadeh_2014 ³	52	RUN, CYCLE, or SWIM	5 training zones (base pace – intervals)	↑ duration	MIX	3–6	30–180	8 ^c
Bates_2013 ^{c33}	16	CYCLE	70–80% VO _{2max} and RPE 12–14 (Borg scale)	↑ intensity Reassess VO _{2max} at 8 wks, work adjusted to achieve THR	CONT	3	30	1.5
Boone_2014 ³⁵	6	CYCLE	4 intensity profiles from 50 to 130% Watts _{max}	DNR	MIX	3.5	60	3.5 ^c
Camargo_2008 ³⁶	12	RUN	70% MHR	DNR	CONT	3	35	1.5
Comelissen_2011 ³⁷	10	WLK, RUN, CYCLE, or STEP	66% HRR	DNR	CONT	3	60	3
Cox_1986 ³⁸	7	RUN, CYCLE	85–90% VO _{2peak} (cycle ergometer) – 100% VO _{2max} (5 × 11 W/week 5 min intervals)	Work adjusted to maintain 70% VO _{2max}	MIX	6	40	4
Dart_1992 ¹²	4	CYCLE	70% VO _{2max}	Work adjusted to maintain 70% HR _{max}	CONT	3	30	1.5
DeMaria_1978 ⁴⁰	11	RUN	70% HR _{max} (ED)	Work adjusted to maintain 70% HR _{max}	CONT	4	60	4
Egelund_2017_1,2 ⁴¹	12	CYCLE	60–96% VO _{2max}	↑ intensity	MIX	3	50	2.5
Ehsani_1991 ¹³	52	RUN, CYCLE	60–100% VO _{2max}	↑ VO _{2max} + supplement with intervals + re-measure VO _{2max} every 3 months	MIX	5	60	5
Esfandiari_2014_A ⁴³	2	CYCLE	95–100% VO _{2max}	↑ #intervals	INT	3	20 ^c	1 ^c
Esfandiari_2014_B ⁴³	2	CYCLE	65% VO _{2max}	↑ duration	CONT	3	105 ^c	5.25 ^c
Eskelinen_2016_A ⁴⁴	2	CYCLE	Maximal effort	↑ #intervals	INT	3	18.5 ^c	1
Eskelinen_2016_B ⁴⁴	2	CYCLE	60% peak workload	↑ duration	CONT	3	50 ^c	2.5 ^c
Fujimoto_2010 ⁴⁷	52	RUN	Below VT (~75–85 HR _{max}) – VT threshold (~85–90% HR _{max}) + intervals (5–10 beats below HR _{max})	↑ intensity + duration + intervals	MIX	5 ^c	25–60	
Fujimoto_2013 ⁴⁶	52	RUN	70–80% HR _{max}	↑ duration	CONT	3–4	25–40	2.3
Grace_2018_1,2 ¹⁴	6	CYCLE	50% PPO	↑ frequency	INT	1	18	0.3
Haykowsky_2005 ⁴⁸	12	CYCLE	60–80% HRR	↑ duration	CONT	3	15–42.5	2.1

Continued

Table 2 Continued

Study (author, year)	Length of study (weeks)	Modality	Intensity	Progressive training	Type of training	Frequency (days/week)	Time per session, min	Total hours (h/week)
Hedman_2017 ⁴⁹	12	CYCLE	Alternating increases in resistance and cadence until peak effort achieved	DNR	CONT	3	45–60	3
Hickson_1982_1,2 ⁵⁰	10	RUN, CYCLE	Up to 100% VO_{2max} by end of exercise session	↑ intensity to reach VO_{2max}	MIX	6	30–45	4
Holloway_2018 ¹⁵	6	CYCLE	90% VO_{2max}	Adjustment of training load	INT	3	13–15	0.7 ^c
Huang_2019_1 ⁵¹	6	CYCLE	60% VO_{2max}	Adjustment of training load	CONT	5	30	2.5
Huang_2019_2 ⁵¹	6	CYCLE	80% VO_{2max}	Adjustment of training load	INT	5	30	2.5
Hulke_2012_1,2 ⁵²	16	RUN	Somewhat hard (RPE scale)	DNR	CONT	8	60	8
Hulke_2012_A ¹⁶	20	RUN	59.5% HR_{max} (age-predicted)	DNR	CONT	8	25–30	4
Hulke_2012_B ¹⁶	20	RUN	74.9% HR_{max} (age-predicted)	DNR	CONT	8	25–30	4
Hwang_2016_1 ⁵³	8	ROW ^b	90% HR_{peak}	DNR	INT	4	25	1.7 ^c
Hwang_2016_2 ⁵³	8	ROW ^b	70% HR_{peak}	DNR	CONT	4	32	2.1 ^c
Kiffom_2022 ⁵⁵	12	RUN, CYCLE, SKIPPING, BALL games	50–75% MHR	DNR	CONT	3	60–80	3.5 ^c
Kivisto_2006 ⁵⁶	12	CYCLE	60–75% HR_{max}	↑ intensity ↑ duration	CONT	3–4	30	1.75 ^c
Krzeminski_1989 ¹⁷	13	CYCLE	50–75% VO_{2max}	↑ intensity	CONT	3	30	1.5
Landry_1985 ⁵⁹	20	CYCLE	60–85% HRR	↑ intensity	CONT	4–5	40–45	3.4
Lane_2014_1,2 ⁶⁰	8	CYCLE or ELL or RUN	>60–90% HR_{max}	DNR	CONT	3	30–60	2.25 ^c
Lusiani_1986 ⁶¹	11	RUN	DNR	DNR	MIX	3	DNR	DNR
Mahdiabadi_2013_A ⁶²	8	RUN	70% HR_{max}	DNR	CONT	3	45	2.25
Mahdiabadi_2013_B ⁶²	8	RUN	70% HR_{max}	DNR	INT	3	45	2.25
Masoomah_2012 ⁵⁷	8	RUN	65–80% HR_{max}	↑ intensity ↑ duration	CONT	3	16–30	1.2 ^c
Marsh_1983 ⁶⁴	DNR	RUN	70% HR_{max}	↑ distance Repeat examinations when mileage changed from 30 and 50 miles from baseline	CONT	5.5	DNR	DNR
Marsh_2021_1,2 ⁶³	12	RUN, CYCLE	60–90% HR_{max}	Periodized progressive macrocycle (x3) plan	MIX	3	60	3
Matsuo_2014_A ⁶⁵	8	CYCLE	120% VO_{2max}	Recalculation of workload (W) as per equation	INT	5	10	0.8
Matsuo_2014_B ⁶⁵	8	CYCLE	50–90% VO_{2max}	Recalculation of workload (W) as per equation	INT	5	18	1.5
Matsuo_2014_C ⁶⁵	8	CYCLE	60–65% VO_{2max}	Recalculation of workload (W) as per equation	CONT	5	45	3.75

Continued

Table 2 Continued

Study (author, year)	Length of study (weeks)	Modality	Intensity	Progressive training	Type of training	Frequency (days/week)	Time per session, min	Total hours (h/week)
Morrison_1986 ⁶⁶	35	RUN	65–75% HRR	Remeasurement of VO ₂ every 2 weeks	CONT	3	40	2
O'Driscoll_2018 ⁶⁷	2	CYCLE	100% effort	↑ intensity to maintain HR	INT	3	5.5	0.3
Park_2003 ⁶⁸	36	RUN, CYCLE	50–60% HRR	Reperform ET and adjust intensity	CONT	3	60	3
Perrault_1982_1 ⁶⁹	20	RUN	80% HR _{max} (ED)	Adjustment of training load to maintain intensity	CONT	3	30	1.5
Perrault_1982_2 ⁶⁹	20	CYCLE	80% HR _{max} (ED)	Adjustment of training load to maintain intensity	CONT	3	30	1.5
Pickering_1997 ⁷⁰	16	CYCLE	50% VO _{2max} – HR @ lactate threshold	↑ intensity	MIX	3	20–55	2.75 ^c
Rahimi_2018_1 ⁷¹	12	SWIM (water aerobics)	60–80% HR _{max}	↑ duration	CONT	3	30–60	2 ^c
Rodrigues_2006 ⁷²	26	RUN	HR _{AT} – HR 10% lower than RCP	↑ intensity	CONT	3	40–45	2.25
Rojek_2015 ⁷³	52	RUN, CYCLE, SWIM	DNR	DNR	DNR	5	> 90	11.2 ^c
Rubal_1987_1,2 ¹⁹	10	RUN	>70% HR _{max}	DNR	CONT	3.5	> 30	2 ^c
Saadatnia_2016 ⁷⁶	10	RUN	Maximal effort	↑ #intervals	INT	3	8–16	0.2 ^c
Sagiv_1989 ⁷⁴	12	RUN	70% VO _{2max}	Adjustment of training load to maintain intensity	CONT	3	30	1.5
Sayevand_2015 ⁷⁶	4	CYCLE	80% VO _{2max}	Adjustment of training power output	INT	3	40	2
Scharf_2015 ⁷⁷	16	RUN	65–90% HR _{max}	↑ frequency	MIX	2–4	DNR	DNR
Shapiro_1983 ²⁰	6	RUN	DNR	DNR	CONT	5	15–30	1.8 ^c
Skattebo_2020 ⁷⁹	10	CYCLE	70–90% HR _{peak}	↑ #intervals	MIX	3	36–60	0.8 ^c
Slordahl_2004 ⁸⁰	8	RUN	50–95% HR _{max}	DNR	INT	3	25	1.25 ^c
Soto_2008 ⁸¹	44	RUN, CYCLE	60–100% VO _{2max}	Re-measure VO ₂ every 3 months	MIX	4–5	60	4.5 ^c
Spence_2011 ⁸²	24	RUN	Individualized (based on VO _{2peak} and time trial performances) periodized programme, included 3 phases [preparatory phase (low–moderate intensity), specific phase (hill running, short intervals), competition phase (intensity maintained, volume reduced)]	Variations in intensity, volume	MIX	3	60	3

Continued

Table 2 Continued

Study (author, year)	Length of study (weeks)	Modality	Intensity	Progressive training	Type of training	Frequency (days/week)	Time per session, min	Total hours (h/week)
Spina_1992 ⁸⁴	12	RUN and CYCLE	50–95% VO _{2max}	Re-measure VO ₂ every 3 weeks	MIX	6	40–45	4.25 ^c
Spina_1997 ⁸⁶	36	RUN, CYCLE	60–95% VO _{2max}	↑ pace and power Re-measure VO ₂ every 3 months and adjust intensity	MIX	5	60	5
Spina_2000 ⁸⁵	48	RUN, CYCLE	60–95% VO _{2max}	Re-measure VO ₂ every 3 months and adjust intensity	MIX	5	60	5
Spina_2004 ⁸³	12	RUN, CYCLE, ROW	65–90% VO _{2max}	DNR	MIX	3	10–60	3 ^c
Vance_2014 ⁸⁷	16	RUN	Half-marathon training	DNR	DNR	DNR	DNR	DNR
Vanhees_1992 ⁸⁸	16	RUN, CYCLE	70% HRR	Encouraged to improve performance	CONT	3	60	3
Wieling_1981_1 ²¹	30	Non-specific endurance, ROW	Non-specific endurance, interval training, rowing training	DNR	MIX	DNR	DNR	9 ^c
Windecker_2002 ⁹²	22	RUN or CYCLE	80% VO _{2peak}	DNR	CONT	≥4	≥60	≥4
Wolfe_1979 ²	26	RUN ^a	60–80% HR _{max}	↑ intensity ↑ duration	CONT	4	10–30	2 ^c
Wolfe_1992 ⁹³	11	RUN	80–85% HR _{max}	↑ frequency ↑ duration	CONT	3–5	20–45	2.2 ^c
Younis_1987 ⁹⁴	26	SWIM, RUN, T/F, basic sports, ball games	≥70% HR _{max}	DNR	CONT	DNR	≥2	12
Trained								
Baggish_2009 ⁴³²	13	ROW	DNR (college training plan followed)	DNR (college training plan followed)	CONT	DNR (college training plan followed)	60–180	11.1 ^c
Bonaduce_1998 ³⁴	20	CYCLE, AER	DNR (followed training plan)	DNR (followed training plan)	CONT	7	180	21
D'Ascenzi_2015 ³⁹	18	SOC, VB, BB	Low training period (high volume/low intensity and sprinting); peak training (70–95% HR _{max})	DNR (followed training plan)	MIX	DNR	DNR	12–20
Ehansi_1978 ¹²	9	SWIM	DNR (followed swim team training plan)	DNR	DNR	6	180	12
Fagard_1983 ⁴⁵	DNR	CYCLE	DNR (followed competitive cycling training plan)	DNR	DNR	DNR	DNR	DNR
Kleinbittelink_2021_1,2 ⁵⁷	36	ROW	DNR (followed elite rowing training programme consisted of high intensity and endurance)	DNR	MIX	DNR	DNR	24–35

Continued

Table 2 Continued

Study (author, year)	Length of study (weeks)	Modality	Intensity	Progressive training	Type of training	Frequency (days/week)	Time per session, min	Total hours (h/week)
Lamont_1980 ⁵⁸	13	SWIM	DNR (followed swim training programme)	DNR	MIX	5	180–240	16
Naylor_2005 ¹⁸	26	ROW, AER	80–90% HR _{max}	DNR (followed rowing training programme)	CONT	6–7	270	29.3
Rahimi_2018_2 ⁷¹	12	SWIM (water aerobics)	60–80% HR _{max}	↑ intensity	CONT	3	30–60	2 ^c
Sareban_2018 ⁷⁵	11	ROW, RUN, CYCLE, SWIM	DNR (followed preparatory training period for elite athletes)	↑ duration	DNR	DNR	DNR	11.8 ^c
Shah_2018 ⁷⁸	13	ROW	DNR (collegiate rowing training programme)	DNR	CONT	5–6	60–180	11.9 ^c
Venckunas_2006 ⁸⁹	52	RUN	Increased training volume ~ 50% (20–125) in first 2 months and then continued at this training volume for remaining 10 months, participated in no more than once every 2 months, allowed to alternate mileage of sessions as desired	↑ volume	CONT	DNR	~120	DNR
Wasfy_2018 ⁹⁰	12	ROW	DNR (collegiate rowing training programme)	DNR	DNR	DNR	DNR	13
Wasfy_2019 ⁹¹	13	SWIM	DNR (followed collegiate training programme)	DNR	CONT	7	60–180	16.3
Wieling_1981_2 ²¹	30	ROW	DNR (followed collegiate rowing training programme)	DNR	MIX	DNR	DNR	12 ^c
Zilinski_2015 ⁹⁵	18	RUN	DNR (followed marathon training programme)	↑ distance	CONT	4–5	DNR	4

Articles assessing independent training groups were evaluated as individual studies and were distinguished by letters (A, B, C, D); articles assessing independent population groups were evaluated as individual studies and were distinguished by number (1, 2, 3, 4); AER, aerobics; BB, basketball; CYCLE, cycling; ED, exercise-determined; ELL, elliptical; ET, exercise test; MHR, maximum heart rate; NR, not reported; PPO, peak power output; RCP, respiratory compensation point; ROW, rowing; RUN, outdoor running or treadmill; SOC, soccer; STEP, stationary stepping; T/F, track and field; THR, target heart rate; VB, volleyball; VT, ventilatory threshold.

^aBicycle ergometer substituted if injury or inclement weather.

^bArm/leg non-weight-bearing ergometer.

^cMean if hours or frequency reported as range, and when mean hours reported for entire study population.

Table 3 Pooled means for left ventricular structure and function

LV parameter (units)	Descriptives (N analyses; N participants)	Pre-intervention (mean \pm SD)	Post-intervention (mean \pm SD)	% change \pm SD (range)
Untrained				
LV structure				
LVM (g)	57; 1035	139.7 \pm 25.6	149.4 \pm 27.0	7.7 \pm 7.3 (–4.6 to 32.1)
LVMi (g/m ²)	39; 824	77.7 \pm 12.6	83.8 \pm 14.0	8.3 \pm 8.3 (–5.13 to 31.7)
IVSd (mm)	65; 1144	9.0 \pm 1.4	9.3 \pm 1.2	3.6 \pm 6.3 (–19.5 to 20.3)
PWTd (mm)	63; 1139	8.8 \pm 1.1	9.2 \pm 1.1	4.6 \pm 6.3 (–5.7 to 28.9)
LVDd (mm)	64; 1125	47.9 \pm 1.9	48.8 \pm 1.8	2.0 \pm 2.8 (–6.1 to 13.2)
LV function				
VO _{2max} (mL/min/kg)	61; 1017	37.9 \pm 4.5	44.1 \pm 4.5	15.0 \pm 7.6 (0.4 to 33.8)
VO _{2max} (L/min)	7; 132	2.4 \pm 2.4	2.6 \pm 2.5	12.2 \pm 4.4 (8.8 to 21.5)
End-diastolic volume (mL)	33; 546	125.5 \pm 18.0	131.6 \pm 42.8	5.5 \pm 5.3 (–1.3 to 21.7)
End-systolic volume (mL)	31; 524	49.2 \pm 9.4	50.0 \pm 8.8	1.2 \pm 7.3 (–17.5 to 14.4)
Stroke volume (mL)	40; 702	76.3 \pm 12.4	82.7 \pm 12.4	9.0 \pm 9.8 (–7.2 to 33.5)
Trained				
LV structure				
LVM (g)	10; 229	196 \pm 43.8	217.7 \pm 50.8	11.2 \pm 5.5 (3.8–19.5)
LVMi (g/m ²)	13; 281	111.2 \pm 17.7	124.4 \pm 18.7	11.5 \pm 7.7 (–0.9 to 29.4)
IVSd (mm)	11; 240	10.0 \pm 1.6	10.8 \pm 2.5	7.6 \pm 4.7 (1.1 to 18.2)
PWTd (mm)	13; 293	9.7 \pm 1.1	10.4 \pm 1.1	6.3 \pm 5.8 (–2.0 to 20.0)
LVDd (mm)	13; 270	52.1 \pm 2.1	53.3 \pm 2.0	2.6 \pm 2.9 (–1.16 to 8.8)
LV function				
VO _{2max} (mL/min/kg)	7; 118	53.5 \pm 3.8	57.3 \pm 4.9	8.9 \pm 5.7 (0 to 15.5)
VO _{2max} (L/min)	1; 17	3.2 \pm 0.6	3.5 \pm 0.5	7.8
End-diastolic volume (mL)	6; 167	133 \pm 29.4	142.2 \pm 30.3	7.5 \pm 10.2 (–8.5 to 21.4)
End-systolic volume (mL)	5; 159	55.6 \pm 16.0	60.4 \pm 15.4	6.9 \pm 9.6 (–6.9 to 18.5)
Stroke volume (mL)	4; 61	89.4 \pm 17.0	92.5 \pm 18.0	3.1 \pm 9.3 (–9.6 to 12.1)

changes in response to ET. In the present study, both males and females demonstrated a significant increase in LVM; however, the SMD was greater in males than females. This is in-fitting with other reports where LVM was shown to increase in both males and females.¹⁰² Additionally, a positive correlation was observed between the length of the study programme and change in LVM in males, but not in females. Few studies have explored cardiac adaptation over time between the sexes. One study investigated sex differences in cardiac structure every 3 months over 1 year of an intensive ET intervention.¹⁰³ They found that both sexes demonstrated an increase in LVM (scaled to fat-free mass); however, the females demonstrated the greatest increase in LVM within the first 3 months of a 12-month training programme, compared with the males where LVM progressively increased throughout the entire 12 months, with the greatest increase in the first 6 months.¹⁰³ Additionally, they found that with an increasing exercise stimulus, LVM was markedly augmented in males, whereas in females, it was attenuated. Analysis of PWTd, IVSd, and LVDd revealed that only PWTd and IVSd significantly increased in males, whereas LVDd was similarly augmented in both males and females. Previous research in athletes demonstrates that when females LVDd is indexed for BSA, they exhibit a higher cavity size. When involved in dynamic sports, females predominantly exhibit eccentric LV hypertrophy, whereas a significant proportion of males in dynamic sports demonstrate concentric LV remodelling.¹⁰⁴ Some potential mechanisms contributing to these sex differences are that males have a higher circulating concentration of

testosterone and a higher density of myocardial testosterone receptors.¹⁰⁵ Additionally, a higher exercise-related systolic blood pressure in males may play a role in the development of LV hypertrophy.¹⁰⁶

Training status

Early studies comparing athletes to inactive individuals demonstrated that exercise leads to increased wall thickness, LVM, and LV cavity size.^{4,7,107} In the present analysis, this was supported by an increase in all LV structural parameters in both the untrained and trained groups. However, the trained group had a greater increase compared with the untrained group, suggesting that prior training and/or cardiorespiratory fitness is an important factor in cardiac adaptation. It can be speculated that previously trained individuals would have less capacity to increase their LVM due to already having a well-adapted heart. Possible explanations for this could be that athletes train at a higher overall volume of exercise eliciting a greater response, have a superior genetic endowment, and have a greater capacity to push their bodies to their limits.¹⁰⁷ Another factor could be the use of prohibited substances; however, this was not reported in any of the studies. Our data support the suggestion that a greater number of training hours per week may stimulate enhanced cardiac adaptation. Additionally, a higher baseline VO_{2max} was significantly correlated with greater changes in LVM, and the post-intervention VO_{2max} in the untrained group was less than that of the baseline VO_{2max} seen in the trained group. Thus, these findings may also represent




Figure 2 Change in left ventricular structure by (A) sex; (B) age group; (C) training type; (D) trained status; and (E) training mode.

in part a genetic component driving the enhanced adaptations seen in trained individuals.

Type of training

The type of training programme may influence the magnitude of cardiac remodelling. Previous research suggests that interval training involving shorter intervals and rest periods primarily improves the oxidative capacity of peripheral muscles, whereas continuous and aerobic interval training involving longer intervals targets central adaptations such as cardiac function.^{65,108} Previous studies have shown continuous and aerobic interval training is characterized by both an increase in LVDd and LV wall thickness, and consequently LVM.^{10,65} Increases in LVM have been

observed in short/sprint interval training and can occur with submaximal heart rates to allow optimal CA^{2+} cycling,¹⁰⁸ increased training volumes, high baseline cardiorespiratory fitness, and longer duration programmes (i.e. >3 months).^{56,65,82} In the present meta-analysis, mixed-type training elicited the greatest increase in LVM, PWTd, and LVDd, compared with interval- and continuous-type training. It is possible that changes in these structural parameters were not seen in interval-type training as they are often shorter duration as a time-efficient method to improve VO_{2max} , and, therefore, the length of study and training volume were not sufficient to induce these cardiac structural changes. Additionally, the present analysis did not report sprint and aerobic interval training separately; therefore, we were not able to determine cardiac structural changes specific to these two interval training types.




Figure 3 Forest plot of standardized mean difference between pre- and post-training left ventricular mass.




Figure 4 Dose–response association between the length of training programme and left ventricular mass in (A) males and (B) females.




Figure 5 Schematic diagram of the influence of moderator variables (sex, age group, training status, training type, mode of training) on left ventricular structures.

Mode of training

Left ventricular structural changes are influenced by the degree of static and dynamic components involved in the sport modality.¹⁰⁹ We found all modes of exercise increased LVM; however, the mode of exercise influenced the adaptation of PWTd and LVDd with swimming and rowing eliciting the greatest increase and cycling conferring the smallest change. Modalities that combine both dynamic and static components (i.e. rowing) require increases in cardiac output, heart rate, SV, systolic blood pressure, and mean arterial pressure, thereby generating the greatest changes in cavity dimensions and LV wall thickness.¹⁰⁹

Conversely, studies that are predominately dynamic (i.e. running) have lower afterload, and therefore will observe smaller LV structural changes.¹⁰⁹ Although there were only three studies that included swimming as the training stimulus, all elicited a significant increase in LVM, PWTd, and LVDd.^{42,58,91} Swimming is a unique sport, due to the physiological response of being immersed in water, and less gravitational forces being exerted on the swimmer.¹¹⁰ The horizontal position of swimming aids venous return, which is increased with the kicking of the legs. There is a concomitant increase in preload, increasing SV, and cardiac output which can generate an increase in both wall thickness and

LVDd.¹¹⁰ In studies that have compared swimming to other sporting disciplines, the results are unequivocal;^{4,111–116} however, when age and BSA are accounted for, swimming is associated with greater cardiac dimensions.^{115,116} Although there is a small sample size of swimmers in this analysis, the present findings confirm previous reports. Interestingly, cycling elicited the smallest changes across LV structures, which does not conform to previous literature where cyclists have been shown to elicit the greatest increases in LVM compared with other sport types.^{10,116–118} In the present analysis the majority (93%) of the studies utilising cycling as their training modality were in previously sedentary individuals, therefore, the disparity seen in this study compared to other studies that investigated the impact of cycling on LVM is likely due to the other studies including elite male and female cyclists.^{116,117} The stimulus in the cycling studies included in the present analysis was likely not intense or long enough to elicit large increases in LVM compared with that observed in previous studies.

Limitations

The present analysis has some limitations. First, it is important to note that these results pertain to healthy populations, and may not translate to clinical populations. Secondly, the impact of ethnicity could not be determined as only four studies reported ethnicity. Thirdly, while we compared the impact of ET in males and females from individual studies, there were very few within-study sex comparisons. Our overall female sample size was much smaller than that of males, and the Egger's test for publication bias was significant in males, warranting further studies exploring sex differences. Fourthly, a possible confounding factor seen in the trained group was that several of these studies reported strength training as part of their training programme for their specific sports which may have impacted LV structural parameters seen in this group. We are unable to isolate the impact of this additional strength training on cardiac adaptations within an ET intervention; however, a previous meta-analysis of cross-sectional studies reported no significant difference in LV structural parameters between athletes that are combined endurance-trained and strength-trained compared with those that are only endurance-trained.¹⁰ Fifthly, we found cycling interventions elicited the smallest change in LVM, contrary to previous cross-sectional studies where cyclists demonstrated greater LVM compared with other sports. This observation may be due to most cycling studies included in this meta-analysis were conducted with untrained participants, therefore may not represent the competitive cyclist response to an ET. Sixthly, swimming elicited the greatest increase in LVM; however there were only three studies and a very small sample size within each one; therefore, these results need to be interpreted with caution and confirmed in additional studies. Lastly, there was inconsistent reporting between studies with respect to whether they indexed for BSA, fat-free mass, or allometric scaling; therefore, we could not determine how the LV adaptations would be altered when indexed following changes in body mass or composition.

Conclusion

From this review, we confirm our hypothesis that LV structure is significantly increased following ET. Males, younger, trained individuals and ET interventions involving mixed training regimes elicit the greatest changes in LVM and other LV structural variables. Understanding these mediating factors during ET is important in developing effective training programmes and can help delineate between physiological adaptations to ET and potential pathology.

Authors' contributions

B.N.M. contributed to the design, screening, data extraction, and analysis, and wrote this review. K.G. contributed to design, data analysis,

and interpretation, and critically reviewed the paper. E.K. and D.D. contributed to the design, search strategy, and performed the literature search. L.R. and R.J.M. contributed to the screening and data analysis. A.T.C. conceived the project, contributed to the design, review, data analysis and interpretation, resolved conflicts, and critically reviewed the paper. All authors reviewed the manuscript and gave final approval for submission.

Supplementary material

Supplementary material is available at *European Journal of Preventive Cardiology*.

Acknowledgements

We would like to thank Natalie Szakun and Andrew Roberts for their assistance with the article screening process.

Funding

A.T.C. was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC, Discovery Grant 2018-06848) and the Canada Research Chairs Program (CRC) Program (950-231915). B.N.M. was supported through A.T.C.s CRC program.

Conflict of interest: None declared.

Data availability

The data sets generated from the review are available from the corresponding author upon reasonable requests.

References

- Hellsten Y, Nyberg M. Cardiovascular adaptations to exercise training. *Compr Physiol* 2015;**6**:1–32.
- Pressler ANJ. *Textbook of Sports and Exercise Cardiology*. Switzerland: Springer; 2020. p3–13.
- Arbab-Zadeh A, Perhonen M, Howden E, Peshock RM, Zhang R, Adams-Huet B, Haykowsky MJ, Levine BC. Cardiac remodeling in response to 1 year of intensive endurance training. *Circulation* 2014;**130**:2152–2161.
- Scharhag J, Schneider G, Urhausen A, Rochette V, Kramann B, Kindermann W. Athlete's heart: right and left ventricular mass and function in male endurance athletes and untrained individuals determined by magnetic resonance imaging. *J Am Coll Cardiol* 2002;**40**:1856–1863.
- Prakken NH, Velthuis BK, Teske AJ, Mosterd A, Mali WP, Cramer MJ. Cardiac MRI reference values for athletes and nonathletes corrected for body surface area, training hours/week and sex. *Eur J Cardiovasc Prev Rehabil* 2010;**17**:198–203.
- Luijckx T, Cramer MJ, Prakken NH, Buckens CF, Mosterd A, Rienks R, Backx FJ, Mali WP, Velthuis BK. Sport category is an important determinant of cardiac adaptation: an MRI study. *Br J Sports Med* 2012;**46**:1119–1124.
- Utomi V, Oxborough D, Whyte GP, Somaroo J, Sharma S, Shave R, Atkinson G, George K. Systematic review and meta-analysis of training mode, imaging modality and body size influences on the morphology and function of the male athlete's heart. *Heart (British Cardiac Society)* 2013;**99**:1727–1733.
- Cote AT, Bredin SS, Phillips AA, Koehle MS, Glier MB, Devlin AM, Warburton DER. Left ventricular mechanics and arterial-ventricular coupling following high-intensity interval exercise. *J Appl Physiol (1985)* 2013;**115**:1705–1713.
- Caselli S, Di Paolo FM, Pisicchio C, Di Pietro R, Quattrini FM, Di Giacinto B, Culasso F, Pelliccia A. Three-dimensional echocardiographic characterization of left ventricular remodeling in Olympic athletes. *Am J Cardiol* 2011;**108**:141–147.
- Pluim BM, Zwinderman AH, van der Laarse A, van der Wall EE. The athlete's heart. A meta-analysis of cardiac structure and function. *Circulation* 2000;**101**:336–344.
- Ardern CL, Büttner F, Andrade R, Weir A, Ashe MC, Holden S, Impellizzeri FM, Delahunt E, Dijkstra HP, Mathieson S, Rathleff MS, Reurink G, Sherrington C, Stamatakis E, Vicenzio B, Whittaker JL, Wright AA, Clarke M, Moher D, Page MJ, Khan KM, Winters M. Implementing the 27 PRISMA 2020 statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the PERSiST (implementing Prisma in Exercise, Rehabilitation, Sport medicine and SporTs science) guidance. *Br J Sports Med* 2022;**56**:175–195.

12. Dart AM, Meredith IT, Jennings GL. Effects of 4 weeks endurance training on cardiac left ventricular structure and function. *Clin Exp Pharmacol Physiol* 1992;**19**:777–783.
13. Ehsani AA, Ogawa T, Miller TR, Spina RJ, Jilka SM. Exercise training improves left ventricular systolic function in older men. *Circulation* 1991;**83**:96–103.
14. Grace F, Herbert P, Elliott AD, Richards J, Beaumont A, Sculthorpe NF. High intensity interval training (HIIT) improves resting blood pressure, metabolic (MET) capacity and heart rate reserve without compromising cardiac function in sedentary aging men. *Exp Gerontol* 2018;**109**:75–81.
15. Holloway K, Roche D, Angell P. Evaluating the progressive cardiovascular health benefits of short-term high-intensity interval training. *Eur J Appl Physiol* 2018;**118**: 2259–2268.
16. Hulke SM, Phatak MS, Vaidya YP. Cardiorespiratory response to aerobic exercise programs with different intensity: 20 weeks longitudinal study. *J Res Med Sci* 2012;**17**: 649–655.
17. Krzeminski K, Miskiewicz Z, Niewiadomski W, Nazar K, Kozłowski S. Effect of endurance training on cardiovascular response to static exercise performed with untrained muscles. *Int J Sports Med* 1989;**10**:363–367.
18. Naylor LH, Arnolda LF, Deague JA, Playford D, Maurogiovanni A, O'Driscoll G, Green DJ. Reduced ventricular flow propagation velocity in elite athletes is augmented with the resumption of exercise training. *J Physiol*. 2005;**563**:957–963.
19. Rubal BJ, Al-Muhailani AR, Rosentswieg J. Effects of physical conditioning on the heart size and wall thickness of college women. *Med Sci Sports Exerc* 1987;**19**:423–429.
20. Shapiro LM, Smith RG. Effect of training on left ventricular structure and function. An echocardiographic study. *Br Heart J* 1983;**50**:534–539.
21. Wieling W, Borghols EA, Hollander AP, Danner SA, Dunning AJ. Echocardiographic dimensions and maximal oxygen uptake in oarsmen during training. *Br Heart J* 1981; **46**:190–195.
22. Wolfe LA, Cunningham DA, Reznitzer PA, Nichol PM. Effects of endurance training on left ventricular dimensions in healthy men. *J Appl Physiol Respir Environ Exerc Physiol* 1979;**47**:207–212.
23. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**:557–560.
24. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**:629–634.
25. Shi L, Lin L. The trim-and-fill method for publication bias: practical guidelines and recommendations based on a large database of meta-analyses. *Medicine (Baltimore)* 2019; **98**:e15987.
26. National Heart, Lung, and Blood Institute. Study quality assessment tools 2021. <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools> (updated July 2021).
27. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, eds. *Cochrane Handbook for Systematic Reviews of Interventions*, version 6.3. Cochrane, 2022. www.training.cochrane.org/handbook (updated February 2022).
28. Adams TD, Yanowitz FG, Fisher AG, Ridges JD, Lovell K, Pryor TA. Noninvasive evaluation of exercise training in college-age men. *Circulation* 1981;**64**:958–965.
29. Aksakal E, Kurt M, Oztürk ME, Tanboğa IH, Kaya A, Nacar T, Semimli S, Gürletoş Y. The effect of incremental endurance exercise training on left ventricular mechanics: a prospective observational deformation imaging study. *Anadolu Kardiyol Derg* 2013;**13**: 432–438.
30. Alves GB, Oliveira EM, Alves CR, Rached HR, Mota GF, Pereira AC, Rondon MU, Hashimoto NY, Azevedo LF, Krieger JE, Negrão CE. Influence of angiotensinogen and angiotensin-converting enzyme polymorphisms on cardiac hypertrophy and improvement on maximal aerobic capacity caused by exercise training. *Eur J Cardiovasc Prev Rehabil* 2009;**16**:487–492.
31. Andersen LJ, Hansen PR, Søgaard P, Madsen JK, Bech J, Krstrup P. Improvement of systolic and diastolic heart function after physical training in sedentary women. *Scand J Med Sci Sports* 2010;**20**:50–57.
32. Baggish AL, Weiner RB, Yared K, Wang F, Kupperman E, Hutter AM Jr, Picard MH, Wood MJ. Impact of family hypertension history on exercise-induced cardiac remodeling. *Am J Cardiol*. 2009;**104**:101–106.
33. Bates MG, Newman JH, Jakovljevic DG, Hollingsworth KG, Alston CL, Zalewski P, Klawe JJ, Blamire AM, MacGowan GA, Keavney BD, Bourke JP, Schaefer A, McFarland R, Newton JL, Turnbull DM, Taylor RW, Trenell MI, Gorman GS. Defining cardiac adaptations and safety of endurance training in patients with m.3243A > G-related mitochondrial disease. *Int J Cardiol* 2013;**168**:3599–3608.
34. Bonaduce D, Petretta M, Cavallaro V, Apicella C, Iannicello A, Romano M, Breglio R, Marciano F. Intensive training and cardiac autonomic control in high level athletes. *Med Sci Sports Exerc* 1998;**30**:691–696.
35. Bonne TC, Doucende G, Flück D, Jacobs RA, Nordsborg NB, Robach P, Walther G, Lundby C. Phlebotomy eliminates the maximal cardiac output response to six weeks of exercise training. *Am J Physiol Regul Integr Comp Physiol* 2014;**306**:R752–R760.
36. Camargo MD, Stein R, Ribeiro JP, Schwartzman PR, Rizzatti MO, Schaan BD. Circuit weight training and cardiac morphology: a trial with magnetic resonance imaging. *Br J Sports Med* 2008;**42**:141–145; discussion 5.
37. Cornelissen VA, Goetschalckx K, Verheyden B, Aubert AE, Arnout J, Persu A, Rademakers F, Fagard RH. Effect of endurance training on blood pressure regulation, biomarkers and the heart in subjects at a higher age. *Scand J Med Sci Sports* 2011;**21**: 526–534.
38. Cox ML, Bennett JB 3rd, Dudley GA. Exercise training-induced alterations of cardiac morphology. *J Appl Physiol (1985)* 1986;**61**:926–931.
39. D'Ascenzi F, Pelliccia A, Alvino F, Solari M, Loffreno A, Cameli M, Focardi M, Bonifazi M, Mondillo S. Effects of training on LV strain in competitive athletes. *Heart* 2015;**101**: 1834–1839.
40. DeMaria AN, Neumann A, Lee G, Fowler W, Mason DT. Alterations in ventricular mass and performance induced by exercise training in man evaluated by echocardiography. *Circulation* 1978;**57**:237–244.
41. Egelund J, Jørgensen PG, Mandrup CM, Fritz-Hansen T, Stallknecht B, Bangsbo J, Nyberg M, Hellsten Y. Cardiac adaptations to high-intensity aerobic training in premenopausal and recent postmenopausal women: the Copenhagen women study. *J Am Heart Assoc* 2017;**6**:e005469.
42. Ehsani AA, Hagberg JM, Hickson RC. Rapid changes in left ventricular dimensions and mass in response to physical conditioning and deconditioning. *Am J Cardiol* 1978;**42**: 52–56.
43. Esfandiari S, Sasson Z, Goodman JM. Short-term high-intensity interval and continuous moderate-intensity training improve maximal aerobic power and diastolic filling during exercise. *Eur J Appl Physiol* 2014;**114**:331–343.
44. Eskelinen JJ, Heinonen I, Löyttyniemi E, Hakala J, Heiskanen MA, Motiani KK, Virtanen K, Pärkkä JP, Knuuti J, Hannukainen JC, Kalliokoski KK. Left ventricular vascular and metabolic adaptations to high-intensity interval and moderate intensity continuous training: a randomized trial in healthy middle-aged men. *J Physiol* 2016;**594**:7127–7140.
45. Fagard R, Aubert A, Lysens R, Staessen J, Vanhees L, Amery A. Noninvasive assessment of seasonal variations in cardiac structure and function in cyclists. *Circulation* 1983;**67**: 896–901.
46. Fujimoto N, Hastings JL, Carrick-Ranson G, Shafer KM, Shibata S, Bhella PS, Abdullah SM, Barkley KW, Adams-Huet B, Boyd KN, Livingston SA, Palmer D, Levine BD. Cardiovascular effects of 1 year of alagebrium and endurance exercise training in healthy older individuals. *Circ Heart Fail* 2013;**6**:1155–1164.
47. Fujimoto N, Prasad A, Hastings JL, Arbab-Zadeh A, Bhella PS, Shibata S, Palmer D, Levine BD. Cardiovascular effects of 1 year of progressive and vigorous exercise training in previously sedentary individuals older than 65 years of age. *Circulation* 2010;**122**: 1797–1805.
48. Haykowsky M, McGavock J, Vonder Muhll I, Koller M, Mandic S, Welsh R, Taylor D. Effect of exercise training on peak aerobic power, left ventricular morphology, and muscle strength in healthy older women. *J Gerontol A Biol Sci Med Sci* 2005;**60**:307–311.
49. Hedman K, Bjarnegård N, Länne T. Left ventricular adaptation to 12 weeks of indoor cycling at the gym in untrained females. *Int J Sports Med* 2017;**38**:653–658.
50. Hickson RC, Kanakis C Jr, Davis JR, Moore AM, Rich S. Reduced training duration effects on aerobic power, endurance, and cardiac growth. *J Appl Physiol Respir Environ Exerc Physiol* 1982;**53**:225–229.
51. Huang Y-C, Tsai H-H, Fu T-C, Hsu C-C, Wang J-S. High-intensity interval training improves left ventricular contractile function. *Med Sci Sports Exercise* 2019;**51**: 1420–1428.
52. Hulke S, Yuganti V, Ratta A. Effects of sixteen weeks exercise training on left ventricular dimensions and functions in young adults. *Natl J Physiol Pharm Pharmacol* 2012;**2**: 152–158.
53. Hwang C-L, Yoo J-K, Kim H-K, Hwang M-H, Handberg E-M, Petersen J-W, Christou DD. Novel all-extremity high-intensity interval training improves aerobic fitness, cardiac function and insulin resistance in healthy older adults. *Exp Gerontol* 2016;**82**: 112–119.
54. Kanakis C, Coehlo A, Hickson R. Left ventricular response to strenuous endurance training and reduced training frequencies. *J Cardiac Rehab* 1982;**2**:141–146.
55. Kiflom S, Enyew D, Ayalew A, Hailu A, Gebretensay M, Gebrehawerya G. Effect of aerobic exercise on physiological and left ventricular echocardiographic characteristics of non-athletic adult males in northern Ethiopia. *J Sports Med Phys Fitness* 2022;**62**: 288–295.
56. Kivistö S, Perhonen M, Holmström M, Lauerma K. Assessment of the effect of endurance training on left ventricular relaxation with magnetic resonance imaging. *Scand J Med Sci Sports* 2006;**16**:321–328.
57. Kleinnibbelink G, Panhuysen-Goedkoop N, Hulshof H, van Dijk A, George K, Somauroo J, Oxborough D, Thijssen DJH. Exercise training induces left- but not right-sided cardiac remodelling in Olympic rowers. *Int J Sports Med* 2022;**43**:151–160.

58. Lamont LS. Effects of training on echocardiographic dimensions and systolic time intervals in women swimmers. *J Sports Med Phys Fitness* 1980;**20**:397–404.
59. Landry F, Bouchard C, Dumessnil J. Cardiac dimension changes with endurance training. Indications of a genotype dependency. *JAMA* 1985;**254**:77–80.
60. Lane AD, Yan H, Ranadive SM, Kappus RM, Sun P, Cook MD, Harvey I, Woods J, Wilund K, Fernhall B. Sex differences in ventricular-vascular coupling following endurance training. *Eur J Appl Physiol* 2014;**114**:2597–2606.
61. Lusiani L, Ronsisvalle G, Bonanome A, Visona A, Castellani V, Macchia C, Pagnan A. Echocardiographic evaluation of the dimensions and systolic properties of the left ventricle in freshman athletes during physical training. *Eur Heart J* 1986;**7**:196–203.
62. Mahdiabadi J, Gaeni AA, Kazemi T, Mahdiabadi MA. The effect of aerobic continuous and interval training on left ventricular structure and function in male non-athletes. *Biol Sport* 2013;**30**:207–211.
63. Marsh CE, Thomas HJ, Naylor LH, Dembo LG, Green DJ. Sex differences in cardiac adaptation to distinct modalities of exercise: a cardiac magnetic resonance study. *Med Sci Sports Exerc* 2021;**53**:2543–2552.
64. Marsh R, Lapan D, Goldman S, Boyden T, Rotkis T, Stanforth P, Wilmore J. Effects of chronic, moderate endurance running on body composition and cardiac structure in women. *J Card Rehabil* 1983;**3**:208–212.
65. Matsuo T, Saotome K, Seino S, Shimojo N, Matsushita A, Iemitsu M, Ohshima H, Tanaka K, Mukai C. Effects of a low-volume aerobic-type interval exercise on VO₂max and cardiac mass. *Med Sci Sports Exerc* 2014;**46**:42–50.
66. Morrison DA, Boyden TW, Pamenter RW, Freund BJ, Stini WA, Harrington R, Wilmore JK. Effects of aerobic training on exercise tolerance and echocardiographic dimensions in untrained postmenopausal women. *Am Heart J* 1986;**112**:561–567.
67. O'Driscoll JM, Wright SM, Taylor KA, Coleman DA, Sharma R, Wiles JD. Cardiac autonomic and left ventricular mechanics following high intensity interval training: a randomized crossover controlled study. *J Appl Physiol* (1985) 2018;**125**:1030–1040.
68. Park S-K, Park J-H, Kwon Y-C, Yoon M-S, Kim C-S. The effect of long-term aerobic exercise on maximal oxygen consumption, left ventricular function and serum lipids in elderly women. *J Physiol Anthropol Appl Hum Sci* 2003;**22**:11–17.
69. Perrault H, Lajoie D, Péronnet F, Nadeau R, Tremblay G, Lebeau R. Left ventricular dimensions following training in young and middle-aged men. *Int J Sports Med* 1982;**3**:141–144.
70. Pickering GP, Fellmann N, Morio B, Ritz P, Amonchot A, Vermorel M, Coudert J. Effects of endurance training on the cardiovascular system and water compartments in elderly subjects. *J Appl Physiol* (1985) 1997;**83**:1300–1306.
71. Rahimi S, Khademvatani K, Zolfaghari MR. Association of circular Klotho and insulin-like growth factor 1 with cardiac hypertrophy indexes in athlete and non-athlete women following acute and chronic exercise. *Biochem Biophys Res Commun* 2018;**505**:448–452.
72. Rodrigues AC, de Melo Costa J, Alves GB, Ferreira da Silva D, Picard MH, Andrade JL, Mathias W Jr, Negrão CE. Left ventricular function after exercise training in young men. *Am J Cardiol* 2006;**97**:1089–1092.
73. Rojek A, Bialy D, Przewlocka-Kosmala M, Negrusz-Kawecka M, Mysiak A, Kosmala W. Biventricular response of the heart to endurance exercise training in previously untrained subjects. *Echocardiography* 2015;**32**:779–786.
74. Sagiv M, Fisher N, Yaniv A, Rudoy J. Effect of running versus isometric training programs on healthy elderly at rest. *Gerontology* 1989;**35**:72–77.
75. Sareban M, Winkert K, Sperlich B, Berger MM, Niebauer J, Steinacker JM, Treff G. Speckle tracking-derived bi-atrial strain before and after eleven weeks of training in elite rowers. *Sci Rep* 2018;**8**:14300.
76. Sayevand Z, Mehrialvar Y, Heidari N, Gheeraat A. Comparison of high-intensity interval training and resistance training on cardiac structure. *Med Sport* 2015;**11**:2526–2531.
77. Scharf M, Schmid A, Kemmler W, von Stengel S, May MS, Wuest W, Achenbach S, Uder M, Lell MM. Myocardial adaptation to high-intensity (interval) training in previously untrained men with a longitudinal cardiovascular magnetic resonance imaging study (Running Study and Heart Trial). *Circul Cardiovasc Imaging* 2015;**8**:e002566.
78. Shah AB, Zilinski J, Brown MG, Neary JH, Weiner RB, Hutter AM, Apple FS, Picard MH, Januzzi JL, Baggish AL. Endurance exercise training attenuates natriuretic peptide release during maximal effort exercise: biochemical correlates of the "athlete's heart". *J Appl Physiol* (1985) 2018;**125**:1702–1709.
79. Skattebo Ø, Bjerring AW, Auensen M, Sarvi SI, Cumming KT, Capelli C, Hallén J. Blood volume expansion does not explain the increase in peak oxygen uptake induced by 10 weeks of endurance training. *Eur J Appl Physiol* 2020;**120**:985–999.
80. Slørdahl SA, Madslien VO, Støylen A, Kjos A, Helgerud J, Wisløff U. Atrioventricular plane displacement in untrained and trained females. *Med Sci Sports Exerc* 2004;**36**:1871–1875.
81. Soto PF, Herrero P, Schechtman KB, Waggoner AD, Baumstark JM, Ehsani AA, Gropler RJ. Exercise training impacts the myocardial metabolism of older individuals in a gender-specific manner. *Am J Physiol Heart Circ Physiol* 2008;**295**:H842–H850.
82. Spence AL, Naylor LH, Carter HH, Buck CL, Dembo L, Murray CP, Watson P, Oxborough D, George K, Green DJ. A prospective randomised longitudinal MRI study of left ventricular adaptation to endurance and resistance exercise training in humans. *J Physiol* 2011;**589**:5443–5452.
83. Spina RJ, Meyer TE, Peterson LR, Villareal DT, Rinder MR, Ehsani AA. Absence of left ventricular and arterial adaptations to exercise in octogenarians. *J Appl Physiol* (1985) 2004;**97**:1654–1659.
84. Spina RJ, Ogawa T, Coggan AR, Holloszy JO, Ehsani AA. Exercise training improves left ventricular contractile response to beta-adrenergic agonist. *J Appl Physiol* (1985) 1992;**72**:307–311.
85. Spina RJ, Rashid S, Dávila-Román VG, Ehsani AA. Adaptations in beta-adrenergic cardiovascular responses to training in older women. *J Appl Physiol* (1985) 2000;**89**:2300–2305.
86. Spina RJ, Turner MJ, Ehsani AA. Exercise training enhances cardiac function in response to an afterload stress in older men. *Am J Physiol* 1997;**272**:H995–1000.
87. Vance DD, Chen GL, Stoutenberg M, Myerburg RJ, Jacobs K, Nathanson L, Perry A, Seo D, Goldschmidt-Clermont PJ, Rampersaud E. Cardiac performance, biomarkers and gene expression studies in previously sedentary men participating in half-marathon training. *BMC Sports Sci Med Rehabil* 2014;**6**:6.
88. Vanhees L, Hespel P, Van Hoof R, Fagard R, Amery A. Effect of physical training on systemic and brachial artery haemodynamics in normal men. *Int J Sports Med* 1992;**13**:145–151.
89. Venckunas T, Stasiulis A, Raugaliene R. Concentric myocardial hypertrophy after one year of increased training volume in experienced distance runners. *Br J Sports Med* 2006;**40**:706–709.
90. Wasfy MM, Bibbo CF, Brown M, DeLuca JR, Wang F, Berkstresser B, Weiner RB, Lewis GD, Hutter AM, Picard MH, Di Carli MF, Baggish AL. Myocardial metabolism in endurance exercise-induced left ventricular hypertrophy. *JACC Cardiovasc Imaging* 2018;**11**:928–930.
91. Wasfy MM, Weiner RB, Wang F, Berkstresser B, Deluca J, Hutter AM Jr, Picard MH, Baggish AL. Myocardial adaptations to competitive swim training. *Med Sci Sports Exerc* 2019;**51**:1987–1994.
92. Windecker S, Allemann Y, Billinger M, Pohl T, Hutter D, Orsucci T, Blaga L, Meier B, Seiler C. Effect of endurance training on coronary artery size and function in healthy men: an invasive followup study. *Am J Physiol Heart Circ Physiol* 2002;**282**:H2216–H2223.
93. Wolfe LA, Laprade A, Burggraf GW, Norman R. Cardiac responses of young women to conditioning for a 10 kilometer race. *Int J Sports Med* 1992;**13**:384–389.
94. Younis L, Poelnitz A, Sturbois X, Detry J-M. Effects of exercise training on left ventricular function in healthy young men: a longitudinal echocardiographic study. *J Cardiopulm Rehabil* 1987;**7**:278–284.
95. Zilinski JL, Contursi ME, Isaacs SK, Deluca JR, Lewis GD, Weiner RB, Hutter AM Jr, d'Hemecourt PA, Troyanos C, Dyer KS, Baggish AL. Myocardial adaptations to recreational marathon training among middle-aged men. *Circ Cardiovasc Imaging* 2015;**8**:e002487.
96. Saadatnia A, Ebrahim K, Rashidlamir A. Echocardiographic evaluation of the effects of high-intensity interval training on cardiac morphology and function. *Arch Cardiovasc Imaging* 2016;**4**:5.
97. Masoomah H, Agha AH, Peeri M, Hajsadeghi S. Effect of endurance, resistance and concurrent training on the heart structure of the female university students. *Biol Sport* 2012;**29**:17–21.
98. Torlasco C, D'Silva A, Bhuya AN, Faini A, Augusto JB, Knott KD, Benedetti G, Jones S, Zalen JV, Scully P, Lobascio I, Parati G, Lloyd G, Hughes AD, Manisty CH, Sharma S, Moon JC. Age matters: differences in exercise-induced cardiovascular remodelling in young and middle aged healthy sedentary individuals. *Eur J Prev Cardiol* 2021;**28**:738–746.
99. Lakatta EG, Levy D. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part I: aging arteries: a "set up" for vascular disease. *Circulation* 2003;**107**:139–146.
100. Wooten SV, Moestl S, Chilibeck P, Alvero Cruz JR, Mittag U, Tank J, Tanaka H, Rittweger J, Hoffmann F. Age- and sex-differences in cardiac characteristics determined by echocardiography in masters athletes. *Front Physiol* 2020;**11**:630148.
101. Aronson D. Cross-linking of glycated collagen in the pathogenesis of arterial and myocardial stiffening of aging and diabetes. *J Hypertens* 2003;**21**:3–12.
102. Diaz-Canestro C, Montero D. The impact of sex on left ventricular cardiac adaptations to endurance training: a systematic review and meta-analysis. *Sports Med* 2020;**50**:1501–1513.
103. Howden EJ, Perhonen M, Peshock RM, Zhang R, Arbab-Zadeh A, Adams-Huet B, Levine BD. Females have a blunted cardiovascular response to one year of intensive supervised endurance training. *J Appl Physiol* (1985) 2015;**119**:37–46.

104. Finocchiaro G, Dhutia H, D'Silva A, Malhotra A, Steriotis A, Millar L, Prakash K, Narain R, Papadakis M, Sharma R, Sharma S. Effect of sex and sporting discipline on LV adaptation to exercise. *JACC Cardiovasc Imaging* 2017;**10**:965–972.
105. Svartberg J, von Muhlen D, Schirmer H, Barrett-Connor E, Sundfjord J, Jorde R. Association of endogenous testosterone with blood pressure and left ventricular mass in men. The Tromso study. *Eur J Endocrinol* 2004;**150**:65–71.
106. Zemva A, Rogel P. Gender differences in athlete's heart: association with 24-h blood pressure. A study of pairs in sport dancing. *Int J Cardiol* 2001;**77**:49–54.
107. Douglas PS. Citius, altius, fortius (the Olympic motto: swifter, higher, stronger). *J Am Coll Cardiol* 2004;**44**:150–151.
108. Wisløff U, Ellingsen Ø, Kemi OJ. High-intensity interval training to maximize cardiac benefits of exercise training? *Exerc Sport Sci Rev* 2009;**37**:139–146.
109. Mitchell JH, Haskell W, Snell P, Van Camp SP. Task force 8: classification of sports. *J Am Coll Cardiol* 2005;**45**:1364–1367.
110. Riding NR. The swimmer's heart—the cardiovascular influence across the spectrum. *Aspetar Sports Med* 2015;**4**:440–446.
111. Pelliccia A, Culasso F, Di Paolo FM, Accettura D, Cantore R, Castagna W, Ciacciarelli A, Costini G, Cuffari B, Drago E, Federici V, Gribaudo CG, Iacovelli G, Landolfi L, Menichetti G, Atzeni UO, Parisi A, Pizzi AR, Rosa M, Santelli F, Santilio F, Vagnini A, Casasco M, Di Luigi L. Prevalence of abnormal electrocardiograms in a large, unselected population undergoing pre-participation cardiovascular screening. *Eur Heart J* 2007;**28**:2006–2010.
112. Pelliccia A, Culasso F, Di Paolo FM, Maron BJ. Physiologic left ventricular cavity dilatation in elite athletes. *Ann Intern Med* 1999;**130**:23–31.
113. Pelliccia A, Sharma S, Gati S, Bäck M, Börjesson M, Caselli S, Collet JP, Corrado D, Drezner JA, Halle A, Hansen D, Heidbuchel H, Myers J, Niebauer J, Papadakis M, Piepoli MF, Prescott E, Roos-Hesselink JW, Graham SA, Taylor RS, Thompson PD, Tiberi M, Vanhees L, Wilhelm M. 2020 ESC guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J* 2021;**42**:17–96.
114. Sharma S, Maron BJ, Whyte G, Firoozi S, Elliott PM, McKenna WJ. Physiologic limits of left ventricular hypertrophy in elite junior athletes: relevance to differential diagnosis of athlete's heart and hypertrophic cardiomyopathy. *J Am College Cardiol* 2002;**40**:1431–1436.
115. Spirito P, Pelliccia A, Proschan MA, Granata M, Spataro A, Bellone P, Caselli G, Biffi A, Vecchio C, Maron BJ. Morphology of the "athlete's heart" assessed by echocardiography in 947 elite athletes representing 27 sports. *Am J Cardiol* 1994;**74**:802–806.
116. Whyte GP, George K, Sharma S, Firoozi S, Stephens N, Senior R, McKenna WJ. The upper limit of physiological cardiac hypertrophy in elite male and female athletes: the British experience. *Eur J Appl Physiol* 2004;**92**:592–597.
117. Pelliccia A, Maron BJ, Spataro A, Proschan MA, Spirito P. The upper limit of physiologic cardiac hypertrophy in highly trained elite athletes. *N Engl J Med* 1991;**324**:295–301.
118. Wundersitz DWT, Gordon BA, Lavie CJ, Nadurata V, Kingsley MIC. Impact of endurance exercise on the heart of cyclists: a systematic review and meta-analysis. *Progr Cardiovasc Dis* 2020;**63**:750–761.