

1 Low fidelity imitation of atypical biological kinematics in autism spectrum disorders is
2 modulated by self-generated selective attention
3

4 Spencer J. Hayes,^{1*} Matthew Andrew,¹ Digby Elliott,^{1,2} Emma Gowen,³ and Simon J.
5 Bennett¹
6

7 ¹Brain and Behaviour Laboratory,
8 Research Institute for Sport and Exercise Sciences,
9 Liverpool John Moores University,
10 Liverpool, UK
11

12 ²Department of Kinesiology,
13 McMaster University,
14 Ontario, Canada
15

16 ³Faculty of Life Sciences,
17 The University of Manchester,
18 Manchester, UK
19

20 ^{*}Corresponding author

21 Brain and Behaviour Laboratory, Faculty of Science, Liverpool John Moores University,
22 Byrom Street, Liverpool, L3 3AF, UK

23 Tel: +44 (0) 151 904 6237, Fax: +44 (0) 151 904 6284

24 s.hayes@ljmu.ac.uk
25

Abstract

We examined whether adults with autism had difficulty imitating atypical biological kinematics. To reduce the impact that higher-order processes have on imitation we used a non-human agent model to control social attention, and removed end-state target goals in half of the trials to minimise goal-directed attention. Findings showed that only neurotypical adults imitated atypical biological kinematics. Adults with autism did, however, become significantly more accurate at imitating movement time. This confirmed they engaged in the task, and that sensorimotor adaptation was self-regulated. The attentional bias to movement time suggests the attenuation in imitating kinematics might be a compensatory strategy due to deficits in lower-level visuomotor processes associated with self-other mapping, or selective attention modulated the processes that represent biological kinematics.

Key words: autism spectrum disorders, imitation, biological motion kinematics, attention

1 Introduction

2
3 Imitation is a powerful mechanism for learning new sensorimotor behaviours (e.g. throwing a
4 Frisbee) as well as for developing socio-cognitive skills such as rapport (Chartrand & Bargh,
5 1999) and affiliation (Lakin & Chartrand, 2003). One way humans acquire these behaviours
6 is by copying a novel movement displayed by another person. This process is defined as ‘true
7 imitation’ because an observer is required to copy the properties of human movement
8 (biological motion) after observing a model, rather than being able to merely reproduce the
9 movement using an already learned movement pattern based on previous experience (Byrne
10 & Russon, 1998). In the context of human movement, biological motion is the visual-sensory
11 information contained in a movement that describes a particular action (Johansson, 1973;
12 Kozlowski & Cutting, 1977). For example, a person can be judged to be walking based on
13 how the arms and legs move in relation to each other. Therefore, during ‘true imitation’
14 (henceforth imitation) attention is directed to the biological motion kinematics (joint
15 configurations; limb velocity) of the observed person/model. Over repeated observations and
16 physical attempts at imitating the model, a new sensorimotor pattern is represented and
17 refined based on the available afferent and efferent sensorimotor feedback.

18 The mechanism underpinning imitation combines higher-order cognitive/attention and
19 lower-level visuomotor processes (Bandura, 1977; Byrne & Russon, 1998; Heyes, 2001)
20 embedded within a system linking perception with action (Prinz, 1997). Although not fully
21 understood, individuals with autism spectrum disorders (henceforth autism) exhibit different
22 neuropsychological processes and behaviour during imitation compared to typically
23 developed individuals (Edwards, 2014; Hamilton, 2013; Vivanti & Hamilton, 2014;
24 Williams, Whiten, & Singh, 2004). That is, people with autism often imitate the end-state
25 goal (to reach a target) of an action (Bird, Leighton, Press, & Heyes, 2007; Hamilton,

Brindley, & Frith, 2007; Wild, Poliakoff, Jerrison, & Gowen, 2012), but show difficulties imitating the style (i.e. a gentle or harsh hand action) in which the movement goal is achieved (Hobson & Lee, 1999; Perra et al., 2008; Rogers, 1999; Rogers, Bennetto, McEvoy, & Pennington, 1996; Rogers, Hepburn, Stackhouse, & Wehner, 2003; Rogers & Pennington, 1991; Salowitz et al., 2013; Smith & Bryson, 1994).

Extending upon original work that used descriptive measures (Bernier, Dawson, Webb, & Murias, 2007; Rogers, et al., 1996; Vivanti, Nadig, Ozonoff, & Rogers, 2008), kinematic analysis has been used to determine what, if any, aspects of movement form (e.g., velocity; timing of peak velocity) are imitated (Stewart, McIntosh, & Williams, 2013; Wild, et al., 2012; Williams, Casey, Braadbaart, Culmer, & Mon-Williams, 2014). Specifically, participants in the study of Wild et al. (2012) observed a human model performing an upper-limb pointing movement that differed in speed, while context was manipulated so the model aimed to targets (dots on a table), or to end space (dots removed). The notion is that, when targets are removed from the environment, the imitator focuses their attention on imitating the model's movement (kinematics; velocity) as opposed to merely reaching the target (dot) goal. The imitation of the model's movement is thought to occur via direct lower-level visuomotor mapping (Heyes, 2001; Southgate & Hamilton, 2008) and is suggested to be compromised in autism (Edwards, 2014; Stewart, et al., 2013; Williams, et al., 2004). When targets are present, the goal is to aim at a target (an action goal), which occurs via goal-directed processes, and are less affected in autism (Hamilton, et al., 2007). The results from Wild et al. (2012) showed only control participants imitated the different speeds when targets were removed. The lack of scaling of movement speed exhibited in participants with autism was accompanied by less time spent smoothly pursuing the hand (with the eyes) and thus more shifts of gaze between the targets. It was suggested the shift in gaze, and thus attention away from the hand, may have modulated the amount of action-based biological motion

information extracted from the model (Vivanti, et al., 2008), which thereby influenced the imitation of movement speed.

Notwithstanding an attentional contribution, reduced imitation of kinematics in individuals with autism has been linked to lower-level visuomotor processes (Stewart, et al., 2013). For instance, imitation in a neurotypical control group was similar after observing a human or non-human model, thus indicating that top-down processes associated with social modulation (Cook & Bird, 2012; Spengler, Bird, & Brass, 2010; Wang & Hamilton, 2012) did not exert an influence on behaviour. However, the autism group exhibited greater path length error and action duration in both observation conditions, which was attributed to impaired lower-level visuomotor processes that compromised self-other mapping in the mirror system (Bernier, et al., 2007; Nishitani, Avikainen, & Hari, 2004; Williams et al., 2006; Williams, et al., 2004; Williams, Whiten, Suddendorf, & Perrett, 2001). These lower-level processes link action-observation to action-execution, and sub-serve imitation by mapping observed biological motion onto the motor system (Buccino et al., 2004; Di Dio et al., 2013; Iacoboni et al., 2001; Iacoboni et al., 1999).

Although previous work has provided novel contributions to understanding specific imitation deficits in autism by isolating the contribution of lower-level processes (Stewart, et al., 2013; Wild, et al., 2012), the examination of biological motion kinematics was undertaken by manipulating only the speed or amplitude of the modelled movement and by evaluating performance based on data from the whole imitation session. In terms of biological kinematics, the aforementioned context requires an imitator to scale an existing motor pattern (upper-limb movement) to meet new task demands (e.g., faster movement), but does not isolate whether the deficit is attributable to imitating specific lower-level properties (e.g., velocity) of biological motion kinematics. Furthermore, because ‘true imitation’ is an active process whereby a novel representation is developed and refined over repeated

observations, it might be the case that important information about imitation adaptation is masked by collapsing the analysis over all trials. An alternative approach that can reveal more about adaptation is to evaluate performance in the early and late stages of imitation, which is a typical in observation learning studies (Bird & Heyes, 2005; Byrne & Russon, 1998; Hayes, Ashford, & Bennett, 2008). To this end, we further examined imitation of biological motion in individuals with and without autism by employing a novel protocol that required participants to imitate movements that had distinctly different, but still biologically plausible, movement kinematics (Hayes, Roberts, Elliott, & Bennett, 2014). The experimental models displayed movement that had exactly the same spatial and temporal outcomes as a control model, but with a velocity profile of either typical or atypical kinematics. The atypical model ensured that an observer had to configure the sensorimotor system to represent the novel movement kinematics, as opposed to the typical model that could be achieved by rescaling an existing representation of a typical upper-limb aiming movement (Hayes, Timmis, & Bennett, 2009; Vivanti, et al., 2008). We controlled top-down influences by using a protocol that minimized social attention (Cook & Bird, 2012; Wang & Hamilton, 2012) by presenting a non-human agent model (white-dot) with limited social context. To control for visual attention towards end-state target goal-directed features of the task environment (Vivanti, et al., 2008; Wild, et al., 2012), we only displayed target goals in half of the imitation trials in order to encourage attention towards the trajectory of the model. Finally, to examine adaption we examined performance at the early and late phases of imitation.

Method

Volunteers

Fifteen typical control participants (14 male; 1 female) and 15 participants with autism (14 male; 1 female) volunteered for the study. The volunteers with autism were recruited from an autistic society in North West England, and the University of Manchester, UK. The volunteers were provided with a participant information sheet and selected if they consented to be part of the study. The typical control participants were recruited from Liverpool John Moores University, UK. All participants had normal or corrected-to-normal vision and were screened via self-report for the following exclusion criteria: dyspraxia, dyslexia, epilepsy and other neurological or psychiatric conditions. The participants with autism had a diagnosis of autism, Asperger's syndrome or autism spectrum disorder by an independent clinician. Diagnosis was confirmed by a researcher trained (with research-reliability status) in the administration of module 4 of the Autism Diagnostic Observation Schedule 2 (ADOS-2) (Lord et al., 2000). All participants with autism met the threshold for a diagnosis of autism spectrum disorder on the ADOS-2 total classification score, and communication and reciprocal social interaction subscales. Moreover, groups were equated for age, and using the Wechsler Abbreviated Scale of Intelligence (WASI-II) (Wechsler, 1999), matched for full-scale IQ, and the verbal and performance subscales. Sample characteristics for these aforementioned control variables are presented in Table 1. The experiment was designed and conducted in accordance with the 1964 Declaration of Helsinki and approved by the research ethics committee of Liverpool John Moores University, UK.

Procedure

Participants were sat so their eyes were located approximately 555 mm from the centre of a 21-inch CRT monitor (Iiyama Vision Master 505), operating with a resolution of 1280 x 1024 pixels and a refresh rate of 85 Hz. Connected to the monitor was a desktop PC (Dell Optiplex GX280), graphics tablet and a hand-held stylus (Wacom Intuos Pro XL) (Fig. 1a).

In-house routines programmed in MATLAB (The Mathworks, Inc.) controlled the experiment, and the visual stimulus, which was generated using the Cogent 2000 toolbox (www.vislabucl.ac.uk/cogent.php).

Participants were provided with general instructions to “watch and then copy the movement displayed by a white dot on the computer monitor”. The model (i.e., a non-human agent model) displayed a single horizontal trajectory that originated from a home-target (diameter = 12.50 mm) positioned on the left-hand side of the screen and ended at a right-hand end-target (diameter = 12.50 mm), or right-hand end-space (i.e., no target; see below). The amplitude of the movement was 200 mm, and the total movement time was 1700 ms. To examine the imitation of biological motion, three non-human agent models were created that displayed typical, atypical or constant velocity profiles. The kinematics of the typical model trajectory were of biological origin as they were created by a human volunteer. To do so, the volunteer practised the task by performing typical goal-directed aiming movements using a hand-held stylus on a graphics tablet until a white dot/cursor (diameter = 6.25 mm), which represented the stylus, moved from the left-hand home-target to the right-hand end-target in 1700 ms. The model displayed a typical (Flash & Hogan, 1985) bell-shaped velocity profile (displacement time-series is displayed as the dark grey trace in Fig. 1b) that had a magnitude of peak velocity that was 0.20 mm/ms and a peak that occurred at 44% of the movement time. The atypical model (black trace in Fig. 1b) was created by the same volunteer, and thus ensured the kinematics of the movement were both biological and achievable by human participants. The volunteer practised performing an atypical movement until the 200 mm amplitude was completed in 1700 ms. The atypical model had a magnitude of peak velocity that was 0.41 mm/ms that occurred at 18% of the movement time. The model displaying constant velocity was created according to the amplitude (200 mm) and time (1700 ms) constraints associated with the task. The model displayed the exact movement time and

moved at constant velocity in the horizontal axis (0.118 mm/ms), with no deviations in the perpendicular axis (see Fig. 1b).

The imitation task comprised 14 blocks of 6 trials (84 trials). A block contained the typical, atypical and constant velocity models, each performed in the target and no-target conditions. Trial order within a block, as well as block order, was fully randomised across participants. The randomised structure reduced predictability of an upcoming model(s) and thus promoted imitation on a trial-by-trial basis. Prior to the experimental trials, all participants completed a familiarisation period that replicated the conditions of the imitation task. Participants performed four familiarisation trials, 2 trials representing a target condition, and 2 trials representing a no-target condition. Each trial commenced with the model being positioned in the centre of the home target at the left-hand side of the display after which it moved to the end-target (target condition), or end-space (no-target condition), with a constant velocity in a time of 1700 ms. A constant velocity trajectory was used to ensure construct validity by preventing participants from experiencing biological motion before the actual imitation trials. Participants were not informed about the time duration of the movement, the different types of stimulus, or the end-state target manipulation. Therefore, after observing a model, participants were only provided with a general instruction to copy the model (not a specific instruction to copy a certain aspect of the model; e.g., the kinematics) by moving the stylus on the tablet so that the cursor moved to the end-target, or end-space, as per the movement of the model. All participants confirmed they understood the model, the target and no-target conditions, the instruction to imitate a model, and the sensorimotor association between the stylus on a graphics tablet and the corresponding movement of cursor on the monitor.

Insert Fig. 1 here

Data Reduction and Analysis

To quantify imitation of motor timing, we extracted movement time from each trial and calculated an accuracy score (*motor timing accuracy*), which reflected the signed (+ or -) difference between a participant's movement time and that of the model (e.g., 1900 ms – 1700 ms = 200 ms). To examine motor timing consistency, we calculated a variability score (*motor timing variability*) that represented the within-participant distribution of movement time within a condition (i.e., standard deviation).

To quantify imitation of movement kinematics we focused the analysis on x-axis data because, much like the constant velocity model that had zero deviation in the y-axis, the perpendicular deviation in the y-axis for the atypical model and typical model was minimal. This was confirmed prior to our analysis by calculating perpendicular deviation using root-mean-square-error (RMSE) with respect to a value of zero (i.e., no deviation). Indeed, we observed the atypical model RMSE was 0.9 mm and typical model RMSE was 1.5 mm. Therefore, and similar to our previous work (Hayes, Andrew, Elliott, Roberts, & Bennett, 2012; Hayes, Elliott, & Bennett, 2010, 2013; Hayes, et al., 2014), we felt it was appropriate to conduct an analysis of dependent measures extracted from the primary movement (x-axis) only. Given such minimal deviation in the y-axis, any displacement in this axis by the participant would have most likely been an incidental result of anatomical constraints rather than intentional imitation. To complete the analysis, we identified within the x-axis position data the start and end of the movement. The start was defined as the moment the centre of the cursor moved beyond the perimeter of the 'home' target, and end equated to the moment the participant clicked the lower-button on the stylus. For each imitation trial, the resulting position data were filtered using a low pass 4th order autoregressive filter with an 8 Hz cut-off. The filtered data were next differentiated using a central difference algorithm to obtain

velocity. A MATLAB routine then extracted *peak velocity* and *percentage-time-to-peak-velocity* from each trial.

For both timing and kinematic dependent variables, intra-participant means were calculated from the first six and last six trials performed following observation of the 3 non-human agent models in the 2 target conditions. These data were submitted to separate 2 Group (autism; control) x 3 Model (atypical; typical; constant velocity) x 2 Target (target; no-target) x 2 Phase (early; late) repeated measures ANOVA. Significant main and/or interactions effects involving more than two means were analysed using Tukey HSD post-hoc procedure and alpha was set at $p < 0.05$. Additional correlation analysis on relevant significant comparisons indicated by ANOVA were then completed to assess whether the dependent measure correlated with autism severity (i.e., ADOS total score).

Results

The analysis of *motor timing accuracy* indicated a main effect of model [$F(2, 56) = 51.267, p < 0.01, \eta_p^2 = 0.647$]. Timing was more accurate after imitating atypical compared to typical ($p = 0.024$; difference = 67 ms) and constant ($p < 0.01$; difference = 285 ms) velocity models, and after imitating the typical compared to constant ($p < 0.01$; difference = 218 ms) velocity model. As illustrated in Fig. 2a, a 2-way interaction involving group x phase [$F(1, 28) = 9.480, p < 0.01, \eta_p^2 = 0.253$] showed *motor timing accuracy* increased by 175 ms from the early to late phase for the autism group ($p < 0.05$), whereas the performance of the control group deteriorated by 139 ms over the training period ($p < 0.05$). Correlation analysis revealed no relationship between *motor timing accuracy* in the early phase and ADOS total score (Pearson's $r(15) = 0.12, p > 0.05$) or late phase and ADOS total score (Pearson's $r(15) = -0.02, p > 0.05$).

The analysis of *motor timing variability* indicated a main effect of model [$F(2, 56) = 4.679, p = 0.01, \eta_p^2 = 0.143$], group [$F(1, 28) = 11.610, p = 0.01, \eta_p^2 = 0.293$], and a 2-way interaction involving group x phase [$F(1, 28) = 4.770, p < 0.05, \eta_p^2 = 0.146$]. As illustrated in Fig. 2b, motor timing was less variable after imitating the typical compared to constant ($p < 0.01$; difference 55 ms) velocity model, and when imitating atypical compared to the constant ($p = 0.01$; difference 51 ms) velocity model. Although the main effect indicated the control group ($M = 240$ ms) was significantly ($p < 0.05$) less variable overall than the autism group (363 ms), the interaction showed that only the autism group significantly ($p < 0.05$) decreased *motor timing variability* (by 99 ms) from the early to late phase of practice. Correlation analysis revealed no relationship between *motor timing variability* in the early phase and ADOS total score (Pearson's $r(15) = -0.25, p > 0.05$) or late phase and ADOS total score (Pearson's $r(15) = -0.33, p > 0.05$).

Insert Fig. 2 here

The analysis of *peak velocity* indicated a main effect of model [$F(2, 56) = 74.405, p < 0.01, \eta_p^2 = 0.727$] and a 2-way interaction involving group x phase [$F(1, 28) = 5.000, p < 0.05, \eta_p^2 = 0.152$]. As illustrated in Fig. 3a, *peak velocity* was greater when imitating atypical (0.238 mm/ms) compared to typical (0.192 mm/ms) and constant (0.162 mm/ms) velocity models ($ps < 0.01$). The interaction indicated *peak velocity* increased by 0.024 mm/ms from the early to late phase for the autism group ($p < 0.05$), whereas the decrease of 0.009 mm/ms for the control group was not significant. Correlation analysis revealed no relationship between *peak velocity* in the early phase and ADOS total score (Pearson's $r(15) = 0.03, p > 0.05$) or late phase and ADOS total score (Pearson's $r(15) = -0.06, p > 0.05$).

Insert Fig. 3 about here

The analysis of *percentage-time-to-peak-velocity* showed a main effect of model [$F(2, 56) = 41.536, p < 0.01, \eta_p^2 = 0.597$] and an interaction involving group x model [$F(2, 56) = 8.569, p < 0.01, \eta_p^2 = 0.234$]. As illustrated in Fig. 3b, although the groups did not differ when imitating the typical and constant velocity models ($ps > 0.05$), there was a significant difference when imitating the atypical velocity model ($p < 0.01$). The control group exhibited a *percentage-time-to-peak-velocity* that occurred significantly ($p < 0.01$) earlier in the movement (24 %), which was more similar to the atypical criterion model (18%: dashed line on Fig. 1b), than the autism group (33 %). These effects can be seen in the exemplar velocity traces illustrated in Fig. 4. When imitating the atypical model (back trace Fig. 4a), peak velocity occurred significantly earlier in the movement for the control group (dark grey trace), than the autism group (light grey trace). Whereas peak velocity occurred toward the midpoint of the movement for both groups (autism = light grey trace; control = dark grey trace) when imitating the typical (Fig. 4b) and constant (Fig. 4c) velocity models. Correlation analysis revealed no relationship between *percentage-time-to-peak-velocity* for atypical model and ADOS total score (Pearson's $r(15) = 0.22, p > 0.05$) or typical model and ADOS total score (Pearson's $r(15) = 0.05, p > 0.05$).

Insert Fig. 4 about here

Discussion

We examined imitation, and imitation adaption (i.e., performance change from the early to late phase on imitation), of biological motion kinematics using a novel behavioural protocol

that required adults with and without autism to observe a model that displayed distinctly different but biologically plausible kinematics. Importantly, the atypical biological motion would not have been represented in the sensorimotor repertoire of observers, and thus could not be imitated by rescaling a typical upper-limb aiming movement. After observing an atypical model, participants in the control group exhibited movements with a *percentage-time-to-peak-velocity* that occurred at 24% of the movement trajectory. This early occurrence of peak velocity was similar to that displayed by the atypical model (*percentage-time-to-peak-velocity* = 18%), and significantly different to the *percentage-time-to-peak-velocity* exhibited after observing typical ($M = 34\%$) and constant ($M = 39\%$) velocity control models. The presence of temporal correspondence between control participants' movements and the atypical model indicates high fidelity imitation of biological motion kinematics based on lower-level sensorimotor processes (Brass, Bekkering, & Prinz, 2001; Gangitano, Mottaghy, & Pascual-Leone, 2001; Hayes, et al., 2014; Heyes, 2001).

Equivalent high fidelity imitation of biological motion kinematics was not found for adults with autism. Although the magnitude of *peak velocity* was similar to control adults, there was a lack of temporal correspondence to the atypical model. The kinematic data showed *percentage-time-to-peak-velocity* occurred at 33% of the movement trajectory, which was significantly different from the control group, but statistically similar to the *percentage-time-to-peak-velocity* exhibited when imitating the typical ($M = 38\%$) and constant velocity ($M = 39\%$) control models. In this respect, our data are consistent with other work that demonstrated differences between those with and without autism in imitating the style (e.g., a gentle or harsh hand action) of a movement (Hobson & Lee, 1999; Rogers, et al., 1996; Smith & Bryson, 1994) or movement speed (Stewart, et al., 2013; Wild, et al., 2012). Importantly, however, the present findings extend understanding by showing differences in imitation are

1 directly related to attenuation in representing the temporal occurrence of peak velocity
 2 associated with the observed biological motion kinematics.

3 Before interpreting this effect, it is important to highlight that we isolated the
 4 examination of biological kinematics using a protocol that controlled higher order factors
 5 known to constrain imitation. First, we displayed an atypical model to ensure imitation was
 6 associated with representing novel biological kinematics, as opposed to presenting a
 7 movement that could be imitated using a pre-existing motor pattern recalled via higher-order
 8 semantic (Rumiati et al., 2005) or action-goal (Bekkering, Wohlschlaeger, & Gattis, 2000;
 9 Southgate & Hamilton, 2008) processes. Second, because imitation is modulated by social
 10 top-down factors (Chartrand & Bargh, 1999; Cook & Bird, 2012; Spengler, et al., 2010;
 11 Wang & Hamilton, 2012), we used a non-human agent model that reduced the influence of
 12 emotional (Grèzes, Wicker, Berthoz, & de Gelder, 2009) and/or theory of mind (Baron-
 13 Cohen et al., 1999) constraints that are inherent in realistic human models. Third, we
 14 controlled the influence of end-state target goal attainment (Bekkering, et al., 2000) by
 15 displaying a movement trajectory that had no targets in half of the trials. In combination, our
 16 use of these control measures minimizes the likelihood that the deficit in imitating biological
 17 motion kinematics in adults with autism is attributable to higher-order processes associated
 18 with reaching a target, or social imitation.

19 One explanation for the attenuation in imitating biological motion kinematics could be
 20 associated with lower-level processes that integrate visuomotor information (Dapretto et al.,
 21 2006; Oberman et al., 2005; Stewart, et al., 2013; Théoret et al., 2005; Williams, et al., 2006;
 22 Williams, et al., 2004). For example, visuomotor integration of biological motion occurs
 23 through specialised visual areas (posterior superior temporal sulcus; (Grossman, Battelli, &
 24 Pascual-Leone, 2005; Grossman et al., 2000) and lower-level sensorimotor processes linked
 25 to the mirror system (Iacoboni, 2005; Southgate & Hamilton, 2008). These processes are part

of a functional network that represents an observed movement by mapping the biological motion characteristics directly onto the motor system (Iacoboni, et al., 1999; Rizzolatti & Craighero, 2004). However, while lower-level processing deficits associated with visuomotor integration during self-other mapping (Stewart, et al., 2013; Williams, et al., 2006; Williams, et al., 2004) could attenuate imitation of atypical biological kinematics, it is notable that adults with autism show intact mapping of biological motion during automatic imitation (Bird, et al., 2007), which is a behavioural protocol that isolates processing to the lower-level mirror system. Moreover, results from neuropsychological work is mixed on whether such a fundamental impairment is present in autism (Hamilton, 2013).

Our data revealed an intriguing adaptation effect whereby adults with autism became significantly more accurate at representing movement time, reducing movement time variability, and increasing the magnitude of peak velocity over trials during imitation. This adaptation must have been self-regulated, as opposed to augmented, because external feedback regarding movement time performance was not provided. This change in behaviour can be ascribed to active and functional true imitation, with sensorimotor adaptation most likely a result of attending to, and comparing against, the observed stimulus using feedforward and feedback processes (Byrne & Russon, 1998; Carroll & Bandura, 1982; Kilner, Friston, & Frith, 2007). Moreover, within the group of high-functioning autism participants recruited in the current study, it would seem this adaptation is a general process that is not related to autism severity as determined by correlations with ADOS total score. In addition to modulating the magnitude of peak velocity, the positive change in accuracy for movement timing also reduced the influence of end-state-target-goals such that timing and kinematics changed similarly for target and no-target conditions. Moreover, we also found no evidence that the adult control group prioritised the attainment of an end-state-target-goal, over the imitation of atypical biological kinematics, when present during observation.

1 Although goal-directed imitation effects have been reported in complex movement sequences
2 (Wild, Poliakoff, Jerrison, & Gowen, 2010; Wild, et al., 2012) or a full body point-light
3 model (Hayes, Hodges, Huys, & Williams, 2007), it seems the target was less constraining
4 when individuals observed a point-light non-human agent model performing a single segment
5 movement.

6 The fact that adults with autism became significantly more accurate at imitating
7 movement time, and exhibited a magnitude of peak velocity that was similar to the control
8 group, suggests visual attention was orientated to the information displayed by the non-
9 human agent model. This effect is consistent with data showing visual attention to action
10 features of a model (Vivanti, et al., 2008), and non-human stimuli (Swettenham et al., 1998),
11 is typical in autism, whereas attention to facial features differs from controls (Bird, Catmur,
12 Silani, Frith, & Frith, 2006; Boucher & Lewis, 1992; Vivanti, et al., 2008). Moreover,
13 because no other attention-distracting stimuli were present in our display, it is unlikely that
14 reduced imitation of atypical biological kinematics was associated with visual attention being
15 drawn away from the non-human agent model (Wild, et al., 2012). A more parsimonious
16 explanation is that the selective attention bias to movement time during imitation was
17 controlled via alternative (and efficient) higher-order processes (Hamilton, et al., 2007;
18 Southgate & Hamilton, 2008; Wild, et al., 2012). A possibility is the movement time goal
19 was imitated using processes associated with action comprehension, which are functional in
20 autism (Dinstein et al., 2010), and as such goal attainment was secured using an efficient pre-
21 existing motor pattern. This interpretation is consistent with our kinematic data, which
22 showed individuals with autism executed movements that exhibited typical [peak velocity
23 occurred towards the mid-point of the trajectory (Elliott et al., 2010)] motor control
24 trajectories when imitating both the atypical and typical models.

1 In addition to a goal-directed and action comprehension interpretation, the selective
2 attention bias to movement time may have modulated input to the lower-level mirror system.
3 Input modulation is suggested to impact the activation, or development, of sensorimotor
4 representations via the intentionally mediated orientation of visual attention (Heyes, 2011;
5 Heyes & Bird, 2007; Liepelt & Brass, 2010; Longo, Kosobud, & Bertenthal, 2008).
6 Therefore, because we did not specify within our task instructions what aspect of the model
7 to imitate, the self-selected focus on movement time may have regulated the lower-level
8 processes such that this temporal variable was placed higher on the embedded hierarchy of
9 imitation goals (Hamilton & Grafton, 2007; Hayes, et al., 2014; Wohlschlagel, Gattis, &
10 Bekkering, 2003) than atypical kinematics. Although it is unclear if such input modulation is
11 operational in autism (Vivanti & Hamilton, 2014), we have differentially modulated how
12 atypical biological kinematics and movement time is imitated in neurotypical volunteers
13 using pre-specified verbal instructions (Hayes, et al., 2014). For example, the imitation of
14 atypical biological kinematics can be modulated if volunteers are instructed to focus attention
15 on imitating the movement time goal. Likewise, imitation accuracy can be enhanced if
16 selective attention is directed to the kinematics. Therefore, we cannot say for certain if the
17 focus on motor timing in individuals with autism is causally related to deficits in lower-level
18 self-other mapping processes and/or motor ability, or whether the attentional effect is a
19 compensatory strategy. One way to determine if the attenuation in imitating atypical
20 biological kinematics is associated with top-down attentional modulation is to present a
21 similar non-human agent model and employ a selective attention protocol that uses explicit
22 instructions to guide observers to attend and imitate the atypical biological kinematics
23 (Stewart, et al., 2013), as opposed to the observers self-selecting which action-based
24 information to imitate.

When considering the findings in respect to the broader context of imitation in autism, it is important to highlight we designed our study to examine ‘true imitation’. True imitation is a fundamental developmental process as it underpins the acquisition of novel social, and important sensorimotor skills that facilitate everyday life such as, tying shoe laces, riding a bicycle, or playing ice hockey. Although our data showed an attenuation in the imitation of biological motion kinematics, we did find that movement time accuracy and variability was significantly improved. The implication is that sensorimotor adaption and representation (Gidley Larson, Bastian, Donchin, Shadmehr, & Mostofsky, 2008) of movement time is intact in high-functioning adults with autism. These are first data to show this adaptation in a ‘true imitation’ context and indicates adults with autism do imitate, but they seem to do so in their own way. Therefore, the challenge is to examine the possibility that adults with autism can learn to imitate and represent biological motion kinematics following specific manipulations to the learning context (e.g., practice type, instructions, feedback). If the results are positive, then social and environmental procedures can be implemented by clinicians and practitioners to facilitate the acquisition of social and sensorimotor behaviors in autism.

In summary, the data presented here demonstrate, for the first time experimentally, that adults with autism have difficulties imitating the velocity characteristics associated with atypical biological motion kinematics. Compared to control participants, adults with autism became significantly more accurate at imitating movement time across trials. The positive change in behaviour confirmed they actively engaged in the task, and that sensorimotor adaptation during imitation is self-regulated in autism. The bias to movement time suggests the attenuation in imitating biological motion kinematics in autism is perhaps a compensatory strategy due to deficits in lower-level visuomotor processes associated with self-other

- 1 mapping and/or motor ability, or that selective attention input to the processes that represent
- 2 atypical biological motion kinematics.
- 3

References

- Bandura, A. (1977). *Social learning theory*: Englewood Cliffs, NJ: Prentice-Hall.
- Baron-Cohen, S., Ring, H. A., Wheelwright, S., Bullmore, E. T., Brammer, M. J., Simmons, A., & Williams, S. C. R. (1999). Social intelligence in the normal and autistic brain: an fMRI study. *European Journal of Neuroscience*, 11(6), 1891-1898. doi: 10.1046/j.1460-9568.1999.00621.x
- Bekkering, H., Wohlschlaeger, A., & Gattis, M. (2000). Imitation of gestures in children is goal-directed. *The Quarterly Journal of Experimental Psychology*, 53(1), 153-164. doi: 10.1080/713755872
- Bernier, R., Dawson, G., Webb, S., & Murias, M. (2007). EEG mu rhythm and imitation impairments in individuals with autism spectrum disorder. *Brain and Cognition*, 64(3), 228-237. doi: dx.doi.org/10.1016/j.bandc.2007.03.004
- Bird, G., Catmur, C., Silani, G., Frith, C., & Frith, U. (2006). Attention does not modulate neural responses to social stimuli in autism spectrum disorders. *Neuroimage*, 31(4), 1614-1624. doi: 10.1016/j.neuroimage.2006.02.037
- Bird, G., & Heyes, C. (2005). Effector-Dependent Learning by Observation of a Finger Movement Sequence. *Journal of Experimental Psychology: Human Perception and Performance*, 31(2), 262-275.
- Bird, G., Leighton, J., Press, C., & Heyes, C. (2007). Intact automatic imitation of human and robot actions in autism spectrum disorders. *Proceedings. Biological sciences / The Royal Society*, 274(1628), 3027-3031. doi: 10.1098/rspb.2007.1019
- Boucher, J., & Lewis, V. (1992). Unfamiliar Face Recognition in Relatively Able Autistic Children. *Journal of Child Psychology and Psychiatry*, 33(5), 843-859. doi: 10.1111/j.1469-7610.1992.tb01960.x
- Brass, M., Bekkering, H., & Prinz, W. (2001). Movement observation affects movement execution in a simple response task. *Acta Psychologica*, 106(1-2), 3-22.

- Buccino, G., Vogt, S., Ritzl, A., Fink, G. R., Zilles, K., Freund, H. J., & Rizzolatti, G. (2004). Neural circuits underlying imitation learning of hand actions: An event-related fMRI study. *Neuron*, 42(2), 323-334. doi: Doi 10.1016/S0896-6273(04)00181-3
- Byrne, R. W., & Russon, A. E. (1998). Learning by imitation: a hierarchical approach. *Behavioral and Brain Sciences*, 21(5), 667-684.
- Carroll, W. R., & Bandura, A. (1982). The role of visual monitoring in observational learning of action patterns: making the unobservable observable. *Journal of Motor Behavior*, 14(2), 153-167.
- Chartrand, T. L., & Bargh, J. A. (1999). The chameleon effect: the perception-behavior link and social interaction. *Journal of Personality and Social Psychology*, 76(6), 893-910.
- Cook, J. L., & Bird, G. (2012). Atypical Social Modulation of Imitation in Autism Spectrum Conditions. *Journal of Autism and Developmental Disorders*, 42(6), 1045-1051.
- Dapretto, M., Davies, M. S., Pfeifer, J. H., Scott, A. A., Sigman, M., Bookheimer, S. Y., & Iacoboni, M. (2006). Understanding emotions in others: mirror neuron dysfunction in children with autism spectrum disorders. [10.1038/nn1611]. *Nature Neuroscience*, 9(1), 28-30.
- Di Dio, C., Di Cesare, G., Higuchi, S., Roberts, N., Vogt, S., & Rizzolatti, G. (2013). The neural correlates of velocity processing during the observation of a biological effector in the parietal and premotor cortex. *Neuroimage*, 64(0), 425-436. doi: dx.doi.org/10.1016/j.neuroimage.2012.09.026
- Dinstein, I., Thomas, C., Humphreys, K., Minshew, N., Behrmann, M., & Heeger, D. J. (2010). Normal Movement Selectivity in Autism. *Neuron*, 66(3), 461-469. doi: doi.org/10.1016/j.neuron.2010.03.034
- Edwards, L. A. (2014). A Meta-Analysis of Imitation Abilities in Individuals With Autism Spectrum Disorders. *Autism Research*, 7(3), 363-380. doi: 10.1002/aur.1379
- Elliott, D., Hansen, S., Grierson, L. E. M., Lyons, J., Bennett, S. J., & Hayes, S. J. (2010). Goal-directed aiming: Two components but multiple processes. *Psychological Bulletin*, 136(6), 1023-1044. doi: 10.1037/a0020958
- Flash, T., & Hogan, N. (1985). The coordination of arm movements: an experimentally confirmed mathematical model. *The Journal of Neuroscience*, 5(7), 1688-1703.

- Gangitano, M., Mottaghy, F. M., & Pascual-Leone, A. (2001). Phase-specific modulation of cortical motor output during movement observation. *Neuroreport*, 12(7), 1489-1492.
- Gidley Larson, J. C., Bastian, A. J., Donchin, O., Shadmehr, R., & Mostofsky, S. H. (2008). Acquisition of internal models of motor tasks in children with autism. *Brain*, 131(11), 2894-2903. doi: 10.1093/brain/awn226
- Grèzes, J., Wicker, B., Berthoz, S., & de Gelder, B. (2009). A failure to grasp the affective meaning of actions in autism spectrum disorder subjects. *Neuropsychologia*, 47(8-9), 1816-1825. doi: dx.doi.org/10.1016/j.neuropsychologia.2009.02.021
- Grossman, E. D., Battelli, L., & Pascual-Leone, A. (2005). Repetitive TMS over posterior STS disrupts perception of biological motion. *Vision Research*, 45(22), 2847-2853. doi: dx.doi.org/10.1016/j.visres.2005.05.027
- Grossman, E. D., Donnelly, M., Price, R., Pickens, D., Morgan, V., Neighbor, G., & Blake, R. (2000). Brain areas involved in perception of biological motion. *Journal of Cognitive Neuroscience*, 12(5), 711-720.
- Hamilton, A. F. d. C. (2013). Reflecting on the mirror neuron system in autism: A systematic review of current theories. *Developmental Cognitive Neuroscience*, 3(0), 91-105.
- Hamilton, A. F. d. C., Brindley, R. M., & Frith, U. (2007). Imitation and action understanding in autistic spectrum disorders: how valid is the hypothesis of a deficit in the mirror neuron system? *Neuropsychologia*, 45(8), 1859-1868.
- Hamilton, A. F. d. C., & Grafton, S. T. (2007). The motor hierarchy: from kinematics to goals and intentions. In P. Haggard, Y. Rosetti & M. Kawato (Eds.), *Sensorimotor foundations of higher cognition: Attention and performance XXII*. (pp. 381-408). Oxford, UK: Oxford University Press.
- Hayes, S. J., Andrew, M., Elliott, D., Roberts, J. W., & Bennett, S. J. (2012). Dissociable contributions of motor-execution and action-observation to intermanual transfer. *Neuroscience Letters*, 506(2), 346-350.
- Hayes, S. J., Ashford, D., & Bennett, S. J. (2008). Goal-directed imitation: The means to an end. *Acta Psychologica*, 127(2), 407-415.
- Hayes, S. J., Elliott, D., & Bennett, S. J. (2010). General motor representations are developed during action-observation. *Experimental Brain Research*, 204, 1-8.
- Hayes, S. J., Elliott, D., & Bennett, S. J. (2013). Visual online control processes are acquired during observational practice. *Acta Psychologica*, 143(3), 298-302. doi: dx.doi.org/10.1016/j.actpsy.2013.04.012

- Hayes, S. J., Hodges, N. J., Huys, R., & Williams, A. M. (2007). End-point focus manipulations to determine what information is used during observational learning. *Acta Psychologica*, 126(2), 120-137.
- Hayes, S. J., Roberts, J. W., Elliott, D., & Bennett, S. J. (2014). Top-Down Attentional Processes Modulate the Coding of Atypical Biological Motion Kinematics in the Absence of Motor Signals. *Journal of Experimental Psychology: Human Perception and Performance*, 40(4), 1641-1653.
- Hayes, S. J., Timmis, M. A., & Bennett, S. J. (2009). Eye movements are not a prerequisite for learning movement sequence timing through observation. *Acta Psychologica*, 131(3), 202-208.
- Heyes, C. (2001). Causes and consequences of imitation. *Trends in Cognitive Sciences*, 5(6), 253-261.
- Heyes, C. (2011). Automatic imitation. *Psychological Bulletin*, 137(3), 463-483. doi: 10.1037/a0022288
- Heyes, C., & Bird, G. (2007). Mirroring, association, and the correspondence problem. In P. Haggard, Y. Rossetti & M. Kawato (Eds.), *Sensorimotor foundations of higher cognition: Attention and performance XX* (pp. 461-479). Oxford, England: Oxford University Press.
- Hobson, R. P., & Lee, A. (1999). Imitation and Identification in Autism. *Journal of Child Psychology and Psychiatry*, 40(4), 649-659. doi: 10.1111/1469-7610.00481
- Iacoboni, M. (2005). Neural mechanisms of imitation. *Current Opinion in Neurobiology*, 15(6), 632-637. doi: DOI 10.1016/j.conb.2005.10.010
- Iacoboni, M., Koski, L. M., Brass, M., Bekkering, H., Woods, R. P., Dubeau, M. C., . . . Rizzolatti, G. (2001). Reafferent copies of imitated actions in the right superior temporal cortex. *Proceedings of the National Academy of Sciences of the United States of America*, 98(24), 13995-13999. doi: DOI 10.1073/pnas.241474598
- Iacoboni, M., Woods, R. P., Brass, M., Bekkering, H., Mazziotta, J. C., & Rizzolatti, G. (1999). Cortical mechanisms of human imitation. *Science*, 286(5449), 2526-2528. doi: DOI 10.1126/science.286.5449.2526
- Johansson, G. (1973). Visual perception of biological motion and a model for its analysis. *Perception & Psychophysics*, 14(2), 201-211. doi: 10.3758/BF03212378

- Kilner, J. M., Friston, K. J., & Frith, C. D. (2007). Predictive coding: an account of the mirror neuron system. *Cognitive Processing*, 8(3), 159-166. doi: 10.1007/s10339-007-0170-2
- Kozlowski, L. T., & Cutting, J. E. (1977). Recognizing the sex of a walker from a dynamic point-light display. *Perception & Psychophysics*, 21(6), 575-580. doi: 10.3758/BF03198740
- Lakin, J. L., & Chartrand, T. L. (2003). Using Nonconscious Behavioral Mimicry to Create Affiliation and Rapport. *Psychological Science*, 14(4), 334-339. doi: 10.1111/1467-9280.14481
- Liepelt, R., & Brass, M. (2010). Automatic Imitation of Physically Impossible Movements. *Social Cognition*, 28(1), 59-73.
- Longo, M. R., Kosobud, A., & Bertenthal, B. I. (2008). Automatic imitation of biomechanically possible and impossible actions: effects of priming movements versus goals. *Journal of Experimental Psychology: Human Perception and Performance*, 34(2), 489-501. doi: 10.1037/0096-1523.34.2.489
- Lord, C., Risi, S., Lambrecht, L., Cook, E., Jr., Leventhal, B., DiLavore, P., . . . Rutter, M. (2000). The Autism Diagnostic Observation Schedule—Generic: A Standard Measure of Social and Communication Deficits Associated with the Spectrum of Autism. *Journal of Autism and Developmental Disorders*, 30(3), 205-223. doi: 10.1023/A:1005592401947
- Nishitani, N., Avikainen, S., & Hari, R. (2004). Abnormal imitation-related cortical activation sequences in Asperger's syndrome. *Annals of Neurology*, 55(4), 558-562. doi: 10.1002/ana.20031
- Oberman, L. M., Hubbard, E. M., McCleery, J. P., Altschuler, E. L., Ramachandran, V. S., & Pineda, J. A. (2005). EEG evidence for mirror neuron dysfunction in autism spectrum disorders. *Cognitive Brain Research*, 24(2), 190-198.
- Perra, O., Williams, J. H. G., Whiten, A., Fraser, L., Benzie, H., & Perrett, D. I. (2008). Imitation and 'theory of mind' competencies in discrimination of autism from other neurodevelopmental disorders. *Research in Autism Spectrum Disorders*, 2(3), 456-468. doi: dx.doi.org/10.1016/j.rasd.2007.09.007

- Prinz, W. (1997). Perception and action planning. *European Journal of Cognitive Psychology*, 9(2), 129-154.
- Rizzolatti, G., & Craighero, L. (2004). The mirror-neuron system. *Annual Review of Neuroscience*, 27, 169-192.
- Rogers, S. J. (1999). An examination of the imitation deficit in autism. In J. a. B. Nadel, G. (Ed.), *Imitation in infancy* (pp. 255-283.). Cambridge: Cambridge University Press.
- Rogers, S. J., Bennetto, L., McEvoy, R., & Pennington, B. F. (1996). Imitation and Pantomime in High-Functioning Adolescents with Autism Spectrum Disorders. *Child Development*, 67(5), 2060-2073. doi: 10.1111/j.1467-8624.1996.tb01843.x
- Rogers, S. J., Hepburn, S. L., Stackhouse, T., & Wehner, E. (2003). Imitation performance in toddlers with autism and those with other developmental disorders. *Journal of Child Psychology and Psychiatry*, 44(5), 763-781. doi: 10.1111/1469-7610.00162
- Rogers, S. J., & Pennington, B. F. (1991). A theoretical approach to the deficits in infantile autism. *Development and Psychopathology*, 3(02), 137-162. doi: doi:10.1017/S0954579400000043
- Rumiati, R. I., Weiss, P. H., Tessari, A., Assmus, A., Zilles, K., Herzog, H., & Fink, G. R. (2005). Common and differential neural mechanisms supporting imitation of meaningful and meaningless actions. *Journal of Cognitive Neuroscience*, 17(9), 1420-1431.
- Salowitz, N. G., Eccarius, P., Karst, J., Carson, A., Schohl, K., Stevens, S., . . . Scheidt, R. (2013). Brief Report: Visuo-spatial Guidance of Movement during Gesture Imitation and Mirror Drawing in Children with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 43(4), 985-995. doi: 10.1007/s10803-012-1631-8
- Smith, I. M., & Bryson, S. E. (1994). Imitation and action in autism: a critical review. *Psychological Bulletin*, 116(2), 259-273.
- Southgate, V., & Hamilton, A. F. d. C. (2008). Unbroken mirrors: challenging a theory of Autism. *Trends in Cognitive Sciences*, 12(6), 225-229. doi: 10.1016/j.tics.2008.03.005
- Spengler, S., Bird, G., & Brass, M. (2010). Hyperimitation of Actions Is Related to Reduced Understanding of Others' Minds in Autism Spectrum Conditions. *Biological Psychiatry*, 68(12), 1148-1155. doi: doi.org/10.1016/j.biopsych.2010.09.017

- Stewart, H. J., McIntosh, R. D., & Williams, J. H. G. (2013). A Specific Deficit of Imitation in Autism Spectrum Disorder. *Autism Research*, n/a-n/a. doi: 10.1002/aur.1312
- Swettenham, J., Baron-Cohen, S., Charman, T., Cox, A., Baird, G., Drew, A., . . . Wheelwright, S. (1998). The Frequency and Distribution of Spontaneous Attention Shifts between Social and Nonsocial Stimuli in Autistic, Typically Developing, and Nonautistic Developmentally Delayed Infants. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 39(05), 747-753. doi: doi:null
- Théoret, H., Halligan, E., Kobayashi, M., Fregni, F., Tager-Flusberg, H., & Pascual-Leone, A. (2005). Impaired motor facilitation during action observation in individuals with autism spectrum disorder. *Current Biology*, 15(3), R84-R85. doi: dx.doi.org/10.1016/j.cub.2005.01.022
- Vivanti, G., & Hamilton, A. F. d. C. (2014). Imitation in Autism Spectrum Disorders *Handbook of Autism and Pervasive Developmental Disorders, Fourth Edition*: John Wiley & Sons, Inc.
- Vivanti, G., Nadig, A., Ozonoff, S., & Rogers, S. J. (2008). What do children with autism attend to during imitation tasks? *Journal of Experimental Child Psychology*, 101(3), 186-205. doi: doi.org/10.1016/j.jecp.2008.04.008
- Wang, Y., & Hamilton, A. F. d. C. (2012). Social Top-down Response Modulation (STORM): A model of the control of mimicry in social interaction. *Frontiers in Human Neuroscience*, 6, 1. doi: 10.3389/fnhum.2012.00153
- Wechsler, D. (1999). *Wechsler Abbreviated Scales of Intelligence (WASI)*. . San Antonio, TX: Psychological Corporation.
- Wild, K. S., Poliakoff, E., Jerrison, A., & Gowen, E. (2010). The influence of goals on movement kinematics during imitation. *Experimental Brain Research*, 204(3), 353-360. doi: 10.1007/s00221-009-2034-8

- 1 Wild, K. S., Poliakoff, E., Jerrison, A., & Gowen, E. (2012). Goal-Directed and Goal-Less
2 Imitation in Autism Spectrum Disorder. *Journal of Autism and Developmental*
3 *Disorders*. doi: 10.1007/s10803-011-1417-4
- 4
- 5 Williams, J. H. G., Casey, J. M., Braadbaart, L., Culmer, P. R., & Mon-Williams, M. (2014).
6 Kinematic Measures of Imitation Fidelity in Primary School Children. *Journal of*
7 *Cognition and Development*, 15(2), 345-362. doi: 10.1080/15248372.2013.771265
- 8
- 9 Williams, J. H. G., Waiter, G. D., Gilchrist, A., Perrett, D. I., Murray, A. D., & Whiten, A.
10 (2006). Neural mechanisms of imitation and 'mirror neuron' functioning in autistic
11 spectrum disorder. *Neuropsychologia*, 44(4), 610-621. doi:
12 doi.org/10.1016/j.neuropsychologia.2005.06.010
- 13
- 14 Williams, J. H. G., Whiten, A., & Singh, T. (2004). A Systematic Review of Action Imitation
15 in Autistic Spectrum Disorder. *Journal of Autism and Developmental Disorders*,
16 34(3), 285-299. doi: 10.1023/B:JADD.0000029551.56735.3a
- 17
- 18 Williams, J. H. G., Whiten, A., Suddendorf, T., & Perrett, D. I. (2001). Imitation, mirror
19 neurons and autism. *Neuroscience & Biobehavioral Reviews*, 25(4), 287-295. doi:
20 doi.org/10.1016/S0149-7634(01)00014-8
- 21
- 22 Wohlschlager, A., Gattis, M., & Bekkering, H. (2003). Action generation and action
23 perception in imitation: an instance of the ideomotor principle. *Philosophical*
24 *Transactions of the Royal Society B: Biological Sciences*, 358(1431), 501-515. doi:
25 10.1098/rstb.2002.1257
- 26
- 27



Figure Captions

Figure 1(a) A schematic representation of the laboratory/experimental set-up for the imitation task. The black outlined rectangle represents a graphics tablet. The white circle displayed on the CRT monitor represents the model. The single-segment movement is depicted by the arrow (i.e., from the start position to the final position) in the target (red target) and no-target conditions. *(b)* Typical (dark-grey trace), atypical (black trace) and constant (light-grey trace) velocity models presented as a function of movement time.

Figure 2 (a) Timing accuracy and *(b)* Timing variability for the imitation task (error bars represent standard error of the mean) presented as a function of Group, Model and Phase.


Figure 3 (a) Peak velocity and *(b)* percentage-time-to-peak-velocity for the imitation task (error bars represent standard error of the mean) presented as a function of Group, Model and Phase. The dashed lines in Fig. 3a represent the magnitude of peak velocity for the typical (i.e., 0.41 mm/ms) and atypical (i.e., 0.20 mm/ms) models (Fig. 3a). In Fig. 3b, they represent the percentage-time-to-peak-velocity for the typical (i.e., 44%) and atypical (i.e., 18%) models.


Figure 4 The black velocity traces represent the atypical (a), typical (b), and constant (c) velocity models. In each panel (a, b, c), the light grey (autism) and dark grey (control) velocity traces display exemplar data from a representative participant imitating each model.




a**b**

1

2





a**b****c**

1

2

1 *Table 1* Characteristics of autism and neurotypical control participants

	Autism (n = 15)		Neurotypical (n = 15)		<i>t</i> test <i>p</i> value
	Mean (SD)	Range	Mean (SD)	Range	
Chronological age in years	26 (8)	18 - 44	26 (9)	18 - 45	<i>p</i> = 0.835
Full scale IQ	106 (10)	89 - 119	109 (7)	98 - 119	<i>p</i> = 0.333
Verbal IQ	104 (11)	88 - 127	108 (8)	95 - 122	<i>p</i> = 0.218
Performance IQ	105 (10)	90 - 128	106 (11)	90 - 124	<i>p</i> = 0.771
ADOS: Total	10 (2)	8 - 16			
ADOS: Communication	4 (1)	2 - 6			
ADOS: Social interaction	6 (2)	5 - 10			
Gender	14M : 1F		14M : 1F		

2