



# Geraniol prevents CCl<sub>4</sub>-induced hepatotoxicity via suppression of hepatic oxidative stress, pro-inflammation and apoptosis in rats

Abdulmohsen I. Algefare<sup>a</sup>, Manal Alfwuaires<sup>a</sup>, Ademola C. Famurewa<sup>b,c,\*</sup>, Hany Elsayy<sup>d</sup>, Azza Sedky<sup>a,e,\*\*</sup>

<sup>a</sup> Department of Biological Sciences, College of Science, King Faisal University, Al-Ahsa 31982, Saudi Arabia

<sup>b</sup> Department of Medical Biochemistry, Faculty of Basic Medical Sciences, College of Medical Sciences, Alex Ekwueme Federal University, Ndufu-Alike, Ikwo, Ebonyi State, Nigeria

<sup>c</sup> Department of Pharmacology, Manipal College of Pharmaceutical Sciences, Manipal Academy of Higher Education, Manipal, Karnataka State, India

<sup>d</sup> Department of Chemistry, Faculty of Science, Tanta University, Tanta 31527, Egypt

<sup>e</sup> Department of Zoology, Faculty of Science, Alexandria University, Alexandria, Egypt

## ARTICLE INFO

Handling Editor: Prof. L.H. Lash

### Keywords:

Geraniol  
Oxidative stress  
Hepatotoxicant  
Apoptosis  
Hepatotoxicity

## ABSTRACT

Carbon tetrachloride (CCl<sub>4</sub>) is a classic chemical hepatotoxicant that triggers liver damage through hepatic exacerbation of oxidative stress. Geraniol (GRL) is a natural bioactive acyclic monoterpene with several pharmacological effects. We thus explored whether GRL could prevent CCl<sub>4</sub>-triggered hepatic toxicity. Rats were divided and administered GRL (100 mg/kg) and/or CCl<sub>4</sub> (1 ml/kg of 1:1 v/v CCl<sub>4</sub>: olive oil) in Control group, GRL group, CCl<sub>4</sub> group, GRL + CCl<sub>4</sub> groups 2 times per week for 4 consecutive weeks. CCl<sub>4</sub> caused significantly ( $p < 0.05$ ) elevated serum activities of alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine aminotransferase (ALT), and total bilirubin (TB), whereas the albumin (ALB) and total protein (TP) levels were significantly ( $p < 0.05$ ) reduced relative to the control group. The liver activities of catalase (CAT), glutathione peroxidase (GPx), and superoxide dismutase (SOD) decreased significantly ( $p < 0.05$ ), while malondialdehyde (MDA) level evidently elevated in comparison to the control group. The CCl<sub>4</sub> exposure caused significant increases in proinflammatory interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF- $\alpha$ ), apoptotic caspase-3 and caspase-9 levels, whereas the anti-inflammatory interleukin-4 (IL-4) and interleukin-10 (IL-10) were reduced in consistent with histopathological changes compared to the control. On the contrary, the GRL administration prevented the hepatic toxicity and lesions through restoration of liver status markers, antioxidant enzyme activities, MDA, cytokines and apoptosis in comparison to the CCl<sub>4</sub> group. Altogether, the findings reveal that GRL could abrogate CCl<sub>4</sub>-provoked hepatic toxicity via inhibition of hepatic oxidative stress, inflammation and apoptosis in rats.

## 1. Introduction

Carbon tetrachloride is an industrial chemical and a classic hepatotoxicant. Carbon tetrachloride (CCl<sub>4</sub>) is still found as a component of insecticides, fire extinguishers and grains exposed to fumigation [22]. Human exposure to CCl<sub>4</sub> is well recognised to cause liver fibrosis and injury to the liver organ; hence, it is an established experimental agent for development of hepatotoxicity model [29]. The target organ of CCl<sub>4</sub> metabolism is the liver, where it is metabolised by cytochrome

P450-enzymes. The hepatic metabolism produces toxic oxidant metabolites like trichloromethyl peroxy ( $\bullet\text{OOCCL}_3$ ) and trichloromethyl ( $\bullet\text{CCl}_3$ ) oxygen free radicals associated with hepatic oxidative degeneration and cell death. Several studies indicate that the oxygen free radicals orchestrate impairment in hepatic membrane organization and antioxidant homeostasis leading to activation of aberrant signalling that favours inflammatory cascades and apoptotic cues [25,3,6]. Specifically, trichloromethyl radical reacts with sulfhydryl (-SH) group of thiol proteins, glutathione and glutathione-based enzymes to overrun the hepatic

\* Corresponding author at: Department of Medical Biochemistry, Faculty of Basic Medical Sciences, College of Medical Sciences, Alex Ekwueme Federal University, Ndufu-Alike, Ikwo, Ebonyi State, Nigeria.

\*\* Correspondence to: Department of Biological Sciences, College of Science, King Faisal University, PO Box 400 Al-Ahsa, 31982 Al-Ahsa, Saudi Arabia.

E-mail addresses: [ademola.famurewa@funai.edu.ng](mailto:ademola.famurewa@funai.edu.ng) (A.C. Famurewa), [asadek@kfu.edu.sa](mailto:asadek@kfu.edu.sa) (A. Sedky).

<sup>1</sup> ORCID: 0000-0002-8795-9964.

<https://doi.org/10.1016/j.toxrep.2024.01.007>

Received 15 May 2023; Received in revised form 10 January 2024; Accepted 11 January 2024

Available online 13 January 2024

2214-7500/© 2024 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

antioxidant system as well as glutathione metabolism [6]. These intracellular alterations in the liver predominantly provoke pathophysiology progressing to hepatic injury, non-alcoholic fatty liver disease, fibrosis, cirrhosis, hepatitis, and even hepatocellular carcinoma depending on the dose and duration [33]. For example, studies have shown that CCl<sub>4</sub> induces hepatotoxicity and mitochondrial damage characterized by redox imbalance, inflammation and apoptosis with consistent histological necrosis and infiltration of inflammatory cells [25,34,7]. In recent studies, however, the role of CCl<sub>4</sub>-induced reactive oxygen species (ROS), pro-inflammation and altered signalling cascades in hepatic fibrosis have been confirmed [22,24,8]. Therefore, antioxidants such as natural polyphenols with known anti-inflammatory potentials may exert inhibitory effects against CCl<sub>4</sub> hepatotoxicity.

Several studies have demonstrated that natural phytoactive compounds exhibiting antioxidant potentials can combat the ROS-mediated hepatotoxicity caused by CCl<sub>4</sub> [6,33]. Geraniol is a natural bioactive acyclic monoterpene with the chemical formula C<sub>10</sub>H<sub>18</sub>O. GRL (3, 7-dimethylocta-trans-2, 6-dien-1-ol) is an important content of palmarosa, lemon, ginger, lavender, orange essential oils, rose, and about 20 % component of GRL oil content [18]. The antioxidant and several pharmacological effects of GRL have been reported [17,20,8], such as the enhancement of antioxidants. GRL enhances antioxidant enzyme activities in dopaminergic neurons against ROS injury and apoptotic cell death in Parkinson's disease [23]. GRL blocks lead-induced or 2-acetylaminofluorene-induced hepatotoxicity through attenuation of oxidative stress, inflammation and apoptosis in rats [13,19]. The report of El Azab et al., [8] was only found on the efficacy of GRL against CCl<sub>4</sub>-induced hepatic oxidative stress and inflammation. Till date, it is unknown whether GRL could halt CCl<sub>4</sub>-induced oxidative stress-mediated apoptosis. Therefore, the study was conceived to explore the possible protective action of GRL against CCl<sub>4</sub>-induced hepatic damage via oxidative stress, inflammation and apoptosis in rats.

## 2. Materials and methods

### 2.1. Chemical and reagents

CCl<sub>4</sub> and GRL were purchased from Aladdin (China). Commercial kits were purchased from Jiancheng Co. (Nanjing, China).

### 2.2. Animals

Wistar rats (weight 200–240 g, eight weeks old,) were procured from the Animal House of Faculty of Science, King Faisal University, Saudi Arabia. The study protocols were in consistent with the research ethics at the King Faisal University (Reference number: KFU-REC-2022-DEC-ETHICS399). Animal cages with wire mesh that allows for good ventilation at 25 ± 2 °C, 50–60 % relative humidity and 12 h night/12 h day were used for this study. The rats were subjected to food and water without any restriction throughout the study period. Acclimatization was conducted by allowing the rats to feed on chow and water for one week.

### 2.3. Experimental design

At the end of acclimatization, the rats (male) were divided randomly into 4 groups (n = 6).

#### 2.3.1. Group 1 (Control)

Rats received olive oil (0.5 ml/kg b.w) intraperitoneally (ip) on the 1st and 4th day of each week for 4 weeks.

#### 2.3.2. Group 2 (GRL)

Rats received only GRL (100 mg/kg/day) orally (os) for 4 weeks.

#### 2.3.3. Group 3 (CCl<sub>4</sub>) group

Rats was administered only CCl<sub>4</sub> (1 ml/kg b.w of 1:1 v/v CCl<sub>4</sub>: olive oil, ip) on the 1st and 4th day of each week for 4 weeks.

#### 2.3.4. Group 4 (GRL+ CCl<sub>4</sub>)

Rats was administered CCl<sub>4</sub> (1 ml/kg b.w of 1:1 v/v CCl<sub>4</sub>: olive oil, ip) on the 1st and 4th day of each week for 4 weeks + GRL (100 mg/kg/day, os) for 4 weeks.

The doses of CCl<sub>4</sub> and GRL were chosen according to previous studies [14,21]. On the 29th day, rats were anesthetized, and blood samples collected into tubes for determination of liver function indices. The clotted blood was centrifuged at 2500g for 10 min at 4 °C. The serum obtained was stored at – 20 °C temperature. The rats were euthanized using 10 mg/kg xylazine and 100 mg/kg ketamine HCl for collection of liver samples for biochemical and histological analyses.

## 2.4. Biochemical analysis

### 2.4.1. Determination of hepatic markers

The serum samples were used for measuring the activities and levels of liver markers, AST, ALT, ALP, total protein (TP), albumin (ALB), and total bilirubin (TB) using standard kits and the kit instructions were followed for the assay steps.

### 2.4.2. Determination of oxidative stress markers

Liver sample (100 mg) was homogenized in PBS and centrifuged at 3000g for 20 min at 4 °C. The supernatant was separated into tubes. The liver activities of catalase (CAT, Cat. No. A007–1-1), superoxide dismutase (SOD, Cat. No. A001–1), and glutathione peroxidase (GPx, Cat. No. A005–1-1), as well as malondialdehyde (MDA, Cat. No. A003–1-2) level were estimated using commercial kits.

### 2.4.3. Determination of inflammatory markers

The liver supernatant samples were used to estimate the liver levels of pro-inflammatory TNF-α (Cat No R019) and IL-6 (Cat No R016), and anti-inflammatory IL-4 (Cat No R013) and IL-10 (Cat No R017). Rat standard ELISA kits were used and the determinations were conducted according to the manufacturer's instructions.

### 2.4.4. Apoptotic caspases estimation

Hepatic caspase 9 (Cat No A069) and caspase 3 (Cat No A064) were quantitatively measured in line with the manufacturer's directions.

### 2.4.5. Histopathology

Hepatic tissues were fixed with 10 % buffered formalin followed by the routine processes of blocking, paraffin and sectioning (5 μm thick) with microtome, and then mounted on glass slides. The processed sections were stained with H&E solution. Any histopathological alterations were viewed and photographed with a light microscope (Nikon 80i, Japan). The histopathological changes were scored semi-quantitatively and mean values were determined: normal histostructure (0), mild (1), moderate (2), and severe alterations consistent with congested central vein and presence of inflammatory cells and vacuolation (3) [9].

### 2.4.6. Statistical analysis

Data were presented as mean ± standard error of mean (SEM) obtained from the 6 rats in each group. The Graph Pad Prism statistical program (version 8; GraphPad Software Inc., San Diego, CA, USA) was used to analyse data by one-way ANOVA followed by Tukey's post-hoc test. The statistical significance was set at p < 0.05.

## 3. Results

### 3.1. Effect of GRL on liver function markers

Table 1 shows the effect of GRL on liver function markers in CCl<sub>4</sub>-

**Table 1**Effect of GRL on liver function indices in CCl<sub>4</sub>-intoxicated rats.

Group	ALT (U/L)	AST (U/L)	ALP (U/L)	TB (mg/dl)	ALB (g/dl)	TP (g/dl)
Control	35.5 ± 0.1	45.3 ± 0.2	25.9 ± 0.2	0.3 ± 0.1	4.30 ± 0.1	5.8 ± 0.11
GRL	36.2 ± 0.2	45.2 ± 0.2	28.4 ± 0.1	0.29 ± 0.1	4.30 ± 0.1	5.7 ± 0.14
CCl <sub>4</sub>	50.3 ± 0.1*	153.3 ± 0.2*	55.0 ± 0.5*	0.66 ± 0.1*	2.11 ± 0.1*	3.1 ± 0.10*
GRL + CCl <sub>4</sub>	40.2 ± 0.2#	84.2 ± 0.2#	30.1 ± 0.3#	0.38 ± 0.1#	3.20 ± 0.1#	4.1 ± 0.10#

Data are presented mean ± SEM (n = 6 rats/group). CCl<sub>4</sub>: carbon tetrachloride; GRL: Geraniol; \*P < 0.05: significant compared to the control group in the same column. #P < 0.05: significant compared to the CCl<sub>4</sub> group in the same column.

exposed rats. The CCl<sub>4</sub> exerted toxicity on the liver; CCl<sub>4</sub> significantly (p < 0.05) increased the serum activities of ALT, AST, ALP and level of TB in comparison to the control. Additionally, CCl<sub>4</sub> markedly reduced (p < 0.05) the levels of TP and ALB in comparison to the control. The administration of GRL prevented the toxic effect of CCl<sub>4</sub> demonstrated by significant reductions in serum activities of AST, ALP, ALT, and level of TB coupled with marked elevated levels of TP and ALB compared to the CCl<sub>4</sub> group.

### 3.2. Effect of GRL on oxidative stress markers


Fig. 1 presents the effect of GRL on oxidative stress indices in CCl<sub>4</sub>-exposed rats. The CCl<sub>4</sub> significantly (p < 0.05) reduced the hepatic activities of CAT, SOD and GPx, while level of MDA was increased in comparison to the control. The GRL administration in GRL + CCl<sub>4</sub> group significantly (p < 0.05) increased the activities of CAT, SOD and GPx, while level of MDA decreased significantly compared to the CCl<sub>4</sub> group.

### 3.3. Effect of GRL on hepatic inflammatory and anti-inflammatory markers


Fig. 2 presents the effect of GRL on hepatic anti-inflammatory and pro-inflammatory marker levels in CCl<sub>4</sub>-injected rats. The CCl<sub>4</sub> exposure induced significant increases (p < 0.05) in hepatic IL-6 and TNF-α levels compared to the control; CCl<sub>4</sub> triggered significant decreases in the anti-inflammatory IL-4 and IL-10 levels in the current study. GRL markedly (p < 0.05) inhibited the alterations in the hepatic TNF-α, IL-6, IL-4 and IL-10 levels compared to the CCl<sub>4</sub> group.

### 3.4. Effect of GRL on hepatic apoptosis caspases

Fig. 3 presents the effect of GRL on hepatic apoptosis marker levels in CCl<sub>4</sub>-injected rats. The CCl<sub>4</sub> exposure induced significant increases (p < 0.05) in the hepatic levels of caspase-3 and caspase-9 compared to the control. GRL significantly (p < 0.05) exerted reduction in the levels



**Fig. 1.** Effect of GRL on hepatic activities of oxidative stress markers in rats. GRL: Geraniol; CCl<sub>4</sub>: Carbon tetrachloride; data are shown as mean ± SEM (6 rats/group). \*p < 0.05: significant in comparison to the control group. #p < 0.05: significant in comparison to the CCl<sub>4</sub> group.



**Fig. 2.** Effect of GRL on liver cytokine levels in CCl<sub>4</sub>-injected rats. NRL: Geraniol; CCl<sub>4</sub>: Carbon tetrachloride; data are shown as mean  $\pm$  SEM (n = 6). \*p < 0.05: significant in comparison to the control group. #p < 0.05: significant in comparison to the CCl<sub>4</sub> group.

of caspase-3 and caspase-9 compared to the CCl<sub>4</sub> group.

### 3.5. Effect of GRL on the liver histology


Fig. 4 depicts the results of the histopathological analysis of the effect of GRL and CCl<sub>4</sub> on the liver of rats. The control and the GRL group liver samples revealed normal ultrastructure with evident normal hepatocyte, blood sinusoid, and central vein (a and b). In the CCl<sub>4</sub> group, histopathological abrasions like congested central vein and inflammatory cell infiltration; the hepatocytes appeared degenerated with vacuolated cytoplasm (c–e). However, in the GRL + CCl<sub>4</sub> group, we observed amelioration regarding the degeneration, congestion and infiltration of inflammatory cells (f).

## 4. Discussion

CCl<sub>4</sub> is a potential carcinogen and an established hepatotoxicant; nevertheless, it remains an industrial solvent, and humans are exposed to it at workplaces [6]. Given the critical biochemical role of the liver and its susceptibility to serious health problems worldwide, its protection from the attacks of exogenous toxicants is very important. Herein, our findings demonstrate that GRL could protect the liver from the oxidative and degenerative effects of CCl<sub>4</sub> via inhibition of oxidative inflammation and hepatic apoptosis.

In the current study, two times injection of CCl<sub>4</sub> per week for 4

consecutive weeks was used following the study of Hussein and Khan Ref. [14]. We thus found considerable liver damage revealed by significant elevations in the serum activities of liver enzymes, AST, ALT, ALP, and TB levels in comparison to the control group. Furthermore, the markers of hepatic protein synthetic capacity, ALB and TP, were markedly reduced in comparison to the control (Table 1). It therefore corroborates the earlier reports that CCl<sub>4</sub> is an hepatotoxicant [1,22,3,8]. The liver is involved in the catabolism of drugs and xenobiotics. These enzymes, ALT and AST are preponderant in the hepatocytes and are used to monitor the effect of drugs and xenobiotics on the liver health status. The damage to the hepatocytes by the CCl<sub>4</sub> caused the outflow of the enzymes into the serum [11], whereas the increased activity of ALP is an indicator of hepatic-cholestatic damage [12]. Liver synthesises serum proteins to main transport and blood osmotic balance. The reduced levels of TP and ALB confirmed the reduced synthetic capacity of the liver occasioned by the CCl<sub>4</sub>. Therefore, our results on the distortions of hepatic functional indices created by the CCl<sub>4</sub> injection are in agreement with the previous report [16,6,7]. In contrast, our concomitant administration of GRL to rats inhibited the CCl<sub>4</sub>-induced hepatic damage in this study. We found that the activities of AST, ALT, ALP, as well as levels of TB, ALB and TP were significantly restored in the serum compared to the CCl<sub>4</sub>-injected rats (Table 1). This demonstrates that GRL prevented the hepatic damage and also preserved the hepatic integrity such that the enzyme leakage into the blood circulation was significantly limited. Literature reveals the hepatoprotective effect of



**Fig. 3.** Effect of GRL on liver caspase levels in CCl<sub>4</sub>-injected rats. GRL: Geraniol; CCl<sub>4</sub>: Carbon tetrachloride; data are shown as mean  $\pm$  SEM (n = 6). \*Significant in comparison to the control (p < 0.05); #significant in comparison to the CCl<sub>4</sub> group (p < 0.05).

GRL in animal studies [13,19]. Its beneficial effect has also been associated with its antioxidant properties [22,30].

Redox homeostasis modulates the physiological production of ROS for benefits in immune responses and balance by antioxidant mechanism [2]. However, exogenous agents are known to cause impaired redox balance provoking oxidative stress-mediated damage [10]. We explored the effect of CCl<sub>4</sub> on the hepatic antioxidant status. CCl<sub>4</sub> impaired the antioxidant mechanism in the liver of rats in this study revealed by significant diminution in the hepatic activities of CAT, SOD and GPx, and the consequent elevation in MDA levels (Fig. 1). Our finding herein corroborates the literature reports that the oxidant metabolites produced upon the hepatic metabolism of CCl<sub>4</sub> do overwhelm hepatic antioxidant milieu [3,32,33,8]. Robust body of evidence shows that covalent binding of CCl<sub>4</sub> metabolites, trichloromethyl free radicals, to cell proteins and antioxidant apparatus is the first step in a chain reaction that eventually leads to membrane lipid peroxidation, liver lesions and oxidative stress [29]. The cooperative effect of these conditions might contribute to the depressed SOD, CAT, and GPx activities and elevated MDA levels shown in this study [7,25]. Evidently, the histological results showed liver lesions such as congested central vein and degenerated cells that could be linked with the oxidative milieu [29]. The antioxidant efficacy of GRL is reported in the literature [21,23]. However, the sulfhydryl groups in glutathione, GPx and other thiol proteins can react with trichloromethyl to form an oxidative complex,


and by extension, exerts deleterious effects on cellular antioxidant apparatus such as SOD and CAT, GPx, and reduced glutathione [6]. In this study, GRL exhibited antioxidant effect via improving the SOD, GPx and CAT activities against CCl<sub>4</sub> herein. These enzyme expressions were significantly elevated in the GRL + CCl<sub>4</sub> group compared to the CCl<sub>4</sub> group. Consequently, the level of MDA was also decimated significantly relative to the CCl<sub>4</sub> group. In oxidative stress-mediated diseases/disorder such as diabetes, diabetic cardiomyopathy, osteoarthritis, neuropathy and drug-induced toxicity, GRL has been underlined as a potent antioxidant agent for improvement of redox cycle balance [17,27,5,8]. Mechanistically, the GRL antioxidant action has been linked to its intracellular potential to activate the redox-sensitive pathway of nuclear factor erythroid 2-related factor 2 (Nrf2)/heme oxygenase-1 (HO-1) that promotes gene expression of antioxidant enzymes such as CAT, SOD and GPx among others via nuclear transcription [30]. Therefore, the GRL action in our study is in conformity with the earlier findings and could be attributed to the underlying activation/expression of antioxidant genes which were manifested via the increased activities of CAT, SOD and GPx.

Following the existing reports, CCl<sub>4</sub> oxidative stress causes activation of inflammatory and apoptotic cascades. CCl<sub>4</sub> action stimulates nuclear factor-kappa B (NF- $\kappa$ B), a nuclear transcriptional factor implicated in the expression of inflammatory cytokines and caspase-dependent apoptosis in liver damage [1,26,31,34]. To this end, we evaluated the effect of CCl<sub>4</sub> on pro-inflammatory (IL-6 and TNF- $\alpha$ ), anti-inflammatory (IL-4 and IL-10) cytokines and caspase markers of apoptosis. It was observed that the levels of TNF- $\alpha$  and IL-6 significantly increased, whereas IL-4 and IL-10 levels significantly reduced in CCl<sub>4</sub>-injected rats relative to the control rats (Fig. 2). This result shows that CCl<sub>4</sub> provoked pro-inflammation and reduces anti-inflammation [25,28]. NF- $\kappa$ B is translocated to the nucleus during cellular oxidative stress in the hepatic tissue [4], and its nuclear effect in terms of cytokine gene expression has been reported during CCl<sub>4</sub> hepatotoxicity [1,25]. In addition, we found histological infiltration of inflammatory cells (Fig. 4). It is therefore conceivable, in consonance with the previous findings, that CCl<sub>4</sub> could induce pro-inflammation in the liver [26,29,31]. Moreover, the cross-talk of inflammation and apoptosis is widely reported in published papers. For example, in a model of liver mitochondrial damage orchestrated by CCl<sub>4</sub>, the network of oxidative stress and hepatic inflammation was found to trigger apoptosis [28,34]. Our findings herein also agree with the earlier papers in that the expression of caspase 3 and caspase-9 were significantly elevated in the liver of CCl<sub>4</sub>-injected rats (Fig. 3). In contrast, the treatment with GRL evidently reversed the aberrant levels of the cytokines as well as the caspases. The hepatic IL-6 and TNF- $\alpha$  levels reduced while IL-4 and IL-10 levels increased significantly in GRL + CCl<sub>4</sub> group compared to the CCl<sub>4</sub> group in this study. We hereby confirm the anti-inflammatory effect of GRL in CCl<sub>4</sub>-mediated inflammatory liver damage. On the other hand, the caspases were also reduced significantly as a result of the antiapoptotic effect of GRL. The anti-inflammatory and antiapoptotic effects of GRL may be associated with the mechanistic inhibitory effect of GRL on the inflammatory factors, activators and mitochondrial integrity as suggested by the literature [15,17,21,30].

In conclusion, the present study has demonstrated the hepatotoxicity of CCl<sub>4</sub> in consistent with the literature. Importantly, the present work unravels that GRL could exert hepatoprotective effect against CCl<sub>4</sub>-triggered hepatic toxicity through mitigation of oxidative stress, inflammation and apoptosis. The effect could be added to the antioxidant, anti-inflammatory and antiapoptotic activities of GRL in this study.

#### CRediT authorship contribution statement

**Azza Sedky:** Conceptualization, Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing. **Hany Elsaywy:** Formal analysis, Funding acquisition, Investigation, Writing – review & editing. **Ademola C. Famurewa:** Conceptualization, Formal



**Fig. 4.** Photomicrographs of liver tissue (H&E). a) Liver tissue of the control group showed normal architecture, hepatocyte (H) with normal structure, sinusoid (arrows) and central vein (CV) (X100); b) GRL-treated group showed normal hepatocytes (H) between blood sinusoid (arrows) and central vein (CV) (X100); c-e) CCl<sub>4</sub>-injected group showing c) absence of hepatic architecture with congested central vein (CV) and inflammatory cell infiltration (arrows) (X100); d) congested central vein (CV) and inflammatory cell infiltration (arrows) (X400); e) congested central vein (CV) and hepatocytes with vacuolated cytoplasm (arrows) (X400); f) GRL + CCl<sub>4</sub> group showing improvement of structure of hepatocytes (H) and central vein (CV) (X400).

analysis, Writing – original draft, Writing – review & editing. **Manal A. Alfwuaires:** Conceptualization, Formal analysis, Writing – review & editing. **Abdulmohsen I. Algefare:** Conceptualization, Formal analysis, Writing – review & editing.

#### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Data availability

Data will be made available on request.

#### Acknowledgements

The authors acknowledge the Deanship of Scientific Research, Vice Presidency for Graduate Studies and Scientific Research at King Faisal University, Saudi Arabia for financial support under the Annual Funding track (Grant No. 5568).

#### References

- [1] H. Alabbad, M. Alfwuaires, A.M. Abdel-moneim, H. Elsway, Hepatoprotective mechanism of apigenin via suppression of oxidative inflammatory signaling and apoptosis against hepatotoxicity induced by CCl<sub>4</sub> in rats, *Indian J. Anim. Res.* (2023) 1–6, <https://doi.org/10.18805/IJAR.BF-1609.Submitted>.
- [2] J.J. Albaqami, H. Hamdi, A. Narayanankutty, N.U. Visakh, A. Sasidharan, A. M. Kuttithodi, A.C. Famurewa, Pathrose B. Chemical composition and biological

activities of the leaf essential oils of *Curcuma longa*, *Curcuma aromatica*, and *Curcuma angustifolia*, *Antibiotics* 11 (11) (2022) 1547.

- [3] S.A. Almatroodi, S. Anwar, A. Almatroudi, A.A. Khan, F. Alrumaihi, M.A. Alsahli, A.H. Rahmani, Hepatoprotective effects of garlic extract against carbon tetrachloride (CCl<sub>4</sub>)-induced liver injury via modulation of antioxidant, anti-inflammatory activities and hepatocyte architecture, *Appl. Sci.* (Switz.) 10 (18) (2020), <https://doi.org/10.3390/APP10186200>.
- [4] C.O. Edeogu, M.E. Kalu, A.C. Famurewa, N.T. Asogwa, G.N. Onyeji, K.O. Ikpemo, Nephroprotective effect of *Moringa oleifera* seed oil on gentamicin-induced nephrotoxicity in rats: biochemical evaluation of antioxidant, anti-inflammatory, and antiapoptotic pathways, *J. Am. Coll. Nutr.* (2019) 1–9, <https://doi.org/10.1080/07315724.2019.1649218>.
- [5] H.M. El-Bassossy, H. Ghaleb, A.A. Elberry, K.S. Balamash, S.A. Ghareib, A. Azhar, Z. Banjar, Geraniol alleviates diabetic cardiac complications: Effect on cardiac ischemia and oxidative stress, *Biomed. Pharmacother.* = *Biomed. Pharmacother.* 88 (2017) 1025–1030, <https://doi.org/10.1016/j.biopha.2017.01.131>.
- [6] A.E. El-Hadary, M.F. Ramadan Hassanien, Hepatoprotective effect of cold-pressed *Syzygium aromaticum* oil against carbon tetrachloride (CCl<sub>4</sub>)-induced hepatotoxicity in rats, *Pharm. Biol.* 54 (8) (2016) 1364–1372, <https://doi.org/10.3109/13880209.2015.1078381>.
- [7] D.H. El-Kashef, R.A. Zaghloul, Ameliorative effect of montelukast against carbon tetrachloride-induced hepatotoxicity: targeting NLRP3 inflammasome pathway, *Life Sci.* 304 (May) (2022) 120707, <https://doi.org/10.1016/j.lfs.2022.120707>.
- [8] E.F. El Azab, N.M. Elguindy, G.A. Yacout, D.A. Elgamal, Hepatoprotective impact of geraniol against CCl<sub>4</sub>-induced liver fibrosis in rats, *Pak. J. Biol. Sci.: PJB* 23 (12) (2020) 1650–1658, <https://doi.org/10.3923/PJB.2020.1650.1658>.
- [9] H. Elsway, A.C. Famurewa, A. Sedky, Resveratrol mitigates diabetic testicular dysfunction, endocrine deficits, and insulin resistance via suppression of sperm-endocrine aberrations and oxidative inflammation in rats, *Andrologia* (2023) 2023.
- [10] A.C. Famurewa, N.T. Asogwa, S.C. Ezea, Antidiabetic drug sitagliptin blocks cyclophosphamide cerebral neurotoxicity by activating Nrf2 and suppressing redox cycle imbalance, inflammatory iNOS/NO/NF-κB response and caspase-3/Bax activation in rats, *Int. Immunopharmacol.* 116 (2023) 109816, <https://doi.org/10.1016/j.intimp.2023.109816>.
- [11] A.C. Famurewa, C.A. Ekeleme-Egedigwe, E.E. David, C.O. Eleazu, A.M. Folawiyi, N.A. Obasi, Zinc abrogates anticancer drug tamoxifen-induced hepatotoxicity by suppressing redox imbalance, NO/iNOS/NF-κB signaling, and caspase-3-dependent

- apoptosis in female rats, *Toxicol. Mech. Methods* 30 (2) (2020), <https://doi.org/10.1080/15376516.2019.1669243>.
- [12] A.C. Famurewa, O.G. Ufebe, C.A. Egedigwe, O.E. Nwankwo, G.S. Obaje, Virgin coconut oil supplementation attenuates acute chemotherapy hepatotoxicity induced by anticancer drug methotrexate via inhibition of oxidative stress in rats, *Biomed. Pharmacother.* 87 (2017), <https://doi.org/10.1016/j.biopha.2016.12.123>.
- [13] S.K. Hasan, S. Sultana, Geraniol attenuates 2-acetylaminofluorene induced oxidative stress, inflammation and apoptosis in the liver of wistar rats, *Toxicol. Mech. Methods* 25 (7) (2015) 559–573, <https://doi.org/10.3109/15376516.2015.1070225>.
- [14] M. Hussein, R. Khan, CCL4-Induced Hepatotoxicity: Study in Rats Intoxicated with Carbon Tetrachloride and Treated with Camel Milk and Urine, *J. Chem. Stud.* 1 (1) (2022) 07–11, <https://doi.org/10.32996/jcs>.
- [15] M. Jayachandran, B. Chandrasekaran, N. Namasivayam, Geraniol attenuates fibrosis and exerts anti-inflammatory effects on diet induced atherogenesis by NF- $\kappa$ B signaling pathway, *Eur. J. Pharmacol.* 762 (2015) 102–111, <https://doi.org/10.1016/j.ejphar.2015.05.039>.
- [16] G.M. Kanawati, I.H. Al-Khateeb, Y.I. Kandil, Arctigenin attenuates CCl4-induced hepatotoxicity through suppressing matrix metalloproteinase-2 and oxidative stress, *Egypt. Liver J.* 11 (1) (2021) 1–7, <https://doi.org/10.1186/S43066-020-00072-6/FIGURES/4>.
- [17] Y. Lei, P. Fu, X. Jun, P. Cheng, Pharmacological properties of geraniol - a review, *Planta Med.* 85 (1) (2019) 48–55, <https://doi.org/10.1055/A-0750-6907>.
- [18] M.C.A. Leite, A.P. De Brito Bezerra, J.P. De Sousa, E. De Oliveira Lima, Investigating the antifungal activity and mechanism(s) of geraniol against *Candida albicans* strains, *Med. Mycol.* 53 (3) (2015) 275–284, <https://doi.org/10.1093/mmy/myu078>.
- [19] A. Ozkaya, Z. Sahin, M. Kuzu, Y.S. Saglam, M. Ozkaraca, M. Uckun, E. Yologlu, V. Comakli, R. Demirdag, S. Yologlu, Role of geraniol against lead acetate-mediated hepatic damage and their interaction with liver carboxylesterase activity in rats, *Arch. Physiol. Biochem.* 124 (1) (2018) 80–87, <https://doi.org/10.1080/13813455.2017.1364772>.
- [20] B. Pavan, A. Dalpiaz, L. Marani, S. Beggiato, L. Ferraro, D. Canistro, M. Paolini, F. Vivarelli, M.C. Valerii, A. Comparone, L.D. De Fazio, E. Spisni, Geraniol pharmacokinetics, bioavailability and its multiple effects on the liver antioxidant and xenobiotic-metabolizing enzymes, *Front. Pharmacol.* 9 (JAN) (2018), <https://doi.org/10.3389/fphar.2018.00018>.
- [21] S.N. Prasad, M. Muralidhara, Protective effects of geraniol (a monoterpene) in a diabetic neuropathy rat model: attenuation of behavioral impairments and biochemical perturbations, *J. Neurosci. Res.* 92 (9) (2014) 1205–1216, <https://doi.org/10.1002/jnr.23393>.
- [22] R. Que, M. Cao, Y. Dai, Y. Zhou, Y. Chen, L. Lin, Decursin ameliorates carbon-tetrachloride-induced liver fibrosis by facilitating ferroptosis of hepatic stellate cells, *Biochem. Cell. Biol. = Biochim. Et. Biol. Cell.* 100 (5) (2022) 378–386, <https://doi.org/10.1139/BCB-2022-0027>.
- [23] K.R. Rekha, G.P. Selvakumar, Gene expression regulation of Bcl2, Bax and cytochrome-C by geraniol on chronic MPTP/probenecid induced C57BL/6 mice model of Parkinson's disease, *Chem. -Biol. Interact.* 217 (2014) 57–66, <https://doi.org/10.1016/J.CBI.2014.04.010>.
- [24] E.A. Sayed, G. Badr, K.A.H. Hassan, H. Waly, B. Ozdemir, M.H. Mahmoud, S. Alameri, Induction of liver fibrosis by CCl4 mediates pathological alterations in the spleen and lymph nodes: the potential therapeutic role of propolis, *Saudi J. Biol. Sci.* 28 (2) (2021) 1272–1282, <https://doi.org/10.1016/j.sjbs.2020.11.068>.
- [25] J.C. Tsai, C.S. Chiu, Y.C. Chen, M. shiou Lee, X.Y. Hao, M.T. Hsieh, C.P. Kao, W. H. Peng, Hepatoprotective effect of *Coreopsis tinctoria* flowers against carbon tetrachloride-induced liver damage in mice, *BMC Complement. Altern. Med.* 17 (1) (2017) 1–9, <https://doi.org/10.1186/s12906-017-1604-8>.
- [26] M. Wang, J. Niu, L. Ou, B. Deng, Y. Wang, S. Li, Zerumbone protects against carbon tetrachloride (CCl4)-induced acute liver injury in mice via inhibiting oxidative stress and the inflammatory response: Involving the TLR4/NF- $\kappa$ B/COX-2 pathway, *Molecules* 24 (10) (2019), <https://doi.org/10.3390/molecules24101964>.
- [27] Y. Wu, Z. Wang, X. Fu, Z. Lin, K. Yu, Geraniol-mediated osteoarthritis improvement by down-regulating PI3K/Akt/NF- $\kappa$ B and MAPK signals: In vivo and in vitro studies, *Int. Immunopharmacol.* 86 (2020) 106713, <https://doi.org/10.1016/J.INTIMP.2020.106713>.
- [28] C. Yang, L. Li, Z. Ma, Y. Zhong, W. Pang, M. Xiong, S. Fang, Y. Li, Hepatoprotective effect of methyl ferulic acid against carbon tetrachloride-induced acute liver injury in rats, *Exp. Ther. Med.* 15 (3) (2018) 2228–2238, <https://doi.org/10.3892/etm.2017.5678>.
- [29] Y.H. Yeh, Y.L. Hsieh, Y.T. Lee, Effects of yam peel extract against carbon tetrachloride-induced hepatotoxicity in rats, *J. Agric. Food Chem.* 61 (30) (2013) 7387–7396, <https://doi.org/10.1021/jf401864y>.
- [30] N.S. Younis, M.S. Abduldaum, M.E. Mohamed, Protective effect of Geraniol on oxidative, inflammatory and apoptotic alterations in isoproterenol-induced cardiotoxicity: Role of the Keap1/Nrf2/HO-1 and PI3K/Akt/mTOR pathways, *Antioxidants* 9 (10) (2020) 1–17, <https://doi.org/10.3390/antiox9100977>.
- [31] S. Yue, N. Xue, H. Li, B. Huang, Z. Chen, X. Wang, Hepatoprotective effect of apigenin against liver injury via the non-canonical NF- $\kappa$ B pathway in vivo and in vitro, *Inflammation* 43 (5) (2020) 1634–1648, <https://doi.org/10.1007/S10753-020-01238-5>.
- [32] Y. Zhang, Y. Zhu, Y. Li, F. Ji, G. Ge, H. Xu, Midazolam ameliorates acute liver injury induced by carbon tetrachloride via enhancing Nrf2 signaling pathway, *Front. Pharmacol.* 13 (July) (2022) 1–12, <https://doi.org/10.3389/fphar.2022.940137>.
- [33] J. Zhao, Z. Zhang, J. Dai, L. Wang, C. Zhang, Y. Ye, L. Li, Synergistic protective effect of chlorogenic acid, apigenin and caffeic acid against carbon tetrachloride-induced hepatotoxicity in male mice, *RSC Adv.* 4 (81) (2014) 43057–43063, <https://doi.org/10.1039/c4ra07261h>.
- [34] J. Zhou, Y. Zhang, S. Li, Q. Zhou, Y. Lu, J. Shi, J. Liu, Q. Wu, S. Zhou, Dendrobium nobile Lindl. alkaloids-mediated protection against CCl4-induced liver mitochondrial oxidative damage is dependent on the activation of Nrf2 signaling pathway, *Biomed. Pharmacother.* 129 (June) (2020) 110351, <https://doi.org/10.1016/j.biopha.2020.110351>.