



LJMU Research Online

Pan, D, Sze, S, Irizar, P, George, N, Chaka, A, Lal, Z, Martin, CA, Nazareth, J, Baggaley, RF, Gray, LJ, Katikireddi, SV, Nellums, LB, Khunti, K and Pareek, M

Are clinical outcomes from COVID-19 improving in ethnic minority groups?

<http://researchonline.ljmu.ac.uk/id/eprint/25664/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Pan, D, Sze, S, Irizar, P, George, N, Chaka, A, Lal, Z, Martin, CA, Nazareth, J, Baggaley, RF, Gray, LJ, Katikireddi, SV, Nellums, LB, Khunti, K and Pareek, M (2023) Are clinical outcomes from COVID-19 improving in ethnic minority groups? eClinicalMedicine. 61.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Are clinical outcomes from COVID-19 improving in ethnic minority groups?

Daniel Pan,^{a,b,c,d} Shirley Sze,^e Patricia Irizar,^f Nisha George,^{a,b} Aasiya Chaka,^a Zainab Lal,^a Christopher A. Martin,^{a,b,c} Joshua Nazareth,^{a,b,c} Rebecca F. Baggaley,^g Laura J. Gray,^{c,g} S. Vittal Katikireddi,^h Laura B. Nellums,ⁱ Kamlesh Khunti,^j and Manish Pareek^{a,b,c,*}

^aDepartment of Respiratory Sciences, University of Leicester, UK

^bDepartment of Infectious Diseases and HIV Medicine, University Hospitals of Leicester NHS Trust, UK

^cLeicester NIHR Biomedical Research Centre, UK

^dLi Ka Shing Centre for Health Information and Discovery, Oxford Big Data Institute, University of Oxford, UK

^eDepartment of Cardiovascular Sciences, University of Leicester, UK

^fDepartment of Sociology, School of Social Sciences, University of Manchester, UK

^gDepartment of Population Health Sciences, University of Leicester, UK

^hMRC/CSO Social and Public Health Sciences Unit, University of Glasgow, UK

ⁱLifespan and Population Health, School of Medicine, University of Nottingham, UK

^jDiabetes Research Centre, University of Leicester, UK

Disproportionately worse COVID-19 clinical outcomes in people from ethnic minority groups have been a concern since early in the pandemic.¹ Now as time progresses, it may be useful to look back at the evolving evidence base. We performed the first systematic review on clinical outcomes in ethnic minority groups in May 2020, where we found across several countries a higher proportion of patients from ethnic minority groups infected with SARS-CoV-2, admitted to intensive care units with COVID-19 and dying in hospitals due to COVID-19. However, the collected data was of too poor quality to allow meaningful data synthesis.² Our findings were used as evidence for debate in UK Parliament in June 2020, resulting in a recommendation to mandate comprehensive ethnicity data collection and recording as part of routine hospital data collection systems.³

After some time, more studies started to emerge. This allowed us to conduct a meta-analysis to disentangle why ethnic minority groups were suffering disproportionately from the pandemic. In our second review, published in November 2020, we found that ethnic minority groups in the UK and USA had an increased risk of SARS-CoV-2 infection compared to those of White ethnicity. However, differences in hospitalization and death rates in this meta-analysis, when synthesised, was less clear.⁴ Whilst many studies reported a higher mortality rate from COVID-19 among ethnic minority groups compared to the majority groups, in the generation of mortality estimates, most did not take into account the number of infected individuals from ethnic minority groups in the community (see Fig. 1, Panel A). In our latest meta-analysis published in March 2023, now including over 200

million study participants globally, the risk of infection remained elevated across nearly all ethnic minority groups studied, compared to the majority group in each country.⁵ However, we observed far smaller differences for hospitalisation, intensive care admission and death following infection, although there was evidence of ethnic inequalities in these outcomes. Therefore, at least initially, it appeared that increased risk of infection was the main driver of the disproportionate outcomes in ethnic minority groups (see Fig. 1, Panel B). Our work was cited by the World Health Organization's living guideline on COVID-19 infection prevention and control, emphasizing the need for healthcare workers from ethnic minority groups to have equal access to personal protective equipment.⁶

Concrete data identifying and quantifying factors relating to SARS-CoV-2 infection remains limited, compared to risk factors for developing severe disease once infected. In an immunologically naïve population, mathematical models have proposed that the increased risk of infection with an airborne pathogen is related to a higher frequency and/or duration of exposure to individuals who emit high quantities of the virus in poorly ventilated spaces.⁷ Infection is therefore most likely to occur in homes and workplaces with poor ventilation and in occupations involving public-facing roles such as healthcare, even during mandated national lockdowns, all of which are common among those from ethnic minority groups. The higher prevalence of multi-generational occupancy within the homes of ethnic minority groups, which are more likely to have poorer ventilation, may also predispose them to a higher risk of infection.

To complicate matters, we now know that past infection with SARS-CoV-2 provides immune protection against severe disease for at least 40 weeks following infection, regardless of variants.⁸ While COVID-19 in immunologically naïve populations may have resulted in worse clinical outcomes for ethnic minority groups compared to the majority groups, a history of previous



eClinicalMedicine
2023;61: 102091

Published Online 11 July
2023

<https://doi.org/10.1016/j.eclinm.2023.102091>

*Corresponding author. Department of Respiratory Sciences, University of Leicester, UK.

E-mail address: Manish.pareek@leicester.ac.uk (M. Pareek).

© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1: Panel A: Dots represent proportions of those from different ethnic groups. A higher number of people infected with SARS-CoV-2 in the total population from ethnic minority groups would mean that a higher number of ethnic minority groups are hospitalised with COVID-19, even if the risk of hospitalisation remains the same across all ethnic groups. Panels B and C are simplified Directed Acyclic Diagrams, which show how hospitalisation and mortality may change between different waves of SARS-CoV-2 infection, depending on the number of people infected from ethnic minority groups.

infection in those who have survived their first infection, and therefore have existing immunity that protects against severe disease may have reduced the proportion of those who are susceptible to hospitalisation, intensive care admission and death from COVID-19 (see Fig. 1, Panel C). This could explain the UK's Office for National Statistics most recent report, which shows that the proportion of deaths from COVID-19 in ethnic minority groups are now comparable, and in some cases lower than that of the White British majority.⁹

Going forwards, we must implement tangible interventions that reduce the likelihood of infection among ethnic minority groups, which may still be ongoing. This will require policy-makers to address the longstanding inequalities that have led to an elevated risk of virus exposure in ethnic minority groups. In 2010, the Marmot review highlighted the need to reduce systemic racial inequalities in the UK and set clear policy objectives to address this. Ten years later, in the peak of the pandemic, the 2020 Marmot Review on Health Inequalities reported that housing affordability, declines in education funding, and an increase in zero-hour contracts are worse for ethnic minority groups compared to the previous report.¹⁰ Over the past three years, one thing has become clear: systemic inequality is likely the root cause of disproportionate deaths from the COVID-19 pandemic in ethnic minority groups. We now have a unique

opportunity to rebuild, plan ahead and work with communities from ethnic minority groups. Only by reducing the systemic gap in the risk of infection can we reduce preventable deaths from the next global pandemic.

Contributors

DP, SS, PI and MP conceived the idea of the manuscript. DP wrote the initial draft of the manuscript. All authors reviewed the manuscript and approved the final version prior to submission.

Declaration of interests

KK is Chair of the Ethnicity Subgroup of the UK Government Scientific Advisory Group for Emergencies (SAGE) and a member of SAGE. SVK was co-chair of the Scottish Government Expert Reference Group on Ethnicity and COVID-19 and a member of the Ethnicity Subgroup of SAGE. MP reports grants from the UKRI-MRC, NIHR, Sanofi and Gilead outside the current work and has received consulting fees from QIAGEN.

Acknowledgements

DP is supported by a NIHR doctoral research fellowship (NIHR302338). SS is supported by a NIHR Academic Clinical Lectureship. RFB is supported by a NIHR advanced fellowship (NIHR302494). MP is funded by a NIHR Development and Skills Enhancement Award and is supported by the Leicester NIHR Biomedical Research Centre (BRC). LJG and KK are supported by the National Institute for Health and Care Research (NIHR) Applied Research Collaboration East Midlands (ARC EM) and Leicester NIHR BRC. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. SVK acknowledges funding from the Medical Research Council (MC_UU_00022/2) and the Scottish Government Chief Scientist Office (SPHSU17).

References

- 1 Pareek M, Bangash MN, Pareek N, et al. Ethnicity and COVID-19: an urgent public health research priority. *Lancet*. 2020;395(10234):1421–1422.
- 2 Pan D, Sze S, Minhas JS, et al. The impact of ethnicity on clinical outcomes in COVID-19: a systematic review. *eClinicalMedicine*. 2020;23:100404.
- 3 Covid-19 and Black, Asian and minority ethnic communities. House of Commons Library, UK Parliament. <https://commonslibrary.parliament.uk/research-briefings/cdp-2020-0074/>. Accessed April 21, 2023.
- 4 Sze S, Pan D, Nevill CR, et al. Ethnicity and clinical outcomes in COVID-19: a systematic review and meta-analysis. *eClinicalMedicine*. 2020;29:100630.
- 5 Irizar P, Pan D, Kapadia D, et al. Ethnic inequalities in COVID-19 infection, hospitalisation, intensive care admission, and death: a global systematic review and meta-analysis of over 200 million study participants. *eClinicalMedicine*. 2023;57:101977.
- 6 World Health Organization. Infection prevention and control in the context of coronavirus disease (COVID-19): a living guideline, 7th March 2022. World Health Organization. <https://apps.who.int/iris/handle/10665/352339>. Accessed May 21, 2023.
- 7 Sze To GN, Chao CY. Review and comparison between the Wells-Riley and dose-response approaches to risk assessment of infectious respiratory diseases. *Indoor Air*. 2010;20(1):2–16.
- 8 COVID-19 Forecasting Team. Past SARS-CoV-2 infection protection against re-infection: a systematic review and meta-analysis. *Lancet*. 2023;401(10379):833–842.
- 9 Updating ethnic and religious contrasts in deaths involving the coronavirus (COVID-19), England: 24 January 2020 to 23 2022. Office for National Statistics, UK Government. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/updatingethniccontrastsindeathsinvolvingthecoronaviruscovid19englandandwales/24january2020to23november2022>. Accessed April 21, 2023.
- 10 *Health equity in England: the Marmot review 10 years on*. Institute of Health Equity; 2020. https://www.health.org.uk/sites/default/files/upload/publications/2020/Health%20Equity%20in%20England_The%20Marmot%20Review%2010%20Years%20On_full%20report.pdf.