

De Moor, D, Skelton, M, MacaqueNet, , Amici, F, Arlet, ME, Balasubramaniam, KN, Ballesta, S, Berghänel, A, Berman, CM, Bernstein, SK, Bhattacharjee, D, Bliss-Moreau, E, Brotcorne, F, Butovskaya, M, Campbell, LAD, Carosi, M, Chatterjee, M, Cooper, MA, Cowl, VB, De la O, C, De Marco, A, Dettmer, AM, Dhawale, AK, Erinjery, JJ, Evans, CL, Fischer, J, García-Nisa, I, Giraud, G, Hammer, R, Hansen, MF, Holzner, A, Kaburu, S, Konečná, M, Kumara, HN, Larrivaz, M, Leca, J-B, Legrand, M, Lehmann, J, Li, J-H, Lezé, A-S, MacIntosh, A, Majolo, B, Maréchal, L, Marty, PR, Massen, JJM, Maulany, RI, McCowan, B, McFarland, R, Merieau, P, Meunier, H, Micheletta, J, Mishra, PS, Sah, SAM, Molesti, S, Morrow, KS, Müller-Klein, N, Ngakan, PO, Palagi, E, Petit, O, Pflüger, LS, di Sorrentino, EP, Raghaven, R, Raimbault, G, Ram, S, Reichard, UH, Riley, EP, Rincon, AV, Ruppert, N, Sadoughi, B, Santhosh, K, Schino, G, Sheeran, LK, Silk, JB, Singh, M, Sinha, A, Sosa, S, Stribos, MS, Sueur, C, Tiddi, B, Tkaczynski, PJ, Trebouet, F, Widdig, A, Whitehouse, J, Wooddell, LJ, Xia, D-P, von Fersen, L, Young, C, Schülke, O, Ostner, J, Neumann, C, Duboscq, J and Brent, LNJ

MacaqueNet: Advancing comparative behavioural research through large-scale collaboration

<http://researchonline.ljmu.ac.uk/id/eprint/25693/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

De Moor, D, Skelton, M, MacaqueNet, , Amici, F, Arlet, ME, Balasubramaniam, KN, Ballesta, S, Berghänel, A, Berman, CM, Bernstein, SK, Bhattacharjee, D, Bliss-Moreau, E, Brotcorne, F, Butovskaya, M, Campbell, LAD, Carosi, M, Chatterjee, M, Cooper, MA, Cowl, VB, De la O, C.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

RESEARCH ARTICLE

MacaqueNet: Advancing comparative behavioural research through large-scale collaboration

Delphine De Moor¹  | Macaela Skelton¹  | MacaqueNet | Federica Amici^{2,3}  |
 Malgorzata E. Arlet⁴  | Krishna N. Balasubramaniam^{5,6}  | Sébastien Ballesta^{7,8}  |
 Andreas Berghänel⁹  | Carol M. Berman¹⁰  | Sofia K. Bernstein¹¹  |
 Debottam Bhattacharjee¹²  | Eliza Bliss-Moreau¹³  | Fany Brotcorne¹⁴  |
 Marina Butovskaya¹⁵  | Liz A. D. Campbell¹⁶  | Monica Carosi¹⁷  |
 Mayukh Chatterjee^{18,19}  | Matthew A. Cooper²⁰  | Veronica B. Cowl²¹  |
 Claudio De la O^{22,23,24}  | Arianna De Marco²⁵  | Amanda M. Dettmer²⁶  |
 Ashni K. Dhawale²⁷  | Joseph J. Erinjery²⁸  | Cara L. Evans²⁹ | Julia Fischer^{30,31,32}  |
 Iván García-Nisa³³  | Gwennan Giraud¹⁴  | Roy Hammer³⁴  |
 Malene F. Hansen^{35,36,37,38}  | Anna Holzner^{39,40,41}  | Stefano Kaburu⁴²  |
 Martina Konečná⁴³  | Honnavalli N. Kumara⁴⁴  | Marine Larrivaz⁴⁵ |
 Jean-Baptiste Leca⁴⁶  | Mathieu Legrand⁸  | Julia Lehmann⁴⁷  | Jin-Hua Li^{48,49}  |
 Anne-Sophie Lezé⁵⁰ | Andrew MacIntosh⁵¹  | Bonaventura Majolo⁵²  |
 Laëtitia Maréchal⁵²  | Pascal R. Marty⁵³  | Jorg J. M. Massen^{12,54}  |
 Risma Illa Maulany⁵⁵  | Brenda McCowan⁶  | Richard McFarland^{56,57}  |
 Pierre Merieau⁵⁰ | Hélène Meunier^{7,8} | Jérôme Micheletta⁵⁸  | Partha S. Mishra^{44,59}  |
 Shahrul A. M. Sah⁴¹  | Sandra Molesti⁶⁰  | Kristen S. Morrow⁶¹  |
 Nadine Müller-Klein⁶²  | Putu Oka Ngakan⁵⁵  | Elisabetta Palagi⁶³  | Odile Petit⁶⁴  |
 Lena S. Pflüger^{34,54}  | Eugenia Polizzi di Sorrentino⁶⁵  | Roopali Raghaven¹⁹ |
 Gaël Rimbault⁸ | Sunita Ram^{19,66} | Ulrich H. Reichard⁶⁷  | Erin P. Riley⁶⁸  |
 Alan V. Rincon⁵⁸  | Nadine Ruppert⁴¹  | Baptiste Sadoughi^{7,69,70,71}  |
 Kumar Santhosh⁴⁴  | Gabriele Schino⁶⁵  | Lori K. Sheeran⁷²  | Joan B. Silk⁷³  |
 Mewa Singh⁷⁴  | Anindya Sinha¹⁹  | Sebastian Sosa⁴⁰  | Mathieu S. Stribos¹² |
 Cédric Sueur⁷⁵  | Barbara Tiddi⁷⁶  | Patrick J. Tkaczynski⁷⁷  | Florian Trebouet⁷⁸  |
 Anja Widdig^{39,40,79}  | Jamie Whitehouse⁵⁶  | Lauren J. Wooddell⁸⁰  |
 Dong-Po Xia^{48,81}  | Lorenzo von Fersen⁸²  | Christopher Young⁵⁶  |
 Oliver Schülke^{32,70,71}  | Julia Ostner^{32,70,71}  | Christof Neumann^{30,32}  |
 Julie Duboscq^{71,83}  | Lauren J. N. Brent¹ 

For affiliations refer to page 11.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Journal of Animal Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

Correspondence

Delphine De Moor

Email: delphinedemoor@gmail.com**Funding information**

H2020 European Research Council,
Grant/Award Number: 864461; Leibniz
ScienceCampus EEGA, Grant/Award
Number: LSC-AF2023_03

Handling Editor: Jean-Michel Gaillard**Abstract**

1. There is a vast and ever-accumulating amount of behavioural data on individually recognised animals, an incredible resource to shed light on the ecological and evolutionary drivers of variation in animal behaviour. Yet, the full potential of such data lies in comparative research across taxa with distinct life histories and ecologies. Substantial challenges impede systematic comparisons, one of which is the lack of persistent, accessible and standardised databases.
2. Big-team approaches to building standardised databases offer a solution to facilitating reliable cross-species comparisons. By sharing both data and expertise among researchers, these approaches ensure that valuable data, which might otherwise go unused, become easier to discover, repurpose and synthesise. Additionally, such large-scale collaborations promote a culture of sharing within the research community, incentivising researchers to contribute their data by ensuring their interests are considered through clear sharing guidelines. Active communication with the data contributors during the standardisation process also helps avoid misinterpretation of the data, ultimately improving the reliability of comparative databases.
3. Here, we introduce MacaqueNet, a global collaboration of over 100 researchers (<https://macaquetnet.github.io/>) aimed at unlocking the wealth of cross-species data for research on macaque social behaviour. The MacaqueNet database encompasses data from 1981 to the present on 61 populations across 14 species and is the first publicly searchable and standardised database on affiliative and agonistic animal social behaviour. We describe the establishment of MacaqueNet, from the steps we took to start a large-scale collective, to the creation of a cross-species collaborative database and the implementation of data entry and retrieval protocols.
4. We share MacaqueNet's component resources: an R package for data standardisation, website code, the relational database structure, a glossary and data sharing terms of use. With all these components openly accessible, MacaqueNet can act as a fully replicable template for future endeavours establishing large-scale collaborative comparative databases.

KEYWORDS

comparative research, data sharing, database, *Macaca*, primates, repository, social networks, team science

1 | INTRODUCTION

Comparative studies are fundamental in understanding the biological basis of traits. They can reveal broad patterns in evolutionary and developmental history, past and current selective pressures and underlying mechanisms (Nunn, 2011; Tinbergen, 1963). Yet, a major limitation to comparative research is reaching a sufficiently large and diverse sample for reliable inference (Borries et al., 2016; Schneider et al., 2019). Comparative studies therefore typically rely on combining independent research on individual populations (Lukas & Clutton-Brock, 2017).

There is a vast and ever-accumulating amount of behavioural data on individually recognised animals (Sheldon et al., 2022). This is an incredible resource for comparative research, that can reveal fundamental principles on the ecological and evolutionary processes that shape animal behaviour. However, combining these rich behavioural data comes with substantial challenges (Pinter-Wollman et al., 2013). The field of animal behaviour historically and currently still consists of many independent research groups collecting and managing data using similar yet slightly different methods (the 'long tail' of data: many small datasets that together represent the vast majority of existing data; Wallis et al., 2013). Choices regarding which behaviours

to observe, how those behaviours are defined, and which methods are used to record them, result in a wide variation in behavioural data types and definitions. This complicates the creation of standardised databases on behaviour, compared with existing databases pooling relatively consistently defined and quantified data, such as life history and demography, morphological and ecological traits, population sizes and geographic ranges, social organisation and measures of biodiversity (SPI-Birds: Culina et al., 2020; The Global Biodiversity Information Facility: Edwards et al., 2000; Co-Breed: Mocha et al., 2024; panTHERIA: Jones et al., 2009; COMADRE: Salguero-Gomez et al., 2016; The Primate Life History Database: Strier et al., 2010; BIDDABA : Lebreton et al., 2010).

Compounding this challenge, most comparative studies in animal behaviour typically consist of one-off comparative research projects, spearheaded by an individual research team and focused on a specific question. The generated databases are usually not conceived for general use, making them difficult to find or repurpose for future studies (O'Dea et al., 2021). For instance, data are often summarised into a single datapoint per species, according to the definitions and decisions of the research team and project. As such, the painstaking effort of searching the literature for suitable data, acquiring such data, and standardising the data and metadata across datasets needs to be repeated for each new comparative study (Poisot et al., 2019). A far more sustainable approach is to create enduring and reusable databases, with well-defined protocols for (meta)data standardisation, archiving and accessing (Culina et al., 2020; Urbano & Cagnacci, 2021). Clear and concise guidelines for such databases are provided by the Findable, Accessible, Interoperable and Reusable (FAIR) guiding principles to ensure data are 'FAIR' (Wilkinson et al., 2016). The development of such databases has become a priority for many scientific fields and has led to the creation of big-team science initiatives, bringing together scientists 'across labs, institutions, disciplines, cultures, and continents' (Forscher et al., 2022, p. 2). The aim of these collaborations is to enable large teams of researchers to collaborate more effectively and to accelerate scientific discoveries by making it easier to integrate and compare data from different sources.

Many of these team-science endeavours follow the 'Many Labs' approach (Klein et al., 2014; e.g. ManyPrimates: ManyPrimates et al., 2019; ManyBabies: Visser et al., 2022), where research teams collaborate to each contribute data collected according to a standardised protocol to maximise comparability. This is a promising approach, and one that ideally should be adopted more broadly across long-term research sites to allow for the identification of broad patterns across sites and study systems (Rubenstein & Abbot, 2017). Yet, it would be a waste not to use the 70+ years' worth of behavioural data that already exist (Purgar et al., 2022; Sheldon et al., 2022). Although complex, there is great potential in standardising existing data and making them usable for comparative research (Pinter-Wollman et al., 2013).

Two big-team initiatives on animal spatial data, Euromammals and Movebank, demonstrate the strength of collaborative efforts in standardising and centralising animal behaviour data. Since their inception

in 2007 and 2008, respectively, these initiatives have grown into key repositories for thousands of animal movement datasets, supporting collaborative research and resulting in numerous publications (Kays et al., 2021; Kranstauber et al., 2011; Urbano & Cagnacci, 2021). More recently, similar initiatives have emerged to aggregate social behavioural data. The Animal Social Network Repository (ASNR) was the first comprehensive effort to consolidate social behaviour data across different animal species using FAIR principles (Sah et al., 2019). Since its publication, the ASNR database has been used for several comparative studies across a broad taxonomic range (e.g. Collier et al., 2022; Gagliardi et al., 2023), demonstrating the value of such databases for the scientific community. However, the ASNR database's diversity also presents challenges for comparative research as it contains unstandardised social data collected in various ways, with substantial variation in observation effort and capturing fundamentally distinct aspects of sociality. A second, recently developed database is DomArchive, which focuses on dominance interactions within groups of various taxa (Strauss et al., 2022). Social dominance is conceptually similar and measured fairly consistently across taxa (Strauss et al., 2022), and therefore lends itself well to comparative questions. However, a comprehensive understanding of social behaviour necessitates data on both affiliative and agonistic social interactions. Both types of interactions play crucial but distinct roles in shaping the dynamics and complexity of social systems. In contrast to dominance interactions, affiliative interactions are typically more varied in terms of both the types of behaviours and methods used to measure them, making it more complex to build a database of truly comparable data (De Moor et al., 2024; Ellis et al., 2019). One approach to overcoming this challenge is to collate data for taxa in which the observed behaviours and recording protocols are relatively uniform. These databases can then be integrated into an archive of databases, such as the EcoEvo data source catalogue (<https://ckan-ecoevo.d4science.org/>, Culina et al., 2018), to ultimately create a comprehensive FAIR database of social behaviour.

Here, we introduce MacaqueNet (<https://macaquetnet.github.io/>), a global grassroots collaboration that provides a platform for big-team social behavioural research on macaque monkeys (*Macaca* spp.), and the MacaqueNet database, a cross-species collaborative FAIR database, facilitating synthetic comparative research (Borries et al., 2016; Coles et al., 2022). To pave the way for future collaborative initiatives in comparative behavioural research, we describe how we initiated MacaqueNet, compiled and standardised data to create a relational database, implemented data sharing protocols and built a global community of collaborating researchers. With all core components openly accessible, MacaqueNet can act as a fully replicable template for other researchers interested in setting up collaborative cross-species databases.

2 | METHODS AND RESULTS

MacaqueNet connects researchers studying the social behaviour of macaques in diverse settings. The aim of MacaqueNet is twofold: to

encourage and facilitate grassroots collaboration between independent research teams and to provide a lasting central platform for data archiving, standardising and accession. Here, we describe the establishment of MacaqueNet, from starting a large-scale collective to the creation of a cross-species collaborative database and the implementation of data entry and retrieval protocols. First, however, we illustrate the exceptional suitability of macaques as a taxon to initiate the construction of cross-species databases of social behavioural data.

2.1 | Why the genus *Macaca* lends itself to the creation of a comparative database of social behaviour

With 25 currently recognised extant species, *Macaca* is the most widely geographically distributed non-human primate genus (Roos et al., 2019). Macaques are one of the most versatile and adaptable primates, exploiting very different environments, from the temperate, mountainous habitats of Morocco and Japan to the tropical forests of Southeast Asia (Thierry, 2007). Throughout this range, they live in areas of varying anthropogenic impact, from cities over natural habitats coming under increasing tourist and agricultural pressure, to pristine forests (Radhakrishna et al., 2012). This variation in habitats and climates is reflected in the species' ecology, mating system and social structure (the content, quality and patterning of social relationships between individuals belonging to the same social unit; Cords, 2012; Kappeler et al., 2019). Macaques are primarily frugivorous, range from being arboreal to semiterrestrial and encounter different levels of predation (Fleagle, 2013). Mating is polygynandrous and reproduction can be seasonal or year-round, with substantial variation in male reproductive skew (biased distribution of offspring sired towards a few, usually high-ranking, males) across species (Schülke & Ostner, 2008).

Despite this variation, macaques have a rather conserved social organisation (the size and composition of a social unit; Kappeler et al., 2019). They typically live in relatively large multi-male multi-female groups of approximately ten to a hundred individuals in the wild and up to several hundred in urban or provisioned populations (Cords, 2012). Females are philopatric and form the core of the group, clustered into multigenerational matriline. Males typically disperse from their birth group around puberty to breed elsewhere and may even change groups several times in their life (De Moor et al., 2020). Both male and female macaques exhibit a wide range of social behaviours, many of which are shared across all species (Thierry, 2007). While social relationships tend to be kin biased and hierarchical to some extent, macaques show substantial variation in the patterns of affiliative and agonistic interactions. This remarkable diversity within unity makes macaques a compelling taxon for establishing a cross-species database aimed at investigating the ecological and evolutionary processes underlying this variability (Balasubramaniam et al., 2018; Thierry, 2007).

Macaques are also excellent as a starting point to address the challenge of cross-species behavioural databases. Many species of

macaques have been extensively studied in the wild or naturalistic semi-free ranging settings for years or even decades, with observed behaviours and recording protocols being relatively uniform across research sites, facilitating comparison (*Macaca assamensis*, Ostner & Schülke, 2018; *Macaca fascicularis*, Van Noordwijk & Van Schaik, 1985; *Macaca fuscata*, Nakagawa et al., 2010; *Macaca leonina*, Albert et al., 2013; *Macaca maura*, Okamoto et al., 2000; Riley et al., 2014; *Macaca mulatta*, Cooper et al., 2022; *Macaca nemestrina*, Ruppert et al., 2018; *Macaca nigra*, Duboscq & Micheletta, 2023; *Macaca radiata*, Sinha, 2005; *Macaca sinica*, Dittus, 1975; *Macaca sylvanus*, McFarland & Majolo, 2013; *Macaca thibetana*, Li et al., 2020). Japanese macaques (*Macaca fuscata*) were one of the first species for which researchers recognised and followed subjects individually, generating unprecedented levels of detail on their behaviour. The first published record on macaque behaviour dates back to 1956 (Kawamura, 1956). Since then, a body of research has leveraged macaque behavioural data, including comparative research across macaque species (Balasubramaniam et al., 2012, 2020; De Waal & Luttrell, 1989; Sueur et al., 2011; Thierry et al., 2000; Thierry, 2021), to greatly contribute to answering fundamental questions on the evolution, selective pressures and adaptive functions of social behaviour. For instance, macaque research has shed light on the evolutionary drivers and consequences of dominance hierarchies (Balasubramaniam et al., 2012, 2018; Bernstein & Sharpe, 1966; Neumann & Fischer, 2023; Simons et al., 2022; Van Noordwijk & Van Schaik, 1985), the influence of kinship on social structure (Berman & Thierry, 2010; Brent et al., 2017; De Moor et al., 2020; Schülke & Ostner, 2008; Widdig et al., 2016; Yamada, 1963), patterns of social ageing (Almeling et al., 2016; Sadoughi et al., 2024; Siracusa et al., 2022), the ultimate function of social relationships (Campbell et al., 2018; Ellis et al., 2019; McFarland & Majolo, 2013; Micheletta et al., 2012; Riley et al., 2014; Schülke et al., 2010; Singh et al., 2010; Young et al., 2014) and the impact of anthropogenic challenges on social structure (Balasubramaniam et al., 2020; Holzner et al., 2019; Kaburu et al., 2019; Morrow et al., 2019; Testard et al., 2021). Building on these strong foundations, MacaqueNet aims to further promote collaborative and comparative research on macaque social behaviour.

2.2 | Creating MacaqueNet

MacaqueNet originated in 2017 as a small-scale collaborative proposal for one specific study. From there, MacaqueNet developed and expanded to become a global grassroots network of macaque researchers. To mitigate the inherent bias towards researchers from Europe and the United States, and to facilitate the involvement of scientists from macaque-range nations and historically underrepresented research communities we identified further potential collaborators through word of mouth and a literature search on Google Scholar. An initial email was sent to all first and last authors of papers that contained or pointed to the existence of social behavioural data on macaques, inviting them to contribute their data to a collaborative database. We asked the contacted researchers to forward the

invitation to their relevant collaborators and to list those collaborators as co-contributors of the data shared with MacaqueNet, ensuring everyone received proper credit. We also encouraged them to suggest other researchers who might be interested in joining the consortium. This inclusive approach allowed us to engage a broader community of researchers. As a result, the MacaqueNet community currently consists of 106 researchers based at 58 institutes across five continents. Beyond our data contributors, MacaqueNet is open to any interested researcher, regardless of whether they have contributed data. All members receive a quarterly newsletter, can join meetings, request data and propose new research directions.

2.3 | The MacaqueNet relational database

We brought together social behavioural data from independent research teams into a standardised cross-species database. The MacaqueNet database currently contains social interaction data from 1981 to the present for 14 of the 25 recognised species of macaques (Figure 1). At the time of writing, the data have been collected on 21 wild, 22 captive and 18 free-ranging populations (some of which have been observed for several study periods) at a total of 61 field sites, zoos and research centres (Figure 2), resulting in a total of 3972 individual behavioural datapoints (Figure 1).

The full extent of the data available can be explored using the search tool on the MacaqueNet website: <https://macaquetnet.github.io/database/>.

The core of the database consists of sociometric matrices that represent the two primary axes of animal social structure: dyadic affiliative and dyadic agonistic behaviour between individuals, aggregated by study period (Figure 3). The behavioural data comprise three categories of affiliative behaviour—grooming, spending time in body contact and spending time in close spatial proximity—and two categories of agonistic behaviour—contact and non-contact aggression. These data represent the most common behavioural interactions expressed in macaques (Thierry, 2007), forming the backbone of affiliative and dominance relationships, and are therefore collected by most researchers studying macaques. Data have been collected observing one focal individual or (part of a) group at a time, using continuous recording or discrete sampling of dyadic behaviours, recorded as counts or durations (see glossary at <https://macaquetnet.github.io/documentation/> for definitions). Each set of sociometric matrices for a particular group-period (i.e. a given study period for a given group) is accompanied by a subject data file, which contains individual attributes on sex, age and observation effort (Figure 3).

In addition to the behavioural data, the database includes rich metadata on methodology, study populations and research teams.




FIGURE 1 Summary of database content at the time of publication. (a) The MacaqueNet database currently contains data for 14 out of 25 recognised macaque species (note that the depicted phylogenetic tree (Arnold, Matthews & Nunn, 2010) does not include relatively newly described species: *Macaca selai*, *Macaca leucogenys* and *Macaca siberu*). (b) Overview of the number of group-periods (i.e. a given study period for a given group), individuals and sociometric matrices for aggressive and affiliative behaviours for each species. As some individuals have been observed over multiple group-periods, the number of individuals represents the number of unique individual datapoints but not necessarily the number of unique individuals. The dot plot on the right illustrates grooming network densities (the proportion of dyads that were observed grooming at least once), with each dot representing the density for one grooming sociometric matrix.




FIGURE 2 Geographical distribution of research sites in which the data currently present in the MacaqueNet database have been collected. Populations in America and Europe (with the exception of Gibraltar) have been introduced. Living conditions are classified as wild, free-ranging or captive (see glossary at <https://macaque.net/github.io/documentation/> for definitions). Group-periods represent the total number of study periods for all groups at a given research site.

The data and metadata are organised as a relational database (Figure 4), with clear definitions for each variable outlined in a glossary (both the relational database structure and the glossary are available at <https://macaque.net/github.io/documentation/>).

To ensure that the data and metadata are comparable across studies, we set up a standardisation pipeline through which all contributed data ('primary data') is run before it enters the database. Figure 5 provides an overview of the workflow from primary data submission, over the data checks and standardisation pipeline to data requests.

In a nutshell, for each dataset contributed to MacaqueNet, we extract the sociometric matrices for each behaviour, the subject data and the metadata on the study population, observational methods and research team. We perform standard data checks on the extracted sociometric matrices and subject data to ensure there are no missing or unexpected values. Next, we standardise the data by establishing uniform units of measurement and clearly defined categories for metadata on the populations and observation protocols. For instance, we classify observational methods based on three descriptors (see glossary at <https://macaque.net/github.io/documentation/> for full definitions): 'focal_data' (whether the data are collected focusing on one individual or a group of individuals), 'sampling_method' (whether the data are recorded continuously or at set time intervals) and 'data_type' (whether the data are collected as counts or as durations). Similarly, we classify groups based on four descriptors (see glossary for full definitions): 'range_management' (the extent to which a group's range is managed), 'medical_intervention' (whether a group receives medical interventions as part of the

population's management plan), 'food_provisioned' (whether a group is routinely provided with food) and 'human_contact' (whether a group is in regular contact with non-researcher humans). This classification process is essential because terms describing observation methods and population settings often have multiple inconsistent definitions (e.g. 'focal sampling' or 'free-ranging'). Our collaborative approach to building the database was particularly powerful in this regard, as it allowed us to standardise data through direct communication with contributors, thus avoiding misinterpretations. The resulting standardisation ensures consistency across datasets, simplifies filtering for data users based on their specific needs and facilitates accurate comparison and analysis.

Both the primary data and the standardised database are currently stored and managed in GitHub. The data standardisation pipeline is available as an R package (<https://github.com/MacaqueNet/macaquetpipeline>), which provides examples and code for how contributed data are processed before entering the database. The available data can be explored and requested using the search tool on the MacaqueNet website (<https://macaque.net/github.io/database/>), which also provides a version history documenting changes and updates to the database.

2.4 | MacaqueNet is a community that facilitates comparative research

MacaqueNet makes macaque data FAIR (Wilkinson et al., 2016), in four main ways. First, by identifying, accessing and bringing

	a	b	c	d	e	f	g	h	i	j	k	l	m	n	o	p	q	r	s	t	u	v	w	x	y	subject	sex	age	age_cat	obseff_duration	
a	0	0	0	69	0	0	0	96	0	0	0	0	0	9	0	0	0	0	0	0	0	9	0	0	0	a	f	20	adult	9.0	
b	0	0	0	7	0	0	0	0	0	4	0	0	21	0	0	0	0	17	0	0	0	0	25	0	0	b	f	7	adult	9.3	
c	0	0	0	14	0	5	0	0	0	0	0	0	0	0	2	2	0	0	0	0	0	36	0	0	26	c	f	8	adult	8.7	
d	8	0	0	0	0	0	22	0	17	24	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12	d	f	19	adult	8.3	
e	0	0	0	0	0	0	11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	e	f	21	adult	6.7	
f	0	4	4	0	0	0	0	12	8	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	f	f	22	adult	8.6	
g	0	0	0	21	0	0	0	0	41	3	0	0	4	0	0	0	0	0	0	0	0	0	0	0	0	g	f	12	adult	8.3	
h	33	0	0	13	0	0	0	0	0	10	0	100	0	0	0	0	0	0	0	0	0	0	0	0	33	29	h	f	15	adult	7.7
i	0	0	0	75	0	0	0	0	0	0	0	0	0	0	0	4	0	0	0	0	0	43	5	0	0	i	f	10	adult	9.0	
j	0	0	0	0	0	0	0	5	0	0	0	7	2	0	0	0	5	2	2	0	0	25	0	0	16	j	f	11	adult	9.0	
k	0	26	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	24	0	0	0	3	0	0	0	k	f	8	adult	9.3	
l	0	0	0	41	0	0	0	65	0	0	0	0	0	0	5	0	0	0	1	0	16	0	0	0	0	l	f	10	adult	9.0	
m	0	27	0	4	6	0	0	4	33	0	0	0	0	0	10	0	6	0	0	0	25	0	2	16	m	f	7	adult	9.0		
n	0	0	0	5	0	0	0	0	0	0	0	16	0	0	0	0	0	0	0	0	0	0	0	0	0	n	f	25	adult	7.7	
o	0	0	12	11	5	0	0	0	0	0	0	0	0	0	25	0	0	0	0	0	1	6	6	17	o	f	7	adult	9.0		
p	0	1	5	0	0	0	0	0	0	0	0	4	0	10	0	0	18	4	0	0	0	0	5	14	p	f	6	adult	9.0		
q	0	13	0	0	0	0	0	0	20	1	0	0	0	2	0	0	0	0	0	0	0	2	0	10	q	f	6	adult	9.6		
r	0	14	0	0	0	0	0	0	0	17	68	0	3	0	0	38	0	0	0	0	7	0	0	28	0	r	f	6	adult	8.7	
s	15	0	0	0	0	0	0	0	0	4	0	0	1	0	0	13	36	57	0	0	0	0	0	13	33	s	f	6	adult	7.6	
t	0	8	0	0	0	10	41	0	43	0	0	20	0	0	0	0	0	0	0	0	11	0	0	22	t	f	16	adult	9.7		
u	0	0	0	0	0	12	0	0	0	3	0	0	6	0	0	25	0	0	0	0	0	0	12	0	0	u	f	5	adult	8.3	
v	4	0	140	0	0	0	0	0	2	0	0	5	0	0	0	0	0	0	0	0	0	0	0	0	0	v	f	18	adult	9.7	
w	0	22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	w	f	20	adult	8.7		
x	0	0	0	0	0	0	0	0	0	0	0	5	0	4	2	0	3	0	0	0	0	0	0	24	x	f	9	adult	9.6		
y	0	7	0	0	0	0	0	0	0	0	0	21	0	0	0	0	0	0	0	0	0	0	32	0	0	y	f	22	adult	8.3	




FIGURE 3 Example of the data of one dataset as stored in the MacaqueNet database. Top left: A sociometric matrix documenting pairwise interactions in a given group-period. For directed matrices, the actors—the individuals who initiate the behaviour—are listed in the rows, while the receivers—the individuals towards whom the behaviour is directed—are listed in the columns. The matrix entries represent either the total number of times (counts, as depicted here) or the total duration (in seconds) for which an individual in the row performed the behaviour towards the individual in the column for a given study period. Depicted here: Counts of dyadic grooming events in a group of Barbary macaques (*Macaca sylvanus*) from La Montagne des Singes in 2017. Top right: A subject table listing all individuals observed in a given group-period, along with individual attributes: Sex, age and observation effort, here duration of observation (in hours). Bottom left: An illustration of the network representing the data in the sociometric matrix. Each blue circle represents a subject, green lines between circles represent dyadic affiliation strength, here quantified as the dyadic rate of grooming. Bottom right: A picture of Barbary macaques grooming (photo credit: Anthony Poynton).

together data from multiple sources into one centralised place <https://macaquetnet.github.io/database/>. Instead of creating a database of summarised traits with one data point per species,

this approach consolidates the original data itself and creates a more flexible resource for future use. Moreover, with several species represented by multiple populations and/or study periods,




FIGURE 4 Set-up of the relational MacaqueNet database. All variables are defined in the glossary (available at <https://macaque.net.github.io/documentation/>). Arrows indicate how each table is linked to at least one other table in the database through unique identifiers (the 'id' columns in each table). Each entry in the 'datakey' table links to a sociometric matrix with all instances of a specific behaviour category for one group period (as depicted in [Figure 3](#) top left). Part of the 'datakey' table is only relevant for the data import pipeline. This figure was made using dbdiagram (<https://www.dbdiagram.io/>).

both between-species and within-species variation are captured (Mocha et al., 2024; Sinha, 2005). The latter is often overlooked in comparative analyses but can be a significant source of error if not accounted for (Sandel et al., 2016; Schradin, 2013). Second, by developing a common vocabulary for the data and metadata in the MacaqueNet database, with openly accessible definitions (Borries et al., 2016; Urbano & Cagnacci, 2021). Third, by standardising data across studies and research sites together with the data contributors who understand the intricacies and subtleties of the data (Schneider et al., 2019; Urbano & Cagnacci, 2021). Fourth, by providing transparent and replicable annotation of how primary data were processed so that the original data can always be recovered and processed in different ways if future researchers want to make other decisions (Schneider et al., 2019; Urbano & Cagnacci, 2021). As a result, MacaqueNet provides a meticulously curated database with data readily available for future comparative studies (Borries et al., 2016; Gomes et al., 2022).

MacaqueNet extends beyond mere data sharing. By linking researchers across different research groups and institutions, MacaqueNet facilitates a global exchange of ideas and fosters the development of new research projects. To provide an opportunity for all the current data contributors to meet, discuss their research and learn about ongoing projects, we organised a virtual global symposium 'Weaving the MacaqueNet' in November 2021. During

the symposium, future directions for MacaqueNet were discussed and agreed on. As a first step in the establishment of an infrastructure that permits researchers to communicate and coordinate their research, we set up a website (<https://macaque.net.github.io>), a mailing list and a group messaging channel on Slack. The second 'Weaving the MacaqueNet' symposium was held in August 2023 at the Joint Conference of the International Primatological Society and Malaysian Primatological Society, where the MacaqueNet community got the opportunity to meet in person. At that meeting, we discussed the expansion of MacaqueNet to include more types of data, thereby increasing MacaqueNet's ability to address a range of questions on social evolution. As a result, efforts are currently underway to include ecological, life history and genetic data to complement the existing behavioural data. To promote transparency, we also agreed to make all meetings associated with MacaqueNet open to all MacaqueNet members, in addition to disseminating a quarterly newsletter outlining new projects and discussion points.

New projects can be easily proposed, and data requested with a brief project proposal via the website (<https://macaque.net.github.io/database/>). The proposal is forwarded to the data contributors, who assess whether they can provide their data for the proposed project and determine their usage terms (see MacaqueNet terms of use at <https://macaque.net.github.io/documentation/>). New types of data to be added to the database can also be suggested




FIGURE 5 Overview of the workflow from primary data submission, over the data checks and standardisation pipeline to data requests. All contributed data contain several sociometric matrices along with corresponding subject data and metadata for at least one group-period for a specific species.

on the website. These ideas and their potential implementation are discussed at each MacaqueNet symposium.

The code for the website, the glossary, the relational database structure, the workflow and the terms of use are openly accessible in a GitHub repository (<https://github.com/MacaqueNet>). Together, these resources provide a fully replicable model for the development of cross-species collaborative databases.

3 | DISCUSSION

In this paper, we advocate for the collaborative building of FAIR cross-species databases as a novel and promising tool in comparative research. Understanding the ecological and evolutionary drivers of variation in social behaviour requires comparing taxa with distinct life histories, ecology and interlinked evolutionary histories (Kappeler et al., 2019; Lukas & Clutton-Brock, 2018; Pinter-Wollman et al., 2013). Yet, despite decades worth of social behavioural data (Sheldon et al., 2022), substantial challenges have impeded systematic comparative research (Albery et al., 2024; De Moor et al., 2024), one of which is the creation of truly comparable databases (Borries et al., 2016). We propose that large-scale collaborative efforts, where data and expertise are shared among researchers, can provide a solution to this issue and help facilitate reliable cross-species comparisons. To inspire and facilitate more collaborative efforts in behavioural research, we share our experience in developing MacaqueNet and its cross-species database.

Building on the foundations of previous big-team projects that have been instrumental in shaping MacaqueNet (including but not restricted to SPI-Birds: Culina et al., 2020; ManyPrimates: ManyPrimates et al., 2019), we highlight three substantial advantages for the scientific community to join forces and pool their data into FAIR databases.

First, by creating an enduring infrastructure for collaboration, big-team science efforts represent a crucial interface linking existing data and data users (Culina et al., 2020; Urbano & Cagnacci, 2021). They ensure that valuable data, which might have otherwise remained largely unused, are easier to discover, repurpose, and synthesise (Gonzalez & Peres-Neto, 2015; Purgar et al., 2022; Wallis et al., 2013) and help pinpoint areas that would benefit most from additional data collection. Moreover, rather than summarising the data into single datapoints, building a database that brings the standardised original data together creates a much more flexible resource. Such a database represents variation at different biological levels, can easily keep growing to include more data as they become available, and can be subset and summarised tailored to specific research questions.

Second, much of the existing social behaviour data are the result of years of effort and funding acquisition to set up the necessary logistics and to collect data on often wild and endangered animals. Big-team science can set up guidelines and regulations to ensure that sharing is done in such a way that it considers individual researchers' interests, and their ability to acquire funding to keep collecting new

data. Transparent but controlled accessibility, where data contributors retain full ownership of their data, best balances benefits to the scientific community while protecting data contributors, as proven successful in other big-team science endeavours (Culina et al., 2020; Strier et al., 2010; Urbano & Cagnacci, 2021). Creating a community rather than just a database also offers data contributors networking opportunities and access to others' data, fostering collaborative research possibilities (Urbano & Cagnacci, 2021).

Third, social behavioural data are particularly complex, with many researcher degrees of freedom involved in how the data are defined and what methods of observation are used (Borries et al., 2016). Taking a collaborative approach to building comparative databases allows for data standardisation to be done through direct interaction with data contributors, thus ensuring clarity around the intricacies of the data and averting inadvertent misinterpretations (Mocha et al., 2024; Schradin, 2017; Urbano & Cagnacci, 2021). This process is essential for providing database users with the necessary information to determine which data are comparable for their specific research questions and how to appropriately account for differences when conducting comparisons (De Moor et al., 2024).

The rise of collaborative databases in a variety of fields is a testament to the enthusiasm of researchers to join forces and share data and resources to facilitate comparative and interdisciplinary science, an emerging and promising way of doing research (Coles et al., 2022). Large-scale collaborative endeavours like MacaqueNet allow for the exploration of innovative and comparative questions no single research group could address individually, make testing the reproducibility of research findings across contexts possible, and make data access more equitable by setting up FAIR (Wilkinson et al., 2016) and transparent data sharing policies. Sustaining such initiatives necessitates continual financial and personnel support for database storage and maintenance as well as community engagement. This underscores the importance for funding agencies and the broader academic system to prioritise and fund these endeavours accordingly (Gomes et al., 2022). While the creation of databases on behavioural data comes with substantial challenges, we truly believe this is the way forward to harness the incredible comparative potential of the wealth of existing behavioural data. We hope that MacaqueNet, along with similar endeavours, can catalyse large-scale collaborative research on animal behaviour.

AUTHOR CONTRIBUTIONS

Julie Duboscq, Christof Neumann, Lauren J. N. Brent, Julia Ostner and Oliver Schülke generated the idea to compile data for comparative research. Julie Duboscq and Christof Neumann instigated the collaborative effort, with Lauren J. N. Brent and Delphine De Moor formalising it into the grassroots network 'MacaqueNet'; Delphine De Moor, Lauren J. N. Brent, Julie Duboscq and Macaela Skelton built a community of researchers. Delphine De Moor, Julie Duboscq and Macaela Skelton gathered the data. Christof Neumann and Delphine De Moor created the database package and standardised the data. Delphine De Moor wrote the manuscript, with input from

Lauren J. N. Brent, Julie Duboscq, Christof Neumann, Julia Ostner, Oliver Schülke and Krishna N. Balasubramaniam. Delphine De Moor, Macaela Skelton and Christof Neumann created the figures. All authors contributed data to the MacaqueNet database, provided critical feedback on the manuscript's structure and content, and gave their final approval for publication.

AFFILIATIONS

¹School of Psychology, Centre for Research in Animal Behaviour, University of Exeter, Exeter, UK; ²Research Group for Human Biology and Primate Cognition, Biology Institute, University of Leipzig, Leipzig, Germany; ³Department of Comparative Cultural Psychology, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany; ⁴Faculty of Biology, Institute of Human Biology and Evolution, Adam Mickiewicz University, Poznań, Poland; ⁵Faculty of Science & Engineering, School of Life Sciences, Anglia Ruskin University, Cambridge, UK; ⁶Department of Population Health & Reproduction, School of Veterinary Medicine, University of California at Davis, Davis, California, USA; ⁷Laboratoire de Neurosciences Cognitives et Adaptatives, Strasbourg, France; ⁸Centre de Primatologie de l'Université de Strasbourg, Niederhausbergen, France; ⁹Domestication Lab, Department of Interdisciplinary Life Sciences, Konrad Lorenz Institute of Ethology, University of Veterinary Medicine Vienna, Vienna, Austria; ¹⁰Program in Evolution, Ecology and Behavior, Department of Anthropology, University at Buffalo, Buffalo, New York, USA; ¹¹Division of Natural Sciences, Engineering and Mathematics, Department of Biology, University of St. Thomas, Houston, Texas, USA; ¹²Animal Behavior & Cognition, Department of Biology, Utrecht University, Utrecht, The Netherlands; ¹³Department of Psychology and the California National Primate Research Center, University of California at Davis, Davis, California, USA; ¹⁴Primate Research Group, Research Unit SPHERES, The University of Liège, Liège, Belgium; ¹⁵Institute of Ethology and Anthropology, Russian Academy of Sciences, Moscow, Russia; ¹⁶Wildlife Conservation Research Unit (WildCRU), University of Oxford, Oxford, UK; ¹⁷Department of Sciences, Roma Tre University, Rome, Italy; ¹⁸Conservation Science and Outreach, North of England Zoological Society, Cheshire, UK; ¹⁹Animal Behaviour and Cognition Programme, National Institute of Advanced Studies, Indian Institute of Science Campus, Bengaluru, India; ²⁰Department of Psychology, University of Tennessee Knoxville, Knoxville, Tennessee, USA; ²¹Science Department, North of England Zoological Society, Chester, UK; ²²FES Zaragoza, National Autonomous University of Mexico, Mexico City, Mexico; ²³National Institute of Psychiatry Ramón de la Fuente Muñiz, Mexico City, Mexico; ²⁴School of Psychology, Mexico City, Mexico; ²⁵Fondazione Ethoikos, Convento dell'Osservanza, Radicondoli, Italy; ²⁶Yale Child Study Center, Yale School of Medicine, New Haven, Connecticut, USA; ²⁷Animal Behaviour and Cognition Programme, National Institute of Advanced Studies, Bengaluru, India; ²⁸Department of Zoology, Kannur University, Kannur, India; ²⁹Department of Anthropology, Durham Cultural Evolution Research Centre, Durham University, Durham, UK; ³⁰Cognitive Ethology Lab, German Primate Center, Göttingen, Germany; ³¹Department for Primate Cognition, Georg-August-University Göttingen, Göttingen, Germany; ³²Primate Cognition, Göttingen, Germany; ³³Department of Anthropology, Durham University, Durham, UK; ³⁴Department of Behavioral and Cognitive Biology, University of Vienna, Vienna, Austria; ³⁵Department of Anthropology, Princeton University, Princeton, New Jersey, USA; ³⁶The Long-Tailed Macaque Project, Sorø, Denmark; ³⁷Behavioural Ecology Group, Department of Biology, University of Copenhagen, Copenhagen, Denmark; ³⁸Oxford Wildlife Trade Research Group, Oxford Brookes University, Oxford, UK; ³⁹Behavioural Ecology Research Group, Institute of Biology, University of Leipzig, Leipzig, Germany; ⁴⁰Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany; ⁴¹School of Biological Sciences, Universiti Sains Malaysia, Gelugor, Malaysia; ⁴²School of Animal, Rural and Environmental Sciences, Nottingham Trent University, Southwell, UK; ⁴³Department of Zoology, Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic; ⁴⁴Sálim Ali Centre for Ornithology and Natural History, Coimbatore, India; ⁴⁵Département d'Anthropologie, Faculté des Arts et des Sciences, Udem, Montréal, Quebec,

Canada; ⁴⁶Department of Psychology, University of Lethbridge, Lethbridge, Alberta, Canada; ⁴⁷Whitelands College, Roehampton University, London, UK; ⁴⁸International Collaborative Research Center for Huangshan Biodiversity and Tibetan Macaque Behavioral Ecology, School of Resource and Environmental Engineering, Anhui University, Hefei, China; ⁴⁹School of Resource and Environmental Engineering, Anhui University, Hefei, China; ⁵⁰ZooParc de Beauval et Beauval Nature, Saint-Aignan, France; ⁵¹Wildlife Research Center, Kyoto University, Inuyama, Japan; ⁵²School of Psychology, University of Lincoln, Lincoln, UK; ⁵³Wildlife Park Goldau, Goldau, Switzerland; ⁵⁴Austrian Research Center for Primatology, Ossiach, Austria; ⁵⁵Forest Conservation Department, Hasanuddin University, Makassar, Indonesia; ⁵⁶NTU Psychology, Nottingham Trent University, Nottingham, UK; ⁵⁷Applied Behavioural Ecology and Ecosystems Research Unit, University of South Africa, Pretoria, South Africa; ⁵⁸Department of Psychology, Centre for Comparative and Evolutionary Psychology, University of Portsmouth, Portsmouth, UK; ⁵⁹Srishti Manipal Institute of Arts Design and Technology, Bengaluru, India; ⁶⁰CLLE, Université de Toulouse, CNRS, Toulouse, France; ⁶¹Department of Anthropology, University of Georgia, Athens, Georgia, USA; ⁶²Institute for Evolutionary Ecology and Conservation Genomics, Ulm University, Ulm, Germany; ⁶³Department of Biology, University of Pisa, Pisa, Italy; ⁶⁴Laboratoire de Psychologie Sociale et Cognitive, Centre National de la Recherche Scientifique et Université Clermont-Auvergne, Clermont-Ferrand, France; ⁶⁵Istituto di Scienze e Tecnologie della Cognizione, Consiglio Nazionale delle Ricerche, Rome, Italy; ⁶⁶Foundation for Ecological Research, Advocacy and Learning Morattandi, Villupuram, India; ⁶⁷Anthropology Program, School of Anthropology, Political Science and Sociology, Southern Illinois University Carbondale, Carbondale, USA; ⁶⁸Department of Anthropology, San Diego State University, San Diego, California, USA; ⁶⁹Department of Life Sciences, University of Roehampton, London, UK; ⁷⁰Primate Social Evolution Group, German Primate Center, Leibniz Institute for Primate Research, Göttingen, Germany; ⁷¹Behavioral Ecology Department, University of Göttingen, Göttingen, Germany; ⁷²Anthropology, Central Washington University, Ellensburg, Washington, USA; ⁷³School of Human Evolution and Social Change and Institute of Human Origins, Arizona State University, Tempe, Arizona, USA; ⁷⁴Biopsychology Laboratory, Institution of Excellence, University of Mysore, Mysuru, India; ⁷⁵Institut pluridisciplinaire Hubert Curien (UMR 7178), Centre national de la recherche scientifique, Université de Strasbourg, Strasbourg, France; ⁷⁶Natural Resources Institute, University of Greenwich, Kent, UK; ⁷⁷Research Centre for Evolutionary Anthropology & Palaeoecology, School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, UK; ⁷⁸Department of Anthropology, Northern Arizona University, Flagstaff, Arizona, USA; ⁷⁹Department of Primate Behavior and Evolution, Max-Planck Institute for Evolutionary Anthropology, Leipzig, Germany; ⁸⁰Department of Neurosurgery, Emory University, Atlanta, Georgia, USA; ⁸¹School of Life Sciences, Anhui University, Hefei, Anhui, China; ⁸²Nuremberg Zoo, Nürnberg, Germany and ⁸³UMR7206 Eco-Anthropology, CNRS-MNHN-Université Paris Cité, Paris, France

ACKNOWLEDGEMENTS

We thank all members of MacaqueNet for thoughtful discussion during the development of this collaborative endeavour. We are especially grateful for the enormous efforts of all current and past researchers who have contributed to generating the data in the MacaqueNet database. We thank Antica Culina for sharing her insights about creating the SPI-Birds Network and Database. This work was supported by a European Research Council Consolidator grant (FriendOrigins—864461) to L.J.N.B. and by an Audacity Fund of the Leibniz ScienceCampus Primate Cognition (LSC-AF2023_03).

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no competing interests.

DATA AVAILABILITY STATEMENT

The code for the data standardisation pipeline on the available datasets, along with resources, such as the Glossary, the workflow and the MacaqueNet Terms of Use, are openly accessible in the MacaqueNet GitHub repository (<https://github.com/MacaqueNet/>). MacaqueNet data can be requested following the requesting protocol through the MacaqueNet website (<https://macaquenet.github.io/database/>).

STATEMENT ON INCLUSION

One of MacaqueNet's primary objectives is to foster collaboration among macaque researchers worldwide. The geographic distribution of MacaqueNet and the authors involved in this manuscript already reflects a relatively diverse group of researchers. However, we remain committed to enhancing the inclusion of researchers from macaque-range countries in Asia and Northern Africa. We aspire to see the MacaqueNet community play a pivotal role in achieving this goal by establishing a network where funding opportunities and access to data are more equitably distributed. Furthermore, we aim to provide a platform within MacaqueNet to showcase the remarkable work conducted by researchers in macaque-range countries and to catalyse collaborations on a global scale.

ORCID

Delphine De Moor  <https://orcid.org/0000-0003-2474-6125>
Macaela Skelton  <https://orcid.org/0009-0002-7166-2995>
Federica Amici  <https://orcid.org/0000-0003-3539-1067>
Malgorzata E. Arlet  <https://orcid.org/0000-0001-9890-5999>
Krishna N. Balasubramaniam  <https://orcid.org/0000-0002-2101-0588>
Sébastien Ballesta  <https://orcid.org/0000-0002-7854-5735>
Andreas Berghänel  <https://orcid.org/0000-0002-3317-3392>
Carol M. Berman  <https://orcid.org/0000-0001-7328-2588>
Sofia K. Bernstein  <https://orcid.org/0000-0003-4093-6922>
Debottam Bhattacharjee  <https://orcid.org/0000-0002-4555-7983>
Eliza Bliss-Moreau  <https://orcid.org/0000-0002-0740-5612>
Fany Brotcorne  <https://orcid.org/0000-0003-2780-3631>
Marina Butovskaya  <https://orcid.org/0000-0002-5528-0519>
Liz A. D. Campbell  <https://orcid.org/0000-0002-8302-7430>
Monica Carosi  <https://orcid.org/0000-0003-1377-2942>
Mayukh Chatterjee  <https://orcid.org/0000-0002-2794-4189>
Matthew A. Cooper  <https://orcid.org/0000-0001-9645-1671>
Veronica B. Cowl  <https://orcid.org/0000-0002-7928-4549>
Claudio De la O  <https://orcid.org/0000-0001-5910-2130>
Arianna De Marco  <https://orcid.org/0000-0002-3681-0254>
Amanda M. Dettmer  <https://orcid.org/0000-0003-1360-3567>
Ashni K. Dhawale  <https://orcid.org/0000-0002-7185-9870>
Joseph J. Erinjeri  <https://orcid.org/0000-0002-5690-838X>
Julia Fischer  <https://orcid.org/0000-0002-5807-0074>
Iván García-Nisa  <https://orcid.org/0000-0003-4150-2480>
Gwennan Giraud  <https://orcid.org/0000-0002-6022-2663>
Roy Hammer  <https://orcid.org/0000-0003-2747-4612>

Malene F. Hansen  <https://orcid.org/0000-0002-4152-7947>
Anna Holzner  <https://orcid.org/0000-0002-1442-0802>
Stefano Kaburu  <https://orcid.org/0000-0001-7456-3269>
Martina Konečná  <https://orcid.org/0000-0002-2650-1597>
Honnnavalli N. Kumara  <https://orcid.org/0000-0002-7958-6395>
Jean-Baptiste Leca  <https://orcid.org/0000-0002-2371-671X>
Mathieu Legrand  <https://orcid.org/0009-0009-5880-3424>
Julia Lehmann  <https://orcid.org/0000-0001-7278-1705>
Jin-Hua Li  <https://orcid.org/0000-0003-4859-5763>
Andrew MacIntosh  <https://orcid.org/0000-0002-9136-7099>
Bonaventura Majolo  <https://orcid.org/0000-0002-0235-3040>
Laëtitia Maréchal  <https://orcid.org/0000-0003-3499-9134>
Pascal R. Marty  <https://orcid.org/0000-0003-3040-464X>
Jorg J. M. Massen  <https://orcid.org/0000-0002-1630-9606>
Risma Illa Maulany  <https://orcid.org/0000-0001-5061-6725>
Brenda McCowan  <https://orcid.org/0000-0002-2278-418X>
Richard McFarland  <https://orcid.org/0000-0001-8245-9269>
Jérôme Micheletta  <https://orcid.org/0000-0002-4480-6781>
Partha S. Mishra  <https://orcid.org/0000-0002-6328-5897>
Shahrul A. M. Sah  <https://orcid.org/0000-0002-7553-6980>
Sandra Molesti  <https://orcid.org/0000-0002-3523-0055>
Kristen S. Morrow  <https://orcid.org/0000-0002-8373-4514>
Nadine Müller-Klein  <https://orcid.org/0000-0002-8194-936X>
Putu Oka Ngakan  <https://orcid.org/0000-0002-1383-2661>
Elisabetta Palagi  <https://orcid.org/0000-0002-2038-4596>
Odile Petit  <https://orcid.org/0000-0002-4340-7199>
Lena S. Pflüger  <https://orcid.org/0000-0003-4494-4970>
Eugenia Polizzi di Sorrentino  <https://orcid.org/0000-0002-5554-9813>
Ulrich H. Reichard  <https://orcid.org/0000-0002-5564-3922>
Erin P. Riley  <https://orcid.org/0000-0002-2679-0595>
Alan V. Rincon  <https://orcid.org/0000-0001-6181-0152>
Nadine Ruppert  <https://orcid.org/0000-0002-9760-0058>
Baptiste Sadoughi  <https://orcid.org/0000-0001-5626-3318>
Kumar Santhosh  <https://orcid.org/0000-0002-0368-6437>
Gabriele Schino  <https://orcid.org/0000-0002-5011-4896>
Lori K. Sheeran  <https://orcid.org/0000-0002-8903-9379>
Joan B. Silk  <https://orcid.org/0000-0002-1286-707X>
Mewa Singh  <https://orcid.org/0000-0002-9198-0192>
Anindya Sinha  <https://orcid.org/0000-0001-5623-711X>
Sebastian Sosa  <https://orcid.org/0000-0002-5087-9135>
Cédric Sueur  <https://orcid.org/0000-0001-8206-2739>
Barbara Tiddi  <https://orcid.org/0000-0002-1832-835X>
Patrick J. Tkaczynski  <https://orcid.org/0000-0003-3207-2132>
Florian Trebouet  <https://orcid.org/0000-0001-9531-1482>
Anja Widdig  <https://orcid.org/0000-0003-1772-8770>
Jamie Whitehouse  <https://orcid.org/0000-0003-2607-5492>
Lauren J. Wooddell  <https://orcid.org/0000-0001-6923-1219>
Dong-Po Xia  <https://orcid.org/0000-0002-1266-2285>
Lorenzo von Fersen  <https://orcid.org/0000-0002-3451-0530>
Christopher Young  <https://orcid.org/0000-0001-8919-2093>
Oliver Schülke  <https://orcid.org/0000-0003-0028-9425>
Julia Ostner  <https://orcid.org/0000-0001-6871-9976>

Christof Neumann  <https://orcid.org/0000-0002-0236-1219>

Julie Duboscq  <https://orcid.org/0000-0002-6342-6709>

Lauren J. N. Brent  <https://orcid.org/0000-0002-1202-1939>

REFERENCES

- Albert, A., Huynen, M., Savini, T., & Hambuckers, A. (2013). Influence of food resources on the ranging pattern of Northern pig-tailed macaques (*Macaca leonina*). *International Journal of Primatology*, 34(4), 696–713. <https://doi.org/10.1007/s10764-013-9690-z>
- Albery, G. F., Bansal, S., & Silk, M. J. (2024). Comparative approaches in social network ecology. *Ecology Letters*, 27(1), e14345. <https://doi.org/10.1111/ele.14345>
- Almeling, L., Hammerschmidt, K., Sennhenn-Reulen, H., Freund, A. M., & Fischer, J. (2016). Motivational shifts in aging monkeys and the origins of social selectivity. *Current Biology*, 26(13), 1744–1749. <https://doi.org/10.1016/j.cub.2016.04.066>
- Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: A new online resource for primate phylogeny. *Evolutionary Anthropology: Issues, News, and Reviews*, 19(3), 114–118. <https://doi.org/10.1002/evan.20251>
- Balasubramaniam, K. N., Beisner, B. A., Berman, C. M., De Marco, A., Duboscq, J., Koirala, S., Majolo, B., MacIntosh, A. J., McFarland, R., Molesti, S., Ogawa, H., Petit, O., Schino, G., Sosa, S., Sueur, C., Thierry, B., de Waal, F. B. M., & McCowan, B. (2018). The influence of phylogeny, social style, and sociodemographic factors on macaque social network structure. *American Journal of Primatology*, 80(1), e22727. <https://doi.org/10.1002/ajp.22727>
- Balasubramaniam, K. N., Dittmar, K., Berman, C. M., Butovskaya, M., Cooper, M. A., Majolo, B., Ogawa, H., Schino, G., Thierry, B., & de Waal, F. B. M. (2012). Hierarchical steepness and phylogenetic models: Phylogenetic signals in *Macaca*. *Animal Behaviour*, 83(5), 1207–1218. <https://doi.org/10.1016/j.anbehav.2012.02.012>
- Balasubramaniam, K. N., Marty, P. R., Samartino, S., Sobrino, A., Gill, T., Ismail, M., Saha, R., Beisner, B. A., Kaburu, S. S. K., Bliss-Moreau, E., Arlet, M. E., Ruppert, N., Ismail, A., Sah, S. A. M., Mohan, L., Rattan, S. K., Kodandaramaiah, U., & McCowan, B. (2020). Impact of individual demographic and social factors on human-wildlife interactions: A comparative study of three macaque species. *Scientific Reports*, 10(1), 21991. <https://doi.org/10.1038/s41598-020-78881-3>
- Berman, C. M., & Thierry, B. (2010). Variation in kin bias: Species differences and time constraints in macaques. *Behaviour*, 147(13–14), 1863–1887. <https://doi.org/10.1163/000579510x539691>
- Bernstein, I. S., & Sharpe, L. G. (1966). Social roles in a rhesus monkey group. *Behaviour*, 26(1–2), 91–104. <https://doi.org/10.1163/156853966X00038>
- Borries, C., Sandel, A. A., Koenig, A., Fernandez-Duque, E., Kamilar, J. M., Amoroso, C. R., Barton, R. A., Bray, J., Di Fiore, A., Gilby, I. C., Gordon, A. D., Mundry, R., Port, M., Powell, L. E., Pusey, A. E., Spriggs, A., & Nunn, C. L. (2016). Transparency, usability, and reproducibility: Guiding principles for improving comparative databases using primates as examples. *Evolutionary Anthropology*, 25(5), 232–238. <https://doi.org/10.1002/evan.21502>
- Brent, L. J. N., Ruiz-Lambides, A., & Platt, M. L. (2017). Family network size and survival across the lifespan of female macaques. *Proceedings of the Royal Society B*, 284(1854), 20170515. <https://doi.org/10.1098/rspb.2017.0515>
- Campbell, L. A. D., Tkaczynski, P. J., Lehmann, J., Mouna, M., & Majolo, B. (2018). Social thermoregulation as a potential mechanism linking sociality and fitness: Barbary macaques with more social partners form larger huddles. *Scientific Reports*, 8(1), 6074. <https://doi.org/10.1038/s41598-018-24373-4>
- Coles, N. A., Hamlin, J. K., Sullivan, L. L., Parker, T. H., & Altschul, D. (2022). Build up big-team science. *Nature*, 601, 505–507. <https://doi.org/10.1038/d41586-022-00150-2>
- Collier, M., Albery, G. F., McDonald, G. C., & Bansal, S. (2022). Pathogen transmission modes determine contact network structure, altering other pathogen characteristics. *Proceedings of the Royal Society B*, 289(1989), 20221389. <https://doi.org/10.1098/rspb.2022.1389>
- Cooper, E. B., Brent, L. J. N., Snyder-Mackler, N., Singh, M., Sengupta, A., Khatiwada, S., Malaivijitnond, S., Qi Hai, Z., & Higham, J. P. (2022). The natural history of model organisms: The rhesus macaque as a success story of the Anthropocene. *eLife*, 11, e78169. <https://doi.org/10.7554/eLife.78169>
- Cords, M. (2012). The behavior, ecology, and social evolution of cercopithecine monkeys. In J. C. Mitani, J. Call, P. M. Kappeler, R. A. Palombit, & J. B. Silk (Eds.), *The evolution of primate societies* (pp. 91–112). Springer.
- Culina, A., Adriaenssens, F., Bailey, L. D., Burgess, M. D., Charmantier, A., Cole, E. F., Eeva, T., Matthyssen, E., Nater, C. R., Sheldon, B. C., Sæther, B.-E., Vriend, S. J. G., Zajkova, Z., Adamik, P., Aplin, L. M., Angulo, E., Artemyev, A., Barba, E., Barišić, S., ... Visser, M. E. (2020). Connecting the data landscape of long-term ecological studies: The SPI-Birds data hub. *Journal of Animal Ecology*, 90(9), 2147–2160. <https://doi.org/10.1111/1365-2656.13388>
- Culina, A., Baglioni, M., Crowther, T. W., Visser, M. E., Woutersen-Windhouwer, S., & Manghi, P. (2018). Navigating the unfolding open data landscape in ecology and evolution. *Nature Ecology & Evolution*, 2(3), 420–426. <https://doi.org/10.1038/s41559-017-0458-2>
- De Moor, D., Brent, L. J. N., Silk, M. J., & Brask, J. B. (2024). Layers of latency in social networks and their implications for comparative analyses. *EcoEvoRxiv*. <https://doi.org/10.32942/X2G894>
- De Moor, D., Roos, C., Ostner, J., & Schülke, O. (2020). Bonds of bros and brothers: Kinship and social bonding in postdispersal male macaques. *Molecular Ecology*, 29(17), 3346–3360. <https://doi.org/10.1111/mec.15560>
- De Waal, F. B. M., & Luttrell, L. M. (1989). Toward a comparative socioecology of the genus *Macaca*: Different dominance styles in rhesus and stump-tail monkeys. *American Journal of Primatology*, 19(2), 83–109. <https://doi.org/10.1002/ajp.1350190203>
- Dittus, W. P. J. (1975). Population dynamics of the toque monkey, *Macaca sinica*. In R. H. Tuttle (Ed.), *Socioecology and psychology of primates* (Vol. 42, pp. 125–151). Mouton Publishers.
- Duboscq, J., & Micheletta, J. (2023). *Macaca nigra* in the spotlight: Accounting for diversity in behavior, ecology, and conservation in primates. *International Journal of Primatology*, 44, 1026–1041. <https://doi.org/10.1007/s10764-023-00386-3>
- Edwards, J. L., Lane, M. A., & Nielsen, E. S. (2000). Interoperability of biodiversity databases: Biodiversity information on every desktop. *Science*, 289(5488), 2312–2314. <https://doi.org/10.1126/science.289.5488.2312>
- Ellis, S., Snyder-Mackler, N., Ruiz-Lambides, A., Platt, M. L., & Brent, L. J. N. (2019). Deconstructing sociality: The types of social connections that predict longevity in a group-living primate. *Proceedings of the Royal Society B*, 286(1917), 20191991. <https://doi.org/10.1098/rspb.2019.1991>
- Fleagle, J. G. (2013). Chapter 6—Old world monkeys. In J. G. Fleagle (Ed.), *Primate adaptation and evolution* (3rd ed., pp. 119–150). Academic Press. <https://doi.org/10.1016/B978-0-12-378632-6.00006-9>
- Forscher, P. S., Wagenmakers, E., Coles, N. A., Silan, M. A., Dutra, N., Basnight-Brown, D., & IJzerman, H. (2022). The benefits, barriers, and risks of big-team science. *Perspectives on Psychological Science*, 18, 607–623. <https://doi.org/10.1177/17456916221082970>
- Gagliardi, B. M., St-Amant, N., & Dakin, R. (2023). Social network structure scales with group size in a multi-species analysis. *bioRxiv*. <https://doi.org/10.1101/2023.08.28.555173>
- Gomes, D. G. E., Pottier, P., Crystal-Ornelas, R., Hudgins, E. J., Foroughirad, V., Sanchez-Reyes, L. L., Turba, R., Martinez, P.

- A., Moreau, D., Bertram, M. G., Smout, C. A., & Gaynor, K. M. (2022). Why don't we share data and code? Perceived barriers and benefits to public archiving practices. *Proceedings of the Royal Society B*, 289(1987), 20221113. <https://doi.org/10.1098/rspb.2022.1113>
- Gonzalez, A., & Peres-Neto, P. R. (2015). Act to staunch loss of research data. *Nature*, 520(7548), 436. <https://doi.org/10.1038/520436c>
- Holzner, A., Ruppert, N., Swat, F., Schmidt, M., Weiss, B. M., Villa, G., Mansor, A., Mohd Sah, S. A., Engelhardt, A., Kuhl, H., & Widdig, A. (2019). Macaques can contribute to greener practices in oil palm plantations when used as biological pest control. *Current Biology*, 29(20), R1066–R1067. <https://doi.org/10.1016/j.cub.2019.09.011>
- Jones, K. E., Bielby, J., Cardillo, M., Fritz, S. A., O'Dell, J., Orme, C. D. L., Safi, K., Sechrest, W., Boakes, E. H., Carbone, C., Connolly, C., Cutts, M. J., Foster, J. K., Grenyer, R., Habib, M., Plaster, C. A., Price, S. A., Rigby, E. A., Rist, J., ... Michener, W. K. (2009). PanTHERIA: A species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology*, 90(9), 2648. <https://doi.org/10.1890/08-1494.1>
- Kaburu, S. S. K., Marty, P. R., Beisner, B., Balasubramaniam, K. N., Bliss-Moreau, E., Kaur, K., Mohan, L., & McCowan, B. (2019). Rates of human-macaque interactions affect grooming behavior among urban-dwelling rhesus macaques (*Macaca mulatta*). *American Journal of Physical Anthropology*, 168(1), 92–103. <https://doi.org/10.1002/ajpa.23722>
- Kappeler, P. M., Clutton-Brock, T., Shultz, S., & Lukas, D. (2019). Social complexity: Patterns, processes, and evolution. *Behavioral Ecology and Sociobiology*, 73(1), 5. <https://doi.org/10.1007/s00265-018-2613-4>
- Kawamura, S. (1956). The spatial distribution test of individuals of a natural group of the Japanese macaque. *The Annual of Animal Psychology*, 6, 1–10. <https://doi.org/10.2502/janip1944.6.1>
- Kays, R., Davidson, S. C., Berger, M., Bohrer, G., Fiedler, W., Flack, A., Hirt, J., Hahn, C., Gauggel, D., Russell, B., Kölsch, A., Lohr, A., Partecke, J., Quetting, M., Safi, K., Scharf, A., Schneider, G., Lang, I., Schaeffelhut, F., ... Wikelski, M. (2021). The Movebank system for studying global animal movement and demography. *Methods in Ecology and Evolution*, 13(2), 419–431. <https://doi.org/10.1111/2041-210x.13767>
- Klein, R. A., Ratliff, K. A., Vianello, M., Adams, R. B., Bahnik, Š., Bernstein, M. J., Bocian, K., Brandt, M. J., Brooks, B., Brumbaugh, C. C., Cemalcilar, Z., Chandler, J., Cheong, W., Davis, W. E., Devos, T., Eisner, M., Frankowska, N., Furrow, D., Galliani, E. M., ... Nosek, B. A. (2014). Investigating variation in replicability. *Social Psychology*, 45(3), 142–152. <https://doi.org/10.1027/1864-9335/a000178>
- Kranstauber, B., Cameron, A., Weinzerl, R., Fountain, T., Tilak, S., Wikelski, M., & Kays, R. (2011). The Movebank data model for animal tracking. *Environmental Modelling & Software*, 26(6), 834–835. <https://doi.org/10.1016/j.envsoft.2010.12.005>
- Lebreton, J., Devillard, S., Popy, S., Desprez, M., Besnard, A., & Gaillard, J. M. (2010). Towards a vertebrate demographic data bank. *Journal of Ornithology*, 152(S2), 617–624. <https://doi.org/10.1007/s10336-010-0582-0>
- Li, J., Sun, L., & Kappeler, P. M. (2020). *The behavioral ecology of the Tibetan macaque*. <https://doi.org/10.1007/978-3-030-27920-2>
- Lukas, D., & Clutton-Brock, T. (2017). Comparative studies need to rely both on sound natural history data and on excellent statistical analysis. *Royal Society Open Science*, 4(11), 171211. <https://doi.org/10.1098/rsos.171211>
- Lukas, D., & Clutton-Brock, T. (2018). Social complexity and kinship in animal societies. *Ecology Letters*, 21(8), 1129–1134. <https://doi.org/10.1111/ele.13079>
- ManyPrimates, Altschul, D. M., Beran, M. J., Bohn, M., Call, J., DeTroy, S., Duguid, S. J., Egelkamp, C. L., Fichtel, C., Fischer, J., Flessert, M., Hanus, D., Haun, D. B. M., Haux, L. M., Hernandez-Aguilar, R. A., Herrmann, E., Hopper, L. M., Joly, M., Kano, F., ... Watzek, J. (2019). Establishing an infrastructure for collaboration in primate cognition research. *PLoS One*, 14(10), e0223675. <https://doi.org/10.1371/journal.pone.0223675>
- McFarland, R., & Majolo, B. (2013). Coping with the cold: Predictors of survival in wild Barbary macaques, *Macaca sylvanus*. *Biology Letters*, 9(4), 20130428. <https://doi.org/10.1098/rsbl.2013.0428>
- Micheletta, J., Waller, B. M., Panggur, M. R., Neumann, C., Dubosq, J., Agil, M., & Engelhardt, A. (2012). Social bonds affect anti-predator behaviour in a tolerant species of macaque, *Macaca nigra*. *Proceedings of the Royal Society B*, 279(1744), 4042–4050. <https://doi.org/10.1098/rspb.2012.1470>
- Mocha, Y. B., Woith, M., Drobnik, S. M., Markman, S., Frisoni, F., Baglione, V., Boersma, J., Cousseau, L., Covas, R., Braga de Miranda, G. H., Dey, C. J., Doutrelant, C., Gula, R., Heinsohn, R., Kingma, S. A., Li, J., Middleton, K., Radford, A. N., Restrepo, C., ... Griesser, M. (2024). Advancing cooperative breeding research with a peer-reviewed and "live" Cooperative-Breeding Database (Co-Breed). *bioRxiv*. <https://doi.org/10.1101/2024.04.26.591342>
- Morrow, K. S., Glanz, H., Ngakan, P. O., & Riley, E. P. (2019). Interactions with humans are jointly influenced by life history stage and social network factors and reduce group cohesion in moor macaques (*Macaca maura*). *Scientific Reports*, 9(1), 20162. <https://doi.org/10.1038/s41598-019-56288-z>
- Nakagawa, N., Nakamichi, M., & Sugiura, H. (2010). *The Japanese macaques*. Springer Science & Business Media.
- Neumann, C., & Fischer, J. (2023). Extending Bayesian Elo-rating to quantify the steepness of dominance hierarchies. *Methods in Ecology and Evolution*, 14(2), 669–682. <https://doi.org/10.1111/2041-210x.14021>
- Nunn, C. L. (2011). *The comparative approach in evolutionary anthropology and biology*. University of Chicago Press.
- O'Dea, R. E., Parker, T. H., Chee, Y. E., Culina, A., Drobnik, S. M., Duncan, D. H., Fidler, F., Gould, E., Ihle, M., Kelly, C. D., Lagisz, M., Roche, D. G., Sanchez-Tojar, A., Wilkinson, D. P., Wintle, B. C., & Nakagawa, S. (2021). Towards open, reliable, and transparent ecology and evolutionary biology. *BMC Biology*, 19(1), 68. <https://doi.org/10.1186/s12915-021-01006-3>
- Okamoto, K., Matsumura, S., & Watanabe, K. (2000). Life history and demography of wild moor macaques (*Macaca maurus*): Summary of ten years of observations. *American Journal of Primatology*, 52(1), 1–11. [https://doi.org/10.1002/1098-2345\(200009\)52:1<1::AID-AJP1>3.0.CO;2-F](https://doi.org/10.1002/1098-2345(200009)52:1<1::AID-AJP1>3.0.CO;2-F)
- Ostner, J., & Schülke, O. (2018). Linking sociality to fitness in primates: A call for mechanisms. In M. Naguib, L. Barrett, S. D. Healy, J. Podos, L. W. Simmons, & M. Zuk (Eds.), *Advances in the study of behavior* (Vol. 50, pp. 127–175). Academic Press. <https://doi.org/10.1016/bs.asb.2017.12.001>
- Pinter-Wollman, N., Hobson, E. A., Smith, J. E., Edelman, A. J., Shizuka, D., de Silva, S., Waters, J. S., Prager, S. D., Sasaki, T., Wittmyer, G., Fewell, J., & McDonald, D. B. (2013). The dynamics of animal social networks: Analytical, conceptual, and theoretical advances. *Behavioral Ecology*, 25(2), 242–255. <https://doi.org/10.1093/beheco/art047>
- Poisot, T., Bruneau, A., Gonzalez, A., Gravel, D., & Peres-Neto, P. (2019). Ecological data should not be so hard to find and reuse. *Trends in Ecology & Evolution*, 34(6), 494–496. <https://doi.org/10.1016/j.tree.2019.04.005>
- Purgar, M., Klanjscek, T., & Culina, A. (2022). Quantifying research waste in ecology. *Nature Ecology & Evolution*, 6(9), 1390–1397. <https://doi.org/10.1038/s41559-022-01820-0>
- Radhakrishna, S., Huffman, M. A., & Sinha, A. (2012). *The macaque connection: Cooperation and conflict between humans and macaques* (Vol. 43). Springer Science & Business Media.
- Riley, E. P., Sagnotti, C., Carosi, M., & Putu Oka, N. (2014). Socially tolerant relationships among wild male moor macaques (*Macaca*

- maura). *Behaviour*, 151(7), 1021–1044. <https://doi.org/10.1163/1568539x-00003182>
- Roos, C., Kothe, M., Alba, D. M., Delson, E., & Zinner, D. (2019). The radiation of macaques out of Africa: Evidence from mitogenome divergence times and the fossil record. *Journal of Human Evolution*, 133, 114–132. <https://doi.org/10.1016/j.jhevol.2019.05.017>
- Rubenstein, D. R., & Abbot, P. (2017). *Comparative social evolution*. Cambridge University Press. <https://doi.org/10.1017/9781107338319>
- Ruppert, N., Holzner, A., See, K. W., Gisbrecht, A., & Beck, A. (2018). Activity budgets and habitat use of wild southern pig-tailed macaques (*Macaca nemestrina*) in oil palm plantation and forest. *International Journal of Primatology*, 39(2), 237–251. <https://doi.org/10.1007/s10764-018-0032-z>
- Sadoughi, B., Mundry, R., Schulke, O., & Ostner, J. (2024). Social network shrinking is explained by active and passive effects but not increasing selectivity with age in wild macaques. *Proceedings of the Royal Society B*, 291(2018), 20232736. <https://doi.org/10.1098/rspb.2023.2736>
- Sah, P., Mendez, J. D., & Bansal, S. (2019). A multi-species repository of social networks. *Scientific Data*, 6(1), 44. <https://doi.org/10.1038/s41597-019-0056-z>
- Salguero-Gomez, R., Jones, O. R., Archer, C. R., Bein, C., de Buhr, H., Farack, C., Gottschalk, F., Hartmann, A., Henning, A., Hoppe, G., Romer, G., Ruoff, T., Sommer, V., Wille, J., Voigt, J., Zeh, S., Viereg, D., Buckley, Y. M., Che-Castaldo, J., ... Vaupel, J. W. (2016). COMADRE: A global data base of animal demography. *Journal of Animal Ecology*, 85(2), 371–384. <https://doi.org/10.1111/1365-2656.12482>
- Sandel, A. A., Miller, J. A., Mitani, J. C., Nunn, C. L., Patterson, S. K., & Garamszegi, L. Z. (2016). Assessing sources of error in comparative analyses of primate behavior: Intraspecific variation in group size and the social brain hypothesis. *Journal of Human Evolution*, 94, 126–133. <https://doi.org/10.1016/j.jhevol.2016.03.007>
- Schneider, F. D., Fichtmueller, D., Gossner, M. M., Güntsch, A., Jochum, M., König-Ries, B., Le Provost, G., Manning, P., Ostrowski, A., Penone, C., Simons, N. K., & Orme, D. (2019). Towards an ecological trait-data standard. *Methods in Ecology and Evolution*, 10(12), 2006–2019. <https://doi.org/10.1111/2041-210x.13288>
- Schradin, C. (2013). Intraspecific variation in social organization by genetic variation, developmental plasticity, social flexibility or entirely extrinsic factors. *Philosophical Transactions of the Royal Society B*, 368(1618), 20120346. <https://doi.org/10.1098/rstb.2012.0346>
- Schradin, C. (2017). Comparative studies need to rely both on sound natural history data and on excellent statistical analysis. *Royal Society Open Science*, 4(9), 170346. <https://doi.org/10.1098/rsos.170346>
- Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance reproductive success in male macaques. *Current Biology*, 20(24), 2207–2210. <https://doi.org/10.1016/j.cub.2010.10.058>
- Schülke, O., & Ostner, J. (2008). Male reproductive skew, paternal relatedness, and female social relationships. *American Journal of Primatology*, 70(7), 1–4. <https://doi.org/10.1002/ajp.20546>
- Sheldon, B. C., Kruuk, L. E. B., & Alberts, S. C. (2022). The expanding value of long-term studies of individuals in the wild. *Nature Ecology & Evolution*, 6, 1799–1801. <https://doi.org/10.1038/s41559-022-01940-7>
- Simons, N. D., Michopoulos, V., Wilson, M., Barreiro, L. B., & Tung, J. (2022). Agonism and grooming behaviour explain social status effects on physiology and gene regulation in rhesus macaques. *Philosophical Transactions of the Royal Society B*, 377(1845), 20210132. <https://doi.org/10.1098/rstb.2021.0132>
- Singh, M., Jeyaraj, T., Prashanth, U., & Kaumanns, W. (2010). Male–male relationships in lion-tailed macaques (*Macaca silenus*) and bonnet macaques (*Macaca radiata*). *International Journal of Primatology*, 32(1), 167–176. <https://doi.org/10.1007/s10764-010-9448-9>
- Sinha, A. (2005). Not in their genes: Phenotypic flexibility, behavioural traditions and cultural evolution in wild bonnet macaques. *Journal of Biosciences*, 30(1), 51–64. <https://doi.org/10.1007/BF02705150>
- Siracusa, E. R., Negron-Del Valle, J. E., Phillips, D., Platt, M. L., Higham, J. P., Snyder-Mackler, N., & Brent, L. J. N. (2022). Within-individual changes reveal increasing social selectivity with age in rhesus macaques. *Proceedings of the National Academy of Sciences of the United States of America*, 119(49), e2209180119. <https://doi.org/10.1073/pnas.2209180119>
- Strauss, E. D., DeCasien, A. R., Galindo, G., Hobson, E. A., Shizuka, D., & Curley, J. P. (2022). DomArchive: A century of published dominance data. *Philosophical Transactions of the Royal Society B*, 377(1845), 20200436. <https://doi.org/10.1098/rstb.2020.0436>
- Strier, K. B., Altmann, J., Brockman, D. K., Bronikowski, A. M., Cords, M., Fedigan, L. M., Lapp, H., Liu, X., Morris, W. F., Pusey, A. E., Stoinski, T. S., & Alberts, S. C. (2010). The Primate Life History Database: A unique shared ecological data resource. *Methods in Ecology and Evolution*, 1(2), 199–211. <https://doi.org/10.1111/j.2041-210X.2010.00023.x>
- Sueur, C., Petit, O., De Marco, A., Jacobs, A. T., Watanabe, K., & Thierry, B. (2011). A comparative network analysis of social style in macaques. *Animal Behaviour*, 82(4), 845–852. <https://doi.org/10.1016/j.anbehav.2011.07.020>
- Testard, C., Larson, S. M., Watowich, M. M., Kaplinsky, C. H., Bernau, A., Faulder, M., Marshall, H. H., Lehmann, J., Ruiz-Lambides, A., Higham, J. P., Montague, M. J., Snyder-Mackler, N., Platt, M. L., & Brent, L. J. N. (2021). Rhesus macaques build new social connections after a natural disaster. *Current Biology*, 31, 1–11. <https://doi.org/10.1016/j.cub.2021.03.029>
- Thierry, B. (2021). Where do we stand with the covariation framework in primate societies? *American Journal of Biological Anthropology*, 178, 5–25. <https://doi.org/10.1002/ajpa.24441>
- Thierry, B., Iwaniuk, A. N., & Pellis, S. M. (2000). The influence of phylogeny on the social behaviour of macaques (Primates: Cercopithecidae, genus *Macaca*). *Ethology*, 106(8), 713–728. <https://doi.org/10.1046/j.1439-0310.2000.00583.x>
- Thierry, B. (2007). Unity in diversity: Lessons from macaque societies. *Evolutionary Anthropology: Issues, News, and Reviews*, 16(6), 224–238. <https://doi.org/10.1002/evan.20147>
- Tinbergen, N. (1963). On aims and methods of ethology. *Zeitschrift für Tierpsychologie*, 20(4), 410–433.
- Urbano, F., & Cagnacci, F. (2021). Data management and sharing for collaborative science: Lessons learnt from the Euromammals initiative. *Frontiers in Ecology and Evolution*, 9, 727023. <https://doi.org/10.3389/fevo.2021.727023>
- Van Noordwijk, M. A., & Van Schaik, C. P. (1985). Male migration and rank acquisition in wild long-tailed macaques (*Macaca fascicularis*). *Animal Behaviour*, 33(3), 849–861. [https://doi.org/10.1016/S0003-3472\(85\)80019-1](https://doi.org/10.1016/S0003-3472(85)80019-1)
- Visser, I., Bergmann, C., Byers-Heinlein, K., Dal Ben, R., Duch, W., Forbes, S., Franchin, L., Frank, M. C., Geraci, A., Hamlin, J. K., Kaldy, Z., Kulke, L., Laverty, C., Lew-Williams, C., Mateu, V., Mayor, J., Moreau, D., Nomikou, I., Schuwerk, T., ... Zettersten, M. (2022). Improving the generalizability of infant psychological research: The ManyBabies model. *Behavioral and Brain Sciences*, 45, e35. <https://doi.org/10.1017/S0140525X21000455>
- Wallis, J. C., Rolando, E., & Borgman, C. L. (2013). If we share data, will anyone use them? Data sharing and reuse in the long tail of science and technology. *PLoS One*, 8(7), e67332. <https://doi.org/10.1371/journal.pone.0067332>
- Widdig, A., Kessler, M. J., Bercovitch, F. B., Berard, J. D., Duggleby, C., Nurnberg, P., Rawlins, R. G., Saueremann, U., Wang, Q., Krawczak, M., & Schmidtke, J. (2016). Genetic studies on the Cayo Santiago rhesus macaques: A review of 40 years of research. *American Journal of Primatology*, 78(1), 44–62. <https://doi.org/10.1002/ajp.22424>
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J. W., da Silva Santos, L. B.,

- Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., ... Mons, B. (2016). The FAIR guiding principles for scientific data management and stewardship. *Scientific Data*, 3, 160018. <https://doi.org/10.1038/sdata.2016.18>
- Yamada, M. (1963). A study of blood-relationship in the natural society of the Japanese macaque—An analysis of co-feeding, grooming, and playmate relationships in Minoo-B-troop. *Primates*, 4(3), 43–65. <https://doi.org/10.1007/bf01733671>
- Young, C., Majolo, B., Heistermann, M., Schülke, O., & Ostner, J. (2014). Responses to social and environmental stress are attenuated by strong male bonds in wild macaques. *Proceedings of the National Academy of Sciences of the United States of America*, 111(51), 18195–18200. <https://doi.org/10.1073/pnas.1411450111>

How to cite this article: De Moor, D., Skelton, M.; Amici, F., Arlet, M. E., Balasubramaniam, K. N., Ballesta, S., Berghänel, A., Berman, C. M., Bernstein, S. K., Bhattacharjee, D., Bliss-Moreau, E., Brotcorne, F., Butovskaya, M., Campbell, L. A. D., Carosi, M., Chatterjee, M., Cooper, M. A., Cowl, V. B., De la O, C., ... Brent, L. J. N. (2025). MacaqueNet: Advancing comparative behavioural research through large-scale collaboration. *Journal of Animal Ecology*, 00, 1–16. <https://doi.org/10.1111/1365-2656.14223>