

LJMU Research Online

Johnson, CL, Krasner, S, Woodward, W, Mao, E, McCourt, A, Dockerill, C, Balkhausen, K, Chandrasekaran, B, Kardos, A, Sabharwal, N, Firoozan, S, Sarwar, R, Senior, R, Sharma, R, Wong, K, Augustine, DX, Paton, M, O'Driscoll, J, Oxborough, D, Pearce, K, Robinson, S, Willis, J, Leeson, P and EVAREST/BSE-NSTEP Investigators

Changes in Coronary Disease Management Decisions in Real World Practice between 2015 and 2023: Insights from the Evarest/BSE-NSTEP observational study

<https://researchonline.ljmu.ac.uk/id/eprint/26072/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)










Johnson, CL, Krasner, S, Woodward, W, Mao, E, McCourt, A, Dockerill, C, Balkhausen, K, Chandrasekaran, B, Kardos, A, Sabharwal, N, Firoozan, S, Sarwar, R, Senior, R, Sharma, R, Wong, K, Augustine, DX, Paton, M, O'Driscoll, J. Oxborough, D. Pearce, K. Robinson, S. Willis, J. Leeson, P and

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

Changes in coronary disease management decisions in real-world practice between 2015 and 2023: insights from the EVAREST/BSE-NSTEP observational study

Casey L. Johnson ¹, Samuel Krasner ¹, William Woodward ^{1,2}, Emma Mao¹, Annabelle McCourt¹, Cameron Dockerill^{1,3}, Katrin Balkhausen⁴, Badrinathan Chandrasekaran ⁵, Attila Kardos ^{6,7}, Nikant Sabharwal ⁸, Soroosh Firoozan⁹, Rizwan Sarwar⁸, Roxy Senior ¹⁰, Rajan Sharma¹¹, Kenneth Wong¹², Daniel X. Augustine¹³, Maria Paton¹⁴, Jamie O'Driscoll^{11,15}, David Oxborough ¹⁶, Keith Pearce¹⁷, Shaun Robinson¹⁸, James Willis¹³, and Paul Leeson ^{1*}; On behalf of the EVAREST/BSE-NSTEP Investigators

¹Cardiovascular Clinical Research Facility, Division of Cardiovascular Medicine, University of Oxford, Oxford, UK; ²Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK; ³Kings College London, London, UK; ⁴Royal Berkshire Hospital, Reading, UK; ⁵Great Western Hospital, Swindon, UK; ⁶Department of Cardiology, Translational Cardiovascular Research Group, Milton Keynes University Hospital NHS Foundation Trust, Milton Keynes, UK; ⁷Faculty of Medicine and Health Sciences, University of Buckingham, Buckingham, UK; ⁸Oxford University Hospitals NHS Foundation Trust, Oxford, UK; ⁹Wycombe Hospital, High Wycombe, UK; ¹⁰Northwick Park Hospital, London, UK; ¹¹St George's University Hospitals NHS Foundation Trust, Tooting, UK; ¹²Lancashire Cardiac Centre, Blackpool Teaching Hospitals NHS Foundation Trust, Blackpool, UK; ¹³Royal United Hospitals, Bath, UK; ¹⁴Leeds Institute of Cardiovascular and Metabolic Medicine, University of Leeds, Leeds, UK; ¹⁵Diabetes Research Centre, College of Life Sciences, University of Leicester, Leicester, UK; ¹⁶Research Institute for Sport and Exercise Sciences and the Liverpool Centre for Cardiovascular Science, Liverpool John Moores University, Liverpool, UK; ¹⁷Manchester University NHS Foundation Trust, Manchester, UK; and ¹⁸Imperial Healthcare NHS Trust, London, UK

Received 24 February 2025; accepted 5 March 2025; online publish-ahead-of-print 22 March 2025

Aims

To assess the real-world impact of updated clinical guidelines and literature on the management of patients undergoing stress echocardiography for the assessment of inducible ischaemia across a national health service.

Methods and results

A total of 13 819 patients from 32 UK hospitals, referred for stress echocardiography between 2015 and 2023, were analysed across two phases: phase 1 (2015–2020) and phase 2 (2020–2023). Follow-up data for 1 year was available for 4920 participants through NHS Digital. Patients in phase 2 were younger and presented with a higher cardiovascular risk profile, although sex distribution remained similar across phases. There was an observed reduction in invasive angiography referrals within 1 year following a positive stress echocardiogram ($P < 0.01$), which appeared to be attributed to changes in the management of patients with moderate ischaemia (3–4 segments; $P < 0.01$). For those who did receive invasive assessment, there were no changes in intervention rate ($P = 0.27$), regardless of ischaemic burden. This trend was most evident in centres performing a higher volume of stress echocardiograms.

Conclusion


Coronary disease management pathways have changed within the UK and fewer patients with moderate ischaemia are undergoing invasive coronary angiography. However, coronary intervention rates are unchanged, suggesting that stress echocardiography is being used to improve patient selection for invasive procedures while minimizing unnecessary referrals. Future work will assess if this reduction in angiography referrals is maintained long term, and if there are any effects on patient outcomes.

* Corresponding author. E-mail: paul.leeson@cardiov.ox.ac.uk

© The Author(s) 2025. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Graphical Abstract



Keywords

echocardiography • coronary artery disease • stress echocardiography • invasive coronary angiography

Introduction

For many years, referral for invasive coronary angiography was common for patients with evidence of more than mild cardiac ischaemia on functional imaging to ensure that patients received the opportunity to have revascularization. However, contradicting evidence from randomized studies has led to recent debate over the appropriate investigation and treatment steps in the care pathway for coronary disease.

Several studies have indicated that the risk of death, myocardial infarction, or other cardiac events may not be reduced by an initial invasive strategy in all patients in the non-acute setting,^{1–5} and the ORBITA trial found limited symptomatic benefit of an invasive approach, with no improvement in exercise capacity.^{6–9} However, the ORBITA-2 trial demonstrated improved patient-reported symptom scores for those who received coronary intervention,¹⁰ and outcome benefits were also evident in long-term follow-up of the ISCHEMIA study¹¹ consistent with a recent meta-analysis.¹²

To take account of this emerging data, guidelines have evolved to focus on better selection of high-risk patients for revascularization, preferring an initial medical management strategy in those with lower ischaemic burden. Referral for invasive angiography is reserved for when guideline-directed medical therapy fails to relieve symptoms.^{13–15} Prior to these recent updates, referral of patients for invasive coronary angiography after diagnosis of coronary artery disease was entrained in medical practice, and patients understood this as 'the best' way to manage their disease.^{16–18} Whether patients and medical staff would adopt recent guideline updates therefore remained unclear.

Stress echocardiography has been used as a first-line test for patients with chest pain for decades.^{14–23} The Echocardiography Value and Accuracy at Rest and Stress (EVAREST) studied the care pathway of up to 18 000 patients undergoing stress echocardiography across 32 UK hospitals between 2015 and 2023. Recruitment spanned the time that key randomized trials and updated guidelines on cardiovascular disease management were published. As such, the study is uniquely placed to provide insight into whether there have been temporal changes in

management decisions for patients presenting with coronary disease in the UK.


Methods

Study design

The EVAREST study is a multi-centre observational study evaluating the use, accuracy, and performance of stress echocardiography in real-world practice. From 2020 to 2023, the study incorporated the British Society of Echocardiography National Review of Stress Echocardiography Practice (BSE-NSTEP). The full methodology and 6-month outcome results from the first phase of the study have been previously reported.²⁴ The study is registered at ClinicalTrials.gov (NCT03674255), and ethical approval was provided by the Health Research Authority South Central Berkshire Research Ethics Committee (Ethics Reference: 14/SC/1437). The study design predates the common widespread use of patient and public involvement, but patient and public feedback was received via the National Institute for Health and Care Research survey of research participants following the conclusion of the study. Written informed consent was obtained from all patients and the study was conducted in accordance with the Declaration of Helsinki principles.

Participants

Participants were recruited in two phases with the first recruitment phase running from March 2015 to March 2020 and included patients who were referred for stress echocardiography to assess inducible ischaemia. Recruitment restarted after the initial period of the COVID pandemic and then ran from October 2020 to September 2023, but was expanded to include patients referred to stress echocardiography for any clinical reason. However, only patients referred to stress echocardiography to assess inducible ischaemia were included in this analysis. As there was variation in participating centres between the two phases, only participants recruited at 26 sites that were recruiting centres during both phases of the study were



included in this analysis. A subgroup of participants provided consent to link their details with follow-up outcome data provided by NHS England.

Data collection

Participant demographics and stress echocardiogram procedure details were collected by the local study teams and entered into an electronic database (Castor EDC, Amsterdam, Netherlands). The annual stress echocardiography volume was self-reported by each hospital. Hospital capacity as measured by the number of beds was retrieved from NHS England.²⁵ This work uses data provided by patients and collected by the NHS as part of their care and support via the NHS Digital Data Access Request Service. Hospital admission data was collected from the Hospital Episode Statistics Admitted Patient Care database. Data collected included the date and reason for admission, and any procedures undertaken during admission such as invasive coronary angiography, percutaneous coronary intervention, and coronary artery bypass grafting. Reasons for admission were defined by the International Classification of Disease-10th revision coding (ICD-10), and interventions and/or procedures were defined by the OPCS Classification of Interventions and Procedures-4th revision coding (OPCS 4.10) which is the procedural classification used within the NHS in the UK. Data on subsequent diagnostic imaging including invasive coronary angiography was obtained from the

Diagnostic Imaging Dataset held by NHS England. Imaging data submitted to NHS England are coded using the Systematized Nomenclature of Medicine-Clinical Terms (SNOMED-CT). Mortality data including date and cause of death were obtained from the Civil Registrations of Death database provided by NHS England. Details of codes used in this analysis are provided in the [Supplementary material](#).

Statistical analysis

Patient demographics and stress echocardiogram procedural details are reported using standard approaches. Variations in hospital size, measured via annual stress echocardiography volume and hospital bed capacity, were separated into quartiles for comparison. Descriptive statistics were investigated as frequencies and medians [interquartile range (IQR)]. A comparison of discrete data between recruitment phases was conducted using Pearson's χ^2 tests. Stress echocardiogram result (positive or negative) was reported as the result by the clinician responsible for each participant's care. Kaplan–Meier time-to-event curves and log-rank tests were used to assess differences in rates of invasive coronary angiography and percutaneous coronary intervention. Additionally, differences in invasive angiography referrals between recruitment phases were examined according to ischaemic burden (mild: 1–2 ischaemic segments, moderate: 3–4 ischaemic segments, or severe: ≥ 5 ischaemic segments). Participants with missing data


	Phase 1 total cohort (n = 7332)	Phase 2 total cohort (n = 6487)	P-value*	Phase 1 NHS Digital subgroup (n = 2451)	Phase 2 NHS Digital subgroup (n = 2469)	P-value**
Participant demographics						
Male (%)	4126/7332 (56.3)	3739/6486 (57.6)	0.10	1440/2451 (58.8)	1429/2469 (57.9)	0.53
Median age (years) (IQR)	66 (57–73)	65 (57–74)	0.20	67 (59–74)	65 (56–73)	<0.001
Median BMI (kg/m ²) (IQR)	28.2 (25.0–31.9)	28.1 (25–31.6)	0.87	27.9 (24.9–31.5)	28.1 (24.9–31.9)	<0.05
Positivity rate (%)	1394/7332 (19.0)	1231/6487 (19.0)	0.96	448/2451 (18.3)	523/2469 (21.2)	<0.05
Current smoker (%)	860/7045 (12.2)	744/6285 (11.8)	0.63	203/2541 (8.3)	374/2469 (15.1)	<0.001
Ex-smoker (%)	2687/7045 (38.1)	2125/6285 (33.8)	<0.001	1025/2451 (41.8)	786/2469 (31.8)	<0.001
Non-smoker (%)	3498/7045 (49.7)	3416/6285 (54.4)	<0.001	1144/2451 (46.7)	1242/2469 (50.3)	<0.05
Hypertension (%)	3346/6931 (48.3)	3445/6455 (53.4)	<0.001	1332/2295 (58.0)	1301/2466 (52.8)	<0.001
Hypercholesterolaemia (%)	2767/6931 (39.9)	3090/6455 (47.9)	<0.001	1064/2295 (46.4)	1187/2466 (48.1)	0.22
Diabetes mellitus (%)	1331/7332 (18.2)	1463/6455 (22.7)	<0.001	506/2451 (20.6)	525/2466 (21.3)	0.57
Peripheral vascular disease (%)	204/6931 (2.9)	99/6453 (1.5)	<0.001	74/2295 (3.2)	40/2467 (1.6)	<0.001
Previous MI (%)	1243/7214 (17.2)	1121/6453 (17.4)	0.83	445/2423 (18.4)	472/2467 (19.1)	0.49
Previous PCI (%)	2256/7228 (31.2)	1323/6453 (20.5)	<0.001	835/2425 (34.4)	548/2467 (22.2)	<0.001
Precious CABG (%)	529/7241 (7.3)	441/6453 (6.8)	0.28	178/2429 (7.3)	156/2467 (6.3)	0.16
Resting RVWA (%)	1080/7328 (14.7)	916/6449 (14.2)	0.37	397/2449 (16.2)	313/2467 (12.7)	<0.001
Stress echocardiogram details						
Exercise (%)	2211/7330 (30.2)	2508/6480 (38.7)	<0.001	778/2451 (31.7)	982/2469 (39.8)	<0.001
Pacemaker (%)	11/7332 (0.2)	39/6480 (0.6)	<0.001	7/2451 (0.3)	18/2469 (0.7)	<0.05
Dobutamine (%)	5108/7330 (69.7)	3919/6480 (60.5)	<0.001	1665/2451 (67.9)	1464/2469 (59.3)	<0.001
Atropine use in DSE (%)	2518/5091 (49.5)	1773/3859 (45.9)	<0.001	734/1665 (44.1)	689/1464 (47.1)	0.10
Contrast used	5258/7309 (71.9)	5279/6407 (82.4)	<0.001	1961/2440 (80.4)	1907/2444 (78.0)	<0.05
SonoVue (%)	4847/7309 (66.3)	4193/6407 (65.4)	0.07	1729/2440 (70.9)	1668/2444 (68.2)	<0.05
Luminity (%)	393/7309 (5.4)	1080/6407 (16.9)	<0.001	220/2440 (9.0)	236/2444 (9.7)	0.44
Other (e.g. Optison) (%)	18/7309 (0.2)	6/6407 (0.1)	0.031	12/2440 (0.5)	3/2444 (0.1)	<0.05

***P-value comparison between recruitment phases (NHS Digital subgroup).

phase 2). Follow-up data were received from NHS Digital for a subgroup of 4920 participants (2451 in phase 1 and 2469 in phase 2). Participant inclusion and exclusion for this analysis are described in *Figure 1*.

Full patient demographics are provided in *Table 1*. Median age was similar between phase 1 and phase 2, 66 (IQR 57–73) and 65 (IQR 57–74) years, respectively. Patient sex was also consistent across both phases (56.3 vs. 57.6%), but patients recruited in phase 2 had a higher incidence of hypertension (48.3 vs. 53.4%), hypercholesterolaemia (39.9 vs. 47.9%) and diabetes (18.2 vs. 22.7%) (all $P < 0.001$). The percentage of patients reporting that current smoking practices remained consistent, but there was an increase in ex-smokers (49.7 vs. 54.4%) ($P < 0.001$). There was a decrease in peripheral vascular disease in phase 2 (2.9 vs. 1.5%) as well as a decrease in previous PCI (31.2 vs. 20.5%) (both $P < 0.001$). There were no significant changes in the percentage of previous MI, CABG, and the presence of resting regional wall motion abnormalities between phases.

Between March 2015 and September 2023, 17 656 patients were recruited into the EVAREST study of which data for 13 819 participants were available for the temporal analysis (7332 in phase 1 and 6487 in

**Table 2** Multivariate Cox regression analysis of invasive coronary angiography in NHS Digital subgroup

Variable	Total subgroup			Positive stress echocardiogram		
	HR	95% CI	P-value	HR	95% CI	P-value
Recruitment phase (phase 2)	0.77	0.65–0.91	<0.01	0.75	0.62–0.92	<0.01
Age	0.99	0.99–1.00	0.08	0.99	0.98–1.00	<0.05
Sex (male)	1.16	0.97–1.38	0.10	1.16	0.94–1.42	0.17
SE outcome (positive)	17.38	14.29–21.15	<0.001	—	—	—
Current smoker	0.90	0.70–1.16	0.41	0.87	0.65–1.16	0.35
Hypertension	1.19	1.00–1.41	<0.01	1.16	0.95–1.41	0.14
Previous peripheral vascular disease	0.91	0.56–1.48	0.71	1.11	0.67–1.84	0.69
Previous PCI	0.79	0.66–0.96	<0.05	0.73	0.59–0.90	<0.01
Baseline RVMA	1.21	1.00–1.46	<0.05	1.12	0.91–1.37	0.29
Hospital demographics						
SE per year (per quartile)	0.75	0.68–0.81	<0.001	0.71	0.64–0.79	<0.001
Bed number (per quartile)	1.10	1.00–1.21	<0.05	1.12	1.00–1.25	0.06

Bold values indicate statistically significant.

Stress echocardiogram positivity was similar between phases at 19.0%. While dobutamine remained the most common stressor used between phases, there was a decrease from 69.7% in phase 1 to 60.5% in phase 2 ($P < 0.001$). This saw a corresponding increase in the use of exercise stress from 30.2 to 38.7% between phases ($P < 0.001$). There was

also a marginal increase in the use of pacemaker stress (0.2 vs. 0.6%, $P < 0.001$). Within dobutamine stress echocardiograms, there was a decrease in the use of atropine (49.5 vs. 45.9%, $P < 0.001$). The use of contrast increased (71.9 vs. 82.4%, $P < 0.001$) with a related increase in the use of Luminy as a contrast agent (5.4 vs. 16.9%, $P < 0.001$).

	Total angiograms (%)*	Angiogram phase 1 (%)*	Angiogram phase 2 (%)*
Stress echocardiography volume			
<400	137 (17.3)	68 (14.4)	69 (21.5)
400–599	116 (10.5)	72 (12.4)	44 (8.4)
600–850	102 (13.0)	44 (13.5)	58 (12.6)
>850	264 (11.8)	145 (13.5)	119 (10.2)
Hospital capacity (number of beds)			
<600	194 (14.9)	104 (14.3)	90 (15.8)
600–799	84 (12.0)	74 (14.5)	10 (5.3)
800–1000	155 (10.4)	35 (10.0)	120 (10.6)
>1000	186 (12.9)	116 (13.5)	70 (12.2)

There are limitations to this analysis. Firstly, due to the nature of the data collection, there are no results on patient symptoms throughout the management period. Some studies have shown that patient-reported symptoms and quality of life are improved with an invasive

management strategy even when a reduction in mortality and adverse events is not identified.^{5,10} However, this improvement in symptoms appears inconsistent⁶ and some investigators have attributed this to a placebo effect.³¹ Secondly, the time horizon used in this analysis is limited to 1 year. Most patients referred for an elective coronary angiogram following their stress echocardiogram will be seen within this timeframe and this appears appropriate to account for further investigational testing such as invasive angiography. However, any invasive angiography performed more than 1 year following uncontrolled symptoms with an initial medical management strategy may be unaccounted for. Thirdly, the study was focused on an evaluation of real-world practice and it is possible associations may differ if other stress echocardiography protocols were applied in practice that used additional measures that may improve predictive accuracy such as heart rate reserve. Fourthly, it should be noted that data received from the data request service from NHS Digital has inherent limitations. If no outcome data was received after supplying NHS Digital with participant identifiers for data linkage, it was assumed that this participant had no follow-up outcomes or events within the requested timeframe. This could, however, also mean that the participant had follow-up data, but was not able to be retrieved by NHS Digital for unknown reasons. Fifthly, while sites remained consistent for the temporal analysis, not all sites began recruiting at the same time and had varying recruitment rates. Therefore, some sites contributed more proportionally to the dataset. Finally, due to the nature of the prospective consented study design, there may be a selection bias among those enrolled towards those with an interest in research participation.

Conclusion

This study provides real-world evidence of a change in coronary disease management decisions within the NHS. Since 2020, there has been a small but significant reduction in the number of patients who are referred for invasive angiography after a positive stress echocardiogram. This can be attributed to a reduced referral to invasive angiography in patients with moderate ischaemia, while those with mild and severe disease have not experienced significant changes in their management pathways. Interestingly, rates of use of percutaneous coronary intervention did not change over the recruitment period, suggesting a better selection of patients for angiography. These results should be considered in the context of the sample size and time horizon, and future work will aim to further confirm these management changes, and establish what effect, if any, this has on patient outcomes long term.

EVAREST/BSE-NSTEP Investigators

Abraheem Abraheem¹⁹, Sanjay Banypersad²⁰, Sadie Bennett²¹, Henry Boardman⁷, Christopher Boos²², Sudantha Bulugahapitiya²³, Jeremy Butts²⁴, Duncan Coles²⁵, Joanna d'Arcy⁸, Jacob Easaw¹³, Sarah Fairbairn²⁶, Patrick Gibson²⁷, Haytham Hamdan²⁸, Shahnaz Jamil-Copley²⁹, Gajen Kanaganayagam³⁰, Guy Lloyd³¹, Ioannis Moukas³², Tom Mwambuingu³³, Thuraia Nageh³⁴, Antonis Pantazis³⁵, Alexandros Papachristidis³⁶, Ronak Rajani³⁷, Muhammad Amer Rasheed³⁸, Naveed A. Razvi³⁹, Sushma Rekhrja²⁹, Joban Sehmi⁴⁰, Azeem Sheikh²², David P. Ripley⁴¹, Kathleen Rose⁴², Michaela Scheuermann-Freestone⁴³, Rebecca Schofield⁴⁴, Ayyaz Sultan²⁸, Spiros Zidros⁴⁵

¹⁹Department of Cardiology, Tameside and Glossop Integrated Care NHS Foundation Trust, Ashton-under-Lyne, UK.

²⁰Department of Cardiology, Royal Blackburn Hospital, East Lancashire Hospitals NHS Trust, Burnley, UK.

²¹Department of Cardiology, Heart & Lung Centre, Royal Stoke University Hospital, University Hospitals of North Midlands NHS Foundation Trust, Stoke, UK.

²²Department of Cardiology, Poole Hospital NHS Foundation Trust, Poole, UK.

²³Department of Cardiology, Bradford Royal Infirmary Hospital, Bradford Teaching Hospitals NHS Foundation Trust, Bradford, UK.

²⁴Department of Cardiology, the Calderdale Royal Hospital and Huddersfield Royal Infirmary, Calderdale and Huddersfield NHS Foundation Trust, Calderdale, UK.

²⁵Department of Cardiology, Broomfield Hospital, Mid and South Essex NHS Foundation Trust, Broomfield, UK.

²⁶Bristol Heart Institute, University Hospitals Bristol and Weston NHS Foundation Trust, Bristol, UK.

²⁷Royal Infirmary Edinburgh and Western General Hospital, NHS Lothian, UK.

²⁸Department of Cardiology, Wrightington, Wigan and Leigh NHS Foundation Trust, Wigan, UK.

²⁹Department of Cardiology, Nottingham University Hospitals NHS Trust, Nottingham, UK.

³⁰Department of Cardiology, West Middlesex University Hospital, Chelsea and Westminster Hospital NHS Foundation Trust, London, UK.

³¹University College London, Barts Health NHS Trust, UK.

³²Department of Cardiology, Warrington and Halton Teaching Hospitals NHS Foundation Trust, Warrington, UK.

³³Department of Cardiology, The Mid Yorkshire Hospitals NHS Trust, Pinderfields, UK.

³⁴Department of Cardiology, Southend University Hospital, Mid and South Essex NHS Foundation Trust, Southend-on-Sea, UK.

³⁵Department of Cardiology, North Middlesex University Hospital NHS Trust, London, UK.

³⁶Department of Cardiology, King's College Hospital NHS Foundation Trust, London, UK.

³⁷Department of Cardiology, Guy's and St Thomas' NHS Foundation Trust, London, UK.

³⁸Department of Cardiology, Yeovil District Hospital, Somerset NHS Foundation Trust, Yeovil, UK.

³⁹Department of Cardiology, East Suffolk and North Essex NHS Foundation Trust, Ipswich, UK.

⁴⁰West Hertfordshire Hospitals NHS Trust, UK.

⁴¹Department of Cardiology, Northumbria Healthcare NHS Foundation Trust, North Tyneside, UK.

⁴²Department of Cardiology, Northampton General Hospital NHS Trust, Northampton, UK.

⁴³Department of Cardiology, Hampshire Hospitals NHS Foundation Trust, Basingstoke, UK.

⁴⁴Department of Cardiology, North West Anglia NHS Foundation Trust, Peterborough, UK.

⁴⁵Department of Cardiology, Bedford Hospital, Bedfordshire Hospitals NHS Foundation Trust, Bedford, UK.

Author contributions

P.L. was responsible for conceptualization of the analysis. P.L., K.B., B.C., A.K., N.S., S.F., R.Sa, R.Se, R.Sh, K.W., D.X.A., M.P., J.O.D., D.O., K.P., S.R., and J.W. developed the overarching research study design. P.L. assisted with funding acquisition. W.W., C.D., C.L.J., A.M., E.M., and S.K. provided project administration and data curation. W.W., C.D., C.L.J., A.M., S.K., D.X.A., J.W., M.P., P.L., R.Se, J.O.D., R.S., and all EVAREST/BSE-NSTEP investigators provided investigation for the study. S.K. supported data analysis and visualization. C.L.J. and P.L. wrote and edited the original manuscript draft. All authors reviewed and edited the manuscript. P.L. accepts the responsibility of guarantor and senior

