



## LJMU Research Online

Tuschick, E, Ferguson, J, Coulton, S, Eberhardt, J, Reneesha, A, Osindeinde, I, Deluca, P, Sumnall, H, Stevens, A, Bray, J and Newbury-Birch, D

**Development of a trial application to assess the effectiveness and cost-effectiveness of adult dRug scrEening and brief interventionS in key hEalth, social care and justice setTings: The RESET PROJECT**

<https://researchonline.ljmu.ac.uk/id/eprint/26502/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Tuschick, E, Ferguson, J, Coulton, S, Eberhardt, J, Reneesha, A, Osindeinde, I, Deluca, P, Sumnall, H, Stevens, A, Bray, J and Newbury-Birch, D (2025) Development of a trial application to assess the effectiveness and cost-effectiveness of adult dRug scrEening and brief**

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)



<http://researchonline.ljmu.ac.uk/>



RESEARCH ARTICLE

# Development of a trial application to assess the effectiveness and cost-effectiveness of adult dRug scrEening and brief interventionS in key hEalth, social care and justice setTings: The RESET PROJECT

[version 1; peer review: 1 approved]

Emma Tuschick<sup>1</sup>, Jennifer Ferguson<sup>1</sup>, Simon Coulton<sup>2</sup>, Judith Eberhardt <sup>1</sup>, Aysha Reneesha<sup>3</sup>, Ifeoluwa Osindeinde<sup>3</sup>, Paolo Deluca<sup>4</sup>, Harry Sumnall<sup>5</sup>, Alex Stevens<sup>6</sup>, Jeremy Bray<sup>7</sup>, Dorothy Newbury-Birch <sup>1</sup>

<sup>1</sup>School of Social Sciences, Humanities & Law, Teesside University, Middlesbrough, TS1 3BX, UK

<sup>2</sup>Centre for Health Services Research, University of Kent, Kent, CT2 7NP, UK

<sup>3</sup>School of Health and Life Sciences, Teesside University, Middlesbrough, TS1 3BX, UK

<sup>4</sup>Institute of Psychiatry, Psychology & Neuroscience, Kings College London, London, WC2R 1AP, UK

<sup>5</sup>School of Psychology, Liverpool John Moores University, Liverpool, L3 5AH, UK

<sup>6</sup>School of Social Policy, Sociology and Social Research, University of Kent, Kent, CT2 7NP, UK

<sup>7</sup>Bryan School of Business and Economics, University of North Carolina at Greensboro, 44 Bryan Building, Greensboro, North Carolina, 26170, USA

**V1** First published: 22 May 2025, 5:49  
<https://doi.org/10.3310/nihropenres.13633.1>  
Latest published: 22 May 2025, 5:49  
<https://doi.org/10.3310/nihropenres.13633.1>

## Abstract

### Background

In England and Wales, alcohol-related crime is estimated to cost society £27.4 billion and drugs £20 billion annually. Effective interventions therefore have the potential to reduce the costs relating to substance use and increase individual social welfare.

Brief drug use interventions are a secondary prevention activity, which are aimed at those individuals who are using substances in a pattern that is likely to be harmful to health and/or well-being. At present there is limited evidence regarding the effects and most of the work has been carried out outside the UK. We examined the evidence to develop a trial to test effectiveness of brief drug interventions.

### Methods

## Open Peer Review

### Approval Status

1	
<hr/>	
version 1	
22 May 2025	<a href="#">view</a>
<hr/>	
1. Emily E. Tanner-Smith  , University of Oregon, Eugene, USA	
Any reports and responses or comments on the article can be found at the end of the article.	

We carried out two systematic reviews of the literature and examined effectiveness, barriers and facilitators, screening tools and active ingredients of interventions. We also carried out qualitative work to examine this issue.

## Results

The quantitative review included 46 papers (mostly from the USA) and the qualitative review included 14. We ascertained that the ASSIST screening tool was the best tool to use for screening. We carried out interviews with practitioners, policy makers and individuals who have had experience of mental health, criminal justice and family services which were identified as the services we should work with for a definitive trial. Key issues community members perceived included the need to address trauma, the efficacy of interventions, how stigma and societal views affect individuals, the importance of support from various institutions, and how community involvement and personal responsibility play a role in the recovery process. Stakeholders emphasised the need for interventions to be brief yet personalised, underlining the importance of trust and effective evaluation.

## Conclusions

We used the findings from the reviews and qualitative work to develop a robust pilot trial application.

## Plain Language Summary

In England and Wales, alcohol-related crime costs society £27.4 billion a year. Drugs cost £20 billion. If we can find interventions that work this will help individuals and could save a lot of money.

We examined evidence around brief drug interventions. We did not include those that are dependent on drugs. At present there is limited evidence regarding the effects. Most of the work has been carried out outside the UK. We looked at what has been done. This was to develop a trial to find out if they do make a difference.

We carried out reviews of literature. One looked at whether brief interventions work for drug use. The other looked at what people thought about them. We then carried out a lot of interviews with people who had been involved in them.

The quantitative review included 46 papers (mostly from the USA). The qualitative review included 14 papers. We found that the ASSIST tool was the best tool to use for screening. We found that mental health, criminal justice and family services were the best sites for the future study.

We carried out interviews with practitioners, policy makers and individuals. Community members want interventions that address

trauma. They wanted to look at how stigma and societal views affect individuals. Finally, they also told us the importance of support from various institutions, and how community involvement and personal responsibility play a role in the recovery process. Stakeholders told us that interventions need to be brief yet personalised. This showed the importance of trust.

We used the findings of our work and have submitted a funding application to NIHR PHR to see if it can make a difference.

### Keywords

Brief interventions; substance use; healthcare settings; criminal justice settings, mental health, mixed methods

**Corresponding author:** Dorothy Newbury-Birch ([d.newbury-birch@tees.ac.uk](mailto:d.newbury-birch@tees.ac.uk))

**Author roles:** **Tuschick E:** Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Ferguson J:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Coulton S:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Eberhardt J:** Conceptualization, Funding Acquisition, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Reneesha A:** Data Curation, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Osindeinde I:** Data Curation, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Deluca P:** Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation; **Sumnall H:** Conceptualization, Funding Acquisition, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Stevens A:** Conceptualization, Funding Acquisition, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Bray J:** Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Newbury-Birch D:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

**Competing interests:** No competing interests were disclosed.

**Grant information:** This project is funded by the National Institute for Health Research (NIHR) under its [‘Research for Patient Benefit (RfPB) Programme’ (Grant Reference Number NIHR156167)- [ Development of a trial application to assess the effectiveness and cost-effectiveness of adult dRug scrEening and brief interventionS in key hEalth, social care and justice setTings: The RESET PROJECT.] The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2025 Tuschick E *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Tuschick E, Ferguson J, Coulton S *et al.* **Development of a trial application to assess the effectiveness and cost-effectiveness of adult dRug scrEening and brief interventionS in key hEalth, social care and justice setTings: The RESET PROJECT [version 1; peer review: 1 approved]** NIHR Open Research 2025, 5:49 <https://doi.org/10.3310/nihropenres.13633.1>

**First published:** 22 May 2025, 5:49 <https://doi.org/10.3310/nihropenres.13633.1>

## Introduction

Between 2010 and 2019 the number of people using drugs increased by 22% globally (United Nations on Drugs and Crime, 2023). In England and Wales around 9% of adults have used an illicit drug with 2% being frequent users (Office for National Statistics, 2023). In the UK general population around 26% of adults are risky drinkers (38% of men, 16% of women) (Drummond *et al.*, 2004) with substance use amongst those involved in the criminal justice system being higher. A recent review found 63% of people in the UK criminal justice system scored positive for risky drinking (Newbury-Birch *et al.*, 2016) and the prevalence of drug use was 81% (Light *et al.*, 2013) with an interlink between both alcohol and drug use. Therefore, it is sensible to include both drugs and alcohol into a single substance use intervention.

In England and Wales, alcohol-related crime is estimated to cost society £27.4 billion (Institute of Alcohol Studies, 2024) and drugs £20 billion annually (Black, 2020). Effective interventions therefore have the potential to significantly reduce the costs relating to substance use as well as increase individual social welfare (Raistrick *et al.*, 2006).

Brief substance use interventions serve as a secondary prevention measure, targeting individuals whose substance use patterns are likely detrimental to their health and well-being. These interventions have demonstrated frequent effectiveness in primary healthcare settings (Kaner *et al.*, 2018; O'Donnell *et al.*, 2014) and have shown some success in hospital environments (Gaume *et al.*, 2021; McQueen *et al.*, 2009). Typically, they are administered by non-specialist practitioners to non-treatment, opportunistic populations (Miller & Rollnick, 1991). Additionally, there is some evidence supporting their efficacy in reducing recidivism within the criminal justice system (Newbury-Birch *et al.*, 2014). Although there is limited evidence regarding the effects of drug-targeted brief interventions on drug use (Halladay *et al.*, 2019; Humeniuk *et al.*, 2018; Tanner-Smith *et al.*, 2022), most of the work has been carried out outside the UK.

It is crucial to implement strategies that utilise a co-production method, involving both practitioners and, where possible, service users, to ensure the findings are effectively implemented (Newbury-Birch *et al.*, 2016; Sherman *et al.*, 2015). While it is often argued that academics and criminal justice practitioners come from vastly different backgrounds, the gap between them may not be as wide as perceived (Wehrens, 2014). A co-production approach, where researchers, practitioners, and community members collaborate, could facilitate genuine translational research (Graham & Tetroe, 2007). This concept is aptly summarised by Shepherd (2014), who emphasised that evidence needs to flow through the ecosystem from generation to end-user, requiring both push and pull mechanisms.

## Objectives

1. To carry out a systematic review of quantitative data relating to brief drug interventions

2. To carry out a systematic review of qualitative data relating to brief drug interventions
3. To carry out qualitative work with practitioners and community members to understand the barriers and facilitators to carrying out brief drug interventions.
4. To develop a research team with appropriate skills and expertise, including community sector
5. To develop an application for conducting a trial on the effectiveness and cost-effectiveness of targeted drug screening and brief interventions (SBIs), including extended brief interventions, in reducing drug use and associated harm across health, social care, and justice settings in the UK, and to compare the effectiveness across these different settings.

## Objective one: systematic review of quantitative data relating to brief drug interventions

The aim of this systematic review was to ascertain the effectiveness of brief interventions in reducing drug use. Secondly, we wanted to explore different screening tools and interventions used in studies. Finally we assessed the key ingredients of included studies using the TIDieR framework (Hoffman *et al.*, 2014).

## Methods

The protocol for this systematic review was registered with PROSPERO (CRD42023429734). The review is reported in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2022 reporting guidelines (Rethlefsen *et al.*, 2021).

## Search strategy

Six electronic databases were searched: MedLine, PsychINFO, EMBASE, CINAHL, SCOPUS, and Cochrane Library, and four grey literature databases were searched; MEDNAR, Google Scholar, Google, and opengrey.au, with the first 100 hits being retrieved (Haddaway *et al.*, 2015).

Searches of all databases were conducted in line with both the PICO (Patient or problem, Intervention or exposure, Comparison or control, and Outcomes) (Rethlefsen *et al.*, 2021) and SPIDER (Sample, Phenomenon of Interest, Design, Evaluation, and Research type) (Cook *et al.*, 2012) frameworks for conducting literature searches for quantitative analysis. Elements of the SPIDER screening tool (Cook *et al.*, 2012) was used to inform the key words and identify relevant papers. Sample included adults, Phenomenon of Interest included brief intervention and substance use screening, Design included Randomised controlled trials (RCTs), and Research type included quantitative. All searches were conducted in September 2023.

## Eligibility criteria

Papers were included if they used quantitative research methods, specifically RCTs, to explore the properties of brief

interventions for adults aged over 18 years old for substance use.

**Inclusion criteria:** Data relating to participants 18 or over could be abstracted; Any language; Data published from 2003 to ensure data was relevant.

**Exclusion criteria** Data related to participants under the age of 18; Data published prior to 2003.

### *Study selection and data management*

All results from the database search were imported into End-Note for storage, duplication detection, and sifting. The lead reviewer (DNB) screened all the titles and abstracts against the inclusion criteria. A second reviewer in the team independently double-screened 20% of the papers. No discrepancies could not be agreed on.

Papers that were identified as potentially relevant went through the second sifting phase of full paper screening as per guidelines (Rethlefsen *et al.*, 2021). All full texts were retrieved and saved on Microsoft Teams for review. One reviewer (ET) sifted all full papers, and a second reviewer (DNB) independently double-screened 20%.

### *Data extraction*

A Microsoft Excel spreadsheet was developed for the data extraction which captured; the authors, year of publication, country of study, the aim of the research, study design, setting of the research, sample size, description and length of intervention, screening tool used, drug type, participant demographics, primary and secondary outcomes, results, effectiveness, and recommendations.

A TIDieR table was also created to ascertain what the ingredients of the interventions were (Hoffman *et al.*, 2014). ET undertook the data extraction with 20% checked by another team member.

### *Assessment of quality*

The CASP (CASP-UK, 2002) screening tool for appraisal of RCTs was used for the quality assessment of the included papers. The purpose of the CASP tool was to assess quality; it was not used to contribute to decisions about whether or not to include studies. A high risk of bias was noted if “no” or “unsure” was recorded for 6 or more of the 11 questions on the tool. A medium risk of bias was assigned if “no” or “unsure” was recorded for 4–5 questions, and a low risk of bias for 1–3 questions, consistent with our previous studies (Newbury-Birch *et al.*, 2018; Newbury-Birch *et al.*, 2022).

### *Results*

The initial searches yielded 17,270 records. In total, 46 met the inclusion criteria and were included in the review (Aharonovich *et al.*, 2017; Assanangkornchai *et al.*, 2015; Bagøien *et al.*, 2013; Baker *et al.*, 2005; Barrowclough *et al.*, 2010; Blow *et al.*, 2017; Boden *et al.*, 2012; Carroll *et al.*, 2009; de Oliveira Christoff & Boerngen-Lacerda 2015; Fischer *et al.*, 2013; Forsberg *et al.*, 2011; Gates *et al.*, 2012;

Gelberg *et al.*, 2015; Gelberg *et al.*, 2017; Gmel *et al.*, 2013; Goodness & Palfai, 2020; Graham *et al.*, 2016; Guan *et al.*, 2015; Heather *et al.*, 2004; Hoch *et al.*, 2014; Humeniuk *et al.*, 2018; Jonas *et al.*, 2012; Jungerman *et al.*, 2007; Kavanagh *et al.*, 2004; Kim *et al.*, 2017; Lerch *et al.*, 2017; McKee *et al.*, 2007; Merchant *et al.*, 2018; Mertens *et al.*, 2014; Mitcheson *et al.*, 2006; Nagel *et al.*, 2009; Palfai *et al.*, 2014; Poblete *et al.*, 2017; Prendergast *et al.*, 2017; Roy-Byrne *et al.*, 2014; Satre *et al.*, 2016; Shekhawat *et al.*, 2023; Sobell *et al.*, 2009; Sorsdahl *et al.*, 2021; Stephens *et al.*, 2004; Thompson *et al.*, 2020; van Emmerik-van Oortmerssen *et al.*, 2019; Wernett *et al.*, 2018; Woodruff *et al.*, 2014; Woolard *et al.*, 2013; Zahradnik *et al.*, 2009) (Figure 1). The articles were published between 2004 and 2023.

The characteristics from the 46 included papers are presented in Extended Data (Table 1) Twenty-one papers reported on research from the United States of America (USA), five from Australia, four from the United Kingdom (UK), three from Germany, two each from Brazil, Canada, South Africa and one each from Sweden, Chile, India, Norway, Switzerland, Thailand, and the Netherlands. The included studies consisted of 14,048 (range 25-1030). Most of the studies had a higher percentage of male participants.

Thirty-six papers were in a health-based setting, three in a criminal justice-based setting, three in a general setting, three in universities, and one in another setting (Extended Data (Table 1)). All papers used RCTs to measure effectiveness of the interventions. Several tools were used to screen participants for substance use with the main tools being, the Timeline Follow Back (TLFB) and versions of the ASSIST (Humeniuk *et al.*, 2010)(Extended Data (Table 1)).


Results showed that screening for substance use was carried out by therapists (n=12), researchers (n=9), clinicians (n=5), psychologists (n=5), counsellors (n=4), health staff (n=2), trained interventionists (n=2), psychiatrists (n=1), GP's (n=1), practitioners (n=1), and social workers (n=1). Three papers did not include this information (Extended Data (Table 1)).

### *Health Settings*

Thirty-six studies delivered a brief intervention in a health-based setting (Aharonovich *et al.*, 2017; Assanangkornchai *et al.*, 2015; Bagøien *et al.*, 2013; Barrowclough *et al.*, 2010; Blow *et al.*, 2017; Boden *et al.*, 2012; Carroll *et al.*, 2009; Gelberg *et al.*, 2017; Gelberg *et al.*, 2015; Gmel *et al.*, 2013; Goodness & Palfai, 2020; Graham *et al.*, 2016; Guan *et al.*, 2015; Heather *et al.*, 2004; Hoch *et al.*, 2014; Humeniuk *et al.*, 2018; Jungerman *et al.*, 2007; Kavanagh *et al.*, 2004; Kim *et al.*, 2017; Mertens *et al.*, 2014; Merchant *et al.*, 2018; McKee *et al.*, 2007; Mitcheson *et al.*, 2006; Nagel *et al.*, 2009; Palfai *et al.*, 2014; Poblete *et al.*, 2017; Roy-Byrne *et al.*, 2014; Satre *et al.*, 2016; Shekhawat *et al.*, 2023; Sobell *et al.*, 2009; Sorsdahl *et al.*, 2021; van Emmerik-van Oortmerssen *et al.*, 2019; Wernett *et al.*, 2018; Woodruff *et al.*, 2014; Woolard *et al.*, 2013; Zahradnik *et al.*, 2009).

These were in; primary care (n=9), outpatient units (n=10), emergency departments (n=5), hospitals/GP surgeries (n=4),





**Figure 1. Quantitative Systematic Review PRISMA.**

health centres (n=4), inpatient units (n=3), and remotely (n=1). Brief interventions lasted between 3–120 minutes. The control conditions consisted of; TU/standard care (n=13), handouts on health or substance use (n=5), videos and booklets (n=4), delayed treatment (n=4), assessments of substance use (n=2), simple advice (n=2), 3 sessions of CBT (n=1), questionnaire (n=1), feedback (n=1), enhanced care as usual and handout (n=1), individual MI (n=1), and an attention placebo (n=1). In terms of quality assessment, eighteen studies were identified as medium quality, ten as low quality, and eight as high quality. The thirty-six studies looked at different primary outcome measures including days of substance use (for example the number of days used any drug; frequency of drug use; and reduction in drug use); substance use severity and changes in substance

use; abstinence and treatment engagement; and specific drug use (for example benzodiazepines and cannabis use). Secondary outcomes included psychological and mental health outcomes; behavioural and treatment engagement; social and legal aspect; re-admission to hospital; and treatment cost effectiveness (Extended Data (Table 2)).

Ten (Gmel *et al.*, 2013; Kim *et al.*, 2017; Merchant *et al.*, 2018; Mertens *et al.*, 2014; Nagel *et al.*, 2009; Palfai *et al.*, 2014; Poblete *et al.*, 2017; Roy-Byrne *et al.*, 2014; van Emmerik-van Oortmerssen *et al.*, 2019; Woodruff *et al.*, 2014) of the studies found no significance in relation to substance-use, however, some of the studies found significance in other areas including; the BI group having lower odds of linking to treatment

(Kim *et al.*, 2017), reductions in alcohol ASSIST scores by 38% (Mertens *et al.*, 2014), BI groups reporting better well-being (Nagel *et al.*, 2009); reduced perceived norms regarding peer marijuana use (Palfai *et al.*, 2014), and reducing ADHD symptoms (van Emmerik-van Oortmerssen *et al.*, 2019). Two studies found significance on the impact on readiness to change drug use including at 12 months (adjusted OR 2.05, 95% CI 1.26 to 3.31;  $P=0.004$ ) (Barrowclough *et al.*, 2010) and a 63% increase in engaging with treatment (Graham *et al.*, 2016).

Twenty-four studies found significantly greater reductions in substance use. Aharonovich *et al.* (2017) reported a 51% reduction in the frequency and quantity of non-injection drug use within the MI-Only group (Aharonovich *et al.*, 2017). Assanangkornchai *et al.* (2015) demonstrated substantial reductions in both ASSIST-SSIS and ASSIST-TSIS scores (Assanangkornchai *et al.*, 2015). Bagøien *et al.* (2013) revealed a 2-year net difference of 7.3 days of substance use per month, favouring the intervention group (95% CI 1.9 to 12.6,  $p < 0.01$ ) (Bagøien *et al.*, 2013) (Bagøien *et al.*, 2013). Blow *et al.* (2017) highlighted the effectiveness of Therapist BI in reducing the number of days using any drug [95% confidence interval (CI)=0.41, 0.07,  $P = 0.0422$ ] and weighted drug-days (95% CI = -0.41, 0.08,  $P = 0.0283$ ). Gelberg *et al.* (2015) and Gelberg *et al.* (2017) showcased significant reductions in the use of the highest scoring drug (HSD) ( $p < 0.042$ , 95% CI: 0.2, 8.7). Goodness and Palfai (2020) observed a small to medium effect on cannabis use frequency ( $f^2 = 0.09$ ), while Guan *et al.* (2015) reported substantial decreases in total InDUC scores in the treatment arm (57.6%).

Heather *et al.* (2004) found larger reductions in benzodiazepine consumption in the letter and consultation groups (24% overall), and Hoch *et al.* (2014) reported a significantly higher abstinence rate in Assertive Treatment (AT) patients compared to Day Treatment Center (DTC) patients. Boden *et al.* (2012) revealed that the BI group was associated with significantly better drug use outcomes ( $P < 0.05$ ). Carroll *et al.* (2009) demonstrated reductions in substance use during the 4-week therapy phase, with significant main effects for time ( $t(5740) = -3.0$ ,  $p < .01$ ), phase ( $t(5740) = -2.79$ ,  $p < .01$ ), and their interaction ( $t(5740) = 2.64$ ,  $p < .01$ ). Humeniuk *et al.* (2018) reported a significant reduction in total illicit substance and Amphetamine-Type Stimulants involvement for participants receiving the ASSIST-linked BI, compared with the control group ( $P < 0.001$ ). Jungerman *et al.* (2007) found that both treatments outperformed Day Treatment Center, with particularly notable results for cannabis use ( $p = 0.0002$ ).

Kavanagh *et al.* (2004) highlighted that all 13 participants in the BI group proceeding to MI reported less substance use at 6 months compared to 58% in the Screening and Consultation alone group. McKee *et al.* (2007) demonstrated that participants receiving MET and CBT attended more drug treatment sessions, reported a greater desire for abstinence, and expected greater difficulty in maintaining abstinence compared to the CBT condition. Satre *et al.* (2016) found that at 6 months, Motivational Interviewing (MI) was more effective than control

in reducing the rate of cannabis use and hazardous drinking. Mitcheson *et al.* (2006) observed a large and statistically significant reduction in heroin use among those in the MI condition (unstandardised regression coefficient = -2.04, 95% confidence interval -4.44 to 0.37,  $p = 0.093$ ). Shekhawat *et al.* (2023) reported that the BI group had fewer days of cannabis use than the simple advice group at 4, 8, and 12 weeks. Sobell *et al.* (2009) demonstrated significant and large reductions in alcohol and drug use during treatment and at the 12-month follow-up ( $F(2, 43) 26.71$ ,  $p .001$ ). Sorsdahl *et al.* (2021) found a significantly lower frequency of methamphetamine use in the treatment group at both the 6-week and 3-month endpoints ( $r = -4.63$ ,  $p < 0.01$ ). Wernett *et al.* (2018) noted consistently high ratings of acceptability for the intervention, and participants reported a significantly larger reduction (54%) in any marijuana or alcohol use compared to the control group at the 4-month follow-up. Woolard *et al.* (2013) reported a significant decrease in binge drinking and conjoint marijuana and alcohol use for the treatment group ( $M=0.72$ ; 95% CI=0.36–1.12) compared to the standard care group ( $M=1.77$ ; 95% CI=1.19–1.57). Lastly, Zahradnik *et al.* (2009) revealed that after 3 months, more participants in the intervention group reduced their Defined Daily Dosage compared to the control group (51.8% versus 30%;  $c2 = 6.17$ ;  $P = 0.017$ ).

#### **Criminal Justice Setting: Brief Interventions**

Three studies delivered a brief intervention in a criminal justice-based setting for adults over the age of 18 (Forsberg *et al.*, 2011; Lerch *et al.*, 2017; Prendergast *et al.*, 2017); prison ( $n=1$ ), probation ( $n=1$ ), and the police ( $n=1$ ). Each brief intervention lasted between 15-45 minutes. The control conditions consisted of; usual planning interview ( $n=1$ ), standard probation intake ( $n=1$ ), and substance use information ( $n=1$ ). In terms of quality assessment all three studies were identified as being of medium quality. The three studies looked at different primary outcome measures including prior 30 days of drug or alcohol use; treatment involvement; and reduction in drug or alcohol use. Secondary outcomes included days of illegal activity; heavy alcohol or drug use; and post-release participation in treatment, rearrest, quality of life, and HIV risk behaviours. There were no significant results in relation to substance use, however, there were significant findings relating to the increase in the number of days working by intervention participants (Forsberg *et al.*, 2011) and the increase in reporting treatment initiation (Lerch *et al.*, 2017).

#### **General Settings: Brief Interventions**

Four studies (Gates *et al.*, 2012; Jonas *et al.*, 2012; Stephens *et al.*, 2004; Thompson *et al.*, 2020) delivered a brief intervention in a general-based setting for adults over the age of 18 including; a cannabis information helpline ( $n=1$ ), a website ( $n=1$ ), a research centre ( $n=1$ ), and in a shelter ( $n=1$ ). Each brief intervention lasted between 20-90 minutes. The control conditions consisted of; TAU ( $n=1$ ), no treatment offered ( $n=1$ ), and delayed feedback ( $n=1$ ). The four studies looked at different primary outcome measures including frequency and quantity of cannabis use; number of days of marijuana and alcohol use; and sexual risk behaviours. Secondary outcomes included



other drug-use; and willingness to change. [Gates et al. \(2012\)](#) found those who were given a BI reported greater reduction in dependence symptoms ( $P < 0.001$ ,  $d = 0.9$  [0.5–1.3]) and greater confidence to reduce cannabis use at 4 weeks ( $P = 0.002$ ,  $d = 0.5$  [0.1–0.9]), reporting more abstinent days at 12 weeks ( $P = 0.019$ ,  $d = 0.6$  [0.2–1.0]). [Stephens et al. \(2004\)](#) found those given the intervention reported fewer days of marijuana use per week, fewer periods of use per day, and fewer dependence symptoms at 7 weeks, 6 months, and 12 months follow-ups ( $P = 0.019$ ;  $d = 0.45$ ). Lastly, [Thompson et al. \(2020\)](#) found intervention participants significantly reduced past two-week number of drinks ( $p=.023$ ), times used marijuana ( $p=.046$ ), times engaged in unprotected sex ( $p=.012$ ), and times used drugs before sexual activity ( $p=.019$ ). [Jonas et al. \(2012\)](#) did not find any significant results.

### University Settings: Brief Interventions

Three studies delivered a brief intervention in a university-based setting for adults over the age of 18 ([Baker et al., 2005](#); [de Oliveira Christoff & Boerngen-Lacerda, 2015](#); [Fischer et al., 2013](#)). Each brief intervention lasted between 5–50 minutes. The control conditions consisted of; a self-help booklet ( $n=1$ ), feedback ( $n=1$ ), and a BI on general health ( $n=1$ ). In terms of quality assessment, all three studies were identified as being of high quality. The three studies looked at different primary outcome measures including change in amphetamine use; drug and alcohol use; and the feasibility and impact of a cannabis BI. Secondary outcomes included changes in other drug use. [Baker et al. \(2005\)](#) found participants given a BI significantly increased in the likelihood of abstinence from amphetamines among those receiving two or more treatment sessions (1.04 units compared to 0.76 units). [de Oliveira Christoff & Boerngen-Lacerda \(2015\)](#) found a small positive effect in the ASSIST/MBII and control groups for marijuana (Q2 and Q4 in ASSIST/MBII,  $p = .05$ ; Q4 in control,  $p = .02$ ). Lastly, [Fischer et al. \(2013\)](#) did not find any significant results in relation to substance use, however, they did find the prevalence of driving after cannabis use fell from 44.44% to 30.65% ( $p = 0.020$ ) in the combined intervention groups (Extended Data (Table 2)).

### Quality assessment

Ten of the included studies were classified as low chance of risk, 25 as medium chance of risks and 11 as high chance of risk (Extended Data (Table 1)).

### TIDieR results

Most of the interventions were conducted face-to-face. The majority of the studies reported the intervention was based on the work of Miller and Rollnick ([Miller & Rollnick, 1991](#)) with many studies giving a lot of information in relation to the studies (Extended Data (Table 3 and Table 4)

Extended data (Table 5)

## Objective two: systematic review of qualitative data relating to brief drug interventions

The aim of this systematic review was to identify, explain and interpret the prominent or recurring themes relating to the barriers and facilitators of embedding SBI across three settings (health, social care and criminal justice).

## Methods

The protocol for this systematic review was registered with PROSPERO (CRD42023429726). The review is reported in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2022 guidelines ([Rethlefsen et al., 2021](#)), ensuring a structured and transparent approach. This includes a comprehensive checklist and flow diagram covering all aspects of the review process. Key elements are detailed, including clear eligibility criteria, thorough search strategies, systematic data extraction, and rigorous risk of bias assessments. The methods section outlines information sources, study selection, and data synthesis procedures. Results are presented with a flow diagram, study characteristics, and bias assessments. The discussion summarises evidence, limitations, and conclusions, and the funding section details sources and their roles. This adherence enhances the reliability, reproducibility, and utility of the review in evidence-based practice and policy.

### Search strategy

Six electronic databases were searched: MedLine, PsychINFO, EMBASE, CINAHL, SCOPUS, and Cochrane Library, and four grey literature databases were searched; MEDNAR, Google Scholar, Google, and opengrey.au, with the first 100 hits being retrieved ([Haddaway et al., 2015](#)).

Searches of all databases were conducted in line with both the PICO (Patient or problem, Intervention or exposure, Comparison or control, and Outcomes) ([Rethlefsen et al., 2021](#)) and SPIDER (Sample, Phenomenon of Interest, Design, Evaluation, and Research type) ([Cook et al., 2012](#)) frameworks for conducting literature searches for quantitative analysis. Elements of the SPIDER tool was used to inform the key words and identify relevant papers. Sample included adults, Phenomenon of Interest included brief intervention and substance use screening, Design included qualitative methods such as interviews and focus groups, and Research type included qualitative. All searches were conducted in September 2023.

### Eligibility criteria

Papers were included if they used qualitative research methods to explore the barriers and facilitators of screening and brief interventions for adults aged over 18 years old for substance use.

**Inclusion criteria:** Data relating to participants 18 or over could be abstracted; any language; data published from 2003 to ensure data was relevant; studies that included a qualitative element (including from survey results).

**Exclusion criteria:** Data related to participants under the age of 18; data published prior to 2003; studies that only included quantitative data.

### Study selection and data management

All results from the database search were imported into End-Note for storage, duplication detection, and sifting. The lead reviewer (JF) screened all the titles and abstracts against the inclusion criteria. A second reviewer in the team independently double-screened 20% of the papers. No discrepancies could not be agreed on.

Papers that were identified as potentially relevant went through the second sifting phase of full paper screening as per guidelines (Rethlefsen *et al.*, 2021). All full texts were retrieved and saved on Microsoft Teams for review. One reviewer (JF) sifted all full papers, and a second reviewer independently double-screened 20%.

### **Data extraction**

A Microsoft Excel spreadsheet was developed for the data extraction which captured; the authors, year of publication, country of study, the aim of the research, study design, methods, setting of the research, sample size, description of intervention, length of intervention, barriers to interventions, facilitators to interventions, participant demographics, recommendations and conclusions. LT undertook the data extraction with 20% checked by another team member (JF).

### **Assessment of quality**

As some of the included studies were mixed methods, the relevant CASP tool for appraisal was used (CASP-UK, 2002) for the quality assessment of the included papers. The purpose of the CASP tool was to assess quality; it was not used to contribute to decisions about whether or not to include studies. High risk of bias was recorded if “no” or “unsure” was recorded for 6 or more of the 11 questions on the tool. Medium risk of bias was assigned if “no” or “unsure” was recorded for 4–5 questions and Low risk for 1–3 questions, as in our previous study (Newbury-Birch *et al.*, 2018; Newbury-Birch *et al.*, 2022). 517 interviews were included (range 1–73).

### **Results**

The initial searches yielded 22,609 records. In total, 14 met the inclusion criteria and were included in the review (Aharonovich *et al.*, 2012; Darker *et al.*, 2016; Dujon, 2021; Fazio *et al.*, 2022; Hunter *et al.*, 2018; Morris *et al.*, 2024; Owens *et al.*, 2018; Roberts & Nuru-Jeter, 2010; Rudzinski *et al.*, 2012; Saunders *et al.*, 2019; Sharma *et al.*, 2023; Starks *et al.*, 2020; Venner *et al.*, 2018; Whiteside *et al.*, 2010) (Figure 2). The articles were published between 2019–2023. Eleven were from the USA and one each from Canada, Ireland and India.

### **Qualitative systematic review: themes**

Using thematic synthesis (Braun & Clarke, 2006), five main themes emerged: 1. Barriers within a health setting; 2. Facilitators within a health setting; 3. Barriers within a social care setting; 4. Facilitators within a social care setting; 5. Barriers within a criminal justice setting. Extended Data (Table 5) contains examples of quotes demonstrating each theme. Perspectives in papers varied and included the perspectives from providers and those receiving the intervention.

#### **Health setting barriers**

Within a health setting, multiple barriers to interventions were identified including physical barriers, participants attitude to the intervention, patient internal conflicts, cost and sustainability, influence of provider, missed opportunity for intervention, unsuitability of the intervention, and influence of drug type.

#### **Physical barriers**

One paper raised issues around physical barriers to accessing interventions, as relocation prevented some from participating in the study. Providers also assisted participants in accessing a telephone, which suggests that without this access, participation would not be possible (Aharonovich *et al.*, 2012).

#### **Attitude to the intervention**

Three papers mentioned that participant attitudes towards the intervention may act as a barrier, as some highlighted difficulties remembering to take part, found taking part ‘annoying’, or would not engage with the materials provided outside of the intervention itself (Aharonovich *et al.*, 2012; Darker *et al.*, 2016; Fazio *et al.*, 2022).

#### **Patient internal conflicts**


Patients raised internal challenges around disclosing their drug use, factors such as a lack of coping skills, a lack of self-esteem, lack of self-interest/knowledge they had a problem and feelings of dismissal or judgement from providers resulted in patients not sharing information around substance use or changing providers (Dujon, 2021). Expected consequences of disclosure were also a barrier for some, with patients concerned around the potential implications of disclosing their substance use to providers. Patients feared this could have legal implications or impact employment, future medical care, or result in adverse psychological or social consequences (Roberts & Nuru-Jeter, 2010; Saunders *et al.*, 2019); Some concealed their drug use to avoid these consequences (Roberts & Nuru-Jeter, 2010). Additionally, concerns around the privacy of patients’ substance use information was highlighted as a barrier, and patients were wary of other providers potential access to their medical record. Attitudes towards screening methods were also influenced by privacy concerns, due to the risk of being ‘hacked’ or unauthorised parties accessing their data (Saunders *et al.*, 2019).

#### **Cost and Sustainability**

The resource cost of implementing and sustaining interventions may be a potential barrier as staff and resources are often already limited within healthcare settings such as the emergency department (Fazio *et al.*, 2022). Capacity and staff training may be a barrier to continued implementation of interventions as staff turnover and the uncertainty surrounding funding for positions may cause the practice to ‘slip away’ (Hunter *et al.*, 2018). The cost of the interventions themselves, specifically drug assisted treatment, was a particular concern for the continued implementation of interventions, with providers questioning how the expense would be sustained (Hunter *et al.*, 2018). The ongoing fidelity of structure given to interventions was also highlighted as a potential barrier, with concerns raised around the impact of time upon the continuation of the approach and how that could be maintained (Hunter *et al.*, 2018).

#### **Influence of provider**

When exploring patients’ substance use, three papers found that patient’s relationship with their provider influenced the



**Figure 2. Qualitative Systematic Review PRISMA.** We confirm we own and have permission to use the figures accompanying our submission.

interaction and could be a barrier to disclosing or discussing substance use. A lack of trust in the physician/provider and fear and experiences of judgement/stigma were highlighted as influential on the relationship between patient and provider which then influenced patients' likelihood of honestly disclosing substance use (Dujon, 2021; Roberts & Nuru-Jeter, 2010; Saunders *et al.*, 2019). Provider constraints were identified as potential barriers, as the providers discomfort, lack of preparedness and lack of time with the patient when addressing substance use negatively impacted the interaction (Saunders *et al.*, 2019). Additionally, two papers mentioned provider shortfalls as potential barriers; a negative approach from the provider or the providers poor communication style were not appreciated by patients and resulted in decisions to not disclose information or change provider (Dujon, 2021). If patients perceived their provider as not listening to general health concerns, some

responded by emotionally disengaging (Roberts & Nuru-Jeter, 2010).

#### *Missed opportunity for intervention*

Three papers highlighted how current practise resulted in missed opportunity for intervention. While some providers assessed substance use annually or at new patient visits, current screening practices for substance use were not systematic (Saunders *et al.*, 2019). Additionally, universal screening frequently did not occur and this rate was even lower for opiates (Venner *et al.*, 2018). This was supported by patients who frequently voiced they had never been or did not recall ever being asked about their substance use during healthcare visits (Dujon, 2021). Providers reported using their clinical judgement to detect substance use and focus their attention on those with established substance use problems rather than screening

universally, resulting in those at the ‘tip of the pyramid’ being seen more than those in the middle (Venner *et al.*, 2018). Venner *et al.* (2018) found that only half of the providers reported that patients felt comfortable seeking treatment for substance use problems, further limiting opportunity for intervention (Venner *et al.*, 2018).

### ***Unsuitability of intervention***

The need for interventions to be contextually and culturally appropriate to the service user within this setting was highlighted by Darker *et al.* (2016) (Darker *et al.*, 2016). Clinicians felt that materials within the intervention, such as reasons to reduce substance use, need to be relevant to the patient and their experiences to be successful and encourage behaviour change. It was felt that unsuitable materials would have minimal impact.

### ***Influence of drug type***

One paper highlighted how the type of drug being used may be a barrier to intervention. Those using marijuana for medicinal purposes may be reluctant to reduce consumption in contrast to those using other substances who while ambivalent, perceived the need to reduce their substance use (Aharonovich *et al.*, 2012). Cannabis was mentioned further by Saunders *et al.* (2019) who felt that the legalisation of cannabis in some areas may result in patients no longer viewing cannabis as a drug and ignoring their problematic use due to perceptions of safety, thus acting as a barrier to intervention and requiring education on the risks of cannabis use to overcome them (Saunders *et al.*, 2019).

### **Health setting facilitators**

Within a health setting, facilitators identified included the influence of provider, suitability of the intervention to the individual, accessible and well understood intervention, sustainability, intervention features, screening type, participant engagement, suitability of the intervention to the provider, and influence of drug type.

### ***Influence of provider***

While the patient’s relationship with the provider could act as a barrier to disclosing information around substance use, four papers mentioned this could also facilitate. Staff training on the use of empathetic, conversational approaches that are less likely to appear confrontational or intimidating would be useful to improve communication between patient and provider and minimise perceptions of judgement from providers (Roberts & Nuru-Jeter, 2010). When the interaction between provider and patient was perceived positively, such as a caring, trustworthy (Fazio *et al.*, 2022), understanding or supportive provider (Roberts & Nuru-Jeter, 2010), patients were encouraged to discuss or reduce substance use. This approach from the provider made patients feel understood, comfortable and important, and they valued the opportunity to ‘get things off their chest’ (Fazio *et al.*, 2022). Provider connectedness and patient familiarity with the provider was identified by providers as ‘critical’ to promote honest disclosure of substance use as it was important for patients to feel comfortable and not judged (Fazio *et al.*, 2022; Saunders *et al.*, 2019). highlighted the

influence of a relatable provider, with some patients feeling that a provider who had been through similar experiences of substance use would be valuable. The provider would be able to understand and provide support and direction based on their first-hand experience of what is effective and where to go. Roberts & Nuru-Jeter (2010) found that while some feared providers identifying their substance use, some felt if their provider had recognised this, it would have made it easier to discuss and seek help.

### ***Suitability of intervention to the user***

The need for the intervention to be suitable to the individual/cohort it is targeting was demonstrated (Darker *et al.*, 2016). Both clinicians and patients mentioned how examples within interventions need to be tailored to be specific to the cohort of users to have the most impact as patient concerns vary across groups. Patients may be more concerned with psychosocial risks than physical health risks for example, and as this holds more meaning, may have greater impact in the context of an intervention. The intervention itself could also be tailored to the patient based upon their individual risk profile using tick boxes for the provider.

### ***Accessible and well understood intervention***

Three papers discussed the need to ensure interventions and associated materials were accessible and well understood by both providers and patients. Accessible language and the use of imagery to help patients with literacy difficulties understand key risks were included, and suggestions were made to simplify the language (Darker *et al.*, 2016). The development of a clear, easy to use manual was valuable to providers due to the time limited environment in which they work, including a sample script and roleplay for training purposes, providers also felt this was helpful (Darker *et al.*, 2016). Within the paper by Aharonovich *et al.* (2012), all patients highlighted the use of telephone calls within the intervention as useful and felt this was easy to do, of which may facilitate participation (Aharonovich *et al.*, 2012). To improve patients understanding of data privacy and potentially facilitate information sharing, clear policies and communication around confidentiality of the information obtained is recommended (Roberts & Nuru-Jeter, 2010).

### ***Sustainability***

Two papers mentioned the need for providers to access training and resources to successfully conduct screening and interventions for substance use. Leadership and staff support, program champions, additional training opportunities for providers, low cost and accessible screening and access to additional funding were all highlighted as facilitators to sustaining the implementation of interventions (Hunter *et al.*, 2018; Saunders *et al.*, 2019). Hunter *et al.* (2018) found that making adaptations to some aspects, such as increasing the time between screenings, was necessary.

### ***Intervention features***

Aharonovich *et al.* (2012) found that participants commented on particular aspects of the intervention positively. Patients found the intervention ‘exciting’ and useful, that it provided



positive reinforcement and aided patients in being focused and alert. Being able to take accountability for their substance use and track achievement using graphs was also valuable to patients (Aharonovich *et al.*, 2012).

### **Screening type**

When screening for substance use, Saunders *et al.* (2019) found that the type of screening utilised was important (Saunders *et al.*, 2019). Universal screening was perceived to be less 'accusatory' to patients and providers felt this was necessary to identify substance use in those that would likely be missed by targeted approaches. Providers agreed that annual screening was most suitable, with an exception for patients with indicators of substance use, as regular screening may frustrate patients. Screening patients once a relationship with the provider had developed, as opposed to meeting for the first time, was mentioned as a facilitator to engaging patients. Self-administered screening was supported by both patients and providers, as this increased patient comfort and increased validity and efficiency in comparison to interviewer administered screening. However, some providers highlighted patients' dislike for completing forms, poor reading comprehension, lack of honesty and privacy concerns, or were concerned that screening not conducted as an interview may not be reviewed.

### **Participant engagement**

Participant engagement with the intervention was highlighted as a facilitator in three papers. Interactive interventions were praised by patients, highlighting how they provided 'attention never received from doctors' (Aharonovich *et al.*, 2012). Patients' personal goals such as access to children may also facilitate participation in interventions, as the goal may act as a motivator to engage (Roberts & Nuru-Jeter, 2010). In one intervention, telephone call reminders to engage with the intervention were mostly appreciated by patients (Aharonovich *et al.*, 2012). Risk was highlighted by Darker *et al.* (2016) as a facilitator to engagement, as materials reminding patients of risk associated with substance use or the use of emotional/impactful imagery demonstrating risk were recommended.

### **Suitability of intervention to provider**

When the treatment was perceived as fitting with the organisation/clinic mission and could be institutionalised within practice, this could facilitate the continuation of provision. The perspective the approach had given could not be undone, and care for alcohol and opioid use disorder was institutionalised into the practice model within the clinic (Hunter *et al.*, 2018). Collaborative care models and increased practice coordination with community resources were endorsed by providers as facilitators to screening and brief intervention (Venner *et al.*, 2018).

### **Influence of drug type**

The type of drug used by the patient may act as a facilitator to intervention. Saunders *et al.* (2019) mentioned how due to the legalisation of cannabis in some areas, open dialogue can occur around use of the drug as patients are more comfortable

discussing their use and there is less concern around potential consequences.

### **Social setting/university barriers**

Barriers to interventions within a social setting were identified as logistical barriers, a lack of interaction/engagement, privacy concerns and implications of disclosure, provider approach and conflict within sessions.

### **Logistical barriers**

Two papers mentioned logistical difficulties as a barrier to intervention. One highlighted the influence of technical difficulties on participant engagement as the participant had to redo the task (Sharma *et al.*, 2023). Another mentioned a reduction in participant numbers may have been due to challenges with physically attending the intervention within the institution as the students who took part predominantly commuted in to attend college (Morris *et al.*, 2024).

### **Lack of interaction or engagement**

Rudzinski *et al.* (2012) found that interventions that did not engage participants or lacked interaction with a provider were viewed less favourably by those who took part (Rudzinski *et al.*, 2012). Being able to ask questions was felt to be important, and printed interventions such as pamphlets were criticised as being 'outdated' and easy to discard. This lack of engagement was expanded on further by Sharma *et al.* (2023) where using language that was difficult for students to understand prevented participants from engaging with the intervention.

### **Privacy concerns and implications of disclosure**

Participants had concerns around drug use disclosure and their privacy, and this was a barrier to recruitment in one study. Participants sought reassurance that their data would be stored securely and could not be accessed by anyone outside of the study and were concerned that sharing their data may result in legal issues (Sharma *et al.*, 2023). Although these were mentioned in the context of recruitment, similar concerns may emerge in relation to participating in an intervention that would require substance use disclosure.

### **Provider approach**

The approach utilised by the provider may become a barrier to engaging with an intervention if the approach is unsuitable for the individual. Whiteside *et al.* (2010) highlighted how a facilitator focusing on increasing an individual's commitment to change stage and feel ambivalent may result in resistance or lowered autonomy in the participant. This emphasizes the need for a balanced approach from the provider.

### **Conflict within session**

One paper exploring an intervention involving couples found that conflict between the two partners may be a barrier through conflation of thoughts and feelings, vague or indirect communication, and incorrect assumptions (Starks *et al.*, 2020). This conflict potentially restricts the extent that partners can



understand and empathise with each other and seek resolutions while increasing the likelihood of defensiveness, blame or further conflict.

### Social setting/university facilitators

Facilitators to interventions within a social setting were identified as the suitability of the intervention to the individual, influence of provider, provider approach, accessibility, intervention features and participant engagement.

#### *Suitability of the intervention to the individual*

Two papers discussed the suitability of the intervention to the individual targeted as a facilitator to the intervention. Interventions are appreciated when they are tailored to the needs of the individual and are appropriate to that population (Morris *et al.*, 2024). By personalising and tailoring the intervention to the individual, advice can be provided that is specific to the individual and their circumstances (Rudzinski *et al.*, 2012). Materials that are not appropriate to the population, such as those using unrelatable language, may not be suitable (Rudzinski *et al.*, 2012).

#### *Influence of provider*

The provider involved in the intervention was influential in the interaction. Providers were valued when they were perceived as being supportive, non-judgemental and caring towards the participant (Morris *et al.*, 2024). Establishing rapport between the provider and the individual receiving the intervention was also highlighted as beneficial for increasing the effectiveness of the intervention, with further appreciation for the providers friendly, reflective, non-judgemental stance without pressure to accept goals (Whiteside *et al.*, 2010). Rudzinski *et al.* (2012) further highlighted the appreciation for interventions that were 'straightforward, unbiased, nonthreatening, nonpatronizing and non-judgemental' (Rudzinski *et al.*, 2012).

#### *Provider approach*

A provider utilising a balanced approach during the intervention can facilitate the clients engagement with the intervention and reduce resistance (Morris *et al.*, 2024), additionally a balanced approach reduces the likelihood that the receiver will feel lectured or as though the intervention is inappropriate to their circumstances (Whiteside *et al.*, 2010). The providers response to conflict within interventions delivered to couples may facilitate behaviour change by achieving consensus between partners around intervention goals and recognising potential barriers to behaviour change. By responding appropriately, the provider can shift focus, reflect, clarify thoughts and feelings, establish effective communication and correct inaccurate assumptions (Starks *et al.*, 2020).

#### *Accessibility*

Three papers highlighted accessibility as a facilitator to interventions. By using simple terms that could be easily understood by the target population, the process is made easier for the recipient (Sharma *et al.*, 2023). Interventions that were 'short' and 'convenient' were enjoyed by participants (Rudzinski *et al.*, 2012) and participants appreciated the convenience of accessing support for drug use from home (Sharma *et al.*, 2023).

When interventions were flexible in terms of their access and implementation, this was useful as the format could be adjusted as needed to suit the participants needs, such as the addition of in person intervention rather than online only (Sharma *et al.*, 2023). In addition, Whiteside *et al.* (2010) mentioned how the social-norms approach used within one intervention had been effective in various formats including in person, group, mailed and computer-based interventions (Whiteside *et al.*, 2010).

#### *Intervention features*

Specific features within interventions were identified as valuable or useful to participants. Rudzinski *et al.* (2012) felt that minor suggestions, such as delaying drug use, facilitated behaviour change and made it seem possible as there were smaller steps that could be taken to progress (Rudzinski *et al.*, 2012). Goal setting, weighing pros and cons and restricting opportunities for use were identified as useful elements for behaviour change as they encouraged reflection and awareness of their use (Rudzinski *et al.*, 2012; Morris *et al.*, 2024). Interventions focused on the individuals' goals with reflection on goals set in the previous sessions were valued (Morris *et al.*, 2024). Other elements of interventions that were beneficial and appreciated by participants included coping mechanisms and mindfulness (Morris *et al.*, 2024), and increased awareness of their substance use (Whiteside *et al.*, 2010; Rudzinski *et al.*, 2012). Facilitators recognising and reinforcing change talk, the use of harm reduction targets, and addressing ambivalence were also mentioned (Whiteside *et al.*, 2010).

#### *Participant engagement*

Participant engagement was identified as a facilitator to interventions. Rudzinski *et al.* (2012) demonstrated how participants felt the intervention was useful for themselves or others and spoke positively about its informative nature (Rudzinski *et al.*, 2012). Through this information, participants felt the benefit of seeing facts and numbers demonstrating the impact of substance use upon them, which they felt 'really brings it home'. Informing participants of the risk associated with their substance use was mentioned in two papers as potential facilitators to change, increased awareness of risk resulted in reduction or positive changes in substance use, potentially due concerns around the impact on their health (Whiteside *et al.*, 2010; Rudzinski *et al.*, 2012). Curiosity and the use of a participatory approach were highlighted as facilitators to recruitment within a college student sample, while the inclusion of novel information and the opportunity for more detailed responses through open ended questions were facilitators to engagement (Sharma *et al.*, 2023). The interactive elements of interventions were also well received (Morris *et al.*, 2024). Rudzinski *et al.* (2012) highlighted how the use of meaningful tone and language and the inclusion of perspectives of active/previous users may facilitate engagement with the materials and provide examples, this was suggested by several participants.

#### *Criminal justice setting barriers*

Only one paper explored barriers within the criminal justice system and this used the perspective of formerly incarcerated adults, these barriers included internal conflicts, absence

of a problem, seeking informal assistance, poor access, and 'other'.

### **Internal conflicts**

Fear of treatment and privacy concerns were mentioned as potential barriers to those within the criminal justice system. Previous unsuccessful experiences of addiction treatment, reluctance to participate, discomfort with speaking in a group setting, and struggles with privacy may all inhibit likelihood of participation (Owens *et al.*, 2018).

### **Absence of a problem**

Individuals not recognising the need for treatment or ambivalence towards treatment may prevent interventions from being accessed. This includes perceptions that they do not need treatment or can manage without it, or enjoy using the drug and lack the motivation to attend (Owens *et al.*, 2018).

### **Seeking informal/alternative assistance**

Some highlighted that treatment had not been accessed due to the desire to explore informal assistance first, such as family support, and attempt to reduce use through these alternative methods (Owens *et al.*, 2018).

### **Poor access**

Poor access to treatment was highlighted as a barrier, with challenges including time conflict, poor treatment availability, and admission difficulty. Participants highlighted pending court dates, reluctance to take time to attend, challenges with transport and time availability, and the impact of previous offences on being accepted into the treatment as barriers to treatment (Owens *et al.*, 2018).

### **Other**

Three barriers came under the 'other' category that were difficult to interpret, including participants feeling the intervention would not provide them with new information, AA meetings, and the frequent advertisement of alcohol (Owens *et al.*, 2018).

### **Quality Assessment**

Seven of the included studies were classified as low chance of risk, five as medium chance of risks and two as high chance of risk (Extended Data (Table 5 and Table 6).

## **Objective three: qualitative work with practitioners and community members to understand the barriers and facilitators to carrying out brief drug interventions**

### **Patient and Public Involvement**

We engaged with all three Health Determinants Research Centres (HDRCs) to explore potential collaborations and develop initiatives through the HDRCs. Each HDRC either had or was in the process of establishing teams of community researchers, with whom we collaborated during this development year. While no community members participated in this particular phase of the study, we leveraged our extensive experience in community engagement from over 20 prior studies in this field. Drawing on this background, we refined our approach

and will continue to use and adapt these insights throughout the next stages of the study.

For the qualitative element of the study, a combination of semi-structured interviews and focus groups were carried out. Our intention was to carry out interviews/focus groups relating to health/social care and criminal justice (36 with stakeholders and 36 with community members). This would consider three geographical areas of the UK.

The research team spoke to both service users and stakeholders across the key areas to ascertain their thoughts about how the interventions could be developed. Information leaflets were sent to potential participants with an invitation email from the gatekeeper at Health Determinants Research Collaborations (HDRCs) as well as on social media and through our links. Participants were asked to contact the research team to express an interest in taking part in the study. Once an interview or focus group had been arranged, written informed consent was obtained, having ascertained that the participant had read the information leaflet. All focus groups were arranged at a time to suit participants and were conducted face to face. They lasted up to one hour. Interviews similarly lasted up to one hour and were conducted either online or face-to-face. All were digitally audio-recorded with written consent, then transcribed, anonymised, and checked for accuracy.

### **Participants and procedure**

Participants were identified through three newly developed HDRCs who acted as gatekeepers to accessing key individuals within each geographical area and each of the three settings. The sample for the qualitative work was identified through the systematic reviews. These reviews revealed that the most suitable areas for further qualitative investigation included, within the healthcare setting, the Improving Access to Psychological Therapies (IAPT) programme, adult family services for social care settings, and within criminal justice settings, both police and probation services

To be eligible to take part, individuals had to be aged 18 or older, able to speak and understand English, and have the capacity to provide informed consent; and have links with one of the three context areas (health, social care, and/or criminal justice), either as a service user (community member) or a stakeholder. Community member participants were given a £20 gift voucher for taking part. Some of these participants had experience across more than one of the settings studied (e.g., both criminal justice and family services).

### **Ethical considerations and consent**

Ethics approval for this research was granted by the SSSH Research Ethics Sub-Committee at Teesside University on 30/01/2024, under reference number 15992. Prior to commencing the study, all essential documents, including consent forms, information leaflets, interview and focus group questions, and invitation emails, were thoroughly reviewed and approved by the ethics committee of the lead author's institution. The study was conducted in full compliance with the ethical principles

outlined in the Declaration of Helsinki (World Medical Association, 2004), ensuring the highest standards of ethical conduct and participant protection were maintained throughout the research process. Ethical approval included participants giving written consent to the study which they all did.

To minimise risk and ensure participants' understanding of participating, only individuals who had the capacity to consent were approached to take part in the study. The research team was guided by the Mental Capacity Act (2005). Thus, consent was given in written form (either digitally or by hand-written signatures on a consent form) before the focus groups or interviews began.

All personal and identifiable information, such as participants' names and any other identifiable information, was redacted and replaced with a unique participant number. Subsequently, the identifiable information was confidentially destroyed, in keeping with the policy of the lead author's institution. Names mentioned in the audio recordings were redacted from the transcripts and changed to a pseudonym in the write-up.

Participants were informed prior to the interviews or focus groups and on the participant information sheet that, should they disclose anything that posed a risk to themselves or others, or anything of a criminal nature, then confidentiality would need to be breached, the chief investigator would be informed, and then take further action if needed. However, this did not occur throughout the study.

### Data analysis

The data was analysed using thematic analysis, a method that identifies and reports patterns within the data. Based on Braun and Clarke's guidelines (2022), the analysis was reflexive and iterative, involving constant moving back and forth between the dataset, the coded extracts of data, and the analysis of the data (Braun & Clarke, 2006). This approach facilitated a deeper engagement with the data and the identification of themes that accurately reflected participants' experiences. The process of analysis started with a line-by-line examination of the transcripts, during which initial codes were produced. These codes were subsequently collated into potential themes through a process of constant comparison and refinement. Coding was conducted using an inductive approach so that the codes and subsequent themes were developed directly from the data without being influenced by pre-existing theories.

We found that most stakeholders and community members had experience of more than one agency. We carried out interviews/focus groups with 18 community members and 16 stakeholders however this related to 33 inputs for community members (13 mental health; 9 CJS and 11 social services). For practitioners this related to 21 inputs (5 mental health; CJS and 7 for social services).

### Community Member Themes

Key issues community members perceived in substance use and mental health interventions included the need to address trauma, the efficacy of interventions, how stigma and societal

views affect individuals, the importance of support from various institutions, and how community involvement and personal responsibility play a role in the recovery process.

### 1. Trauma and Coping Mechanisms

Participants consistently linked trauma, especially from childhood to the development of addiction. They saw addiction as a method of coping with unresolved trauma. One described, "*most of the lives of people in hardcore addiction are spent in trauma... without them even knowing it*" (C\_0004), highlighting the profound and often unrecognised impact of early adversities on substance use behaviours.

### 2. The Efficacy of Interventions

Community members drew attention to the critical need for interventions that are not just timely but align with the individual's journey and readiness for change. They described how the requirements for support change during the stages of addiction and recovery, from initial prevention efforts to active intervention and ultimately, rehabilitation. As one participant put it, "*I think the intervention is one thing, prevention is another thing we have to look at it in different stages*" (C\_0004), indicating a holistic view of addressing substance use, from early prevention to rehabilitation.

### 3. Stigma and Social Perceptions

Stigma emerged as a significant barrier, with participants expressing how societal perceptions, particularly around gender and addiction, influenced the willingness to seek help. As one stated, "*Right, we lose our kids. We're the pieces of s\*\*t as mothers, and the men don't even get a look at. Well, they're the parent as well. Do you know what I mean? It's not just one-sided.*" (FG31). This shows how, in situations of addiction that affect family dynamics, mothers may face harsher criticism and consequences, while fathers or male partners may not be scrutinised or held accountable to the same extent.

### 4. Institutional Roles and Accessibility

Community members' critiques of institutions highlighted the challenges within systems meant to support recovery. Long waiting times, perceived lack of understanding, and inadequate support were recurrent themes. C\_011 expressed, "*It's really weird, so they actually won't take people that have been sectioned because apparently, you're too high risk, so you sometimes fall through a massive gap, like at the moment I have nothing.*" This theme depicts the challenges community members faced when interacting with institutions meant to support recovery, including issues with the timing and appropriateness of interventions, systemic gaps in support for high-risk individuals, a perceived lack of empathy and understanding, and inconsistencies in the quality of support provided.

### 5. Community, Support, and Personal Agency

Participants' narratives around community engagement, financial strain, and the importance of personal growth evidenced a perceived interaction of external factors and individual agency in the context of addiction. Interviewees spoke of their desire for healthier social connections and the challenges of socioeconomic issues. C\_0015, for example, explained, "*I am looking*

forward to building a more healthier social connection in a new community”, expressing a need for supportive environments conducive to recovery and personal development.

### Stakeholder Themes

Stakeholders emphasised the need for interventions to be brief yet personalised, underlining the importance of trust, effective evaluation, and the specific role of police in crisis interventions. These themes underline the necessity of adapting support strategies to individual needs and systemic challenges, highlighting the complexity of effective intervention in high-stress environments.

#### 1. The Importance of Tailored and Brief Interventions

The significance of designing interventions that are both brief and tailored to meet individual needs was a recurring theme. Stakeholders advocated for brief interventions due to their potential to prompt immediate reflection and action. One stakeholder observed, “If we do something because it makes them think, well, actually I need to go to that. Otherwise, I could commit an offence. I could be arrested or trying to sort myself out now” (SH\_0001), highlighting the efficacy of brief interventions in encouraging individuals to seek help or reconsider their actions promptly. Similarly, SH\_0002 stated, “I would rather have a brief conversation, deliver a brief intervention so somebody is making informed choices...I believe they do [work]”. Furthermore, stakeholders’ emphasis on tailored approaches highlighted the need for personalised care, moving away from generic solutions. As another respondent pointed out, “People need different options that will suit them so that they can better engage and that you can as individuals have the hook that suits their mind and way of working and what makes them tick as opposed to this usual one cap fits all” (SH\_0001), advocating for interventions that align with individual learning styles and preferences, thereby enhancing engagement and effectiveness.

#### 2. Trust and Connection in Brief Interventions

Confidentiality and the complexity of relational work were both discussed by interviewees. Ensuring a safe and private environment was seen to be important, as highlighted by a stakeholder’s caution against rushing interventions: “Don’t offer a brief intervention if you only got 5 minutes because you just never know. [The client] might want to disclose something” (SH\_0002). This underlines the necessity of a trusted space for meaningful interactions. Moreover, the skill of relational work within such constraints was perceived to present challenges, with empathic confrontation in brief sessions demanding advanced skills and a thoughtful approach, as another stakeholder observed: “The only concern I have about brevity...is that empathic confrontation piece is a really advanced clinical skill. I think it’s very hard to get right” (SH\_0006). Thus, in brief interventions, establishing trust and forging deep connections, despite time limitations, were seen to be fundamental for impactful outcomes.

#### 3. Evaluation and Sharing Best Practices

Stakeholders called for interventions to be evidence-based, evaluated, and continuously improved based on feedback and outcomes. Learning from past experiences and sharing best practices among professionals and agencies was seen as essential

for refining intervention strategies. “The issue again goes back to first you understanding what already exists and what’s working. So, if that could be built on and more people do it, that’s gonna help, hasn’t it? Which goes back to that sharing practice” (SH\_0015). This emphasises the value of learning from existing practices and the continuous improvement of services through feedback, underlining the importance of evidence-based interventions and adopting what has been proven effective.

#### 4. The Police’s Role in Brief Interventions

Stakeholders highlighted the police’s crucial role in delivering brief interventions at critical moments. As SH\_0015 remarked, “It’s about how then we can identify and signpost out from the need to actually make an arrest. If you can really avoid that, that doesn’t actually help the situation, but with the view is we’ve got a suite of interventions that firstly goes back to proper assessment.” However, they also pointed out the challenges in aligning these interventions with broader service networks, pointing out hurdles in information sharing and coordination. Additionally, concerns were raised about securing consent and the effectiveness of follow-up for drug and alcohol support within police custody, highlighting areas for improvement in the system.

#### Objective four: to develop a research team with appropriate skills and expertise, including community sector

It was important that the study included academics with expertise and knowledge of the topic area. The research team are international experts in the field of carrying out RCTs in the field, including the SIPS alcohol screening and brief intervention trials (Drummond *et al.*, 2014; Kaner *et al.*, 2013; Newbury-Birch *et al.*, 2014). Further to the original team we have brought in others such as Professor Jeremy Bray and Professor Tom Phillips who bring economics/health economics expertise and nursing/intervention expertise. We have also carried out a lot of work with the community sector and Dr Andrew Divers who is an expert in community research will lead this work in co-production with HDRC-South Tees and Junction 42. We have had a number of meetings with community representatives who were key to our funding application.

#### Objective five: to develop a funding application for a future definitive trial

We have used the information gathered in the work we have done and have submitted a stage one application to NIHR PHR for funding.

#### AIM

To conduct three two-arm feasibility randomised controlled trial to assess the feasibility of developing a definitive trial application for RCTs of adult drug screening and brief interventions in key health, social care and justice settings.

#### Ethics and consent

##### Ethical considerations and consent

Ethics approval for this research was granted by the SSSH Research Ethics Sub-Committee at Teesside University on 30/01/2024, under reference number 15992. Prior to commencing



the study, all essential documents, including consent forms, information leaflets, interview and focus group questions, and invitation emails, were thoroughly reviewed and approved by the ethics committee of the lead author's institution. The study was conducted in full compliance with the ethical principles outlined in the Declaration of Helsinki (World Medical Association, 2004), ensuring the highest standards of ethical conduct and participant protection were maintained throughout the research process. Ethical approval included participants giving written consent to the study which they all did.

### Data (and software) availability

No data associated with this article other than what is available at <https://osf.io/2nqvm/> as this was not requested from the ethical committee. To discuss data that is not available at please contact [d.newbury-birch@tees.ac.uk](mailto:d.newbury-birch@tees.ac.uk)

### Extended data

Materials used in this study, including the interview schedule, participant information sheets, and consent forms can be found in the repository 'Reset Report' at <https://osf.io/2nqvm/> (Tuschick *et al.*, 2025)

**OSF:** Reset Report: *NIHR PHR REPORT DOCUMENTS FOR RESET*

This project contains the following extended data: <https://osf.io/2nqvm/>

### Checklists- sys reviews

- *PRISMA\_2020\_checklist-QUAL*
- *PRISMA\_2020\_checklist-QUANT*

### Qualitative work

- *Consent Form*
- *Debrief*
- *Interview schedule*
- *Community Participant Information Sheet*
- *Stakeholder Participant Information Sheet*

### Tables and figures:

- *Figure 1*
- *Figure 2*
- *Table 1*
- *Table 2*
- *Table 3*
- *Table 4*
- *Table 5*
- *Table 6*

### Details of license:

Data are available under the terms of the Creative Commons Zero "No rights reserved" data waiver (CC0 1.0 Public domain dedication) (<http://creativecommons.org/publicdomain/zero/1.0/>)

### Reporting guidelines

PRISMA Checklist for, "Development of a trial application to assess the effectiveness and cost-effectiveness of adult dRug screening and brief interventionS in key hEalth, social care and justice setTings: The RESET PROJECT ". <https://osf.io/2nqvm/>

## References

- Aharonovich E, Greenstein E, O'Leary A, *et al.*: **HealthCall: technology-based extension of motivational interviewing to reduce non-injection drug use in HIV primary care patients – a pilot study.** *AIDS Care.* 2012; **24**(12): 1461–1469. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Aharonovich E, Sarvet A, Stohl M, *et al.*: **Reducing non-injection drug use in HIV primary care: a randomized trial of brief motivational interviewing, with and without HealthCall, a technology-based enhancement.** *J Subst Abuse Treat.* 2017; **74**: 71–79. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Assanangkornchai S, Nima P, McNeil EB, *et al.*: **Comparative trial of the WHO ASSIST-linked brief intervention and simple advice for substance abuse in primary care.** *Asian J Psychiatr.* 2015; **18**: 75–80. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bagøien G, Bjørngaard JH, Østensen C, *et al.*: **The effects of motivational interviewing on patients with comorbid substance use admitted to a psychiatric emergency unit—A randomised controlled trial with two year follow-up.** *BMC Psychiatry.* 2013; **13**: 93. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Baker A, Lee NK, Claire M, *et al.*: **Brief cognitive behavioural interventions for regular amphetamine users: a step in the right direction.** *Addiction.* 2005; **100**(3): 367–378. [PubMed Abstract](#) | [Publisher Full Text](#)
- Barrowclough C, Haddock G, Wykes T, *et al.*: **Integrated motivational interviewing and cognitive behavioural therapy for people with psychosis and comorbid substance misuse: randomised controlled trial.** *BMJ.* 2010; **341**: c6325. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Black C: **Review of drugs (Review of Drugs Executive Summary).** London, 2020. [Reference Source](#)
- Blow FC, Walton MA, Bohnert ASB, *et al.*: **A randomized controlled trial of brief interventions to reduce drug use among adults in a low-income urban emergency department: the HealthIER You study.** *Addiction.* 2017; **112**(8): 1395–1405. [PubMed Abstract](#) | [Publisher Full Text](#)
- Boden MT, Kimerling R, Jacobs-Lentz J, *et al.*: **Seeking safety treatment for male veterans with a substance use disorder and Post-Traumatic Stress Disorder symptomatology.** *Addiction.* 2012; **107**(3): 578–586. [PubMed Abstract](#) | [Publisher Full Text](#)
- Braun V, Clarke V: **Using thematic analysis in Psychology.** *Qualitative Research Psychology.* 2006; **3**(2): 77–101. [Publisher Full Text](#)
- Carroll KM, Martino S, Ball SA, *et al.*: **A multisite randomized effectiveness trial of motivational enhancement therapy for Spanish-speaking substance users.** *J Consult Clin Psychol.* 2009; **77**(5): 993–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)



CASP-UK: **Critical Appraisal Skills Programme (CASP)**. London Oxford, 2002. [Reference Source](#)

Cooke A, Smith D, Booth A: **Beyond PICO: the SPIDER tool for qualitative evidence synthesis**. *Qual Health Res*. 2012; **22**(10): 1435–1443. [PubMed Abstract](#) | [Publisher Full Text](#)

Darker C, Sweeney B, Keenan E, *et al.*: **Tailoring a brief intervention for illicit drug use and alcohol use in Irish methadone maintained opiate dependent patients: a qualitative process**. *BMC Psychiatry*. 2016; **16**(1): 373. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

de Oliveira Christoff A, Boerngen-Lacerda R: **Reducing substance involvement in college students: a three-arm parallel-group randomized controlled trial of a computer-based intervention**. *Addict Behav*. 2015; **45**: 164–171. [PubMed Abstract](#) | [Publisher Full Text](#)

Drummond C, Deluca P, Coulton S, *et al.*: **The effectiveness of alcohol screening and brief intervention in emergency departments: a multicentre pragmatic cluster randomized controlled trial**. *PLoS One*. 2014; **9**(6): e99463. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Drummond C, James D, Coulton S, *et al.*: **The effectiveness and cost-effectiveness of screening and stepped-care interventions for alcohol use disorders in the primary care setting**. Final report to the Welsh Office for Research and Development. Cardiff, Wales Office for Research and Development, 2004.

Dujon CS: **Screening for alcohol and substance misuse in BB: how numerous factors affect participation**. Ph.D., Northcentral University, 2021. [Reference Source](#)

Fazio D, Zuiderveen S, Guyet D, *et al.*: **ED-Home: pilot feasibility study of a targeted homelessness prevention intervention for emergency department patients with drug or unhealthy alcohol use**. *Acad Emerg Med*. 2022; **29**(12): 1453–1465. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Fischer B, Dawe M, McGuire F, *et al.*: **Feasibility and impact of brief interventions for frequent cannabis users in Canada**. *J Subst Abuse Treat*. 2013; **44**(1): 132–138. [PubMed Abstract](#) | [Publisher Full Text](#)

Forsberg LG, Ernst D, Sundqvist K, *et al.*: **Motivational interviewing delivered by existing prison staff: a randomized controlled study of effectiveness on substance use after release**. *Subst Use Misuse*. 2011; **46**(12): 1477–1485. [PubMed Abstract](#) | [Publisher Full Text](#)

Gates PJ, Norberg MM, Copeland J, *et al.*: **Randomized controlled trial of a novel cannabis use intervention delivered by telephone**. *Addiction*. 2012; **107**(12): 2149–2158. [PubMed Abstract](#) | [Publisher Full Text](#)

Gaume JVS, Grazioli S, Paroz C, *et al.*: **Developing a brief motivational intervention for young adults admitted with alcohol intoxication in the emergency department - Results from an iterative qualitative design**. *PLoS One*. 2021; **16**(2): e0246652. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Gelberg L, Andersen RM, Afifi AA, *et al.*: **Project QUIT (Quit Using Drugs Intervention Trial): a randomized controlled trial of a primary care-based multi-component brief intervention to reduce risky drug use**. *Addiction*. 2015; **110**(11): 1777–1790. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Gelberg L, Andersen RM, Rico MW, *et al.*: **A pilot replication of QUIT, a randomized controlled trial of a brief intervention for reducing risky drug use, among Latino primary care patients**. *Drug Alcohol Depend*. 2017; **179**: 433–440. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Gmel G, Gaume J, Bertholet N, *et al.*: **Effectiveness of a brief integrative multiple substance use intervention among young men with and without booster sessions**. *J Subst Abuse Treat*. 2013; **44**(2): 231–240. [PubMed Abstract](#) | [Publisher Full Text](#)

Goodness TM, Palfai TP: **Electronic screening and brief intervention to reduce cannabis use and consequences among graduate students presenting to a student health center: a pilot study**. *Addict Behav*. 2020; **106**: 106362. [PubMed Abstract](#) | [Publisher Full Text](#)

Graham HL, Copello A, Griffith E, *et al.*: **Pilot randomised trial of a brief intervention for comorbid substance misuse in psychiatric in-patient settings**. *Acta Psychiatr Scand*. 2016; **133**(4): 298–309. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Graham ID, Tetroe J: **How to translate health research knowledge into effective healthcare action**. *Healthcare Q*. 2007; **10**(3): 20–22. [PubMed Abstract](#) | [Publisher Full Text](#)

Guan W, Liu T, Baird JR, *et al.*: **Evaluation of a brief intervention to reduce the negative consequences of drug misuse among adult emergency department patients**. *Drug Alcohol Depend*. 2015; **157**: 44–53. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Haddaway NR, Collins AM, Coughlin D, *et al.*: **The role of google scholar in evidence reviews and its applicability to grey literature searching**. *PLoS One*. 2015; **10**(9): e0138237. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Halladay J, Scherer J, MacKillop J, *et al.*: **Brief interventions for cannabis use in emerging adults: a systematic review, meta-analysis, and evidence map**.

*Drug Alcohol Depend*. 2019; **204**: 107565.

[PubMed Abstract](#) | [Publisher Full Text](#)

Heather N, Bowie A, Ashton H, *et al.*: **Randomised controlled trial of two brief interventions against long-term benzodiazepine use: outcome of intervention**. *Addiction Research & Theory*. 2004; **12**(2): 141–154. [Publisher Full Text](#)

Hoch E, Bühringer G, Pixa A, *et al.*: **CANDIS treatment program for cannabis use disorders: findings from a randomized multi-site translational trial**. *Drug Alcohol Depend*. 2014; **134**(1): 185–193. [PubMed Abstract](#) | [Publisher Full Text](#)

Hoffman T, Glasziou P, Boutron I, *et al.*: **Better reporting of interventions: Template for Intervention Description and Replication (TIDieR) checklist and guide**. *BMJ*. 2014; **348**: g1687. [PubMed Abstract](#) | [Publisher Full Text](#)

Humeniuk R, Henry-Edward S, Ali R, *et al.*: **The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): manual for use in primary care**. Geneva, World Health Organization, 2010. [Reference Source](#)

Humeniuk R, Newcombe D, Dennington V, *et al.*: **A randomised controlled trial of a brief intervention for illicit drug use linked to ASSIST screening in a primary healthcare setting: results from the Australian component of the World Health Organization Phase III ASSIST studies**. *Aust J Prim Health*. 2018; **24**(2): 149–154. [PubMed Abstract](#) | [Publisher Full Text](#)

Hunter SB, Ober AJ, McCullough CM, *et al.*: **Sustaining alcohol and opioid use disorder treatment in primary care: a mixed methods study**. *Implement Sci*. 2018; **13**(1): 83. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Institute of Alcohol Studies: **Economy**. 2024; Retrieved 22/05/24. [Reference Source](#)

Jonas B, Tossmann P, Tensil M, *et al.*: **Effekte einer einmaligen Chat-Intervention auf problematischen Substanzkonsum = Efficacy of a single-session online-intervention on problematic substance use**. *Sucht: Zeitschrift für Wissenschaft und Praxis*. 2012; **58**(3): 173–182. [Publisher Full Text](#)

Jungerman FS, Andreoni S, Laranjeira R: **Short term impact of same intensity but different duration interventions for cannabis users**. *Drug Alcohol Depend*. 2007; **90**(2–3): 120–127. [PubMed Abstract](#) | [Publisher Full Text](#)

Kaner EFS, Beyer FR, Muirhead C, *et al.*: **Effectiveness of brief alcohol interventions in primary care populations**. *Cochrane Database Syst Rev*. 2018; **2**(2): CD004148. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kaner E, Bland M, Cassidy P, *et al.*: **Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): pragmatic cluster randomised controlled trial**. *BMJ*. 2013; **346**: e8501. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kavanagh DJ, Young R, White A, *et al.*: **A brief motivational intervention for substance misuse in recent-onset psychosis**. *Drug Alcohol Rev*. 2004; **23**(2): 151–155. [PubMed Abstract](#) | [Publisher Full Text](#)

Kim TW, Bernstein J, Cheng DM, *et al.*: **Receipt of addiction treatment as a consequence of a brief intervention for drug use in primary care: a randomized trial**. *Addiction*. 2017; **112**(5): 818–827. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lerch J, Walters ST, Tang L, *et al.*: **Effectiveness of a computerized Motivational Intervention on treatment initiation and substance use: results from a randomized trial**. *J Subst Abuse Treat*. 2017; **80**: 59–66. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Light M, Grant E, Hopkins K: **Gender differences in substance misuse and mental health amongst prisoners: results from the Surveying Prisoner Crime Reduction (SPCR) longitudinal cohort study of prisoners**. London, Ministry of Justice Analytical Services, 2013. [Reference Source](#)

McKee SA, Carroll KM, Sinha R, *et al.*: **Enhancing brief Cognitive-Behavioral Therapy with motivational enhancement techniques in cocaine users**. *Drug Alcohol Depend*. 2007; **91**(1): 97–101. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

McQueen J, Howe T, Allan L, *et al.*: **Brief interventions for heavy alcohol users admitted to general hospital wards**. *Cochrane Database Syst Rev*. 2009; **(3)**: CD005191. [PubMed Abstract](#) | [Publisher Full Text](#)

Mental Capacity Act: **Mental capacity act 2005**. London Station Office, 2005.

Merchant RC, Zhang Z, Zhang Z, *et al.*: **Lack of efficacy in a randomised trial of a Brief Intervention to reduce drug use and increase drug treatment services utilisation among adult Emergency Department patients over a 12-month period**. *Emerg Med J*. 2018; **35**(5): 282–288. [PubMed Abstract](#) | [Publisher Full Text](#)

Mertens JR, Ward CL, Bresick GF, *et al.*: **Effectiveness of nurse-practitioner-delivered brief Motivational Intervention for young adult alcohol and drug use in primary care in South Africa: a randomized clinical trial**. *Alcohol Alcohol*. 2014; **49**(4): 430–438. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

- Miller W, Rollnick S: **Motivational interviewing; preparing people to change addictive behavior.** New York, Guildford Press, 1991.  
[Reference Source](#)
- Mitcheson L, McCambridge J, Byrne S: **Pilot cluster-randomised trial of adjunctive motivational interviewing to reduce crack cocaine use in clients on methadone maintenance.** *Eur Addict Res.* 2006; **13**(1): 6–10.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Morris SL, Langwerden RJ, Wagner EF, *et al.*: **Implementation of a brief motivational intervention for alcohol and other drug using Latinx college students.** *J Am Coll Health.* 2024; **72**(6): 1785–1795.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nagel T, Robinson G, Condon J, *et al.*: **Approach to treatment of mental illness and substance dependence in remote indigenous communities: results of a mixed methods study.** *Aust J Rural Health.* 2009; **17**(4): 174–182.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Newbury-Birch D, Coulton S, Bland M, *et al.*: **Alcohol screening and brief interventions for offenders in the probation setting (sips trial): a pragmatic multicentre cluster randomised controlled trial.** *Alcohol Alcohol.* 2014; **49**(5): 540–548.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Newbury-Birch D, Ferguson J, Connor N, *et al.*: **A rapid systematic review of worldwide alcohol use disorders and brief alcohol interventions in the criminal justice system.** *Front Psychiatry.* 2022; **13**: 900186.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Newbury-Birch D, Ferguson J, Landale S, *et al.*: **A systematic review of the efficacy of alcohol interventions for incarcerated people.** *Alcohol Alcohol.* 2018; **53**(4): 412–425.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Newbury-Birch D, McGovern R, Birch J, *et al.*: **A rapid systematic review of what we know about alcohol use disorders and brief interventions in the criminal justice system.** *Int J Prison Health.* 2016; **12**(1): 57–70.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- O'Donnell A, Kaner E, Newbury-Birch D, *et al.*: **The impact of brief alcohol interventions in primary healthcare: a systematic review of reviews.** *Alcohol Alcohol.* 2014; **49**(1): 66–78.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Office for National Statistics: **Drug misuse in England and Wales: year ending March 2023.** London, 2023.  
[Reference Source](#)
- Owens MD, Chen JA, Simpson TL, *et al.*: **Barriers to addiction treatment among formerly incarcerated adults with substance use disorders.** *Addict Sci Clin Pract.* 2018; **13**(1): 19.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Palfai TP, Saitz R, Winter M, *et al.*: **Web-based screening and brief intervention for student marijuana use in a university health center: pilot study to examine the implementation of eCHECKUP TO GO in different contexts.** *Addict Behav.* 2014; **39**(9): 1346–1352.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Poblete F, Barticevic NA, Zuzulich MS, *et al.*: **A randomized controlled trial of a brief intervention for alcohol and drugs linked to the Alcohol, Smoking and Substance Involvement Screening Test (ASSIST) in primary health care in Chile.** *Addiction.* 2017; **112**(8): 1462–1469.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Prendergast ML, McCollister K, Warda U: **A randomized study of the use of Screening, Brief Intervention, and Referral to Treatment (SBIRT) for drug and alcohol use with jail inmates.** *J Subst Abuse Treat.* 2017; **74**: 54–64.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Raistrick D, Heather N, Godfrey C: **Review of the effectiveness of treatment for alcohol problems.** *National Treatment Agency for Substance Misuse UK.* 2006.  
[Reference Source](#)
- Rethlefsen ML, Kirtley S, Waffenschmidt S, *et al.*: **PRISMA-S: an extension to the PRISMA statement for reporting literature searches in systematic reviews.** *Syst Rev.* 2021; **10**(1): 39.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Roberts SCM, Nuru-Jeter A: **Women's perspectives on screening for alcohol and drug use in prenatal care.** *Womens Health Issues.* 2010; **20**(3): 193–200.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Roy-Byrne P, Bumgardner K, Krupski A, *et al.*: **Brief intervention for problem drug use in safety-net primary care settings: a randomized clinical trial.** *JAMA.* 2014; **312**(5): 492–501.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rudzinski K, McGuire F, Dawe M, *et al.*: **Brief intervention experiences of young high-frequency Cannabis users in a Canadian setting.** *Contemp Drug Probl.* 2012; **39**(1): 49–72.  
[Publisher Full Text](#)
- Satre DD, Leibowitz A, Sterling SA, *et al.*: **A randomized clinical trial of motivational interviewing to reduce alcohol and drug use among patients with depression.** *J Consult Clin Psychol.* 2016; **84**(7): 571–579.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Saunders EC, Moore SK, Gardner T, *et al.*: **Screening for substance use in rural primary care: a qualitative study of providers and patients.** *J Gen Intern Med.* 2019; **34**(12): 2824–2832.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sharma K, Ghosh A, Krishnan NC, *et al.*: **Digital screening and brief intervention for illicit drug misuse in college students: a mixed methods, pilot, cluster, randomized trial from India.** *Asian J Psychiatr.* 2023; **81**: 103432.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shekhawat AS, Mathur R, Sarkar S, *et al.*: **A randomized controlled trial of brief intervention for patients with Cannabis use disorder.** *J Neurosci Rural Pract.* 2023; **14**(4): 710–716.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Shepherd J: **How to achieve more effective services: the evidence ecosystem.** Cardiff, Cardiff University, 2014.  
[Reference Source](#)
- Sherman LW, Strang H, Barnes G, *et al.*: **Twelve experiments in restorative justice: the Jerry Lee program of randomized trials of restorative justice conferences.** *J Exp Criminol.* 2015; **11**(4): 501–540.  
[Publisher Full Text](#)
- Sobell LC, Sobell MB, Agrawal S: **Randomized controlled trial of a cognitive-behavioral motivational intervention in a group versus individual format for substance use disorders.** *Psychol Addict Behav.* 2009; **23**(4): 672–683.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sorsdahl K, Stein DJ, Pasche S, *et al.*: **A novel brief treatment for methamphetamine use disorders in South Africa: a randomised feasibility trial.** *Addict Sci Clin Pract.* 2021; **16**(1): 3.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Starks TJ, Robles G, Doyle KM, *et al.*: **Motivational Interviewing with male couples to reduce substance use and HIV risk: manifestations of partner discord and strategies for facilitating dyadic functioning.** *Psychotherapy (Chic).* 2020; **57**(1): 58–67.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stephens RS, Roffman RA, Fearer SA, *et al.*: **The Marijuana check-up: reaching users who are ambivalent about change.** *Addiction.* 2004; **99**(10): 1323–1332.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tanner-Smith EE, Parr NJ, Schweer-Collins M, *et al.*: **Effects of brief substance use interventions delivered in general medical settings: a systematic review and meta-analysis.** *Addiction.* 2022; **117**(4): 877–889.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Thompson RG, Aivadyan C, Stohl M, *et al.*: **Smartphone application plus brief motivational intervention reduces substance use and sexual risk behaviors among homeless young adults: results from a randomized controlled trial.** *Psychol Addict Behav.* 2020; **34**(6): 641–649.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tuschick E, Ferguson J, Coulton S, *et al.*: **RESET REPORT. [DATA SET].** London, OSFHOME, 2025.
- United Nations on Drugs and Crime: **World drug report 2023.** UONCD Research, 2023.  
[Reference Source](#)
- van Emmerik-van Oortmerssen K, Vedel E, Kramer FJ, *et al.*: **Integrated cognitive behavioral therapy for ADHD in adult substance use disorder patients: results of a randomized clinical trial.** *Drug Alcohol Depend.* 2019; **197**: 28–36.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Venner KL, Sánchez V, García J, *et al.*: **Moving away from the tip of the pyramid: screening and Brief Intervention for risky alcohol and opioid use in underserved patients.** *J Am Board Fam Med.* 2018; **31**(2): 243–251.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wehrens R: **Beyond two communities – from research utilization and knowledge translation to co-production?** *Public Health.* 2014; **128**(6): 545–551.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wernett GT, Plegue M, Kahler CW, *et al.*: **A pilot randomized controlled trial of a computer-delivered Brief Intervention for substance use and risky sex during pregnancy.** *J Womens Health (Larchmt).* 2018; **27**(1): 83–92.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Whiteside U, Cronce JM, Pedersen ER, *et al.*: **Brief motivational feedback for college students and adolescents: a harm reduction approach.** *J Clin Psychol.* 2010; **66**(2): 150–163.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Woodruff SI, Clapp JD, Eisenberg K, *et al.*: **Randomized clinical trial of the effects of screening and Brief Intervention for illicit drug use: the life shift/shift gears study.** *Addict Sci Clin Pract.* 2014; **9**(1): 8.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Woolard R, Baird J, Longabaugh R, *et al.*: **Project reduce: reducing alcohol and marijuana misuse: effects of a Brief Intervention in the Emergency Department.** *Addict Behav.* 2013; **38**(3): 1732–1739.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- World Medical Association: **Declaration of Helsinki: World Medical Association. Ethical principles for medical research on human beings.** 2004; **64**: 2013.  
[Reference Source](#)
- Zahradnik A, Otto C, Crackau B, *et al.*: **Randomized controlled trial of a Brief Intervention for problematic Prescription Drug use in non-treatment-seeking patients.** *Addiction.* 2009; **104**(1): 109–117.  
[PubMed Abstract](#) | [Publisher Full Text](#)

# Open Peer Review

Current Peer Review Status: 

---

## Version 1

Reviewer Report 09 June 2025

<https://doi.org/10.3310/nihropenres.14803.r35827>

© 2025 Tanner-Smith E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Emily E. Tanner-Smith** 

University of Oregon, Eugene, Oregon, USA

This manuscript reports findings from two systematic reviews synthesizing evidence about the effectiveness, barriers/facilitators, screening tools, and core components of brief drug interventions, along with findings from a qualitative interview study exploring interest-holders' perspectives on barriers/facilitators of brief drug intervention implementation. The three projects each used rigorous and transparent methods, resulting in a comprehensive summary of current best available evidence that informed the development of an application for a new trial examining the (cost)effectiveness of targeted drug screening and brief interventions in diverse settings in the UK. The three projects followed pre-registered protocols, and the manuscript provides links to supplemental project materials to further ensure replicability. Below are minor comments that might be considered to further strengthen reproducibility and utility.

1. Given that the first objective involves a systematic review of quantitative data related to brief drug interventions, but does not involve a quantitative meta-analysis, it might be useful to adhere to the SWiM reporting guidelines (<https://www.bmj.com/content/368/bmj.l6890>) in addition to the PRISMA reporting guidelines. Specifically, it would be useful to see more transparent discussion around the standardised metric(s) used and the synthesis methods used for summarizing evidence. For instance, it appears that synthesis was conducted using vote counting methods (which is acceptable for addressing questions about directions of effects, but which does not focus on variability or distribution of effects), although the exact approach is not stated explicitly.
2. For the second objective of conducting a systematic review of qualitative data related to brief drug interventions, it would be useful to see more detail regarding the methods for implementing the thematic synthesis. This might be achieved through the application of the ENTREQ reporting guidelines (<https://pmc.ncbi.nlm.nih.gov/articles/PMC3552766/>), specifically providing more detail on the appraisal process, study comparison methods, and derivation of themes.
3. Given that numerous prior systematic reviews have been conducted on this topic (which are cited in the manuscript), and given the explicit focus on the UK context, it could be useful to see objectives 1 and 2 provide supplementary information focused solely on the evidence available from the UK. Although the manuscript transparently reports the country for each included study, no separate syntheses are conducted for UK-based studies. Such subgroup analyses may be quite useful for further underscoring the current gaps in available evidence, particularly for health,

social care, and justice settings in the UK.

4. I would encourage the authors to consider using the more inclusive term of "interest-holder" throughout (REFER 1).

### References

1. Akl E, Khabisa J, Petkovic J, Magwood O, et al.: "Interest-holders": A new term to replace "stakeholders" in the context of health research and policy. *Cochrane Evidence Synthesis and Methods*. 2024; **2** (11). [Publisher Full Text](#)

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Evidence synthesis methods, evaluation methods, substance use prevention

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---