

Sabti, MA and Shamsaldeen, YA

Correcting hypophosphataemia in a paediatric patient with Sanjad-Sakati syndrome through a single oral dose of potassium phosphate intravenous solution

<https://researchonline.ljmu.ac.uk/id/eprint/26773/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)


Sabti, MA and Shamsaldeen, YA (2021) Correcting hypophosphataemia in a paediatric patient with Sanjad-Sakati syndrome through a single oral dose of potassium phosphate intravenous solution. SAGE Open Medical Case Reports. 9. pp. 1-4. ISSN 2050-313X

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

Correcting hypophosphataemia in a paediatric patient with Sanjad–Sakati syndrome through a single oral dose of potassium phosphate intravenous solution

SAGE Open Medical Case Reports
Volume 9: 1–4
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2050313X20988412
journals.sagepub.com/home/sco


Mnaff A Sabti¹ and Yousif A Shamsaldeen^{2,3} 

Abstract

Sanjad–Sakati syndrome is an autosomal recessive disorder that is quite common in Kuwait. Among a wide range of complications in Sanjad–Sakati syndrome patients is the vulnerability to infections and subsequent hypophosphataemia. Hypophosphataemia is a metabolic alteration that contributes to numerous consequences such as cardiac arrhythmia. Therefore, if hypophosphataemia is left unresolved, it may culminate in death. A 20-month-old boy of 2.5 kg body weight diagnosed with Sanjad–Sakati syndrome was initially admitted to the paediatric intensive care unit after recovering from COVID-19, and then shifted to the general ward. He was diagnosed with recurrent pneumonia and urinary tract infection. After 9 days, the patient showed severe hypophosphataemia with serum phosphate concentration reaching 0.33 mmol/L. Despite the availability of potassium phosphate intravenous solution, it was difficult to administer potassium phosphate intravenously because of the small body size and low body weight of the patient. Therefore, 0.6 mL potassium phosphate containing 2.4 mEq of potassium and 5.3 mEq of phosphate was administered through a nasogastric tube. The patient showed rapid response after a single dose through the nasogastric tube. Such an intervention in Sanjad–Sakati syndrome patients shows possible advantages of shifting drug administration from intravenous to oral route that includes a convenient route of administration, whether in the intensive care unit or in the general ward. Moreover, shifting drug administration from the intravenous to oral route overcomes the risk of cannula-induced infection and reduces nurses' workload.

Keywords

Sanjad–Sakati syndrome, hypophosphataemia, paediatrics

Date received: 12 October 2020; accepted: 28 December 2020

Introduction

Sanjad–Sakati syndrome is an autosomal recessive disorder, originating mainly in the Middle East and specifically the Arabian Gulf countries, where it was first reported in 1988.¹ In Kuwait, the estimated prevalence of Sanjad–Sakati syndrome is 7–18 per 100,000 live births.² Sanjad–Sakati syndrome starts during early stages of embryogenesis and is characterised by intrauterine growth retardation. Children with Sanjad–Sakati syndrome present with hypocalcaemia tetany and seizures due to hypoparathyroidism at an early stage in their lives that is associated with hyperphosphataemia.^{1,3} Furthermore, Sanjad–Sakati syndrome patients are prone to infections such as pneumonia due to their compromised immunity.^{2,3} Therefore, despite the characteristic feature of hyperphosphataemia in Sanjad–Sakati syndrome patients,

their vulnerability towards infection may lead to severe hypophosphataemia.⁴ Hypophosphataemia is a metabolic alteration that contributes to a wide range of consequences such as suppressing the production of cellular energy adenosine triphosphate (ATP).⁵ Physiological phosphate level ranges from 0.8 to 1.3 mmol/L. However, hypophosphataemia

¹Paediatric Department, Al-Adan Hospital, Ahmadi Medical Governorate, Hadiya, Kuwait

²Kuwait-Al-Adan Joint Hospital, Al-Adan Paediatric Pharmacy, Kuwait Hospital, Sabah Alsalem, Kuwait

³Department of Pharmacy, Kuwait Hospital, Sabah Alsalem, Kuwait

Corresponding Author:

Yousif A Shamsaldeen, Department of Pharmacy, Kuwait Hospital, Mohammad Tahoos Nasser bin Tahoos Street, Sabah Alsalem 44001, Kuwait.
Email: yahmad@kuwaithospital.com.kw



Table 1. Patient's electrolyte concentration upon admission.

| Electrolyte | Concentration (mmol/L) | Normal range (mmol/L) |
|-------------|------------------------|-----------------------|
| Phosphate | 0.87 | 1–1.65 |
| Magnesium | 0.82 | 0.75–0.95 |
| Calcium | 2.19 | 2.21–2.61 |
| Sodium | 152 | 135–150 |
| Potassium | 3.3 | 3.5–5 |
| Chloride | 123 | 98–110 |

is defined as serum phosphate concentration below 0.8 mmol/L.⁵ According to recent studies, the prevalence of hypophosphataemia in the paediatric intensive care unit (PICU) is approximately 50%.⁶ Symptoms of hypophosphataemia are of a wide range and include cardiac arrhythmia, fatigue, irritability and reduced diaphragm contractility.⁶ Such a wide range of symptoms renders hypophosphataemia less diagnosable and hence may lead to silent and progressive complications, which, if left unresolved, may culminate in death.⁷

Case report

Here, we report a 20-month-old boy of 2.5 kg body weight diagnosed with Sanjad–Sakati syndrome alongside other comorbidities and complications. These included epilepsy that is controlled through phenobarbitone syrup 5 mg twice daily (4 mg/kg/day), gastro-oesophageal reflux disease that is controlled through domperidone 1 mg twice daily (0.8 mg/kg/day) and omeprazole 5 mg once daily (2 mg/kg/day), and congenital heart disease (atrial septum defect, ventricular septum defect, and patent ductus arteriosus) that is managed with furosemide 2 mg twice a day (1.6 mg/kg/day).

The patient recovered from COVID-19 at the Jaber Hospital and then was transferred to Al-Adan-Kuwait Hospital, where he was initially admitted to the PICU because of recurrent pneumonia and urinary tract infection (UTI), which were treated through amikacin administered intramuscularly 37.5 mg once daily (15 mg/kg/day). The main challenge of controlling our patient's condition was in the inability to administer drugs through intravenous route due to his compromised body growth. Therefore, we administered injectable drugs through the intramuscular route and oral drugs through a nasogastric tube (NGT). Upon his admission to the PICU, the initial serum electrolyte concentrations were as shown in Table 1.

Based on the electrolyte levels at admission, and since potassium concentration was low, the patient was administered 2.5 mL of potassium chloride (1 M) containing 2.4 mEq potassium once daily by NGT. The patient was stable and hence was moved to the general ward within 24 h of admission. After 9 days, the patient showed severe hypophosphataemia with serum phosphate concentration reaching 0.33 mmol/L, while the concentrations of other electrolytes were normal, as shown in Table 2.

Table 2. Patient's electrolytes concentration 9 days after admission.

| Electrolyte | Concentration (mmol/L) | Normal range (mmol/L) |
|-------------|------------------------|-----------------------|
| Phosphate | 0.33 | 1–1.65 |
| Magnesium | 0.93 | 0.75–0.95 |
| Calcium | 2.35 | 2.21–2.61 |
| Sodium | 132 | 135–150 |
| Potassium | 3.8 | 3.5–5 |
| Chloride | 89 | 98–110 |

Table 3. Patient's electrolytes 15 h after administering potassium phosphate.

| Electrolyte | Concentration (mmol/L) | Normal range (mmol/L) |
|-------------|------------------------|-----------------------|
| Phosphate | 1.28 | 1–1.65 |
| Magnesium | 0.86 | 0.75–0.95 |
| Calcium | 2.08 | 2.21–2.61 |
| Sodium | 132 | 135–150 |
| Potassium | 5.5 | 3.5–5 |
| Chloride | 93 | 98–110 |

Accordingly, it was essential to correct phosphate levels. However, as mentioned, the main obstacle towards such a vital step was the inability to administer potassium phosphate intravenously. In discussing possible alternatives with a hospital pharmacist, he suggested administering potassium phosphate through the NGT route, rather than the designated intravenous route of administration. Such a suggestion was based on calculating the milliequivalents (mEq) of potassium in potassium chloride (KCl) solution and calculating the volume of potassium phosphate based on mEq of potassium in potassium phosphate.

Since KCl (1 M) is of 2.4 mEq potassium, we aimed to provide potassium phosphate through delivering 2.4 mEq potassium. After detailed calculations, it was concluded that administering 0.6 mL from potassium phosphate sterile solution for injection that contains 170.3 mg/mL potassium and 285 mg/mL phosphate provides 2.4 mEq of potassium and 5.3 mEq of phosphate. Therefore, 0.6 mL of potassium phosphate sterile solution was administered through NGT to treat such a case of acute severe hypophosphataemia. 15 h after administering 0.6 mL potassium phosphate through NGT, serum phosphate concentration was normalised as it reached 1.28 mmol/L, while other serum electrolytes concentrations were preserved at normal levels as shown in Table 3.

Discussion

Sanjad–Sakati syndrome is a rare autosomal recessive disorder that exists in the Gulf area. Therapeutic strategies mainly focus on correcting electrolytes and treating infections.¹ Hypophosphataemia is a common complication of infection.

Although hypophosphataemia may be attributed to the use of furosemide or omeprazole, if not both, our patient is using both medicines chronically and has been admitted with normal phosphate concentration (Table 1). Moreover, a previous study revealed that the prevalence of hypophosphataemia in septic patients is approximately 80%.⁴ The percentage of paediatric patients suffering from pneumonia-induced hypophosphataemia is approximately 45%, and children with UTI contribute to approximately 18%.⁸ Since our patient was diagnosed with both pneumonia and UTI, the observed hypophosphataemia is likely attributed to the infections he was suffering from. Such a metabolic complication, if left untreated, may lead to further complications, including cardiac arrhythmia, which is considered a lethal prognosis for our patient, who is also diagnosed with congenital heart disease.

The restricted and limited ability to administer drugs intravenously is considered a major obstacle in managing infection-induced hypophosphataemia in Sanjad–Sakati syndrome. Such a pathological factor that limits the intravenous accessibility of medicines can be solved through changing the route of administration. Converting the route of administration is a challenging skill that requires careful calculations and considerations to provide the optimal benefit from the administered medicine. The challenges of converting the route of administration stem from the pharmacokinetic and pharmacodynamic differences. Thus, the oral route of administration exerts numerous barriers that contribute to reducing drug bioavailability, whereas these barriers are overcome through the intravenous route in which bioavailability is 100%. These barriers include the rate and percentage of absorption as well as intestinal and hepatic metabolism (first-pass effect).⁹ The conversion from intravenous to the oral route is classified mainly into three categories: (1) sequential therapy, where a drug is replaced with its oral counterpart, such as administering esomeprazole (20–40 mg) orally instead of intravenously;⁹ (2) switch therapy, where an intravenously administered drug is replaced with an orally administered drug that belongs to the same pharmacological class and is expected to exert similar therapeutic outcome, such as oral administration of cefuroxime (500 mg) twice daily rather than intravenous administration of ceftriaxone (1 g) once daily;¹⁰ and (3) step-down therapy, through which an injectable drug is replaced with an oral agent that belongs to a different pharmacological class of different frequency, dose and the spectrum of activity. A common example of step-down therapy is the shifting from injectable heparin to orally administered warfarin.¹¹

Here, we have shown an example of sequential therapy through the administration of potassium phosphate through NGT rather than intravenously. This approach provided rapid correction of serum phosphate concentration. Such a therapeutic intervention provides numerous advantages. Mainly, it reduces the risk of cannula-related infections. Thus, shifting the administration of potassium phosphate from the intravenous to oral route may provide additional

benefit through reducing cannula-related infections, hence reducing the need for any additional use of antibiotics, especially in Sanjad–Sakati patients.⁹ Moreover, switching the route of administration reduces the workload on nurses as shown in a previous study which revealed that an intravenous administration of antibiotics takes approximately 10 min and therefore, switching to oral route reduces the nurses' workload by approximately 350 h/year.¹² In addition, we have shown the importance of hospital pharmacists as part of a multidisciplinary team, providing important and vital recommendations to provide optimal healthcare service with the available therapeutic resources.¹³

Conclusion

Sanjad–Sakati syndrome patients are prone to recurrent infections, such as pneumonia and UTI, which contribute to severe hypophosphataemia. Correcting hypophosphataemia in those patients is challenging since intravenous cannulation is extremely difficult because of small body size and low body weight. Therefore, shifting the route of administration from intravenous to oral provides a prompt and essential solution for severe hypophosphataemia in Sanjad–Sakati syndrome patients.

Acknowledgements

The authors acknowledge the Kuwait Hospital management for their support and cooperation during the COVID-19 pandemic, nursing staff in the paediatric ward for their cooperation and continuous help and support in managing this case, and Al-Adan Hospital and Kuwait Hospital pharmacy staff for their support, particularly pharmacist Haya Alfaiakawi for her support during the work shift in the pharmacy.

Author contributions

M.A.S. conducted patient supervision and drug administration. Y.A.S. provided drug option and calculations. M.A.S. and Y.A.S. wrote the manuscript.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship and/or publication of this article.

Ethical approval

Our institution does not require ethical approval for reporting case reports or case series.

Informed consent

Written informed consent was obtained from a legally authorised representative (patient's father) for anonymised patient information to be published in this article.

ORCID iD

Yousif A Shamsaldeen  <https://orcid.org/0000-0001-8203-0067>

References

1. Elhassanien AF and Alghaiaty HA-A. Neurological manifestations in children with Sanjad–Sakati syndrome. *Int J Gen Med* 2013; 6: 393–398.
2. Naguib K, Gouda S, Elshafey A, et al. Sanjad–Sakati syndrome/Kenny–Caffey syndrome type 1: a study of 21 cases in Kuwait. *East Mediterr Health J* 2009; 15(2): 345–352.
3. Sanjad S, Sakati NA, Abu-Osba Y, et al. A new syndrome of congenital hypoparathyroidism, severe growth failure, and dysmorphic features. *Arch Dis Child* 1991; 66(2): 193–196.
4. Barak V, Schwartz A, Kalickman I, et al. Prevalence of hypophosphatemia in sepsis and infection: the role of cytokines. *Am J Med* 1998; 104(1): 40–47.
5. Bugg N and Jones J. Hypophosphataemia. *Pathophysiology, effects and management on the intensive care unit. Anaesthesia* 1998; 53(9): 895–902.
6. Rady HI and Khalek M. Prevalence and risk factors of hypophosphatemia in pediatric intensive care unit. *J Anesth Crit Care Open Access* 2014; 1: 1–5.
7. de Menezes FS, Leite HP, Fernandez J, et al. Hypophosphatemia in children hospitalized within an intensive care unit. *J Intens Care Med* 2006; 21(4): 235–239.
8. Antachopoulos C, Margeli A, Giannaki M, et al. Transient hypophosphataemia associated with acute infectious disease in paediatric patients. *Scand J Infect Dis* 2002; 34(11): 836–839.
9. Cyriac JM and James E. Switch over from intravenous to oral therapy: a concise overview. *J Pharmacol Pharmacother* 2014; 5(2): 83–87.
10. Wilson R, Langan C, Ball P, et al. Oral gemifloxacin once daily for 5 days compared with sequential therapy with iv ceftriaxone/oral cefuroxime (maximum of 10 days) in the treatment of hospitalized patients with acute exacerbations of chronic bronchitis. *Respir Med* 2003; 97(3): 242–249.
11. Büller HR and Prins MH. Secondary prophylaxis with warfarin for venous thromboembolism. *N Engl J Med* 2003; 349(7): 702–704.
12. Mertz D, Koller M, Haller P, et al. Outcomes of early switching from intravenous to oral antibiotics on medical wards. *J Antimicrob Chemother* 2009; 64(1): 188–199.
13. Kaboli PJ, Hoth AB, McClimon BJ, et al. Clinical pharmacists and inpatient medical care: a systematic review. *Arch Int Med* 2006; 166(9): 955–964.