

Personalising renal function monitoring and interventions in people living with heart failure: protocol for co-designing a care pathway in the RENAL-HF programme

Hargreaves, Suzy C; Armitage, Christopher J; Brown, Benjamin C; Dowding, Dawn; Downing, Jennifer; Goodall, Mark; Gummery, Alison; Lees, Carolyn; Sowden, Emma; Williams, Nefyn H; Young, Bridget; Pirmohamed, Munir

DOI: <https://doi.org/10.3399/BJGPO.2025.0099>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 19 May 2025

Accepted 12 July 2025

© 2025 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by BJGP Open. For editorial process and policies, see: <https://bjgpopen.org/authors/bjgp-open-editorial-process-and-policies>

When citing this article please include the DOI provided above.

Author Accepted Manuscript

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in BJGP Open, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

Title

Personalising renal function monitoring and interventions in people living with heart failure: protocol for co-designing a care pathway in the RENAL-HF programme

Authors

¹Ms Suzy C Hargreaves, BA(Hons), MPhil, FRSPH*

²Professor Christopher J Armitage, BA(Hons), PhD, C Psychol., AFBPsS, FAcSS

³Dr Benjamin C Brown, MB ChB, MRCP, MSc, MPH, PhD, FFCI

³Professor Dawn Dowding, PhD, RN, FAAN, FIAHSI

¹Dr Jennifer Downing, BA(Hons), MA, PhD

¹Dr Mark Goodall, BSc(Hons), DipPH, PhD

⁴Ms Alison Gummery, BSc(Hons)

⁵Dr Carolyn Lees, SFHEA, PhD, MSc, MEd, DN, RGN

³Dr Emma Sowden, BSc(Hons), PhD

¹Professor Nefyn H Williams, PhD, FRCGP

⁶Professor Bridget Young, BA(Hons), PhD,

⁴Professor Sir Munir Pirmohamed, MB ChB (Hons), MRCP, PhD, FRCP, FRCP(E),
FBPh, FMedSci

The RENAL-HF Consortium

¹*Department of Primary Care and Mental Health, Institute of Population Health,
University of Liverpool*

²*Manchester Centre for Health Psychology. University of Manchester and NIHR
Greater Manchester, Patient Safety Research Collaboration, University of
Manchester*

³*School of Health Sciences, University of Manchester*

⁴*Department of Pharmacology and Therapeutics, Institute of Systems, Molecular &
Integrative Biology, University of Liverpool*

⁵*School of Allied Health Professions and Nursing, Institute of Population Health,
University of Liverpool*

⁶*Department of Public Health, Policy and Systems, Institute of Population Health,
University of Liverpool*

*Corresponding author s.c.hargreaves@liverpool.ac.uk

Abstract

Background: Heart failure affects almost one million people in the UK and is increasing in prevalence. Many drugs used to treat heart failure impair renal function and can lead to hospitalisation. Adverse drug problems can be partially mitigated through regular renal monitoring and optimising of drug dose and choice to prevent deterioration of kidney function. This protocol describes part of a wider research programme: personalising **renal** function monitoring and interventions in people living with heart failure (RENAL-HF). **Aim:** The aim of RENAL-HF is to develop improved processes in primary care to manage kidney health in people living with heart failure. **Method:** The protocol covers gathering views of healthcare

professionals, patients and carers, to co-develop a care pathway for use in primary care. Using a mixed methods approach, the work comprises six stages: 1) understanding current practice of optimising heart failure treatment while preserving renal function, 2) co-designing a care pathway including personalised renal function monitoring, thresholds for intervention and clinical guidelines, 3) decision-making to identify elements that will support the care pathway, 4) developing training materials for primary care to enable use of the care pathway, 5) testing the useability of the prototype care pathway, and 6) a feasibility and acceptability study to inform the pre-clinical development and usability of the care pathway ahead of a cluster randomised control trial. **Conclusion:** All stages will elicit evidence from primary care practices, practitioners, and patients with which to assess and refine the care pathway. The evidence will inform how algorithm-guided individualised treatment can be implemented to improve outcomes of patients with heart failure.

Keywords

Renal function, optimising care, evaluation, intervention

MeSH headings: Heart failure, primary health care, United Kingdom, critical pathways, kidney

How this fits in

- Many drugs used to treat heart failure impair renal function.
- There is some guidance on monitoring renal function in people with heart failure following changes in medication, but this does not consider individuals' clinical factors, and there is a marked variation in clinical practice.
- This protocol describes a work package that forms part of a wider programme of research (personalising **renal** function monitoring and interventions in people living with heart failure (RENAL-HF)). RENAL-HF involves various integrated studies to develop better processes in primary care to manage kidney health in people with heart failure, through the creation of a standardised care pathway based on individual patient data, predicted by an algorithm.
- This protocol paper describes the understanding of current practice around the optimisation of heart failure treatment whilst preserving renal function, the co-creation of a care pathway, and a feasibility and acceptability study, ahead of a future cluster randomised control trial (RCT).

Main text

Introduction

Heart Failure is a condition where the heart is unable to pump blood effectively to meet the body's oxygen needs (1). It affects almost one million people in the UK and its prevalence is increasing (2,3). Despite modern treatment reducing morbidity and improving survival for this population, heart failure is a common contributing factor to

acute and chronic kidney disease (4). Renal impairment associated with heart failure drugs is the second most common adverse drug reaction resulting in hospitalisation (4,5), meaning even a small decline in renal function may precipitate costly hospital admission putting further pressure on the healthcare system (6).

Clear guidance on the timing and frequency of reviews is lacking (4,7). There is a need to standardise care processes for this population to reduce variability in clinical practice and improve outcomes by developing an improved care pathway within primary care. It is recommended that identifying patients who are at most risk of declining renal function would benefit from standardised care processes, including monitoring, alongside improving health outcomes and a reduction in hospital admissions (4).

Overview of the wider programme (RENAL HF)

The NIHR Programme Grant for Applied Research (ISRCTN75284771)(8), personalising **renal** function monitoring and interventions in people living with **heart failure** (RENAL-HF) consists of five interconnected work packages (WPs). The study described in this protocol forms a work package (WP2) (comprising stages 1-6) in this wider programme of research. **Figure 1** shows an overview of RENAL-HF and the five WPs and **Figure 2** shows the overall logic model for the research programme, including inputs, activity, response, and short- and longer-term outcomes.

The overall aim of RENAL-HF is to develop better processes in primary care to manage kidney health in people with heart failure. We will use medical records to a) develop technology to predict how often each person with heart failure needs a renal function blood test and b) inform the development of expert advice for healthcare practitioners on how best to adjust medication. In WP1 we will use advanced analytical methods and electronic healthcare records to refine the accuracy of an algorithm that predicts the change in renal function in patients with heart failure. WP3 comprises an RCT to test clinical effectiveness of the pathway developed in WPs1 and 2. We will determine if this approach is more effective or better value for money than the current standard of care by conducting a cost-effectiveness analysis of the pathway (WP4). WP5 will develop and manage patient and public involvement and will integrate public advisors throughout the whole programme to ensure the patient voice is present at all stages of the work.

Working with patients, primary care practitioners and specialists (including cardiologists and nephrologists), we will find the best way of implementing this personalised approach to renal function monitoring and interventions through the co-design of a care pathway.

Method

WP2 will focus on the co-creation of a care pathway, using mixed methods, with six stages (see **Figure 3**). All quantitative analyses will be a mix of descriptive and more in-depth statistical models, such as multi-variable regression to predict behaviour change. Microsoft Excel and IBM SPSS Statistics 29.0.2.0 will be used to analyse the data. Qualitative analyses will be underpinned by the behaviour change wheel (BCW) theory (9,10). Following inductive and reflexive thematic analysis and Framework analysis (11,12) using NVivo 14 data will be deductively mapped onto the Capability, Motivation, Opportunity and Behaviour model (COM-B) and the Theoretical Domains Framework (TDF) (9,10,13). The methods for each stage are outlined below:

1. Understanding current practice

Stage 1 aims to understand current practice and establish patient and practitioner views on standards of care through a survey of healthcare professionals in England, including GPs, pharmacists, and nurses. This will be co-developed by the research team and PPIE and delivered through YouGov. A sample of 600 participants; 115 GPs, 115 pharmacists, and 370 nurses will be recruited. Data will be analysed quantitatively. Open-ended questions will be thematically analysed and mapped onto the TDF (9,10,13). Furthermore, this stage aims to identify barriers and facilitators to optimising the treatment of patients with heart failure while preserving renal function. This will be achieved by conducting brief focused qualitative interviews with GPs, nurses and pharmacists recruited from primary care practices to generate options for intervention functions. The interview guide will be informed by the BCW (9,10) and will be used to understand drivers of optimising treatment of people living with heart failure while preserving renal function. There will be a sample of up to 51 participants. Analysis includes inductive thematic coding and mapping onto the TDF (13).

2. Co-designing the care pathway

Researchers will work closely with stakeholders, the WP1 team and an expert clinical panel (comprised of cardiologists, nephrologists, clinical pharmacologists, and GPs) to develop prototype training material to ensure optimal design and uptake of the RENAL-HF care pathway. This will include personalised renal function monitoring, thresholds for intervention, and clinical guidelines.

The material generated in WP1 and outputs from WP2 stage 1 will be synthesised during series of meetings to generate possible elements of the care pathway. Proposals will be rated in stakeholder workshops. The clinical expert panel will identify clinical parameters for key outputs and treatment changes that may be required based on renal function deterioration. Stakeholder consensus workshops will be held to ensure views of key stakeholders are included in design of the

intervention. Five stakeholder groups of nine individuals each, including patients, pharmacists, nurses, GPs and other key informants, will engage in three rounds, rating the proposals created from the synthesis of stage 1 and of WP1. The RAND/UCLA Appropriateness Methods (RAM) will be used (14,15) to reach group consensus.

3. Decision-making

Elements that will support optimal implementation of the evidence-based system created in stage 2 will be considered in stage 3. The BCW will be used to identify intervention functions and behaviour change techniques that will be most likely to achieve the change required (10). Evaluation of the intervention will be conducted according to the acceptability, practicability, effectiveness/cost-effectiveness, affordability, and safety/side-effects (APPEASE) defined in the BCW (10).

4. Training materials

Prototype training material will be developed for primary care teams (GPs, pharmacists, nurses, care navigators) and refined through beta testing (stage 5) and feasibility study (stage 6), ready for the cluster RCT in WP3. Co-design of the behaviour change interventions will be completed in a series of up to three workshops made up of the wider RENAL-HF research team. Findings stages 1-3 and WP1 will help determine the type of content to be included in training material. Materials will include rationale for the study, how the algorithm and interventions were developed, evidence bases for undertaking interventions (increased renal function monitoring, changing drugs or doses etc.), how to use the care pathway, and how it is being used. The format for the materials will be informed by stakeholders and will include videos hosted on a dedicated website and integrated self-guided learning materials.

5. Beta testing

Evaluation of the usability of the prototype materials (care pathway and clinical guidance tool) developed in previous stages will be conducted through a series of rapid rounds of beta testing. Evaluation will use 'think aloud' interviews with 36 primary care staff (nine from each sub-group; GPs, pharmacists, nurses, and care navigators). This will provide insight into how primary care staff interact with the systems and refinements will be made where appropriate following this (16). During the interviews, participants will be instructed to verbalise their thoughts while conducting predefined tasks with the care pathway within a dummy environment. The prototype training materials will be refined according to the findings of this stage in preparation for the feasibility/acceptability study. Data will be analysed using thematic analysis (11).

6. Feasibility and acceptability sub-study

**Personalising renal function monitoring and interventions in people living with heart failure:
protocol for co-designing a care pathway in the RENAL-HF programme**

Stages 1-5 will inform the pre-clinical development and usability testing of an algorithm-guided care pathway to improve the kidney health of people with heart failure. Stage 6, a feasibility and acceptability study of the whole process in a 'real world environment', will be conducted to inform final refinements before commencement of the cluster RCT (WP3). The algorithm will be installed in five GP practices and will be used for a minimum of six weeks. Quantitative data will be gathered on the number of patients with heart failure, the number of patients with heart failure who already have renal impairment, and those that develop renal impairment.

Patients will be identified by the RENAL-HF care pathway and analytics will be used to understand the number of times the pathway is used, by whom (e.g., job role) and in what circumstance. Assessment of the number of alerts triggered and ignored, actions taken (e.g., patient contacted for blood test performed in accordance with, and/or contrary to the suggested schedule), and changes in treatment after using the pathway. Data on referrals to cardiology and renal services in secondary care and emergency hospital admissions will also be collected. To assess the health inequalities impact of this intervention, data related to gender, ethnicity, ages, and postcodes for patients and the general practice will be collected. This information will contribute to a health inequalities impact assessment informed by the Health Inequalities Assessment Toolkit (HIAT) (17).

Qualitative evaluation will be undertaken in two parts:

- a) In-depth semi-structured interviews will be conducted with a purposive sample of primary care practitioners from each of the five participating practices. Interview guides will be informed by the BCW and the COM-B model to explore facilitators of and barriers to using the pathway. We will recruit a purposive sample of up to 34 practitioners, including those who did and did not engage with the care pathway. Data analysis will use thematic coding and mapping on to the TDF (13) and will be informed by the HIAT (17).
- b) In-depth, semi-structured interviews with a purposive sample of up to 25 people living with heart failure and their informal carers will take place to examine the acceptability and health inequalities impact of the care pathway from the patient perspective. The sampling will take place across the participating practices, aiming for a balanced coverage in terms of patient socio-demographics (gender, ethnicity, and socio-economic status), those with normal renal function, and those with chronic kidney disease stage three or higher. Interviews will be informed by the Theoretical Framework of Acceptability (18). Interviews will be contextualised around the patient's recent experiences of heart failure care, such as monitoring visits, treatment decision-making, medicine-taking, information and support needs, and communication with healthcare professionals. Data analysis will initially be inductive, drawing on reflexive thematic approaches (11), then deductive, using the Theoretical Framework of Acceptability (18) and informed by the HIAT (17).

Data oversight

The Chief Investigator will preserve the confidentiality of the participants taking part in the study and will abide by the Data Protection Act 2018 and the UK GDPR. The University of Liverpool is the data controller.

Discussion

Summary: Individualised identification of patients most at risk of declining renal function through monitoring, enabled by the algorithm and a clear standardised care pathway, should improve patient care and outcomes (4). Stages 1-6 of WP2 described in this protocol paper, will provide strong evidence from primary care practices, practitioners, and patients on which to assess and refine the pathway, before it is tested in a cluster RCT in WP3.

Strengths and limitations: Strengths include the use of behaviour change models, such as the COM-B and TDF in the design of research tools, the use of mixed quantitative and qualitative methods, and embedded patient and public involvement and engagement (PPIE) across the whole research programme.

Anticipated challenges include delivering a complex research programme, within the timescales available, and managing collaborative partnerships across sectors, including contractual delays with external providers adapting electronic health records and providing access to clinical data.

Implications for research and practice: This WP will co-design a new primary care intervention using behaviour change theory to incorporate an algorithm-guided individualised treatment pathway to maintain the renal function of patients treated for heart failure. This new pathway will then be tested in a cluster RCT.

Additional information

Funding: This project is funded by the National Institute for Health and Care Research (NIHR) under its Programme Grants for Applied Research (PGfAR) (Grant Reference Number NIHR202349). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. ISRCTN reference 75284771.

Competing interests: Armitage is supported by the NIHR Manchester Biomedical Research Centre and the NIHR Greater Manchester Patient Safety Research Collaboration. Downing and Goodall are supported by the NIHR Applied Research Collaborative, North West Coast.

Acknowledgements: The RENAL-HF team would like to dedicate this paper to the hard work and dedication of Dr Sudeh Cheraghi-Sohi at the University of Manchester, who passed away at the end of 2024. Her contribution to the

development of the protocol was vital to its success and her influence has been felt throughout the work.

The authors also wish to acknowledge the contribution to this protocol of the wider collaborative research team members who have developed the RENAL-HF project, the design of the research tools, and methodology, including:

- The rest of the RENAL-HF consortium: Jay Wright, Mark Devonald, Darren M Ashcroft, Niels Peek, Bertram Müller-Myhsok, David Jenkins, Alexander Vincent-Paulraj, Matthew Carr, Girven Burnside, Dyfrig Hughes, Sayem Ahmed, Will Hardy, and Naila Khan.
- PPIE representatives Lynn Hedgecoe, Saiqa Ahmed, and Robbie Jones.


References

1. Bozkurt B, Coats AJS, Tsutsui H, Abdelhamid CM, Adamopoulos S, Albert N, et al. Universal definition and classification of heart failure: a report of the Heart Failure Society of America, Heart Failure Association of the European Society of Cardiology, Japanese Heart Failure Society and Writing Committee of the Universal Definition of Heart Failure: Endorsed by the Canadian Heart Failure Society, Heart Failure Association of India, Cardiac Society of Australia and New Zealand, and Chinese Heart Failure Association. *Eur J Heart Fail*. 2021 Mar;23(3):352–80.
2. Ponikowski P, Anker SD, AlHabib KF, Cowie MR, Force TL, Hu S, et al. Heart failure: preventing disease and death worldwide. *ESC Heart Fail*. 2014 Sep;1(1):4–25.
3. Conrad N, Judge A, Canoy D, Tran J, Pinho-Gomes AC, Millett ERC, et al. Temporal Trends and Patterns in Mortality After Incident Heart Failure: A Longitudinal Analysis of 86 000 Individuals. *JAMA Cardiol*. 2019 Nov 1;4(11):1102.
4. Al-Naher A, Wright D, Devonald MAJ, Pirmohamed M. Renal function monitoring in heart failure – what is the optimal frequency? A narrative review. *Br J Clin Pharmacol*. 2018 Jan;84(1):5–17.
5. Pirmohamed M, James S, Meakin S, Green C, Scott AK, Walley TJ, et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ*. 2004 Jul 3;329(7456):15–9.
6. Duffy D. The NHS must break the cycle on heart failure. *Integr Care J* [Internet]. 2023 [cited 2025 Apr 24];February 23 2023. Available from: <https://integratedcarejournal.com/the-nhs-must-break-the-cycle-on-heart-failure/#:~:text=It%20is%20also%20estimated%20that,50%20per%20cent%20in%20the>
7. Al-Naher A. The PERMIT Project: Personalised Renal Function Monitoring via Information Technology. University of Liverpool; 2020.

8. Pirmohamed M. Personalising renal function monitoring and interventions in people living with heart failure: RENAL-HF [Internet]. NIHR; 2022. Available from: <https://fundingawards.nihr.ac.uk/award/NIHR202349>
9. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implement Sci.* 2011 Dec;6(1):42.
10. Michie S, Atkins L, West R. The behaviour change wheel: a guide to designing interventions. London: Silverback Publishing; 2014. 329 p.
11. Braun V, Clarke V. Thematic analysis: a practical guide. Los Angeles London New Delhi Singapore Washington DC Melbourne: SAGE; 2022. 338 p.
12. Ritchie J, editor. Qualitative research practice: a guide for social science students and researchers. 2. ed. Los Angeles, Calif.: Sage; 2014. 430 p.
13. Atkins L, Francis J, Islam R, O'Connor D, Patey A, Ivers N, et al. A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implement Sci.* 2017 Dec;12(1):77.
14. Fitch K, editor. The Rand/UCLA appropriateness method user's manual. Santa Monica: Rand; 2001. 109 p. (MR / Rand DG-XII/RE).
15. Berian JR, Baker TL, Rosenthal RA, Coleman J, Finlayson E, Katlic MR, et al. Application of the RAND-UCLA Appropriateness Methodology to a Large Multidisciplinary Stakeholder Group Evaluating the Validity and Feasibility of Patient-Centered Standards in Geriatric Surgery. *Health Serv Res.* 2018 Oct;53(5):3350–72.
16. Kaklamanou D, Armitage CJ, Jones CR. A further look into compensatory health beliefs: A think aloud study. *Br J Health Psychol.* 2013 Feb;18(1):139–54.
17. Porroche-Escudero A, Popay J. The Health Inequalities Assessment Toolkit: supporting integration of equity into applied health research. *J Public Health.* 2021 Sep 22;43(3):567–72.
18. Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and development of a theoretical framework. *BMC Health Serv Res.* 2017 Dec;17(1):88.


Tables and Figures

Figure 1 Co-creation of the care pathway within context of wider RENAL-HF programme



Personalising renal function monitoring and interventions in people living with heart failure: protocol for co-designing a care pathway in the RENAL-HF programme

Figure 2 Logic model describing the RENAL-HF programme and care pathway



Personalising renal function monitoring and interventions in people living with heart failure: protocol for co-designing a care pathway in the RENAL-HF programme

Figure 3 Stages 1-6 of WP2 within context of wider RENAL-HF programme

