

LJMU Research Online

Liew, BXW, Senden, R, Rugamer, D, Sommer, E, Meijer, K, Mei, Q, Foster, R and Taylor, M

Predicting normative walking biomechanics across the lifespan using seven simple features

https://researchonline.ljmu.ac.uk/id/eprint/27507/

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Liew, BXW, Senden, R, Rugamer, D, Sommer, E, Meijer, K, Mei, Q, Foster, R ORCID logoORCID: https://orcid.org/0000-0003-2410-9839 and Taylor, M (2025) Predicting normative walking biomechanics across the lifespan using seven simple features. Journal of Neuroengineering and

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

RESEARCH Open Access



Predicting normative walking biomechanics across the lifespan using seven simple features

Bernard X. W. Liew^{1*}, Rachel Senden², David Rugamer³, Emanuel Sommer³, Kenneth Meijer⁴, Qichang Mei^{5,6}, Richard Foster⁷ and Matthew Taylor¹

Abstract

The assessment of gait impairments requires a normative reference for comparison. For a fair assessment, comparisons must be made against a reference standing after controlling for sex, anthropometry, and walking characteristics. This study aimed to develop statistical models that predict the lower-limb kinematics and kinetics of walking across the lifespan of healthy participants, using seven simple covariates. Sixteen statistical models predicted 16 joint kinematics and kinetics during walking using the covariates of sex, age, height, mass, side (laterality), walking speed, and cadence, which were developed based on 301 participants between three to 91 years old. The root mean squared error (RMSE) ranged from 4.71° to 7.97° for joint angles, within 0.07 N/kg for ground reaction forces, 0.09 to 0.15 Nm/kg for joint moments, and 0.33 to 0.39 W/kg for joint powers. We provide both online and local apps which can be easily used by clinicians and scientists to generate normative walking data with uncertainty values, which can be used for movement impairment analysis (https://github.com/EmanuelSommer/ShinyFOSR).

Keywords Functional regression, Kinematics, Kinetics, Lifespan, Walking speed

Bernard X. W. Liew

bl19622@essex.ac.uk; liew_xwb@hotmail.com

Introduction

Movement impairments are common in many neurological [1], musculoskeletal [2], cardiovascular [3] disorders, and even manifest naturally due to ageing [4]. The traditional approach to assessing movements occurs in a lab, involves three-dimensional motion capture using multiple optical cameras and force plates [5]. With rapid advancements in computer vision and sensor technologies, movement assessments can now be easily undertaken in clinical environments, using ubiquitous technologies, requiring no specialised training [6]. Assessing movement impairments in patients requires the comparison with typical reference standards from a healthy person or cohort [7, 8]. To ensure a correct interpretation, comparisons must be made with healthy controls after controlling for sex, anthropometry (e.g. body



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

^{*}Correspondence:

¹School of Sport, Rehabilitation and Exercise Sciences, University of Essex, Colchester, Essex, UK

²Department of Physical Therapy, Maastricht University Medical Center, Maastricht, Limburg, The Netherlands

 $^{^3\}mbox{Department}$ of Statistics, Ludwig-Maximilians-Universität München, Munich, Germany

⁴Department of Nutrition and Movement Sciences, NUTRIM Research Institute of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre+. Maastricht. The Netherlands

⁵Faculty of Sports Science, Ningbo University, Ningbo, China

⁶Auckland Bioengineering Institute, The University of Auckland, Auckland,

⁷Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK

mass index, height), and spatiotemporal walking characteristics (e.g. walking speed) [9]. To our knowledge, there are very few tools that non-specialists can leverage to provide that baseline reference to make clinical judgements.

The most common method to compare movement impairments is based on tools, termed "gait indices", such as the Gait Deviation Index [10–12], Gait Profile Score [13], and the Gillette Gait Index [10]. A challenge when using traditional gait indices is that it requires clinicians or researchers to collect data on individuals defined as "normative", from scratch, to develop a database upon which a normative index is quantified and compared. This process can be very time-consuming and expensive. To our knowledge, no open-source "normative" dataset or normative gait indices have been published to enable researchers to compare. Another limitation of traditional gait indices is that typical movements can be affected by movement-specific factors, like the speed of walking [14], and potentially individual-specific factors, like age. This means that if the gait index was developed based on a sample with a different characteristic from the target patient, or performed a movement with different task requirements, traditional gait indices may classify the target patient as impaired.

An alternative to the use of traditional gait indices is a direct population-based statistical approach towards quantifying an average with a measure of uncertainty of any biomechanical parameters of interest [9, 15, 16]. Population-based statistical approaches in biomechanics, such as Statistical Shape Modelling, are not new and have been largely used to scale the morphology of differential anatomical region(s) from a generic model to fit the anthropometry of the measured individual [17]. Less research has focused on the statistical prediction of normal human motion, particularly via predictive factors that are simple, quick, and cheap to measure clinically. A previous study reported that the sagittal plane joint angles during walking of the ankle, knee, and hip could be predicted within < 2°, based only on knowing an individual's walking speed, gender, age, and body mass index [9]. Another study reported that lower-limb joint angles can be predicted within a root-mean-squared error (RMSE) of 4.45° to 6.61° based on age, height, and weight, or based on the shape of the lower-limb bones [18].

A limitation of prior studies [9, 18] is the inclusion of a restricted age spectrum of participants (e.g. 39 ± 13 years in [9], and 22 ± 2 years in [18]), the assumption of statistical linearity between the covariates (also known as predictors) and the biomechanical outcomes and the inclusion only of kinematic features as outcomes for prediction. It is well known that kinematic and kinetic gait differences can be observed across the human lifespan [19]. For example, in a study, only children reported that

ankle push-off power (A2) appeared to mature by 4 years old [20]. Separate research only on adults (i.e. ≥ 18 years old) reported that A2 power peaked again at 60 years [19]. A population-based statistical model of gait biomechanics must include participants from a wider age spectrum of young children to very old adults. Previous studies also assumed a linear relationship between the included covariates and biomechanical outcomes using statistical techniques like partial least squares regression [18] and linear stepwise regression [9]. However, previous studies have reported nonlinear relationships between covariates such as age and biomechanical features like joint power [19]. In other words, the association between a covariate and a predictor varies depending on the value of the covariates.

This study aims to develop statistical models that nonspecialists can use to generate normative kinematic and kinetic data during walking, to (1) provide a baseline reference for comparison, and (2) use it as a tool for clinical and academic education. We will do so by creating models that predict the lower-limb kinematics and kinetics of walking across the lifespan of healthy participants, using simple covariates that can be easily measured in clinical settings, such as sex, age, height, mass, side (laterality), walking speed, and cadence. We hypothesised that the root-mean-squared error (RMSE) for the predicted lower-limb joint angles would be less than the previously defined minimal detectable change threshold of $\leq 5^{\circ}$ [21]. For joint kinetics, we also hypothesise that the RMSE of the predicted joint moments would be less $\leq 0.4 \text{ Nm/Kg}$ [21].

Methods

Participants

Normative data collected from previous studies of Senden et al. were used for this study, and the two studies included 55 typically developed children [22] and 246 healthy adults [23].

Data collection

Three-dimensional (3D) gait analysis was performed at the motion lab of the MUMC+using the Computer Assisted Rehabilitation Environment (CAREN, Motek Medical BV, Amsterdam) system. Participants walked on an instrumented split-belt treadmill (ForceLink, Culemborg, 1000 Hz), while marker trajectories were captured with a 12-camera optical motion capture system (Vicon, Oxford, 100 Hz). Participants wore standardised gymnastic shoes and a safety harness to prevent falls.

Reflective markers were placed on the following landmarks according to the Human Body Lower Limb Model (HBM-II) [24]: Xyphoid process, sternum, C7, T10, bilateral anterior and posterior superior iliac spines, and for both legs on the thigh, medial and lateral femoral condyles and malleoli, anterior of the tibia, second toe, 5th metatarsal, and mid-calcaneus [24]. Markers on medial femoral condyles and malleoli were used for calibration only. The pelvis had 6 degrees of freedom (DOF) (3 global translations of pelvis origin and rotation about Z (yaw), Y (pitch), X (roll)), the hip had three rotational DOFs (sagittal, frontal, rotation), the knee had one sagittal DOF (sagittal), and the ankle had two DOF (sagittal and frontal) [24]. The position and orientation of the segments were resolved using an inverse kinematics algorithm [24].

After a six-minute familiarisation period at comfortable walking speed, participants walked for 250 steps at a comfortable, slow (30% slower than comfortable), and fast (30% faster than comfortable) speed, which were performed in a random order. To determine comfortable walking speed, children performed repeated overground walking trials over a nine-meter walkway, while speed was measured using two movement detection ports. For the healthy adults, the RAMP protocol was used where participants started to walk on the treadmill at 0.5 m/s while the speed was gradually increased to 0.01 m/s every second until a comfortable speed was reached. This was repeated three times and the average of three repetitions was used as the comfortable speed.

The force plate configurations for the analog data were set at 10 Hz for the low-pass prefilter frequency and 20 N for the force threshold. Marker trajectories and force plate data were filtered with a unidirectional 2nd order Butterworth filter at 6 Hz. Gait event detection was defined based on a combination of heel marker kinematics and force plate data (exceeding the threshold of 50 N) [25]. Custom Matlab scripts were used to check data quality and calculate spatiotemporal parameters, kinematics, and kinetics. Kinematic and kinetic data were time-normalised to 100 data points between two consecutive initial contacts for each side. GRF, joint moment and power data were subsequently normalised to body mass (kg). The kinematic and kinetic trajectories have been uploaded and are freely available in the Open Science Framework (OSF) (children: https://osf.io/3xqew/; adults: https://osf.io/t72cw/).

Statistical analysis

All analyses were conducted in R software (v4.4.2) using the *refund* package (v0.1-37). We developed 16 statistical models to predict 16 kinematic and kinetic trajectories – angles (hip flexion, abduction, rotation; knee flexion; ankle plantarflexion and inversion), GRF (vertical and anterior-posterior), moments (hip flexion, abduction, rotation; knee flexion; ankle plantarflexion), and powers (hip, knee, ankle flexion). For all models, the same covariates were included: sex (male or female), age (years), walking speed (m/s), body mass (kg), height (m), cadence

(steps/min), side (left or right), and the random effect of the subject. We use a function-on-scalar model for each of the three joints — specifically the *pffr* function in the *refund* package. As a functional additive model, we model the expectation of each outcome trajectory yi(t) for every subject i at gait point t.

$$y_{i}(t) = \beta_{0}(t) + f_{age}(age_{i}, t) + f_{speed}(speed_{i}, t)$$

$$+ f_{cadence}(cadence_{i}, t) + f_{ht}(height_{i}, t)$$

$$+ f_{mass}(mass_{i}, t) + sex_{i}\beta_{sex}$$

$$+ side_{i}\beta_{side} + b_{i}(t) + \varepsilon_{ij}$$

where $\beta_0(t)$ is the model's time-varying intercept, $\beta_{\rm sex}$ a time-independent effect of sex, $\beta_{\rm side}$ a time-independent effect of side, f(, t) indicate functional effects that are estimated to be non-linear both in the direction of the covariate and across time, bi(t) are time-varying random intercepts, and ε_i and independent Gaussian error term.

An 80:20 split of the data was performed to create a training set (80%, n=240 participants) to develop the models, and a testing set (n=61 participants) to validate the accuracy of the outcome predictions. The prediction performance of the models was determined by comparing the 16 predicted outcomes in the test set, against their original values using the Root Integrated Mean Squared Error (BW), relative Root Integrated Mean Squared Error (relRMSE, %) [26], and Pearson correlation coefficient (cor) [27, 28].

$$RMSE = \sqrt{\frac{\int_{0}^{T} \left[u_{obs}\left(t\right) - u_{pred}\left(t\right)\right]^{2} dt}{T}}$$
 (1)

$$relRMSE = \frac{RMSE}{0.5 \left[\sum_{i=1}^{2} \left(\max_{0 < t < T} \left(u_i \left(t \right) \right) - \min_{0 < t < T} \left(u_i \left(t \right) \right) \right) \right]} \times 100\%$$

where T represents the stance duration between initial contact and toe-off, $u_{obs}\left(t\right)$ represents the value at the t^{th} time point of the observed outcome, $u_{pred}\left(t\right)$ represents the value at the t^{th} time point of the predicted outcome, and i represents either the observed or predicted outcomes. We also report the association (point estimate and 95% confidence interval [CI]) between the outcomes and each of the fixed effect covariates. For the continuous covariates of age, speed, cadence, height and mass, we selected the 25th, 50th, and 75th quantiles of the values to plot the smooth association estimate.

To make the model open source and easily accessible, we retrained the models using the entire dataset (n = 301) and created a shiny app (https://github.com/EmanuelSommer/ShinyFOSR) for readers to input different covariate values to predict the mean waveform value of different kinematic and kinetic outcomes. Readers can use

the app to predict, plot, and export the values as a figure and table. We also included a reference with a detailed and easy set of instructions to download and use the app locally on their computer. This local version can predict the mean values and the 95%CI or the standard deviation.

Results

Basic descriptive summaries of the cohort can be found in Figure. 1. Figures 2 and 3 illustrate the average predicted kinematic and kinetic trajectories, alongside the observed trajectories in the test dataset. RMSE ranged from 4.71° to 7.97° for kinematic data, while relRMSE ranged from 9.85% to 50.72% (Table 1). Accuracy was the best generally for sagittal plane joint angles and the poorest for transverse plane angles (Table 1). For kinetic data, the statistical models predicted GRFs to be within 0.07 N/kg and <7% for RMSE, and relRMSE (Table 1). The statistical models were better at predicting joint moments than joint powers, with an accuracy for the former ranging from 0.09 to 0.15 Nm/kg for RMSE and 9.23% to 24.71% for relRMSE; and for the latter ranging from 0.33 to 0.39 W/kg for RMSE and 12.98 to 17.91% for relRMSE (Table 1).

Smooth effect plots

The smooth effect plots support the nonlinear association between the included covariates of age and speed and the kinematic and kinetic outcomes. For the covariate age, the greatest effect is on the hip sagittal plane angle at 26% of the gait cycle, where at the age of 46 years it was associated with greater hip extension angle by -16.9° (95%CI -17.3° to -16.5°) (Fig. 4). For joint moments, age had the greatest effect on the hip sagittal moment, where at age 46 years, it was associated with greater hip flexion moment by -0.35Nm/kg (95%CI -0.36 to -0.33Nm/kg) at 21% of the gait cycle (Fig. 5). Age also had the greatest effect on hip flexion power, where at age 46 years, it was associated with greater hip power absorption by -0.35 W/kg (95%CI -0.39 to -0.31 W/kg) at 19% of the gait cycle (Fig. 5).

For the covariate speed, the greatest effect is on the hip sagittal angle at 1% of the gait cycle, where with a speed of 1.84 m/s, it was associated with an increase in hip flexion angle 13.4° (95%CI 13.4° to 14.2°) (Fig. 6). For joint moments, speed had the greatest effect on the hip sagittal moment, where at a speed of 1.84 m/s, it was associated with an increase in hip extensor moment by 0.43Nm/kg (95%CI 0.42 to 0.45Nm/kg) at 1% of the gait cycle (Fig. 7). Speed had the greatest effect on ankle plantarflexion




Fig. 1 Descriptive characteristics of the included participants (n = 301). Point estimate reflect means with error bars representing one standard deviation (a – e), apart from the number of participants (f). Red (male), green (female)




Fig. 2 Average predicted (blue) kinematic trajectories against the observed (black) trajectories for the test dataset




Fig. 3 Average predicted (blue) kinetic trajectories against the observed (black) trajectories for the test dataset

Table 1 Accuracy of predicted kinematic and kinetic trajectories

Variables	RMSE	relRMSE (%)	Correla- tion
Hip Sagittal Angle (°)	7.97 (3.48)	16.59 (8.45)	0.94 (0.06)
Hip Frontal Angle (°)	3.44 (1.45)	21.38 (9.7)	0.89 (0.09)
Hip Transverse Angle (°)	7.94 (3.32)	50.72 (22.1)	0.44 (0.44)
Knee Sagittal Angle (°)	6.21 (2.78)	9.85 (4.7)	0.97 (0.04)
Ankle Sagittal Angle (°)	4.71 (4.13)	18.96 (19.22)	0.88 (0.1)
Ankle Frontal Angle (°)	7.71 (5.85)	45.81 (43.46)	0.38 (0.35)
GRF AP (N/kg)	0.02 (0.01)	6.63 (3.85)	0.96 (0.06)
GRF Vertical (N/kg)	0.07 (0.03)	6.15 (2.7)	0.99 (0.02)
Hip Sagittal Moment (Nm/kg)	0.13 (0.08)	11.07 (7.56)	0.93 (0.11)
Hip Frontal Moment (Nm/kg)	0.12 (0.04)	11.18 (3.8)	0.96 (0.04)
Hip Transverse Moment (Nm/kg)	0.09 (0.03)	24.71 (10.16)	0.79 (0.2)
Knee Sagittal Moment (Nm/kg)	0.15 (0.06)	17.26 (8.38)	0.83 (0.21)
Ankle Sagittal Moment (Nm/kg)	0.15 (0.07)	9.23 (5.13)	0.96 (0.06)
Hip Sagittal Power (W/kg)	0.33 (0.18)	17.61 (7.22)	0.84 (0.15)
Knee Sagittal Power (W/kg)	0.34 (0.15)	17.91 (5.18)	0.78 (0.16)
Ankle Sagittal Power (W/kg)	0.39 (0.18)	12.98 (6.18)	0.79 (0.25)

RMSE – root mean squared error; relRMSE – relative root mean squared error; AP – anterior-posterior; GRF – ground reaction force.

power, where at a speed of 1.84 m/s it was associated with an increase in power generation by 0.82 W/kg (95%CI 0.79 to 0.85 W/kg) at 49% of the gait cycle (Fig. 7).

Discussion

Determination of impaired movement depends on knowing normal movement. Statistical modelling based on simple clinical measures represents a simple way that scientists and clinicians can predict key walking biomechanics of healthy individuals. The RMSE averaged across the stride cycle for the lower-limb joint angles was $< 8^{\circ}$, poorer than our hypothesis of $< 5^{\circ}$, and also marginally poorer than prior studies $< 7^{\circ}$ [18]. In support of our hypothesis, the RMSE was between 0.09 and 0.15 Nm/kg for joint moments, and this was also better than prior studies [29].

The predicted RMSE for joint angles in this study was approximately 5° greater than previous studies, which reported that the RMSE for sagittal plane joint angles was within 2° [9, 30]. Previous studies have reported that the minimal detectable change values of lower-limb joint angles during walking of $\sim 5^\circ$, but can increase up to 7° [21]. RMSE values greater than minimal detectable change values suggest some biomechanical outcomes should be used with caution. For example, readers should have more confidence in using our models to generate normative ankle plantarflexion angles ($\leq 5^\circ$), and use more caution when using them to create hip axial

rotation angles (\geq 5°) (Table 1). However, even though the error metrics presented reflect the waveform-averaged, the accuracy of our prediction models is not uniform across the stride cycle, as observed in Figures. 2 and 3, with better accuracy in some phases of the stride cycle than in other phases. Also, there is no consensus on a single clinically meaningful threshold kinematic and kinetic value, which is likely to vary depending on the clinical condition and joints investigated. Users of the present prediction models in defining normative walking gait biomechanics should be aware of our reported RMSE values and determine clinically if these errors exceed a clinically meaningful threshold.

The differences in the predicted accuracy of our biomechanical waveforms between studies may be due to different validation methods [9, 30]. Previous studies used leave-one-subject-out cross-validation (LOSOCV) as their validation approach [9, 30], whereas the present study used a train-test split approach. LOSOCV uses more data, apart from one subject, for training. This means that a greater percentage of the data was used for training, and a smaller percentage of data was used to validate the model, compared to a train-test split validation approach. For example, the study of Moissenet et al. [9] used 1325 observations (53 subjects, 5 speeds, 5 trials) for each training loop, whereas we used only 720 observations (240 subjects, 3 speeds) for training. The present findings thus provide a more conservative estimate of gait kinematics predictive modelling, which may be more transferable to unseen real-world applications.

The present study demonstrates that normative joint kinetics during walking can be predicted using simple clinical measures, without having to rely on more complex biomechanical inputs as predictors. The predicted RMSE for GRF, moments, and powers in this study ranged between 0.02 and 0.07 N/kg, 0.09 to 0.15Nm/ kg, and 0.33 to 0.39 W/kg, respectively. Surprisingly, these error ranges were less than machine learning prediction models of joint moments using 3D motion capture inputs in running (RMSE: 0.07 to 0.54Nm/kg) [29], and less than models using inertial measurement units (IMUs) as inputs for walking (RMSE: 0.07 to 0.24 Nm/ kg) [31]. Wouda et al. predicted vertical GRF to a RMSE of < 0.27BW using three IMU inputs for running [32]. Another study predicted ankle joint power for walking using IMU sensors with a RMSE of < 0.21 W/kg [33]. Statistical models like the present and others [9], are useful for population average prediction - which cannot capture variation between movement cycles, whereas more complex prediction models built upon biomechanical input predictors may be more useful for capturing cyclespecific predictions.

Whilst the present study did not specify apriori a linear relation, previous studies have assumed a linear




Fig. 4 Smooth plots with error clouds as 95% confidence interval for the covariate of age and joint kinematics. Black (23 years old), blue (46 years old), green (68 years old)

relationship between the predictors investigated (e.g. age, speed) and the biomechanical outcomes [9, 16], whilst another study assumed a quadratic association with the outcomes [30]. The present study reported a non-linear association between age and ankle plantar flexion, given that ankle plantar flexion increased between 23 and 48 years old and decreased from 48 to 68 years old (Fig. 4e). The non-linear association between age and our 16 biomechanical outcomes investigated was evident in our smooth plots (Figs. 4 and 5). Studies which used linear regression methods reported non-significant linear associations between age and ankle plantar flexion during push-off [9], while other studies reported a decreased ankle plantar flexion during push-off in older adults compared to younger adults [34]. This study provides evidence that it may not be appropriate to assume linearity across all values of the covariates and the outcome across a gait cycle. Hence, whilst linear-regression-based approaches provide very interpretable results (e.g. a beta coefficient), they may not adequately model the relationships between the covariates and the biomechanical outcomes.

A strength of the present study is the fact that we included 301 participants between three and 91 years old, which represents a five times greater sample size than recent studies [9]. Apart from the age category of >80 years old, the current study included >20 participants in each decade of years for adults, included in the modelling, making our statistical model more representative of a lifespan population cohort. Non-biomechanical specialists can use our models and online tool to quickly generate data of normal walking joint kinematics and kinetics using simple clinically oriented variables. This is more advantageous than existing metrics like the Gait Deviation Index, which still requires data collection to establish a reference, nor can such metrics be tuned to key personal and gait characteristics easily. For clinicians, the normative data can be used to compare biomechanical




Fig. 5 Smooth plots with error clouds as 95% confidence interval for the covariate of age and joint kinetics. Black (23 years old), blue (46 years old), green (68 years old)

values from patients collected either using traditional motion capture or more recent portable technologies, like OpenCap. Movement impairment assessments may be used as an outcome measure, for prognosis, and even for treatment-decision making. Our lifespan approach also means that clinicians can now account for maturation changes when making clinical judgements in children and adolescents. Lastly, clinical and academic educators may use our models for teaching purposes to visualise normal walking biomechanics.

This study has some limitations. First, the included participants were collected whilst walking on a treadmill, which might differ from overground walking [35]. Differences between treadmill and overground walking ranged from 0.84° (pelvic tilt) to 6.42° (peak knee flexion during swing) for joint angles, and 0.02 Nm/kg (peak knee extension moment) to 0.32 Nm/kg (peak hip extension moment) for joint moment [35]. Second, although we included covariates which have previously been shown to have a significant association with walking biomechanics, more variables may play an important role when developing population-based models which were not currently measured, like ethnicity [36] and psychological factors [37]. Third, our biomechanical model had only one DOF and two DOFs at the knee and ankle, respectively. This means that statistical models to predict the constrained knee and ankle kinematics and kinetics were not generated. Fourth, segment kinematics, like that of the pelvis, were not reported, given that the focus of this paper was on joint kinematics and kinetics. Lastly, our biomechanical model was not scaled geometrically to account for factors like tibial torsion, which has a significant effect on lower-limb joint kinematics and kinetics [38, 39]. Also, we did not assess anthropometric measurements like femoral and tibial torsion angles to use within our statistical models. Although this may increase the performance of our models, it would also make them more complex to use clinically.

Conclusions

Statistical models using seven clinical covariates can predict most normative walking kinematics and kinetics, apart from non-sagittal plane kinematics, to a level comparable to more complex machine learning models using biomechanical covariates. Both our online and local apps can be used by clinicians and scientists to generate normative data with uncertainty values, which can be plotted against patient data for reporting.




Fig. 6 Smooth plots with error clouds as 95% confidence interval for the covariate of speed and joint kinematics. Black (0.92 m/s), blue (1.38 m/s), green (1.64 m/s)




Fig. 7 Smooth plots with error clouds as 95% confidence interval for the covariate of speed and joint kinetics. Black (0.92 m/s), blue (1.38 m/s), green (1.64 m/s)

Acknowledgements

Not applicable.

Author contributions

Data collection: Rachel Senden and Kenneth Meijer. Data processing: Bernard Liew, David Rugamer, Rachel Senden and Kenneth Meijer. Statistical modelling: Bernard Liew, David Rugamer, Emanuel Sommer. Model Validation: All authors. Visualisation: Bernard Liew, David Rugamer, Emanuel Sommer, Qichang Mei, Matthew Taylor. Manuscript writing: All authors. Manuscript revision: All authors. Supervision: Kenneth Meijer, Richard Foster, Matthew Taylor.

Funding

Bernard Liew is funded by the Medical Research Council (MR/Y013557/1) and Innovate UK (10093679).

Data availability

The data is fully open access as reported in the manuscript hyperlinks.

Declarations

Ethics approval and consent to participate

No ethical approval was required for this study as we used open-source data.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 2 April 2025 / Accepted: 29 September 2025 Published online: 24 October 2025

References

- Noorkoiv M, Lavelle G, Theis N, Korff T, Kilbride C, Baltzopoulos V, et al. Predictors of walking efficiency in children with cerebral palsy: Lower-Body joint Angles, Moments, and power. Phys Ther. 2019;99:711–20.
- Resende RA, Fonseca ST, Silva PL, Magalhães CMB, Kirkwood RN. Power at hip, knee and ankle joints are compromised in women with mild and moderate knee osteoarthritis. Clin Biomech (Bristol Avon). 2012;27:1038–44.
- Panizzolo FA, Maiorana AJ, Naylor LH, Dembo L, Lloyd DG, Green DJ, et al. Gait analysis in chronic heart failure: the calf as a locus of impaired walking capacity. J Biomech. 2014;47:3719–25.
- Boyer KA, Johnson RT, Banks JJ, Jewell C, Hafer JF. Systematic review and meta-analysis of gait mechanics in young and older adults. Exp Gerontol. 2017;95:63–70
- Benedetti MG, Catani F, Leardini A, Pignotti E, Giannini S. Data management in gait analysis for clinical applications. Clin Biomech. 1998;13:204–15.
- Uhlrich SD, Falisse A, Kidziński Ł, Muccini J, Ko M, Chaudhari AS, et al. OpenCap: 3D human movement dynamics from smartphone videos. BioRxiv. 2022;2022(2007):2007–499061.
- McKay MJ, Baldwin JN, Ferreira P, Simic M, Vanicek N, Wojciechowski E, et al. Spatiotemporal and plantar pressure patterns of 1000 healthy individuals aged 3-101 years. Gait Posture. 2017;58:78–87.
- 8. Cimolin V, Galli M. Summary measures for clinical gait analysis: a literature review. Gait Posture. 2014;39:1005–10.
- Moissenet F, Leboeuf F, Armand S. Lower limb sagittal gait kinematics can be predicted based on walking speed, gender, age and BMI. Sci Rep. 2019:9:9510.
- Schutte LM, Narayanan U, Stout JL, Selber P, Gage JR, Schwartz MH. An index for quantifying deviations from normal gait. Gait Posture. 2000;11:25–31.
- Schwartz MH, Rozumalski A. The gait deviation index: A new comprehensive index of gait pathology. Gait Posture. 2008;28:351–7.
- 12. Rozumalski A, Schwartz MH. The GDI-Kinetic: A new index for quantifying kinetic deviations from normal gait. Gait Posture. 2011;33:730–2.
- Baker R, McGinley JL, Schwartz MH, Beynon S, Rozumalski A, Graham HK, et al. The gait profile score and movement analysis profile. Gait Posture. 2009;30:265–9.

- Schreiber C, Armand S, Moissenet F. Influence of normative data's walking speed on the computation of conventional gait indices. J Biomech. 2018;76:68–73.
- Røislien J, Skare Ø, Gustavsen M, Broch NL, Rennie L, Opheim A. Simultaneous Estimation of effects of gender, age and walking speed on kinematic gait data. Gait Posture. 2009;30:441–5.
- Chehab EF, Andriacchi TP, Favre J. Speed, age, sex, and body mass index provide a rigorous basis for comparing the kinematic and kinetic profiles of the lower extremity during walking. J Biomech. 2017;58:11–20.
- Nolte D, Ko ST, Bull AMJ, Kedgley AE. Reconstruction of the lower limb bones from digitised anatomical landmarks using statistical shape modelling. Gait Posture. 2020;77:269–75.
- De Roeck J, Duquesne K, Van Houcke J, Audenaert EA. Statistical-Shape prediction of lower limb kinematics during Cycling, Squatting, Lunging, and Stepping-Are bone geometry predictors helpful? Front Bioeng Biotechnol. 2021;9:696360.
- Liew BXW, Rugamer D, Duffy K, Taylor M, Jackson J. The mechanical energetics of walking across the adult lifespan. PLoS ONE. 2021;16:e0259817.
- Samson W, Van Hamme A, Desroches G, Dohin B, Dumas R, Chèze L. Biomechanical maturation of joint dynamics during early childhood: updated conclusions. J Biomech. 2013;46:2258–63.
- Wilken JM, Rodriguez KM, Brawner M, Darter BJ. Reliability and minimal detectible change values for gait kinematics and kinetics in healthy adults. Gait Posture. 2012;35:301–7.
- Senden R, Marcellis R, Meijer K, Willems P, Lenssen T, Staal H, et al. Dataset of 3D gait analysis in typically developing children walking at three different speeds on an instrumented treadmill in virtual reality. Data Brief. 2023;48:109142.
- Senden R, Marcellis R, Willems P, Witlox M, Meijer K. Normative 3D gait data of healthy adults walking at three different speeds on an instrumented treadmill in virtual reality. Data Brief. 2024;53:110230.
- van den Bogert AJ, Geijtenbeek T, Even-Zohar O, Steenbrink F, Hardin EC.
 A real-time system for Biomechanical analysis of human movement and muscle function. Med Biol Eng Comput. 2013;51:1069–77.
- Zeni JA Jr., Richards JG, Higginson JS. Two simple methods for determining gait events during treadmill and overground walking using kinematic data. Gait Posture. 2008:27:710–4.
- Ren L, Jones RK, Howard D. Whole body inverse dynamics over a complete gait cycle based only on measured kinematics. J Biomech. 2008;41:2750–9.
- Johnson WR, Alderson J, Lloyd D, Mian A. Predicting athlete ground reaction forces and moments from Spatio-Temporal driven CNN models. IEEE Trans Biomed Eng. 2019;66:689–94.

- Johnson WR, Mian A, Lloyd DG, Alderson JA. On-field player workload exposure and knee injury risk monitoring via deep learning. J Biomech. 2019;93:185–93.
- Liew BXW, Rügamer D, Zhai X, Wang Y, Morris S, Netto K. Comparing shallow, deep, and transfer learning in predicting joint moments in running. J Biomech. 2021;129:110820.
- Koopman B, van Asseldonk EHF, van der Kooij H. Speed-dependent reference joint trajectory generation for robotic gait support. J Biomech. 2014;47:1447–58.
- 31. Altai Z, Boukhennoufa I, Zhai X, Phillips A, Moran J, Liew BXW. Performance of multiple neural networks in predicting lower limb joint moments using wearable sensors. Front Bioeng Biotechnol. 2023;11:1215770.
- 32. Wouda FJ, Giuberti M, Bellusci G, Maartens E, Reenalda J, van Beijnum B-JF et al. Estimation of vertical ground reaction forces and sagittal knee kinematics during running using three inertial sensors. Front Physiol. 2018;9.
- 33. Jiang X, Gholami M, Khoshnam M, Eng JJ, Menon C. Estimation of ankle joint power during walking using two inertial sensors. Sensors. 2019;19:2796.
- 34. Oliveira CF, Vieira ER, Machado Sousa FM, Vilas-Boas JP. Kinematic changes during prolonged Fast-Walking in old and young adults. Front Med. 2017;4.
- Senden R, Marcellis R, Meijer K, Willems P, Lenssen T, Staal H, et al. Comparison of sagittal plane gait characteristics between the overground and treadmill approach for gait analysis in typically developing children. PeerJ. 2022;10:e13752.
- 36. Hill CN, Reed W, Schmitt D, Sands LP, Queen RM. Racial differences in gait mechanics. J Biomech. 2020;112:110070.
- Nagano H, Sarashina E, Sparrow W, Mizukami K, Begg R. General mental health is associated with gait asymmetry. Volume 19. Basel, Switzerland: Sensor: 2019
- Passmore E, Graham HK, Pandy MG, Sangeux M. Hip- and patellofemoraljoint loading during gait are increased in children with idiopathic torsional deformities. Gait Posture. 2018;63:228–35.
- Aiona M, Calligeros K, Pierce R. Coronal plane knee moments improve after correcting external tibial torsion in patients with cerebral palsy. Clin Orthop Relat Research*. 2012;470:1327–33.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.