



LJMU Research Online

Razavi, SM, Nahar, L, Talichi, H and Sarker, SD

Ferulone A and ferulone B: two new coumarin esters from *Ferula orientalis* L. roots, Natural Products Research

<http://researchonline.ljmu.ac.uk/id/eprint/2851/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Razavi, SM, Nahar, L, Talichi, H and Sarker, SD (2016) Ferulone A and ferulone B: two new coumarin esters from *Ferula orientalis* L. roots, Natural Products Research. Natural Product Research. ISSN 1478-6419

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Ferulone A and ferulone B: two new coumarinesters from *Ferula orientalis* L. roots

Seyed Mehdi Razavi^{a*}, Lutfun Nahar^b, Hamideh Talischi^a, and Satyajit Dey Sarker^b

^a*Department of Biology, Faculty of Sciences, University of Mohaghegh Ardabili, Ardabil, Iran*

^b*Medicinal Chemistry and Natural Products Research Group, School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3 3AF, England, UK*

Abstract

Ferula orientalis (Apiaceae) is a well known perennial herb growing wild in Iran used in traditional medicine. To perform phytochemical studies, dried ground roots of *F. orientalis* were sequentially Soxhlet-extracted using *n*-hexane, dichloromethane and methanol. A combination of vacuum liquid chromatography and preparative thin layer chromatographic analyses were performed to isolate coumarin esters. The structures of the isolated compounds were elucidated by spectroscopic means, and *in vitro* free-radical-scavenging property was determined by the DPPH assay. Two new coumarin esters, 7-*O*-(4,8,12,16-tetrahydroxy-4,8,12,16-tetramethyl-heptadecanoyl)-coumarin and 7-*O*-(4-hydroxy-4,8,12-trimethyl-trideca-7,11-dienoyl)-coumarin, named ferulone A and ferulone B, respectively, were isolated from the *n*-hexane extract of the roots of *F. orientalis*. Both compounds showed a low level of free-radical-scavenging property with the RC₅₀ values of 0.252 and 0.556 mg/mL for compounds 1 and 2, respectively, as opposed to that of the positive control (quercetin) 0.004 mg/mL. This is the first report on the purification of coumarin esters from the genus *Ferula*.

Keywords: *Ferula orientalis*, ferulone, free-radical-scavenging, DPPH, umbelliferone,

*Corresponding author. Email: m_razavi@uma.ac.ir

1. Introduction

Ferula orientalis L. (Apiaceae) of the subgenus *Peucedanoides* (Boiss.) Korovin is a well known medicinal plant (height: 100-150 cm), and one of the *ca.* 150 species of the genus *Ferula* L., which are widely distributed throughout the central and western Asia and the Mediterranean region (Miskiet al. 1987). It grows on rocky steps at 1600-2900 m, and has distinguishable yellow flowers, which bloom during late May and June (Davis 1971). Different species of *Ferula* were commonly named as Anghozeh, Koma, Sekbineh and Barijeh in Iran where they are used as adulterant, culinary spice and medicinal plants (Mozefferian 2003; Iranshahy & Iranshahi 2011). *Ferula* species are used in traditional medicine to treat flatulence and as an anticonvulsant, stimulant, expectorant and cancer chemopreventer (Iranshahi et al. 2008; Bagheriet al. 2010).

Previous studies on *F. orientalis* revealed its antioxidant (Kartalet al. 2007) and antifungal (Alinezhadet al. 2011) properties, and the presence of daucane and germacrane-type sesquiterpenes (Miskiet al. 1987). We previously described a ester coumarin from *F. persica* roots (Razavi & Janani 2015).

In continuation of our work on the genus *Ferula* (Auziet al. 2008; Geroushiet al. 2010), we now report on the isolation, identification and free-radical-scavenging property of two new coumarin esters, named ferulone A (**1**) and ferulone B (**2**) (Figure 1), from the roots of *F. orientalis*.

2. Results and discussion

Chromatographic analyses of the *n*-hexane extract of the roots of *F. orientalis* afforded two new coumarin esters, which were identified as 7-*O*-(4,8,12,16-tetrahydroxy-4,8,12,16-tetramethyl-heptadecanoyl)-coumarin (**1**, named ferulone A) and 7-*O*-(4-hydroxy-4,8,12-trimethyl-trideca-7,11-dienoyl)-coumarin (**2**, named ferulone B) by spectroscopic means (UV-Vis, IR, MS and 1D and 2D NMR).

Compound **1** was isolated as a yellowish-brown color oily substance, while compound **2** was a brown oil. The UV absorption maxima at 320, 270, 265, 259 and 252 nm suggested that compounds **1** and **2** were coumarins. In the ¹H NMR spectrum of **1** (Table S1), the doublets (integrating for 1H each) at δ_H6.26 and 7.64 with the coupling constant *J* = 9.5 Hz were the signature peaks for a coumarin nucleus without any oxygenation at C-5 (Murray et al. 1982). There were three aromatic signals, a doublet at δ_H7.37 (*J* = 8.5 Hz), a doublet of a doublet at δ_H6.84 (*J* = 2.0, 8.5 Hz) and a doublet at δ_H6.82 (*J* = 2.0 Hz), corresponding to *ortho*, *ortho-meta* and *meta* couplings among the aromatic protons suggesting a usual oxygenation at C-7 on the coumarin nucleus, i.e., a classic umbelliferone (**3**) skeleton (Figure 1) (Murray 1982; Kauret et al., 2012). In addition to the ¹H NMR signals assignable to a 7-oxygenated coumarin nucleus (**3**), there were signals for five methyl groups in the region of δ_H1.17-1.22, and overlapped peaks in the region of δ_H1.11-2.29, integrated for 22 protons and assignable to 11 aliphatic methylene groups, suggesting the presence of a long-chain fatty acid moiety in the molecule. In the ¹³C NMR (Table S1, Figure S2) spectrum of **1**, in addition the usual signals attributable to an umbelliferone (**3**) skeleton, there were signals for an ester carbonyl group at δ_C 170.5, five methyl groups at δ_C18.0, 21.0, 22.7, 22.8 and 21.7, ten methylene signals in the region of δ_C24.2-31.4, and a methylene signal at δ_C43.6, assignable to the methylene next to the ester carbonyl functionality, and four oxygenated quaternary carbons in the region of δ_C67.7-74.7. All these signals further supported the presence of a long-chain fatty acid moiety. With the help of a combination of ¹H-¹H COSY and ¹H-¹³C HMBC (Figure S1, Table S2) and ¹H-¹³C HSQC, these signals could be assigned unambiguously to a 4,8,12,16-tetrahydroxy-4,8,12,16-tetramethyl-heptadecanoyl moiety, which could only be ester-linked to C-7 of the umbelliferone (**3**) nucleus. The ESIMS spectrum of **1** revealed the *pseudo*-molecular ion [M+H]⁺ at *m/z* 535, and in the HRMS, this ion was observed at *m/z*535.3270 (calculated 535.3271 for C₃₀H₄₇O₈) confirming the molecular formula C₃₀H₄₆O₈ for coumarin **1**. Thus, taking all these spectroscopic data into account, coumarin **1** could be identified unequivocally as 7-*O*-(4,8,12,16-tetrahydroxy-

4,8,12,16-tetramethyl-heptadecanoyl)-coumarin, which, to the best of our knowledge, is a new natural product.

The ^1H NMR and ^{13}C NMR signals (Table S1) for compound **2** were quite similar to those of coumarin **1**, especially the signals associated with the coumarin nucleus, suggesting that this compound also had an umbelliferone (**3**) skeleton (Figure 1). However, the signals which were not associated with the umbelliferone moiety, were significantly different from those of compound **1**. In the ^1H NMR spectrum, in addition to signals assignable to an umbelliferone (**3**) nucleus, there were signals for four methyl groups in the region of δ_{H} 1.23-1.71, three of which [δ_{H} 1.45, 1.66 and 1.71] could be assigned to methyl groups directly linked to olefinic quaternary carbons, and overlapped peaks in the region of δ_{H} 1.20-2.29, integrated for 12 protons and assignable to six aliphatic methylene groups, and signals for two olefinic methine at δ_{H} 5.00-5.55, suggested the presence of an unsaturated long-chain fatty acid moiety in the molecule. In the ^{13}C NMR (Table S1, Figure S3) spectrum of **2**, there were signals for an ester carbonyl group at δ_{C} 170.5, four methyl groups at δ_{C} 22.9, 27.9, 28.9 and 29.7, and five methylene signals in the region of δ_{C} 26.2-32.4, and a methylene signal at δ_{C} 43.7 assignable to the methylene next to the ester carbonyl functionality, two olefinic methine signals at δ_{C} 119.2 and 124.5, and two olefinic quaternary carbons at δ_{C} 132.5 and 131.6. All these signals further supported the presence of a long-chain unsaturated fatty acid moiety in the molecule. With the help of a combination of ^1H - ^1H COSY and ^1H - ^{13}C HMBC (Figure S1, Table S2) and ^1H - ^{13}C HSQC, these signals could be assigned to a 4-hydroxy-4,8,12-trimethyl-trideca-7,11-dienoyl moiety, which could only be ester-linked to C-7 of the umbelliferone (**3**) nucleus as in compound **1**. The ESIMS spectrum of **2** revealed the *pseudo*-molecular ion $[\text{M}+\text{H}]^+$ at m/z 413, and in the HRMS, this ion was observed at m/z 413.2326 (calculated 413.2328 for $\text{C}_{25}\text{H}_{33}\text{O}_5$) confirming the molecular formula $\text{C}_{25}\text{H}_{32}\text{O}_5$ for coumarin **2**. Thus, coumarin **2** could be identified unambiguously as 7-*O*-(4-hydroxy-4,8,12-trimethyl-trideca-7,11-dienoyl)-coumarin, which, to the best of our knowledge, is a new natural product.

The *in vitro* free-radical-scavenging activities of the isolated compounds were evaluated by the 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay (Takao et al., 1994; Kumarasamy et al. 2002; Chima et al. 2014). Compounds **1** and **2** exhibited low level of free-radical-scavenging property, with the RC_{50} values of 0.252 and 0.556 mg/mL as opposed to that of the positive control (quercetin) 0.004 mg/mL.

A review of literature showed that several coumarins with various sesquiterpenyl moieties have already been reported from different species of the genus *Ferula* (El-Razek et al. 2001; Iranshahiet al. 2010a,b; Bashir et al. 2014; Kasaian et al. 2014; Asghari et al. 2015). We previously described a ester coumarin from *F. persica* roots, another common *Ferula* species in Iran (Razavi & Janani 2015). The identified ester coumarin, ferulone c, has similar structure with two ester coumarins 2w3 isolated from *F. orientalis* in the present work. It has a ester moiety like ferulone B ones with just three hydroxyl group. It can be point out that beside of sesquiterpen coumarins, ester coumarins may be attribute of *Ferula* genus, as well as. It was previously well known that sesquiterpene coumarins from *Ferula* species indicated various bioactivity like cytotoxic and anti-inflammatory properties (Nazari & Iranshahi 2011; Zarei et al. 2012; Kasaian et al. 2013; Kasaian et al. 2015). Although ester coumarins have not considerable antioxidant potential, It can be assumed that ester coumarins might be exhibit different biological activities. Further investigations are need to confirm the hypothesis.

3. Experimental

3.1. General procedure

PG instrument T80+ spectrometer was used to determine UV spectra of compounds in MeOH. NMR Spectra were obtained in a Bruker AVANCE 400 MHz NMR spectrometer (¹H: 400 and ¹³C: 100 MHz) in CDCl₃, and the residual solvent peaks were used as internal standard; HMBC spectra were optimized for a long-range J(H,C) of 9 Hz. A Finnigan MAT95 spectrometer was used to obtain mass spectra of the compounds.

3.2. Plant material

The roots of *Ferula orientalis* L. were collected from Khalkhal in the province of Ardabil, in May 2011, and a voucher specimen (voucher no. 1390-2) representing this collection has been deposited with the Herbarium of the Department of Biology, Faculty of Sciences, Mohaghegh Ardabili University, Iran.

3.3. Extraction and isolation

Dried ground roots of *F. orientalis* (150 g) were extracted sequentially with *n*-hexane, dichloromethane and methanol (MeOH), 500 mL each, using a Soxhlet apparatus. The extracts were dried under vacuum. The *n*-hexane extract (3 g) was subjected to Vacuum Liquid Chromatography (VLC) on silica gel, eluting with solvent mixtures of increasing

polarity: 100% *n*-hexane-ethyl acetate (EtOAc), 100% EtOAc and 100% MeOH, to yield a number of fractions, which upon initial Thin Layer Chromatographic (TLC) analyses, were grouped into 13 main fractions. A mixture of fractions 7 and 8 (50% EtOAc in *n*-hexane, and 60% EtOAc in *n*-hexane) was further analysed by preparative-TLC (mobile phase = 5% acetone in chloroform) to yield compound **2** (13.3 mg, R_f = 0.62, blue fluorescent). The fraction of 100% MeOH was further analysed by preparative-TLC (mobile phase 10% acetone in chloroform) to yield compound **1** (12.1 mg, R_f = 0.40, blue fluorescent). The structures of the isolated compounds were elucidated by spectroscopic means.

Ferulone A [7-*O*-(4,8,12,16-tetrahydroxy-4,8,12,16-tetramethyl-heptadecanoyl)-coumarin, **1**, 12.1 mg]. Yellow-brown oil. UV λ_{max} (MeOH) nm: 320, 270, 265, 259 and 252. IR ν_{max} (CHCl₃) cm⁻¹: 3437, 2925, 2853, 1734, 1563, 1428, 1248, 1125, 968, 893, 836, 486; ¹H NMR (400 MHz, CD₃OD): 6.26(1H, d, J = 9.5 Hz, H-3), 7.64(1H, d, J = 9.5 Hz, H-4), 7.37(1H, d, J = 8.5 Hz, H-5), 6.84(1H, d, J = 8.5, 2.0 Hz, H-6), 6.82(1H, d, J = 2, H-8), 2.28(2H, dd, J = 7.5, 4.5 Hz, H-2'), 1.11-2.29 (overlapes peaks, H-3', H-5', H-6', H-7', H-9', H-10', H-11', H-13', H-14', H-15'), 1.22(3H, s, H-4'-Me), 1.19(3H, s, H-8'-Me), 1.19(3H, s, H-12'-Me), 1.17(3H, s, H-16'-Me), 1.20(3H, s, H-16'-Me); ¹³C NMR (100 MHz, CD₃OD): 161.8 (C-2), 112.9 (C-3), 143.3 (C-4), 128.7 (C-5), 113.3 (C-6), 160.1 (C-7), 101.6 (C-8), 156.0 (C-9), 112.6 (C-10), 170.5 (C-1'), 43.6 (C-2'), 31.4 (C-3'), 73.7 (C-4'), 22.8 (C-4'-Me), 31.4 (C-5'), 28.2 (C-6'), 30.8 (C-7'), 73.7 (C-8'), 22.7 (C-8'-Me), 30.7 (C-9'), 24.2 (C-10'), 30.7 (C-11'), 73.6 (C-12'), 21.7 (C-12'-Me), 30.2 (C-13'), 24.2 (C-14'), 30.2 (C-15'), 67.7 (C-16'), 21.0, 18.0 (C-16'-Me), see also Table S1; ¹H-¹H Cosy, HMBC and HSQC correlations: see Table S2; HR-ESI-MS: 535.3270 ([M+H]⁺, C₃₀H₄₇O₈; calc. 535.3271).

Ferulone B [7-*O*-(4-hydroxy-4,8,12-trimethyl-trideca-7,11-dienoyl)-coumarin, **2**, 13.3 mg]. Brown oil. UV λ_{max} (MeOH) nm: 316, 260, 264, 257, 251 and 246 nm. IR ν_{max} (CHCl₃) cm⁻¹: 3448, 2924, 2958, 2851, 1737, 1568, 1187, 830, 719, 639, 518; 476; ¹H NMR (400 MHz, CD₃OD): 6.28(1H, d, J = 9.5 Hz, H-3), 7.66(1H, d, J = 9.5 Hz, H-4), 7.56(1H, d, J = 8.5 Hz, H-5), 7.15(1H, d, J = 8.5, 2.0 Hz, H-6), 7.02(1H, s, H-8), 2.27(2H, dd, J = 7.5, 4.5 Hz, H-2'), 1.20-2.29 (overlapes peaks, H-3', H-5', H-6', H-9', H-10'), 5.02(2H, bt, H-7'), 5.53(2H, bt, H-11'), 1.23(3H, s, H-4'-Me), 1.66(3H, s, H-8'-Me), 1.458(3H, s, H-12'-Me), 1.71(3H, s, H-12'-Me); ¹³C NMR (100 MHz, CD₃OD): 162.0 (C-2), 113.1 (C-3), 143.5 (C-4), 128.4 (C-5), 113.4 (C-6), 161.2 (C-7), 101.7 (C-8), 155.9 (C-9), 112.6 (C-10), 170.5 (C-1'), 43.7 (C-2'), 31.4 (C-3'), 68.2 (C-

4'), 22.9(C-4'-Me), 31.3(C-5'), 32.0(C-6'), 119.2(C-7'), 132.5(C-8'), 27.9(C-8'-Me), 30.6(C-9'), 32.3(C-10'), 124.5(C-11'), 131.6(C-12'), 28.9, 29.7(C-12'-Me), see also Table S1; ¹H-¹H Cosy, HMBC and HSQC correlations: see Table S2; HR-ESI-MS: 413.2326 ([M+H]⁺, C₂₅H₃₃O₅; calc. 413.2328).

3.4. Free-radical-scavenging activity (the DPPH assay)

1,1-Diphenyl-2-picrylhydrazyl (DPPH), molecular formula C₁₈H₁₂N₅O₆, was obtained from FlukaChemie AG, CH-Buchs. The method of Takao *et al.* (1994) was adopted with suitable modifications (Kumarasamy *et al.*, 2002; Chima *et al.*, 2014). A solution of DPPH (80 µg/mL) in MeOH was used. The test samples were dissolved in MeOH to obtain a concentration of 1 mg/mL. Ten-fold dilutions were conducted with the stock solutions of test compounds to obtain concentrations of 0.1, 0.01, 0.001, 0.0001 and 0.00001 mg/mL. Diluted solutions (1 mL each) were mixed with DPPH (1 mL) and allowed 30 min for any reaction to occur. The absorbance was recorded at 517 nm. The experiment was performed in triplicate, and average absorption was noted for each concentration. Data were processed using the EXCEL, and the concentration that caused a 50% reduction in DPPH absorbance at 517 nm (RC₅₀) was calculated. The same procedure was applied for the positive control, quercetin.

4. Conclusion

To the best of our knowledge, ferulone A and B, are new natural products of ester coumarin class. It has previously been demonstrated that coumarins display several biological activities (Murray *et al.* 1982; Razavi & Zarrini 2010; Razavi *et al.* 2010; Razavi 2011). Therefore, it is reasonable to assume that reported pharmacological properties of *F. orientalis* might be, at least partly, owing to presence of coumarins in its roots.

Acknowledgment

We thank EPSRC National Mass Spectrometry Service Centre (Department of Chemistry, University of Wales Swansea, Swansea, Wales, UK) for MS analyses.

References

- Alinezhad S, Kamalzadeh A, Shams-Ghahfarokhi M, Rezaee MB, Jaimand K, Kawachi M, Zamani Z, Tolouei R, Razzagji-Abyaneh M. 2011. Search for novel antifungals from 49 indigenous medicinal plants: *Foeniculumvulgare* and *Platycladusorientalis* as strong inhibitors of aflatoxin production by *Aspergillusparasiticus*. *Ann Microbiol.* 61: 673-81.
- Auzi AA, Gray AI, Saleem MM, Badwan AA, Sarker SD. 2008. Feruhermonins A-C: three daucane esters, from the seeds of *Ferula hermonis* (Apiaceae). *J Asian Nat Prod Res.* 10: 701-7.
- Bagheri SM, Sahebkar A, Gohar AR, Saeidnia S, Malmir M, Iranshahi M. 2010. Evaluation of cytotoxicity and anticonvulsant activity of some Iranian medicinal *Ferula* species. *Pharm Biol.* 48: 242-6.
- Bashir S, Alam M, Adhikari A, Shrestha RL, Yousuf S, Ahmad B, Parveen S, Aman A, Choudhury MI. 2014. New antileishmanialsesquiterpenecoumarins from *Ferula narthex* Boiss. *Phytochem Letts.* 9: 46-50.
- Chima NK, Nahar L, Majinda RRT, Celik S, Sarker SD. 2014. Assessment of free-radical scavenging activity of the extracts and fractions of *Gypsophila pilulifera*: Assay-guided isolation of the active component. *Braz J Pharmacog.* 24: 38-43.
- Davis PH. 1972. 'Flora of Turkey and the East Aegean Islands', Vol 4. Edinbuegh: Edinburgh University Press; p. 445-6.
- El- Razek MHA, Ohta S, Ahmed AA, Hitara T. 2001. Sesquiterpenecoumarins from the roots of *Ferula assa-foetida*. *Phytochemistry*, 58: 1289-95.
- Geroushi A, Auzi AA, Elhwuegi AS, Elzawam F, Elsherif A, Nahar L, Sarker SD. 2011 .Anti-inflammatory sesquiterpenes from the root oil of *Ferula hermonis*. *Phytother Res.* 25: 774-7.
- Geroushi A, Auzi AA, Elhwuegi AS, Elzawam F, Elsherif A, Nahar L, Sarker SD. . Antinociceptive and anti-inflammatory activity of *Ferula hermonis* root oil in experimental animals. *Latin Am J Pharmacy.* 29: 1436-9.
- Iranshahy M, Iranshahi, M. 2011. Traditional uses, phytochemistry and pharmacology of asafoetida (*Ferula assa-foetida* oleo-gum-resin)- a review. *J Ethnopharmacol.* 134: 1-10. DOI: 10.1016/j.jep.2010.11.067.

- Iranshahi M, Kalategi F, Rezaee R, Shahverdi AR, Ito C, Tokuda H, Itoigawa M. 2008. Cancer chemopreventive activity of terpenoid coumarins from *Ferula* species. *Planta Med.* 74: 147-150.
- Iranshahi M, Kalategi F, Sahebkar A, Sardashti A, Schneider B. 2010. New sesquiterpenecoumarins from the roots of *Ferula flabelliloba*. *Pharm Biol.* 48:217-20.
- Iranshahi M, Masull M, Asili A, Hamedzadeh A, Jahanbin B, Festa M, Capasso A, Piacente S. 2010. Sesquiterpene Coumarins from *Ferula gumosa*. *J Nat Prod.* 73:1958-62.
- Kartal N, Sokmen M, Tepe B, Daferera D, Polissiou M, Sokmen A. 2007. Investigation of the antioxidant properties of *Ferula orientalis* L. using a suitable extraction procedure. *Food Chem.* 100: 584-9.
- Kasaian J, Iranshahi M, Masullo M, Piacente S, Ebrahimi F, Iranshahi M. 2014. Sesquiterpene lactones from *Ferula oopoda* and their cytotoxic properties. *J. Asian Nat Prod Res.* 16: 248-253. DOI: 10.1080/10286020.2013.866099.
- Kasaian J, Mosaffa F, Behravan J, Masollu M, Piacente S, Ghandadi M, Iranshahi M. 2015. Reversal of P-glycoprotein-mediated multidrug resistance in MCF-7/Adr cancer cells by sesquiterpene coumarins. *Fitoter.* 103: 149-154. DOI: 10.1016/j.fitote.2015.03.025.
- Kaur P, Kumar M, Singh B, Kumar S, Kaur S. 2012. Amelioration of oxidative stress induced by oxidative mutagens and COX-2 inhibitory activity of umbelliferone isolated from *Glycyrrhizaglabra* L. *Asian Pac J Trop Biomed.* S:120-6.
- Kumarasamy, Fergusson ME, Nahar L, Sarker SD. 2002. Biological activity of moschamindole from *Centaureamoschata*. *Pharm Biol.* 40: 307-10.
- Miski M, Mabry TJ, Saya O. 1987. Newdaucane and germacrane esters from *Ferula orientalis* var. *orientalis*. *J Nat Prod.* 50: 829-34.
- Mozaffarian V. 2003. *A Dictionary of Iranian Plant Names*. Tehran: Farhang e Moaser publication.
- Nazari ZE, Iranshahi M. 2011. Biologically active sesquiterpene coumarins from *Ferula* species. *Phytoterap Res.* 25: 315-323. DOI: 10.1002/ptr.3311.
- Murray RDH, Mendez J, Brown SA. 1982. *The natural coumarins: occurrence, chemistry and biochemistry*. Chichester: John Wiley and Sons Ltd.
- Razavi SM. 2011. Plant coumarins as allelopathic agents. *Int J Biol Chem.* 5: 86-90.
- Razavi, S.M., Janani, M. 2015. A new ester coumarin from *Ferula Persica* wild, indigenous to Iran. *Nat Prod Res.* 29: 717-721.

- Razavi SM, Zarrini G. 2010. Bioactivity of aviprin and aviprin-3aEuro(3)-*O*-glucoside, two linear furanocoumarins from Apiaceae. *Russ J Biorg Chem.* 36: 359-62.
- Razavi SM, Zahri S, Motamed Z, Ghasemi G. 2010. Bioscreening of oxypeucedanin, a known furanocoumarin. *Iran J Basic Med Sci.* 13: 133-8.
- Sghari J, Atabaki V, Bahar E, Mazaheritehrani M. 2015. Identification of sesquiterpene coumarins of oleo-gum resin of *Ferula assa-foetida* L. from the Yasuj region. *Nat Prod Res.* 2: 1-4. DOI: 10.1080/14786419.20.2015.
- Takao T, Kitatani F, Watanabe N, Yagi I, Sakata K. 1994. A simple screening method for antioxidants and isolation of several antioxidants produced by marine bacteria from fish and shellfish. *Biosci Biotechnol Biochem.* 58: 1780-3.
- Zarei H, Rezaee R, Behravan E, Soltani F, Mosaffa F, Iranshahi M, Behravan J. 2012. Diversin, from *Ferula diversivittata* protects human lymphocytes against oxidative stress induced by H₂O₂. *Nat Prod Res.* 27: 1016-1019.





Figure 1. Structures of umbelliferone (3) and its esters (1 and 2)

Figure1. Structures of compound 1 and 2