

Test purchase, synthesis and characterization of 2-methoxydiphenidine (MXP) and differentiation from its *meta*- and *para*-substituted isomers

Gavin McLaughlin,^{a,b} Noreen Morris,^a Pierce V. Kavanagh,^b John D. Power,^b John O'Brien,^c Brian Talbot,^d Simon P. Elliott,^e Jason Wallach,^f Khoa Hoang,^g Hamilton Morris,^h Simon D. Brandt^{i*}

^a Department of Life and Physical Sciences, School of Science, Athlone Institute of Technology, Dublin Road, Westmeath, Ireland

^b Department of Pharmacology and Therapeutics, School of Medicine, Trinity Centre for Health Sciences, St. James Hospital, Dublin 8, Ireland

^c School of Chemistry, Trinity College, Dublin 2, Ireland

^d School of Pharmacy and Pharmaceutical Sciences, Trinity College, Dublin 2, Ireland

^e ROAR Forensics, Malvern Hills Science Park, Geraldine Road, WR14 3SZ, UK


^f Department of Pharmaceutical Sciences, Philadelphia College of Pharmacy, University of the Sciences, Philadelphia, PA 19104, USA

^g Department of Chemistry and Biochemistry, University of the Sciences, Philadelphia, PA 19104, USA


^h The New School for Social Research, Department of Anthropology, 66 West 12th Street, NY 10011, New York, USA

ⁱ School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, Liverpool L3 3AF, UK


*Correspondence to: Simon D. Brandt, School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, Liverpool, L3 3AF, UK. E-Mail: s.brandt@ljmu.ac.uk



2-MXP
(2-MeO-diphenidine)



3-MXP



4-MXP

Contents:

Page 2: Example of an MXP product obtained from an online vendor

Page 3: GC-El-quadrupole mass spectra obtained from all three synthesized MXP isomers

Page 4: HPLC-SIM-MS traces of powdered 2-MXP samples vs. synthesized standards

Page 5: ESI-QqQ-MS/MS of synthesized MXP isomers

Page 6: ESI-QTOF-MS/MS of synthesized MXP isomers (via UHPLC)



Page 7: TLC analysis of powdered 2-MXP samples vs. synthesized standards

Page 8: Matrix assisted inlet ionization mass spectra of 2-MXP tablet vs. 2-MXP standard


Page 9: ^1H -NMR of 2-MXP extracted from tablet vs. MXP isomers

Page 10: ^{13}C -NMR of 2-MXP extracted from tablet vs. MXP isomers

Pages 11 - 13: ATR-FT-IR spectra of three MXP isomers







Drug Testing and Analysis – McLaughlin *et al.* – Supplementary data



GC-EI-quadrupole mass spectra obtained from all three synthesized MXP isomers

Agilent 6980 GC coupled to an Agilent 5973 MSD (HP-5ms column, 30 m x 0.25 mm x 0.25 μ m). Helium carrier gas at a constant flow of 1 mL/min in splitless mode. Injection port and transfer line set at 250 °C and 280 °C. Oven temperature: 40 °C held for 1 min, ramped at 12 °C/min to 280 °C, held for 5 minutes, then ramped again at 20 °C/min to 300 °C and held for 3 min. The total run time was 30 min.


Drug Testing and Analysis – McLaughlin *et al.* – Supplementary data




HPLC-SIM-MS traces of powdered 2-MXP sample s vs. synthesized standards

Drug Testing and Analysis – McLaughlin *et al.* – Supplementary data


2-MXP MRM R 68 (0.686) Cn (Cen,2, 80.00, Ht); Cm (1:197)



3-MXP MRM R 133 (1.342) Cn (Cen,2, 80.00, Ht); Sm (SG, 2x1.00); Cm (1:197)






4-MXP MRM R 137 (1.382) Cn (Cen,2, 80.00, Ht); Sm (SG, 2x1.00); Cm (2:198)




ESI-QqQ-MS/MS of synthesized MXP isomers



Drug Testing and Analysis – McLaughlin *et al.* – Supplementary data




UHPLC-ESI-QTOF-MS/MS of synthesized MXP isomers




TLC analysis of powdered 2-MXP samples vs. synthesized standards



Matrix assisted inlet ionization mass spectra of 2-MXP tablet vs. 2-MXP standard





¹³C-NMR of 2-MXP extracted from tablet vs. MXP isomers

