

1  
2  
3 **Quantification of seasonal long physical load in soccer**  
4 **players with different starting status from the English**  
5 **Premier League: implications for maintaining squad**  
6 **physical fitness**  
7

8 Liam Anderson<sup>1,2</sup>, Patrick Orme<sup>1,2</sup>, Rocco Di Michele<sup>3</sup>,  
9 Graeme L Close<sup>2</sup>, Jordan Milsom<sup>1</sup>, Ryland Morgans<sup>1</sup>, Barry  
10 Drust<sup>1,2</sup> and James P Morton<sup>2</sup>  
11  
12

13 <sup>1</sup>Liverpool Football Club  
14 Melwood Training Ground  
15 Deysbrook Lane  
16 Liverpool  
17 L12 8SY  
18 UK  
19

20 <sup>2</sup>Research Institute for Sport and Exercise Sciences  
21 Liverpool John Moores University  
22 Tom Reilly Building  
23 Byrom St Campus  
24 Liverpool  
25 L3 3AF  
26 UK  
27

28 <sup>3</sup>Department of Biomedical and Neuromotor Sciences  
29 University of Bologna  
30 Bologna  
31 Italy  
32

33 **Running head:** Starting status and seasonal workload in soccer  
34

35 **Address for correspondence:**

36 Dr James Morton  
37 Research Institute for Sport and Exercise Sciences  
38 Liverpool John Moores University  
39 Tom Reilly Building  
40 Byrom St Campus  
41 Liverpool  
42 L3 3AF  
43 United Kingdom  
44 Email: [J.P.Morton@ljmu.ac.uk](mailto:J.P.Morton@ljmu.ac.uk)  
45 Tel: +44 151 904 6233

## 46    **Abstract**

47    **Purpose.** To quantify the accumulative training and match load  
48    during an annual season in English Premier League soccer  
49    players classified as starters (n=8, started  $\geq 60\%$  of games),  
50    fringe players (n=7, started 30-60% of games) and non-starters  
51    (n=4, started  $<30\%$  of games). **Methods.** Players were  
52    monitored during all training sessions and games completed in  
53    the 2013-2014 season with load quantified using GPS and  
54    Prozone technology, respectively. **Results.** When including  
55    both training and matches, total duration of activity ( $10678 \pm$   
56     $916$ ,  $9955 \pm 947$ ,  $10136 \pm 847$  min;  $P=0.50$ ) and distance  
57    covered ( $816.2 \pm 92.5$ ,  $733.8 \pm 99.4$ ,  $691.2 \pm 71.5$  km;  $P=0.16$ )  
58    was not different between starters, fringe and non-starters,  
59    respectively. However, starters completed more (all  $P<0.01$ )  
60    distance running at 14.4-19.8 km/h ( $91.8 \pm 16.3$  v  $58.0 \pm 3.9$   
61    km;  $ES=2.5$ ), high speed running at 19.9-25.1 km/h ( $35.0 \pm 8.2$   
62    v  $18.6 \pm 4.3$  km;  $ES=2.3$ ) and sprinting at  $>25.2$  km/h ( $11.2 \pm$   
63     $4.2$ , v  $2.9 \pm 1.2$  km;  $ES=2.3$ ) than non-starters. Additionally,  
64    starters also completed more sprinting ( $P<0.01$ .  $ES=2.0$ ) than  
65    fringe players who accumulated  $4.5 \pm 1.8$  km. Such differences  
66    in total high-intensity physical work done were reflective of  
67    differences in actual game time between playing groups as  
68    opposed to differences in high-intensity loading patterns during  
69    training sessions. **Conclusions.** Unlike total seasonal volume of  
70    training (i.e. total distance and duration), seasonal high-  
71    intensity loading patterns are dependent on players' match  
72    starting status thereby having potential implications for training  
73    programme design.

74    **Key Words:** GPS, Prozone, high-intensity zones, training load

75

76

77

78

79

80

81

82

83

84

85

## 86    **Introduction**

87    Soccer match play is characterized by brief bouts of high-  
88    intensity linear and multidirectional activity interspersed with  
89    longer recovery periods of lower intensity.<sup>1</sup> Elite players  
90    typically cover 10-14 km in total distance per game.<sup>2-6</sup> where  
91    both high intensity (speeds > 14.4 km · h<sup>-1</sup>) and very high-  
92    intensity running distance (speeds > 19.8 km · h<sup>-1</sup>) contribute  
93    ~25 and ~8% of the total distance covered, respectively.<sup>7,8</sup> Top-  
94    class soccer players also perform 150-250 intense actions per  
95    game<sup>9</sup> and complete a very high-intensity run approximately  
96    every 72 s.<sup>8</sup>

97            In order to successfully meet these demands, the  
98    physical preparation of elite players has become an  
99    indispensable part of the professional game, with high fitness  
100   levels required to cope with the ever-increasing demands of  
101   match play.<sup>10,11</sup> Nonetheless, despite nearly four decades of  
102   research examining the physical demands of soccer match  
103   play,<sup>12</sup> the quantification of the customary training loads  
104   completed by elite professional soccer players are not currently  
105   well known. For players of the English Premier League, such  
106   reports are limited to a 4-week winter fixture schedule,<sup>13</sup> a 10-  
107   week period,<sup>14</sup> seasonal long analysis<sup>15</sup> and most recently, an  
108   examination of the effects of match frequency in a weekly  
109   microcycle.<sup>16</sup> It is noteworthy that the absolute physical loads  
110   of total distance (e.g. < 7 km), high intensity distance (e.g. <  
111   600 m) and very high intensity distance (e.g. < 400 m)  
112   collectively reported in these studies do not near recreate those  
113   completed in matches. As such, although the typical current  
114   training practices of professional players may be sufficient in  
115   order to promote recovery and readiness for the next game  
116   (thus reducing risk of over-training and injury), it could also be  
117   suggested that it is the participation in match play itself that is  
118   the most appropriate stimulus for preparing players for the  
119   physical demands of match play. This point is especially  
120   relevant considering previous evidence demonstrating  
121   significant positive correlations between individual in season  
122   playing time and aspects of physical performance including  
123   sprint performance and muscle strength.<sup>17</sup>

124            Such differences between match and training load can  
125   be particularly challenging for fitness and conditioning staff  
126   given that players in a first team squad are likely to receive  
127   different loading patterns, depending on whether they regularly  
128   start matches or not. In this way, discrepancies in physical  
129   loads between players could lead to differences in important  
130   components of soccer-specific fitness which may subsequently  
131   present itself on match day when players not accustomed to  
132   match loads are now required to complete the habitual physical  
133   loads performed by regular starting players. The challenge of

134 maintaining squad physical fitness is also technically difficult,  
135 given both organisational and traditional training practices  
136 inherent to professional soccer. For example, in the English  
137 Premier League, it is not permitted for players to train on the  
138 same pitch where the game was played for >15 minutes post-  
139 match. Furthermore, it is often common practice for the entire  
140 playing squad to be given 1-2 days of recovery following each  
141 game (consisting of complete inactivity or light recovery  
142 activities only), especially in those instances where the fixture  
143 schedule consists of the traditional Saturday-to-Saturday  
144 schedule.<sup>16</sup>

145         With this in mind, the aim of the present study was to  
146 quantify the accumulative training and match load (hence total  
147 accumulative physical load) across an annual season in those  
148 players considered as regular starters, fringe players and non-  
149 starters. To this end, we monitored outfield players from the  
150 English Premier League (who competed in the 2013-2014  
151 season) who were classified as starters (starting  $\geq 60\%$  of  
152 games), fringe players (starting 30-60% of games) and non-  
153 starters (starting  $< 30\%$  of games). We specifically hypothesised  
154 that both fringe and non-starting players would complete  
155 significantly less total physical load (especially in high-  
156 intensity zones) than starting players, thereby providing  
157 practical applications for the development of soccer-specific  
158 conditioning programme designed to maintain squad physical  
159 fitness.

160

## 161 **Methods**

### 162 **Subjects**

163 Nineteen professional outfield soccer players from an English  
164 Premier League team (mean  $\pm$  SD: age  $25 \pm 4$  years, body mass  
165  $79.5 \pm 7.8$  kg, height  $180.4 \pm 6.4$  cm) took part in the study.  
166 When quantifying data from the entire “in-season analysis”  
167 there were 8 starters (mean  $\pm$  SD: age  $25 \pm 5$  years, body mass  
168  $80.6 \pm 8.3$  kg, height  $178.8 \pm 6.3$  cm), 7 fringe (mean  $\pm$  SD: age  
169  $26 \pm 4$  years, body mass  $79.7 \pm 7.4$  kg, height  $181.0 \pm 7.3$  cm)  
170 and 4 non-starters (mean  $\pm$  SD: age  $23 \pm 3$  years, body mass  
171  $74.5$  kg, height  $181.5 \pm 6.9$  cm). Players with different position  
172 on the field were tested: 5 wide defenders, 4 central defenders,  
173 6 central midfielders, 2 wide midfielders and 3 attackers. Long-  
174 term injuries were excluded from this study if they were absent  
175 for on field training for duration  $> 4$  weeks. The study was  
176 conducted according to the requirements of the Declaration of  
177 Helsinki and was approved by the university ethics committee  
178 of Liverpool John Moores University.

179

## 180     **Design**

181     Training and match data were collected over a 39-week period  
182     during the 2013-2014 competitive season from August 2013  
183     until May 2014. The team used for data collection competed in  
184     3 official domestic competitions across the season. For the  
185     purposes of this current study, training sessions included for  
186     analysis consisted of all of the ‘on pitch’ training each player  
187     was scheduled to undertake. Sessions that were included in the  
188     analysis were team training sessions, individual training  
189     sessions, recovery sessions and rehabilitation training sessions.  
190     A total number of 181 team-training sessions (2182 individual),  
191     159 rehab sessions (213 individual), 28 recovery sessions (179  
192     individual), 43 competitive matches including substitute  
193     appearances (531 individual) and 12 non-competitive games  
194     including substitute appearances (33 individual) were observed  
195     during this investigation. All data reported are for outdoor field  
196     based sessions only. We can confirm that in the season of  
197     analysis, the players studied did not do any additional aerobic /  
198     high-intensity conditioning in the gym or an indoor facility.  
199     However, all players did complete 1-3 optional gym based  
200     sessions per week (typically consisting of 20-30 minute long  
201     sessions comprising upper and/or lower body strength based  
202     exercises). When expressed as ‘total time’ engaged in training  
203     activities (i.e. also inclusive of gym training) and games, the  
204     data presented in the present paper therefore represent  $78\pm10$ ,  
205      $79\pm6$  and  $86\pm7\%$  of ‘total time’ for starters, fringe players and  
206     non-starters, respectively. This study did not influence or alter  
207     any session or game in any way nor did it influence the  
208     inclusion of players in training sessions and/or games. Training  
209     and match data collection for this study was carried out at the  
210     soccer club’s outdoor training pitches and both home and away  
211     grounds in the English Football League, respectively.

212             The season was analyzed both as a whole and in 5  
213     different in-season periods consisting of 4x8 weeks (periods 1-  
214     4) and 1x7 week period (period 5). Players were split into 3  
215     groups for the entire in season analysis and individually for  
216     each in season period. The 3 groups consisted of “starters”,  
217     “fringe” and “non-starters” and were split based on the  
218     percentage of games started for the entire in season ( $n=8$ , 7 and  
219     4, respectively) and during the individual period 1 ( $n=8$ , 5 and  
220     6, respectively), period 2 ( $n=9$ , 5 and 5, respectively), period 3  
221     ( $n=6$ , 8 and 5, respectively), period 4 ( $n=8$ , 5 and 6,  
222     respectively) and period 5 ( $n=11$ , 2 and 6, respectively).  
223     Starting players started  $\geq 60\%$  competitive games, fringe  
224     players started 30-60% of games and non-starting players  
225     started  $<30\%$  of games. The first day of data collection period  
226     began in the week commencing (Monday) of the first Premier  
227     League game (Saturday) and the last period ended after the  
228     final Premier League game. Data for the entire in season and

each individual period was further divided into training and matches. As outlined previously, training consisted of all ‘on pitch’ training sessions that were organised and planned by the clubs coaches and staff and match data consisted of both competitive and non-competitive games. No data from training or games from when players were on International camps were collected.

## **Methodology**

Players’ physical activity during each training, rehabilitation, recovery sessions and non-competitive game was monitored using portable global positioning system (GPS) units (Viper pod 2, STATSports, Belfast, UK). This device provides position velocity and distance data at 10 Hz. Each player wore the device across the upper back between the left and right scapula inside a custom made vest supplied by the manufacturer. This position on the player allows the GPS antenna to be exposed for a clear satellite reception. This type of system has previously been shown to provide valid and reliable estimates of some of the movements related to soccer, although it should be noted that fast, more instantaneous, and more multidirectional movements are measured less accurately.<sup>18-21</sup> All devices were activated 30-minutes before data collection to allow acquisition of satellite signals, and synchronize the GPS clock with the satellite’s atomic clock.<sup>22</sup> Following each training session, GPS data were downloaded using the respective software package (Viper PSA software, STATSports, Belfast, UK) and were clipped to involve the “main” organised session i.e. the beginning of the warm up to the end of the last organized drill for each player, the initiation of exercise to the cessation of exercise on individual training, recovery and rehab sessions or the start of the game until the end of the game with any distances and times covered and undergone during the half-time period removed. In order to avoid inter-unit error, players wore the same GPS device for each training sessions.<sup>23,24</sup>

Players’ match data were examined using a computerized semi-automatic video match-analysis image recognition system (Prozone Sports Ltd®, Leeds, UK) and were collected using the same methods as Bradley et al.<sup>8</sup> This system has previously been independently validated to verify the capture process and subsequent accuracy of the data.<sup>25</sup>

Variables that were selected for analysis included duration, total distance and 3 different speed categories that were divided into the following thresholds: running (14.4-19.7

275 km · h<sup>-1</sup>), high-speed running (19.8-25.1 km · h<sup>-1</sup>), and sprinting  
 276 (>25.1 km · h<sup>-1</sup>). High-intensity running consists of running,  
 277 high-speed running and sprinting (running speed >14.4 km · h<sup>-1</sup>  
 278 1). Very high-intensity running consists of high-speed running  
 279 and sprinting (running speed > 19.8 km · h<sup>-1</sup>). The speed  
 280 thresholds for each category are similar to those reported  
 281 previously in match analysis research<sup>7,8</sup> and are commonly used  
 282 day to day in professional soccer clubs.

283

284

## 285 **Statistical Analysis**

286 All of the data are presented as mean ± standard deviation  
 287 (SD). Data were analysed using between-group one-way  
 288 ANOVAs for independent samples. When the F-test was  
 289 significant (p<0.05), post-hoc pairwise comparisons were  
 290 performed, in which the significance level was adjusted to  
 291 0.017 (Bonferroni correction). Cohen's d indices were  
 292 calculated for all pairwise differences to determine an effect  
 293 size (ES). The absolute ES value was evaluated according to  
 294 the following thresholds: < 0.2 = trivial, 0.2-0.6 = small, 0.7-  
 295 1.2 = moderate, 1.3-2.0 = large, and > 2.0 = very large.

296

## 297 **Results**

### 298 **Seasonal long comparison of “total” physical load**

299 A comparison of seasonal physical load (inclusive of both  
 300 training and matches) is presented in Table 1. Although there  
 301 was no significant difference in total duration (P=0.502) and  
 302 distance covered (P=0.164) between player categories, non-  
 303 starters completed significantly less running (P=0.002;  
 304 ES=2.5), high-speed running (P=0.004; ES=2.3) and sprinting  
 305 (P=0.003; ES=2.3) than starters. Additionally, fringe players  
 306 completed significantly less sprinting than starters (P=0.002;  
 307 ES=2.0) though no differences were apparent in running  
 308 (P=0.062) and high-speed running (P=0.038) between these  
 309 groups.

### 310 **Seasonal long comparison of total “training” and “match” 311 physical load**

312 A comparison of seasonal long training and match load is  
 313 presented in Figure 1A and B (for duration and total distance).  
 314 In relation to matches, both fringe and non-starters completed  
 315 less duration of activity (both P<0.01; ES=2.7 and 5.7,  
 316 respectively) and total distance (both P<0.01; ES=5.4 and 2.5,  
 317 respectively) compared with starters. Additionally, non-starters

318 also completed less duration ( $P=0.001$ ;  $ES=0.7$ ) and total  
319 distance than fringe players ( $P=0.001$ ;  $ES=0.7$ ). In relation to  
320 training, differences were only apparent between non-starters  
321 and starters where non-starters spent longer time training  
322 ( $P=0.003$ ;  $ES=2.4$ ) and covered greater total distance ( $P=0.003$ ;  
323  $ES=2.3$ ).

#### 324 **Seasonal long comparison of “training” and “match”** 325 **physical load in high-intensity speed zones**

326 Seasonal long distance covered in running, high-speed running  
327 and sprinting in both training and matches is displayed in  
328 Figure 2A-C. In relation to matches, both fringe and non-  
329 starters completed significantly less distance in running (both  
330  $P<0.01$ ;  $ES=1.7$  and  $4.0$ , respectively), high-speed running  
331 (both  $P<0.01$ ;  $ES=2.0$  and  $3.4$ , respectively) and sprinting (both  
332  $P<0.01$ ;  $ES=2.2$  and  $2.6$ , respectively) compared with starters.  
333 In addition, fringe players covered significantly more distance  
334 in running than non-starters ( $P=0.008$ ;  $ES=0.7$ ). However, no  
335 differences were apparent between fringe and non-starters for  
336 high-speed running and sprinting ( $P=0.026$  and  $0.045$ ;  $ES=0.7$   
337 and  $0.5$ , respectively). In contrast to match load, no differences  
338 were observed between groups for distance completed in  
339 running, high-speed running and sprinting during training  
340 ( $P=0.297$ ,  $0.658$  and  $0.802$ , respectively).

#### 341 **Comparison of “total” physical load within specific in-** 342 **season periods**

343 Total duration, total distance and distance completed in high-  
344 intensity speed zones within 5 in-season periods of the season  
345 are presented in Table 2. For duration of total activity,  
346 significant differences were only observed in periods 4  
347 ( $P=0.004$ ;  $ES=1.9$ ) and 5 ( $P=0.001$ ;  $ES=2.2$ ) where non-starters  
348 completed less total duration of activity than starters,  
349 respectively. Similarly, non-starters also completed less total  
350 distance than starters in periods 3-5 (all  $P<0.01$ , respectively;  
351  $ES=1.9$ ,  $3.1$  and  $3.4$ , respectively), less running in periods 1, 3,  
352 4 and 5 (all  $P<0.01$ , respectively;  $ES=1.0$ ,  $2.3$ ,  $3.6$  and  $3.6$ ,  
353 respectively), less high-speed running in periods 3-5 (all  
354  $P<0.01$ , respectively;  $ES=2.1$ ,  $2.6$  and  $3.0$ , respectively) and  
355 less sprinting in periods 2-5 (all  $P<0.01$ , respectively;  $ES=1.6$ ,  
356  $2.5$ ,  $3.0$  and  $2.5$ , respectively). Furthermore, starters completed  
357 more sprinting distance than fringe in periods 3 and 4 (both  
358  $P<0.01$ , respectively;  $ES=2.2$  and  $1.6$ , respectively) but fringe  
359 only differed from non-starters in period 4 only where they  
360 completed more sprinting ( $P=0.006$ ;  $ES=1.2$ ).

361

362



## **Comparison of “training” and “match” physical load within specific in-season periods**

Duration of activity, total distance, running, high-speed running and sprinting in matches are displayed in Figure 3A-E. As expected, in periods 1-5, starters had higher duration and than both non-starters (all  $P<0.01$ ; ES=2.7, 2.6, 13.2, 11.9 and 5.6, respectively) and fringe (all  $P<0.01$ ; ES=1.9, 1.6, 4.0, 5.5 and 2.5, respectively) whilst fringe players also exhibited higher durations than non-starters in periods 3-5 (all  $P<0.01$ ; ES=0.9, 1.3 and 2.3). Similarly, starters covered higher total distances in periods 1-5 than both non-starters (all  $P<0.01$ ; ES=2.6, 2.5, 9.5, 12.8 and 5.9, respectively) and fringe (all  $P<0.01$ ; ES=1.9, 1.6, 3.0, 5.1 and 2.4, respectively) and fringe players covered higher total distances than non-starters in periods 3-5 (all  $P<0.01$ ; ES=0.9, 1.3 and 2.3, respectively).

In relation to specific speed zones, starters completed more running in periods 1-5 than non-starters (all  $P<0.01$ ; ES=2.2, 2.1, 5.1, 7.2 and 4.7, respectively), more high-speed running in periods 1-5 (all  $P<0.01$ ; ES=1.8, 1.9, 3.5, 5.5 and 3.8) and more sprinting in periods 2-5 (all  $P<0.01$ ; ES=1.7, 2.8, 3.2 and 2.5). Moreover, starters completed more running than fringe players in periods 3 ( $P=0.009$ ; ES=1.7) and 4 ( $P=0.001$ ; ES=2.6), more high-speed running in periods 3 ( $P=0.003$ ; ES=2.0) and 4 ( $P=0.004$ ; ES=2.1) and more sprinting in periods 3 ( $P=0.001$ ; ES=2.2) and 4 ( $P=0.012$ ; ES=1.7). Fringe players also covered more running distance in periods 3-5 (all  $P<0.01$ ; ES=0.9, 1.3 and 2.3, respectively), more high-speed running in periods 4 ( $P=0.002$ ; ES=1.3) and 5 ( $P=0.008$ ; ES=2.2) and more sprinting in period 4 ( $P=0.003$ ; ES=1.3) than non-starters.

Duration of activity, total distance, running, high-speed running and sprinting in training are displayed in Figure 4A-E. In contrast to matches, total duration of activity was only different in period 3 ( $P=0.014$ ; ES=1.8) where non-starters trained for longer durations than starters. In addition, starters completed less total distance in periods 3 and 4 compared to non-starters (both  $P<0.01$ ; ES=2.5, 1.8, respectively) and non-starters also covered more total distance in period 3 than fringe players ( $P=0.007$ ; ES=0.4). Non-starters also covered more running than starters and fringe players in period 3 (both  $P<0.01$ ; ES=2.1 and 0.6, respectively) and more high-speed running than starters in period 4 ( $P=0.015$ ; ES=1.5). Finally, no differences were apparent between groups for sprinting during periods 1-5 ( $P=0.506, 0.361, 0.605, 0.521$  and  $0.487$ ).

## 409 Discussion

410 The aim of the present study was to quantify the accumulative  
411 training and match load (and total accumulative physical load)  
412 during an annual season in those players considered as regular  
413 starters, fringe players and non-starters. Contrary to our  
414 hypothesis, we observed that starting status had no effect on the  
415 apparent total volume completed, as reflected by total duration  
416 of activity and total distance covered during the season.  
417 Perhaps more important, however, was the observation of  
418 significant differences in the pattern of activity completed  
419 within specific high-intensity speed zones. In this regard, we  
420 report that starters generally completed more distance in  
421 running, high-speed running and sprinting zones than both  
422 fringe and non-starting players. This effect was largely due to  
423 differences in game time between groups as opposed to  
424 differences in training loading patterns. Given the role of  
425 training intensity in promoting soccer-specific fitness,<sup>10, 26-28</sup>  
426 our data therefore suggest that the training practices of those  
427 players not deemed to be receiving appropriate game time  
428 should be altered to include more emphasis on recreating the  
429 high-intensity demands of match play, so as to potentially  
430 maintain overall squad fitness, game readiness and reduce  
431 injury risk.

432 To the authors' knowledge, this is the first study to  
433 report seasonal long physical loads completed by elite  
434 professional soccer players. In our seasonal long accumulation  
435 analysis, we observed no evidence of starting status affecting  
436 total duration of activity or total distance covered across the  
437 entire in-season period (see Table 1). For example, total  
438 duration and total distance were similar in starters, fringe and  
439 non-starters. These distances are substantially higher (e.g.  
440 approximately 400 km) than that observed in a competitive in-  
441 season in other team sports such as Australian Football<sup>29</sup> likely  
442 due to shorter seasons in the latter i.e. 22 weeks (18 weeks in  
443 the study) versus 39 weeks in the English Premier League.

444 Although we observed no differences in the seasonal  
445 long profile between groups (i.e. duration and total distance  
446 covered), the proportion of this volume made up from training  
447 and game is, as expected, significantly different between  
448 groups. For example, in relation to training, starters displayed  
449 lower duration and total distances than non-starters but not  
450 fringe players. This fact is, of course, due to the fact that  
451 starting players engage in "recovery" training activities and  
452 days after games as opposed to traditional training sessions.<sup>13,16</sup>  
453 When quantifying match load, however, starters displayed  
454 higher duration and total distance than both fringe players and  
455 non-starters. Given the obvious difference between the physical  
456 and physiological demands between training and matches,<sup>13,16</sup>

457 such data could potentially suggest that the long-term  
458 physiological adaptations arising within these playing groups  
459 are likely very different. This point is especially apparent when  
460 considering the large discrepancy between intensity specific  
461 physical loads between groups. For example, starters covered  
462 higher distances in running and high-speed running speed  
463 zones, respectively, when compared with non-starters, but not  
464 fringe players (see Table 1). In addition, seasonal long distance  
465 covered whilst sprinting was also higher in starters compared to  
466 both fringe players and non-starters. As such, these data  
467 demonstrate that although players are able to maintain similar  
468 volume across the in-season period, distance covered in high-  
469 intensity zones is considerably greater in starters.

470         The differences in high-intensity loading patterns  
471 between groups is also especially relevant when considering  
472 that such differences were not due to alterations in training  
473 loads but rather, merely due to starters engaging in the high-  
474 intensity activity associated with match play. Indeed, we  
475 observed no difference in running, high-speed running and  
476 sprinting in training *per se* between starters, fringe players and  
477 non-starters. In contrast, starters displayed higher distance in  
478 matches when running, high-speed running and sprinting  
479 compared to fringe and non-starters (see Figure 2A-C). Such  
480 data clearly highlight that it is the participation in match play  
481 *per se* which represents the most appropriate opportunity to  
482 achieve high-intensity loading patterns. The practical  
483 implications of such discrepancies are important for designing  
484 training programmes to maintain overall squad physical fitness  
485 and game readiness. Indeed, the distances covered at these  
486 speeds during games display strong associations to physical  
487 capacity<sup>30,31</sup> and thus, players not consistently exposed to such  
488 stimuli during the season may eventually display de-training  
489 effects when compared to that displayed in the pre-season  
490 period.<sup>10,17</sup> Indeed, completion of high-intensity activity (even  
491 at the expense of total physical load done) is both sufficient and  
492 necessary to activate the molecular pathways that regulate  
493 skeletal muscle adaptations related to both aerobic<sup>32,33</sup> and  
494 anaerobic<sup>34</sup> performance. Additionally, when those players  
495 classified as fringe or non-starters are then required to start  
496 games, a potential for injury also exists due to the necessity to  
497 complete uncustomary loading patterns.<sup>35</sup>

498         In addition to the seasonal long physical loads, we also  
499 quantified the training and match load within 5 discrete periods  
500 of the in-season period. In this analysis, we observed that  
501 variations in physical load between groups were especially  
502 evident in periods 3, 4 and 5, an effect that was especially  
503 apparent between starters and both non-starters and fringe  
504 players for total duration, total distance and total zone 6 activity  
505 (i.e. sprinting). Similar to the seasonal long analysis, these

506 differences between groups were also largely reflective of  
507 differences in game time as opposed to training time. Such  
508 differences in loading within specific in-season periods are  
509 likely due to tactical and technical differences associated with  
510 specific fixture schedules. For example, in the present study,  
511 period 3 was the winter fixture schedule<sup>13</sup> whereas periods 4  
512 and 5 were reflective of a period where the team under  
513 investigation were challenging for domestic honours. In all of  
514 these periods, the management and coaching staff displayed  
515 little squad rotation policies and hence, differences in loading  
516 inevitably ensued.

517         Despite the novelty and practical application of the  
518 current study, our data are not without limitations, largely a  
519 reflection of currently available technology and the practical  
520 demands of data collection in an elite football setting. Firstly,  
521 the simultaneous use of both GPS and Prozone® to quantify  
522 training and competitive match demands, respectively, has  
523 obvious implications for the comparability of data between  
524 systems.<sup>38,39</sup> Nevertheless, during the chosen season of study, it  
525 was against FIFA rules to wear GPS in competitive matches.  
526 Whilst it is now within the rules to wear GPS in competitive  
527 games, it is still not common policy due to managers'  
528 preferences, players' comfort issues and poor signal strength  
529 due to the roofing in many stadiums in the English Premier  
530 League. Secondly, we also chose to not report data from games  
531 or training from International camps given that the loads of  
532 these practices were not controlled by the current research team  
533 or club's tactical and coaching staff. Finally, this study is only  
534 reflective of one team (albeit reflective of a top English Premier  
535 League team) and hence may not be representative of the  
536 customary training and match demands of other domestic teams  
537 or teams from other countries. When taken together, the  
538 simultaneous use of GPS in training and games, quantification  
539 of load in additional settings and the use of wider based  
540 samples all represent fruitful areas for future research.

541

## 542 **Practical Applications**

543         Given that we observed distinct differences in high-  
544 intensity distance completed throughout the season, our data  
545 have obvious practical implications for training programme  
546 design. In this regard, data suggest that players classified as  
547 fringe and non-starters should engage in additional high-  
548 intensity training practices and/or complete relevant time in  
549 non-competitive friendlies and U21 games in an attempt to  
550 recreate the high-intensity physical load typically observed in  
551 competitive first team games. This point is especially important  
552 given the relevance and importance of high-intensity activity in

553 both building and maintaining aspects of soccer specific  
 554 fitness.<sup>10,36,37</sup> Furthermore, our observation of more marked  
 555 differences in periods 3, 4 and 5 of the season also suggest that  
 556 specific attention should be given to those periods of the season  
 557 when tactical choices dictate low-squad rotation policies.  
 558 Future studies should now correlate changes in physical load  
 559 during the season to seasonal variation in soccer-specific fitness  
 560 components as well as introducing soccer-specific training  
 561 interventions at the relevant in-season periods (e.g. Iaia et al.  
 562 <sup>37</sup>).

563

564

## 565 **Conclusions**

566 In summary, we quantify for the first time the accumulative  
 567 training and match load (and total accumulative physical load)  
 568 during an annual season in those players considered as regular  
 569 starters, fringe players and non-starters. Importantly, although  
 570 we report that total duration of activity and total distance  
 571 covered was not different between playing groups, we observed  
 572 that starters generally completed more time in high-intensity  
 573 zones than fringe and non-starters players. Our data  
 574 demonstrate the obvious importance of participation in game  
 575 time for completing such high-intensity physical load. Such  
 576 data suggest that the training practices of these latter groups  
 577 should potentially be manipulated in order to induce  
 578 comparable seasonal workloads.

579

580

## 581 **Acknowledgements**

582 The authors would like to thank all of the team's coaches and  
 583 players for the cooperation during all data collection  
 584 procedures. We also acknowledge the support of Dave Rydings  
 585 in helping to collate and quantify the duration of resistance  
 586 training during the season of analysis.

587

588

## 589 **References**

- 590 1. Varley MC, Aughey RJ. Acceleration Profiles in Elite  
 591 Australian Soccer. *Int J Sports Med.* 2013;34:34-39.
- 592 2. Dellal A, Chamari K, Wong DP, et al. Comparisons of  
 593 physical and technical performance in European soccer

- 594 match-play: FA Premier League and La Liga. *Eur J*  
595 *Sport Sci.* 2011;11:51-59.
- 596 3. Di Salvo V, Baron R, Tschan H, Calderon Montero FJ,  
597 Bechl N, Pigozzi, F. Performance characteristics  
598 according to playing position in elite soccer. *Int J Sports*  
599 *Med.* 2007;28:222-227.
- 600 4. Bloomfield J, Polman R, O'Donoghue P. Physical  
601 demands of different positions in FA Premier League  
602 Soccer. *J Sports Sci Med.* 2007;6:63-70.
- 603 5. Bangsbo J, Mohr M, Krstrup P. Physical and  
604 metabolic demands of training and match-play in the  
605 elite football player. *J Sports Sci.* 2006;24:665-674.
- 606 6. Fernandes O, Caixinha P, Malta P. Techno-tactics and  
607 running distance analysis using one camera. *J Sports Sci*  
608 *Med.* 2007;6:204-205.
- 609 7. Rampinini E, Coutts AJ, Castagna C, Sassi R,  
610 Impellizzeri FM. Variation in top level soccer match  
611 performance. *Int J Sports Med.* 2007;28:1018-1024.
- 612 8. Bradley PS, Sheldon W, Wooster B, Olsen P, Boanas P,  
613 Krstrup P. High-intensity running in English FA  
614 Premier League soccer matches. *J Sports Sci.*  
615 2009;27:159-168.
- 616 9. Mohr M, Krstrup P, Bangsbo J. Match performance of  
617 high-standard soccer players with special reference to  
618 development of fatigue. *J Sports Sci.* 2003;21:439-449.
- 619 10. Iaia FM, Rampinini E, Bangsbo J. High-intensity  
620 training in football. *Int J Sports Physiol Perform.*  
621 2009;4:291-306.
- 622 11. Barnes C, Archer DT, Hogg B, Bush M, Bradley, PS.  
623 The Evolution of Physical and Technical Performance  
624 Parameters in the English Premier League. *Int J Sports*  
625 *Med.* 2014;35:1095-1100.
- 626 12. Reilly T, Thomas V. A motion analysis of work-rate in  
627 different positional roles in professional football match-  
628 play. *Journal of Human Movement Studies.* 1976;2:87-  
629 97.
- 630 13. Morgans R, Orme P, Anderson L, Drust B, Morton JP.  
631 An intensive winter fixture schedule induces a transient  
632 fall in salivary IgA in English premier league soccer  
633 players. *Res Sports Med.* 2014;22:346-354.
- 634 14. Gaudino P, Iaia FM, Alberti G, Strudwick AJ, Atkinson  
635 G, Gregson W. Monitoring training in elite soccer  
636 players: systematic bias between running speed and  
637 metabolic power data. *Int J Sports Med.* 2013;34:963-  
638 968.
- 639 15. Malone JJ, Di Michele R, Morgans R, Burgess D,  
640 Morton JP, Drust B. Seasonal training load  
641 quantification in elite English Premier League soccer  
642 players. *Int J Sports Physiol Perform.* 2015;10:489-497.

- 643 16. Anderson L, Orme P, Di Michele R, et al.  
644 Quantification of training load during one, two and  
645 three game week schedules in professional soccer  
646 players from the English Premier League: implications  
647 for carbohydrate periodization. *J Sports Sci.* In Press.
- 648 17. Silva JR, Magalhães JF, Ascensão AA, Oliveira EM,  
649 Sebba AF, Rebelo AN. Individual match playing time  
650 during the season affects fitness related parameters of  
651 male professional soccer players. *J Strength Cond Res.*  
652 2011;25: 2729-2739.
- 653 18. Coutts AJ, Duffield R. Validity and reliability of GPS  
654 devices for measuring movement demands of team  
655 sports. *J Sci Med Sport.* 2010;13:113-135.
- 656 19. Castellana J, Casamichana D, Calleja-Gonzalez J,  
657 Roman JS, Ostojic SM. Reliability and accuracy of 10  
658 Hz GPS devices for short-distance exercise. *J Sports Sci*  
659 *Med.* 2011;10:233-234.
- 660 20. Varley MC, Fairweather IH, Aughey RJ. Validity and  
661 reliability of GPS for measuring instantaneous velocity  
662 during acceleration, deceleration, and constant motion.  
663 *J Sports Sci.* 2012;30:121-127.
- 664 21. Rampinini E, Alberti C, Florenza M, et al. Accuracy of  
665 GPS devices for measuring high-intensity running in  
666 field-based team sports. *Int J Sports Med.* 2015;36:49-  
667 53.
- 668 22. Maddison R, Ni Mhurchu C. Global positioning system:  
669 a new opportunity in physical activity measurement. *Int*  
670 *J Behav Nutri Phys Act.* 2009;6,73.
- 671 23. Buchheit M, Al Haddad H, Simpson BM, Palazzi D,  
672 Bourdon PC, Di Salvo V, Mendez-Villanueva A.  
673 Monitoring accelerations with GPS in football: Time to  
674 slow down? *Int J Sports Physiol Perform.* 2014;9:442-  
675 445
- 676 24. Jennings D, Cormack S, Coutts AJ, Boyd LJ, Aughey  
677 RJ. Variability of GPS units for measuring distance in  
678 team sport movements. *Int J Sports Physiol Perform.*  
679 2010;5:565-569.
- 680 25. Di Salvo V, Collins A, McNeil B, Cardinale M.  
681 Validation of ProZone®: A new video-based  
682 performance analysis system. *Int J Perform Anal Sport.*  
683 2006;6:108-109.
- 684 26. Bangbo J. *Aerobic and Anaerobic training in soccer:*  
685 *with special emphasis on training of youth players.*  
686 Fitness training in soccer I. Bagsvaerd: Ho & Storm;  
687 2008.
- 688 27. Dupont G, Akakpo K, Berthoin S. The effect of in-  
689 season, high-intensity interval training in soccer players.  
690 *J Strength Cond Res.* 2004;18:584-589.
- 691 28. Wells C, Edwards A, Fysh M, Drust B. Effects of high-  
692 intensity running training on soccer-specific fitness in

- 693 professional male players. *Appl Physiol Nutr Metab*.  
694 2014;39:763-769.
- 695 29. Colby MJ, Dawson B, Heasman J, Rogalski B, Gabbett  
696 TJ. Accelerometer and GPS-derived running loads and  
697 injury risk in elite Australian Footballers. *J Strength*  
698 *Cond Res*. 2014;28:2244-2252.
- 699 30. Krstrup P, Mohr M, Amstrup T, et al. The yo-yo  
700 intermittent recovery test: physiological response,  
701 reliability, and validity. *Med Sci Sports Exerc*.  
702 2003;35:697-705.
- 703 31. Krstrup P, Mohr M, Ellingsgaard H, Bangsbo J.  
704 Physical demands during an elite female soccer game:  
705 importance of training status. *Med Sci Sports Exerc*.  
706 2005;37:1242-1248.
- 707 32. Egan B, Carson BP, Garcia-Roves PM, et al. Exercise  
708 intensity-dependent regulation of PGC-1 $\alpha$  mRNA  
709 abundance is associated with differential activation of  
710 upstream signalling kinases in human skeletal muscle. *J*  
711 *Physiol*. 2010;588:1779-1790.
- 712 33. Gillen JB, Gibala MJ. Is high-intensity interval training  
713 a time-efficient exercise strategy to improve health and  
714 fitness? *Appl Physiol Nutri Metab*. 2014;39:409-412.
- 715 34. Iaia FM, Thomassen M, Kolding H, et al. Reduced  
716 volume but increased training intensity elevates muscle  
717 Na<sup>+</sup>-K<sup>+</sup> pump  $\alpha$ 1-subunits and NHE1 expression as  
718 well as short-term work capacity in humans. *Am J*  
719 *Physiol Regul Integr Comp Physiol*. 2008;294:966-974.
- 720 35. Gabbett TJ. Influence of training and match intensity on  
721 injuries in rugby league. *J Sports Sci*. 2004;22:409-417.
- 722 36. Laursen PB, Jenkins DG. The scientific basis for high-  
723 intensity interval training: optimising training  
724 programmes and maximising performance in highly  
725 trained endurance athletes. *Sports Med*. 2002;32:53-73.
- 726 37. Iaia FM, Fiorenza M, Perri E, Alberti G, Millet GP,  
727 Bangsbo J. The effect of two speed endurance training  
728 regimes on performance of soccer players. *PLoS One*.  
729 2015;10:1-16.
- 730 38. Harley JA, Lovell RJ, Barnes CA, Portas MD, Weston  
731 M. The interchangeability of global positioning system  
732 and semiautomated video-based performance data  
733 during elite soccer match play. *J Strength Cond Res*.  
734 2011;25:2334-2338.
- 735 39. Buchheit M, Allen A, Poon TK, Modonutti M, Gregson  
736 W, Di Salvo V. Integrating different tracking systems in  
737 football: multiple camera semi-automatic system, local  
738 position measurement and GPS technologies. *J Sports*  
739 *Sci*. 2014;32:1844-1857.



742

743

744

745

746

747

748

749

750 TABLE 1 – Total duration (minutes), total distance (km),  
751 running distance (km), high-speed running distance (km) and  
752 sprinting distance (km) covered across the entire in-season  
753 period, as inclusive of both training and matches. \* denotes  
754 difference from starters,  $P < 0.05$  (Bonferroni corrected).

755

756 TABLE 2 – Total duration (minutes), total distance (km),  
757 running distance (km), high-speed running distance (km) and  
758 sprinting distance (km) within 5 specific in-season periods. \*  
759 denotes difference to starters, # denotes difference to fringe  
760 players,  $P < 0.05$  (Bonferroni corrected).

761

762 FIGURE 1 – Accumulative season long A) duration and B)  
763 total distance in both training and matches. Shaded bars =  
764 training and open bars = matches. \* denotes difference to  
765 starters (matches), # denotes difference to fringe players  
766 (matches), <sup>a</sup> denotes difference to starters (training),  $P < 0.05$   
767 (Bonferroni corrected).

768

769 FIGURE 2 – Accumulative season long A) running distance, B)  
770 high-speed running distance and C) sprinting distance in both  
771 training and matches. Shaded bars = training and open bars =  
772 matches. \* denotes difference to starters,  $P < 0.05$  (Bonferroni  
773 corrected).

774

775 FIGURE 3 – Within period accumulative A) duration, B) total  
776 distance, C) running distance, D) high-speed running distance

777 and E) sprinting distance in match per se. \* denotes difference  
778 to starters, # denotes difference to fringe players,  $P < 0.05$   
779 (Bonferroni corrected).

780

781 FIGURE 4 – Within period accumulative A) duration, B) total  
782 distance, C) running distance, D) high-speed running distance  
783 and E) sprinting distance in training per se. \* denotes difference  
784 to starters, # denotes difference to fringe players,  $P < 0.05$   
785 (Bonferroni corrected).

786