

**Rat eradication comes within a whisker! A case study of a failed project from the South Pacific**

W. Amos<sup>1</sup>, H.J. Nichols<sup>1,2</sup>, T. Churchyard<sup>3</sup> and M. de L. Brooke<sup>1</sup>

<sup>1</sup>Department of Zoology, Downing Street, Cambridge, UK

<sup>2</sup>Liverpool John Moores University, School of Natural Science & Psychology, Liverpool, UK

<sup>3</sup>RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, UK

**Author for correspondence:** M. de L. Brooke email: [m.brooke@zoo.cam.ac.uk](mailto:m.brooke@zoo.cam.ac.uk)

**Keywords:**

Brodifacoum, genetic diversity, Henderson Island, heterozygosity, Pitcairn Islands, *Rattus exulans*

Number of words: Abstract – Figure Legends 5086

Number of references: 33

Three figures, three tables, two supplementary figures, one supplementary table, one supplementary word document of Visual Basic Code

Correspondence: M. de L. Brooke, Department of Zoology, Downing Street, Cambridge CB2 3EJ, UK. Tel: +44-1223-336610, Fax: +44-1223-336676, Email: [m.brooke@zoo.cam.ac.uk](mailto:m.brooke@zoo.cam.ac.uk)

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

To enhance their conservation value, several hundred islands worldwide have been cleared of invasive alien rats *Rattus* spp. One of the largest projects yet undertaken was on 43 km<sup>2</sup> Henderson Island in the Pitcairn group, South Pacific, in August 2011. Following massive immediate mortality, a single *R. exulans* was observed in March 2012 and, subsequently, rat numbers have recovered. The survivors show no sign of resistance to the toxicant used, brodifacoum. Using pre- and post-operation rat tissue samples from Henderson, plus samples from around the Pacific, we exclude re-introduction as the source of continued rat presence. Microsatellite analysis of 18 loci enabled comparison of genetic diversity of Henderson rats before and after the bait drop. The fall in diversity measured by allele frequency change indicated that the bottleneck ( $N_e$ ) through which the breeding population passed was probably around 50 individuals, representing a census population of about 60-80 animals. This is the first failed project that has estimated how close it was to success.

## 56 1. Introduction

57

58 Remote oceanic islands harbour a wide range of species found nowhere else in the world, yet  
59 many of these species are being driven to extinction by the introduction of alien species, most  
60 notably rats *Rattus* spp. [1,2]. Recent conservation efforts have led to rodent eradication  
61 operations on 719 islands across the world (<http://diise.islandconservation.org/>, accessed 10 April  
62 2015). For larger islands, the eradication method of choice usually involves helicopter distribution  
63 of poison bait. The great majority, 80% (578 of 719) of such operations are successful but, despite  
64 meticulous planning, a minority fail and some correlates of failure are emerging [3]. A more  
65 refined understanding of the reasons behind failures will, in part, depend on knowing how close a  
66 project was to success. More specifically, in the aftermath of a failure, it would be valuable for  
67 conservation planners to know whether a mere handful of rats survived the poison bait, in which  
68 case minor tweaking of the baiting protocol might ensure the success of a second attempt, or  
69 whether survivors numbered hundreds or more, suggesting the need for major revision of the  
70 eradication protocol [4]. The study we report here is, to our knowledge, the first attempt to use  
71 genetic data to estimate the number of survivors of a failed rat eradication operation (but see [5]).

72 The project in question was undertaken in August 2011 on the 43 km<sup>2</sup> World Heritage site  
73 of Henderson Island (24°20'S, W 128°19'W) in the Pitcairn group, South Pacific. The island is  
74 home, inter alia, to four species of endemic landbird and four breeding species of surface-nesting  
75 *Pterodroma* petrel. The latter's chicks are known to be victims of predation by Pacific rats *Rattus*  
76 *exulans* which arrived with Polynesian colonists some 800 years ago [6,7]. To rid the island of  
77 rats, 75 tonnes of bait, laced with the anti-coagulant poison brodifacoum, were dropped from  
78 helicopters in an operation costing about £1.5 million organised by the Royal Society for the  
79 Protection of Birds (RSPB) [8]. Massive immediate mortality was achieved and no rats were seen  
80 on the island for the first three months when personnel were on-island continuously. However, in  
81 March 2012, seven months after the bait drop, a single *R. exulans* was observed by a temporary

visitor. Since then rat numbers (which may vary with season) have recovered fully to 50,000 – 100,000 individuals [9-11].

Independent reviews of the project in 2013 did not identify any operational flaws [12]. However, in the absence of genetic data, the reviews could neither exclude re-invasion of rats from elsewhere nor brodifacoum resistance among Henderson rats.

By sampling rats from other island groups in the region, we confirmed that the project's failure was indeed a failure of eradication as opposed to an untimely re-introduction from neighbouring islands [13]. Furthermore, a comparison of microsatellite data from rats sampled before the operation, in 2009, and after, in 2012 and 2013, allowed us to estimate the number of animals to which the population was reduced. Finally, on-island tests of the survivors in 2013 yielded no evidence that the failure was due to brodifacoum resistance among those survivors.

## 2. Methods

### (a) Study sites and sample acquisition

Henderson Island is roughly 9 km north-to-south by 5 km east-to-west. All rat samples were obtained in the northern one-third of the island (electronic supplementary material, figure S1) because movement over the terrain is so difficult for people due to the thickness of the vegetation. However the vegetation would pose no barrier to a rat, and there are no obvious topographical features that would impede rat dispersal across the island. Most of the samples were obtained by snap-trapping, either on the plateau, a former lagoon on this raised coral island now lying 30 m above sea level, or in the vegetation immediately behind the North and East Beaches. In each case about 2cm of tail was removed and stored in 95% ethanol. Henderson samples were obtained as follows:

(i) Pre-eradication samples (henceforth pre-samples) – 50 animals, September 2009

(ii) Post-eradication samples (henceforth post-samples)

108 (a) 1 animal, May 2012

109 (b) 63 animals, November 2012

110 (c) 19 animals, August 2013

111 The density of the vegetation across the island, coupled with the considerable cost of  
112 keeping a large vessel, two helicopters and personnel on station, meant that island-wide  
113 monitoring for rats surviving after the August 2011 bait drop was simply impractical. These twin  
114 problems, remoteness and onshore conditions, are likely to preclude sustained post-drop  
115 monitoring at many islands of high conservation value where eradication may be under  
116 consideration for the future.

117 To test the possibility that the continuing presence of rats on Henderson is not due to an  
118 introduction after the eradication attempt, samples were obtained from other islands. The nearest  
119 island to Henderson, and therefore the most likely source of a rat re-introduction is Pitcairn, ~200  
120 km to the south-west. Pitcairn rat samples ( $n = 30$ , July 2011;  $n = 18$ , June 2014) were obtained  
121 and preserved as for the Henderson samples in July 2011. Additional samples were obtained from  
122 more distant island groups, namely the Gambier archipelago in south-east French Polynesia 600  
123 km west of Henderson ( $n = 38$  successful genotypings from 26 rats obtained in April 2010 and 23  
124 in January 2013), and the Cook Islands 3000 km west of Henderson ( $n = 10$  amplified from  
125 Anchorage and Motu Tou islets off Suvarrow, May 2013).

126

## 127 (b) Laboratory analysis

128 DNA was extracted from most tail-tips using a glass milk extraction method, adapted for use in  
129 microtitre plates rather than individual tubes. Briefly, tissue was digested in proteinase K and the  
130 liberated DNA adsorbed to flint glass particles in the presence of a 3X excess of 6M NaI. After  
131 two ethanol washes the DNA was eluted in 100ul of low TE buffer. In a brief study commissioned  
132 from Landcare Research, New Zealand, to assess whether the single rat caught post-eradication in  
133 May 2012 was a re-introduction, samples from this individual, from 30 pre-eradication Henderson

134 rats, and from 30 Pitcairn rats were extracted and analysed for eight microsatellite markers. These  
135 DNA samples were used by us without re-extraction.

136

137 Table 1 about here

138

139 Samples were genotyped at 19 microsatellite loci isolated from the *Rattus norvegicus*  
140 genome (table 1). The 19 included seven of the eight loci used by Landcare, extended by testing a  
141 further 31 markers from *R. norvegicus* on *R. exulans*. In order to minimise the probability of  
142 linkage between microsatellites, the markers chosen for testing were selected from throughout the  
143 *R. norvegicus* genome. Genotyping was conducted in 10µl multiplex PCR reactions (Qiagen®  
144 Multiplex PCR Kit, UK) with fluorescent-labelled forward primers containing approximately 20ng  
145 DNA, following the manufacturer's protocol. Three sets of multiplex reactions were carried out,  
146 each with between five and seven primer pairs (table 1). PCR conditions comprised an initial  
147 denaturing step of 95°C for 15 min, followed by 35 cycles of 94°C denaturation for 30s, 57°C  
148 annealing for 90s and 72°C extension for 60 s, followed by a final extension of 30 min at 60°C.  
149 Following PCR, 1µl of PCR product was mixed with 10µl of loading mix (1mL Hi-Di™  
150 Formamide plus 20µl of Genescan LIZ 500 ladder (ABI)) before visualisation on an ABI3730  
151 DNA Analyzer.

152

### 153 (c) Statistical analysis

154 Alleles were called using GeneMapper 3.7. The Henderson pre-bottleneck data were tested for the  
155 presence of null alleles using Cervus [14] with two having low frequency nulls (3% for D15Rat77  
156 and 7% for D7Rat13). Another (D2Rat234) had an inferred null allele frequency of 36% and was  
157 excluded from all the analyses we report, which are therefore based on 18 loci. To explore the  
158 level of differentiation seen among our sampling regions and potentially exclude other  
159 neighbouring islands as a source of a re-introduction, we used the program STRUCTURE [15],

160 with burn-in length 20,000 and 50,000 steps, varying the possible number of putative sub-groups  
161 (k) from one to 15 with 10 repetitions of each. We did not use sampling locality priors, allowed  
162 admixture and assumed that loci were unlinked. In addition, for the two focal islands, Henderson  
163 and Pitcairn, we constructed a joint individual-based neighbour-joining tree using a pairwise  
164 relatedness matrix, constructed following the methods of Queller & Goodnight [16]. The tree was  
165 calculated using Phylip [17].

166 Classical methods of estimating the effective population size ( $N_e$ ) of a bottleneck may  
167 struggle in our study. First, bottleneck size can be estimated through the associated loss of  
168 heterozygosity. However, this offers a rather blunt tool because detectable losses of  
169 heterozygosity require either very low numbers of survivors (<10) or a longer duration (i.e. no  
170 immediate recovery). Further, loss of heterozygosity could conceivably be distorted by the  
171 preferential survival of more heterozygous individuals [18]. Some other methods exploit features  
172 of the way microsatellites evolve and compare observed test statistics with those based on  
173 simulations assuming a strict stepwise mutation model (SMM) or related models. For example,  
174 the program 'Bottleneck' [19] is based on transient changes in the ratio of allele range to  
175 heterozygosity associated with the loss of rare alleles. However, the SMM is over-simple because  
176 it is now known that microsatellite mutations are centrally directed, with long alleles contracting  
177 and short alleles expanding. Moreover, cross-species markers can show strong departures from the  
178 SMM due to interruption mutations within the microsatellite itself [20]. Finally, our cross-species  
179 markers include several that may or actually do carry non-amplifying alleles.

180 To address these issues and to estimate the likely size of the bottleneck we used two  
181 approaches. First, we used the program NeEstimator v2 [21] to implement the standard temporal  
182 method [22]. Second, we used stochastic simulations implemented using custom code written in  
183 Visual Basic (Supplementary Material). For the latter, semi-realistic allele frequency distributions  
184 are established by allowing microsatellites to evolve under a strict stepwise mutation model in a  
185 homogeneous population of size 1000 diploid individuals with mutation rate chosen such that

heterozygosity was similar to that seen in our empirical data for pre-eradication attempt rats on Henderson. Each population is then reduced to bottleneck size  $X$  ( $X=5$  to 100, with equal sex ratio) and then allowed to expand rapidly at three fold per generation over a maximum of six generations back to a maximum size of 1000. During this realistically short time period the contribution of new mutations can safely be ignored. Samples of 50 and 83 were drawn at random both immediately before the bottleneck and at the end of the simulation and used to assess heterozygosity, calculated as expected heterozygosity at Hardy-Weinberg equilibrium, and allele number. All simulations were repeated 100 times to obtain a mean and standard deviation.

As mentioned above, changes in heterozygosity are likely too slight to be informative. Consequently, we focused on the average change in frequency of individual alleles [23]. There are two potential issues with this approach. First, our estimated allele frequencies are based on modest samples of an unknown underlying distribution. Second, even if we did know the underlying frequency distributions, the chance that these are replicated closely during simulation is remote. Consequently, we systematically explored the fate of single alleles at all possible frequencies. Specifically, individual populations of 1000 individuals were founded carrying just two alleles (frequency of minor allele = 0.01 to 0.5, step 0.01). The bottleneck, recovery and sampling were performed as above, yielding a complete spectrum of frequency trios: unobserved true pre-bottleneck, pre-bottleneck in a sample the size we collected and post-bottleneck of the size we collected with, on average, 100 replicates of each observed pre-bottleneck frequency. Using these data we can then translate our empirically observed allele frequencies into the expected average change in frequency for a bottleneck of any given size. To explore the impact of null alleles, all simulations were repeated two further times, adding null alleles to the pre-bottleneck population at the observed frequencies seen at two loci of (3%, 7%).

(d) Testing post-eradication pesticide resistance



211 To address the question of whether the rats surviving on Henderson were resistant to brodifacoum,  
212 58 rats were captured in July and August 2013 using Elliot live traps. On capture the weight and  
213 sex of each rat were recorded prior to placing it in individual 70cmx50cmx50cm cages, made from  
214 wire mesh and plywood board. Each cage contained a food bowl and water dispenser, while  
215 plastic screens placed between cages prevented visual interaction.

216 Each rat had three days of acclimatisation on a diet of guinea-pig food (a non-toxic food  
217 source obtained from a pet food supplier; Animates Rabbit and Guinea Pig Food, Animates,  
218 Wellington, New Zealand). After the three days, each rat was re-weighed (to ensure it was healthy  
219 and not losing body mass) and randomly assigned to one of eight trial groups. There were a control  
220 group (no poison) and seven experimental groups exposed to progressively higher concentrations  
221 of brodifacoum in their diet. Each group had eight rats except for the two groups receiving the  
222 highest dose where  $n = 5$ .

223 Each trial group was then presented with brodifacoum-laced peanut butter (brodifacoum  
224 0.002% wet weight), weighed to deliver the dose of brodifacoum required by the rat's weight,  
225 along with some guinea-pig food (table 3, below). The control group animals each received 1g of  
226 non-toxic peanut butter. All rats had consumed all the poison within 24 hours. After toxicant  
227 consumption, the rats' diet returned to guinea-pig food. Each rat was monitored twice daily until  
228 death or 14 days after the poison dose was ingested. Any rat clearly suffering and close to death  
229 was euthanized by cervical dislocation (this occurred twice); rats surviving after 14 days were  
230 released close to where captured.

231

### 232 3. Results

233

#### 234 (a) Genetic substructure

235 The program STRUCTURE revealed the best-fit number of sub-groups to be 10 (electronic  
236 supplementary material, figure S2). Group membership is summarised in a bar plot (figure 1).

237 This reveals a number of interesting features. First, rats from the Cook Islands and from four  
238 sampling localities within the Gambier archipelago are all assigned to their own groups with near  
239 100% confidence and minimal overlap with either Henderson or Pitcairn (NB one Cook Islands  
240 sample came from a different region from the others and was clearly genetically different, failing  
241 to place in any of the 10 groups). Despite being genetically very similar ‘by eye’, in the sense that  
242 the same alleles occur in both samples at ostensibly similar frequencies, Pitcairn and Henderson  
243 samples are also largely separable. Within the islands, the two main groups on Pitcairn (yellow  
244 and blue) correspond well with the two different sampling efforts, while there are three main  
245 groups on Henderson (pink, white and grey). On Henderson, there is a weaker but still rather clear  
246 separation between pre-eradication samples (mainly pink) and post-eradication samples (largely  
247 white and grey).

248 The neighbour-joining tree of all Henderson and Pitcairn rats is shown in figure 2. Two  
249 important features are apparent. First, mirroring the STRUCTURE results, the Pitcairn rats fall  
250 mainly in a single, almost pure clade, indicating genetic separation (figure 2a). Second, the  
251 Henderson rats exhibit some degree of clustering, in that multiple rats from the same sampling  
252 location often cluster together (figure 2b). This may well be due to the sampling of related  
253 individuals on a scale of tens of metres. On the other hand, each of the primary sampling areas (E.  
254 Beach, N. Beach, Plateau; electronic supplementary material, figure S1) contributes clusters spread  
255 right across the tree, suggesting appreciable gene flow between the different regions. Also, there  
256 is no strong tendency for pre-eradication and post-eradication attempt samples to cluster strongly.  
257 We conclude that, while mild sub-structure is present, this is probably not strong enough to  
258 indicate strongly restricted gene flow between the island regions.

259

260 Figs 1, 2 and 3 and Table 2 about here

261

262 (b) Estimating the likely minimum number of rats

263 Unbiased heterozygosity reveals negligible change while expected heterozygosity actually  
264 increases slightly. However, as discussed in Methods, changes in overall heterozygosity provide  
265 a relatively crude measure and would require a very severe bottleneck to show a detectable  
266 change. Changes in allele frequency may provide a more robust estimate because each locus  
267 carries multiple alleles, each of which provide a point estimate. Using this approach, we find  
268 excellent agreement between three related estimators, implemented in NeEstimator v2, and our  
269 own stochastic simulations. In all cases the best-fit number of survivors was an effective  
270 population size of approximately  $50 \pm 30$  rats (table 2, figure 3).

271

#### 272 (c) Toxicity tests

273 Mortality was 100% for dose levels of 0.2 mg/Kg (bw) and above; only the control group had zero  
274 mortality (table 3). The median lethal dose LD<sub>50</sub>, calculated in R3.0.1, was  $0.061 \pm 0.02$  mg/Kg  
275 (95% CI).

276

277 Table 3 about here

278

#### 279 4. Discussion

280 We genotyped samples collected from other Pacific islands for 18 polymorphic microsatellites and  
281 found no evidence that re-introduction from these sources is the reason for the continuing presence  
282 of rats on Henderson Island. We therefore conclude that rats remain on Henderson because of a  
283 failure to eradicate. We then compared the microsatellite genotypes obtained from rats sampled  
284 before and after the eradication attempt, to estimate the number to which the population was  
285 reduced. Heterozygosity was unchanged but the average change in frequency of individual alleles  
286 suggests an effective population size ( $N_e$ ) of around 50 rats at the deepest part of the bottleneck.

287 Early studies of the relationship between effective population size ( $N_e$ ) and census size ( $N$ )  
288 suggested that most species have a ratio  $N_e / N$  in the approximate range 0.25 to 0.75 [24]. A more

289 recent study [25] suggests that mammals have an average ratio of  $0.749 \pm 0.117$  standard  
290 deviation, implying that our estimate of  $N_e = 50$  rats will translate into a census size of 58 to 80  
291 rats.

292 An estimate of approximately 60-80 actual survivors seems plausible. Following the  
293 attempted eradication, no rats were seen over the first three months when fieldworkers were  
294 continuously present. Even with the dense vegetation, this seems unlikely unless the numbers left  
295 alive were very small. On the other hand, once the first rat was seen by a temporary visitor seven  
296 months after the bait-drop, numbers did rebuild rapidly [10], something that seems less likely if  
297 the population had been reduced below 10. Under such circumstances inbreeding depression and  
298 the difficulty of finding a mate might have slowed recovery [26].

299 Our analysis reveals subtle population substructure but probably not enough to impact our  
300 conclusions. Thus, Pitcairn was sampled at two different locations and times and these are  
301 reasonably well resolved by STRUCTURE (figure 1). Similarly, while the pre-eradication sample  
302 from Henderson appears fairly homogeneous, STRUCTURE suggests three weak groups among  
303 animals sampled after the eradication attempt. These groups may well reflect a small number of  
304 pockets of survivors which interbred before dispersing more widely. On the other hand the  
305 individual-based tree shows no obvious relationship with geography beyond the presence of small,  
306 possibly family clusters (figure 2), and STRUCTURE places some Pitcairn rats on Henderson and  
307 *vice versa*. Importantly, appreciable substructure should result in weak homozygote excess across  
308 loci and we find a weak heterozygote excess (data not shown).

309 Logistic constraints meant we did not sample rats across the full extent of Henderson  
310 (figure S1). However we did sample from sites that are reasonably distant in the context of the  
311 entire island and find very little (arguably negligible) structure beyond that which might be  
312 expected from the sampling of relatives. Moreover, Pitcairn shows only marginal distinctiveness.  
313 If the Henderson recovery was based on immigration from Pitcairn, presumably by  $<10$   
314 individuals, we would expect extremely low diversity, significantly lower than we observe, and the

315 post-eradication rats should form a reasonably distinct sub-clade within the Pitcairn clade. That  
316 post-eradication rats have higher diversity and co-cluster with pre-eradication Henderson rats  
317 therefore argues strongly that this is survival rather than reintroduction. Combined, this suggests  
318 that our sampling has captured the majority of Henderson diversity: it would be truly remarkable if  
319 an unsampled clade existed on Henderson that was more different than rats from Pitcairn.

320 Our toxicity tests revealed that only the control rats receiving a zero brodifacoum dose  
321 showed no mortality. The calculated LD<sub>50</sub> of 0.061 mg/Kg among rats on Henderson in 2013 was  
322 considerably lower than the acute LD<sub>50</sub> of 0.32 mg/Kg found by Conor & Booth [27] in another  
323 population of *R. exulans*. These results therefore provide no support for the possibility that failure  
324 of the 2011 operation was due to brodifacoum resistance among the Henderson rats.

325 Since 0.2mg/kg (or higher) was lethal for all rats tested, an average rat of 80g (own data)  
326 would certainly die after ingesting 0.016mg of toxin. Given that the active toxin, brodifacoum,  
327 constituted 0.002% (20 mg/kg) of pelleted bait weight in the 2011 eradication attempt [8], the fatal  
328 amount of toxicant would be ingested by a rat eating 0.8 g of bait. Since each bait pellet weighed  
329 roughly 1.6 g [11], an average weight rat would die after consuming less than a single pellet. The  
330 amount of bait broadcast on the island, averaging almost 20kg/ha, was equivalent to hundreds of  
331 pellets per rat.

332 Our genetic results show that the eradication attempt was very nearly successful, and the  
333 brodifacoum tests provide no evidence of resistance to the toxicant used. The genetic results also  
334 argue against any re-introduction which is in any case highly unlikely given our knowledge of  
335 inter-island boat movements. It appears then that failure was neither caused by a tiny number of  
336 survivors nor by several hundred survivors, the latter indicating some systematic fault in the  
337 baiting approach used. Rather, the failure could have been due to a small number of rats either not  
338 encountering poison pellets or showing a preference for natural food over bait pellets. In tropical  
339 or sub-tropical latitudes, it may always be more difficult than in temperate latitudes to ensure that  
340 operations are undertaken when the availability of natural food is very low [28], especially when

critical ‘postpone or proceed’ decisions have of necessity to be taken months before the operation. Further plant phenological studies that enhance understanding of the ebb and flow of natural food on Henderson would be useful [29]. Arguably even more useful and of widespread value to other projects would be work directed towards increasing the attractiveness of bait pellets over natural food [30].

Ethics. Having considered the ethical and welfare implications, RSPB Council gave ethical approval for the brodifacoum experiment we report.

Data accessibility. See Supplementary Material.

Authors’ contributions. WA analysed the genetic data and ran the simulations, HJN conducted the lab analysis, TC performed the brodifacoum tests, and MdeLB assembled the non-Henderson rat samples and drafted the paper. All authors helped design the study and revise the drafts.

Competing interests. We have no competing interests.

Funding. The Darwin Initiative funded much of the RSPB’s work on Henderson Island and therefore sample collection and analysis, the latter also supported by the Sir Peter Scott Commemorative Expedition to the Pitcairn Islands.

Acknowledgements. Crucial fieldwork help on Henderson came from Richard Cuthbert, Gavin Harrison, Tara Proud, Mike Rodden and Nick Torr, while travel to and from the island hinged on the wonderful service provided the crews of *Braveheart*, *Claymore* and *Teba*. Rat samples from other islands were kindly supplied by Steve Cranwell and Kelvin Passfield (Cook Islands), the Reasin family, Laurent Yan and the staff of Manu (Gambier Islands), and Sue O’Keefe, Jay Warren and Pawl Warren (Pitcairn Island). We also thank Robyn Hewitt and Frank Molinia of Landcare Research for extracting DNA samples. Referee input saved us from grievous errors; we offer sincere thanks.

## 367 REFERENCES

368


- 369 1. Atkinson IA. 1985 The spread of commensal species of *Rattus* to oceanic islands and their  
 370 effects on island avifaunas. In *Conservation of Island Birds* (ed Moors PJ), pp. 35-81.  
 371 Cambridge UK: ICBP Technical Publication No. 3.
- 372 2. Howald G, Donlan CJ, Galvan JP, Russell JC, Parkes J, Samaniego A. *et al.* 2007 Invasive  
 373 rodent eradication on islands. *Conserv. Biol.* **21**, 1258-1268. (doi: 10.1111/j.1523-  
 374 1739.2007.00755.x)
- 375 3. Holmes ND, Griffiths, R, Pott, M, Alifano A, Will D, Wegmann AS *et al.* 2015 Factors  
 376 associated with rodent eradication failure. *Biol. Conserv.* **185**, 8-16. (doi:  
 377 10.1016/j.biocon.2014.12.018)
- 378 4. Pott M, Wegmann A, Griffiths R, Samaniego-Herrera A, Cuthbert RJ, Brooke MdeL. *et al.*  
 379 2015 Improving the odds: assessing bait availability before rodent eradications to inform bait  
 380 application rates. *Biol. Conserv.* **185**, 27-35. (doi: 10.1016/j.biocon.2014.09.049)
- 381 5. Abdelkrim J, Pascal M, Samadi S. 2007 Establishing causes of eradication failure based on  
 382 genetics: Case study of ship rat eradication in Ste. Anne archipelago. *Conserv. Biol.* **21**, 719-  
 383 730. (doi: 10.1111/j.1523-1739.2007.00696.x)
- 384 6. Brooke MdeL. 1995 The breeding biology of the gadfly petrels *Pterodroma* spp. of the Pitcairn  
 385 Islands: characteristics, population sizes and controls. *Biol. J. Linn. Soc.* **56**, 213-231.
- 386 7. Weisler MI. 1994 The settlement of marginal Polynesia – new evidence from Henderson  
 387 Island. *J. Field Archaeol.* **21**, 83-102. (doi: 10.2307/530246)
- 388 8. Torr N, Brown D. 2012 *Henderson Island Restoration Project: Post-Operational Report.*  
 389 Sandy, UK: RSPB.
- 390 9. Brooke MdeL, Cuthbert RJ, Henricson A, Torr N, Warren P, O’Keefe S. 2010 *Towards Rat*  
 391 *Eradication on Henderson Island: Fieldwork Report August-September 2009.* Sandy, UK:  
 392 RSPB.

- 393 10. Churchyard T, Proud T, Brooke MdeL, O’Keefe S, Warren P, Rodden M. 2015 *Henderson*  
394 *Island trip report 30th June – 31st August 2013*. Sandy, UK: RSPB.
- 395 11. Cuthbert RJ, Brooke MdeL, Torr N. 2012 Overcoming hermit crab interference during  
396 rodent baiting operations: a case study from Henderson Island, South Pacific. *Wildlife Res.*  
397 **39**, 70-77. (doi: 10.1071/WR11033)
- 398 12. Island Eradication Advisory Group, Island Conservation, Wildlife Management  
399 International. 2013 *Three independent reviews of the Henderson Island eradication*  
400 *operation*. Sandy, UK: RSPB.
- 401 13. Russell JC, Miller SD, Harper GA, MacInnes HE, Wylie MJ, Fewster RM. 2010 Survivors  
402 or reinvaders? Using genetic assignment to identify invasive pests following eradication.  
403 *Biol. Invasions* **12**, 1747-1757. (doi: 10.1007/s10530-009-9586-1)
- 404 14. Marshall TC, Slate JBKE, Kruuk LEB, Pemberton JM. 1998. Statistical confidence for  
405 likelihood-based paternity inference in natural populations. *Mol. Ecol.* **7**, 639-655. (doi:  
406 10.1046/j.1365-294x.1998.00374.x)
- 407 15. Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus  
408 genotype data. *Genetics* **155**, 945-959.
- 409 16. Queller DC, Goodnight KF 1989 Estimating relatedness using genetic markers. *Evolution* **43**, 258-  
410 275. (doi: 10.2307/2409206)
- 411 17. Felsenstein J. 2005 *PHYLIP (phylogeny inference package)*, version 3.6. University of Washington,  
412 Seattle: Department of Genome Sciences.
- 413 18. Coltman DW, Pilkington JG, Smith JA, Pemberton JM. 1999 Parasite-mediated selection against  
414 inbred Soay sheep in a free-living island population. *Evolution* **53**, 1259-1267. (doi:  
415 10.2307/2640828)
- 416 19. Luikart G, Allendorf FW, Cornuet J-M, Sherwin WB. 1998 Distortion of allele frequency  
417 distributions provides a test for recent population bottlenecks. *J. Hered.* **89**, 238-247. (doi:  
418 10.1093/jhered/89.3.238).



- 419 20. Jin L, Macaubas C, Hallmayer J, Kimura A, Mignot E. 1996. Mutation rate varies among  
420 alleles at a microsatellite locus: phylogenetic evidence. *Proc. Natl. Acad. Sci. USA* **93**,  
421 15285-15288. (doi: 10.1073/pnas.93.26.15285)
- 422 21. Do C, Waples RS, Peel D, Macbeth GM, Tillett BJ, Ovenden JR. 2014. NeEstimator v2: re-  
423 implementation of software for the estimation of contemporary effective population size ( $N_e$ )  
424 from genetic data. *Mol. Ecol. Resour.* **14**, 209-214. (doi: 10.1111/1755-0998.1215 )
- 425 22. Waples RS. 1989. A generalized approach for estimating effective population size from  
426 temporal changes in allele frequency. *Genetics* **121**, 379-391.
- 427 23. Pollak E. 1983. A new method for estimating the effective population size from allele  
428 frequency changes. *Genetics* **104**, 531-548.
- 429 24. Nunney L. 1996 The influence of variation in female fecundity on effective population size.  
430 *Biol. J. Linn. Soc.* **59**, 411-425
- 431 25. Waples RS, Luikart G, Faulkner JR, Tallmon DA 2013 Simple life-history traits explain key  
432 effective population size ratios across diverse taxa. *Proc. Roy. Soc. B*, **280**. Article 20131339.  
433 (doi: 10.1098/rspb.2013.1339)
- 434 26. Nathan HW, Clout MN, MacKay JWB, Murphy EC, Russell JC. 2015. Experimental island  
435 invasion of house mice. *Popul. Ecol.* **57**, 363-371. (doi: 10.1007/s10144-015-0477-2)
- 436 27. Connor CEO, Booth LH. 2001 Palatability of rodent baits to wild house mice. *Science for*  
437 *Conservation* **184**, NZ Department of Conservation, New Zealand.
- 438 28. Russell JC, Holmes ND. 2015 Tropical island conservation: Rat eradication for species  
439 recovery. *Biol. Conserv.* **185**, 1-7. (doi: 10.1016/j.biocon.2015.01.009)
- 440 29. Brooke MdeL, Jones PJ, Vickery JA, Waldren S. 1996 Seasonal patterns of leaf growth and  
441 loss, flowering and fruiting on a sub-tropical Central Pacific Island. *Biotropica* **28**, 164-179.
- 442 30. Keitt B, Griffiths R, Boudjelas S, Broome K, Cranwell S, Millett J. *et al.* 2015 Best practice  
443 guidelines for rat eradication on tropical islands. *Biol. Conserv.* **185**, 17-26. (doi:  
444 10.1016/j.biocon.2014.10.014)

- 445 31. Laulerkind SJ, Hayman GT, Wang SJ, Smith JR, Lowry TF, Nigam R. *et al.* 2013 The Rat  
446 Genome Database 2013 – data, tools and users. *Briefings in Bioinformatics* **14**, 520-526.
- 447 32. Nei M, Tajima F. 1981. Genetic drift and estimation of effective population size. *Genetics*  
448 **98**, 625-640.
- 449 33. Jorde PE, Ryman N. 2007. Unbiased estimator for genetic drift and effective population size.  
450 *Genetics* **177**, 927-935. (doi: 10.1534/genetics.107.075481)
- 451




453    C    Gam    H-post    H-pre    Pit

454

455    **Figure 1.** STRUCTURE plot of rats sampled from the Cook Islands (C), the Gambier archipelago

456    (Gam), Henderson Island pre- and post-eradication attempt, and Pitcairn (Pit)



458


459    **Figure 2.** (a) Neighbour-joining tree of all Henderson and Pitcairn rats. The tree was constructed

460    using a distance matrix of  $1-R$ , where  $R$  is relatedness sensu Queller and Goodnight. For clarity,

461    individual rats are colour coded: blue = Pitcairn; black = Henderson pre-eradication attempt; red =

462    Henderson post-eradication attempt.

463 (b) The neighbour-joining tree of Henderson rats, according to trapping location and  
 464 period. Red = North Beach pre-eradication, orange = North Beach post eradication, dark blue =  
 465 East Beach pre-eradication, light blue = East Beach post eradication, black = plateau.  
 466



467  
 468 **Figure 3.** Change in allele frequency among Henderson rats. The heavy horizontal line shows the  
 469 estimated average change in individual microsatellite allele frequency between rats sampled before  
 470 and after the eradication program. The errors on the estimate are one standard error of the mean,  
 471 obtained by averaging across 18 loci. Solid dots represent the mean change in allele frequency of  
 472 microsatellites when a simulated population passes through a bottleneck of the size given on the x-  
 473 axis.

474  
 475  
 476

**Table 1.** A summary of the 19 microsatellite loci, amplified in three multiplex PCR reactions, used to genotype Pacific rats *Rattus exulans* samples. All primer sequences are available in the Rat Genome Database [31].

Locus	Fluorescent label	Multiplex number	Product size range (bp)	Number of alleles
D7Arb16	HEX	1	87-104	9
D2Rat234	HEX	1	114-121	4
D8Mgh4	6-FAM	1	114-156	9
D12Rat36	HEX	1	182-195	8
D19Rat75	6-FAM	1	195-207	6
D10Rat20	6-FAM	2	97-128	12
D9Mit3	HEX	2	98-109	6
D11Rat7	HEX	2	131-149	5
D6Rat99	6-FAM	2	133-158	10
D5Rat83	6-FAM	2	166-180	6
D7Rat13	HEX	2	201-209	3
D15Rat77	6-FAM	2	240-261	11
D2Rat312	HEX	3	94-112	9
D17Mgh1	6-FAM	3	112-136	13
D1Rat313	6-FAM	3	153-167	8
D4Rat106	HEX	3	155-178	7
D6Rat100	6-FAM	3	174-186	7
D8Rat162	HEX	3	202-215	6
D14Rat39	6-FAM	3	223-257	14

488  
489  
490  
491  
492  
493  
494  
495  
496

**Table 2.** Estimated bottleneck size based on the temporal method. The program NeEstimator v2 was used to estimate the likely size of the bottleneck caused by the attempted eradication of rats on Henderson Island. The program implements three related methods (Pollak [23], Nei/Tajima [32], Jorde/Ryman [33]) and sets the minimum allele frequency accepted to four thresholds (5%, 2%, 1% and 0%), with resulting number of alleles considered in brackets. Resulting estimates of Ne are in bold with 95% confidence intervals in brackets. Confidence intervals are obtained both as parametric approximations and by jackknifing across loci. We present the parametric estimates which are generally a little tighter, particularly for the Jorde/Ryman method.

Minimum frequency	0.05 (61)	0.02 (75)	0.01 (85)	0 (97)
Pollak	<b>42.2</b> (23.5, 79.1)	<b>44.9</b> (26.1, 80.2)	<b>48.8</b> (29.0, 86.4)	<b>59.2</b> (35, 107.7)
Nei/Tajima	<b>45.6</b> (25.0, 87.7)	<b>47.4</b> (27.4, 86.3)	<b>50.8</b> (29.9, 91.0)	<b>59.7</b> (35.3, 109.0)
Jorde/Ryman	<b>35.8</b> (24.2, 49.6)	<b>35.3</b> (24.9, 47.4)	<b>35.1</b> (25.4, 46.5)	<b>35.3</b> (26.1, 45.0)

497  
498  
499  
500

501  
502 **Table 3.** The eight trial groups and the dose of brodifacoum presented in mg/kg of rat body weight  
503 (bw). Each dose was calculated individually for each rat based on its post-acclimatisation weight.  
504  
505

Poison Dose (mg/Kg bw)	Number of rats	Number surviving
0	8	8
0.025	8	7
0.05	8	5
0.1	8	1
0.2	8	0
0.35	8	0
0.55	5	0
0.80	5	0

506  
507