

# **A systematic review and meta-analysis on the effects of exercise training *versus* hypocaloric diet: distinct effects on body weight and visceral adipose tissue**

SHORT TITLE: Distinct effects of exercise training *versus* diet on weight and visceral fat

REBECCA J.H.M. VERHEGGEN, MD, MSc<sup>1</sup>

MARTIJN F.H. MAESSEN, MSc<sup>1</sup>

DANIEL J. GREEN, PhD<sup>2,3</sup>

AD R.M.M. HERMUS, MD, PhD<sup>4</sup>

MARIA T.E. HOPMAN, MD, PhD<sup>1</sup>

DICK H.T. THIJSEN, PhD<sup>1,2</sup>

<sup>1</sup>*Department of Physiology, Radboud University Medical Centre, Nijmegen, the Netherlands;*

<sup>2</sup>*Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, United Kingdom;* <sup>3</sup>*School of Sport Science, Exercise and Health, the University of Western Australia, Crawley, Western Australia, Australia.* <sup>4</sup>*Department of Internal Medicine, Division of Endocrinology, Radboud University Medical Centre, Nijmegen, the Netherlands*

***Total Word Count: 3505***

***Word Count Abstract: 199***

***Total Number of Tables: 2***

## **Author for correspondence:**

Rebecca Verheggen, MD, Msc, Dept. of Physiology (392), Radboud University Medical Center, PO Box 9101, 6500 HB Nijmegen, The Netherlands

e-mail: rebecca.verheggen@radboudumc.nl

Tel. (+31) (0)24 36 14906

Fax. (+31) (0)24 354 0535

## **Conflict of interest:**

All authors have no potential conflict of interest to report.

## ABSTRACT

Exercise training (“exercise”) and hypocaloric diet (“diet”) are frequently prescribed for weight loss in obesity. Whilst body weight changes are commonly used to evaluate lifestyle interventions, visceral adiposity (VAT) is a more relevant and stronger predictor for morbidity and mortality. We performed a meta-analysis to assess the effects of exercise training or hypocaloric diet on VAT (quantified by radiographic imaging). Relevant databases were searched through May 2014. 117 Studies (n=4,815) were included. We found that both exercise and diet cause VAT loss ( $P<0.0001$ ). When comparing diet *versus* training, diet caused a larger weight loss ( $P=0.04$ ). In contrast, a trend was observed towards a larger VAT decrease in exercise ( $P=0.08$ ). Changes in weight and VAT showed a strong correlation after diet ( $R^2=0.737$ ,  $P<0.001$ ), and a modest correlation after exercise ( $R^2=0.451$ ,  $P<0.001$ ). In the absence of weight loss, exercise is related to 6.1% decrease in VAT, whilst diet showed virtually no change (1.1%). In conclusion, both exercise and diet reduce VAT. Despite a larger effect of diet on total body weight loss, exercise tends to have superior effects in reducing VAT. Finally, total body weight loss does not necessarily reflect changes in VAT and may represent a poor marker when evaluating benefits of lifestyle-interventions.

**Keywords:** obesity, visceral adipose tissue, exercise training, hypocaloric diet

## INTRODUCTION

The worldwide prevalence of obesity, characterized by an excess in adipose tissue, has grown to pandemic proportions.<sup>1, 2</sup> Multiple reviews have demonstrated that accumulation of adipose tissue in general, and in the visceral area in particular, is strongly and positively correlated with all-cause morbidity and mortality.<sup>3</sup> Since obesity is an important, but also modifiable, risk factor for cardiovascular and metabolic diseases<sup>4, 5</sup> the WHO has recommended lifestyle interventions to aim at 5-10% reduction in body weight as treatment for obesity.<sup>6</sup>

Caloric restriction and exercise training cause a reduction in body weight by inducing a negative energy balance in which energy expenditure exceeds caloric intake. When comparing hypocaloric diet and exercise training, previous meta-analyses revealed that dietary restriction has superior effects on weight reduction.<sup>7, 8</sup> However, a growing body of evidence shows that excess visceral adipose tissue (VAT) may result in more detrimental obesity-related health effects than excess body weight.<sup>9</sup> Indeed, increased VAT is strongly associated with insulin resistance, atherogenic dyslipidemia, and cardiovascular disease.<sup>3, 10, 11</sup> Moreover, a reduction in VAT improves cardiovascular and metabolic risk.<sup>3, 12</sup> Hence, changing VAT is considered to be more important than weight reduction in the management of obesity.

In patients with obesity, physical exercise training leads to a healthier metabolic and cardiovascular phenotype.<sup>13-15</sup> Whilst exercise training does not always aim to reduce body weight, exercise training in general and aerobic exercise training in particular, have potent effects on reducing VAT.<sup>16-18</sup> Previous meta-analyses have evaluated only the effects of caloric restriction and aerobic exercise on weight loss. The effects of these interventions on VAT have not yet been compared. Therefore, we aim to conduct a systematic review and

meta-analysis to investigate the effect of caloric restriction *versus* aerobic exercise training on visceral adiposity loss in overweight and obese adults. For this purpose, we included studies that examined VAT after: 1. Caloric restriction only, 2. Exercise training only, and 3. Aerobic exercise training *versus* caloric restriction. We hypothesize that, in marked contrast to body weight loss, caloric restriction and exercise training have comparable effects on reducing VAT. With the use of a meta-regression analysis we aim to further explore the impact of intervention (e.g. duration, intensity, frequency) and subject (e.g. age, sex, baseline body weight) characteristics on the magnitude of changes in VAT.

Several international guidelines recommend lifestyle interventions aimed at a reduction in body weight of 5-10% as treatment for obesity.<sup>6, 19, 20</sup> Previous work, however, demonstrated that a reduction in body weight is a poor marker for VAT change.<sup>9</sup> Accordingly, changes in VAT may occur irrespective of changes in body weight. A hypocaloric diet causes a reduction in skeletal muscle mass, which along with a reduction in fat mass, contributes to weight loss.<sup>21, 22</sup> Aerobic exercise training, however, may be associated with an increase in lean body mass and/or plasma volume.<sup>23-25</sup> Assuming that fat mass decreases with exercise training, training may still not lead to weight loss<sup>24, 26</sup> Therefore, we hypothesize that the relation between changes in body weight and changes in VAT differs between caloric restriction and exercise interventions.

## **METHODS**

### **Data sources and searches**

The systematic literature search and documentation of literature was performed with the use of the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA)

statement.<sup>27</sup> Databases systematically searched were Pubmed, Cochrane, Web of Science and Embase. The following search strategy was used, with adaption for each database:

```
((("Energy Intake"OR"Diet Therapy"OR"(calori*AND restrict*)"OR(low AND
calori*)OR"dietary intervention*"OR"diet intervention*)"AND
("Overweight"OR"obes*")AND("Abdominal Fat"OR("Adipose Tissue"AND("intra-
abdom"*ORintraabdom*ORabdom*ORvisceral*))OR"Body Composition"OR"abdominal
adipos*ORvisceral adipos*ORintra-abdominal fat"OR"abdominal fat"OR"total body
fat"OR"adipose tissue distribution))OR(("Overweight"OR"obes*")AND("Motor
Activity"OR"Exercise"OR"Running"OR"Swimming"OR"Walking"OR"Warm-Up
Exercise"OR"Exercise Therapy"OR"Motion Therapy, Continuous Passive"OR"Sports"OR
"Athletic Performance"OR"Bicycling"OR"Physical Exertion"OR"running"
OR"bicycling"OR"cycling"OR"walking"OR"swimming"OR"training"OR"physical
activity"OR"exercis*"OR"cardio-training"))AND("Abdominal Fat"OR("Adipose Tissue"
AND (intra-abdom*OR"intraabdom*"OR"abdom*"OR"visceral*"))OR"Body Composition"
OR"abdominal adipos*"OR"visceral adipos*"OR"intra-abdominal fat"OR"abdominal
fat"OR"total body fat"OR"adipose tissue distribution")). Randomized Controlled Trials
(RCTs), Non-randomized Controlled Trials (non-RCTs) OR Clinical Trials (CT) published in
English, German and Dutch were included from January 1th, 1987 to May 5th, 2014.
Reference lists of included articles were manually checked by RV for possible eligible studies
that were missed during the literature search (Figure 1). This represents a valid and frequently
used method to further increase the number of potentially eligible studies.
```

## Study selection

To standardize the selection procedure by two independent reviewers, investigators received the local review protocol previous to the selection of studies,. After the elimination of

duplicates, one investigator (RV) screened study titles for eligibility with use of the in- and exclusion criteria in the review protocol, which are listed below. Two reviewers (RV, MM) independently screened the abstracts of the remaining studies. 389 studies were assessed in full text (Figure 1). Inter-reviewer disagreements were resolved through consensus or by consulting a third reviewer (MH). When study characteristics or viable information was missing, an attempt was made to request missing information from the authors by email (n=6 studies; authors of n=2 studies provided requested information). Studies were included when the mean age at entry was  $\geq 18$  years and mean BMI was  $\geq 25$  kg/m<sup>2</sup>. Studies of HIV-infected individuals were excluded because of the interference of anti-retroviral drugs with abdominal adipose tissue.<sup>16</sup> Because spinal cord injuries associated with changes in body composition, studies conducted in spinal cord injured individuals were also excluded.<sup>28</sup> Studies with one or more arms assigned to an aerobic exercise intervention or a hypocaloric diet were eligible for inclusion. For the first aim, Clinical Trials and Randomized Controlled Trials (RCTs) with one arm assigned to exercise *or* caloric restriction were selected. Furthermore, in order to directly compare duration- and energy deficit- matched exercise training with caloric restriction, RCT's with an exercise- *and* a diet-arm were included. To identify exercise and subject characteristics that predict the magnitude of change in VAT using the meta-regression analysis, clinical trials and RCT's with one arm assigned to exercise *or* caloric restriction were selected. Finally, diet and exercise studies that provided baseline and post-intervention results for VAT and weight were included for the correlation analysis. Exercise training was defined as a program including voluntary aerobic exercise at a low to vigorous intensity for at least two times per week during a minimum period of four weeks and with a minimum duration of 20 minutes per session. Caloric restriction was defined as a daily reduction in energy (caloric) intake of at least 10% of the habitual intake (2000 kcal for women, 2500 kcal for men) during a minimum period of four weeks. Interventions combining exercise and diet

therapy or adding resistance exercise or bariatric surgery to an intervention arm were excluded. Studies in which a pharmacological dietary supplement was used were excluded from our analysis. Studies were eligible when VAT was measured with the use of Computerized Tomography (CT) or Magnetic Resonance Imaging (MRI), which are both considered to be the gold standard for the quantitative measurement of VAT<sup>29</sup>. Studies that used another measurement technique were excluded.

### **Data extraction and Quality assessment**

Baseline and post-intervention mean VAT area or volume and standard deviation or standard error was independently recorded by two authors (RV, MM). When VAT was measured at multiple sites, the measurement at the 4<sup>th</sup> and/or 5<sup>th</sup> lumbar vertebrae was recorded for further analysis, since this region is most strongly correlated with body adiposity.<sup>28</sup> Based on changes in visceral abdominal fat area or volume, percentage change in VAT for each study was calculated by one of the authors (RV) for the correlation analysis. Percentage weight loss was also calculated based on pre- and post-intervention values. Furthermore, publication year, journal, study design, sample size, age, sex, weight, BMI, and intervention details (duration, intensity and frequency (exercise studies), caloric deficit (diet studies) were extracted from all included studies. When results were depicted in figures only (n=14 studies), data were extracted with the use of GetData Graph Digitizer. A request by email was sent to the authors, when key information was not included in the published manuscript (n=6 studies). Two out of six authors responded to our repeated email requests, thus the remaining 4 studies were excluded from further analysis.

The quality of each eligible study was independently assessed by two authors (RV and MM), with the use of a modified version of The Critical Review Form for Quantitative Studies, from

Law *et al.*<sup>30</sup>. One item (“contamination was avoided”) was not applicable for the studies included in this meta-analysis and was therefore removed for analysis. Only studies with a minimum score of 10 out of 14 items were eligible for inclusion (Figure 1).

### **Data synthesis and analysis**

To account for potential heterogeneity between studies, a random-effects model (specified *a priori*) was used to determine the overall effect size of the intervention (exercise training or hypocaloric diet) on VAT. Effect sizes for RCTs and clinical trials were calculated as the standardized mean difference (SMD) with corresponding 95%-CI. A correlation of 0.5 between the outcomes measured in each study arm (*i.e.* exercise, diet, or control) was assumed. When a study contained multiple study arms, all were included in the statistical analysis, whereby the different intervention groups were individually compared against the control group. Analyses to assess the following comparisons: (1) diet *versus* control; (2) exercise *versus* control; (3) diet *versus* exercise were performed. The Cochrane’s Q statistic and  $I^2$  were calculated to assess the degree of heterogeneity across studies. Publication bias was assessed using visual analysis of the funnel plot asymmetry using the ‘trim and fill’ and the ‘Classic fail ‘n safe’ algorithms. All calculations and plots were performed in CMA-2 (Comprehensive Meta-Analysis 2nd version, Biostat, Englewood, NJ, USA).

### **Meta-regression analysis**

To assess the effects of subject and intervention characteristics on VAT loss, random-effects meta-regression analysis with SMD as dependent variable was calculated. The weighted inverse variance (with correction for total n) was used as weight factor. For the purpose of meta-regression analysis, the aerobic exercise arms (n=86) were separated from the hypocaloric diet arms (n=87). In both study types, duration of the exercise training or diet



intervention (weeks), measurement technique (CT or MRI), body weight at baseline, age and sex were defined as a covariate. Duration was assessed as a categorical variable (duration <16 weeks *versus* duration of  $\geq 16$  weeks). In the exercise studies, intensity of the intervention was examined as a covariate. Intensity was categorized in ‘vigorous intensity’ (i.e. largely performed at 70% of maximal heart rate (maxHR) or >55% of maximal oxygen uptake (VO<sub>2</sub>max) or 60-80% of the heart rate reserve (HRR)), ‘moderate intensity’ (60-70% of maxHR, 45-55% of VO<sub>2</sub>max or at the lactate threshold), and ‘low intensity’ (<60% of HRmax or <45% of VO<sub>2</sub>max) based on previous work.<sup>18</sup> This categorization is somewhat different from the often used and more practical categories based on METs as proposed in the ACSM and AHA guidelines.<sup>31, 32</sup> However, only two studies included in our meta-analysis provided data on METs. Since translating percentages of maximum heart rate, heart rate reserve and VO<sub>2</sub>max into METs proves difficult with the use of average data on group level, we adopted the aforementioned strategy to divide studies based on intensity. In hypocaloric diet studies, “intensity” was divided in ‘very low calorie diets’ (VLCDs; reduction to maximal 800 kcal/day) and ‘low calorie diets (LCDs; caloric restriction to 800-2000 kcal/day). Lastly, frequency (times spent in training per week) was added as covariate in exercise studies.

### *Correlation analysis*

To examine correlations between weight loss and VAT improvement a Pearson correlation coefficient was calculated. The formula of the corresponding trend line was retrieved with the use of linear regression. Meta-regression analyses and correlation analysis were conducted with use of SPSS version 20.0.

## **RESULTS**

### *Selection of studies for the meta-analysis*

The original search resulted in 15,964 studies. Eleven more studies were found from the reference lists of the included full text papers. After removal of duplicates and elimination of papers based on the eligibility criteria and quality assessment, 50 aerobic exercise studies and 59 hypocaloric diet studies were included (Figure 1). For the analysis of a direct comparison between caloric restriction and exercise training, 8 studies were included. (Table 1) One study, which directly compared exercise training with caloric restriction was excluded as duration and energy deficit did not match between the two intervention arms. This study was included for the separate analysis of diet or exercise training only.

#### *Cohort characteristics*

A total number of 4,815 individuals (2,404 in the exercise studies, 2,411 in the hypocaloric diet studies) participated in the interventions. (Table 1) In the 8 RCT's that directly compared exercise training and caloric restriction, a total of 400 individuals were included (200 in the exercise arm, 200 in the diet arm). (Table 2) 28 studies exclusively included male subjects, whereas in 39 studies females were exclusively included. 55 studies included both sexes. Some studies recruited specific populations, which included older (aged 50-80 years) individuals (n=4); patients with type 2 diabetes (n=11), impaired glucose tolerance (n=3) and metabolic syndrome (n=3). (Table 1)

#### *Meta-analysis*

The SMD of change in VAT after exercise training was -0.47 (95%CI -0.56 to -0.39,  $P<0.0001$ ). (Figure S1) Heterogeneity analysis showed significant heterogeneity (Cochran's  $Q=265.4$ ;  $P=68.0$ ). Through a funnel plot of standard error by Hedge's  $g$  and the Trim 'n Fill algorithm, publication bias was assessed. With the use of the Classic Fail 'n Safe approach, it became clear that there was no significant publication bias since 7427 missing studies would

be required to achieve a p-value above 0.05. The SMD of change in VAT after caloric restriction was  $-0.63$  (95%CI  $-0.71$  to  $-0.55$ ,  $P<0.0001$ ) (Figure S2), whilst significant heterogeneity was present (Cochran's  $Q=236.0$ ;  $I^2=63.6$ ). In these studies, no publication bias was present since there would be 4096 studies required to achieve a p-value above 0.05. Based on the studies that directly compared exercise training and caloric restriction, exercise training caused a non-significantly larger decrease in VAT ( $-0.59$ , 95%CI  $-1.248$  to  $0.071$ ;  $P=0.08$ ), whilst caloric restriction caused a significantly larger weight loss than exercise training (SMD  $0.308$ , 95%CI  $0.02$  to  $0.60$ ;  $P=0.04$ ) (Figure 2). Heterogeneity analysis showed significant heterogeneity (Cochran's  $Q=51.9$ ;  $I^2=86.5$ ). Publication bias was assessed with the Trim 'n Fill method and showed no change in SMD when adding trimmed studies, for both the weight loss as VAT loss data.

#### *Meta-regression analysis*

No effect of measurement technique on the SMD was observed for studies that performed exercise training or diet (data not shown). In the exercise studies, univariate analysis revealed that the SMD for VAT improvement was significantly influenced by sex ( $R^2=0.11$ ; 95%CI  $0.06 - 0.472$ ;  $P=0.012$ ); duration ( $R^2=0.073$ ; 95%CI  $-0.449$  to  $-0.055$ ;  $P=0.013$ ) and frequency ( $R^2=0.084$ ; 95%CI  $-0.157$  to  $-0.030$ ;  $P=0.004$ ). The multivariate regression analysis, which included the factors that revealed a significant impact in the univariate analysis, identified an impact of male sex on SMD ( $R^2=0.20$ ; 95%CI  $0.066$  to  $0.467$ ;  $P=0.01$ ). In hypocaloric diet studies, univariate analysis showed a significant effect of male sex only ( $R^2=0.09$ ; 95%CI  $0.116$  to  $0.632$ ;  $P=0.005$ ).

#### *Correlation analysis*

For exercise studies, a moderate correlation was found between changes in weight versus changes in VAT after exercise training ( $R^2=0.453$ ,  $P<0.001$ ), whilst diet-interventions showed a strong correlation between the change in weight versus change in VAT ( $R^2=0.737$ ,  $P<0.001$ ) after caloric restriction. Exercise training showed a somewhat steeper slope compared to diet ( $-3.04x$  versus  $-2.41x$ , respectively), and a larger Y-axis intercept ( $-6.1$  versus  $-1.1$ , respectively, Figure 3).

## DISCUSSION

The present work is the first meta-analysis to compare the effect of caloric restriction and aerobic exercise training on visceral adipose tissue (VAT) loss in overweight and obese individuals. We present the following findings. First, our results confirm that both caloric restriction and exercise training successfully reduce VAT. Second, in studies that provided a direct comparison of caloric restriction and exercise training, a hypocaloric diet resulted in significantly larger weight loss. Interestingly, these studies reveal a different story for VAT. Exercise training tends to show a larger decrease in VAT compared to caloric restriction. The distinct effects of both interventions on total body weight and VAT are supported by the correlation analysis. Only a moderate correlation was found for the exercise training cohort between changes in weight and VAT. Furthermore, in the absence of weight loss, exercise training results in a 6.1% decrease in VAT, whilst a hypocaloric diet leads to virtually no change (1.1%). This suggests that evaluating only total body weight changes could lead to spurious conclusions when evaluating the efficacy of a lifestyle intervention in overweight and obese individuals since health benefits occur independent of body weight changes. Indeed, even in the absence of weight loss after exercise training, health benefits such as a reduction in VAT are present.

In line with previous meta-analyses, we found caloric restriction to have a larger effect on weight loss than exercise training.<sup>7, 8</sup> We extended this finding by a direct comparison of studies with matched duration and energy deficit in order to more accurately compare the impact of both interventions. In marked contrast to the superior effect of caloric restriction on weight loss, no difference in VAT reduction was observed between caloric restriction and exercise training. In fact, exercise training tended to have a superior effect on VAT reduction compared to caloric restriction. A possible mechanism underlying these different effects on weight and VAT could relate to distinct changes in body composition during these lifestyle interventions. During caloric restriction, both muscle mass and fat mass are lost, resulting in a marked decline in weight.<sup>21, 22</sup> During exercise training, however, lean body mass and circulating plasma volume increase, whilst fat mass decreases.<sup>21, 23, 25, 26, 33</sup> Previous work that directly measured these factors indeed showed that an increase in lean body mass counteracts loss of fat mass after 8 weeks of exercise training.<sup>34</sup> These opposing effects resulted in the absence of total body weight loss.<sup>34</sup> Appreciating and understanding these effects are important to acknowledge that exercise training effectively reduces VAT, despite the absence of a reduction in body weight.

In this meta-analysis, a large number of studies were included. Multivariate meta-regression analysis on these data showed that male sex is associated with a larger decrease in VAT, in both exercise and diet interventions. Other subject and intervention characteristics did not influence the magnitude of VAT loss in the multivariate model. A possible explanation that underlies the larger effect of lifestyle interventions on VAT in men is that men typically have larger VAT stores than women. As a result, this makes male participants more likely to lose VAT than female participants.<sup>35</sup> However, our meta-regression analysis showed no effect of

baseline VAT area on the magnitude of VAT decrease. The exact underlying mechanisms should be subject for future research.

The distinct effects of diet and exercise training on weight and VAT suggest the presence of a different correlation between changes in body weight and VAT after caloric restriction in comparison to exercise training. Indeed, whilst a strong correlation between changes in body weight and VAT was found after caloric restriction, this correlation was only moderate for exercise training studies. This means that a change in weight after hypocaloric diet predicts a substantial effect on VAT, whereas changes in weight after exercise training only modestly predict the change in VAT. Furthermore, the trend lines for these correlations show important differences. The Y-intercept for the correlation of exercise studies is 6.1%, meaning that the absence of weight loss after exercise training is still correlated with a significant and meaningful reduction in VAT of 6.1%. In marked contrast, studies examining the impact of hypocaloric diet revealed a Y-intercept of only 1.1%, which means that in the absence of weight loss only 1.1% of VAT is lost. Furthermore, the steepness of the correlation for exercise training is slightly higher than that observed after hypocaloric diet. Taken together, these data strongly indicate that a change in weight, which is currently recommended by international guidelines for the management of obesity, does not necessarily reflect changes in VAT.

### *Limitations*

The presence of heterogeneity of the included studies may represent a potential limitation when interpreting the results of this review. However, to correct for this heterogeneity a random effect approach was selected to perform the meta-analyses, which was specified *a priori*. Furthermore, analysis of publication bias with use of the Classic Fail 'n Safe method

showed that an unrealistically large number of studies is needed to influence the significant results obtained in our meta-analyses. Therefore, we are confident that the heterogeneity observed in the studies included in this analysis does not impact the major conclusions of our study. Another limitation might be that information about the ethnicity of participants in the included studies is lacking. Often, this information was not available in the original papers. However, we included studies that were conducted on all different continents, with exception for the African continent. We therefore believe that this meta-analysis provides conclusions applicable for every ethnic group.

### *Clinical relevance*

As treatment for obesity, international guidelines including WHO and ACSM guidelines, recommend a minimum of 5% loss of body weight loss.<sup>4, 6, 20</sup> Although in common clinical practice a combination of training and hypocaloric diet is often prescribed, it is highly relevant to understand the separate effects of these interventions. Indeed, our study reveals that effects on weight loss and VAT loss are different in training and diet interventions. For example, a 5% reduction in body weight after hypocaloric diet has a different effect on VAT than a similar reduction in body weight after exercise training. Indeed, 5% loss in body weight is associated with 21.3% reduction in VAT after exercise training, but only with 13.4% reduction in VAT after a hypocaloric diet. To reduce VAT by 13.4% after exercise training, weight loss of only 2.4% is needed. Moreover, the absence of a reduction in body weight after exercise training may lead physicians to *incorrectly* conclude that the intervention has failed. This is in accordance with the ACSM position statement on appropriate physical intervention strategies for weight loss, which also emphasized that exercise training entails health benefits in the absence of weight loss.<sup>20</sup> In fact, it is likely that a clinically relevant VAT reduction (of 6.1%) is present in the absence of weight loss after exercise training, which may lead to

reductions in cardiovascular risk and improvement in metabolic health. Therefore, it seems incorrect to recommend a 5% weight loss for all lifestyle interventions.

In conclusion, our systematic review and meta-analysis provide evidence that exercise training, despite smaller effects on reducing body weight, tends to have superior effects on reducing visceral adipose tissue compared to diet interventions in overweight and obese subjects. This suggests that changes in body weight represent a poor marker for adaptation in visceral adipose tissue, especially when performing exercise training. Our data therefore strongly indicate that, in clinical practice, caution should be taken when interpreting (lack in) changes of body weight after exercise training interventions. Incorrect conclusions can potentially lead to recommendations or suggestions that the exercise intervention was unsuccessful, despite the presence of a marked effect on body composition. Setting the correct targets for evaluating the health benefits of lifestyle interventions is therefore recommended.

### **Acknowledgements**

We would like to thank Alice Tillema, BSc. for her assistance during the literature search. We would also like to thank Hans Groenewoud, MSc. for his assistance with the statistical analyses.

### **Authors' contributions statement**

RJV, DJG, MTH, DHT: conception and design of research. RJV and MFM: data acquisition and analysis. RJV, MFM, ARH, MTH, DHT: interpretation results of research. RJV and MFM: preparation figures. RJV drafted manuscript. All authors edited and revised manuscript. All authors approved final version of manuscript and agree to be accountable for all aspects of the work.



**Supporting Information:**

**Figure S1.** Forest plot of the effect size (SMD) of exercise training on VAT loss. The effect size (SMD) and 95% CI for individual studies and the pooled estimate (assessed with the use of Random Effects Model) are depicted.

**Figure S2.** Forest plot of the effect size (SMD) of hypocaloric diet on VAT loss. The effect size (SMD) and 95% CI for individual studies and the pooled estimate (assessed with the use of Random Effects Model) are depicted.

**FIGURE LEGENDS**

**FIGURE 1.** PRISM Flowchart of outcomes of search strategy

**FIGURE 2.** Forest plot of the effect size (SMD) of (A) exercise training versus caloric restriction on weight loss and (B) exercise training versus caloric restriction on VAT loss. The effect size (SMD) and 95% CI for individual studies and the pooled estimate (assessed with the use of Random Effects Model) are depicted.

**FIGURE 3.** Correlation between % VAT improvement and % weight loss for exercise studies ( $R^2=0.4531$ ,  $P<0.001$ ; trendline:  $y = -3.03x - 6.1$ ), and caloric restriction studies ( $R^2=0.737$ ,  $P<0.001$ ; trendline:  $y = -2.46x - 1.1$ ).

Table 1 Overview of the characteristics of the included Exercise training (n=50) and Hypocaloric diet (n=59) studies.

Exercise Training Studies										
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Dura- tion (weeks)	Assess- ment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Baria et al. <sup>36</sup>	Centre-based exercise	10 (10/0)	52.1 ± 11.4	30.8 ± 5.1	Personal ventilatory threshold	3x/week / 30-60min	12	CT L4-L5 (mm)	113.1 ± 24.1 / 106.6 ± 22.8	86.2 ± 19.4 / 86.1 ± 20.7
	Home-based exercise	8 (8/0)	50.8 ± 7.7	30.9 ± 3.9					115.2 ± 20.5 / 107.4 ± 17.0	90.9 ± 12.4 / 89.3 ± 11.9
	Control	10 (10/0)	53.4 ± 9.6	29. ± 1.9					92.1 ± 25.9 / 97.0 ± 23.9	84.8 ± 7.8 / +1.5kg
Boudou et al. <sup>37</sup>	Exercise	8 (8/0)	42.9 ± 5.2	28.3 ± 3.9	75% of VO <sub>2</sub> peak & 5 x 2 min 85% of VO <sub>2</sub> peak alternated by 3x 50% of VO <sub>2</sub> peak	2x/week / 45 min & 1x/week / 19 min	8	MRI L4-L5 (cm <sup>2</sup> )	153.25± 38.55/ 84.20±21.30	86.90 ± 13.4/ 85.00 ±13.8
	Control	8 (8/0)	47.9±8.35	30.85±5.2					156.85±23.40 / 150.35±23.3	85 ± 13.75 / 88.75 ± 1.30
Cho et al. <sup>38</sup>	Low-intensity exercise	15 (0/15)	42.4 ± 7.6	25.6 ± 1.7	40-50% of VO <sub>2</sub> max 70-75% of VO <sub>2</sub> max	3/week (duration depending on energy expenditure)	12	CT L4-L5 (cm <sup>2</sup> )	99 ± 41 / 79 ± 40	64.4 ± 6.0 / 62.3 ± 5.5
	High-intensity exercise	15 (0/15)	45.6 ± 4.6	25.1 ± 2.0					90 ± 26 / 83 ± 30	63.2 ± 6.4 / 60.4 ± 6.4
	Control	15 (0/15)	49.2 ± 8.7	26.1 ± 2.7					106 ± 33 / 103 ± 28	63.0 ± 7.8 / 64.5 ± 7.0
Cuff et al. <sup>39</sup>	Aerobic exercise	9 (0/9)	59.4 ± 1.9	32.5 ± 1.4	60-75% maxHR	3x /week / 75 min	16	CT L4-L5 (cm <sup>2</sup> )	215.7 (25.8) / -8.8 (5.4)	81.2 ± 3.8 / -1.2 (0.7)
	Control	9 (0/9)	60.0 ± 2.9	36.7 ± 2.0					225.8 (8.9) / -0.4 (12.0)	95.6 ± 6.5 / +2.0 (1.2)
Davidson et al. <sup>40</sup>	Aerobic exercise	37 (17/20)	68.8±6.0(m) 69.1±6.5 (f)	29.9 ± 3.0 29.2 ± 3.7	60-75% of VO <sub>2</sub> peak	7x/week / 30 min	26	MRI (kg)	-11.0 (1.9)%	NR / -2.7 (3.1)%
	Control	28 (11/17)	67.4±3.8(m) 66.7 ± 3.7(f)	30.5 ± 2.0 30.4 ± 3.2					-0.7 (2.5)%	NR / -0.1 (0.7)%
Dekker et al. <sup>41</sup>	Obese with T2D	8 (8/0)	51.0 ( 3.0)	29.9 (1.2)	60% VO <sub>2</sub> max	5x/week / 60 min	12	MRI L4-L5 (kg)	3.8 (0.3) / 3.1(0.4)	93.5 (2.9) / 93.9 (3.2)
	Obese	8 (8/0)	47.1 (3.1)	32.4 (0.6)					4.0 (0.4) /3.4 (0.4)	97.6 (3.4) / 97.2 (0.6)
Despres et al. <sup>42</sup>	Aerobic exercise	13 (0/13)	38.8 ± 5.3	34.5 ± 4.3	55% of VO <sub>2</sub> max	4-5x/week / 90 min	12	CT L4-L5 (cm <sup>2</sup> )	124.7 ± 48.6 / 121.3 ± 45.5	90.0 ± 11.8 / 86.3 ± 9.6

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Dura- tion (weeks)	Assess- ment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Dipietro et al.</b> <sup>43</sup>	Aerobic exercise	9 (2/7)	72 (1)	27.5 (2.7)	55% of maxHR during 1 month 75% during 3 months	4x/week / 40-60min	17	CT L4-L5 (cm <sup>2</sup> )	116 (31) / 106 (24)	65 (5) / 64(4)
	Control	7 (1/6)	73 (1)	26.8 (1.5)					136(28) / 118 (27)	69 (4) / 69(4)
<b>Donges et al.</b> <sup>44</sup>	Aerobic exercise	13 (13/0)	45.4 (1.7)	32.0 (1.3)	75% of age-predicted maxHR during first 4 weeks, thereafter 80% of maxHR	3x/week / 40-50min	12	CT L4 (cm <sup>2</sup> )	1371 (113) / 1222 (100)	103.1 (4.6) / -1.9 (0.7)%
	Control	8 (8/0)	49.5 (2.6)	29.6 (2.1)					1383 (164) / 1349 (145)	92.2 (6.9) / + 0.1(0.6)%
<b>Donges et al.</b> <sup>45</sup>	Aerobic	41 (16/25)	NR	30.0 ± 5.5	70-75% of max HR	3x/week / 30-50min	10	DEXA (kg)	1.49 ± 0.55 / 1.38 ± 0.58	84.8 ± 18.6 / -0.8 ± 1.9
	Control	26 (13/13)		28.3 ± 4.1					1.44 ± 0.43 / 1.44 ± 0.45	83.2 ± 13.4 / +0.6 ± 1.3kg
<b>Donnelly et al.</b> <sup>46</sup>	Exercise Group men	16 (16/0)	22 ± 4	29.7 ± 2.9	60% of HRR with a gradual increase to 75% at 6 months	5x/week / 20-45min	65	CT L4-L5 (cm <sup>2</sup> )	97.9 ± 22.5 / 75.5 ± 18.3	94.0 ± 12.6 / 88.8 ± 9.5
	Control Group men	15 (15/0)	24 ± 4	29.0 ± 3.0					91.7 ± 29.7 / 85.4 ± 39.7	94.1 ± 11.4 / 93.16 ± 11.6
	Exercise group women	25 (25/0)	24 ± 5	28.7 ± 3.2					60.6 ± 25.5 / 57.4 ± 28.4	77.0 ± 11.4 / 77.6 ± 12.8
	Control group women	18 (18/0)	21 ± 4	29.3 ± 2.3					62.9 ± 21.8 / 66.0 ± 13.9	79.9 ± 8.1 / 82.8 ± 9.2
<b>Friedenreich et al.</b> <sup>47</sup>	Aerobic exercise	160 (0/160)	61.2±5.4	29.14.5	70-80% of maxHR	3.6x/week / 45min	52	CT umbilicus (cm <sup>2</sup> )	101.4 ± 55.4 / -16.5	75.6 ± 13.0 / -2.3
	Control	160 (0/160)	60.6±5.7	29.3 ± 4.3					103.2 ± 56 / -1.6	76.3 ± 12.7 / -0.5
<b>Gan et al.</b> <sup>48</sup>	Aerobic exercise	18 (0/18)	37.4 (1.3)	30.9 (0.7)	55-70% VO2max	4-5x/week / 40min	9.7	MRI L4-L5 (l)	2.23 (0.12) / 2.11 (0.12)	94.1 (2.0) / 92.8 (2.0)
<b>Giannopoulos et al.</b> <sup>49</sup>	Aerobic exercise	11 (0/11)	57 (entire study)	35.9 (2.2)	70% HRR	3x/week / 60-75min	12	MRI (cm <sup>3</sup> )	5204 (598) / 4675 (550)	92.9 (6) / 91.2 (5.6)

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Halverstad t et al. <sup>50</sup>	Aerobic exercise	83 (34/49)	57.9 ± 0.6	36.0 ± 1.1 (% body fat)	50% of VO <sub>2</sub> max with a gradual increase to 70% of VO <sub>2</sub> max (for at least 14 w)	3x/week / 20-40min and addition of one extra low intensity exercise session	24	CT (cm <sup>2</sup> )	127.8 (4.5) / -14.4 (2.4)	80.6 ± 1.6 / -1.1 (0.3)
Haus et al. <sup>51</sup>	Aerobic exercise	16 (5/11)	65 ± 1	33 ± 1	60-65% max HR with a gradual increase to 80-85% week 4	5x/week / 50- 60min	12	CT (cm <sup>2</sup> )	182.4 ± 21.5 / 134.5 ± 15.9	95.7 ± 4.1 / 91.9 ± 3.8
Heydari et al. <sup>52</sup>	Aerobic exercise	25 (25/0)	24.7 ± 4.8	28.4 ± 0.5	80-90% of max HR during 8 sec sprint, whereafter 12 sec recovery	3x/week / 20min	12	CT L4/L5 (g)	62.6 (6.2) / 51.8 (5.1)	87.8 ± 2.7 / 86.3 ± 2.7
	Control	21 (21/0)	25.1 ± 3.9	29 ± 0.9					69.7 (9.7) / 67.3 (8.4)	89 ± 2.9 / 89.4 (3.1)
Hutchison et al. <sup>53</sup>	Obese	8 (0/8)	NR	36.9 (2.1)	75-85% of maxHR OR HIIT: 6-8 x 5 minutes at 95- 100% of maxHR – 1-2 min recovery	3x/week / 60 min (alternating between HIIT and continuous)	12	CT L4-L5 (cm <sup>2</sup> )	135.1 (15.7) / 132.7 (18.1)	99.4 (5.4) / 96.9 (4.5)
	PCOS	14 (0/14)							119.5 (16.1) / 107.6 (15.1)	96.9 (4.8) / 95.3 (4.8)
Irving et al. <sup>54</sup>	Low-intensity exercise	11 (0/11)	51 ± 9 (entire group)	34.7 ± 7.5	At lactate threshold Midway between lactate threshold and VO <sub>2</sub> max (3 days) ; lactate threshold (2 days)	5x/week / duration depended on energy expenditure	16	CT L4/L5 (cm <sup>2</sup> )	153 ± 51 / 146 ± 49	97.2 ± 22 / 95.1 ± 19.3
	High-intensity exercise	9 (0/9)		34.7 ± 6.8					173 ± 73 / 148 ± 59	93.5 ± 18.3 / 90.0 ± 15.6
	Controls	7 (0/7)		32.7 ± 3.8					157 ± 71 / 155 ± 71	89.6 ± 11.2 / 88.7 ± 10.6

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Duration (weeks)	Assess- ment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Irwin et al.<sup>55</sup></b>	Exercise	87 (0/87)	60.7 ± 6.7	30.4 ± 4.1	Start 40% of maxHR with a gradual increase to 60-75% by week8	5x/week / 45min	52	CT L4-L5 (g/cm <sup>2</sup> )	147.6 (134.3-161) / -8.5	81.4 ± 14.1 / -1.3%
	Control	86 (0/86)	60.6 ± 6.8	30.5 ± 3.7					147.6 (135.4-159.8) / +0.1	81.7 ± 12.1 / 0.1%
<b>Janssen et al.<sup>56</sup></b>	Aerobic exercise in black men	84 (84/0)	Not depicted for entire groups	27.0 ± 4.8	75% of VO2max	3x/week / 50 min	20	CT L4-L5 (cm <sup>2</sup> )	77.5 ± 5.1 / 71.9 ± 52.2	83.9 ± 16.3 / -0.5 ± 2.4
	Aerobic exercise in white men	255 (255/0)		26.7 ± 4.9					109.5 ± 63.6 / 102.4 ± 61.2	84.3 ± 16.3 / -0.3 ± 2.1
	Aerobic exercise in black women	160 (0/160)		28.2 ± 6.1					69.1 ± 40.8 / 65.4 ± 37.9	73.8 ± 16.3 / -0.4 ± 3.0
	Aerobic exercise in white women	243 (0/243)		24.9 ± 4.8					75.4 ± 52.7 / 72.2 ± 49.1	67.0 ± 13.6 / -0.1 ± 2.1
<b>Johnson et al.<sup>57</sup></b>	Aerobic exercise	12 (N.R.)	49.1 (2.3)	32.2 (0.8)	Week 1: 50% of VO2p Week 2: 60% VO2p Week 3 and 4: 70% VO2p	3x/week / 30-45 min (interval:15 min training, 5 min rest)	4	MRI L4-L5 (cm <sup>2</sup> )	154.3 (18.3) / 143.6 (18.7)	94.4 (3.8) / 94.1 (4.0)
	Stretching control	7 (N.R.)	47.3 (3.6)	31.1 (1.1)					154.3 (21.2) / 158.6 (23.9)	98.8 (6.0) / 98.6 (6.3)
<b>Jung et al.<sup>58</sup></b>	Moderate intensity	8 (0/8)	56.8 ± 8.2	25.5 ± 1.5	Goal: intensity at 3.5- 5.2 METs Goal: intensity at > 5.3 METs	5x/week / 60min (moderate intensity) vs. 30min (vigourous intensity)	12	CT L4-L5 (cm <sup>2</sup> )	15784.6±4662.7 / 13262.5 ± 3217.8	63.7 ± 5.0 / -2.9% ± 2.5%
	Vigorous intensity	8 (0/8)	48.4 ± 6.1	25.9 ± 1.6					13726.6±3011.8 / 12447.4 ± 2252.6	62.9 ± 4.4 / -2.5% ± 2.3%
	Control	12 (0/12)	55.5 ± 7.6	27.7 ± 3.4					17790.2±5621.7 / 17372.7 ± 5235.7	67.3 ± 9.8 / -1.5% ± 1.6%
<b>Karstoft et al.<sup>59</sup></b>	Continuous walking group	12 (8/4)	60.8 ± 2.2	29.9 ± 1.6	>55% of peak energy expenditure Walking at 70% of peak energy expenditure for 3 minutes, alternated with 3 minutes of slow walking	5x/week / 60min	17	MRI, below diaphragm (l)	4.5 ± 0.3 / 4.2 ± 0.4	88.2 ± 4.7 / 87.5 ± 4.8
	Interval walking group	12 (7/5)	57.5 ± 2.4	29.0 ± 1.3					4.7 ± 0.8 / 4.2 ± 0.7	84.9 ± 4.9 / 80.7 ± 4.1
	Control group	8 (5/3)	57.1 ± 3.0	29.7 ± 1.9					4.7 ± 0.4 / 4.6 ± 0.4	88.5 ± 4.7 / 89.2 ± 5.2

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Kim et al. <sup>60</sup>	Aerobic exercise	24 (24/0)	49.4 ± 9.6	30.7 ± 3.3	Gradual increase of 50-60% of maxHR to 60-70%	3x/week / 60min	12	CT L4-L5 (cm <sup>2</sup> )	197.1 ± 61.9 / 165.7 ± 57.0	87.7 ± 11.2 / -4.2%
Ku et al. <sup>61</sup>	Aerobic exercise	15 (0/15)	55.7 ± 7.0	27.1 ± 2.4	40-50% of maximal exercise capacity	5x/week / 60min	12	CT (g)	15890 ± 4593 / 15038 ± 3369	66.3 ± 6.0 / -1.9 ± 1.2
	Control	16 (0/16)	57.8 ± 8.1	27.4 ± 2.8					17530 ± 4747 / 17362 ± 4728	67.6 ± 7.5 / -0.6 ± 0.7
Kwon et al. <sup>62</sup>	Aerobic exercise	13 (0/13)	55.5 ± 7.5	27.0 ± 2.5	Anaerobic threshold	5x/week / 60min	12	CT L4-L5	16291.5±4808/ 14682.7±3494	66.3 ± 6.5 / NR
	Control	14 (0/14)	57.5 ± 8.6	27.5 ± 3.0					17204.5±4674/ 17216.3±4560	68.0 ± 7.9 / NR
Lee et al. <sup>63</sup>	Obese	8 (8/0)	47.1 ± 8.1	32.4 ± 1.6	~60% of VO <sub>2</sub> peak	5x/week / 60min	13	MRI 5cm below to 15 cm above L4-L5 (kg)	9.2 ± 1.2 / 8.3 ± 1.1	97.6 ± 8.9 / 97.2 ± 8.9
	T2D	9 (9/0)	51.0 ± 8.0	29.9 ± 3.2					7.5 ± 1.3 / 6.7 ± 1.5	93.5 ± 7.6 / 93.9 ± 8.5
Liao et al. <sup>64</sup>	Aerobic exercise	32 (9/23)	55.8 (1.8)	25.6 (0.8)	50% of HRR, with a gradual increase to 70%	3x/week / 60min	26	CT (cm <sup>2</sup> )	86.3 (8.1) / -16.1	66.1 (2.9) / -2.7 (0.4)
	Stretching control group	32 (17/15)	52.2 (1.8)	26.6 (0.8)					112.3 (9.9) / -14.5	69.7 (2.6) / -0.9 (0.3)
Malin et al. <sup>65</sup>	Aerobic exercise	35 (16/19)	66.8 ± 0.8	35.1 ± 0.7	60-65% of maxHR first four weeks, thereafter increase to 80-85%	5x/week / 50-60min	12	CT (cm <sup>3</sup> )	151.4 (14) / -30.6	99.0 ± 2.4 / - 8.1 ± 0.7
Malin et al. <sup>66</sup>	Impaired fasting glucose	12 (8/4)	65.1 ± 0.6 (entire group)	33.8 ± 1.0	60-65% of maxHR first four weeks, thereafter increase to 80-85%	5x/week / 50-60min	12	CT (cm <sup>2</sup> )	139.9 ± 16.8 / 86.6 ± 14.4	100 / 89.9
	Impaired glucose tolerance	9 (4/5)		32.7 ± 1.1					215.9 ± 76.6 / 140.9 ± 45.2	94.5 / 87.3
	Combined glucose intolerance	22 (7/15)		35.6 ± 1.0					187.7 ± 19.1 / 172.2 ± 19.9	96.9 / 90.1
	Normal glucose	15 (4/11)		32.3 ± 1.2					137.4 ± 23.9 /	90.1 / 84.5

	tolerant Type 2 diabetes	18 (7/11)		34.1 ± 1.3					90.5 ± 21.2 139.9 ± 23.7 / 109.0 ± 18.1	94 / 90.4
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>McKenzie et al.<sup>67</sup></b>	Males, GG genotype	29 (29/0)	58 ± 1	28.7 ± 0.7	50-70% of HRR	3-4x/week / 20-40min	24	CT (cm <sup>2</sup> )	150 (129-175) / - 20.1 (5.6)	91.1 ± 2.7 / -1.3
	Males, GT + TT genotype	21 (21/0)	61 ± 1	27.3 ± 0.8					131 (110-156) / - 29.9 (9.2)	86.7 ± 3.1 / -2.3
	Females, GG genotype	38 (0/38)	57 ± 1	27.7 ± 0.7					110 (100-121) / - 7.9 (3.1)	73.8 ± 2.0 / -0.4
	Females, GT + TT genotype	20 (0/20)	58 ± 1	27.9 ± 1.0					111 (97-127) / -0.2 (5.6)	76.0 ± 2.8 / -1.1
<b>McTiernan et al.<sup>68</sup></b>	Women, aerobic exercise	49 (0/49)	54.4 ± 7.1	28.9 ± 5.5	60-85% of maxHR	6x/week / 60min	52	CT L4-L5 (cm <sup>2</sup> )	105.9 ± 60.8 / 100.1 ± 58.8	78 ± 17.8 / -1.4 ± -1.8%
	Women, controls	51 (0/51)	53.7 ± 5.6	28.5 ± 4.5					102.6 ± 55.8 / 104.2 ± 59.6	77.9 ± 12.8 / +0.7 ± 0.9%
	Men, exercisers	51 (51/0)	56.2 ± 6.7	29.7 ± 3.7					161.8 ± 66.3 / 149.6 ± 76.6	94.8 ± 14.9 / -1.8 ± -1.9%
	Men, controls	51 (51/0)	56.6 ± 7.6	30.1 ± 4.8					176.7 ± 79.1 / 170.5 ± 73.3	97.4 ± 18.2 / +0.7 ± 0.9%
<b>Miyatake et al.<sup>69</sup></b>	Aerobic exercise	25 (25/0)	NR	28.5 ± 2.3	60% of maximum HR and walking of an extra 1000 steps/day	1x/week supervised and daily walking / duration NR	52	CT (cm <sup>2</sup> )	109.8 ± 57.2 / 82.7 ± 42.6	81.3 ± 7.9 / 78.1 ± 7.4
<b>Moghadesi et al.<sup>70</sup></b>	Aerobic exercise	8 (8/0)	NR	30.3 ± 2.1	Walking 2 miles on 40-59% VO <sub>2</sub> max	4x/week / 30min	12	MRI L4- L5 (cm <sup>3</sup> )	651.1 ± 31.8 / 602.2 ± 13.7	86.1 ± 4.6 / 84.1 ± 4.3
	Control	8 (8/0)		32.0 ± 5.3					688.4 ± 106.2 / 692.2 ± 108.8	90.4 ± 13.9 / 90.6 ± 14.1
<b>O'Leary et al.<sup>71</sup></b>	Aerobic exercise	16 (5/11)	63 (1)	33.2 (1.4)	Start at 60-65% of maxHR with a gradual increase to 80-85%	5x/week / 50-60min	12	CT L4- L5 (cm <sup>2</sup> )	175.6 (20.2) / 136.2 (16.9)	94.1 (4.3) / 90.9 (4.0)



Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Duration (weeks)	Assess- ment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Park et al.<sup>72</sup></b>	Aerobic exercise	10 (0/10)	42.2 ± 1.91	25.3 ± 1.74	60-70% of maxHR	6x/week / 60min	24	CT umbilicus	195.0 ± 12.55 / 112.4±10.50	63.7 ± 2.58 / - 4.7 kg
	Control	10 (0/10)	43.1 ± 1.67	25.5 ± 0.86					182.9 ± 16.81 / 190.4±15.74	65.2 ± 1.87 / +0.6kg
<b>Prior et al.<sup>73</sup></b>	Aerobic exercise	34 (34/0)	62 ± 1	28.9 ± 0.7	50-70% of VO2max	3x/week / 20-45min	26	CT L4-L5 (cm <sup>2</sup> )	154 (13) / 138.3	91.4 ± 2.4 / -1.6%
<b>Pritchard et al.<sup>74</sup></b>	Aerobic exercise	14 (14/0)	21.0 ± 0.8	26.2 ± 5.5	50-55% of VO2max	7x/week / 57min	13	CT L4-L5 (cm <sup>2</sup> )	80.8 ± 19.0 / 52.1 ± 22.4	82.1 ± 19.9 / 77.1 ± 19.0
<b>Redman et al.<sup>75</sup></b>	Aerobic exercise	8 (0/8)	25 ± 1	32.0 ± 1.6	55% of VO2max	5x/week / 23min week 1- 4; gradual increase to 58min week 12-16	16	MRI (kg)	1.3(0.9-1.9) / 1.2 (0.7-1.7)	84.6 ± 5.8 -1 ± 2%
<b>Reichkender et al.<sup>76</sup></b>	Moderate dose aerobic exercise	18 (18/0)	30 ± 2	28.6 ± 0.4	VO2max > 70% VO2max 50-70%	3x/week / duration depended on EE	11	MRI L4- L5 (kg)	2.2 ± 0.8 kg / 1.9 ± 0.6 kg	93.2 ± 1.9 / 89.6 ± 2.0
	High dose aerobic exercise	18 (18/0)	28 ± 1	27.6 ± 0.3		4x/week / duration depended on EE			2.0 ± 0.7 kg / 1.6 ± 0.4 kg	91.3 ± 1.7 / 88.8 ± 1.6
	Control	17 (17/0)	31 ± 1	28.0 ± 0.6					2.0 ± 0.6kg / 2.1 ± 0.6 kg	92.8 ± 2.1 / 92.9 ± 2.1
<b>Sasai et al.<sup>77</sup></b>	Moderate intra- abdominal fat	33 (33/0)	52.9 ± 10.6	29.2 ± 3.1	Anaerobic treshold	3x/week / 90min	12	CT (cm <sup>2</sup> )	149.7 ± 35.4 / 134.6 ± 43.1	80.9 ± 10.1 / -2.3 ± 2.2
	High intra- abdominal fat	24 (24/0)	53.5 ± 9.5	30.3 ± 3.1					242.4 ± 34.4 / 199.1 ± 39.7	88.8 ± 11.3 / -3.2 ± 3.0
<b>Sasai et al.<sup>78</sup></b>	Low volume exercise	19 (19/0)	49.7 ± 8.2	31.0 ± 4.1	65-80% of maxHR	3x/week / 30-60min	12	CT (cm <sup>2</sup> )	188.1 ± 53.9 / 170.3 ± 46.6	89.8 ± 13.4 / -2.7 ± 3.1
	High volume exercise	18 (18/0)	45.4 ± 8.6	29.3 ± 2.0					167.9 ± 44.3 / 137.9 ± 40.6	85.7 ± 9.6 / -3.4 ± 2.6
<b>Schwartz et al.<sup>79</sup></b>	Young men	13 (13/0)	28.2 ± 2.4	26.0 ± 3.5	50-60% of HRR with a gradual increase to 85%	5x/week / 45min	27	CT (cm <sup>2</sup> )	66.3 ± 37.1 / 54.8 ± 33.6	85.1 ± 15.0 / 84.6 ± 13.4
	Older men	15 (15/0)	67.5 ± 5.8	26.2 ± 2.7					144.5 ± 49.4 /	79.6 ± 7.9 /

									109.0 ± 44.9	77.1 ± 7.8
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Shojaee-Moradie et al.<sup>80</sup></b>	Aerobic exercise	10 (10/0)	47 ± 3	27.6 ± 0.6	60-85% of VO <sub>2</sub> max	3x/week / >20min	6	CT L4-L5 (cm <sup>2</sup> )	169.8 ± 13.1 / 139.2 ± 10.0	87.4 ± 2.8 / 87.6 ± 2.6
	Control	7 (7/0)	55 ± 4	27.6 ± 0.9					197.0 ± 25.6 / 181.4 ± 26.7	84.1 ± 2.5 / 83.3 ± 2.4
<b>Sigal et al.<sup>81</sup></b>	Aerobic training group	60 (39/21)	53.9 ± 6.6	35.6 ± 10.1	60-75% of max HR	3x/week /15- 45min	22	CT L4-L5 (cm <sup>2</sup> )	257 ± 161 / 244 ± 161	103.5 ± 31.0 /100.9 ± 30.2
	Control	63 (41/22)	54.8 ± 7.2	35.0 ± 9.5					252 ± 147 / 250 ± 147	101.3 ± 28.6 /101.0 ± 27.8
<b>Slentz et al.<sup>82</sup></b>	Low Amount, moderate intensity	40 (22/18)	54.0 ± 5.5	29.8 ± 3.2	40-55% of VO <sub>2</sub> max in order to reach walking 19.2km/week	Duration and frequency depended on set goal for distance/intens ity	34-39	CT at L4 pedicle	173 ± 72 / +1.7 ± 19.7%	88.0 ± 16.3 / -0.7%
	Low Amount, vigorous intensity	46 (23/23)	53.0 ± 7.0	29.7 ± 3.1	65-80% of VO <sub>2</sub> max in order to reach jogging 19.2km/week				154 ± 55 / +2.5 ± 21.3%	85.0 ± 13.4 / -0.8%
	High amount, vigorous intensity	42 (23/19)	51.5 ± 5.3	29.1 ± 2.4	65-80% of VO <sub>2</sub> max in order to reach levels of jogging 32.0km per week				168 ± 64 / -6.9 ± 20.8%	85.7 ± 12.2 / -2.6%
	Control	47 (23/24)	52.3 ± 7.7	29.8 ± 3.0					165 ± 68 / +8.6 ± 17.2%	86.9 ± 14.2 / -1.0%
<b>Solomon et al.<sup>83</sup></b>	Aerobic exercise and low glycemic index isocaloric diet	10 (2/7)	67 (2)	34.9 (1.1)	~85% of maximum heart rate	5x/week / 60min	12	CT (cm <sup>2</sup> )	106.9 (12.7) / 78.7 (12.1)	97.4 (3.8) / 89.6 (3.4)
	Aerobic exercise and high glycemic index isocaloric diet	12 (5/7)	64 (1)	34.1 (1.1)					117.5 (26.3) / 73.0 (18.5)	94.7 (4.4) / 85.7 (4.1)

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity	Frequency/ duration per session	Dura- tion (weeks)	Assess- ment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Yassine et al. <sup>84</sup>	Aerobic exercise	12 (NR)	64 ± 2	35.3 ± 5.8	Initially 60-65 of maxHR with a gradual increase to 80-85%	5x/week / 50-60min	12	CT (cm <sup>2</sup> )	192.3 ± 104.3 / 158.4 ± 87.0	99.7 ± 15.7 / 95.9 ± 14.6
Yoshimura et al. <sup>85</sup>	High liver fat group	13 (5/8)	NR	30.2 ± 5.7	Lactate treshold	3/week / 60min	12	CT L4-L5 (cm <sup>2</sup> )	213 ± 63 / 187 ± 66	78.3 ± 17.1 / -3.6%
	Low liver fat group	14 (6/8)		25.5 ± 3.2					139 ± 59 / 116 ± 61	66.0 ± 11.8 / -3.3%
Hypocaloric diet studies										
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)	
Alvarez et al. <sup>86</sup>	Obese, old Obese, young	6 (6/0)	60 (2.7)	28.9 (1.1)	Reduction of 500-800 kcal/day	13	CT (cm <sup>2</sup> )	184 (27) / 140 (31)	91.2 (4.1) / 83.9 (4.0)	
		6 (10/0)	32.9 (2.3)	30.4 (1.0)				135 (17) / 107 (14)	97.9 (4.3) / 90.2 (3.8)	
Banasik et al. <sup>87</sup>	VLCD	15 (2/13)	39.6 ± 13.4	36.2 ± 6.3	Restriction to 800 kcal/day	4	CT L4-L5 (cm <sup>2</sup> )	139.8 ± 82 / 120.8±85.9	104.1 ± 26.8 / 97.3 ± 26.4	
Bosy-Westphal et al. <sup>88</sup>	LCD	30 (0/30)	31.4 ± 6.0	35.5 ± 4.9	Restriction to 800-1000 kcal/day	14.2	MRI (cm <sup>3</sup> )	1757 ± 826 / 1530 ± 755	101.0 ± 18.3/ 91.2 ± 17.4	
Brochu et al. <sup>89</sup>	LCD	71 (0/71)	58.0 ± 4.7	32.2 ± 4.6	Reduction of 500-800 kcal of baseline resting metabolic rate (determined by indirect calorimetry)	26	CT L4-L5 (cm <sup>2</sup> )	186 ± 56 / -23 ± 30	83.6 ± 14.4 / - 5,1 ± 4,7	
Chan et al. <sup>90</sup>	Hypocaloric diet	20 (20/0)	46 ± 8 (entire group)	35 ± 1.0	reduction in energy intake by ~33%	16	MRI (kg)	7.1 ( 0.5) / 5.4 (0.4)	109 (2) / 96 (3)	
	Isocaloric diet	15 (15/0)		31 ± 0.7				6.9 (0.4) / 6.7 (0.4)	105 (3) / 109 (2)	
Colles et al. <sup>91</sup>	VLCD	32 (19/13)	47.5 ± 8.3	47.3 ± 5.5	Restriction to 456-680 kcal/day	12	CT and MRI L2-L3 (cm <sup>2</sup> )	346.3 ± 103.3 / 285.1 ± 89.3	139.8 ± 11.0/ 125.0 ± 11.7	

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Collins et al.</b> <sup>92</sup>	LCD	30 (3/27)	53	56.0 (1.0)	Restriction to 800kcal/day	9	CT (cm <sup>2</sup> )	388.0 (31.2) / 342.1 (23)	NR
<b>Conway et al.</b> <sup>93</sup>	VLCD and LCD in black women	8 (0/8)	34.8 ± 7.2	40.0 ± 5.0	During first twelve weeks: restriction to 800 kcal/day	24	CT L4-L5(cm <sup>2</sup> )	105 (25) / 74 (23)	NR
	VLCD and LCD in white women	10 (0/10)	38.6 ± 6.3	38.2 ± 8.1	During week 12-24: restriction to 1200-1500 kcal/day			160 (70) / 105 (63)	
<b>Cooper et al.</b> <sup>94</sup>	LCD	(2/43)	47.5 ± 6.2	44.0 ± 6.6	Restriction to 1200-2100 kcal/day	52	CT (cm <sup>2</sup> )	186.9 ± 62.9 / -28.7 ± 46.	118.6 ± 16.6 / -8.8 ± 5.9
<b>Dengo et al.</b> <sup>95</sup>	LCD	36 (15/11) (combined groups)	61.2 (0.8)	30.0 (0.6)	Restriction to 1200-1500 kcal/day	12	CT (cm <sup>2</sup> )	177 (15) / 133 (12)	84.6 (2.6) / 77.5 (2.2)
	Control		66.1 (1.9)	31.8 (1.4)				188 (18) / 186 (17)	91.0 (4.8) / 90.4 (4.9)
<b>Trussardi Fayh et al.</b> <sup>96</sup>	LCD only	18 (6/12)	30.1 ± 5.5	34.7 ± 2.4	Reduction of 500-1000 kcal/day	11.4	CT L4-L5 (cm <sup>2</sup> )	136.1 ± 64.0 / 112.5 ± 54.0	95.8 ± 13.7 / 91.5 ± 14.2
<b>Fisher et al.</b> <sup>97</sup>	LCD	29 (0/29)	NR	28 ± 1	Restriction to 800 kcal/day	8	CT L4-L5 (cm <sup>2</sup> )	93 ± 35 / 58 ± 26	78 ± 8 / 66 ± 7
<b>Fujioka et al.</b> <sup>98</sup>	LCD in visceral fat obesity	14 (0/14)	39.6 ± 9.4	34.3 ± 3.2	Gradual decrease over 8 weeks to 800 kcal/day restriction, and rise to ~1100 kcal/day before discharge	8	CT (l)	6.9 ± 3.1 / 4.3 ± 2.9	83.9 ± 12.8 / 71.9 ± 10.4
	LCD in subcutaneous fat obesity	26 (0/26)	37.1 ± 9.9	36.0 ± 5.7				3.9 ± 1.7 / 2.6 ± 1.1	87.6 ± 17.3 / 75.3 ± 15.1
<b>Gasteyger et al.</b> <sup>99</sup>	LCD in women	85 (0/85)	Median 43 (21-67)	Median 37.3 (31.4 - 48.8)	Restriction to 800-1000 kcal/day	8	MRI L4-L5 (cm <sup>2</sup> )	123 (44-288) / -23.7%	Only %loss: -6 ± 5% 0 ± 2%
	LCD in men	26 (26/0)	41 (20-61)	36.6 (33.5 - 41.9)				162 (73-265) / -38.4%	
<b>Giannopoulos et al.</b> <sup>49</sup>	LCD	11 (0/11)	57 (no SE)	35.9 (2.2)	Reduction of 600 kcal/day	14	MRI (cm <sup>3</sup> )	4785 (480) / 4425 (435)	92.4 (5.9) / 88.8 (5.7)

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Goss et al.</b> <sup>100</sup>	High glycemic load LCD	29 (14/15)	34.6 ± 8.1	30.9 ± 4.5	8 weeks eucaloric diet 8 weeks hypocaloric diet with a 1000 kcal-deficit	8	CT L4-L5 (cm <sup>2</sup> )	80.6 ± 48.3 / 82.4 ± 57.9	94.3 ± 20.4 / 89.4 ± 20.9
	Low glycemic load LCD	40 (17/23)	35.6 ± 4.3	32.4 ± 4.1				89.5 ± 46.3 / 81.5 ± 49.4	98.4 ± 17.9 / 92.9 ± 18.1
<b>Gray et al.</b> <sup>101</sup>	VLCD	10 (0/10)	37 ± 4	35.1 ± 2.1	Restriction to 650 kcal/day	10	MRI (cm <sup>2</sup> )	96 ± 36 / 70 ± 26	90.6 ± 8.1 / -10.6 ± 3.8
<b>Gu et al.</b> <sup>102</sup>	VLCD	46 (27/19)	NR	32.6 ± 0.6	Restriction to < 800 kcal/diet	8	MRI L4-L5 (cm <sup>2</sup> )	113.9 (5.8) / 79.8 (3.7)	96.1 (2.7) / 87.4 (2.5)
<b>Haufe et al.</b> <sup>103</sup>	Reduced carbohydrate LCD	52 (8/44)	Subgroups: 42 ± 9 and 45 ± 8	Subgroups: 32.0 ± 3.3 and 35.6 ± 4.7	Reduction of ~30% baseline food (to a minimum of 1200 kcal)	26	MRI (kg)	1.8 ± 1.1 / 1.4 ± 0.9	95.0 ± 15.9 / 89.5 ± 14.9
	Reduced fat LCD	50 (10/40)	44 ± 9 and 46 ± 9	31.9 ± 3.9 and 33.9 ± 3				1.9 ± 1 / 1.5 ± 0.9	93.6 ± 17.3 / 89.4 ± 17.0
<b>Ibanez et al.</b> <sup>104</sup>	LCD	12 (0/12)	51.4 ± 5.5	34.6 ± 3.4	Reduction of 500 kcal/day	16	MRI (cc)	3340 ± 977 / 2724 ± 1052	88.0 ± 15.2 / 82.3 ± 14.0
	Control	9 (0/9)	50.2 ± 6.8	35.0 ± 3.6				3175 ± 1122 / 3157 ± 1073	88.9 ± 11.4 / 88.8 ± 10.5
<b>Jang et al.</b> <sup>105</sup>	LCD	177 (NR)	40.0 (1.04)	27.1 (0.22)	Reduction of 300 kcal/day	12	CT L4 (cm <sup>2</sup> )	88.3 (2.81) / 77.8 (2.58)	71.1 (0.69) / 67.8 (0.58)
<b>Janssen et al.</b> <sup>106</sup>	Men, LCD	10 (10/0)	45.6 (2.1)	31.6 (0.9)	Reduction of 1000 kcal/day from baseline isocaloric diet	16	MRI 5 cm below L4-L5 to 15cm above L4-L5 (cm <sup>2</sup> )	188 (22) / -58 (10)	98.1 (3.5) / -12%
	Women, LCD	10 (0/10)	39.6 (2.4)	34.5 (1.4)				142 (17) / -51 (7)	92.9 (5.0) / -12%
<b>Kanai et al.</b> <sup>107</sup>	LCD	26 (0/26)	50 ± 13	33.7 ± 3.1	Restriction to 1200 kcal/day	12	CT umbilicus (cm <sup>2</sup> )	168 ± 12 / 124 ± 65	81.3 ± 12.1 / 71.9 ± 10.0
<b>Kim et al.</b> <sup>108</sup>	LCD in wild type	224 (144/110)	52.7 (1.31)	25.9 (0.29)	Reduction of 300 kcal/day	12	CT L1 (cm <sup>2</sup> )	274.1 (10.4) / 254.0 (10.6)	69.3 (1.17) / 65.9 (1.17)
	LCD in only UCP3 variant	(entire study)	52.4 (1.05)	25.8 (0.29)				296.8 (10.5) / 276.7 (9.6)	69.4 (1.06) / 66.1 (1.04)

	LCD in only $\beta 3$ -AR variant LCD in both variants		55.4 (1.52) 54.3 (1.65)	25.9 (0.64) 25.4 (0.51)				281.9 (10.5) / 273.2 (9.6) 281.0 (10.5) / 272.3 (10.5)	67.4 (1.69) / 63.9 (1.68) 70.0 (2.13) / 66.7 (2.09)
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Kim et al. <sup>109</sup>	LCD	27 (27/0)	45.8 (1.7)	30.5 (0.7)	Average restriction to 1547 kcal/day	12	CT umbilicus (cm <sup>2</sup> )	195.1 (14.2) / 129.4 (10.9)	89.4 (2.4) / 79.9 (2.7)
Kockx et al. <sup>110</sup>	LCD	50 (25/25)	38.4 $\pm$ 5.5	31.3 $\pm$ 4.5	Reduction of 1000 kcal/day	13	MRI (cm <sup>2</sup> )	98 $\pm$ 31 / 66 $\pm$ 26	85.9 $\pm$ 8.8 / 74.9 $\pm$ 8.9
Laaksone et al. <sup>111</sup>	VLCD	20 (9/11)	46.7 $\pm$ 8.7	35.8 $\pm$ 9.5	Restriction to 800 kcal/day	9	CT L4 (cm <sup>2</sup> )	216 $\pm$ 49 / 148 $\pm$ 31	101.3 $\pm$ 12.0 / 86.4 $\pm$ 9.6
Langendonk et al. <sup>112</sup>	VLCD in lower body obese VLCD in upper body obese	8 (0/8) 8 (0/8)	35.0 (1.7) 38.3 (2.9)	33.2 (1.6) 33.9 (1.1)	Restriction to 478 kcal/day	17	MRI L4-L5 (cm <sup>2</sup> )	303 (37) / 155 (25) 583 (77) / 359 (47)	93.4 (5.0) / 79.2 (4.7) 94.1 (3.0) / 79.7 (2.3)
Larson-Meyer et al. <sup>113</sup>	Control LCD	11 (5/6) 12 (6/6)	37 (7) 39 (5)	27.8 (2.0) 27.8 (1.4)	Reduction of -25% from baseline energy requirements	24	CT L4-L5 (kg)	2.9 (0.4) / 2.8 (0.4) 3.2 (0.5) / 2.3 (0.4)	81.8 (2.8) / 81.9 (2.8) 81.0 (3.3) / 72.6 (3.1)
Lee et al. <sup>114</sup>	LCD	33 (0/33)	32.4 $\pm$ 8.5	27.1 $\pm$ 2.3	Restriction to 1200 kcal/day	12	CT L4-L5 (cm <sup>2</sup> )	79.6 $\pm$ 28.3 / 76.9 $\pm$ 29.1	70.2 $\pm$ 8.2 / 68.2 $\pm$ 6.4
Leenen et al. <sup>115</sup>	LCD in women LCD in men	33 (0/33) 27 (27/0)	39 $\pm$ 5 40 $\pm$ 6	31.3 $\pm$ 2.2 30.7 $\pm$ 2.2	Reduction of 1000 kcal/day	13	MRI (cm <sup>2</sup> )	103 $\pm$ 35 / -33 $\pm$ 21 155 $\pm$ 38/ -61 $\pm$ 26	86.9 $\pm$ 7.6 / -12.4 $\pm$ 4.3 97.4 $\pm$ 8.0 / -13.5 $\pm$ 3.5
Maki et al. <sup>116</sup>	LCD with diacylglycerol supplements LCD with triacylglycerol supplements	65 (25/40) 62 (25/38)	49.9 $\pm$ 11.4 48.1 $\pm$ 11.2	34.5 $\pm$ 3.7 33.9 $\pm$ 3.7	Individual diet with reduction of 500-800 kcal/day	24	CT L4-L5 (cm <sup>2</sup> )	150.4 $\pm$ 10.7 / -38 $\pm$ 3 160.6 $\pm$ 9.9 / -17 $\pm$ 8	NR
Murakami et al. <sup>117</sup>	LCD	18 (10/8)	48.2 (1.9)	27.8 (0.5)	Restriction to 1000-1500 kcal/day (women); 1500-1700 kcal/day (men)	12	CT (cm <sup>2</sup> )	130.6 (16.1) / 97.9 (11.4)	72.5 (2.2) / 66.4 71.5 (2.0) /

									62.9 (1.8)
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Ng et al. <sup>118</sup>	LCD	20 (20/0)	NR	35.2 (1.0)	Restriction to 1467 kcal/day	14	MRI (kg)	7.1 (0.5) / 5.4 (0.4)	109.3 (2.3) / 96.0 (2.7)
Nicklas et al. <sup>119</sup>	LCD	34 (0/100)	58.4 ± 6.0	33.9 ± 4.0	Reduction of 400 kcal/day	20	CT L4-L5 (cm <sup>3</sup> )	2369 ± 870 / -612 ± 338	91.8 ± 10.4 / -11.8 ± 4.1
Okhawara et al. <sup>120</sup>	LCD	9 (9/0)	50.1 ± 12.9	27.9 ± 2.3	Restriction to 1680 kcal/day	13	CT umbilicus (cm <sup>2</sup> )	186 ± 41.9 / 97 ± 17.7	81.1 ± 5.6 / 69.4 ± 4.4
Okura et al. <sup>121</sup>	LCD in intra-abdominal fat obesity LCD in subcutaneous fat obesity	31 (0/31) 34 (0/34)	NR	29.4 ± 3.2 27.8 ± 2.0	Restriction to 1130 kcal/day	14	CT L4-L5 (cm <sup>2</sup> )	148 ± 41 / -37 ± 19 68 ± 24 / -23 ± 17	71.5 ± 8.8 / -7.0 ± 2.4 67.5 ± 5.9 / -7.9 ± 3.6
Pierce et al. <sup>122</sup>	LCD Control	26 (15/11) 14 (9/5)	49.5 (2.5) 40.8 (3.3)	29 (1) 31 (1)	Individualized diet with pre-set weight loss goal (minimum calories 1200 kcal/day)	12	CT L4-L5 (cm <sup>2</sup> )	128 (10) / 84 (7) 150 (19) / 154 (19)	85 (3) / 76(2) 94 (3) / 95(3)
Purnell et al. <sup>123</sup>	LCD	21 (21/0)	65 (60-75)	31 (27-37)	Restriction to 1200 kcal/day	13	CT umbilicus (cm <sup>2</sup> )	201 ± 51 / 153 ± 49	96 ± 11 / 86 ± 11
Purnell et al. <sup>124</sup>	LCD	13 (5/8)	NR	35 ± 4.8	Restriction to 1000 kcal/day for 3 months, thereafter gradual transition during 2 weeks to a solid diet	13	CT umbilicus (cm <sup>2</sup> )	146 ± 57 / 77 ± 47	99 (no SD) / 82
Riches et al. <sup>125</sup>	LCD Isocaloric diet	12 (12/0) 14 (14/0)	NR	34.1 (1.0) 34.6 (0.7)	Restriction to 1200 kcal/day	14	MRI L3 (cm <sup>2</sup> )	322.8 (23.4) / 222.1 (22.1) 309.6 (20.3) / 296.7 (15.1)	106.3 (4.1) / 95.9 (4.0) 108.2 (2.4) / 109.1 (2.6)
Ross et al. <sup>26</sup>	LCD	11 (11/0)	46.8 ± 7.6	31.6 ± 2.7	Reduction of 1000 kcal/day	16	MRI (l)	4.7 ± 1.6 / -1.5 ± 0.8	Only %loss -11.5%
Rossi et al. <sup>126</sup>	LCD	24 (13/11)	46.7 ± 14.3	35.4 ± 4.5	Reduction of 500 kcal below daily energy expenditure	13-26	MRI L4-L5 (cm <sup>2</sup> )	174.8 ± 94.7 / 118.9 ± 76.3	98.4 ± 15.9 / 89.7 ± 14.8
Ryan et al. <sup>127</sup>	LCD in NGT	29 (0/29)	60 (1)	32.8 (0.9)	Reduction of 500 kcal/day	26	CT L4-L5	146.9 (12.6) / 127.1 (11.1)	88.3 (2.8) / 81.9 (2.9)

	LCD in IGT	17 (0/17)	65 (2)	32.7 (1.2)			(cm <sup>2</sup> )	148.7 (11.6) / 126.5 (9.7)	84.4 (3.7) / 77.0 (3.3)
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
Ryan et al. <sup>128</sup>	LCD	23 (0/23)	56 (1)	Range: 25-48	Reduction of 250-350 kcal/day	26	CT L4-L5 (cm <sup>2</sup> )	140.4 (12.1) / 115.1 (11.5)	88.8 (3.8) / 83.6 (3.7)
Saiki et al. <sup>129</sup>	LCD	22 (16/6)	53.6 ± 8.4	30.4 ± 5.3	Restriction to 740 or 970 kcal/day	4	CT L4-L5 (cm <sup>2</sup> )	233.1 ± 66.5 / 191.0 ± 67.0	85.2 ± 17.0 79.0 ± 17.2
Shin et al. <sup>130</sup>	LCD in MAO	106 (0/106)	39.8 ± 12.2	28.0 ± 2.6	Reduction of 300 kcal/day	12	CT L4 (cm <sup>2</sup> )	95.1 ± 34.0 / 89.5 ± 33.4	71.2 ± 8.3 / -3.16±4.08%
	LCD in MHO	23 (0/23)	36.4 ± 11.2	27.2 ± 1.94				69.0 ± 18.5 / 63.6 ± 15.5	70.5 ± 5.1 / -2.83±2.74%
Snel et al. <sup>131</sup>	VLCD	14 (8/6)	53 (2)	35.2 (1.1)	Restriction to 450 kcal/day	16	MRI L5 (ml)	553 (47) / 228 (46)	107 (4) / 83 (4)
Stallone et al. <sup>132</sup>	LCD	11 (0/11)	52 (no SD)	37.0 ± 4.5	3 months restriction 400-800 kcal/day, 2 months refeeding, 1 month 1200-1500 kcal/day	26	CT L4 (cm <sup>2</sup> )	148 ± 75.4 / -52.9 ± 38.0	94.8 ± 10.8 / -18.8 ± 6.9
Svendsen et al. <sup>133</sup>	VLCD	10 (0/10)	Median (range) 34 (28-27)	NR(minimum for each subject: 28)	Restriction to 500-600 kcal/day	8	CT umbilicus (cm <sup>2</sup> )	125.9 ± 115.2 / 109.8 ± 90.3	Only %loss: -11%
Tchernof et al. <sup>134</sup>	LCD	25 (0/25)	57.2 ± 5.5	35.3 ± 4.0	Restriction to 1200 kcal/day	13.9	CT L4-L5 (cm <sup>2</sup> )	202 ± 73 / 128 ± 57	93.0 ± 10.7 / 79.5 ± 11.0
Tiikainen et al. <sup>135</sup>	Women with high liver fat	11 (0/11)	37 (1)	33 (1)	Reduction of 600-800 kcal/day	18-19	MRI (cm <sup>3</sup> )	1665 ± 141 / -383 ± 67	Only %loss: -8.4 (0.2)% -8.3 (0.2)%
	Women with low liver fat	12 (0/12)	37 (2)	32 (10)				1497 ± 167 / -441 ± 122	
Toledo et al. <sup>136</sup>	LCD	7 (3/7)	46.1 (2.0)	33.4 (1.2)	Reduction of 25% of calorie intake (both groups)	19.2	CT L4-L5 (cm <sup>2</sup> )	207.9 (24.7) / 172.1 (32)	95.0 (4.3) / 84.4 (2.7)
Van Dam et al. <sup>137</sup>	VLCD in ovalutory responders	9 (0/9)	30 (2.5)	37.5 (1.6)	Restriction to 470 kcal/day	8	MRI L4-L5 (cm <sup>2</sup> )	138 (17) / 91 (18)	NR
	VLCD in ovalutory non-responders	6 (0/6)	30 (1.8)	41.9 (3.6)				166 (29) / 114 (217)	



Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Caloric restriction	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Weight (kg) (pre/post)
<b>Van der Kooy et al.</b> <sup>138</sup>	Obese women	40 (0/40)	39 ± 6	31.3 ± 2.3	Reduction of 1000 kcal/day	13	MRI (cm <sup>2</sup> )	106 ± 50 / -37 ± 29	86.5 ± 8.7 / -12.6 ± 3.9
	Obese men	38 (38/0)	40 ± 6	30.7 ± 2.3				154 ± 40 / -61 ± 25	98.3 ± 7.2 / -13.3 ± 3.0
<b>Viljanen et al.</b> <sup>139</sup>	VLCD	16 (4/12)	45 (2.5)	33.3 (1.1)	Restriction to 550 kcal/day	6	MRI L2/L3 (kg)	1.6 (0.2) / 1.2 (0.1)	95.7 (3.3) / 84.6 (2.9)
<b>Vissers et al.</b> <sup>140</sup>	LCD	20 (5/15)	45.5 ± 13.1	32.9 ± 3.1	Reduction of -600 kcal/day (for all diet groups)	26	CT L4-L5 (cm <sup>2</sup> )	134.8 ± 57.3 / -26/3 ± 29.2	92.1 ± 11.1 / - 6.1 ± 4.6
	Control	21 (5/16)	44.8 ± 11.4	30.8 ± 3.4				111.5 ± 47.6 / -3.6 ± 20.5	88.6 ± 15.9 / + 0.9 ± 3.4
<b>Wahlroos et al.</b> <sup>141</sup>	VLCD (n=13)	13 (0/13)		45 ± 7	Restriction to 450-800 kcal/day	6	MRI L4-L5 (mm <sup>2</sup> )	22400±11300/ 18300±8700	118.8 ± 16.6/ 110.0 ± 17.5
<b>Weinsier et al.</b> <sup>142</sup>	LCD in white women	23 (0/23)	37.0 ± 5.9	29.0 ± 1.5	Restriction to 800 kcal/day	22	CT L4-L5 (cm <sup>2</sup> )	113.0 ± 39.2 / 67.0 ± 23.8	79.1 ± 5.0 / 66.0 ± 4.8
	LCD in black women	23 (0/23)	35.5 ± 5.9	28.7 ± 1.8				67.6 ± 18.0 / 41.8 ± 16.9	78.2 ± 8.9 / 65.6 ± 7.7
<b>Zamboni et al.</b> <sup>143</sup>	VLCD and LCD	16 (0/16)	38.8 ± 14.1	38.2 ± 6.9	First two weeks restriction to 307 kcal/day (VLCD) LCD for a mean duration of 14 weeks with restriction to 1003 kcal/day	16	CT L4 (cm <sup>2</sup> )	167 ± 80.3 / 93.3 ± 61.6	104.3 ± 18.1 / 88.1 ± 11.6

Table 2. Characteristics of included studies (n=8) that directly compared exercise training with hypocaloric diet

Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity (exercise studies)  Caloric restriction (hypocaloric diet studies)	Frequency/ duration per session (for exercise only)	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Results weight (kg) (pre / post)
<b>Christian sen et al.</b> <sup>144</sup>	Aerobic exercise	19 (9/10)	37.2 ± 7	33.3 ± 4	70% of HRR	60-75 min per session, 3x/week	12	MRI	3038.3±1086.1 / -18.4% ± 2.8	100.9 / -3.5kg
	LCD	19 (10/9)	35.6 ± 7	35.3 ± 4	Restriction to 600 kcal during 8 weeks		8 weeks diet, 4 weeks weight maintenance		3437.5±1516.2 / -30.2% ± 3.2	107.8 / -12.3
<b>Coker et al.</b> <sup>145</sup>	Aerobic exercise	6 (2/4)	55 (2)	32 (1)	50% of VO <sub>2</sub> peak	Depending on energy expenditure (week 1: 1000 kcal with a gradual increase to 2500 kcal/week)	12	CT (cm <sup>2</sup> )	245 (31) / 228 (24)	91 (3) / 91 (3)
	LCD	6 (3/3)	58 (2)	30 (0)	Reduction of 1000 kcal/week in week 1, and a further addition of 500 kcal each week until a reduction of 2500 kcal/week was reached				199 (12) / 170 (11)	86 (2) / 81 (2)
	Control	5 (3/2)	59 (2)	31 (1)					198 (17)/ 170 (11)	89 (4) / 91 (4)
<b>Koo et al.</b> <sup>146</sup>	LCD	19 (0/19)	57 ± 8	27.1 (no SD)	Restriction 1200 kcal/day	7 days/week, 120 minutes	12	CT L4-L5 (cm <sup>2</sup> )	157.8 / 151.7	67.4 (no SD) / 62.4
	Aerobic exercise	13 (0/13)	59 ± 4	25.5	Depending on energy expenditure				162.4 / 146.9	64.0 / 62.4
	Control	18 (0/18)	57 ± 8	28.5					172.4 / 163.4	66.0 / 65.8
<b>Nordby et al.</b> <sup>147</sup>	Aerobic training	12 (12/0)	28 (1)	28.3 (0.3)	65% HRR, alternated with HIIT (bouts at 85% HRR)	7 days/week, duration depended on energy expenditure (600 kcal/day)	12	MRI T11-L5 (L)	1.60 (0.12) / -0.53	94.5 (2.3) / 88.6 (2.3)
	LCD	12 (12/0)	32 (2)	28.0 (0.4)	Reduction of 600 kcal/day				1.83 (0.18) / -0.25	91.2 (1.8) / 85.9 (2.2)*

	Control	12 (12/0)	31 (2)	28.0 (0.4)		kcal per session)			2.12 (0.21) / -0.02	92.2 (2.7) / 92.1 (2.5)
Reference	Groups	N (M/F)	Age (years)	BMI (kg/m <sup>2</sup> )	Intensity (exercise studies)  Caloric restriction (hypocaloric diet studies)	Frequency/ duration per session (for exercise only)	Duration (weeks)	Assessment VAT	Results VAT (pre / post)	Results weight (kg) (pre / post)
Oh et al. <sup>148</sup>	Aerobic exercise  LCD	108 (108/0) 104 (104/0)	NR (adults)	29.2 (0.3) 29.4 (0.4)	60-85% of maxHR  Restriction to 1680 kcal/day	3 days/week, 40-60 minutes	12	CT (cm <sup>2</sup> )	178.1 (5.5) / 156.4 (5.0) 159.0 (6.2) / 123.3 (5.2)	85.2 (1.0) / 82.6 (1.0) 84.9 (1.3) / 77.7 (1.2)
Racette et al. <sup>149</sup>	LCD  Aerobic exercise  Control	19 (7/12)  19 (7/12)  10 (4/6)	55.6 (0.8)  58.8 (0.6)  56.0 (0.9)	27.2 (0.6)  27.2 (0.4)  27.9 (0.4)	Reduction of 16% of caloric intake 3 months, reduction of 20% 9 months Depending on energy deficit (same reduction as diet groups)	7 days /week, duration depending on energy deficit	52	MRI (cm <sup>3</sup> )	824.7 ± 143.4 / 633.5 ± 95.6  1123.5 ± 131.5 / 513.9 ± 107.6  1159.4 ± 203.4 / 1004 ± 155	78.5 (2.3) / 70.5 (2.3)  77.5 (2.4) / 71.0 (2.4)  81.9 (3.7) / 80.0 (3.7)
Ross et al. <sup>150</sup>	LCD  Aerobic exercise  Control	15 (0/15)  17 (0/17)  10 (0/10)	43.9 ± 4.9  43.2 ± 5.1  43.7 ± 6.4	31.9 ± 2.8  32.8 ± 3.9  32.4 ± 2.8	Reduction of 500 kcal/day ~80% maxHR	7 days/week, 63 minutes	14	MRI (kg)	2.4 ± 1.2 / 1.9 ± 1.0 2.3 ± 0.8 / 1.6 ± 0.7 2.3 ± 0.9 / 2.2 ± 0.9	86.6 ± 10.9/ 80.1 ± 11.2 86.8 ± 10.9/ 80.9 ± 10.8 88.1 ± 8.2 / 88.6 ± 7.4
Ross et al. <sup>151</sup>	LCD  Aerobic exercise  Control	14 (14/0)  16 (16/0)  8 (8/0)	42.6 ± 9.7  45.0 ± 7.5  46.0 ± 10.9	30.7 ± 1.9  32.3 ± 1.9  30.7 ± 1.6	Reduction of 700 kcal/day ~75% max HR	7 days per week, 60.4 minutes	14	MRI L4-L5 (kg)	3.2 ± 1.0 / -25.2 (2.0)% 3.9 ± 1.0 / -27.5 (1.9)% 4.1 ± 1.7 / -1.9 (2.7)%	96.1 ± 8.7 / -7.7 (0.2)% 101.5 ± 7.7 / -7.5 (0.3)% 96.7 ± 9.0 / -0.2 (0.4)%

Data depicted as: Mean ± standard deviation or Mean (standard error). Post value –(x) represents absolute decrease in VAT or weight (unless stated otherwise).

Abbreviations: M=male; F=female; BMI=body mass index; NR= not reported; maxHR = maximum heart rate; HRR = heart rate reserve; min = minutes; CT = Computed Tomography; MRI = magnetic resonance imaging; LCD=low calorie diet

- 1 Haslam DW, James WP. Obesity. *Lancet*. 2005; 366: 1197-209.
- 2 World Health Organisation. Global database on body mass index. <http://apps.who.int/bmi/index.jsp>. 2014.
- 3 Tchernof A, Despres JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev*. 2013; 93: 359-404.
- 4 Eckel RH, Krauss RM. American Heart Association call to action: obesity as a major risk factor for coronary heart disease. AHA Nutrition Committee. *Circulation*. 1998; 97: 2099-100.
- 5 Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature*. 2006; 444: 840-6.
- 6 World Health Organisation Western Pacific Region International Association for the Study of Obesity International Obesity Task Force. Redefining Obesity and its treatment. [http://www.who.int/nutrition/publications/obesity/09577082\\_1\\_1/en/](http://www.who.int/nutrition/publications/obesity/09577082_1_1/en/). 2000.
- 7 Franz MJ, VanWormer JJ, Crain AL, et al. Weight-loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc*. 2007; 107: 1755-67.
- 8 Miller WC, Koceja DM, Hamilton EJ. A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. *Int J Obes Relat Metab Disord*. 1997; 21: 941-7.
- 9 Ross R, Bradshaw AJ. The future of obesity reduction: beyond weight loss. *Nature reviews Endocrinology*. 2009; 5: 319-25.
- 10 Jensen MD. Role of body fat distribution and the metabolic complications of obesity. *J Clin Endocrinol Metab*. 2008; 93: S57-63.
- 11 Mathieu P, Poirier P, Pibarot P, Lemieux I, Despres JP. Visceral obesity: the link among inflammation, hypertension, and cardiovascular disease. *Hypertension*. 2009; 53: 577-84.
- 12 Despres JP, Lemieux I, Bergeron J, et al. Abdominal obesity and the metabolic syndrome: contribution to global cardiometabolic risk. *Arterioscler Thromb Vasc Biol*. 2008; 28: 1039-49.
- 13 Farrell SW, Cheng YJ, Blair SN. Prevalence of the metabolic syndrome across cardiorespiratory fitness levels in women. *Obes Res*. 2004; 12: 824-30.
- 14 Katzmarzyk PT, Church TS, Janssen I, Ross R, Blair SN. Metabolic syndrome, obesity, and mortality: impact of cardiorespiratory fitness. *Diabetes Care*. 2005; 28: 391-7.
- 15 Wildman RP, Muntner P, Reynolds K, et al. The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: prevalence and correlates of 2 phenotypes among the US population (NHANES 1999-2004). *Arch Intern Med*. 2008; 168: 1617-24.
- 16 Ismail I, Keating SE, Baker MK, Johnson NA. A systematic review and meta-analysis of the effect of aerobic vs. resistance exercise training on visceral fat. *Obesity Reviews*. 2012; 13: 68-91.
- 17 Kay SJ, Fiatarone Singh MA. The influence of physical activity on abdominal fat: A systematic review of the literature. *Obesity Reviews*. 2006; 7: 183-200.
- 18 Vissers D, Hens W, Taeymans J, Baeyens JP, Poortmans J, Van Gaal L. The Effect of Exercise on Visceral Adipose Tissue in Overweight Adults: A Systematic Review and Meta-Analysis. *PLoS ONE*. 2013; 8.
- 19 Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation*. 2014; 129: S102-38.
- 20 Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc*. 2009; 41: 459-71.
- 21 Janssen I, Fortier A, Hudson R, Ross R. Effects of an energy-restrictive diet with or without exercise on abdominal fat, intermuscular fat, and metabolic risk factors in obese women. *Diabetes Care*. 2002; 25: 431-38.

- 22 Weiss EP, Racette SB, Villareal DT, et al. Lower extremity muscle size and strength and aerobic capacity decrease with caloric restriction but not with exercise-induced weight loss. *Journal of applied physiology (Bethesda, Md : 1985)*. 2007; 102: 634-40.
- 23 Brooks GA. Exercise physiology: Human bioenergetics and its applications 1996.
- 24 Boule NG, Haddad E, Kenny GP, Wells GA, Sigal RJ. Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials. *Jama*. 2001; 286: 1218-27.
- 25 Scholten RR, Hopman MT, Lotgering FK, Spaanderman ME. Aerobic Exercise Training in Formerly Preeclamptic Women: Effects on Venous Reserve. *Hypertension*. 2015; 66: 1058-65.
- 26 Ross R, Rissanen J, Pedwell H, Clifford J, Shragge P. Influence of diet and exercise on skeletal muscle and visceral adipose tissue in men. *Journal of applied physiology (Bethesda, Md : 1985)*. 1996; 81: 2445-55.
- 27 Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annals of Internal Medicine*. 2009; 151: 264-69.
- 28 Spungen AM, Adkins RH, Stewart CA, et al. Factors influencing body composition in persons with spinal cord injury: a cross-sectional study. *Journal of applied physiology (Bethesda, Md : 1985)*. 2003; 95: 2398-407.
- 29 Shuster A, Patlas M, Pinthus JH, Mourtzakis M. The clinical importance of visceral adiposity: a critical review of methods for visceral adipose tissue analysis. *The British journal of radiology*. 2012; 85: 1-10.
- 30 Law M SD, Letts L, Pollock N, Bosch J, et al. Critical Review Form, Quantitative Studies. *McMaster University*: 1998.
- 31 Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc*. 2000; 32: S498-504.
- 32 Haskell WL, Lee IM, Pate RR, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc*. 2007; 39: 1423-34.
- 33 Convertino VA. Blood volume: its adaptation to endurance training. *Med Sci Sports Exerc*. 1991; 23: 1338-48.
- 34 Watts K, Beye P, Siafarikas A, et al. Exercise training normalizes vascular dysfunction and improves central adiposity in obese adolescents. *J Am Coll Cardiol*. 2004; 43: 1823-7.
- 35 Micklesfield LK, Goedecke JH, Punyanitya M, Wilson KE, Kelly TL. Dual-energy X-ray performs as well as clinical computed tomography for the measurement of visceral fat. *Obesity (Silver Spring)*. 2012; 20: 1109-14.
- 36 Baria F, Kamimura MA, Aoike DT, et al. Randomized controlled trial to evaluate the impact of aerobic exercise on visceral fat in overweight chronic kidney disease patients. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2014.
- 37 Boudou P, Sobngwi E, Mauvais-Jarvis F, Vexiau P, Gautier JF. Absence of exercise-induced variations in adiponectin levels despite decreased abdominal adiposity and improved insulin sensitivity in type 2 diabetic men. *European Journal of Endocrinology*. 2003; 149: 421-24.
- 38 Cho JK, Lee SH, Lee JY, Kang HS. Randomized controlled trial of training intensity in adiposity. *International journal of sports medicine*. 2011; 32: 468-75.
- 39 Cuff DJ, Meneilly GS, Martin A, Ignaszewski A, Tildesley HD, Frohlich JJ. Effective Exercise Modality to Reduce Insulin Resistance in Women With Type 2 Diabetes. *Diabetes Care*. 2003; 26: 2977-82.
- 40 Davidson LE, Hudson R, Kilpatrick K, et al. Effects of exercise modality on insulin resistance and functional limitation in older adults: a randomized controlled trial. *Arch Intern Med*. 2009; 169: 122-31.

- 41 Dekker MJ, Lee S, Hudson R, et al. An exercise intervention without weight loss decreases circulating interleukin-6 in lean and obese men with and without type 2 diabetes mellitus. *Metabolism: Clinical and Experimental*. 2007; 56: 332-38.
- 42 Despres JP, Pouliot MC, Moorjani S, et al. Loss of abdominal fat and metabolic response to exercise training in obese women. *American Journal of Physiology - Endocrinology and Metabolism*. 1991; 261: E159-E67.
- 43 DiPietro L, Seeman TE, Stachenfeld NS, Katz LD, Nadel ER. Moderate-intensity aerobic training improves glucose tolerance in aging independent of abdominal adiposity. *Journal of the American Geriatrics Society*. 1998; 46: 875-79.
- 44 Donges CE, Duffield R, Guelfi KJ, Smith GC, Adams DR, Edge JA. Comparative effects of single-mode vs. duration-matched concurrent exercise training on body composition, low-grade inflammation, and glucose regulation in sedentary, overweight, middle-aged men. *Appl Physiol Nutr Metab*. 2013; 38: 779-88.
- 45 Donges CE, Duffield R, Drinkwater EJ. Effects of resistance or aerobic exercise training on interleukin-6, C-reactive protein, and body composition. *Medicine and Science in Sports and Exercise*. 2010; 42: 304-13.
- 46 Donnelly JE, Hill JO, Jacobsen DJ, et al. Effects of a 16-month randomized controlled exercise trial on body weight and composition in young, overweight men and women: the Midwest Exercise Trial. *Arch Intern Med*. 2003; 163: 1343-50.
- 47 Friedenreich CM, Woolcott CG, McTiernan A, et al. Adiposity changes after a 1-year aerobic exercise intervention among postmenopausal women: a randomized controlled trial. *International journal of obesity (2005)*. 2011; 427-35.
- 48 Gan SK, Kriketos AD, Ellis BA, Thompson CH, Kraegen EW, Chisholm DJ. Changes in aerobic capacity and visceral fat but not myocyte lipid levels predict increased insulin action after exercise in overweight and obese men. *Diabetes Care*. 2003; 26: 1706-13.
- 49 Giannopoulou I, Ploutz-Snyder LL, Carhart R, et al. Exercise is required for visceral fat loss in postmenopausal women with type 2 diabetes. *Journal of Clinical Endocrinology and Metabolism*. 2005; 90: 1511-18.
- 50 Halverstadt A, Phares DA, Ferrell RE, Wilund KR, Goldberg AP, Hagberg JM. High-density lipoprotein-cholesterol, its subfractions, and responses to exercise training are dependent on endothelial lipase genotype. *Metabolism*. 2003; 52: 1505-11.
- 51 Haus JM, Solomon TP, Marchetti CM, et al. Decreased visfatin after exercise training correlates with improved glucose tolerance. *Med Sci Sports Exerc*. 2009; 41: 1255-60.
- 52 Heydari M, Freund J, Boutcher SH. The effect of high-intensity intermittent exercise on body composition of overweight young males. *Journal of Obesity*. 2012; 2012.
- 53 Hutchison SK, Stepto NK, Harrison CL, Moran LJ, Strauss BJ, Teede HJ. Effects of exercise on insulin resistance and body composition in overweight and obese women with and without polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism*. 2011; 96: E48-E56.
- 54 Irving B, Weltman J, Davis C, et al. Effects of exercise training intensity on abdominal fat in abdominally obese individuals with the metabolic syndrome. *Diabetes*. 2006; 55: A238-A38.
- 55 Irwin ML, Yasui Y, Ulrich CM, et al. Effect of exercise on total and intra-abdominal body fat in postmenopausal women: A randomized controlled trial. *Journal of the American Medical Association*. 2003; 289: 323-30.
- 56 Janssen I, Katzmarzyk PT, Ross R, et al. Fitness alters the associations of BMI and waist circumference with total and abdominal fat. *Obesity Research*. 2004; 12: 525-37.
- 57 Johnson NA, Sachinwalla T, Walton DW, et al. Aerobic exercise training reduces hepatic and visceral lipids in obese individuals without weight loss. *Hepatology (Baltimore, Md)*. 2009; 1105-12.
- 58 Jung JY, Han KA, Ahn HJ, et al. Effects of aerobic exercise intensity on abdominal and thigh adipose tissue and skeletal muscle attenuation in overweight women with type 2 diabetes mellitus. *Diabetes & metabolism journal*. 2012; 36: 211-21.

- 59 Karstoft K, Winding K, Knudsen SH, *et al.* The effects of free-living interval-walking training on glycemic control, body composition, and physical fitness in type 2 diabetic patients: A randomized, controlled trial. *Diabetes Care*. 2013; 36: 228-36.
- 60 Kim MK, Tomita T, Kim MJ, Sasai H, Maeda S, Tanaka K. Aerobic exercise training reduces epicardial fat in obese men. *Journal of Applied Physiology*. 2009; 106: 5-11.
- 61 Ku Y, Kwon HR, Seok HG, *et al.* Effect of resistance exercise on muscular subfascial adipose tissue and retinol-binding protein-4 concentration in individuals with diabetes. *Diabetologia*. 2009; 52 (S1): S270-S271.
- 62 Kwon HR, Min KW, Ahn HJ, *et al.* Effects of aerobic exercise on abdominal fat, thigh muscle mass and muscle strength in type 2 diabetic subject. *Korean diabetes journal*. 2010; 34: 23-31.
- 63 Lee S, Kuk JL, Davidson LE, *et al.* Exercise without weight loss is an effective strategy for obesity reduction in obese individuals with and without Type 2 diabetes. *Journal of Applied Physiology*. 2005; 99: 1220-25.
- 64 Liao D, Asberry PJ, Shofer JB, *et al.* Improvement of BMI, body composition, and body fat distribution with lifestyle modification in Japanese Americans with impaired glucose tolerance. *Diabetes Care*. 2002; 25: 1504-10.
- 65 Malin SK, Solomon TPJ, Blaszcak A, Finnegan S, Filion J, Kirwan JP. Pancreatic beta-cell function increases in a linear dose-response manner following exercise training in adults with prediabetes. *American Journal of Physiology - Endocrinology and Metabolism*. 2013; 305: E1248-E54.
- 66 Malin SK, Kirwan JP. Fasting hyperglycaemia blunts the reversal of impaired glucose tolerance after exercise training in obese older adults. *Diabetes, Obesity and Metabolism*. 2012; 14: 835-41.
- 67 McKenzie JA, Witkowski S, Ludlow AT, Roth SM, Hagberg JM. AKT1 G205T genotype influences obesity-related metabolic phenotypes and their responses to aerobic exercise training in older Caucasians. *Experimental Physiology*. 2011; 96: 338-47.
- 68 McTiernan A, Sorensen B, Irwin ML, *et al.* Exercise effect on weight and body fat in men and women. *Obesity (Silver Spring)*. 2007; 15: 1496-512.
- 69 Miyatake N, Nishikawa H, Morishita A, *et al.* Evaluation of exercise prescription for hypertensive obese men by ventilatory threshold. *Journal of the Chinese Medical Association*. 2003; 66: 572-78.
- 70 Moghadasli M, Mohebbi H, Rahmani-Nia F, Hassan-Nia S, Noroozi H. Effects of short-term lifestyle activity modification on adiponectin mRNA expression and plasma concentrations. *European journal of sport science*. 2013; 13: 378-85.
- 71 O'Leary VB, Marchetti CM, Krishnan RK, Stetzer BP, Gonzalez F, Kirwan JP. Exercise-induced reversal of insulin resistance in obese elderly is associated with reduced visceral fat. *Journal of Applied Physiology*. 2006; 100: 1584-89.
- 72 Park SK, Park JH, Kwon YC, Kim HS, Yoon MS, Park HT. The effect of combined aerobic and resistance exercise training on abdominal fat in obese middle-aged women. *Journal of Physiological Anthropology and Applied Human Science*. 2003; 22: 129-35.
- 73 Prior SJ, Joseph LJ, Brandauer J, Katznel LI, Hagberg JM, Ryan AS. Reduction in mid-thigh low-density muscle with aerobic exercise training and weight loss impacts glucose tolerance in older men. *Journal of Clinical Endocrinology and Metabolism*. 2007; 92: 880-86.
- 74 Pritchard J, Despres JP, Gagnon J, *et al.* Plasma adrenal, gonadal, and conjugated steroids following long-term exercise-induced negative energy balance in identical twins. *Metabolism: Clinical and Experimental*. 1999; 48: 1120-27.
- 75 Redman LM, Elkind-Hirsch K, Ravussin E. Aerobic exercise in women with polycystic ovary syndrome improves ovarian morphology independent of changes in body composition. *Fertility and Sterility*. 2011; 95: 2696-99.
- 76 Reichkendler MH, Auerbach P, Rosenkilde M, *et al.* Exercise training favors increased insulin-stimulated glucose uptake in skeletal muscle in contrast to adipose tissue: A randomized study using FDG PET imaging. *American Journal of Physiology - Endocrinology and Metabolism*. 2013; 305: E496-E506.

- 77 Sasai H, Katayama Y, Nakata Y, Ohkubo H, Tanaka K. Obesity phenotype and intra-abdominal fat responses to regular aerobic exercise. *Diabetes Research and Clinical Practice*. 2009; 84: 230-38.
- 78 Sasai H, Katayama Y, Nakata Y, et al. The effects of vigorous physical activity on intra-abdominal fat levels: A preliminary study of middle-aged Japanese men. *Diabetes Research and Clinical Practice*. 2010; 88: 34-41.
- 79 Schwartz RS, Shuman WP, Larson V, et al. The effect of intensive endurance exercise training on body fat distribution in young and older men. *Metabolism: Clinical and Experimental*. 1991; 40: 545-51.
- 80 Shojaee-Moradie F, Baynes KC, Pentecost C, et al. Exercise training reduces fatty acid availability and improves the insulin sensitivity of glucose metabolism. *Diabetologia*. 2007; 50: 404-13.
- 81 Sigal RJ, Kenny GP, Boule NG, et al. Effects of aerobic training, resistance training, or both on glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med*. 2007; 147: 357-69.
- 82 Slentz CA, Aiken LB, Houmard JA, et al. Inactivity, exercise, and visceral fat. STRRIDE: A randomized, controlled study of exercise intensity and amount. *Journal of Applied Physiology*. 2005; 99: 1613-18.
- 83 Solomon TP, Haus JM, Kelly KR, et al. A low-glycemic index diet combined with exercise reduces insulin resistance, postprandial hyperinsulinemia, and glucose-dependent insulinotropic polypeptide responses in obese, prediabetic humans. *Am J Clin Nutr*. 2010; 92: 1359-68.
- 84 Yassine HN, Marchetti CM, Krishnan RK, Vrobel TR, Gonzalez F, Kirwan JP. Effects of exercise and caloric restriction on insulin resistance and cardiometabolic risk factors in older obese adults--a randomized clinical trial. *The journals of gerontology*. 2009; Series A, Biological sciences and medical sciences. 64: 90-95.
- 85 Yoshimura E, Kumahara H, Tobina T, et al. A 12-week aerobic exercise program without energy restriction improves intrahepatic fat, liver function and atherosclerosis-related factors. *Obesity Research and Clinical Practice*. 2011; 5: e249-e57.
- 86 Alvarez GE, Davy BM, Ballard TP, Beske SD, Davy KP. Weight loss increases cardiovagal baroreflex function in obese young and older men. *American Journal of Physiology - Endocrinology and Metabolism*. 2005; 289: E665-E69.
- 87 Banasik JL, Walker MK, Randall JM, Netjes RB, Foutz MS. Low-calorie diet induced weight loss may alter regulatory hormones and contribute to rebound visceral adiposity in obese persons with a family history of type-2 diabetes. *Journal of the American Association of Nurse Practitioners*. 2013; 25: 440-8.
- 88 Bosy-Westphal A, Kossel E, Goele K, et al. Association of pericardial fat with liver fat and insulin sensitivity after diet-induced weight loss in overweight women. *Obesity*. 2010; 18: 2111-17.
- 89 Brochu M, Malita MF, Messier V, et al. Resistance training does not contribute to improving the metabolic profile after a 6-month weight loss program in overweight and obese postmenopausal women. *J Clin Endocrinol Metab*. 2009; 94: 3226-33.
- 90 Chan DC, Watts GF, Ng TWK, Yamashita S, Barrett PHR. Effect of weight loss on markers of triglyceride-rich lipoprotein metabolism in the metabolic syndrome. *European Journal of Clinical Investigation*. 2008; 38: 743-51.
- 91 Colles SL, Dixon JB, Marks P, Strauss BJ, O'Brien PE. Preoperative weight loss with a very-low-energy diet: Quantitation of changes in liver and abdominal fat by serial imaging. *American Journal of Clinical Nutrition*. 2006; 84: 304-11.
- 92 Collins J, McCloskey C, Titchner R, et al. Preoperative weight loss in high-risk superobese bariatric patients: A computed tomography-based analysis. *Surgery for Obesity and Related Diseases*. 2011; 7: 480-85.
- 93 Conway JM, Yanovski SZ, Avila NA, Hubbard VS. Visceral adipose tissue differences in black and white women. *Am J Clin Nutr*. 1995; 61: 765-71.
- 94 Cooper JN, Columbus ML, Shields KJ, et al. Effects of an intensive behavioral weight loss intervention consisting of caloric restriction with or without physical activity on common carotid



artery remodeling in severely obese adults. *Metabolism: Clinical and Experimental*. 2012; 61: 1589-97.

95 Dengo AL, Dennis EA, Orr JS, et al. Arterial destiffening with weight loss in overweight and obese middle-aged and older adults. *Hypertension*. 2010; 855-61.

96 Trussardi Fayh AP, Lopes AL, Fernandes PR, Reischak-Oliveira A, Friedman R. Impact of weight loss with or without exercise on abdominal fat and insulin resistance in obese individuals: a randomised clinical trial. *Br J Nutr*. 2013; 110: 486-92.

97 Fisher G, Hyatt TC, Hunter GR, Oster RA, Desmond RA, Gower BA. Effect of diet with and without exercise training on markers of inflammation and fat distribution in overweight women. *Obesity*. 2011; 19: 1131-36.

98 Fujioka S, Matsuzawa Y, Tokunaga K, et al. Improvement of glucose and lipid metabolism associated with selective reduction of intra-abdominal visceral fat in premenopausal women with visceral fat obesity. *International Journal of Obesity*. 1991; 15: 853-59.

99 Gasteyger C, Larsen TM, Vercruysse F, Pedersen D, Toubro S, Astrup A. Visceral fat loss induced by a low-calorie diet: A direct comparison between women and men. *Diabetes, Obesity and Metabolism*. 2009; 11: 596-602.

100 Goss AM, Goree LL, Ellis AC, et al. Effects of diet macronutrient composition on body composition and fat distribution during weight maintenance and weight loss. *Obesity*. 2013; 21: 1133-38.

101 Gray DS, Fujioka K, Colletti PM, et al. Magnetic-resonance imaging used for determining fat distribution in obesity and diabetes. *Am J Clin Nutr*. 1991; 54: 623-7.

102 Gu Y, Yu H, Li Y, et al. Beneficial effects of an 8-week, very low carbohydrate diet intervention on obese subjects. *Evidence-based Complementary and Alternative Medicine*. 2013; 2013.

103 Haufe S, Kast P, Engeli S, et al. The influence of a low fat versus Low carbohydrate hypocaloric diet on intrahepatic fat content in overweight and obese human subjects. *Obesity Reviews*. 2011; 12: 216.

104 Ibáñez J, Izquierdo M, Martínez-Labari C, et al. Resistance training improves cardiovascular risk factors in obese women despite a significative decrease in serum adiponectin levels. *Obesity (Silver Spring, Md)*. 2010; 535-41.

105 Jang Y, Kim OY, Lee JH, et al. Genetic variation at the perilipin locus is associated with changes in serum free fatty acids and abdominal fat following mild weight loss. *International Journal of Obesity*. 2006; 30: 1601-08.

106 Janssen I, Ross R. Effects of sex on the change in visceral, subcutaneous adipose tissue and skeletal muscle in response to weight loss. *International Journal of Obesity*. 1999; 23: 1035-46.

107 Kanai H, Tokunaga K, Fujioka S, Yamashita S, Kameda-Takemura K, Matsuzawa Y. Decrease in intra-abdominal visceral fat may reduce blood pressure in obese hypertensive women. *Hypertension*. 1996; 27: 125-29.

108 Kim OY, Cho EY, Park HY, Jang Y, Lee JH. Additive effect of the mutations in the beta3-adrenoceptor gene and UCP3 gene promoter on body fat distribution and glycemic control after weight reduction in overweight subjects with CAD or metabolic syndrome. *Int J Obes Relat Metab Disord*. 2004; 28: 434-41.

109 Kim MK, Tanaka K, Kim MJ, et al. Comparison of epicardial, abdominal and regional fat compartments in response to weight loss. *Nutrition, Metabolism and Cardiovascular Diseases*. 2009; 19: 760-66.

110 Kockx M, Leenen R, Seidell J, Princen HMG, Kooistra T. Relationship between visceral fat and PAI-1 in overweight men and women before and after weight loss. *Thrombosis and Haemostasis*. 1999; 82: 1490-96.

111 Laaksonen DE, Kainulainen S, Rissanen A, Niskanen L. Relationships between changes in abdominal fat distribution and insulin sensitivity during a very low calorie diet in abdominally obese men and women. *Nutrition, Metabolism and Cardiovascular Diseases*. 2003; 13: 349-56.

- 112 Langendonk JG, Kok P, Frolich M, Pijl H, Meinders AE. Decrease in visceral fat following diet-induced weight loss in upper body compared to lower body obese premenopausal women. *European Journal of Internal Medicine*. 2006; 17: 465-69.
- 113 Larson-Meyer DE, Heilbronn LK, Redman LM, et al. Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care*. 2006; 29: 1337-44.
- 114 Lee HO, Yim JE, Lee JS, Kim YS, Choue R. The association between measurement sites of visceral adipose tissue and cardiovascular risk factors after caloric restriction in obese Korean women. *Nutrition research and practice*. 2013; 7: 43-8.
- 115 Leenen R, Van Der Kooy K, Seidell JC, Deurenberg P, Koppeschaar HPF. Visceral fat accumulation in relation to sex hormones in obese men and women undergoing weight loss therapy. *Journal of Clinical Endocrinology and Metabolism*. 1994; 78: 1515-20.
- 116 Maki KC, Davidson MH, Tsushima R, et al. Consumption of diacylglycerol oil as part of a reduced-energy diet enhances loss of body weight and fat in comparison with consumption of a triacylglycerol control oil. *American Journal of Clinical Nutrition*. 2002; 76: 1230-36.
- 117 Murakami T, Horigome H, Tanaka K, Nakata Y, Katayama Y, Matsui A. Effects of diet with or without exercise on leptin and anticoagulation proteins levels in obesity. *Blood Coagulation and Fibrinolysis*. 2007; 18: 389-94.
- 118 Ng TW, Watts GF, Barrett PH, Rye KA, Chan DC. Effect of weight loss on LDL and HDL kinetics in the metabolic syndrome: associations with changes in plasma retinol-binding protein-4 and adiponectin levels. *Diabetes Care*. 2007; 30: 2945-50.
- 119 Nicklas BJ, Wang X, You T, et al. Effect of exercise intensity on abdominal fat loss during calorie restriction in overweight and obese postmenopausal women: a randomized, controlled trial. *The American journal of clinical nutrition*. 2009; 89: 1043-52.
- 120 Ohkawara K, Nakata Y, Numao S, et al. Response of coronary heart disease risk factors to changes in body fat during diet-induced weight reduction in Japanese obese men: A pilot study. *Annals of Nutrition and Metabolism*. 2010; 56: 1-8.
- 121 Okura T, Nakata Y, Lee DJ, Ohkawara K, Tanaka K. Effects of aerobic exercise and obesity phenotype on abdominal fat reduction in response to weight loss. *International Journal of Obesity*. 2005; 29: 1259-66.
- 122 Pierce GL, Beske SD, Lawson BR, et al. Weight loss alone improves conduit and resistance artery endothelial function in young and older overweight/obese adults. *Hypertension*. 2008; 52: 72-79.
- 123 Purnell JQ, Kahn SE, Albers JJ, Nevin DN, Brunzell JD, Schwartz RS. Effect of weight loss with reduction of intra-abdominal fat on lipid metabolism in older men. *J Clin Endocrinol Metab*. 2000; 85: 977-82.
- 124 Purnell JQ, Cummings D, Weigle DS. Changes in 24-h area-under-the-curve ghrelin values following diet-induced weight loss are associated with loss of fat-free mass, but not with changes in fat mass, insulin levels or insulin sensitivity. *International Journal of Obesity*. 2007; 31: 385-89.
- 125 Riches FM, Watts GF, Hua J, Stewart GR, Naoumova RP, Barrett PH. Reduction in visceral adipose tissue is associated with improvement in apolipoprotein B-100 metabolism in obese men. *J Clin Endocrinol Metab*. 1999; 84: 2854-61.
- 126 Rossi AP, Fantin F, Zamboni GA, et al. Effect of moderate weight loss on hepatic, pancreatic and visceral lipids in obese subjects. *Nutrition and Diabetes*. 2012; 2.
- 127 Ryan AS, Ortmeier HK, Sorkin JD. Exercise with calorie restriction improves insulin sensitivity and glycogen synthase activity in obese postmenopausal women with impaired glucose tolerance. *American Journal of Physiology - Endocrinology and Metabolism*. 2012; 302: E145-E52.
- 128 Ryan AS, Nicklas BJ, Berman DM. Aerobic exercise is necessary to improve glucose utilization with moderate weight loss in women. *Obesity*. 2006; 14: 1064-72.
- 129 Saiki A, Nagayama D, Ohhira M, et al. Effect of weight loss using formula diet on renal function in obese patients with diabetic nephropathy. *Int J Obes (Lond)*. 2005; 29: 1115-20.

- 130 Shin MJ, Hyun YJ, Kim OY, Kim JY, Jang Y, Lee JH. Weight loss effect on inflammation and LDL oxidation in metabolically healthy but obese (MHO) individuals: low inflammation and LDL oxidation in MHO women. *Int J Obes (Lond)*. 2006; 30: 1529-34.
- 131 Snel M, Jonker JT, Hammer S, et al. Long-term beneficial effect of a 16-week very low calorie diet on pericardial fat in obese type 2 diabetes mellitus patients. *Obesity*. 2012; 20: 1572-76.
- 132 Stallone DD, Stunkard AJ, Wadden TA, Foster GD, Boorstein J, Arger P. Weight loss and body fat distribution: a feasibility study using computed tomography. *Int J Obes*. 1991; 15: 775-80.
- 133 Svendsen PF, Jensen FK, Holst JJ, Haugaard SB, Nilas L, Madsbad S. The effect of a very low calorie diet on insulin sensitivity, beta cell function, insulin clearance, incretin hormone secretion, androgen levels and body composition in obese young women. *Scandinavian Journal of Clinical and Laboratory Investigation*. 2012; 72: 410-19.
- 134 Tchernof A, Nolan A, Sites CK, Ades PA, Poehlman ET. Weight loss reduces C-reactive protein levels in obese postmenopausal women. *Circulation*. 2002; 105: 564-69.
- 135 Tiikkainen M, Bergholm R, Vehkavaara S, et al. Effects of identical weight loss on body composition and features of insulin resistance in obese women with high and low liver fat content. *Diabetes*. 2003; 52: 701-07.
- 136 Toledo FG, Menshikova EV, Azuma K, et al. Mitochondrial capacity in skeletal muscle is not stimulated by weight loss despite increases in insulin action and decreases in intramyocellular lipid content. *Diabetes*. 2008; 57: 987-94.
- 137 van Dam EW, Roelfsema F, Veldhuis JD, et al. Retention of estradiol negative feedback relationship to LH predicts ovulation in response to caloric restriction and weight loss in obese patients with polycystic ovary syndrome. *American journal of physiology Endocrinology and metabolism*. 2004; 286: E615-20.
- 138 van der Kooy K, Leenen R, Seidell JC, Deurenberg P, Droop A, Bakker CJ. Waist-hip ratio is a poor predictor of changes in visceral fat. *Am J Clin Nutr*. 1993; 57: 327-33.
- 139 Viljanen AP, Lautamaki R, Jarvisalo M, et al. Effects of weight loss on visceral and abdominal subcutaneous adipose tissue blood-flow and insulin-mediated glucose uptake in healthy obese subjects. *Ann Med*. 2009; 41: 152-60.
- 140 Vissers D, Verrijken A, Mertens I, et al. Effect of long-term whole body vibration training on visceral adipose tissue: A preliminary report. *Obesity Facts*. 2010; 3: 93-100.
- 141 Wahlroos S, Phillips ML, Lewis MC, et al. Rapid significant weight loss and regional lipid deposition: Implications for insulin sensitivity. *Obesity Research and Clinical Practice*. 2007; 1: 7-16.
- 142 Weinsier RL, Hunter GR, Gower BA, Schutz Y, Darnell BE, Zuckerman PA. Body fat distribution in white and black women: different patterns of intraabdominal and subcutaneous abdominal adipose tissue utilization with weight loss. *Am J Clin Nutr*. 2001; 74: 631-6.
- 143 Zamboni M, Armellini F, Turcato E, et al. Effect of weight loss on regional body fat distribution in premenopausal women. *Am J Clin Nutr*. 1993; 58: 29-34.
- 144 Christiansen T, Paulsen SK, Bruun JM, et al. Comparable reduction of the visceral adipose tissue depot after a diet-induced weight loss with or without aerobic exercise in obese subjects: a 12-week randomized intervention study. *European journal of endocrinology / European Federation of Endocrine Societies*. 2009; 160: 759-67.
- 145 Coker RH, Williams RH, Yeo SE, et al. The impact of exercise training compared to caloric restriction on hepatic and peripheral insulin resistance in obesity. *Journal of Clinical Endocrinology and Metabolism*. 2009; 94: 4258-66.
- 146 Koo BK, Min KW, Kim H-J, et al. Changes of Abdominal, Thigh Adipose Tissue and Skeletal Muscle Attenuation in Response to Energy Restriction or Exercise in Overweight Type 2 Diabetes. *Obesity*. 2008; 16: S228-S28.
- 147 Nordby P, Auerbach PL, Rosenkilde M, et al. Endurance training per se increases metabolic health in young, moderately overweight men. *Obesity*. 2012; 20: 2202-12.
- 148 Oh S, Tanaka K, Warabi E, Shoda J. Exercise reduces inflammation and oxidative stress in obesity-related liver diseases. *Med Sci Sports Exerc*. 2013; 45: 2214-22.

- 149 Racette SB, Weiss EP, Villareal DT, *et al.* One year of caloric restriction in humans: feasibility and effects on body composition and abdominal adipose tissue. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2006; 61: 943-50.
- 150 Ross R, Janssen I, Dawson J, *et al.* Exercise-induced reduction in obesity and insulin resistance in women: a randomized controlled trial. *Obesity Research*. 2004; 12: 789-98.
- 151 Ross R, Dagnone D, Jones PJ, *et al.* Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med*. 2000; 133: 92-103.