

1 **Short title:** Variability of Rubisco kinetics in crops

2

3 **Corresponding author details:** Jeroni Galmés, Research Group on Plant Biology
4 under Mediterranean Conditions. Universitat de les Illes Balears, Ctra. Valldemossa
5 km. 7.5; 07122 Palma, Balearic Islands, Spain; Tel: +34971259720; Fax:
6 +34971173168; e-mail: jeroni.galmes@uib.cat

7

8 **Article title:** Rubisco catalytic properties and temperature response in crops

9

10 **All author names and affiliations:** Carmen Hermida-Carrera¹, Maxim V. Kapralov²,
11 Jeroni Galmés^{1*}

12 ¹Research Group on Plant Biology under Mediterranean Conditions. Universitat de les
13 Illes Balears, Balearic Islands, Spain.

14 ²School of Natural Sciences and Psychology, Liverpool John Moores University,
15 Byrom Street, Liverpool, L3 3AF, United Kingdom

16 *Corresponding author; Research Group on Plant Biology under Mediterranean
17 Conditions. Universitat de les Illes Balears, Ctra. Valldemossa km. 7.5; 07122 Palma,
18 Balearic Islands, Spain; Tel: +34971259720; Fax: +34971173168; e-mail:
19 jeroni.galmes@uib.cat

20

21 **One sentence summary:**

22 Variability in Rubisco kinetic parameters and their temperature dependency determine
23 differences in the photosynthetic efficiency in the most important crops worldwide.

24

List of author contributions: J.G. conceived the project; C.H-C. and J.G. designed the experiment; C.H-C. performed the experiments; C.H-C., M.V.K. and JG analyzed the data and wrote the article.

Funding information: This study was financed by the Spanish Ministry of Science and Innovation (MICINN) – projects AGL2009-07999 and AGL2013-42364R awarded to JG. CH-C was supported by a FPI Fellowship of the Spanish Ministry of Education – BES-2010-030796.

Corresponding author, e-mail: jeroni.galmes@uib.cat

ABSTRACT

Rubisco catalytic traits and their thermal dependence are two major factors limiting the CO₂ assimilation potential of plants. In this study, we present the profile of Rubisco kinetics for twenty crop species at three different temperatures. The results largely confirmed the existence of significant variation in the Rubisco kinetics among species. Although some of the species tended to present Rubisco with higher thermal sensitivity (e.g., *Oryza sativa*) than others (e.g., *Lactuca sativa*), interspecific differences depended on the kinetic parameter. Comparing the temperature response of the different kinetic parameters, the Rubisco Michaelis-Menten constants for CO₂ (K_c and K_c^{air}) presented higher energy of activation (ΔH_a) than the maximum carboxylation rate (k_{cat}^c) and the CO₂ compensation point in the absence of mitochondrial respiration (Γ^*). The analysis of the Rubisco large subunit sequence revealed the existence of some sites under adaptive evolution in branches with

specific kinetic traits. Because Rubisco kinetics and their temperature dependency were species-specific, they largely affected the assimilation potential of Rubisco from the different crops, especially under those conditions (i.e., low CO₂ availability at the site of carboxylation and high temperature) inducing Rubisco-limited photosynthesis. As an example, at 25 °C, Rubisco from *Hordeum vulgare* and *Glycine max* presented, respectively, the highest and lowest potential for CO₂ assimilation at both high and low chloroplastic CO₂ concentrations. In our opinion, this information is relevant to improve photosynthesis models and should be considered in future attempts to design more efficient Rubiscos.

INTRODUCTION

The reported stagnation in the annual gains of cereal yields in the last decade clearly indicates that the expected demand for increased yield - at least 50% by 2050 (FAO forecasts) - will not be met by conventional breeding (Zhu et al., 2010). Future improvements will come from novel bioengineering approaches specifically focussed on processes limiting crop productivity that have not been addressed so far (Parry et al., 2012; Ort et al., 2015). A number of specific modifications to the primary processes of photosynthesis that could increase canopy carbon assimilation and production through step changes include the modification of the catalytic properties of Rubisco (Murchie et al., 2009; Whitney et al., 2011; Parry et al., 2013; Ort et al., 2015). First, biochemical models indicate that CO₂ fixation rates are limited by Rubisco activity under physiologically relevant conditions (Farquhar et al., 1980; von Caemmerer, 2000; Rogers, 2014). Second, Rubisco's catalytic mechanism exhibits important inefficiencies which compromise photosynthetic productivity: it is a slow

catalyst – forcing plants to accumulate large amounts of the protein – and unable to distinguish between CO₂ and O₂ – starting a wasteful side reaction with oxygen that leads to the release of previously fixed CO₂, NH₂ and energy (Roy and Andrews, 2000). These inefficiencies not only limit the rate of CO₂ fixation, but also the capacity of crops for an optimal use of resources, principally water and nitrogen (Flexas et al., 2010; Parry et al., 2012).

Rubisco kinetic parameters has been described *in vitro* at 25 °C for about 250 species of higher plants, of which only c.a. 8% are crop species (e.g., Yeoh et al., 1981; Sage, 2002; Bird et al., 1982; Ishikawa et al., 2009; Prins et al., 2016). This amount of data revealed the existence of significant variability in the main Rubisco kinetic parameters both among C₃ (Yeoh et al., 1980, 1981; Bird et al., 1982; Jordan and Ogren, 1983; Parry et al., 1987; Castrillo, 1995; Delgado et al., 1995; Kent and Tomany, 1995; Balaguer et al., 1996; Bota et al., 2002; Galmés et al., 2005, 2014a, 2014c; Ghannoum et al., 2005; Ishikawa et al., 2009) and between C₃ and C₄ species (Kane et al., 1994; Sage, 2002; Kubien et al., 2008; Perdomo et al., 2014). The existence of Rubiscos with different catalytic performance implies that the success – in terms of photosynthetic improvement – of Rubisco engineering approaches in crops will depend on the specific performance of the native enzyme from each crop species. Nevertheless, our knowledge on the actual variability in Rubisco kinetics is still narrow, not only because of the limited number of species that have been examined so far, but mainly because complete Rubisco kinetic characterization (including the main parameters) has been performed in very few species.

Recent modelling confirmed that Rubisco is not perfectly optimized to deliver maximum rates of photosynthesis, and indicated that Rubisco optimization depends on the environmental conditions under which the enzyme operates (Galmés et al.,

2014b). In particular, Rubisco catalytic parameters are highly sensitive to changes in temperature. For instance, the maximum carboxylase turnover rate ($k_{\text{cat}}^{\text{c}}$) increases exponentially with temperature (Sage, 2002; Galmés et al., 2015). However, at temperatures higher than the photosynthetic thermal optimum, the increases in $k_{\text{cat}}^{\text{c}}$ are not translated into increased CO₂ assimilation because of the decreased affinity of Rubisco for CO₂, i.e., higher Michaelis-Menten constant for CO₂ (K_{c}) and lower specificity factor ($S_{\text{c/o}}$), and the decreased CO₂/O₂ concentration ratio in solution (Hall and Keys, 1983; Jordan and Ogren, 1984). These changes favour the RuBP oxygenation by Rubisco relative to carboxylation, increasing the flux through photorespiration and, ultimately, reducing the potential growth at high temperatures (Jordan and Ogren, 1984).

Beyond the discernment of the existing variability in Rubisco kinetics at a standard temperature, the knowledge on the temperature dependence of Rubisco kinetics, and the existence of variability in the thermal sensitivity among higher plants is of key importance for modelling purposes. The number and diversity of plant species for which Rubisco kinetic parameters have been tested *in vitro* at a range of physiologically relevant temperatures are still very scarce (e.g. Laing et al., 1974; Badger and Collatz, 1977; Badger, 1980; Monson et al., 1982; Hall and Keys, 1983; Jordan and Ogren, 1984; Lehnher et al., 1985; Uemura et al., 1997; Zhu et al., 1998; Sage et al., 2002; Galmés et al., 2005; Haslam et al., 2005; Yamori et al., 2006; Perdomo et al., 2015; Prins et al., 2016), and mostly restricted to a few kinetic parameters – actually, there is no study examining the temperature dependencies of the main kinetic constants on the same species. The limited number of data reported so far suggests the existence of interspecific differences in the temperature dependence of some Rubisco kinetic parameters, like $k_{\text{cat}}^{\text{c}}$ (Chabot et al., 1972; Weber

et al., 1977; Sage, 2002) or $S_{c/o}$ (Zhu et al., 1998; Galmés et al., 2005). Actually, differences in the energy of activation of k_{cat}^c and $S_{c/o}$ seems to be ascribed to the thermal conditions typically encountered by the species in their native habitat (Galmés et al., 2005), as well as to the photosynthetic mechanism (Perdomo et al., 2015).

The variability in the response of Rubisco kinetics to changes in temperature, if confirmed, is of paramount importance. The mechanistic models of photosynthesis at leaf, canopy and ecosystem levels are based on the kinetic properties of Rubisco (Farquhar et al., 1980; von Caemmerer, 2000; Bernacchi et al., 2002) and the accuracy of these photosynthetic models depends on knowing the Rubisco kinetic parameters and their species-specific equations for the Rubisco-temperature dependencies (e.g. Niinemets et al., 2009; Yamori and von Caemmerer, 2009; Bermúdez et al., 2012; Díaz-Espejo, 2013; von Caemmerer, 2013; Walker et al., 2013). The need for estimations of the temperature dependencies of Rubisco kinetic parameters becomes timely as modellers try to predict the impact of increasing temperatures on global plant productivity (Sage et al., 2008; Gornall et al., 2010). Ideally, surveying variations in Rubisco kinetics and their temperature dependence should incorporate a correlative analysis with variations in the L- and/or S-subunit amino acid sequence. Such a complementary research would permit deciphering what residue substitutions determine the observed variability in Rubisco catalysis.

In the present study, we examined Rubisco catalytic properties and their temperature dependence in twenty crop species, thereby constituting the largest published data set of its kind. The aims of this work were: i) to compare the Rubisco kinetic parameters among the most economically important crops, ii) to search for differences in the temperature response of the main kinetic parameters among these

species, iii) to test whether crop Rubiscos are optimally suited for the conditions encountered in plant chloroplasts, and iv) to unravel key amino acid replacements putatively responsible for differences in Rubisco kinetics in crops.

RESULTS

The variability in Rubisco kinetics at 25 °C among the most relevant crop species

When considering exclusively the 18 C₃ crop species, at 25 °C, the Rubisco Michaelis-Menten constant for CO₂ under non-oxygenic (K_c) and 21% O₂ (K_c^{air}) varied c.a. two-fold and three-fold, respectively, and the maximum rates of Rubisco carboxylation (k_{cat}^c) varied c.a. two-fold (Table 1). For K_c and K_c^{air} , *Manihot esculenta* presented the lowest values ($K_c = 6.1 \mu\text{M}$, and $K_c^{\text{air}} = 10.8 \mu\text{M}$) and *Spinacia oleracea* the highest ($K_c = 14.1 \mu\text{M}$, and $K_c^{\text{air}} = 26.9 \mu\text{M}$). Values for k_{cat}^c varied between 1.4 s^{-1} (*Manihot esculenta*) and 2.5 s^{-1} (*Ipomoea batatas*). The Rubisco CO₂/O₂ specificity ($S_{c/o}$) was the kinetic parameter with the lowest variation among the C₃ species (Fig. S1), and ranged between $92.4 \text{ mol mol}^{-1}$ (*Solanum lycopersicum*) and $100.8 \text{ mol mol}^{-1}$ (*Beta vulgaris* and *Manihot esculenta*) (Table 1). *Brassica oleracea* and *Glycine max* presented the lowest value for the Rubisco carboxylase catalytic efficiency, calculated as k_{cat}^c/K_c ($0.17 \text{ s}^{-1} \mu\text{M}^{-1}$), and *Coffea arabica* presented the lowest value for the $k_{\text{cat}}^c/K_c^{\text{air}}$ ratio ($0.08 \text{ s}^{-1} \mu\text{M}^{-1}$). With regard to the oxygenase catalytic efficiency (calculated as k_{cat}^o/K_o), *Spinacia oleracea* displayed the lowest value ($1.76 \text{ s}^{-1} \text{ nM}^{-1}$). *Hordeum vulgare* presented the highest values for the Rubisco carboxylase and oxygenase catalytic efficiencies ($k_{\text{cat}}^c/K_c = 0.28 \text{ s}^{-1} \mu\text{M}^{-1}$, $k_{\text{cat}}^c/K_c^{\text{air}} = 0.17 \text{ s}^{-1} \mu\text{M}^{-1}$ and $k_{\text{cat}}^o/K_o = 3.01 \text{ s}^{-1} \text{ nM}^{-1}$).

When data from the two C₄ species (*Saccharum × officinarum* and *Zea mays*) were included in the comparison at 25 °C, the range of variability increased for all parameters (Table 1). Rubisco from the two C₄ species presented higher k_{cat}^c but lower affinity for CO₂ (i.e., higher K_c and K_c^{air} , and lower $S_{c/o}$) than Rubisco from C₃ crops. On average, k_{cat}^c/K_c and k_{cat}^o/K_o of C₄ Rubiscos were 62 % and 70 % of those of C₃ crop Rubiscos, respectively.

The temperature response of Rubisco kinetics in crops and trade-offs between catalytic traits

Both the range of variation and the species showing the extreme values of Rubisco kinetics at 15 °C and 35 °C were similar to those described at 25 °C, with some exceptions. As at 25 °C, among the C₃ crops, Rubisco from *Manihot esculenta* presented the lowest values for K_c and K_c^{air} at 15 °C and 35 °C, while the highest values were measured on Rubisco from *Spinacia oleracea* (Table S1). The lowest and highest values for k_{cat}^c at 15 °C were those of Rubisco from *Cucurbita maxima* and *Hordeum vulgare*, respectively. The degree of dispersion of the data and the range of variation between the maximum and the minimum values for K_c , K_c^{air} and k_{cat}^c increased with the increment in the assay temperature (Table S1 and Fig. S1). Regarding $S_{c/o}$, values ranged between 116.1 mol mol⁻¹ (*Brassica oleracea*) and 132.2 mol mol⁻¹ (*Cucurbita maxima*) at 15 °C, and between 74.2 mol mol⁻¹ (*Oryza sativa*) and 85.0 mol mol⁻¹ (*Manihot esculenta*) at 35 °C (Table S1). As for $S_{c/o}$, the range of variation for k_{cat}^c/K_c and k_{cat}^o/K_o was also narrowed with the increase in the assay temperature (Table S1 and Fig. S1).

Integrating all data across three assay temperatures, k_{cat}^c correlated positively with K_c for both C₃ ($r^2 = 0.82$, $p < 0.001$) and C₄ species ($r^2 = 0.94$, $p < 0.001$), with

Rubisco from C₄ species showing higher K_c for a given k_{cat}^c than that from C₃ species (Fig. 1A). The low interspecific variability in $S_{c/o}$ within each assay temperature determined a non-linear relationship between k_{cat}^c and $S_{c/o}$ when considering data from all temperatures together (Fig. 1B). At each temperature individually, Pearson's correlations between k_{cat}^c and K_c and $S_{c/o}$ were highly significant (Table 2) when considering both C₃ and C₄ together. The results from the Phylogenetically Independent Contrasts (PICs) analyses were in general more conservative compared to Pearson's correlations (Table 2), and some significant correlations were lost with PICs (e.g., $S_{c/o}$ vs. K_c or K_c^{air} at 25 °C). Notably, when excluding the two C₄ species, PCCs decreased in almost all correlations (Table 2). Hence, the PCC between k_{cat}^c and K_c was no longer significant at 15 °C, and the PCC between k_{cat}^c and $S_{c/o}$ was significant only at 15 °C. Furthermore, when considering only C₃ species, the unique significant PICs between k_{cat}^c and K_c and $S_{c/o}$ were those found between k_{cat}^c and $S_{c/o}$ at 15 °C and 25 °C.

The energy of activation (ΔH_a) for K_c varied between 38.2 kJ mol⁻¹ (*Solanum tuberosum*) and 83.1 kJ mol⁻¹ (*Oryza sativa*; Table 3). *Ipomoea batatas* (40.7 kJ mol⁻¹) and *Manihot esculenta* (75.4 kJ mol⁻¹) were the species showing the lowest and highest values for ΔH_a of K_c^{air} . As for k_{cat}^c , ΔH_a varied between 27.9 kJ mol⁻¹ (*Hordeum vulgare*) and 60.5 kJ mol⁻¹ (*Medicago sativa*). Although the range of variation across C₃ species was similar for the energies of activation of both K_c and k_{cat}^c (2.2-fold), non-significant correlation was observed between ΔH_a for K_c and ΔH_a for k_{cat}^c in both conventional and phylogenetically independent analyses ($r^2 = 0.11$ and 0.15, respectively; $P > 0.05$). The lowest and highest values for ΔH_a of the CO₂ compensation point in the absence of mitochondrial respiration (Γ^* , calculated from $S_{c/o}$) were measured in *Beta vulgaris* (19.8 kJ mol⁻¹) and *Glycine max* (26.5 kJ mol⁻¹),

respectively. On average, Rubisco from C₃ crops presented significantly higher ΔH_a for K_c (60.9 ± 1.5 kJ mol⁻¹) and k_{cat}^c (43.7 ± 1.5 kJ mol⁻¹) than Rubisco from C₄ species ($K_c = 52.4 \pm 5.0$ kJ mol⁻¹, $k_{cat}^c = 30.6 \pm 1.6$ kJ mol⁻¹). By contrast, non-significant differences were observed in the average ΔH_a for Γ^* between C₃ (22.9 ± 0.4 kJ mol⁻¹) and C₄ species (25.0 ± 0.7 kJ mol⁻¹).

The CO₂ assimilation potential of Rubisco kinetics in crops

The CO₂ assimilation potential of Rubisco ($A_{Rubisco}$) was modelled at varying temperature and CO₂ availability at the catalytic site (C_c) using the species-specific kinetic data measured at each temperature (from Tables 1 and S2). The simulated value of $C_c = 250$ μ bar is representative of that encountered in the chloroplast stroma of C₃ species under well-watered conditions (e.g., Bermúdez et al., 2012; Scafaro et al., 2012; Galmés et al., 2013). Under mild to moderate water stress, when no metabolic impairment is present, the decrease in the stomatal and leaf mesophyll conductances to CO₂ provokes a decrease in the concentration of CO₂ in the chloroplast (Flexas et al., 2006). We selected a value of 150 μ bar to simulate the chloroplastic CO₂ concentration in water stressed plants.

Differences in $A_{Rubisco}$ across species were largely dependent on the temperature and the availability of CO₂ for carboxylation (Fig. 2). This fact was due to the different prevalence of RuBP-saturated (A_c) and RuBP-limited (A_j) rates governing $A_{Rubisco}$ under the contrasting temperature and C_c , assuming an invariable concentration of active Rubisco sites of 25 μ mol m⁻² for all species. At 15 °C, A_c limited $A_{Rubisco}$ at C_c of 150 μ bar in nine species (indicated by asterisks in Fig. 2). At 15 °C and C_c of 250 μ bar, only six species were A_c limited (*Capsicum annuum*,

248 *Cucurbita maxima*, *Medicago sativa*, *Oryza sativa*, *Solanum tuberosum* and *Spinacia*
249 *oleracea*). At 25 and 35 °C, A_{Rubisco} was A_c -limited in all C_3 species irrespective of C_c .

250 At 25 °C, the *best* Rubisco was that from *Hordeum vulgare* at both C_c , while
251 Rubisco from *Glycine max* yielded the lowest A_{Rubisco} (Fig. 2). Rubisco from *Beta*
252 *vulgaris* presented the best performance at 35 °C irrespective of the CO_2 availability,
253 while *Capsicum annuum* and *Saccharum × officinarum* Rubisco gave the lowest
254 A_{Rubisco} at C_c of 250 and 150 μbar , respectively. At 15 °C and C_c of 250 μbar , the
255 highest potential for CO_2 assimilation was found in Rubisco from *Glycine max*,
256 *Manihot esculenta* and *Triticum aestivum*, while Rubisco from *Manihot esculenta*
257 gave the highest A_{Rubisco} at 15 °C and C_c of 150 μbar . Rubisco from *Cucurbita maxima*
258 displayed the lowest A_{Rubisco} at 15 °C, regardless of the CO_2 availability. It is
259 interesting to note that Rubisco from the two C_4 species, in particular from
260 *Saccharum × officinarum*, performed better than the average C_3 Rubiscos when
261 A_{Rubisco} was simulated according to the photosynthesis model for C_3 leaves (Farquhar
262 et al., 1980), at 15 °C and 25 °C under C_c of 250 μbar (Fig. 2A). At lower C_c (150
263 μbar), the C_4 Rubiscos yielded higher A_{Rubisco} values than the average C_3 Rubiscos at
264 15 °C, and lower values at 35 °C, being similar at 25 °C (Fig. 2B).

265 To test the performance of the different Rubiscos in the context of C_4
266 photosynthesis, A_c was also modelled assuming C_c of 5000 μbar and E of 15 $\mu\text{mol m}^{-2}$
267 s^{-1} . Under these conditions, the advantage of C_4 -type Rubisco kinetics of *Saccharum ×*
268 *officinarum* and *Zea mays* - characterised by higher k_{cat}^c and K_c^{air} - became evident as
269 providing higher A_c values at the three temperatures (data not shown). On average, at
270 saturating CO_2 and lower concentration of Rubisco catalytic sites, C_4 Rubiscos
271 yielded A_c of 35, 49 and 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at 15, 25 and 35 °C, respectively, compared
272 to C_3 -Rubiscos average (10, 23 and 42 $\mu\text{mol m}^{-2} \text{s}^{-1}$, respectively).

Positively selected L-subunit residues: relationship with Rubisco kinetics

The phylogeny obtained with *rbcL*, *matK* and *ndhF* genes matched currently accepted angiosperm classification (Fig. S2) (Bremer et al., 2009).

When considering all species together, 10 L-subunit residues were under positive selection: 94, 262, 281, 309, 439, 446, 449, 470, 477 and amino acid insert between residues 468 and 469. Moreover, positive selection was identified in specific L-subunit residues along branches leading to species with high and low K_c , high k_{cat}^c and low $S_{c/o}$ at 25 °C and low ΔH_a for K_c (Table 4). The residues under positive selection were located at different positions within the Rubisco tertiary structure and included functionally diverse sites participating in L-subunit intradimer and dimer-dimer interactions, interactions with small subunits (S-subunit) and with Rubisco activase (Table 4). No residue under positive selection was associated with ΔH_a for K_c^{air} , ΔH_a for k_{cat}^c or ΔH_a for $S_{c/o}$.

DISCUSSION

Main crops possess Rubiscos with different performance at 25 °C

The kinetic data reported in the present study are consistent with the range previously reported for higher plants at 25 °C (e.g., Yeoh et al., 1980, 1981; Bird et al., 1982; Jordan and Ogren, 1983; Kent and Tomany, 1995; Galmés et al., 2005, 2014a, 2014c; Ishikawa et al., 2009; Prins et al., 2016) (Table 1), and showing the existence of significant variation among species in the carboxylase catalytic efficiency under non-oxygenic (k_{cat}^c/K_c) and atmospheric conditions (k_{cat}^c/K_c^{air}). Recent reports related k_{cat}^c/K_c variation with the growth capacity in a group of closely related species with

similar ecology (Galmés et al., 2014a), suggesting that improving this ratio would be an effective way to engineer a better Rubisco. Nevertheless, such an improvement becomes constrained by the trade-offs between $k_{\text{cat}}^{\text{c}}$, K_{c} and $S_{\text{c/o}}$ (Tcherkez et al., 2006; Savir et al., 2010; Galmés et al., 2014a, 2014c). Here, we demonstrate that these trade-offs, in particular $k_{\text{cat}}^{\text{c}}$ vs. K_{c} , are hold when considering C_3 and C_4 species together, even after accounting for the phylogenetic signal in the data, and that they generally strengthen at increasing assay temperatures (Table 2). However, most of these trade-offs were lost when considering exclusively the C_3 species (Table 2), indicative that the broad-scale patterns of covariation between the Rubisco kinetic parameters may not hold at smaller scales, as previously observed in other angiosperm species (Galmés et al., 2014c).

The maximum carboxylase turnover rate of Rubisco ($k_{\text{cat}}^{\text{c}}$) from *Zea mays* and *Saccharum × officinarum* was 2-fold higher than that of the C_3 species, albeit at the expenses of 3 times less affinity for CO_2 (Table 1). This finding agrees with previously described trends between C_3 and C_4 species (Kubien et al., 2008; Ghannoum et al., 2005; Ishikawa et al., 2009), and with the fact that C_4 species present lower $k_{\text{cat}}^{\text{c}}/K_{\text{c}}$ (Kubien et al., 2008; Perdomo et al., 2015).

Unlike other reports (Sage 2002; Ishikawa et al., 2009), the observed variation in the kinetic parameters at 25 °C among C_3 species was apparently not related to the thermal climate of their respective domestication regions (data not shown). It should be noted that the origin, and hence the climatic conditions, of the selected varieties could be different to the species centre of domestication, and that the different crop varieties may have accumulated adaptive changes to local conditions by means of artificial selection (Meyer et al., 2012). Intraspecific variability in Rubisco catalytic traits has been reported in *Triticum aestivum* (Galmés et al., 2014c) and *Hordeum*

vulgare (Rinehart et al., 1983), but how this variability among genotypes is related to adaptation of Rubisco to local environments remains elusive.

The Rubisco kinetic parameters of the main crops present different thermal sensitivity

The observed temperature response of the Rubisco kinetics parameters confirms well-described trends consisting in increases in k_{cat}^c and K_c and a decrease in $S_{c/o}$ with increasing assay temperature (Table 1 and Table S1) (Jordan and Ogren, 1984; Brooks and Farquhar, 1985; Uemura et al., 1997; Galmés et al., 2005; Prins et al., 2016).

The temperature dependency of full Rubisco catalytic constants was first provided for *Nicotiana tabacum*, using *in vivo*-based leaf gas exchange analysis (Bernacchi et al., 2001). After this report, all studies dealing with the temperature response of photosynthesis assumed the temperature dependency parameters of tobacco Rubisco, irrespective of the modelled species, from annual herbs to trees, and from cold to warm adapted species (e.g., Pons et al., 2009; Keenan et al., 2010; Yamori et al., 2010; Galmés et al., 2011; Bermúdez et al., 2012; Scafaro et al., 2012). Importantly, the present dataset constitutes the most unequivocal confirmation that different temperature sensitivities of Rubisco kinetic parameters exist among different species, and that extrapolating the temperature response of a unique model species to other plants induces errors when modelling the temperature response of photosynthesis. In this sense, the *in vitro* results of the present study support *in vivo* data showing different temperature dependency of Rubisco catalytic constants in *Arabidopsis thaliana* and *Nicotiana tabacum* (Walker et al., 2013).

In general, the Rubisco constant affinities for CO₂ (K_c and K_c^{air}) were more sensitive to changes in assay temperature (i.e., presented higher energies of activation, ΔH_a) than k_{cat}^c and I^* (Table 3), in agreement with a recent study (Perdomo et al., 2015). This fact is explained by the increase in the oxygenase catalytic efficiency (k_{cat}^o/K_o) at increasing temperature. However, it should be remarked that k_{cat}^o/K_o ratio was calculated from the measured parameters K_c , k_{cat}^c and $S_{c/o}$, and that direct measurements of the oxygenase activity of Rubisco, e.g., by mass spectrometry (Cousins et al., 2010), should be undertaken to confirm this trend.

As at 25 °C, the differences in the temperature dependencies of Rubisco kinetic parameters among C₃ species were not related to the thermal environment of the species' domestication regions (data not shown). This finding contrasts with previous evidences suggesting that the temperature sensitivity of Rubisco kinetic properties have evolved to improve the enzyme's performance according to the prevailing thermal environment to which species are adapted (Sage, 2002; Galmés et al., 2005, 2015).

Although only two C₄ species were included in the present study, they presented lower ΔH_a for K_c and for k_{cat}^c than most of the C₃ species, in close agreement with trends recently observed by Perdomo et al. (2015) in *Flaveria* species (Table 3). A larger number of C₄ species need to be surveyed to verify the existence of differences in the temperature dependence of Rubisco kinetics between C₃ and C₄ species.

How do the species-specific properties of Rubisco kinetics and their temperature sensitivity impact the potential capacity of Rubisco to assimilate CO₂?

Modelling the effect of the species-specific Rubisco kinetics and temperature dependencies of Rubisco kinetics resulted in significant differences in the Rubisco CO₂ assimilation potential (A_{Rubisco}) among the studied C₃ crops (Fig. 2). This modelling exercise highlighted which species would mostly benefit from the genetic replacement of their native version of Rubisco by other foreign versions with improved performance. Notably, the modelling results clearly indicate that the performance of specific Rubiscos cannot be evaluated without considering the environmental conditions during catalysis, specifically the temperature and the CO₂ availability at the site of carboxylation (C_c). This fact results from the different temperature dependence of Rubisco kinetics among crops, and from the different impact that Rubisco kinetics have on the RuBP-saturated (A_c) and RuBP-limited (A_j) rates governing A_{Rubisco} . Hence, at 15 °C and C_c of 250 µbar, A_{Rubisco} was limited by A_j in most of the C₃ species (twelve out of eighteen), while it was limited by A_c in all C₃ species at 25 and 35 °C irrespective of the C_c value.

Detailed examination of modelled A_{Rubisco} suggests that future efforts to enhance Rubisco efficiency should be directed on the following C₃ species displaying the poorest performance: *Cucurbita maxima* and *Medicago sativa* at 15 °C and both C_c , *Glycine max*, *Capsicum annuum* and *Coffea arabica* at 25 °C and 250 µbar; *Glycine max*, *Spinacia oleracea*, *Capsicum annuum* and *Coffea arabica* at 25 °C and 150 µbar; *Capsicum annuum*, *Solanum lycopersicum* and *Lactuca sativa* at 35 °C and 250 µbar; and *Capsicum annuum* and *Solanum lycopersicum* at 35 °C and 150 µbar.

In order to focus on the Rubisco catalytic traits, the modelling assumed invariable values for the concentration of active Rubisco sites ($E = 25 \mu\text{mol m}^{-2} \text{s}^{-1}$) and specific values for the rate of photosynthetic electron transport (J) and C_c . However, species adapt and plants acclimate to the prevailing thermal environment

through changes in the concentration and/or activation of Rubisco and the rate of photosynthetic electron transport (Yamasaki et al., 2002; Yamori et al., 2011). Similarly, stomatal (g_s) and leaf mesophyll (g_m) conductances to CO_2 also vary in response to temperature (von Caemmerer and Evans, 2015). Considering the growth temperature effects on these parameters would have altered the equilibrium between A_c and A_j , and indirectly, the consequences of different Rubisco kinetic traits on the CO_2 assimilation potential. In the next future, we aim to increase the accuracy of the present simulation by examining and including the species-specific values for g_s , g_m , E and J at varying environmental conditions.

The analysis of positive selection in branches leading to specific Rubisco traits may reveal lineage specific amino acid substitutions

We found ten Rubisco L-subunit residues under positive selection (94, 262, 281, 309, 439, 446, 449, 469, 470, and 477; Table 4). With the exceptions of residues 469 and 477, these residues have been reported previously in other groups of plants, implying a relatively limited number of residues responsible for the Rubisco ‘fine-tuning’ (Kapralov and Filatov, 2007; Christin et al., 2008; Iida et al., 2009; Kapralov et al., 2011; Kapralov et al., 2012; Galmés et al., 2014a, 2014c). However, despite widespread parallel evolution of amino acid replacements in the Rubisco sequence, solutions found in particular groups of plants may be quite different. For instance, there are only two common residues under positive selection out of ten between this study and methodologically similar work with different sampling design published earlier (Galmés et al., 2014c). This fact raises questions of epistatic interactions and residue co-evolution within Rubisco (Wang et al., 2011) as well as residue co-evolution and complementarity between Rubisco and its chaperones (Whitney et al.,

2015), which both may prevent evolution of identical amino acid replacements because of different genetic backgrounds.

We have not examined the species differences in the sequence of the Rubisco small subunit (S-subunit). Some of the species included in the present survey, like *Triticum aestivum*, possess a large number of S-subunit genes (*rbcS*) encoding different S-subunits (Galili et al., 1998). Previous reports have showed that species with identical L-subunits might have different Rubisco kinetics (Rosnow et al., 2015) as well as directly demonstrated that differences in the S-subunits might affect Rubisco catalytic traits (Ishikawa et al., 2011; Morita et al., 2014). Therefore, we cannot discard that the observed differences in Rubisco kinetics, and their temperature dependence, among the studied crops are partially due to differences in the S-subunits.

Conclusions

The present study confirms the significant variation in carboxylation efficiency and parameters that contribute to it among plant species, and for the first time provides full Rubisco kinetic profiles for the twenty most important crop species. Our dataset could be used as an input for the next generation of species-specific models of leaf photosynthesis and its response to climate change, leading to more precise forecasts of changes in crop productivity and yield. These data could help to decide in which crops CO₂ assimilation potential and carboxylation efficiency of Rubisco might be improved via re-engineering of native enzymes or by replacement with foreign ones as there is no a one size fits all solution. The design of future attempts of Rubisco engineering in crops should be based on surveys of Rubisco catalytic and genetic diversity with a particular stress on the relatives of crops in question. Growing

knowledge of the Rubisco catalytic spectrum combined with the existing engineering toolkits for Rubisco (Whitney and Sharwood 2008) and its chaperones (Whitney et al., 2015) give us a hope that Rubisco efficiency and hence photosynthetic capacity of crops could be improved in a near future.

MATERIALS AND METHODS

Species selection and growth conditions

The following twenty crop species were selected for study: *Avena sativa* L. cv. Forridena, *Beta vulgaris* L. cv. Detroit, *Brassica oleracea* L. var. *italica* cv. Calabres, *Capsicum annuum* L. cv. Picante, *Coffea arabica* L. cv ‘Catuai’ Vermelho IAC 44, *Cucurbita maxima* D. cv. Totanera, *Glycine max* (L.) Merr cv. Ransom, *Hordeum vulgare* L. subsp. *vulgare* cv. Morex, *Ipomoea batatas* (L.) Lam var. *Rosa de Málaga*, *Lactuca sativa* L. cv. Cogollo de Tudela, *Manihot esculenta* C., *Medicago sativa* L. cv. Aragón, *Oryza sativa* L. cv. Bomba, *Phaseolus vulgaris* L. cv. Contender, *Saccharum* × *officinarum* (hybrid between *Saccharum officinarum* and *S. spontaneum*), *Solanum lycopersicum* L. cv. Roma VF, *Solanum tuberosum* L. cv. Erlanger, *Spinacia oleracea* L. cv. Butterfly, *Triticum aestivum* L. cv. Cajeme, *Zea mays* L. cv. Carella. These species represent the most important crops in terms of worldwide production (FAOSTAT, 2010). *Coffea arabica* was selected as being the most important commodity in the international agricultural trade (DaMatta 2004). Plants were grown from seeds under natural photoperiod in a glasshouse at the University of the Balearic Islands (Spain) during 2011 and 2012. Plants were grown in soil-based compost supplemented with slow-release fertilizer and frequently

watered to avoid water stress. The air temperature in the glasshouse during the growth period was maintained between 15°C and 30°C.

Determination of the Rubisco Michaelis-Menten constant for CO₂ (K_c) and the maximum carboxylase turnover rate (k_{cat}^c)

The Rubisco Michaelis-Menten constant for CO₂ under 0% O₂ (K_c) and 21% O₂ (K_c^{air}) were determined in crude extracts obtained as detailed in Galmés et al., (2014a). Rates of ¹⁴CO₂-fixation were measured at 15 °C, 25 °C and 35 °C using activated protein extracts in 7 mL septum capped scintillation vials containing reaction buffer (100 mM Bicine-NaOH pH 8.0, 20 mM MgCl₂, 0.4 mM RuBP and about 100 W-A units of carbonic anhydrase) previously equilibrated either with nitrogen (N₂) or a mixture of O₂ and N₂ (21:79). Nine different concentrations of H¹⁴CO₃⁻ (0.1 to 9.4 mM, each with a specific radioactivity of 3.7×10^{10} Bq mol⁻¹) were prepared in the scintillation vials as described previously (Galmés et al., 2014a). Assays at 35 °C using Rubisco from C₄ species required increasing H¹⁴CO₃⁻ up to 17.7 mM to reach saturating CO₂ concentration in the aqueous-phase. Assays were started by the addition of 10 µL of protein extract and stopped after 1 min by injection of 0.1 mL 10 M formic acid. Acid-stable ¹⁴C was determined by liquid scintillation counting (LS 6500 Multi-Purpose Scintillation Counter, Beckman Coulter, USA) following removal of acid-labile ¹⁴C by evaporation. The Michaelis-Menten constants for CO₂ under 0% O₂ (K_c) and 21% O₂ (K_c^{air}) were determined from the fitted data as described elsewhere (Bird et al., 1982). Replicate measurements ($n = 3-6$) were made using different biological replicates for each species.

To obtain k_{cat}^c , the maximum rate of carboxylation was extrapolated from the Michaelis-Menten fit and divided by the number of Rubisco active sites in solution, quantified by [^{14}C] CABP binding (Yokota and Calvin 1985).

Additional control assays undertaken as detailed in Galmés et al. (2014a) confirmed that the observed acid stable ^{14}C signal was uniquely the result of Rubisco catalytic activity.

Determination of the Rubisco specificity for CO_2/O_2 ($S_{c/o}$)

The Rubisco CO_2/O_2 specificity ($S_{c/o}$) was measured on purified extracts obtained as in Gago et al., (2013). On the day of $S_{c/o}$ measurement, highly concentrated Rubisco solutions were desalted by centrifugation through G25 Sephadex columns previously equilibrated with CO_2 -free 0.1 M Bicine (pH 8.2) containing 20 mM MgCl_2 . The desalted solutions were made 10 mM with $\text{NaH}^{14}\text{CO}_3$ (1.85×10^{12} Bq mol^{-1}) and 4 mM NaH_2PO_4 , to activate Rubisco by incubation at 37.5°C for 40 min. Reaction mixtures were prepared in oxygen electrodes (Oxygraph, Hansatech instruments Ltd., Norfolk, UK) by first adding 0.95 mL of CO_2 -free assay buffer (100 mM Bicine pH 8.2, 20 mM MgCl_2 , containing 0.015 mg of carbonic anhydrase). After the addition of 0.02 mL of 0.1 M $\text{NaH}^{14}\text{CO}_3$ (1.85×10^{12} Bq mol^{-1}), the plug was fitted to the oxygen electrode vessel and enough activated Rubisco (20 μL) was added. The reaction was started by the injection of 10 μL of 25 mM RuBP to be completed between 2 and 7 min depending on the assay temperature. RuBP oxygenation was calculated from the oxygen consumption and carboxylation from the amount of ^{14}C incorporated into PGA when all the RuBP had been consumed (Galmés et al., 2014a). Measurements were performed at 15 $^\circ\text{C}$, 25 $^\circ\text{C}$ and 35 $^\circ\text{C}$, with 3-9 biological replicates per each species and assayed temperature.

For all Rubisco assays, pH of the assay buffers was accurately adjusted at each temperature of measurement. The concentration of CO₂ in solution in equilibrium with HCO₃⁻ was calculated assuming a pK_a for carbonic acid of 6.19, 6.11 and 6.06 at 15 °C, 25 °C and 35 °C, respectively. The concentration of O₂ in solution was assumed to be 305.0, 253.4 and 219.4 (nmol mL⁻¹) at 15 °C, 25 °C and 35 °C, respectively (Truesdale and Downing 1954).

Temperature dependence parameters of Rubisco kinetics

To determine the temperature response of the Rubisco kinetic parameters from each species, values for K_c , K_c^{air} and $S_{c/o}$ were first converted from concentrations to partial pressures. For this, solubilities for CO₂ were considered to be 0.0450, 0.0340 and 0.0262 mol L⁻¹ bar⁻¹ at 15 °C, 25 °C and 35 °C, respectively. In turn, solubilities for O₂ of 0.0016, 0.0013 and 0.0011 mol L⁻¹ bar⁻¹ were used at 15 °C, 25 °C and 35 °C, respectively. The CO₂ compensation point in the absence of mitochondrial respiration (I^*) was obtained from $S_{c/o}$ as in von Caemmerer (2000) using the above solubilities for O₂. Thereafter, values of K_c , I^* and k_{cat}^c at the three temperatures were fitted to an Arrhenius-type equation (Badger and Collatz 1977; Harley and Tenhunen 1991):

$$Parameter = \exp \left[c - \frac{\Delta H_a}{RT_k} \right]$$

where c is a scaling constant, ΔH_a is the energy of activation, R is the molar gas constant (8.314 J K⁻¹ mol⁻¹) and T_k is the absolute assay temperature.

CO₂ assimilation potential of crop Rubiscos at varying temperatures and CO₂ availability

According to the biochemical model of C₃ photosynthesis (Farquhar et al., 1980), the Rubisco CO₂ assimilation potential (A_{Rubisco}) is defined as the minimum of the RuBP-saturated (A_c) and RuBP-limited (A_j) CO₂ assimilation rates:

$$(1) A_{\text{Rubisco}} = \min (A_c, A_j),$$

$$(2) A_c = \frac{k_{\text{cat}}^c \cdot E \cdot (C_c - \Gamma^*)}{C_c + K_c^{\text{air}}}$$

$$(3) A_j = \frac{(C_c - \Gamma^*) J}{4C_c + 8\Gamma^*}$$

A_{Rubisco} was obtained for each species at three different temperatures, 15 °C, 25 °C and 35 °C, and two different concentrations of CO₂ in the chloroplast stroma (C_c), 150 and 250 μbar , simulating situations of moderate water-stress and well-watered conditions in C₃ plants, respectively (Flexas et al., 2006). The Rubisco catalytic traits k_{cat}^c , Γ^* and K_c^{air} were taken from the species- and temperature-specific data obtained in the present study. The concentration of active Rubisco sites (E) was assumed invariable at 25 $\mu\text{mol m}^{-2}$. Values of the CO₂-saturated photosynthetic electron transport rates (J) were assumed 60, 150 and 212 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at 15 °C, 25 °C and 35 °C, respectively, for all species. At 25 °C, $J = 150 \mu\text{mol m}^{-2} \text{s}^{-1}$ matches very well with a $J/(k_{\text{cat}}^c \cdot E)$ ratio of 1.5 (Egea et al., 2011). Values for J at 15 °C and 35 °C were obtained from the J temperature response described for tobacco in Walker et al. (2013).

Analysis of Rubisco L-subunit sites under positive selection

Full length DNA sequences of the Rubisco large subunit (L-subunit) encoding gene, *rbcL* (Fig. S3), and two additional chloroplast genes (*matK* and *ndhF*) were obtained from GenBank (<http://www.ncbi.nlm.nih.gov/genbank/>) for the twenty studied species. Accession numbers information is given in the Table S2.

DNA sequences were translated into protein sequences for alignment using MUSCLE (Edgar 2004). The software MODELTEST 3.7 (Posada and Crandall 1998, Posada and Buckley 2004) was used to check for the best model before running the phylogenetic analyses. The species phylogeny was reconstructed using concatenated alignment of all three chloroplast genes and maximum-likelihood inference conducted with RAxML version 7.2.6 (Stamatakis 2006).

Amino acid residues under positive selection were identified using codon-based substitution models in comparative analysis of protein-coding DNA sequences within the phylogenetic framework (Yang 1997). Given the conservative assumption of no selective pressure at synonymous sites, codon-based substitution models assume that codons with the ratio of nonsynonymous/synonymous substitution rate (d_N/d_S) less than one evolve under purifying selection to keep protein function and properties, while codons with $d_N/d_S > 1$ evolve under positive Darwinian selection to modify properties of the given protein (Yang 1997).

The codeml program in the PAML v4.7 package (Yang 2007) was used to perform branch-site tests of positive selection along pre-specified foreground branches (Yang et al., 2005, Yang 2007). The codeml A model allows $0 \leq d_N/d_S \leq 1$ and $d_N/d_S = 1$ for all branches. The $d_N/d_S > 1$ is permitted only along pre-specified foreground branches and $0 \leq d_N/d_S \leq 1$ and $d_N/d_S = 1$ on background branches. Branches leading to species with high or low K_c , k_{cat}^c , $S_{c/o}$ and ΔH_a were marked as foreground branches. For the purpose of these tests, high or low K_c , k_{cat}^c and $S_{c/o}$ ranges were taken only at 25 °C because of high correlation between values for these kinetic parameters obtained at three different temperatures. ΔH_a for these kinetic parameters were also considered. The A model was used to identify the amino acid sites under positive selection and to calculate the posterior probabilities of an amino

acid belongs to a class with $d_N/d_S > 1$ using the Bayes empirical Bayes (BEB) approach implemented in PAML (Yang et al., 2005).

The Rubisco L-subunit residues were numbered based on the spinach sequence. The location of sites under positive selection was done using Rubisco protein structure from spinach (*Spinacia oleracea* L.) obtained from the RCSB Protein Data Bank (<http://www.rcsb.org>; file 1RCX; Karkehabadi et al., 2003).

Statistical analysis

Statistical analysis consisted of one-way ANOVA and correlation for linear regressions. For all the parameters studied, a univariate model of fixed effects was assumed. The univariate general linear model for unbalanced data (Proc. GLM) was applied and significant differences among species and groups of species were revealed by Duncan tests using IBM SPSS Statistics for Macintosh, Version 21.0. (Armonk, NY: IBM Corp software package). The relationships among the kinetic parameters and the temperature dependence parameters were tested with the square of the correlation coefficient observed for linear regressions using the tool implemented in R 3.1.1 (R Development Core Team 2014, <http://www.R-project.org>). All statistical tests were considered significant at $p < 0.05$.

The Pearson correlation coefficient was calculated between pairwise combinations of the kinetic parameters K_c , K_c^{air} , k_{cat}^c and $S_{c/o}$ at the three temperatures of measurement. However, correlations arising within groups of related taxa might reflect phylogenetic signal rather than true cause-effect relationships, because closely related taxa are not necessarily independent data points and could violate the assumption of randomized sampling employed by conventional statistical methods (Felsenstein 1985). To overcome this issue, tests were performed for the presence of

phylogenetic signal in the data and trait correlations were calculated with phylogenetically independent contrasts using the AOT module of PHYLOCOM (Webb et al., 2008) using the species phylogeny based on the three chloroplast genes (see below). All these tests were considered significant at $p < 0.05$.

ACKNOWLEDGEMENTS

Trinidad Garcia is acknowledged for her technical help and organization of the radioisotope installation at the Serveis Científic-Tècnics of UIB while running these experiments. We would like to thank Mr. Miquel Truyols and collaborators of the UIB Experimental Field and Greenhouses, which are supported by the UIBGrant 15/2015. We would like to thank Dra. Arantxa Molins for her experimental help. We are also very grateful to Dr. Cyril Douthe, Dr. Josep Cifre, Mrs. Patricia González and Dr. Jorge Gago for their help in improving different parts of the manuscript. Mr. Jaume Jaume and Dr. Sebastià Martorell are acknowledged for providing some plant material used in this experiment.

Supplemental Material

Table S1. The Rubisco kinetic parameters measured at 15 °C and 35 °C for the selected crop species.

Table S2. List of crop species and GenBank accession numbers for *rbcL*, *matK* and *ndhF*.

Figure S1. Box plots depiction of Rubisco kinetic parameters (K_c , K_c^{air} , k_{cat}^c and $S_{c/o}$) at 15 °C, 25 °C and 35 °C when considering the 18 C_3 species alone.

Figure S2. Maximum likelihood phylogeny created using *rbcL*, *matK* and *ndhF* for the selected crop species.

Figure S3. Rubisco L-subunit amino acid alignment for the 20 crops species used in this study.

646 **Table 1.** Kinetic parameters of crop Rubiscos measured at 25 °C: the Michaelis-Menten constants for CO₂ under non-oxygenic (K_c) and 21% O₂ (K_c^{air}), the
647 maximum carboxylation rate (k_{cat}^c), the specificity factor ($S_{c/o}$), and the carboxylation (k_{cat}^c/K_c and $k_{\text{cat}}^c/K_c^{\text{air}}$), and the oxygenation catalytic efficiencies
648 (k_{cat}^o/K_o). The k_{cat}^o/K_o ratio was calculated as $[(k_{\text{cat}}^c/K_c)/S_{c/o} \times 1000]$. For each species, data are mean \pm standard error (n = 3-9). Group averages were obtained
649 from individual measurements on each species. Different letters denote statistical differences ($p < 0.05$) by Duncan analysis between C₃ and C₄ groups.
650

Species	K_c (μM)	K_c^{air} (μM)	k_{cat}^c (s^{-1})	$S_{c/o}$ (mol mol^{-1})	k_{cat}^c/K_c ($\text{s}^{-1} \mu\text{M}^{-1}$)	$k_{\text{cat}}^c/K_c^{\text{air}}$ ($\text{s}^{-1} \mu\text{M}^{-1}$)	k_{cat}^o/K_o ($\text{s}^{-1} \text{nM}^{-1}$)
C ₃ species							
<i>Avena sativa</i>	10.8 \pm 0.9	18.1 \pm 2.0	2.3 \pm 0.3	99.9 \pm 3.0	0.21 \pm 0.01	0.13 \pm 0.03	2.14 \pm 0.06
<i>Beta vulgaris</i>	10.8 \pm 1.2	18.6 \pm 1.1	2.0 \pm 0.3	100.8 \pm 2.0	0.19 \pm 0.02	0.10 \pm 0.01	1.94 \pm 0.31
<i>Brassica oleracea</i>	11.8 \pm 0.1	19.2 \pm 0.3	2.1 \pm 0.3	96.2 \pm 1.3	0.17 \pm 0.03	0.11 \pm 0.02	1.81 \pm 0.28
<i>Capsicum annuum</i>	9.6 \pm 0.3	19.8 \pm 1.5	1.9 \pm 0.1	96.0 \pm 4.5	0.20 \pm 0.01	0.10 \pm 0.01	1.98 \pm 0.15
<i>Coffea Arabica</i>	11.0 \pm 0.4	22.9 \pm 2.4	2.1 \pm 0.2	98.7 \pm 3.8	0.19 \pm 0.02	0.08 \pm 0.01	1.98 \pm 0.18
<i>Cucurbita maxima</i>	9.0 \pm 0.5	19.2 \pm 1.0	2.2 \pm 0.2	98.4 \pm 0.4	0.25 \pm 0.04	0.12 \pm 0.01	2.45 \pm 0.31
<i>Glycine max</i>	8.6 \pm 0.2	16.2 \pm 0.7	1.5 \pm 0.1	97.0 \pm 1.1	0.17 \pm 0.02	0.09 \pm 0.01	1.76 \pm 0.21
<i>Hordeum vulgare</i>	9.0 \pm 0.6	14.9 \pm 1.6	2.4 \pm 0.2	99.2 \pm 3.8	0.28 \pm 0.02	0.17 \pm 0.03	3.01 \pm 0.19
<i>Ipomoea batatas</i>	12.0 \pm 0.7	21.1 \pm 1.0	2.5 \pm 0.1	98.5 \pm 6.6	0.20 \pm 0.00	0.12 \pm 0.01	1.96 \pm 0.08
<i>Lactuca sativa</i>	11.1 \pm 0.3	18.2 \pm 1.4	2.2 \pm 0.1	94.0 \pm 1.9	0.19 \pm 0.00	0.12 \pm 0.01	2.06 \pm 0.07

<i>Manihot esculenta</i>	6.1 ± 0.2	10.8 ± 0.6	1.4 ± 0.1	100.8 ± 0.9	0.23 ± 0.02	0.13 ± 0.01	2.24 ± 0.17
<i>Medicago sativa</i>	9.7 ± 1.6	16.4 ± 1.9	1.7 ± 0.1	95.6 ± 2.2	0.20 ± 0.02	0.11 ± 0.01	2.23 ± 0.36
<i>Oryza sativa</i>	8.0 ± 0.4	17.3 ± 2.4	2.1 ± 0.3	93.1 ± 1.2	0.26 ± 0.04	0.14 ± 0.03	2.73 ± 0.43
<i>Phaseolus vulgaris</i>	7.8 ± 0.3	14.0 ± 1.0	1.7 ± 0.2	99.7 ± 2.7	0.22 ± 0.02	0.13 ± 0.02	2.11 ± 0.17
<i>Solanum lycopersicum</i>	9.7 ± 0.4	16.6 ± 1.4	2.3 ± 0.2	92.4 ± 2.3	0.24 ± 0.02	0.14 ± 0.01	2.48 ± 0.20
<i>Solanum tuberosum</i>	9.6 ± 0.2	18.0 ± 0.8	2.0 ± 0.3	95.4 ± 2.3	0.22 ± 0.05	0.12 ± 0.03	2.32 ± 0.46
<i>Spinacia oleracea</i>	14.1 ± 0.8	26.9 ± 0.8	2.4 ± 0.1	97.0 ± 1.2	0.18 ± 0.01	0.09 ± 0.01	1.76 ± 0.13
<i>Triticum aestivum</i>	11.3 ± 0.4	16.0 ± 0.6	2.2 ± 0.2	100.1 ± 1.8	0.20 ± 0.02	0.14 ± 0.01	2.08 ± 0.24
<i>C₃ average</i>	10.0 ± 0.3 ^a	18.0 ± 0.5 ^a	2.1 ± 0.1 ^a	97.5 ± 0.6 ^a	0.21 ± 0.01 ^a	0.12 ± 0.01 ^a	2.17 ± 0.07 ^a
C ₄ species							
<i>Saccharum × officinarum</i>	26.3 ± 4.0	31.7 ± 2.1	3.9 ± 0.3	82.2 ± 1.8	0.15 ± 0.02	0.13 ± 0.01	1.82 ± 0.35
<i>Zea mays</i>	31.6 ± 1.8	42.0 ± 2.8	4.1 ± 0.6	87.3 ± 1.4	0.11 ± 0.02	0.07 ± 0.01	1.22 ± 0.20
<i>C₄ average</i>	27.6 ± 2.3 ^b	36.1 ± 2.6 ^b	4.0 ± 0.5 ^b	84.4 ± 1.5 ^b	0.13 ± 0.02 ^b	0.10 ± 0.01 ^a	1.52 ± 0.23 ^b

651

652

653 **Table 2.** Phylogenetically independent contrasts (PICs, upper part of the diagonals) and Pearson's correlation coefficients (PCCs, lower part of the diagonals)
654 between the Rubisco kinetic parameters (K_c , K_c^{air} , k_{cat}^c and $S_{c/o}$) at 15 °C, 25 °C and 35 °C when considering the 20 C₃ and C₄ species together and the 18 C₃
655 species alone. Significant correlations are marked: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$.

Data from C ₃ and C ₄ species analysed together											
15 °C				25 °C				35 °C			
K_c	K_c^{air}	k_{cat}^c	$S_{c/o}$	K_c	K_c^{air}	k_{cat}^c	$S_{c/o}$	K_c	K_c^{air}	k_{cat}^c	$S_{c/o}$
K_c	0.826***	0.502*	-0.314	K_c	0.913***	0.819***	-0.202	K_c	0.960***	0.710***	-0.775***
K_c^{air}	0.927***	0.036	-0.099	K_c^{air}	0.946***	0.683***	0.037	K_c^{air}	0.962***	0.707***	-0.660**
k_{cat}^c	0.810***	0.645**	-0.660**	k_{cat}^c	0.941***	0.890***	-0.450*	k_{cat}^c	0.894***	0.858***	-0.634**
$S_{c/o}$	-0.498*	-0.361	-0.673**	$S_{c/o}$	-0.772***	-0.699***	-0.749***	$S_{c/o}$	-0.806***	-0.737***	-0.736***
Data from C ₃ species alone											
15 °C				25 °C				35 °C			
K_c	K_c^{air}	k_{cat}^c	$S_{c/o}$	K_c	K_c^{air}	k_{cat}^c	$S_{c/o}$	K_c	K_c^{air}	k_{cat}^c	$S_{c/o}$
K_c	0.900***	0.194	0.120	K_c	0.646**	0.256	0.057	K_c	0.907***	0.363	-0.357
K_c^{air}	0.892***	-0.118	0.394	K_c^{air}	0.829***	0.173	-0.006	K_c^{air}	0.816***	0.401	-0.155
k_{cat}^c	0.268	0.137	-0.787**	k_{cat}^c	0.698***	0.587*	-0.470*	k_{cat}^c	0.613**	0.476*	-0.285
$S_{c/o}$	0.025	0.145	-0.496*	$S_{c/o}$	-0.049	-0.162	-0.083	$S_{c/o}$	-0.386	-0.157	-0.259

Table 3. The energy of activation (ΔH_a , kJ mol⁻¹) and c (dimensionless) values of the Rubisco Michaelis-Menten constants for CO₂ under non-oxygenic (K_c , $\mu\text{mol mol}^{-1}$) and 21% O₂ (K_c^{air} , $\mu\text{mol mol}^{-1}$), the maximum carboxylation rate (k_{cat}^c , s⁻¹) and the CO₂ compensation point in the absence of mitochondrial respiration (Γ^* , $\mu\text{mol mol}^{-1}$) for the twenty crop species. For each species, data are mean \pm standard error (n = 3-9). Group averages were calculated from individual measurements on each species. Different letters denote statistical differences ($p < 0.05$) by Duncan analysis between C₃ and C₄ groups. Parameter concentrations of K_c (μM) and K_c^{air} (μM) in liquid phase (Table 1 and S2) were converted to gaseous phase partial pressures [K_c and/or K_c^{air} ($\mu\text{mol mol}^{-1}$) = parameter (μM) $\times K_h \times \text{Air Volume (L)} / \text{RT}$]. K_h is the hydrolysis constant (15 °C = 22.2, 25 °C = 29.4, 35 °C = 38.2). For the Air Volume (L): 15 °C = 23.7, 25 °C = 24.5, 35 °C = 25.4. The term Γ^* ($\mu\text{mol mol}^{-1}$) is derived from $0.5O / S_{c/o}$.

Species	K_c		K_c^{air}		k_{cat}^c		Γ^*	
	c	ΔH_a	c	ΔH_a	c	ΔH_a	c	ΔH_a
C ₃ species								
<i>Avena sativa</i>	31.3 \pm 0.7	63.4 \pm 2.0	26.0 \pm 0.4	48.9 \pm 1.2	17.6 \pm 2.2	41.5 \pm 5.5	13.3 \pm 0.5	23.6 \pm 1.4
<i>Beta vulgaris</i>	28.7 \pm 1.7	57.0 \pm 4.4	27.2 \pm 0.7	51.8 \pm 1.8	21.5 \pm 3.7	51.2 \pm 9.6	11.7 \pm 0.4	19.8 \pm 1.0
<i>Brassica oleracea</i>	28.1 \pm 1.1	55.3 \pm 2.8	26.5 \pm 0.9	50.1 \pm 2.4	18.8 \pm 2.6	45.7 \pm 6.5	12.6 \pm 0.2	21.8 \pm 0.5
<i>Capsicum annum</i>	26.6 \pm 1.5	51.8 \pm 3.7	27.0 \pm 1.6	51.2 \pm 3.7	16.3 \pm 2.8	39.2 \pm 6.9	13.4 \pm 0.7	24.1 \pm 1.8
<i>Coffea arabica</i>	34.7 \pm 0.3	71.5 \pm 0.9	27.6 \pm 1.8	52.2 \pm 4.3	16.5 \pm 2.6	39.0 \pm 6.1	13.1 \pm 0.5	23.4 \pm 1.1
<i>Cucurbita maxima</i>	28.6 \pm 0.8	57.0 \pm 1.8	29.2 \pm 1.1	56.8 \pm 2.8	20.2 \pm 1.0	48.7 \pm 2.7	12.2 \pm 0.9	21.1 \pm 2.2

<i>Glycine max</i>	34.2 ± 0.5	71.1 ± 1.4	28.4 ± 1.2	55.3 ± 2.9	22.7 ± 2.5	55.2 ± 5.8	14.4 ± 1.7	26.5 ± 4.1
<i>Hordeum vulgare</i>	31.1 ± 1.1	63.4 ± 3.0	30.7 ± 1.9	60.9 ± 5.0	12.2 ± 1.6	27.9 ± 4.0	12.3 ± 0.2	21.2 ± 0.6
<i>Ipomoea batatas</i>	23.0 ± 0.7	42.4 ± 1.6	22.7 ± 1.2	40.7 ± 3.1	14.3 ± 1.5	33.4 ± 3.8	13.0 ± 0.3	22.8 ± 0.8
<i>Lactuca sativa</i>	28.3 ± 1.3	55.8 ± 3.2	29.0 ± 2.1	56.5 ± 5.2	14.1 ± 0.7	33.3 ± 1.7	12.3 ± 0.3	21.2 ± 0.9
<i>Manihot esculenta</i>	33.7 ± 1.4	70.8 ± 3.4	36.1 ± 1.1	75.4 ± 2.8	19.8 ± 1.6	47.4 ± 4.1	12.2 ± 0.2	21.1 ± 0.5
<i>Medicago sativa</i>	29.2 ± 1.3	58.8 ± 3.6	26.1 ± 0.4	49.5 ± 1.0	24.8 ± 1.1	60.5 ± 2.8	11.8 ± 0.2	20.1 ± 0.4
<i>Oryza sativa</i>	38.9 ± 0.8	83.1 ± 1.8	30.5 ± 1.2	60.5 ± 3.1	19.2 ± 1.8	46.4 ± 4.7	13.7 ± 0.5	24.6 ± 1.3
<i>Phaseolus vulgaris</i>	31.5 ± 0.8	64.6 ± 2.0	30.9 ± 2.7	61.7 ± 6.8	19.8 ± 2.1	47.7 ± 5.3	13.4 ± 0.6	24.1 ± 1.5
<i>Solanum lycopersicum</i>	30.8 ± 2.5	62.1 ± 6.3	36.0 ± 2.5	73.8 ± 6.4	14.7 ± 1.4	34.6 ± 3.6	12.5 ± 0.2	21.8 ± 0.5
<i>Solanum tuberosum</i>	21.1 ± 0.2	38.2 ± 0.5	24.4 ± 0.8	44.9 ± 1.9	19.2 ± 0.5	46.2 ± 1.1	13.7 ± 0.9	24.7 ± 2.2
<i>Spinacia oleracea</i>	34.3 ± 0.8	69.9 ± 2.2	25.1 ± 0.5	45.6 ± 1.1	20.2 ± 0.7	48.0 ± 1.8	13.5 ± 0.3	25.2 ± 1.0
<i>Triticum aestivum</i>	30.1 ± 0.5	60.4 ± 2.2	34.4 ± 2.2	70.1 ± 5.4	17.4 ± 1.7	41.2 ± 4.3	13.5 ± 0.2	24.2 ± 0.4
<i>C₃ average</i>	30.2 ± 0.6 ^a	60.9 ± 1.5 ^a	28.8 ± 0.6 ^a	55.9 ± 1.5 ^a	18.3 ± 0.6 ^a	43.7 ± 1.5 ^a	13.0 ± 0.2 ^a	22.9 ± 0.4 ^a
C ₄ species								
<i>Saccharum × officinarum</i>	30.2 ± 1.9	58.3 ± 5.0	32.0 ± 1.0	62.3 ± 2.7	13.6 ± 1.5	30.2 ± 3.5	14.3 ± 0.6	25.8 ± 1.4
<i>Zea mays</i>	24.7 ± 3.4	44.5 ± 8.5	24.7 ± 3.4	44.5 ± 8.5	14.0 ± 0.9	31.0 ± 1.9	13.6 ± 0.1	24.3 ± 0.2
<i>C₄ average</i>	27.9 ± 2.0 ^a	52.4 ± 5.0 ^b	28.9 ± 0.8 ^a	53.7 ± 1.9 ^a	13.7 ± 0.7 ^b	30.6 ± 1.6 ^b	14.0 ± 0.3 ^a	25.0 ± 0.7 ^a

664

665

Table 4. Amino acid replacements in the Rubisco large subunit (L-subunit) identified under positive selection by the Bayes Empirical Bayes (BEB) analysis implemented in the PAML package (Yang et al., 2005; Yang 2007) along branches of the phylogenetic tree leading to species with particular Rubisco properties.

Residue ^a	Amino acid changes	Location of residue	Interaction ^b
Branches leading to species with $K_c \geq 26.0 \mu\text{M}$ and $k_{\text{cat}} \geq 3.9 \text{ s}^{-1}$ at 25 °C (C ₄ species)			
94**	D, E, K → P		ID, RA
446**	R → K	C-terminus	
469**	Insert of G or T before resi 469	C-terminus	ID
Branches leading to species with $k_{\text{cat}} \geq 2.5 \text{ s}^{-1}$ at 25 °C			
281**	A → S	Helix 4	DD, SS
Branches leading to species with $K_c \geq 10.8 \mu\text{M}$ at 25 °C			
439***	A → T, V	Helix G	
469*	Insert of G or T before residue 469	C-terminus	ID
470*	A, E → K, P, Q	C-terminus	ID
477**	S → E, G, P, Q	C-terminus	
Branches leading to species with $S_{\text{c/o}} \leq 94.0 \text{ mol mol}^{-1}$ at 25 °C			
309**	M → I	βF Strand	ID
Branches leading to species with ΔH_a for $K_c \leq 56.0 \text{ kJ mol}^{-1}$			
262**	V → A, T	Loop 3	S-subunit
439*	R → T, V	Helix G	
449**	C, S, T → A	C-terminus	
477**	K → E, G, P, Q	C-terminus	

^a Residue numbering is based on the spinach sequence. Values for Bayesian Posterior Probabilities are: * > 0.90, ** > 0.95, *** > 0.99.

^b Interactions in which the selected residues and/or residues within 5 Å of them are involved. ID – intradimer interactions; DD – dimer-dimer interactions; RA – interface for interactions with Rubisco

675 activase; SS – interactions with small subunits; interactions based on literature survey only are given in
676 italics; after (Spreitzer and Salvucci 2002; Ott et al., 2000; Du et al., 2003).
677

Figure legends

Figure 1. The relationship between the turnover rate for the Rubisco carboxylase reaction ($k_{\text{cat}}^{\text{c}}$) with (A) the Michaelis–Menten affinity constant for CO_2 (K_{c}) and (B) the CO_2/O_2 specificity factor ($S_{\text{c/o}}$). Filled symbols correspond to C_3 species at 15 °C (\blacktriangle), 25 °C (\bullet) and 35 °C (\blacktriangledown); open symbols correspond to C_4 species at 15 °C (\triangle), 25 °C (\circ) and 35 °C (\triangledown). Each symbol represents the average value of a single species per temperature interaction.

Figure 2. Simulated CO_2 assimilation potential of Rubisco (A_{Rubisco}) for the C_3 and C_4 species at 15 °C, 25 °C and 35 °C and at values for the chloroplastic CO_2 concentration (C_{c}) of (A) 250 μbar and (B) 150 μbar . Equations used to calculate A_{Rubisco} were those described in the biochemical model of C_3 photosynthesis (Farquhar et al., 1980), as explained in Materials and Methods. The bars represent the minimum value of A_{c} - and A_{j} -limited A_{Rubisco} . Asterisks (*) above the bars indicate A_{c} -limited A_{Rubisco} (absence of * indicate A_{j} -limited A_{Rubisco}). The rate of electron transport was considered 60, 150 and 212 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at 15 °C, 25 °C and 35 °C, respectively. The concentration of active Rubisco sites was assumed invariable at 25 $\mu\text{mol m}^{-2}$ for all the species and environmental conditions. The values used for the Rubisco kinetic parameters ($k_{\text{cat}}^{\text{c}}$, I^* and $K_{\text{c}}^{\text{air}}$) are those shown in Tables 1 and S2.

REFERENCES

Badger MR, Collatz GJ (1977) Studies on the kinetic mechanism of ribulose-1,5-bisphosphate carboxylase and oxygenase reactions, with particular reference to the effect of temperature on kinetic parameters. *Carnegie Inst Wash Year b* 76:355–361

703 **Badger MR** (1980) Kinetic properties of ribulose1,5-bisphosphate
 704 carboxylase/oxygenase from *Anabaena variabilis*. Arch Biochem Biophys
 705 201: 247–254

706 **Balaguer L, Afif D, Dizengremel P, Dreyer E** (1996) Specificity factor of ribulose
 707 bisphosphate carboxylase/oxygenase of *Quercus robur*. Plant Physiol Bioch
 708 34: 879–883

709 **Bermúdez MA, Galmés J, Moreno I, Mullineaux PM, Gotor C, Romero LC**
 710 (2012) Photosynthetic adaptation to length of day is dependent on S-
 711 sulfocysteine synthase activity in the thylakoid lumen. Plant Physiol 160: 274–
 712 288

713 **Bernacchi CJ, Singaas EL, Pimentel C, Portis Jr AR, Long SP** (2001) Improved
 714 temperature response functions for models of Rubisco- limited
 715 photosynthesis. Plant Cell Environ 24: 253–259

716 **Bernacchi CJ, Portis AR, Nakano H, von Caemmerer S, Long SP** (2002)
 717 Temperature response of mesophyll conductance. Implications for the
 718 determination of Rubisco enzyme kinetics and for limitations to
 719 photosynthesis *in vivo*. Plant Physiol 130: 1992–1998

720 **Bird IF, Cornelius MJ, Keys AJ** (1982) Affinity of RuBP carboxylases for carbon
 721 dioxide and inhibition of the enzymes by oxygen. J Exp Bot 33:1004–1013

722 **Bota J, Flexas J, Keys A J, Loveland J, Parry MAJ, Medrano H** (2002) CO₂/O₂
 723 specificity factor of ribulose-1,5-bisphosphate carboxylase/oxygenase in
 724 grapevines (*Vitis vinifera* L.): first *in vitro* determination and comparison to *in*
 725 *vivo* estimations. Vitis 41: 163.

726 **Bremer B, Bremer K, Chase M, Fay M, Reveal J, Soltis D, Stevens P** (2009) An
 727 update of the Angiosperm Phylogeny Group classification for the orders and
 728 families of flowering plants: APG III. *Bot J Linn Soc* 161: 105–121

729 **Brooks A, Farquhar GD** (1985) Effect of temperature on the CO₂/O₂ specificity of
 730 ribulose-1,5-bisphosphate carboxylase/oxygenase and the rate of respiration in
 731 the light. *Planta* 165: 397–406.

732 **Castrillo M** (1995) Ribulose-1,5-bis-phosphate carboxylase activity in altitudinal
 733 populations of *Espeletia schultzii* Wedd. *Oecologia* 101: 193–196

734 **Chabot BF, Chabot JF, Billings WD** (1972) Ribulose-1,5-diphosphate carboxylase
 735 activity in arctic and alpine populations of *Oxyria digyna*. *Photosynthetica* 6:
 736 364–369

737 **Christin PA, Salamin N, Muasya AM, Roalson EH, Russier F, Besnard G** (2008)
 738 Evolutionary switch and genetic convergence on *rbcL* following the evolution
 739 of C₄ photosynthesis. *Mol Biol Evol* 25: 2361–2368

740 **Cousins AB, Ghannoum O, von Caemmerer, S, Badger MR** (2010) Simultaneous
 741 determination of Rubisco carboxylase and oxygenase kinetic parameters in
 742 *Triticum aestivum* and *Zea mays* using membrane inlet mass spectrometry.
 743 *Plant Cell Environ* 33: 444–452

744 **DaMatta FM** (2004) Exploring drought tolerance in coffee: a physiological approach
 745 with some insights for plant breeding. *Braz J Plant Physiol* 16: 1–6

746 **Delgado E, Medrano H, Keys AJ, Parry MAJ** (1995) Species variation in Rubisco
 747 specificity factor. *J Exp Bot* 46: 1775–1777

748 **Díaz-Espejo A** (2013) New challenges in modelling photosynthesis: temperature
 749 dependencies of Rubisco kinetics. *Plant Cell Environ* 36: 2104–2107

750 **Du YC, Peddi SR, Spreitzer RJ** (2003) Assessment of structural and functional
 751 divergence far from the large subunit active site of Ribulose-1,5-bisphosphate
 752 carboxylase/oxygenase. *J Biol Chem*, 278: 49401–49405.

753 **Edgar RC** (2004) MUSCLE: multiple sequence alignment with high accuracy and
 754 high throughput. *Nucleic Acids Res* 32: 1792–1797

755 **Egea G, González-Real MM, Baille A, Nortes PA, Díaz-Espejo A.** (2011)
 756 Disentangling the contributions of ontogeny and water stress to photosynthetic
 757 limitations in almond trees. *Plant Cell Environ* 34: 962–979

758 FAO (Food and Agriculture Organization of the United Nations) Available at
 759 <http://www.fao.org/economic/ess/ess-home/es/>

760 FAOSTAT (2010) Available at <http://www.faostat.fao.org/>

761 **Farquhar GD, von Caemmerer S, Berry JA** (1980) A biochemical model of
 762 photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* 149: 78–90

763 **Felsenstein J** (1985) Phylogenies and the comparative method. *Am Nat* 125: 1–15

764 **Flexas J, Ribas- Carbó M, Bota J, Galmés J, Henkle M, Martínez- Cañellas S,**
 765 **Medrano H** (2006) Decreased Rubisco activity during water stress is not
 766 induced by decreased relative water content but related to conditions of low
 767 stomatal conductance and chloroplast CO₂ concentration. *New Phytol* 172: 73-
 768 82.

769 **Flexas J, Galmés J, Gallé A, Gulías J, Pou A, Ribas-Carbó M, Tomás M,**
 770 **Medrano H** (2010) Improving water use efficiency in grapevines: potential
 771 physiological targets for biotechnological improvement. *Aust J Grape Wine*
 772 *Research*, 16: 106–121

773 **Gago J, Coopman RE, Cabrera HM, Hermida C, Molins A, Conesa MÀ, Galmés**
774 **J, Ribas-Carbó M, Flexas J** (2013) Photosynthesis limitations in three fern
775 species. *Physiol Plantarum* 149: 599–611

776 **Galili S, Avivi Y, Feldman M** (1998) Differential expression of three *RbcS*
777 subfamilies in wheat. *Plant Sci*, 139: 85-193

778 **Galmés J, Flexas J, Keys AJ, Cifre J, Mitchell RAC, Madgwick PJ, RP Haslam,**
779 **Medrano H, Parry MAJ** (2005) Rubisco specificity factor tends to be larger
780 in plant species from drier habitats and in species with persistent leaves. *Plant*
781 *Cell Environ* 28: 571–579

782 **Galmés J, Ribas-Carbó M, Medrano H, Flexas J** (2011) Rubisco activity in
783 Mediterranean species is regulated by the chloroplastic CO₂ concentration
784 under water stress. *J Exp Bot* 62: 653–665

785 **Galmés J, Perdomo JA, Flexas J, Whitney SM** (2013) Photosynthetic
786 characterization of Rubisco transplasmic lines reveals alterations on
787 photochemistry and mesophyll conductance. *Photosynth Res* 115: 153-166.

788 **Galmés J, Andralojc PJ, Kapralov MV, Flexas J, Keys A J, Molins A, Parry**
789 **MAJ, Conesa MÀ** (2014a) Environmentally driven evolution of Rubisco and
790 improved photosynthesis and growth within the C₃ genus *Limonium*
791 (Plumbaginaceae). *New Phytol* 203: 989–999

792 **Galmés J, Conesa MÀ, Díaz-Espejo A, Mir A, Perdomo JA, Niinemets Ü, Flexas**
793 **J.** (2014b) Rubisco catalytic properties optimized for present and future
794 climatic conditions. *Plant Sci* 226: 61–70

795 **Galmés J, Kapralov MV, Andraloj P, Conesa MÀ, Keys AJ, Parry MAJ, Flexas**
796 **J.** (2014c) Expanding knowledge of the Rubisco kinetics variability in plant

797 species: environmental and evolutionary trends. *Plant Cell Environ* 37: 1989–
 798 2001

799 **Galmés J, Kapralov MV, Copolovici LO, Hermida-Carrera C, Niinemets Ü**
 800 (2015) Temperature responses of the Rubisco maximum carboxylase activity
 801 across domains of life: phylogenetic signals, trade-offs, and importance for
 802 carbon gain. *Photosynth Res* 123: 183–201.

803 GenBank database. Available at <http://www.ncbi.nlm.nih.gov/genbank/>

804 **Ghannoum O, Evans JR, Chow WS, Andrews TJ, Conroy JP, von Caemmerer S.**
 805 (2005) Faster Rubisco is the key to superior nitrogen-use efficiency in NADP-
 806 malic enzyme relative to NAD-malic enzyme C₄ grasses. *Plant Physiol* 137:
 807 638–650

808 **Gornall J, Betts R, Burke E, Clark R, Camp J, Willett K, Wiltshire A** (2010)
 809 Implications of climate change for agricultural productivity in the early
 810 twenty-first century. *Philos T Soc B* 365: 2973–2989

811 **Hall NP, Keys AJ** (1983) Temperature dependence of the enzymic carboxylation and
 812 oxygenation of ribulose 1,5-bisphosphate in relation to effects of temperature
 813 on photosynthesis. *Plant Physiol* 72: 945–948

814 **Harley PC, Tenhunen JD** (1991) Modeling the photosynthetic response of C₃ leaves
 815 to environmental factors. In: Boote KJ, Loomis RS, eds. *Modeling crop*
 816 *photosynthesis—from biochemistry to canopy*. Madison: Crop Science
 817 Society of America, Inc. and American Society of Agronomy, Inc. pp 17–39.

818 **Haslam RP, Keys AJ, Andralojc PJ, Madgwick PJ, Inger A, Grimsrud A,**
 819 **Eilertsen HC, Parry MAJ** (2005) Specificity of diatom Rubisco. In: Omasa
 820 K, Nouchi I, De Kok LJ, eds. *Plant Responses to Air Pollution and Global*
 821 *Change*. Springer Japan. pp 157–164

822 IBM SPSS Statistics for Macintosh, Version 21.0. Armonk, NY: IBM Corp. Released
823 2012

824 **Iida S, Miyagi A, Aoki S, Ito M, Kadono Y, Kosuge K** (2009) Molecular adaptation
825 of *rbcL* in the heterophyllous aquatic plant *Potamogeton*. PLoS One, 4: e4633

826 **Ishikawa C, Hatanaka T, Misoo S, Fukayama H** (2009) Screening of high k_{cat}
827 Rubisco among Poaceae for improvement of photosynthetic CO₂ assimilation
828 in rice. Plant Prod Sci 12: 345–350

829 **Ishikawa C, Hatanaka T, Misoo S, Miyake C, Fukayama H** (2011) Functional
830 incorporation of sorghum small subunit increases the catalytic turnover rate of
831 Rubisco in transgenic rice. Plant Physiol 156: 1603–1611.

832 **Jordan DB, Ogren WL** (1983) Species variation in kinetic properties of ribulose 1,5-
833 bisphosphate carboxylase/oxygenase. Arch Biochem Biophys 227: 425–433

834 **Jordan DB, Ogren WL** (1984) The CO₂/O₂ specificity of ribulose 1,5-bisphosphate
835 carboxylase/oxygenase. Planta 161: 308–313

836 **Kane HJ, Viil J, Entsch B, Paul K, Morell MK, Andrews TJ** (1994) An improved
837 method for measuring the CO₂/O₂ specificity of ribulosebisphosphate
838 carboxylase-oxygenase. Funct Plant Biol 21: 449–461

839 **Kapralov MV, Filatov DA** (2007) Widespread positive selection in the
840 photosynthetic Rubisco enzyme. BMC Evolutionary Biology 7: 73

841 **Kapralov MV, Kubien DS, Andersson I, Filatov DA** (2011) Changes in Rubisco
842 kinetics during the evolution of C₄ photosynthesis in *Flaveria* (Asteraceae) are
843 associated with positive selection on genes encoding the enzyme. Mol Biol
844 Evol 28: 1491–1503

845 **Kapralov MV, Smith JAC, Filatov DA** (2012) Rubisco Evolution in C₄ Eudicots:
846 An Analysis of Amaranthaceae *Sensu Lato*. PloS One 7 (12): e52974.

847 **Karkehabadi S, Taylor TC, Andersson I** (2003) Calcium supports loop closure but
848 not catalysis in Rubisco. *J M Biol* 334: 65–73

849 **Keenan T, Sabate S, Gracia C** (2010) Soil water stress and coupled photosynthesis–
850 conductance models: Bridging the gap between conflicting reports on the
851 relative roles of stomatal, mesophyll conductance and biochemical limitations
852 to photosynthesis. *Agr Forest Meteorol* 150: 443–453

853 **Kent SS, Tomany MJ** (1995) The differential of the ribulose 1,5-bisphosphate
854 carboxylase/oxygenase specificity factor among higher plants and the potential
855 for biomass enhancement. *Plant Physiol Biochem* 33: 71–80

856 **Kubien DS, Whitney SM, Moore PV, Jesson LK** (2008) The biochemistry of
857 Rubisco in *Flaveria*. *J Exp Bot* 59: 1767–1777

858 **Laing WA, Ogren WL, Hageman RH** (1974) Regulation of soybean net
859 photosynthetic CO₂ fixation by the interaction of CO₂, O₂, and ribulose 1,5-
860 diphosphate carboxylase. *Plant Physiol* 54: 678–685

861 **Lehnherr B, Mächler F, Nösberger J** (1985) Influence of temperature on the ratio
862 of ribulose bisphosphate carboxylase to oxygenase activities and on the ratio
863 of photosynthesis to photorespiration of leaves. *J Exp Bot* 36: 1117–1125

864 **Meyer RS, DuVal AE, Jensen HR** (2012) Patterns and processes in crop
865 domestication: an historical review and quantitative analysis of 203 global
866 food crops. *New Phytol* 196: 29–48

867 **Monson RK, Stidham MA, Williams GJ, Edwards GE, Uribe EG** (1982)
868 Temperature dependence of photosynthesis in *Agropyron smithii* Rydb. I.
869 Factors affecting net CO₂ uptake in intact leaves and contribution from
870 ribulose-1,5-bisphosphate carboxylase measured *in vivo* and *in vitro*. *Plant*
871 *Physiol* 69: 921–928

872 **Morita K, Hatanaka T, Misoo S, Fukayama, H** (2014) Unusual small subunit that
873 is not expressed in photosynthetic cells alters the catalytic properties of
874 Rubisco in rice. *Plant Physiol* 164: 69–79

875 **Murchie EH, Pinto M, Horton P** (2009) Agriculture and the new challenges for
876 photosynthesis research. *New Phytol* 181: 532–552

877 **Niinemets Ü, Wright IJ, Evans JR** (2009) Leaf mesophyll diffusion conductance in
878 35 Australian sclerophylls covering a broad range of foliage structural and
879 physiological variation. *J Exp Bot* 60: 2433–2449

880 **Ort DR, Merchant SS, Alric J, Barkan A, Blankenship RE, Bock R, Croce R,**
881 **Hanson MR, Hibberd JM, Long SP, Moore TA, Moroney J, Niyogi KK,**
882 **Parry MAJ, Peralta-Yahya PP, Prince RC, Redding KE, Spalding MH,**
883 **van Wijk KJ, Vermaas WFJ, von Caemmerer S, Weber APM, Yeates TO,**
884 **Yuan JS, Zhu XG** (2015) Redesigning photosynthesis to sustainably meet
885 global food and bioenergy demand. *P Natl Acad Sci USA* 112: 8529–8536

886 **Ott CM, Smith BD, Portis AR Jr, Spreitzer RJ** (2000) Activase region on
887 chloroplast Ribulose-1,5-bisphosphate carboxylase/oxygenase. *J Biol Chem*
888 275: 26241–26244

889 **Parry MAJ, Schmidt CNG, Cornelius MJ, Millard BN, Burton S, Gutteridge, S,**
890 **Dyer TA, Keys AJ** (1987) Variations in properties of ribulose-1, 5-
891 bisphosphate carboxylase from various species related to differences in amino
892 acid sequences. *J Exp Bot* 38: 1260–1271

893 **Parry MA, Hawkesford MJ** (2012) An Integrated Approach to Crop Genetic
894 Improvement. *J Integr Plant Biol* 54: 250–259

895 **Parry MA, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H,**
896 **Whitney SM** (2013) Rubisco activity and regulation as targets for crop
897 improvement. *J Exp Bot* 64: 717–730

898 **Perdomo JA, Conesa MÀ, Medrano H, Ribas-Carbó M, Galmés J** (2014) Effects
899 of long-term individual and combined water and temperature stress on the
900 growth of rice, wheat and maize: relationship with morphological and
901 physiological acclimation. *Physiol Plantarum* doi: 10.1111/ppl.12303

902 **Perdomo JA, Cavanagh AP, Kubien DS, Galmés J** (2015) Temperature
903 dependence of *in vitro* Rubisco kinetics in species of *Flaveria* with different
904 photosynthetic mechanisms. *Photosynth Res* 124: 67–75

905 **Pons TL, Flexas J, von Caemmerer S, Evans JR, Genty B, Ribas-Carbó M,**
906 **Brugnoli E** (2009) Estimating mesophyll conductance to CO₂: methodology,
907 potential errors, and recommendations. *J Exp Bot* doi:10.1093/jxb/erp081

908 **Posada D, Crandall KA** (1998) Modeltest: testing the model of DNA substitution.
909 *Bioinformatics* 14: 817–818

910 **Posada D, Buckley TR** (2004) Model selection and model averaging in
911 phylogenetics: advantages of Akaike information criterion and Bayesian
912 approaches over likelihood ratio tests. *Syst Biol* 53: 793–808

913 **Prins A, Orr DJ, Andralojc PJ, Reynolds MP, Carmo-Silva E, Parry MAJ** (2016)
914 Rubisco catalytic properties of wild and domesticated relatives provide scope
915 for improving wheat photosynthesis. *J Exp Bot* doi: 10.1093/jxb/erv574

916 **R Core Team** (2014) R: A language and environment for statistical computing
917 (version 3.1.1). R Foundation for Statistical Computing, Vienna, Austria.
918 Available at <http://www.R-project.org/>.

919 RCSB Protein Data Bank. Available at
 920 <http://www.rcsb.org/pdb/explore.do?structureId=1RCX>

921 **Rinehart CA, Tingey SV, Andersen WR** (1983) Variability of reaction kinetics for
 922 ribulose-1,5-bisphosphate carboxylase in a barley population. *Plant Physiol*
 923 72: 76–79

924 **Rogers A** (2014) The use and misuse of $V_{c,max}$ in Earth System Models. *Photosynth*
 925 Res 119: 15–29

926 *Rosnow JJ, Evans MA, Kapralov MV, Cousins AB, Edwards GE, Roalson EH* (2015)
 927 Kranz and single-cell forms of C₄ plants in the subfamily Suaedoideae show
 928 kinetic C₄ convergence for PEPC and Rubisco with divergent amino acid
 929 substitutions. *J Exp Bot* 66: 7347–7358.

930 **Roy H, Andrews TJ** (2000) Rubisco: assembly and mechanism. In: Leegood RC,
 931 Sharkey TD, von Caemmerer S, eds. *Photosynthesis: Physiology and*
 932 *Metabolism*. Dordrecht: Kluwer. pp 53–83

933 **Sage RF** (2002) Variation in the k_{cat} of Rubisco in C₃ and C₄ plants and some
 934 implications for photosynthetic performance at high and low temperature. *J*
 935 *Exp Bot* 53: 609–620

936 **Sage RF, Cen Y, Meirong L** (2002) The activation state of Rubisco directly limits
 937 photosynthesis at low CO₂ and low O₂ partial pressures. *Photosynth Res* 71:
 938 241–250.

939 **Sage RF, Way DA, Kubien DS** (2008) Rubisco, Rubisco activase, and global climate
 940 change. *J Exp Bot* 59: 1581–1595

941 **Savir Y, Noorb E, Milob R, Tlustya T** (2010) Cross-species analysis traces
 942 adaptation of Rubisco toward optimality in a low-dimensional landscape. *P*
 943 *Natl Acad Sci USA* 107: 3475–3480

944 **Scafaro AP, Yamori W, Carmo- Silva AE, Salvucci ME, von Caemmerer S,**
 945 **Atwell B J** (2012) Rubisco activity is associated with photosynthetic
 946 thermotolerance in a wild rice (*Oryza meridionalis*). *Physiol Plantarum* 146:
 947 99–109
 948 **Spreitzer RJ, Salvucci ME** (2002) RUBISCO: structure, regulatory interactions, and
 949 possibilities for a better enzyme. *Annu Rev Plant Biol* 53: 449–475.
 950 **Stamatakis A** (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic
 951 analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–
 952 2690
 953 **Tcherkez GG, Farquhar GD, Andrews TJ** (2006) Despite slow catalysis and
 954 confused substrate specificity, all ribulose biphosphate carboxylases may be
 955 nearly perfectly optimized. *P Natl Acad Sci USA* 103: 7246–7251
 956 **Truesdale GA, Downing AL** (1954) Solubility of oxygen in water. *Nature, Lond*
 957 173: 1236
 958 **Uemura K, Miyachi S, Yokota A** (1997) Ribulose-1,5-bisphosphate
 959 carboxylase/oxygenase from thermophilic red algae with a strong specificity
 960 for CO₂ fixation. *Biochem Bioph Res Co* 233: 568–571
 961 **von Caemmerer S** (2000) *Biochemical models of leaf photosynthesis*. Collingwood,
 962 Australia: CSIRO Publishing. ISBN 0 643 06379 X
 963 **von Caemmerer S** (2013) Steady- state models of photosynthesis. *Plant Cell*
 964 *Environ* 36: 1617–1630
 965 **von Caemmerer S, Evans JR** (2015) Temperature responses of mesophyll
 966 conductance differ greatly between species. *Plant Cell Environ* 38: 629–637
 967 **Walker B, Ariza LS, Kaines S, Badger MR, Cousins AB** (2013) Temperature
 968 response of *in vivo* Rubisco kinetics and mesophyll conductance in

969 *Arabidopsis thaliana*: comparisons to *Nicotiana tabacum*. Plant Cell Environ
970 36: 2108–2119

971 **Wang M, Kapralov MV, Anisimova M** (2011) Coevolution of amino acid residues
972 in the key photosynthetic enzyme Rubisco. BMC Evolutionary Biology 11:
973 266

974 **Webb CO, Ackerly DD, Kembel SW** (2008) Phylocom: software for the analysis of
975 phylogenetic community structure and trait evolution. Bioinformatics 24:
976 2098–2100

977 **Weber DJ, Andersen WR, Hess S, Hansen DJ, Gunasekaran M** (1977) Ribulose-
978 1,5-bisphosphate carboxylase from plants adapted to extreme environments.
979 Plant Cell Physiol 18: 693–699

980 **Whitney SM, Sharwood RE** (2008) Construction of a tobacco master line to
981 improve Rubisco engineering in chloroplasts. J Exp Bot 59:1909–1921

982 **Whitney SM, Houtz RL, Alonso H** (2011) Advancing our understanding and
983 capacity to engineer nature's CO₂-sequestering enzyme, Rubisco. Plant
984 Physiol 155: 27–35

985 **Whitney SM, Birch R, Kelso C, Beck JL, Kapralov MV** (2015) Improving
986 recombinant Rubisco biogenesis, plant photosynthesis and growth by
987 coexpressing its ancillary RAF1 chaperone. P Natl Acad Sci USA 112: 3564–
988 3569.

989 **Yamasaki T, Yamakawa T, Yamane Y, Koike H, Satoh K, Katoh S** (2002)
990 Temperature acclimation of photosynthesis and related changes in
991 photosystem II electron transport in winter wheat. Plant Physiol 128: 1087–
992 1097

- 993 **Yamori W, Suzuki K, Noguchi KO, Nakai M, Terashima I** (2006) Effects of
 994 Rubisco kinetics and Rubisco activation state on the temperature dependence
 995 of the photosynthetic rate in spinach leaves from contrasting growth
 996 temperatures. *Plant Cell Environ* 29: 1659–1670
- 997 **Yamori W, von Caemmerer S** (2009) Effect of Rubisco activase deficiency on the
 998 temperature response of CO₂ assimilation rate and Rubisco activation state:
 999 insights from transgenic tobacco with reduced amounts of Rubisco activase.
 1000 *Plant Physiol* 151: 2073–2082
- 1001 **Yamori W, Noguchi K, Hikosaka K, Terashima I** (2010) Phenotypic plasticity in
 1002 photosynthetic temperature acclimation among crop species with different
 1003 cold tolerances. *Plant Physiol* 152: 388–399
- 1004 **Yamori W, Nagai T, Makino A** (2011) The rate- limiting step for CO₂ assimilation
 1005 at different temperatures is influenced by the leaf nitrogen content in several
 1006 C₃ crop species. *Plant Cell Environ* 34: 764–777
- 1007 **Yang Z** (1997) PAML: a program package for phylogenetic analysis by maximum
 1008 likelihood 13: 555–556
- 1009 **Yang Z, Wong WS, Nielsen R** (2005) Bayes empirical Bayes inference of amino
 1010 acid sites under positive selection. *Mol Biol Evol* 22: 1107–1118
- 1011 **Yang Z** (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol*
 1012 *Evol* 24: 1586–1591
- 1013 **Yeoh HH, Badger MR, Watson L** (1980) Variations in Km (CO₂) of ribulose-1,5-
 1014 biphosphate carboxylase among grasses. *Plant Physiol* 66: 1110–1112
- 1015 **Yeoh HH, Badger MR, Watson L** (1981) Variations in kinetic properties of
 1016 ribulose-1,5-biphosphate carboxylases among plants. *Plant Physiol* 67: 1151–
 1017 1155

1018 **Yokota A, Canvin DT** (1985) Ribulose biphosphate carboxylase/oxygenase content
1019 determined with [¹⁴C]carboxypentitol biphosphate in plants and algae. Plant
1020 Physiol 77: 735–739

1021 **Zhu XG, Jensen RG, Bohnert HJ, Wildner GF, Schlitter J** (1998) Dependence of
1022 catalysis and CO₂/O₂ specificity of Rubisco on the carboxy-terminus of the
1023 large subunit at different temperatures. Photosynth Res 57: 71–79

1024 **Zhu XG, Long SP, Ort DR** (2010) Improving photosynthetic efficiency for greater
1025 yield. Annu Rev Plant Biol 61: 235–261

1026