



LJMU Research Online

Nakouti, I, Sihanonth, P, Palaga, T and Hobbs, G

Effect of a siderophore producer on animal cell apoptosis: a possible role as anti-cancer agent.

<http://researchonline.ljmu.ac.uk/id/eprint/4147/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Nakouti, I, Sihanonth, P, Palaga, T and Hobbs, G (2013) Effect of a siderophore producer on animal cell apoptosis: a possible role as anti-cancer agent. International Journal of Pharma Medicine and Biological Sciences. 2 (3). pp. 1-5. ISSN 2278-5221

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



Research Paper

EFFECT OF A SIDEROPHORE PRODUCER ON ANIMAL CELL APOPTOSIS: A POSSIBLE ROLE AS ANTI-CANCER AGENT

Ismiini Nakouti¹, Prakitsin Sihanonth², Tanapat Palaga² and Glyn Hobbs^{1*}

*Corresponding Author: **Glyn Hobbs**, ✉ g.hobbs@ljmu.ac.uk

Iron plays an essential role in the proliferation of aggressive tumors therefore it represents an ideal target for cancer therapy. Cell free supernatants from a siderophore producing actinobacter species previously isolated from Thailand were tested against six human cancer cell lines including malignant melanoma A375 (ATCC no.: CRL-1619), colorectal adenocarcinoma SW620 (ATCC no.: CCL-227), gastric carcinoma Katob! (ATCC no.: HTB-103), liver hepatoblastoma HepG2 (ATCC no.: HB-8065), breast carcinoma BT474 (ATCC no.: HTB-20) and Acute T cell leukemia Jurkat (ATCC no.: CRL-2063). Following treatment of cells with the bacterial culture supernatant the cell viability of A375 cells was dramatically decreased with cell survival of less than 34 % within 48 h. The rest of the cell lines were largely unaffected. Therefore it is suggested that the actinobacterium produced a cytotoxic compound responsible for the cell death by inducing apoptotic activity. We further speculate that this compound was desferioxamine E as the bacterium is known to produce this compound under the culture conditions used.

Keywords: Siderophores, Actinobacteria, Human cell cancer lines, Apoptosis

INTRODUCTION

Cancer cells have an increased requirement for iron compared with 'healthy' cells due to their rapid division. They also possess a higher rate of iron uptake and storage (Elford *et al.*, 1970; Vaughn, 1987). Moreover the ribonucleotide reductase (RR) enzyme involved in DNA synthesis is upregulated in cancerous cells leading to increased DNA replication and cell proliferation (Buss *et al.*, 2003; Dayani *et al.*, 2004,

Richardson, 1997). Iron is fundamental for the activity of RR making iron an obvious anti-cancer candidate (Thelander and Gräslund, 1983; Whitnall *et al.*, 2006).

Siderophores are low molecular weight compounds (600-1500 Daltons) that chelate ferric complexes and other ions with a high affinity. Due to the role of iron in cellular proliferation, iron chelators, such as siderophores, are agents that may be beneficial for the treatment of cancer

¹ School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, Liverpool, L3 3AF, UK.

² Department of Microbiology, Faculty of Science, Chulalongkorn University, Bangkok, 10300, Thailand.

(Wandersman and Delepelaire, 2004). Indeed some data demonstrated that desferrioxamines (iron-free siderophores) can significantly reduce the growth of aggressive tumours in patients suffering from neuroblastom (NB) or leukemia (Buss *et al.*, 2003; Lovejoy and Richardson, 2003). Clinical studies have demonstrated that following a 4 h incubation desferrioxamine can inhibit DNA replication of NB cells (Blatt *et al.*, 1988). In addition a 72 h exposure of NB cells to 60 **Figure 1(a)**M of desferrioxamine reduces the cells' viability by 80 % (Blatt and Stitely, 1987).

An actinobacterium (GenBank accession number EF585403), previously isolated from Tak province in Thailand, was tested positive for siderophore production and its main chelating agent released was identified as desferrioxamine E by HPLC analysis (Nakoutiet *al.*, 2012). The aim of this work was to screen the culture supernatant for anti-cancer activity using a range of cancerous cell lines and to investigate if a possible response mechanism was due to apoptosis.

MATERIALS AND METHODS

Growth Media

Starch casein media: Casein (0.4 g), starch (1.0 g), KNO₃ (0.5 g), K₂HPO₄ (0.2 g), MgPO₄ (0.1 g), CaCO₃ (0.1 g) and bacteriological agar (20 g) were dissolved in 1 litre of distilled H₂O and sterilised by autoclaving at 121°C for 15 min (solid media). Liquid media were prepared by omitting the bacteriological agar.

Culture Conditions

Spores of the strain were stored in a bacterial preservation system (Technical Service Consultants Ltd, Lancashire) at -80°C and 10 μ l were spread onto starch casein agar plates and

incubated at 30° C for a week until sporulation occurred. Fresh spores were collected aseptically (in 20 ml of sterile distilled H₂O) and an aliquot (100 μ l) was introduced into a 250 ml conical flask containing 50 ml of sterile starch casein media (liquid). The cultures were supplemented with 5 g/L lysine prior to incubation on a rotary shaker (200 rpm) at 30° C for 4 days. The conditions were previously optimised for maximum desferrioxamine E production (Nakouti and Hobbs, 2012). Following incubation the medium was centrifuged for 20 min at 8000 g to separate the mycelia. The supernatant was filter-sterilised by using a 0.20 **Figure 1(a)** μ m syringe filter (Corning) and stored at 4°C.

Biological Assay of Anticancer Activity

The effect of the culture supernatant on the cell viability was assessed using the MTT assay (Palaga *et al.*, 1996) against 6 human cancer cell lines including malignant melanoma A375 (ATCC no. CRL-1619), colorectal adenocarcinoma SW620 (ATCC no. CCL-227), gastric carcinoma Katob! (ATCC no. HTB-103), liver hepatoblastoma HepG2 (ATCC no. HB-8065), breast carcinoma BT474 (ATCC no. HTB-20) and Acute T cell leukemia Jurkat (ATCC no. CRL-2063). Cell suspensions in PRMI-1640 medium supplemented with 10% (v/v) fetal bovine serum and 100 mg/ml gentamicin were seeded into flat-bottomed 96-well plates at a density of 2-3 x 10⁴ cells/well and incubated at 37°C in a 5 % CO₂ atmosphere for 24 h. An aliquot (50 μ l) of the culture supernatant was added to the well and incubated (48 h contact time) under the same conditions.

Cell viability was calculated as $(A_{\text{sample}} - A_{\text{blank}}) / (A_{\text{control}} - A_{\text{blank}}) \times 100 \%$ while absorbance (540 nm)

obtained from cells incubated with sterile RPMI-1640 medium was used as the control (positive) and cell free medium as the blank (negative). Ferrichrome (iron-free) from *Ustilago sphaerogena* (5 μ l of 1 mg/ml concentration) was also tested against the cell lines for anti-cancer activity. The reason behind this was that desferrioxamine E was not available in the market; therefore ferrichrome, also a hydroxamate, was purchased from Sigma-Aldrich.

Apoptosis Tests

A cell culture dish (35 x 10 mm) containing a sterile glass coverslip in RPMI-1640 medium was seeded with a cell suspension of A375 cells (final concentration of 1×10^6 cells/ml). The cells were allowed to attach for 24 h, followed by treatment for 48 h. They were then washed with phosphate buffer saline (PBS), fixed with 1% (v/v) glutaraldehyde in the dark for 2 h and washed with PBS. Cells which had attached to the glass coverslips, were stained with DNA dye Hoechst 33342. After being left for 5 min in the dark, apoptotic cell death was determined by observing fragmented nuclei cells under the fluorescent microscope (modified from [14]).

RESULTS AND DISCUSSION

Anticancer Activity

The effect of the strain's supernatant on cellular viability was assessed using the MTT assay. The isolate was found to produce strong cytotoxic substances specific to malignant melanoma cells (A375). Following treatment of cells with the culture supernatant (50 μ l) the percentage cell viability of A375 cells was dramatically decreased with cell survival of less than 34 % (33.193 %) within 48 h (Table 1); whereas the remaining cancer cell lines exhibited similar cell viability. The experiment was an average of 12 replicates. Therefore it was suggested that the organism exhibited selective activity against the skin cancer cell line. Ferrichrome produced by *Ustilago sphaerogena* had little effect on any of the cancer cell lines (cell viability after treatment was measured to be 94 %).

Induction of Apoptosis by *Streptomyces* sp. Strain 23F

The supernatant (50 μ l) of strain 23F (GenBank accession number EF585403) was tested for induction of apoptosis with malignant melanoma cells (A375). Addition of the supernatant to A375 cell culture resulted in the induction of apoptotic

Table 1: MTT Assay Data Demonstrating the Anticancer Activity of the Siderophore Producer Supernatant

| Samples | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | Mean | Viability (%) | STD (%) |
|--------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------------|----------|
| Blank | 0.038 | 0.035 | 0.035 | 0.035 | 0.035 | 0.038 | 0.038 | 0.035 | 0.035 | 0.035 | 0.038 | 0.038 | 0.036 | 0 | N/A |
| Control | 1.178 | 1.226 | 1.293 | 1.190 | 1.198 | 1.227 | 1.210 | 1.293 | 1.193 | 1.224 | 1.194 | 1.290 | 1.226 | 100 | N/A |
| Treated with Supernatant | 0.421 | 0.444 | 0.454 | 0.422 | 0.440 | 0.424 | 0.422 | 0.443 | 0.423 | 0.423 | 0.441 | 0.421 | 0.431 | 33.19 | +/-0.957 |
| Treated with ferrichrome | 1.166 | 1.101 | 1.109 | 1.190 | 1.192 | 1.105 | 1.170 | 1.163 | 1.199 | 1.169 | 1.196 | 1.166 | 1.160 | 94 | +/-2.877 |

activity. Cells from the culture were stained with Hoechst dye and apoptotic nuclei were visualised under the fluorescent microscope (Figure 1). The cancer cells (A375) became smaller through shrinking and the chromatin became agglutinated resulting in nuclear condensation followed by breakdown of the nuclear membrane. The

remaining 'attached' cells appeared very similar to the untreated ones. Cells treated with sterile starch casein media (negative control) did not display any DNA fragmentation and appeared similar to untreated cells (Figure 2). The results suggested that the cytotoxic compound has induced apoptosis in the A375 cell line.

Figure 1: A375 Cancer Cells Stained with Hoechst, the Apoptotic Nuclei were Visualised Under the Fluorescent Microscope (X 40 Magnification)





Figure 2: A375 Cells Treated with Starch Casein Media (Negative Control), Cell Division is Evident (x 40 Magnification)



CONCLUSION

The actinobacterium (GenBank accession number EF585403) released a cytotoxic substance that induced apoptosis to malignant melanoma A375 (ATCC no. CRL-1619).

ACKNOWLEDGMENT

I would like to that the Society For General Microbiology for awarding me the President's Fund. I would also like to thank Prof Prakitsin Sihanonth and Dr. Tanapat Palaga for allowing me to conduct my research in their laboratories.

REFERENCES

- Blatt J, Taylor SR and Stitely S (1988), "Mechanism of antineuroblastoma activity of deferoxamine in vitro", *J. Lab. Clin. Med.*, vol. 112, pp. 433-436.
- Blatt J and Stitely S (1987), "Antineuroblastoma activity of desferoxamine in human cell lines", *Cancer Res.*, Vol. 47, pp. 1749-1750.
- Buss J L, Torti F M and Torti SV (2003), "The role of iron chelation in cancer therapy", *Curr. Med. Chem.*, Vol. 10, pp. 1021-1034.
- Dayani PN, Bishop MC, Black K and Zeltzer PM (2004), "Desferoxamine (DFO)-mediated iron chelation: rationale for a novel approach to therapy for brain cancer", *J. Neurooncol.*, Vol. 67, pp. 367-377.
- Elford HL, Freese M, Passamani E and Morris H P (1970), "Ribonucleotide reductase and cell proliferation", *J. Biol. Chem.*, Vol. 245, pp. 5228-5233.
- Lovejoy D B and Richardson D R (2003), "Iron chelators as anti-neoplastic agents: current developments and promise of the PIH class of chelators", *Cur. Med. Chem.*, Vol. 10, pp. 1035-1049.
- Nakouti I, Sihanonth P and Hobbs G (2012), "A new approach to isolating siderophore producing actinobacteria", *Lett. Appl. Microbiol.*, Vol. 55, pp. 68-72.
- Nakouti I and Hobbs G (2012), "A new approach to studying ion uptake by actinomycetes", *J. Basic. Microbiol.*, in press.
- Palaga T, Kataoka T, Woo J T and Nagai K (1996), "Suppression of apoptotic cell death of IL-3-dependent cell line by ER/ SR Ca²⁺-ATPase inhibitors upon IL-3 deprivation", *Exp. Cell. Res.*, Vol. 228, pp. 92-97.
- Richardson D R (1997), "Iron chelators as effective anti-proliferative agents", *Can. J. Physiol. Pharmacol.*, Vol. 75, pp.164-1180.
- Thelander L and Gräslund A (1983), "Mechanism of inhibition of mammalian ribonucleotide reductase by the iron chelate of 1-formylisoquinoline thiosemicarbazone", *J. Biol. Chem.*, Vol. 258, pp. 4063-4066.
- Vaughn C B, Weinstein R, Bond B, Rice R, Vaughn R W, McKendrick A, Ayad G, Rockwell MA and Rocchio R (1987), "Ferritin content in human cancerous and noncancerous colonic tissue", *Cancer Invest.*, Vol. 5, pp. 7-10.
- Wandersman C and Delepelaire P (2004), "Bacterial iron sources: from siderophores to hemophores", *Annu. Rev. Microbiol.*, Vol. 58, pp. 611-647.
- Whitnall M, Howard J, Ponka P and Richardson D R (2006), "A class of iron chelators with a wide spectrum of potent antitumor activity that overcomes resistance to chemotherapeutics", *Proc. Natl. Acad. Sci.*, vo. 103, pp. 14901-14906.