

Brown dwarfs and low-mass stars in the Pleiades and Praesepe:- Membership and binarity

D. J. Pinfield,¹ P. D. Dobbie,² R. F. Jameson², I. A. Steele¹, H. R. A. Jones¹,
and A. C. Katsiyannis³

1. *Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead, CH41 1LD*

2. *Dept. of Physics & Astronomy, University of Leicester, University Road, Leicester, LE1 7RH*

3. *Dept. of Pure & Applied Physics, The Queen's University of Belfast, BT7 1NN, Northern Ireland*

Accepted 2003 ?; Received 2003 ?; in original form 2002 August

ABSTRACT

We present near infrared J-, H- and K-band photometry and optical spectroscopy of low-mass star and brown dwarf (BD) candidates in the Pleiades and Praesepe open clusters. We flag non-members from their position in K, I-K and J, J-K colour-magnitude diagrams (CMDs), and J-H, H-K two-colour diagrams. In general, the theoretical NextGen isochrones of the Lyon Group fit the K, I-K CMDs well for stars with $I-K \sim 1.5-3.5$. However Pleiades stars with $K \sim 10.5-13$ ($M_K \sim 5-7.5$) are rather redder than the isochrones. We also identify this effect amongst α Per sources from the literature, but find no evidence of it for field stars from the literature. The NextGen isochrones fit the J, J-K CMDs of both clusters very well in this photometric range. It is possible that the I-K colour of youthful stars is affected by the presence of magnetic activity. The Lyon Group's Dusty isochrones fit both K, I-K and K, J-K Pleiades CMDs well for $I-K \sim 4.3-6$ / $J-K \sim 1.1-1.4$. In between these colour ranges the Pleiades cluster sequence comprises three portions. Starting at the bluer side, there is a gap where very few sources are found (the gap size is $\Delta I \sim 0.5$, $\Delta J \sim \Delta K \sim 0.3$), probably resulting from a sharp local drop in the magnitude-mass relation. Then the sequence is quite flat from $I-K \sim 3.5-4$. Finally, the sequence turns over and drops down to join the Dusty isochrone. We also compare model atmosphere colours to the two-colour diagrams of the clusters. The NextGen models are seen to be ~ 0.1 too blue in H-K, and ~ 0.1 too red in J-H for $T_{eff} > 4000$ K. However, they are in reasonable agreement with the data at $T_{eff} \sim 3200$ K. For $T_{eff} \sim 2800-3150$ K, the colour of Pleiades and Praesepe sources are significantly different, where Praesepe sources are ~ 0.1 bluer in J-H and up to ~ 0.1 redder in H-K. These differences could result from gravity sensitive molecular opacities. Cooler Praesepe sources then agree well with the dusty models, suggesting that dust is beginning to form in Praesepe sources around 2500K. However, Pleiades sources remain consistent with the NextGen models (and inconsistent with the dusty models) down to T_{eff} s of ~ 2000 K. It is possible that dust formation does not begin until lower T_{eff} s in sources with lower surface gravities (and hence lower atmospheric pressures). We also identify unresolved binaries in both clusters, and estimate mass ratios (q) for Pleiades BDs. Most of these have $q > 0.7$, however, 3/18 appear to have lower q values. We determine the binary fraction (BF) for numerous mass ranges in each cluster, and find that it is generally rising towards lower masses. We find a BD BF of $50^{+11}_{-10}\%$. We also find some evidence suggesting that the BF- q distribution is flat for $0.5-0.35M_{\odot}$, in contrast to solar type stars.

Key words: stars: low-mass brown dwarfs - open clusters and associations: individual: Pleiades Praesepe

1 INTRODUCTION

Low-mass star and brown dwarf (BD) populations in open clusters make excellent test-beds for our theoretical under-

standing of these objects. With a well constrained cluster age, and assuming uniform composition, we can effectively compare theoretical isochrones to observed colour magnitude diagrams (CMDs) and two-colour diagrams without the complications introduced by a spread in metallicity and a range in surface gravities for a given mass. Furthermore, for most open clusters, the depth effect is small, and the single star sequence is therefore quite tight on CMDs. It is thus possible to study unresolved binarity amongst cluster members.

The T_{eff} range of young to intermediate age low-mass stars and BDs corresponds to M and L spectral types, and modeling the atmospheres of such objects is very complicated. The convection zone penetrates deeply into the optically thin outer layers due to H_2 dissociation (Burrows et al. 1993). Molecular opacities such as TiO and H_2O comprise millions of lines that must be accurately modeled (Allard et al. 1997). Also, dust begins to form in atmospheres at around 2800K (Allard et al. 2001), which depletes opacities as well as affecting the atmospheric temperature gradient. There are many different varieties of dust (Sharp & Huebner 1990), and their formation rates and grain size distributions will depend not only on temperature, but on surface gravity and metallicity. Despite these complexities, much progress has been made in modeling these atmospheres. The Lyon Group has combined structural models (using interior structure physics) with model atmospheres (Hauschildt, Allard & Baron 1999; Allard et al. 2001) to calculate evolutionary models for low-mass stars and BDs, both with and without atmospheric dust condensates (Baraffe et al. 1998, hereafter NextGen; Chabrier et al. 2000, hereafter Dusty). Calculations by other groups include those of Burrows et al. (1997), and D’Antona & Mazzitelli (1994). However, only the Lyon group models are based on consistent non-grey evolutionary calculations relating the mass and age of an object to its observational properties (colour and magnitude).

The Pleiades and Praesepe are two nearby, rich open clusters with near solar composition (Crawford & Perry 1976; Hambly et al. 1995b). With distances of ~ 125 pc and 160pc respectively, both clusters have small depth effects, and are fairly compact on the sky (a few degrees across). The Pleiades is ~ 125 Myrs old (Stauffer, Schultz & Kirkpatrick 1998), and Praesepe is ~ 0.6 – 1.4 Gyrs old (Hambly et al. 1995b). As a consequence, Pleiades low-mass stars ($le 0.3M_{\odot}$) and BDs are not fully contracted, and will have lower surface gravities than their equal T_{eff} counterparts in Praesepe.

In this paper, we derive membership criteria of Pleiades and Praesepe low-mass star and BD candidates using near infrared photometry and optical spectroscopy. We compare NextGen and Dusty theoretical isochrones to our observed CMDs and two-colour diagrams, and identify photometric ranges where the isochrones agree well with observations. We then join these isochrone sections together using additional empirical points, and discuss the overall form of the cluster sequences. We also use our CMDs and two-colour diagrams to identify unresolved binaries. We derive binary fractions over several mass ranges in each cluster, and estimate mass ratios for the Pleiades BD binaries. Finally, we discuss future work.

2 OBSERVATIONS & DATA REDUCTION

2.1 The Pleiades & Praesepe samples

Our Pleiades and Praesepe samples were taken from several sources. For the Pleiades, the highest mass sources ($I < 13.5$) were taken from Hambly, Hawkins & Jameson (1993; HHJ hereafter). Fainter sources were taken from, the ITP survey (Zapatero-Osorio et al. 1999), the CFHT survey (Bouvier et al. 1998), the Burrell Schmidt survey (Pinfield et al. 2000; BPL hereafter) and the INT WFC survey (Dobbie et al. 2002b). The fainter surveys cover a combined area of 7.6 square degrees, and range in photometric depth from $I=19.5$ – 21.8 (unless otherwise stated, I refers to Cousins I). These fainter surveys are summarised in more detail in Jameson et al. (2002).

Our Praesepe sample was taken from four surveys. Proper motion members from the 19 square degree survey of Hambly et al. (1995a, hereafter HSHJ) represent our highest mass sources ($I=10$ – 17.5). Lower mass sources come from Pinfield et al. (1997), Pinfield (1997) and Magazzu et al. (1998). The Pinfield surveys covered 1 and 6 square degrees, and reached $I=21$ and 19.5 respectively. The Magazzu survey covered 800 square arcminutes down to $I=21.2$, and identified 1 BD candidate which they called Roque Pr 1.

2.2 Photometry

For the brighter BPL and HSHJ candidates we obtained J-H- and K-band magnitudes from the 2MASS 2nd incremental point-source catalogue at the NASA/IPAC InfraRed Science Archive (Skrutskie et al. 1995). We used a search radius of 3 arcseconds to allow for positional uncertainties. For the ITP, CFHT and INT WFC surveys, we obtained photometry from the literature for as many sources as was available (see Jameson et al. 2002). Most of the fainter BPL candidates had K-band measurements from Pinfield et al. (2000), and several Praesepe candidates had K-band measurements from Hodgkin et al. (1999). For the remainder of these samples we measured J-, H- and K-band photometry using UFTI on the United Kingdom Infrared Telescope (UKIRT) on Mauna Kea, Hawaii, from 3–5 January 2000, and in service time on 8 October 2001. The sky conditions were photometric. Our observing strategy for candidates without a K-band magnitude was to observe them first at K, and only go on to measure J- and H-band magnitudes if the targets had a sufficiently red I-K colour to be consistent with membership of the cluster. For the remaining candidates, we simply measured J- and H-band magnitudes, as well as K-band if the previous K uncertainty was >0.1 . Each observation comprised a five-point dither pattern, with a windowed 512×512 pixel readout. Total integration times at each of J, H and K were 150s, 300s and 540s, for objects with $I=17$ – 18.5 , $I=18.5$ – 20 and $I>20$ respectively. The raw images were de-biased, dark subtracted, flat-fielded, and combined into mosaics using ORACDR. Aperture photometry was then extracted using GAIA. As the UKIRT photometry was obtained with the “Mauna Kea Observatory” (MKO) filter set (Simons & Tokunaga 2002), the photometry from the literature was transformed into the MKO system using the transforms of Hawarden et al. (2001) and Carpenter (2001) (old UKIRT system and 2MASS respectively). The photometry is given in the appendices. Where there were multiple

Table 1. Spectral indices and equivalent widths of the Pleiades BD candidates.

Source	SpT	PC3	A	VO	I3	I2	H α EW (Å)	KI 7655 EW (Å)	KI 7699 EW (Å)	NaI 8183+8195 EW (Å)
BPL 327	dM7.1	1.55	1.52	2.45	2.33	1.03	-5 \pm 2	-12 \pm 3	-10 \pm 3	-7 \pm 1
BPL 45	dM8.1	1.76	1.69	—	—	—	—	—	—	-7 \pm 2

measurements available of an object, we selected those with small errors (generally trying to avoid measurements with errors larger than ~ 0.1), and took an average.

Our optical photometry was taken from the literature. All I-band photometry was transformed into the Cousins system. Photographic I_N magnitudes from HSHJ were transformed using Bessell (1986). To transform the I_{KP} magnitudes of Pinfield (1997) and Pinfield et al. (2000) we derived the following relation; $I_C - I_{KP} = 0.049(I_{KP} - K) + 0.069$ (for $2.3 < I_C - K < 5.8$). This relation was obtained using synthetic photometry derived from spectra, using the method and data from Dobbie et al. (2002a).

2.3 Spectroscopy

We obtained spectra of 2 BD candidates using ISIS on the 4.2 m William Herschel Telescope (WHT) during 29 Jan 2001. The seeing was ~ 1.2 arcseconds and the weather was clear. Our instrumental setup comprised the TEK4 CCD and the R158R grating on the ISIS red arm, with the mirror in place and no order blocking filters, so as to minimize flux losses (we expect second order contamination for these red objects to be negligible). We used a 1 arcsecond slit, and integrated for 1800s. This gave us a wavelength coverage of 6213-9193Å with a dispersion of 2.91 Å pixel $^{-1}$.

We used IRAF to reduce our spectra. The CCD frames were bias-subtracted and flat-fielded using standard routines, and the spectra extracted and sky-subtracted using the *apall* routine. We calibrated the wavelength scale using the CuAr arc lamp exposures, and the flux using observations of flux standard stars (available in the IRAF environment) on the same night, and with the same instrumental setup.

3 RESULTS & DISCUSSION

3.1 Pleiades spectra

Figure 1 shows the spectra of the two Pleiades BD candidates BPL 327 and BPL 45, where they have been normalised to unity at 7500Å. Table 1 shows the spectroscopic information extracted from these data. Our adopted spectral types were derived using the Pseudo-continuum ratio PC3 from Martín, Rebolo & Zapatero-Osorio (1996), and have uncertainties of ± 0.5 . For BPL 327 the VO index of Kirkpatrick, Henry & Simons (1995) agrees with the PC3 index. Also, the A-index of Kirkpatrick, Henry & McCarthy (1991) implies a slightly earlier spectral type for both sources (dM6.5 and dM7.5 for BPL 327 and BPL 45 respectively), which is what we expect for young sources that are not fully contracted. We also note that the BPL 327 spectra shows that the I2 and I3 indices of Martín & Kun (1996) (which




Figure 1. The 2 Pleiades BD candidate spectra (normalised to unity at 7500Å). The noisier spectra of BPL 45 has been binned by 2 pixels in this plot, and offset for clarity.

measure the CaH and the TiO absorption bands respectively) continue to increase out to \sim dM7, before turning over and decreasing to lower values by dM8. By comparing the I magnitudes and derived spectral types with the spectral type-magnitude diagram of Martín et al. (2000), we found that both BPL 327 and BPL 45 lie on the Pleiades sequence, and appear to be single BDs.

3.2 The IK CMDs

Figures 2 and 3 show the K, I-K CMDs (IK CMDs) for the Pleiades and Praesepe respectively. Small filled circles are higher mass candidate members with 2MASS photometry, and proper motion candidates from HSHJ in the Pleiades and Praesepe respectively. We have several additional fainter Praesepe candidates with 2MASS photometry which are not plotted, since their errors are sizable (see Table A.4). Larger filled symbols are lower mass candidates (candidate BDs in the Pleiades), where circles have J- and H-band photometry (triangles do not). Open circles (small and large) look like background sources in these plots. Open diamonds and open squares are likely old-disk non-members from the K, J-K CMDs (JK CMDs) and the J-H, H-K 2-colour diagrams (JHK diagrams) respectively (see Sections 3.4 and 3.5). Circled objects look like unresolved binaries in these plots (see below). Objects over-plotted with an additional square or triangle are likely unresolved binaries from the JK

© 2003 RAS, MNRAS 000, 1–14




Figure 3. The K, I-K CMD of Praesepe. Symbols are the same as in Figure 2.




Figure 4. The K, I-K CMD of Alpha-Per (filled circles) and field stars (open circles). Sources are taken from Prosser (1992), Barrado y Navascués (2002), and Reid & Cruz (2002).

seem surprising that photospheric bands (and the photometric colours they influence) should be affected by the presence of a chromosphere. However, modeling the correct form of the magnetic field that threads the atmospheres of these stars may allow us to account for these effects.

Below the kink the NextGen isochrone fits both cluster

CMDs well out to $I-K \sim 3.5$, showing the expected decreasing T_{eff} s as degeneracy sets in. For Praesepe, we defined our cluster sequence out to $I-K \sim 3.5$ with the NextGen model isochrone, since it fits the data very well in this range. We chose to do the same for the Pleiades despite the sources being slightly redder than the isochrones for $K=10.5-13$, since we do not expect the reddening to significantly effect the mass-magnitude model predictions.

At redder colours we confine our discussion to the Pleiades, since we only have a few good Praesepe candidates in this range. It can be seen that there is a fairly obvious gap in the Pleiades sequence from $K \sim 14.65-14.95$ (at $I-K \sim 3.5$) where the number of sources is severely reduced. The spectral type corresponding to the gap is M7 (see Figure 4 of Martín et al. 2000), and we refer to this feature as the “M-dwarf gap”. It is discussed in more detail in Dobbie et al. (2002c). However, to summarise, we believe that it results from a sharp local drop in the magnitude-mass relation, possibly caused by the formation of dust in the atmospheres of objects in this T_{eff} regime. We measured the size of the gap at 0.3 in K which compares to ~ 0.5 in I (Dobbie et al. 2002c).

Below the Pleiades M-dwarf gap ($K \sim 14.95-15.3$) the single source candidates follow a fairly flat path in the CMD out to $I-K \sim 4.1$. There is then a turn down, where the candidates drop onto the Dusty isochrone, joining it somewhere from $I-K \sim 4.1-4.5$. We estimated the path followed by the flat part of the sequence by averaging the colours and K-band magnitudes of two groups of sources; the 3 bluest single sources and the 3 reddest single sources from $K=14.95-15.3$ (below which the turn down is obvious). And we estimated the colour where the sequence joins the Dusty isochrone as $I-K \sim 4.3$. The Dusty isochrone then agrees well with the 5 faintest Pleiades candidates out to $I-K \sim 6$, so in this range we used the Dusty isochrones for our sequence.

For Praesepe, we have extended the cluster sequence beyond the NextGen isochrones by assuming it has the same form as the Pleiades sequence; the same M-dwarf gap size, and the same flat portion that turns over and falls onto the Dusty isochrone by $I-K \sim 4.3$. This may not be entirely accurate, but should act as a reasonable approximation.

We have flagged non-members in the Pleiades and Praesepe using our cluster sequences as a guide, and allowing for photometric uncertainties as well as the depth effect of the clusters ($\pm 0.2^{\text{mag}}$ for the Pleiades, and $\pm 0.15^{\text{mag}}$ for Praesepe; Pinfield, Jameson & Hodgkin 1998, Holland et al. 2000). We have also flagged likely unresolved binary members with UKIRT photometry, if they lie significantly ($> 0.25^{\text{mag}}$) above the single star sequence. We duly identified 13 Pleiades binaries, and 8 Praesepe binaries in this way. We refer to them as IK binaries, and they are overplotted with an open circle in the figures. We note that there is a degeneracy between mass and binarity where the IK sequence is dropping down onto the Dusty isochrone ($I-K=3.9-4.5$), and it is not possible to identify all binaries in this region from the IK CMD alone.

3.3 Model J-K colours

In the previous section we found that the NextGen isochrones fit the IK CMDs very well for $I-K \sim 1-3.5$, except for Pleiades sources from $K=10.5-13$, which are slightly red-




Figure 5. The J-K, I-K two-colour diagram of field star sources. Filled circles are young disk, and open circles are old disk. Points have no age information available. 125 Myr and 1 Gyr NextGen and Dusty isochrones are shown as dashed and dotted lines respectively.

der than the isochrones. We also found that Dusty isochrones fit the Pleiades IK CMD well for $I-K=4.3-6$. In this section we test the J-K model predictions for these ranges, for a sample of field stars with kinematics and well measured I-K and J-K colours.

Figure 5 shows J-K plotted against I-K for sources taken from Leggett (1992), Leggett, Allard & Hauschildt (1998) and Dahn et al. (2002). All sources classified as young or old disk (from their kinematics) in the Leggett papers are shown as filled or open circles respectively. In addition, we have assigned 4 L dwarfs as young (<0.5 Gyrs), based on lithium detections (Martín et al. 1999). These are also shown as filled circles. The uncertainties are generally ± 0.05 in colour. We have also overplotted the NextGen and Dusty model isochrones for 120 Myrs (dashed lines) and 1 Gyrs (dotted lines).

Comparison of the two populations shows that for bluer sources ($I-K < 3.5$, $J-K < 1$), the old disk stars have a bigger spread in J-K and on average are slightly bluer in J-K than their young disk counterparts. The two NextGen isochrones essentially lie on top of each other in this range, and fit the data very well. For redder sources ($I-K=4.3-6$, $J-K=1.1-2$), the Dusty isochrones lie on top of each other, and are in excellent agreement with the young disk data. In between these two ranges the isochrones are all too blue in J-K. The NextGen models are too blue because they do not account for dust, and the Dusty models are too blue because they use a less sophisticated treatment of H_2O and TiO molecular opacities. This is all consistent with the gradual change-over from NextGen to Dusty isochrones that was seen in the IK CMDs, and suggests that these models should provide accurate NIR isochrones in the blue and red photometric ranges.




Figure 6. The K, J-K CMD of the Pleiades. Symbols are the same as in Figure 2.




Figure 7. The K, J-K CMD of Praesepe. Symbols are the same as in Figure 2.

3.4 The JK CMDs

Figures 6 and 7 show the JK CMDs for the Pleiades and Praesepe respectively. Only good candidates from the IK CMDs are shown. The symbols mean the same as in Figure 2. The location of several magnitude bins are indicated with dotted lines (see Section 3.5). Isochrones are overplotted as dashed lines, with mass points shown in units of $0.001 M_{\odot}$. Typical photometric errors for UKIRT photometry are indicated.

We constructed cluster sequences for the JK CMDs based on our analysis in Sections 3.2 and 3.3. We used

NextGen isochrones for $I-K < 3.5$ ($\text{mass} < 0.07$ and $0.09 M_{\odot}$ for the Pleiades and Praesepe respectively), and Dusty isochrones for $J-K=1.1-2$ ($I-K=4.3-6$). In between the NextGen and Dusty isochrones we added two additional sequence points, based on our Pleiades sources; one to bridge the M dwarf gap, and the other to represent the turn down point, where the sequences begins to fall onto the Dusty isochrone (see Section 3.2). An average of the J-K colours of the Pleiades sources above and below the gap shows that there is essentially no significant J-K colour change across the gap ($\Delta J-K=0.01$). Averaging of the 3 single Pleiades sources in the turn down region gave a turn down colour of $J-K=1.03$. The flat region seen in the IK CMDs is thus not very apparent in the JK CMDs. For Praesepe we assumed that the JK colour and magnitude changes are the same as in the Pleiades (as we did previously for the IK CMDs). The cluster sequences we constructed are shown as solid lines in the JK CMDs, and a dotted line indicates the spread we expect in the sequence from unresolved binarity.

The NetGen part of the sequence fits the observations very well in both clusters, and there is certainly no evidence for the Pleiades sources being redder than the isochrones from $K=10.5-13$, as was seen in the IK CMDs. The J-K colour is clearly not affected in the way that the I-K colour is. Below the M-dwarf gap, it can be seen that there is a spread in the Pleiades sources, where several are significantly redder than expected for single objects. A spread is also seen for a small number of Praesepe sources. Such a spread could result from several possibilities. Transient surface features such as spots could be responsible. The 2 most obvious red sources (Teide 1 and Calar 3; BPL 137 and 235) have been tested for variability. Bailer-Jones & Mundt (2001) found no evidence of I-band variability in Teide 1 over a 100 hour time scale, and only marginal evidence (just above the noise) for sporadic variability ($\Delta I < 0.03$) in Calar 3 over a 29 hour time scale. Our photometry of Teide 1 comprises 3 well separated ($> a$ year) J- and H-band measurements which are consistent to within their errors. For Calar 3 we have 1 K-band measurement which is consistent with the measurement presented by Martín et al. (2000), and 2 H-band measurements which agree to within their errors. We also have 3 J-band measurements which show marginal evidence (at the 2σ level) of variability. The level of this variation is not in itself sufficient to account for the redness of Calar 3. Photometric errors will also cause some spread in the sources. However, the general consistency (at the $1-\sigma$ level) between our multiple measurements makes it unlikely that these alone are responsible. The third option, and the one we chose to consider in more detail, is that additional unresolved binaries (not identified in the IK CMD) are present. A substantially redder unresolved companion could redden a source sufficiently to account for the spread. These binaries would either be background sources, or cluster binaries that come from the turn down region of the IK CMD, where binarity is not always apparent. We have thus flagged 4 additional candidate unresolved binaries in the Pleiades, and 2 in Praesepe. We refer to these as JK binaries, and they are overplotted with an additional open square in the figures. We note that the bluest of the Pleiades JK binaries is a known background binary (with out lithium; Martín et al. 2000), resolved by the HST (Martín et al. 1998). The Pleiades binaries will be discussed in more detail in Section

3.7. Of the 2 Praesepe JK binaries, the bluer one (RIZ Pr 23) is located just above where we expect the missing M dwarf gap to be, and could be a binary from below the gap. The redder one (RIZ Pr 23) comes from the turn down region of the IK CMD.

We have flagged non-members in the Pleiades and Praesepe using our cluster sequences as a guide. Objects that look too blue are presumably background stars from an old-disk population (see Section 3.3), and we flagged them as such if their J-K colours were at least $2-\sigma$ blueward of the cluster sequence.

3.5 The J-H, H-K 2-colour diagrams

Figures 8 and 9 show the J-H, H-K 2-colour diagrams (JHK diagrams) for the Pleiades and Praesepe respectively. Only good candidates from the IK CMDs are shown. The source symbols mean the same as in Figure 2. The positions of 4 field L-dwarfs (from Dahn et al. 2002) are also shown as asterisks. The photometric error bar shown in the lower right of the plots are typical of the photometry of the BPL sources. The fainter sources have slightly larger errors. Solar metallicity model atmosphere predictions (used for the NextGen and Dusty models) are overplotted as dashed and solid lines respectively, with labels indicating surface gravities. For the Pleiades we plot NextGen models for $T_{\text{eff}} > 2000$ and 3000K , and Dusty models for $T_{\text{eff}} < 2000\text{K}$. For Praesepe, we plot NextGen models for $T_{\text{eff}} > 2500$ and 3000K , and Dusty models for $T_{\text{eff}} < 2500\text{K}$.

For the highest mass stars, it can be seen that the NextGen models are ~ 0.1 too blue in H-K up to the 2-colour turn-over, where convection begins ($\sim 4000\text{K}$). The data turns over at $H-K=0.27$, $J-H=0.57$, and $\log g$ for such objects should be ~ 4.5 . The theoretical J-H colour of the turn over is therefore ~ 0.1 too red. These discrepancies are thought to be caused by inaccuracies in the modeling of water vapour opacities in the H-band (with its overall low opacity) as well as an underestimation of the mixing length parameter. After the turn-over, the 2MASS photometry follows a short path in the JHK diagram, reaching $J-H=0.52$, $H-K=0.38$ by $\sim 3200\text{K}$, in reasonable agreement with the $\log g=5$ NextGen model. In order to look for any non-evolutionary differences between the colours of Pleiades and Praesepe stellar members, we selected Pleiades members from $K=11.5-12$, and Praesepe members from $K=12-12.5$. These two sets of sources should be fully contracted (according to the models), and should represent the $T_{\text{eff}} \sim 3500-3600\text{K}$ range (since we have corrected for distance). We see no significant difference in the colours, with $J-H=0.53 \pm 0.03$ and 0.54 ± 0.03 , and $H-K=0.35 \pm 0.03$ and 0.33 ± 0.04 for the Pleiades and Praesepe respectively. Praesepe is known to be slightly metal rich compared to the Pleiades, with $[\text{Fe}/\text{H}]=0.04 \pm 0.04-0.14$ (Friel & Boesgaard 1992; Reglero & Fabregat 1991). However, this does not appear to significantly affect the stellar NIR colours, consistent with the model predictions in this T_{eff} range.

In order to show what is happening at cooler T_{eff} s, we grouped the sources into a number of K magnitude bins. Above the gap we selected Pleiades bins of $K=13.6-14.1$ and $K=14.1-14.65$, and Praesepe bins of $K=14.5-15.3$ and $K=15.3-15.85$. We labeled these bins A and B respectively, and they should correspond to approximately the same T_{eff}




Figure 8. The J-H, H-K 2-colour diagram of the Pleiades. Symbols are the same as in Figure 2. Individual sources are labeled in italic (R =Roque).




Figure 9. The J-H, H-K 2-colour diagram of Praesepe. Symbols are the same as in Figure 2.

ranges in each cluster (~ 3150 – 3000 K for A, and 3000 – 2800 K for B). Below the gap, we considered 3 additional Pleiades bins. The $K=14.9$ – 15.13 range (bin C) contains objects on the flat section of the IK CMD (see Section 3.2). The $K=15.13$ – 15.5 range (bin D) contains objects at the top of the turn down, where the sequence begins to drop towards the Dusty isochrone in the IK CMD. Finally, the $K=15.5$ – 16.1 range (bin E) contains objects from near the base of the turn down region, where the IK sequence is joining onto the Dusty isochrone. These bins are indicated on the IK and JK CMDs with dotted lines, and in the JHK diagrams by

boxes drawn with dotted lines. In the Pleiades, bin A contains 2MASS sources, and we defined the box dimensions using the standard deviation of the average bin colours. For the remaining bins we defined the box dimensions by attempting to include all non-binary member sources within the box dimensions.

When we compare the colour-colour location of bins A and B in each of the clusters, it can be seen that they differ. In the Pleiades, bins A and B have similar colours, and lie very close to the NextGen model predictions for their T_{eff} range. However, in Praesepe these bins have J-H colours that are ~ 0.1 bluer than this, and bin B has an H-K colour ~ 0.1 redder. The surface gravities expected for these bins are $\log g=5$ for the Pleiades, and $\log g=5.15$ – 5.25 for Praesepe, and these differing surface gravities may provide the explanation for the differing colours. We do not expect any atmospheric dust formation for objects in these bins since their T_{eff} s are too high (see Tsuji 2002 figure 7, and Jones & Tsuji 1997), so gravity sensitive dust formation cannot provide the explanation. However, molecular opacities (such as H_2O) could be more sensitive to gravity (or pressure) than the NextGen models predict and future generations of models with updated EOS and opacity data may account for these colour changes.

Moving to fainter cooler objects in Praesepe, it can be seen that the faint binary candidate below bin B lies on the dusty isochrone, consistent with dust formation beginning in Praesepe at around $T_{eff} \sim 2500$ K. However, in the Pleiades we are faced with a rather different story. Moving across the gap from bin B to C produces no significant change in colour, and between bins C and D there is then an increase of ~ 0.1 in J-H. These changes are all consistent with the NextGen isochrone. Pleiades sources eventually move onto the dusty isochrone between bins D and E, and the 2 faintest Pleiades sources (Roque 25 and 30) both lie on the Dusty isochrones. A possible explanation for these near infrared colours is that dust does not begin to form in Pleiades sources until $T_{eff} \sim 2000$ K (between bins D and E). This might occur because in lower gravity atmospheres the lower pressures could suppress dust formation until lower T_{eff} s. However, one would also expect that lower gravity atmospheres would be more extended, and have lower outer temperatures, which would have the opposite effect.

Assuming that the differences in colour between Pleiades and Praesepe sources result from differing surface gravities, then one might expect even greater differences for sources found in star forming regions. This would have implications for the derreddening process that is commonly used for such sources (eg. Lucas & Roche 2000).

We have flagged likely non-members using the location of bins A to E and the Dusty model atmosphere predictions (for the faintest sources) as a guide. Old disk sources appeared too blue in J-H, and several non-stellar sources were also identified amongst our data by their extremely red colours. One object in Figure 7 (Roque 33) appears to be too blue in J-H. This object is a proper motion member however (see Table A.1), and we note that its photometric errors are ~ 0.2 in colour making them not inconsistent with membership. We also used the Pleiades JHK diagram to identify some additional unresolved binaries. In the turn down region of both the IK and the JK CMDs (where the sequence is dropping steeply down towards the Dusty isochrones), there

remains some degeneracy between mass and binarity. However, we were able to break this degeneracy using our JHK diagram. A single source with $K=15.13-15.5$ should lie in bin D. However, 3 sources in this magnitude range lie in (or very near to) bin E. These are presumably near equal mass binaries, where both components of each binary come from bin E (we refer to them as JHK binaries). These binaries are overplotted with an additional open triangle in the figures.

3.6 Summary of membership criteria

Using the IK and JK CMDs, as well as the JHK diagrams, we have assigned photometric membership criteria to our Pleiades and Praesepe samples (see Appendices). In the Pleiades, we flagged 5 of the 30 BPL BD candidates as non-members. The remaining 25 BPL candidates, and the 9 additional Pleiades candidates from the literature look like cluster BDs. Examination of Table A1 shows that our criteria for these sources are consistent with existing spectroscopic and astrometric measurements, where they exist. It is particularly encouraging that we were able to flag CFHT-PL-18 (BPL 283) as a background binary from its photometry alone. This source was a suspected BD binary, and was only confirmed as a background source when Martín et al. (2000) failed to detect lithium in its spectrum. BPL 45 has been flagged as the same type of object based on its measured J- and K-band photometry alone. Of the Pleiades low-mass star candidates, we flag 14 non-members out of the 194 candidates. This represents only 7% spurious sources in the sample. In Praesepe, we identify higher levels of contamination for our faintest sources. We flag 20 non-members out of the 26 RIZ-Pr candidates, and 37 non-members out of the 89 IZ-Pr candidates. However, for the brighter HSHJ sources we only flag 21 non-members out of 458. The position of Roque-Pr 1 in the IK and JK CMDs is consistent with membership. However, if it is a member, it must be an unresolved BD binary.

3.7 BD binary mass-ratios

Figure 10 shows two regions of the Pleiades IK and JK CMDs, focusing on the candidate BDs. The symbols mean the same as in Figure 2. NextGen and Dusty isochrones are shown with dashed lines, and our cluster sequences (see Sections 3.2 and 3.4) are shown as solid lines. We also show the effects of unresolved binarity, with the use of binary tracks (shown as dotted lines). To construct these binary tracks we started with the single star point which is to form the primary. Then, by combining the magnitude of this primary with that of the faintest point on the sequence (lowest mass), we obtained a new binary point which we joined to our primary point by a dotted line. We then repeated this process for higher mass secondaries, joining each new point to its predecessor until we reach an equal mass binary, 0.75 magnitudes above the original primary. We also show 5 mass points on the CMDs (in units of $0.001M_{\odot}$). The $0.07M_{\odot}$ point just above the M dwarf gap is the NextGen model prediction for a 125Myr isochrone. Since we believe that the M dwarf gap results from a sharp drop in the luminosity-mass relation, we assumed that the mass just below the M dwarf gap is very similar to the mass just above it. Hence, we

have a $0.07M_{\odot}$ point just below the M dwarf gap also. The $0.06M_{\odot}$ mass point was estimated from the Pleiades mass-I relation of Dobbie et al. (2002c), which accounts for the M dwarf gap, and was derived using the Pleiades I-band BD LF, and an assumed power-law MF. We note that this agrees with the NextGen and Dusty model J-band predictions, if they are offset by the J magnitude width of the M dwarf gap. The final two mass points are Dusty model predictions for a 125Myr isochrone. The $0.04M_{\odot}$ mass point should be reasonably representative of the point where the sequence joins the Dusty isochrone. We also note that Dobbie et al. (2002c) suggest that model predictions for the masses of Pleiades BDs from $I=18.5-19.5$ are too low (by up to $0.01M_{\odot}$), which may have some bearing on the Dusty mass estimates.

When analysing these CMDs, we must consider the depth effect of the cluster. Sources near the front and back of the cluster will appear slightly brighter and fainter respectively, than they would in the cluster centre. The tidal radius of the Pleiades is 13.1pc (Pinfield, Jameson & Hodgkin 1998) giving a full depth of $\pm 0.2^{mag}$. However, the majority of sources will be found closer to the cluster centre. Jameson et al. (2002) fit a King profile (King 1962) to the spatial distribution of Pleiades BDs, and derived a core radius of 5pc. We would expect $\sim 80\%$ of members to lie within this core radius, giving a more likely depth of $\pm 0.1^{mag}$. In order to estimate binary mass-ratios (q) for our unresolved binary candidates, we compared their positions to the binary tracks in both CMDs, and estimated q ranges consistent with the photometric errors and the depth effect of the Pleiades. We did this for a depth effect of ± 0.1 and ± 0.2 . We also estimated q ranges on the basis that the Dusty isochrone masses may be higher by $0.01 M_{\odot}$. And finally, we made a small allowance for the known (or suspected) variables (PPL1 and Calar 3). Our results are summarised in Table 2. Column 3 indicates if the source is an IK, a JK or a JHK binary (identified from the IK CMD, the JK CMD or the JHK diagram). Column 4 gives the estimated q range from the mass points in Figure 9 and a ± 0.1 depth effect. Column 5 gives a larger q range, estimated with the full ± 0.2 depth effect, and accounting for the uncertainty in the Dusty isochrone mass points (assuming they may be $0.01M_{\odot}$ too low).

Where a source appeared in both CMDs, we found q ranges that were consistent (to within photometric errors) for all but 2 candidates; CFHT-PL-18 (BPL 283) and BPL 45 appear red in the JK CMD, but do not appear sufficiently red in the IK CMD. The component sources of these binaries would therefore be too blue in I-K to be cluster members. This is confirmed for CFHT-PL-18, which is a known background binary (see Section 3.4). We have also flagged BPL 45 as a probable non-member, based on its measured photometry alone. CFHT-PL-12 (BPL 294) and INT-PL-IZ-42 could have primaries that are either just above or just below the M dwarf gap. However, CFHT-PL-12 would need to be near the front of the cluster if its primary was from below the gap. Roque 4 lies quite high in the JK CMD suggesting that it may be a foreground object. However, it could still be a binary BD if it is at the front of the cluster, and may alternatively be a BD triple system. We were unable to estimate good q ranges for the 2 faintest binaries, since we do not know the form of the cluster sequence for masses below $0.03M_{\odot}$.

It is clear that the majority of the unresolved binary




Figure 10. K, I-K and K, J-K CMDs of Pleiades BDs.

Table 2. Pleiades BD binary mass-ratios (q). The q_1 ranges were estimated assuming a depth effect of $\pm 0.1^{mag}$, and masses from Figure 9. q_2 ranges were estimated assuming a depth effect of $\pm 0.2^{mag}$, and accounting for the fact that the Dusty isochrone masses could be underestimated by $\sim 0.01 M_\odot$.

Binaries	Other names	Binary type	q_1	q_2	Notes
BPL 45		JK			Probable BG source
BPL 66	Roque 4	IK		~ 1	Near front of cluster?/Triple BD?
BPL 79	Roque 13	IK	0.7-1	0.65-1	
BPL 108	Roque 14	IK	0.8-1	0.75-1	
BPL 137	Teide 1	JK	0.55-0.65	0.5-0.9	
BPL 142	Roque 17	IK	0.7-1	0.65-1	
BPL 235	Calar 3	JK	~ 0.5	0.5-0.7	
BPL 249		JHK	~ 1		
BPL 283	CFHT-PL-18	JK			Known BG source
BPL 294	CFHT-PL-12	IK	0.6-0.9	0.4-1	Primary probably above gap
BPL 303	CFHT-PL-25	JHK	< 0.75	< 1	
PPL 1	Roque 15	IK	0.8-1	0.75-1	
PPL 15	IPMBD 23, NPL35	IK	0.7-1	0.65-1	Known binary with $q=0.85$
CFHT-PL-16		IK	0.8-1	0.75-1	
CFHT-PL-23		JHK	~ 1		
INT-PL-IZ-20		IK		< 0.8	
INT-PL-IZ-25		IK	~ 0.8		
INT-PL-IZ-42		IK	0.4-0.6 or ~ 1	0-0.7 or ~ 1	Primary above or below gap?
INT-PL-IZ-43		IK	0.4-0.65	0-0.8	
INT-PL-IZ-76		IK	0.9-1	0.8-1	

candidates have fairly high q values. Indeed, 15 of the 18 probable binary BDs appear to have $q > 0.7$. However, the photometry of 3 sources is inconsistent with them being either single or $q > 0.7$ binaries. One of these is without J- or H-band measurements, and has not been confirmed as a cluster member (by either a proper motion measurement or the lithium test). The other 2 are Teide 1 and Calar 3, which are bona fide cluster BDs. We have previously discussed (Section 3.4) other possible causes for the red J-K colours seen in these sources, and concluded that binarity was the most likely. We note in addition that the position of these sources in the JHK diagram is also consistent with binarity, since

they appear partway between bins D/E and the location of Roque 25. These few objects therefore suggest that BD binaries are not all high q systems. They could consist of two populations; the majority in high q systems, and a smaller fraction in low q systems. Theoretical simulations are not yet at a level to make such predictions. However, the hydrodynamical simulation of Bate, Bonnell & Bromm (2002) shows BDs forming from the collapse and fragmentation of a turbulent molecular cloud. They determined that $\sim 75\%$ of BDs form by the fragmentation of gravitationally unstable circum-stellar disks, with subsequent ejection removing the BD from the region of dense gas. The remaining 25%

Table 3. Binary fractions for the Praesepe and Pleiades samples. Spectral types were estimated using Leggett (1992).

Praesepe			
I-K	Mass/SpT	q-sensitivity	Binary fraction
1–1.9	1.0–0.6M _⊙ G/K stars	0.5–1	17 ⁺⁶ _{–4} %
1.9–2.5	0.6–~0.2M _⊙ M0–M5	0.35–1	41±5%
1.9–2.2	0.6–~0.35M _⊙ M0–M3	0.3–1	31 ⁺⁷ _{–6} %
2.2–2.5	~0.35–~0.2M _⊙ M3–M5	0.4–1	44±6%
3–3.6	~0.11–~0.09M _⊙ M6–M7	0.65–1	47 ⁺¹³ _{–11} %
Pleiades			
1–1.9	1.0–0.6M _⊙ G/K stars	0.5–1	23 ⁺⁶ _{–5} %
2.6–3	0.15–0.09M _⊙ M5–M6	~0.35–1	36±5%
2.6–2.8	0.15–0.11M _⊙ M5	0.35–1	27 ⁺⁶ _{–5} %
2.8–3	0.11–0.09M _⊙ M6	0.4–1	51±8%
3.3–4.3	0.07–0.06M _⊙ M7–M8	0.5–1	50 ⁺¹¹ _{–10} %

were found to form from the fragmentation of dynamically collapsing filamentary gas, where the forming BDs fell into, and were rapidly ejected from unstable multiple systems (see also Reipurth & Clarke 2001). However, this simulation only produced 1 BD binary, but lacked the spatial resolution to predict very close systems. Future simulations with higher spatial resolution may reveal a link between the two different formation mechanisms and the resulting *q* distribution of BD binaries.

3.8 Binary fractions

We determined binary fractions (BF; defined as the number of binary systems in some photometric range divided by the total number of systems in that range) using the IK CMDs in general, for a number of photometric ranges. The results are given in Table 3, where the uncertainties are determined assuming binomial statistics (see Burgasser et al. 2002). The mass ranges were estimated using our IK cluster sequences (see Section 3.2). The stellar *q* ranges were estimated by simulating the effect of unresolved binarity in the IK CMDs (as was done in Section 3.7 for the Pleiades BDs), and finding the *q* value when the binary track crosses midway between the single star sequence and the binary sequence 0.75 magnitudes above it.

Starting in Praesepe, the first range we considered was from I-K=1–1.9 (G and K stars). The single stars all bunch around the predicted sequence, and we flagged binaries by their proximity to the binary star sequence. The next range we chose was from I-K=1.9–2.5 (where the upper limit represents the completeness of 2MASS and the HSHJ survey). The sources lie in a fairly straight path down the CMD in this range, and we therefore chose to separate binaries and

single stars using a straight line. We defined this separation line such that the average spread of members above and below it was the same (as one would expect for a binary induced spread). And by doing this for two sub-ranges (I-K=1.9–2.2 and 2.2–2.5) we ensured that the gradient of the line was correct. We would expect a spread of ± 0.375 to represent the full spread induced from binarity, and we obtained spreads of ± 0.36 and ± 0.43 for the two ranges respectively. The slightly larger value for the redder range is accounted for by larger photometric errors. This approach has the added benefit that it is independent of any assumed distance modulus. Table 3 gives the binary fraction for both sub-ranges, as well as for the full range. For the lowest-mass Praesepe stars we chose to use I-K=3–3.6 (where we just have UKIRT photometry). This avoids the CMD overlap region between the photographic and CCD surveys, where different survey areas and photometric errors complicate matters. Binaries in this range were flagged in Section 3.2. Unlike the higher mass ranges, these candidates do not have proper motions. However, using the disk luminosity function of Kirkpatrick et al. (1994), we expect no more than ~ 1 and ~ 0.5 field star to contaminate the single and binary star sequences respectively. Such numbers will not significantly affect our statistics.

For the Pleiades, the first range we considered was from I-K=2.6–3. We took the same approach as for Praesepe, and divided this range in two (I-K=2.6–2.8 and 2.8–3), defining a selection line such that the scatter above and below (in each region) was identical. The spread about the selection line was ± 0.57 for each of the sub-regions. This is somewhat more than the expected ± 0.375 , but the additional spread will be predominantly due to photometric errors and cluster depth. Finally, we considered a BD range. We counted single BDs below the M dwarf gap that have $K < 15.5$, since our sample is spatially complete in this range. The mass range for the single stars is ~ 0.07 – 0.05 M_⊙, where we have interpolated between the 0.06 and the 0.04M_⊙ points in Figure 9. We then counted binary BDs that could have primaries amongst these single sources, including objects from $K=14.2$ – 15.2 . We note that we ignored CFHT-PL-12 (BPL 294) since it is likely that its primary comes from above the gap. In this way, we selected 10 single BDs, and 10 binary BDs. We estimated the *q* sensitivity range for our BD binaries using Table 2.

We have already discussed how the cluster depth affects BD binaries. However, it will also be a source of error for our stellar BFs. Objects whose position on the CMD is within the cluster depth (eg. 0.2 and 0.15 magnitudes for the Pleiades and Praesepe respectively) of the binary selection line could potentially spill over this line if they were near the front or the back of the cluster. This represents 53% of Pleiades sources, and 40% of Praesepe sources. However, the cluster spatial distributions (Pinfield, Jameson & Hodgkin 1998; Holland et al. 2000) indicate that of these stars, we would only expect ~ 5 – 15% and ~ 5 – 10% of Pleiades and Praesepe stars respectively to actually spill across the line. Further more, an error will only occur if the BF itself is different from 50% (for BF=50% misidentified binaries and single stars will cancel out). Combining these together, we would expect no more than $\sim 1\%$ and 2% BF errors in Praesepe and the Pleiades respectively, due to depth effects.

Another source of error comes from variability, resulting




Figure 11. The binary fractions of the Pleiades (triangles) and Praesepe (Circles). Three points from the literature are shown as open diamonds. At $0.1M_{\odot}$ the BF error bars of two points overlap, so we plot the Praesepe error bars as a dotted line for clarity. q ranges are indicated next to each point, and an A range for the Burgasser and Close points.

from chromospheric activity such as spots or flares pushing some single stars intermittently above the binary selection line. Lockwood et al. (1984) found that $\sim 40\%$ of Hyades F G and K dwarfs varied at a level of $\sim 1-2\%$. For later types, Stauffer et al. (1991) reported H_{α} emission in $\sim 60\%$ of late K–mid M Hyades members, as well as in the vast majority of G–mid M Pleiades members, and Scholz, Eisloffel & Mundt (2000) found photometric variations among young M-dwarfs similar to those of solar type stars. Therefore, although many of our members may be variable, we expect this variability to be generally low (a few %). More violent flares are occasionally seen in late M dwarfs (Leibert et al. 1999), but are rare. We estimate that our stellar BF values should not be increased by more than $\sim 5\%$ as a result of variability.

The BF values are plotted in Figure 11, as a function of mass. Filled triangles and circles represent Pleiades and Praesepe data respectively. Additional points from the literature are shown as open diamonds. The highest mass point agrees well with the field star results of Duquennoy & Mayor (1991), which increases to 70% for $q=0.1-1$. From $0.6-0.35M_{\odot}$ we find a BF of $\sim 30\%$. The BF of NGC 2516 agrees with this value for $\sim 0.6M_{\odot}$, but drops to $\sim 15\%$ for $0.5-0.35M_{\odot}$ (Jeffries, Thurston & Hambly 2001). This difference could be explained by the fact that the NGC 2516 CMD was only sensitive to $q=0.6-1$ binaries, whereas our CMD is sensitive to $q=0.3-1$. If the clusters have intrinsically similar binary populations, then this implies that there is a BF of 15% from $q=0.3-0.6$ as well as from $q=0.6-1$, suggesting a flat BF– q distribution in this mass range. This represents a departure from the BF– q distribution of solar type field stars, which rises towards lower q . If the BF– q distribution were like that of solar type stars we would ex-

pect a BF of 30% for $q=0.3-0.6$, and thus 45% for $q=0.3-1$. Our result is 1.5σ lower than this value. A flat binary– q distribution would give a total BF of $\sim 45\%$ in this mass range. Next, with the exception of the Pleiades point at $0.13M_{\odot}$, there is a general trend of increasing BF towards lower mass. The q ranges of these points are fairly similar, and so should not significantly bias this trend. Moving into the sub-stellar regime, it can be seen that the BD binary fraction is around 50% . Even if we are extremely conservative, and only select the most obvious BD binaries from the IK CMD (assuming that JK and JHK binaries are in fact single), we find a BF of $32^{+11}_{-8}\%$. Figure 11 also shows the T dwarf BF resolved by the HST (Burgasser et al. 2003), and the late M/early L dwarf BF resolved by Hokupa’a on Gemini (Close et al. 2003). Mass range estimates were taken from Close et al. (2003), and estimated for the T dwarfs assuming an age range of 1–5 Gyrs. Both these BFs ($9^{+15}_{-4}\%$ and $15\pm 7\%$ respectively) are significantly lower than our BF for Pleiades BDs. However, these observations were only sensitive to binaries with separations $>1\text{AU}$. It is possible that the total BF of these populations is identical to that of Pleiades BDs if $70-80\%$ of BD binaries have separations $<1\text{AU}$. It may be that many BD binaries are like PPL15 (Basri & Martín, 1999) which has a separation of 0.03AU . We also note that the N-body simulations of Kroupa (2001), and Kroupa, Aarseth & Hurley (2001) suggest that $\sim 50\%$ of cluster BD binaries with $\log P/\text{days}=1-6$ (separations of $\sim 0.04-90\text{AU}$) would be disrupted during the first few Myrs of proto-cluster dynamical evolution. This would require an even higher initial BD BF unless most BD binaries are extremely tight. However, the hydrodynamical simulation of Bate, Bonnell & Bromm (2002) predicts an initial BF of $\sim 5\%$, since only 1 BD binary was produced in their simulation among 20 single BDs. But the spatial resolution of this simulation was not high enough to model the formation of extremely close BD binaries. Such binaries may form during a secondary collapse phase of individual pressure supported fragments (Bonnell & Bate 1994). If this is not the case (Bate 1998), then post fragmentation processes (such as dynamical interactions) might be responsible for creating extremely tight BD binaries (by hardening).

4 FUTURE WORK

Proper motion measurements have already been made for 15 of the BD candidates discussed here. Work is currently underway by some of us to measure proper motions of the remaining Pleiades BDs. It will be particularly useful to confirm all possible low- q BD binaries as Pleiades members. Measuring proper motions of the Praesepe candidates is also important, and is in progress (Chappelle, Pinfield & Steele 2002). Such measurements will allow us to define the Praesepe cluster sequence with more confidence. Also, the faintest Praesepe candidates could be BDs, and would be the first identified in an intermediate age cluster.

Measuring NIR spectra is a natural next step if we are to understand the nature of the cluster sequences. These spectra could be compared to the predictions of both NextGen and Dusty model atmospheres, to determine when dust is beginning to form in the atmospheres of Pleiades and Praesepe sources, and what effects it has in the different surface

gravities. Furthermore, the best way to improve on our analysis of binarity is from NIR spectra. We could combine the spectra of likely single sources, and compare the results with spectra of suspected unresolved binaries, to confirm or not, if these sources really are binary in nature.

The nature of the M-dwarf gap could be further investigated by measuring longer wavelength photometry (eg. L-band) of sources on either side of the M dwarf gap, to look for evidence (or lack of it) of dust formation.

Deeper and larger scale surveys of the clusters will provide much larger samples of members to work with. The 16 square degree survey recently obtained as part of the WFS with the INT and WFC should identify several hundred Pleiades BDs. Also, the WFC survey of Praesepe made by Chappelle, Pinfield & Steele (2002) should identify large numbers of very low-mass stars and possibly BDs. With these larger samples we will be able to more accurately measure the M dwarf gap in both clusters, as well as obtain a better statistical measure of the BD binary fraction and binary-q distribution.

ACKNOWLEDGMENTS

DJP and PDD acknowledge funding from PPARC. AK acknowledges funding from the Leverhulme Trust. The United Kingdom Infrared Telescope is operated by the Joint Astronomy Centre on behalf of the U.K. Particle Physics and Astronomy Research Council. Some of the data reported here were obtained as part of the UKIRT Service Program. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the Aeronautics and Space Administration and the National Science Foundation.

REFERENCES

- Allard F., Hauschildt P. H., Alexander D. R., Starrfield S., 1997, *ARA& A*, 35, 137
- Allard F., Hauschildt P. H., Alexander D. R., Tamanai A., Schweitzer A., 2001, *ApJ*, 556, 357
- Bailer-Jones C. A. L., Mundt R., 2001, *A& A*, 367, 218
- Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1998, *A& A*, 337, 403
- Barrado y Navascués D., Bouvier J., Stauffer J. R., Lodieu N., McCaughrean M. J., 2002, *A& A*, 395, 813
- Basri G., Martín E. L., 1999, *AJ*, 118, 2460
- Bate M. R., 1998, *ApJ*, 508, L95
- Bate M. R., Bonnell I. A., Bromm V., 2002, *MNRAS*, 332, 65
- Bessell M. S., 1986, *PASP*, 98, 1303
- Bonnell I. A., Bate M. R., 1994, *MNRAS*, 271, 999
- Bouvier J., Stauffer J. R., Martín E. L., Barrado y Navascués D., Wallace B., Bejar V. J. S., 1998, *A& A* 336, 490
- Burgasser A. J., Kirkpatrick J. D., Reid I. N., Brown M. E., Miskey C. L., Gizis J. E., 2003, *ApJ*, in press
- Burrows A., Hubbard W. B., Saumon D., Lunine J. I., 1993, *ApJ*, 406, 158
- Burrows A., Marley M., Hubbard W. B., Lunine J. I., Guillot T., Saumon D., Freedman R., Sudarsky D., Sharp C., 1997, *ApJ*, 491, 856
- Carpenter J. M., 2001, *AJ*, 121, 2851
- Chabrier G., Baraffe I., Allard F., Hauschildt P. H., 2000, *ApJ*, 542, 464
- Chabrier G., Baraffe I., 1997, *A& A*, 327, 1039
- Chappelle R. J., Pinfield D. J., Steele I. A., 2002, in proceedings of the IAU Symposium 211 on Brown Dwarfs, in press
- Close L. M., Siegler N., Freed M., Biller B., 2003, *ApJ*, 587, 407
- Crawford D. L., Perry C. L., 1976, *AJ*, 81, 419
- Dahn C. C., Harris H. C., Vrba F. J., Guetter H. H., Canzian B., Henden A. A., Levine S. E., Luginbuhl C. B., Monet A. K. B., Monet D. G., Pier J. R., Stone R. C., Walker R. L., Burgasser A. J., Gizis J. E., Kirkpatrick J. D., Leibert J., Reid I. N., 2002, *AJ*, 124, 1170
- D'Antona F., Mazzitelli I., 1994, *ApJS*, 90, 467
- Dobbie P. D., Kenyon F., Jameson R. F., Hodgkin S. T., 2002a, *MNRAS*, 331, 445
- Dobbie P. D., Kenyon F., Jameson R. F., Hodgkin S. T., Pinfield D. J., Osbourne S. L., 2002b, *MNRAS*, 335, 687
- Dobbie P. D., Pinfield D. J., Jameson R. F., Hodgkin S. T., 2002c, *MNRAS*, 335, 79L
- Duquennoy A., Mayor M., 1991, *A& A*, 248, 485
- Friel E. D., Boesgaard A. M., 1992, *ApJ*, 387, 170
- Hambly N. C., Hawkins M. R. S., Jameson R. F., 1993, *A& AS*, 100, 607
- Hambly N. C., Steele I. A., Hawkins M. R. S., Jameson R. F., 1995a, *A& AS*, 109, 29 (HSHJ)
- Hambly N. C., Steele I. A., Hawkins M. R. S., Jameson R. F., 1995b, *MNRAS*, 273, 505
- Hauschildt P. H., Allard F., Baron E., 1999, *ApJ*, 512, 377
- Hawarden T. G., Leggett S. K., Letawsky M. B., Ballantyne D. R., Casali M. M., 2001, *MNRAS*, 325, 563
- Hawley S. L., Tourtellot J. G., Reid I. N., 1999, *AJ*, 117, 1341
- Hodgkin S. T., Pinfield D. J., Jameson R. F., Steele I. A., Cossburn M. R., Hambly N. C., 1999, *MNRAS*, 310, 87
- Holland K., Jameson R. F., Hodgkin S. T., Davies M. B., Pinfield D. J., 2000, *MNRAS*, 319, 956
- Jameson R. F., Dobbie P. D., Hodgkin S. T., Pinfield D. J., 2002, *MNRAS*, 335, 853
- Jeffries R. D., Thurston M. R., Hambly N. C., 2001, *A& A*, 375, 863
- Jones H. R. A., Tsuji T., 1997, *ApJ*, 480, L39
- King I. R., 1962, *AJ*, 67, 471
- Kirkpatrick J. D., Henry T. J., McCarthy D. W. Jr., 1991, *ApJS*, 77, 417
- Kirkpatrick J. D., Henry T. J., Simons D. A., 1995, *AJ*, 109, 797
- Kirkpatrick J. D., McGraw J. T., Hess T. R., Liebert J., McCarthy D. W. Jr., 1994, *ApJS*, 94, 749
- Kroupa P., 2001, *MNRAS*, 322, 231
- Kroupa P., Aarseth S., Hurley J., 2001, *MNRAS*, 321, 699
- Kroupa P., Tout C. A., Gilmore G., 1990, *MNRAS*, 244, 76
- Leggett S. K., 1992, *ApJS*, 82, 351
- Leggett S. K., Allard F., Hauschildt P. H., 1998, *ApJ*, 509, 836
- Liebert J., Kirkpatrick J. D., Reid I. N., Fisher M. D., 1999, *ApJ*, 519, 345
- Lockwood G. W., Thompson D. T., Radick R. R., Osborn W. H., Baggett W. E., Duncan D. K., Hartmann L. W., 1984, *PASP*, 96, 714
- Lucas P. W., Roche P. F., 2000, *MNRAS*, 314, 858
- Lynga G., 1987, Catalogue, 5th edition (Strasbourg)
- Magazzu A., Rebolo R., Zapatero-Osorio M. R., Martín E. L., Hodgkin S. T., 1998, *ApJ*, 497, 47
- Makarov, V. V., 2002, *AJ*, 124, 3299
- Martín E. L., Basri G., Brandner W., Bouvier J., Zapatero-Osorio M. R., Rebolo R., Stauffer J. R., Allard F., Baraffe I., Hodgkin S. T., 1998, *ApJ*, 509, 113
- Martín E. L., Brandner W., Bouvier J., Luhman K. L., Stauffer J., Basri G., Zapatero-Osorio M. R., Barrado y Navascués D., 2000, *ApJ*, 543, 299

- Martín E. L., Delfosse X., Basri G., Goldman B., Forveille T., Zapatero-Osorio M. R., 1999, *AJ*, 118, 2466
- Martín E. L., Kun M., 1996, *A& AS* 116, 467
- Martín E. L., Rebolo R., Zapatero-Osorio M. R., 1996, *ApJ*, 469, 706
- Mermilliod J. -C., Turon C., Robichon N., Arenou F., Lebreton Y., 1997, in proceedings of the ESA Symposium ‘Hipparcos - Venice ’97’, 643
- Nissen P. E., 1988, *A& A*, 199, 146
- Pinfield D. J., 1997, PhD Thesis, Univ. Leicester
- Pinfield D. J., Hodgkin S. T., Jameson R. F., Cossburn M. R., Hambly N. C., Devereux N., 2000, *MNRAS*, 313, 347
- Pinfield D. J., Hodgkin S. T., Jameson R. F., Cossburn M. R., von Hippel T., 1997, *MNRAS*, 287, 180
- Pinfield D. J., Jameson R. F., Hodgkin S. T., 1998, *MNRAS*, 299, 955
- Prosser C. F., 1992, *AJ*, 103, 488
- Reglero V., Fabregat J., 1991, *A& AS*, 90, 25
- Reid I. N., Cruz K. L., 2002, *AJ*, 123, 2806
- Reipurth B., Clarke C., 2001, *AJ*, 122, 432
- Scholz A., Eisloffel J., Mundt R., 2000, *Astronomische Gesellschaft Abstract Series*, 17, A05
- Scrutskie M. F., Beichman C., Capps R., Carpenter J., Chester T., Cutri R., Elias J., Elston R., Huchra J., Liebert J., Lonsdale C., Monet D., Price S., Schneider S., Seitzer P., Stiening R., Strom S., Weinberg M., 1995, *AAS*, 187, 7507
- Sharp C. M., Huebner W. F., 1990, *ApJS*, 72, 417
- Simons D. A., Tokunaga A., 2002, *PASP*, 114, 169
- Stauffer J. R., Barrado y Navascués D., Bouvier J., Morrison H. L., Harding P., Luhman K. L., Stanke T., McCaughrean M., Terndrup D. M., Allen L., Assoud P., 1999, *ApJ*, 527, 219
- Stauffer J. R., Giampapa M. S., Herbst W., Vincent J. M., Hartmann L. W., Stern R. A., 1991, *ApJ*, 374, 142
- Stauffer J. R., Schultz G., Kirkpatrick J. D., 1998, *ApJ*, 499, L199
- Tsuji T., 2003, *ApJ*, in press (astro-ph/0204401)
- van Leeuwen F., 1999, *A& A*, 341, 71L
- Zapatero-Osorio M. R., Rebolo R., Martín E. L., Hodgkin S. T., Cossburn M. R., Magazzu A., Steele I. A., Jameson R. F., 1999, *A& A Supp.*, 134, 537

APPENDIX A: TABLES OF PHOTOMETRY

Table A1. Photometry (c3-6), membership criteria (c7-9,c11-13) and binarity (c10) of Pleiades BD candidates. Membership is from the IK CMD, the JK CMD and the JHK diagram. Refs; (a) Martín, Rebolo & Zapatero-Osorio (1996), (b) Rebolo et al. (1996), (c) Zapatero Osorio et al. (1997), (d) Martín et al. (1998) (e) Stauffer et al. (1998), (f) Martín et al. (2000), (g) Pinfield et al. (2000), (h) Moraux, Bouvier & Stauffer (2001), (i) Dobbie et al. (2002), (j) Dobbie (2002) private comm. A * indicates a non-member by virtue of binarity.

Name	Other names	I_C	J_{MKO}	H_{MKO}	K_{MKO}	IK	JK	JHK	Bin	SpT	pm	Li
BPL 45		18.61 ^g	16.07±0.05	15.42±0.03	14.93±0.04	y	N*	y	b	y		
BPL 49	CFHT-PI-17	18.55 ^g	16.12±0.04 ^g 16.14±0.10 ⁱ 15.99±0.04	15.49±0.07 ^g 15.51±0.11 ⁱ 15.48±0.05	15.07±0.06 ^g	y	y	y			y ^h	
BPL 62	Roque 7 CFHT-PI-24	19.44 ^g	16.53±0.03 ^g 16.55±0.18 ⁱ	15.85±0.02 ^g 15.88±0.14 ⁱ	15.47±0.02 ^g	y	y	y			y ^h	
BPL 66	Roque 4	19.81 ^g	16.66±0.05 16.62±0.04 16.69±0.12 ⁱ	15.93±0.03 15.92±0.05 16.04±0.09 ⁱ	15.26±0.04 15.26±0.04	y	y	y	b	y ^c		
BPL 76		17.83 ^g	15.41±0.04	14.85±0.03	14.57±0.06 ^g	y	y	y				
BPL 78		18.98 ^g	16.02±0.05	15.45±0.03	15.01±0.04	y	y	y				
BPL 79	Roque 13	18.69 ^g	15.63±0.03 ^g 15.65±0.09 ⁱ	15.09±0.02 ^g 15.11±0.08 ⁱ	14.56±0.03 ^g	y	y	y	b	y ^c		
BPL 81		18.79 ^g	15.97±0.03	15.39±0.05	14.98±0.02 ^g	y	y	y				
BPL 100	Roque 9	19.33 ^g	16.28±0.03 16.30±0.19 ⁱ	15.63±0.05 15.72±0.15 ⁱ	15.22±0.02 ^g	y	y	y				
BPL 108	Roque 14	18.53 ^g	15.50±0.02 ^g 15.53±0.10 ⁱ 15.59±0.03	14.89±0.02 ^g 14.96±0.09 ⁱ 14.96±0.03	14.47±0.03 ^g	y	y	y	b	y ^c		
BPL 124		17.78 ^g	15.43±0.05	14.85±0.03	14.46±0.04	y	y	y			y ^g	
BPL 132	Roque 11 NPL 37	19.06 ^g	16.08±0.03 ^g	15.64±0.03 ^g	15.13±0.04 ^g	y	y	N		y ^b		
BPL 137	Teide 1 NPL 39	19.22 ^g	16.32±0.05 ^g 16.19±0.11 ⁱ 16.24±0.05	15.62±0.05 ^g 15.65±0.09 ⁱ 15.62±0.05	15.08±0.05 ^g	y	y	y	b	y ^b	y ^h	y ^c
BPL 142	Roque 17	18.11 ^g	15.29±0.03 ^g 15.31±0.09 ⁱ	14.77±0.03 ^g 14.79±0.07 ⁱ	14.36±0.05 ^g	y	y	y	b	y ^c	y ^g	
BPL 152	Roque 16 CFHT-PI-11	17.95 ^g	15.49±0.03 ^g 15.51±0.11 ⁱ	14.97±0.02 ^g 14.99±0.09 ⁱ	14.62±0.03 ^g	y	y	y		y ^c	y ^h	y ^e
BPL 163		17.83 ^g	15.31±0.03	14.78±0.05	14.53±0.04 ^g	y	N	y				
BPL 168		17.65 ^g	—	—	16.02±0.10	N	—	—				
BPL 172	Roque 12 NPL 36	18.53 ^g	15.91±0.03 15.93±0.10 ⁱ	15.36±0.05 15.44±0.08 ⁱ	15.10±0.03 ^g 14.97±0.07 ⁱ	y	y	y		y ^c		
BPL 201		19.60 ^g	16.40±0.03 ^g	15.75±0.03 ^g	15.35±0.04 ^g	y	y	y				
BPL 202	CFHT-PI-9	17.75 ^g	15.44±0.03 15.40±0.04	14.88±0.05 14.84±0.02	14.49±0.04 ^g	y	y	y		y ^e	y ^h	N ^e
BPL 235	Calar 3 CFHT-PI-21	18.91 ^g	16.24±0.03 ^g 16.09±0.04	15.53±0.03 ^g 15.49±0.05	14.91±0.03 ^g	y	y	y	b	y ^a	y ^h	y ^b
BPL 240		18.68 ^g	15.99±0.03	15.44±0.05	15.11±0.03 ^g	y	y	y				
BPL 249		20.02 ^g	16.66±0.03 ^g	16.13±0.03 ^g	15.46±0.03 ^g	y	y	y	b			
BPL 254	Teide 2 CFHT-PI-13	17.81 ^g	15.53±0.04 15.53±0.08 ⁱ	14.94±0.03 14.97±0.07 ⁱ	14.57±0.06 ^g 14.54±0.06 ⁱ	y	y	y		y ^d	y ^h	y ^{d,e}
BPL 283	CFHT-PI-18	18.58 ^g	16.04±0.03 ^g	15.43±0.02 ^g	14.94±0.06 ^g	y	N*	y	b	y ^f	N ^h	N ^f
BPL 294	CFHT-PI-12	17.85 ^g	15.16±0.03	14.55±0.02	14.20±0.02 ^g	y	y	y	b	y ^e	y ^h	y ^e
BPL 303	CFHT-PI-25	19.69 ^g	16.63±0.04 16.64±0.15 ⁱ	16.01±0.02 16.05±0.13 ⁱ	15.47±0.06 ^g	y	y	y	b		y ^h	
BPL 306		19.09 ^g	16.16±0.04	15.53±0.02	15.15±0.02 ^g	y	y	y				
BPL 316		18.30 ^g	15.97±0.04	15.41±0.02	15.01±0.03 14.93±0.04	y	y	y				
BPL 327	IPMBD 11	17.94 ^g	15.52±0.03	14.99±0.02	14.60±0.02 ^g	y	y	y		y	y ^g	
PPL 1	Roque 15	18.21 ⁱ	15.37±0.08 ⁱ	14.82±0.06 ⁱ	14.34±0.05 ⁱ	y	y	y	b	y		y
PPL 15	IPMBD 23 NPL 35	18.24 ⁱ	15.34 ⁱ	14.77 ⁱ	14.41 ⁱ	y	y	y	b	y	y	y
CFHT-PI-23		19.33 ⁱ	16.38±0.08 ⁱ	15.79±0.08 ⁱ	15.25±0.05 ⁱ	y	y	y	b			
Roque 5		20.19 ⁱ	16.89±0.14 ⁱ	16.33±0.15 ⁱ	15.74±0.10 ⁱ	y	y	y				
Roque 8		19.55 ⁱ	16.68±0.11 ⁱ	16.14±0.10 ⁱ	15.57±0.06 ⁱ	y	y	y				
Roque 25		21.80 ⁱ	17.7±0.3 ⁱ	17.0±0.3 ⁱ	16.31±0.19 ⁱ	y	y	y		y		
Roque 30		20.82 ⁱ	17.49±0.12 ⁱ	16.77±0.11 ⁱ	16.14±0.07 ⁱ	y	y	y				
Roque 33	NPL 40	20.44 ⁱ	17.07±0.12 ⁱ	16.62±0.18 ⁱ	16.02±0.08 ⁱ	y	y	?		y	y ^j	y
Roque 36		20.25 ⁱ	17.17±0.20 ⁱ	16.58±0.12 ⁱ	16.09±0.08 ⁱ	y	y	y				

Table A2. Photometry (c2-5) and membership (c6-8) of Pleiades low-mass star candidates (higher mass BPL candidates). Membership is from the IK CMD, the JK CMD and the JHK diagram. J- H- and K-band photometry is from 2MASS, and has been transformed into the MKO system.

BPL	I _C	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK
3	14.81	13.25±0.03	12.74±0.03	12.36±0.03	y	y	y
4	14.17	12.67±0.03	12.12±0.02	11.77±0.03	y	y	y
5	14.25	12.66±0.03	12.17±0.03	11.80±0.03	y	y	y
6	15.74	14.01±0.03	13.53±0.03	13.09±0.04	y	y	y
7	16.28	14.44±0.03	13.92±0.03	13.48±0.04	y	y	y
16	17.02	15.20±0.04	14.67±0.05	14.24±0.06	N	y	y
18	14.39	12.89±0.03	12.35±0.03	11.97±0.03	y	y	y
19	16.39	14.60±0.03	14.09±0.04	13.76±0.04	y	y	y
20	14.81	13.21±0.03	12.70±0.03	12.28±0.03	y	y	y
26	17.00	14.67±0.03	14.16±0.04	13.76±0.05	y	y	y
30	15.90	14.15±0.03	13.58±0.03	13.22±0.04	y	y	y
34	13.72	12.01±0.02	11.45±0.03	11.10±0.03	y	y	y
38	16.12	14.27±0.03	13.78±0.04	13.42±0.04	y	y	y
43	15.81	14.11±0.03	13.56±0.03	13.18±0.04	y	y	y
44	16.09	14.30±0.03	13.76±0.03	13.37±0.04	y	y	y
56	14.90	13.29±0.03	12.74±0.02	12.40±0.03	y	y	y
58	17.69	15.37±0.05	14.77±0.06	14.23±0.07	y	y	y
61	14.86	13.11±0.02	12.58±0.03	12.23±0.03	y	y	y
68	14.33	12.85±0.03	12.34±0.03	12.03±0.04	y	y	y
69	17.59	16.61±0.15	16.10±0.20	15.56±0.18	N	y	y
70	15.40	13.33±0.02	12.76±0.03	12.45±0.03	y	y	y
71	15.14	13.18±0.02	12.58±0.03	12.23±0.03	y	y	y
72	16.18	14.04±0.03	13.52±0.03	13.20±0.03	y	y	y
73	15.09	13.08±0.02	12.60±0.03	12.21±0.03	y	y	y
74	17.44	15.19±0.05	14.68±0.06	14.23±0.06	y	y	y
75	16.28	14.03±0.03	13.46±0.03	13.06±0.03	y	y	y
84	17.21	14.81±0.04	14.29±0.04	13.85±0.04	y	y	y
89	13.89	12.01±0.03	11.46±0.03	11.14±0.03	y	y	y
91	14.74	12.73±0.03	12.21±0.03	11.88±0.03	y	y	y
92	15.44	13.32±0.03	12.83±0.03	12.44±0.03	y	y	y
93	14.63	12.71±0.03	12.18±0.03	11.81±0.03	y	y	y
94	15.26	13.58±0.03	13.08±0.03	12.74±0.03	y	y	y
95	14.02	12.51±0.03	11.96±0.03	11.65±0.03	y	y	y
96	16.38	14.05±0.03	13.51±0.04	13.13±0.03	y	y	y
97	14.55	12.82±0.03	12.32±0.03	11.96±0.03	y	y	y
98	15.82	13.95±0.03	13.39±0.03	13.11±0.04	y	y	y
99	15.30	13.60±0.03	13.08±0.03	12.74±0.03	y	y	y
101	16.70	14.74±0.04	14.19±0.05	13.82±0.06	y	y	y
102	15.11	13.40±0.03	12.87±0.03	12.55±0.03	y	y	y
103	14.57	12.90±0.03	12.38±0.03	12.05±0.03	y	y	y
104	16.26	14.09±0.03	13.54±0.04	13.19±0.03	y	y	y
105	15.49	13.33±0.03	12.84±0.03	12.43±0.03	y	y	y
106	17.44	15.22±0.06	14.62±0.07	14.15±0.07	y	y	y
107	16.18	14.04±0.03	13.54±0.04	13.16±0.04	y	y	y
109	16.02	14.18±0.03	13.70±0.04	13.28±0.05	y	y	y
110	16.70	14.40±0.03	13.83±0.04	13.47±0.05	y	y	y
111	15.98	14.19±0.03	13.62±0.04	13.23±0.04	y	y	y
112	17.30	14.98±0.05	14.37±0.06	14.21±0.08	y	y	y
113	13.73	12.02±0.02	11.45±0.03	11.12±0.02	y	y	y
114	13.85	12.34±0.02	11.82±0.03	11.48±0.03	y	y	y
115	17.00	14.90±0.05	14.37±0.05	13.98±0.06	y	y	y
116	15.79	13.75±0.03	13.27±0.03	12.87±0.03	y	y	y
117	14.64	13.00±0.03	12.49±0.03	12.12±0.03	y	y	y
118	14.82	13.07±0.02	12.53±0.03	12.17±0.03	y	y	y
119	15.10	13.23±0.03	12.67±0.03	12.29±0.04	y	y	y
120	15.06	13.11±0.03	12.59±0.03	12.23±0.03	y	y	y
121	15.03	13.20±0.03	12.67±0.03	12.33±0.03	y	y	y

BPL	I _C	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK
122	17.13	14.96±0.05	14.46±0.06	14.04±0.07	y	y	y
123	17.32	15.74±0.08	14.98±0.09	14.90±0.15	N	y	N
125	15.80	13.98±0.03	13.40±0.04	13.08±0.03	y	y	y
126	14.34	12.74±0.02	12.18±0.03	11.87±0.03	y	y	y
127	17.71	15.37±0.06	14.84±0.07	14.32±0.08	y	y	y
128	16.45	14.44±0.03	13.91±0.04	13.64±0.05	y	y	y
129	16.23	14.23±0.03	13.76±0.04	13.43±0.04	y	y	y
130	17.27	14.81±0.04	14.27±0.05	13.97±0.06	y	y	y
131	16.12	14.00±0.03	13.55±0.04	13.18±0.04	y	y	y
133	14.02	12.34±0.03	11.80±0.03	11.45±0.03	y	y	y
134	15.60	13.62±0.02	13.12±0.03	12.75±0.03	y	y	y
135	15.22	13.21±0.04	12.73±0.03	12.36±0.03	y	y	y
136	15.29	13.37±0.03	12.84±0.03	12.46±0.04	y	y	y
138	15.09	13.14±0.02	12.65±0.03	12.29±0.03	y	y	y
139	16.23	14.07±0.03	13.53±0.03	13.17±0.04	y	y	y
140	16.44	14.22±0.04	13.71±0.04	13.32±0.04	y	y	y
141	14.27	12.64±0.03	12.13±0.03	11.78±0.03	y	y	y
143	15.70	13.84±0.03	13.34±0.03	13.00±0.03	y	y	y
144	14.06	12.30±0.02	11.75±0.03	11.41±0.03	y	y	y
145	15.34	13.68±0.03	13.17±0.03	12.86±0.04	y	y	y
146	16.77	14.97±0.05	14.40±0.06	14.21±0.07	N	y	y
147	15.50	13.69±0.03	13.22±0.03	12.87±0.03	y	y	y
148	14.93	12.97±0.04	12.42±0.03	12.07±0.03	y	y	y
149	15.45	13.67±0.03	13.10±0.03	12.75±0.04	y	y	y
150	15.46	13.64±0.03	13.12±0.03	12.74±0.03	y	y	y
151	14.04	12.55±0.03	12.03±0.03	11.68±0.03	y	y	y
153	14.76	13.03±0.03	12.49±0.03	12.13±0.03	y	y	y
154	16.94	14.72±0.04	14.20±0.05	13.78±0.05	y	y	y
155	15.52	13.49±0.03	12.88±0.03	12.54±0.03	y	y	y
156	15.30	13.41±0.03	12.91±0.03	12.56±0.04	y	y	y
157	14.78	12.95±0.03	12.42±0.03	12.07±0.03	y	y	y
158	15.27	13.48±0.03	13.01±0.03	12.62±0.03	y	y	y
159	13.78	12.30±0.03	11.77±0.03	11.45±0.03	y	y	y
160	15.06	12.96±0.03	12.47±0.04	12.12±0.03	y	y	y
161	15.78	13.72±0.04	13.20±0.04	12.87±0.03	y	y	y
162	16.07	14.15±0.04	13.62±0.05	13.25±0.05	y	y	y
164	16.38	14.43±0.04	13.96±0.05	13.50±0.06	y	y	y
165	14.04	11.81±0.03	11.37±0.04	10.97±0.03	y	y	y
166	14.09	12.35±0.04	11.82±0.03	11.48±0.03	y	y	y
167	13.74	12.13±0.03	11.55±0.05	11.24±0.02	y	y	y
169	17.60	15.32±0.06	14.71±0.07	14.43±0.08	y	y	y
170	15.68	13.89±0.04	13.37±0.05	12.97±0.03	y	y	y
171	15.43	13.41±0.04	12.90±0.03	12.59±0.03	y	y	y
173	14.16	12.42±0.03	11.92±0.06	11.58±0.02	y	y	y
174	15.55	13.72±0.04	13.26±0.04	12.92±0.03	y	y	y
175	14.05	12.31±0.03	11.77±0.03	11.48±0.03	y	y	y
176	14.42	12.87±0.04	12.39±0.04	12.00±0.04	y	y	y
177	17.06	14.86±0.05	14.40±0.05	14.00±0.06	y	y	y
178	14.65	12.82±0.03	12.31±0.03	11.93±0.03	y	y	y
179	15.61	15.07±0.05	14.77±0.07	14.56±0.09	N	N	N
180	13.72	12.20±0.03	11.65±0.03	11.34±0.03	y	y	y
181	17.33	14.73±0.04	14.16±0.05	13.70±0.05	y	y	y
182	16.06	14.05±0.03	13.52±0.04	13.18±0.04	y	y	y
183	16.84	14.85±0.04	14.33±0.05	13.84±0.05	y	y	y
184	14.21	12.68±0.03	12.14±0.03	11.81±0.03	y	y	y
185	14.86	13.18±0.04	12.66±0.06	12.32±0.03	y	y	y
186	16.59	14.57±0.04	14.06±0.05	13.56±0.04	y	y	y

BPL	I _C	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK
187	14.09	12.55±0.03	12.02±0.03	11.67±0.02	y	y	y
188	16.26	14.31±0.04	13.80±0.06	13.44±0.04	y	y	y
189	15.22	13.45±0.03	12.98±0.03	12.57±0.03	y	y	y
190	15.86	14.02±0.04	13.50±0.06	13.11±0.04	y	y	y
191	17.74	15.06±0.05	14.51±0.06	14.08±0.06	y	y	y
192	17.46	15.09±0.05	14.56±0.06	14.08±0.06	y	y	y
193	13.73	12.10±0.03	11.55±0.03	11.27±0.02	y	y	y
194	15.89	13.93±0.04	13.45±0.04	13.09±0.04	y	y	y
197	16.42	14.17±0.04	13.66±0.04	13.33±0.04	y	y	y
198	16.95	14.71±0.04	14.16±0.05	13.74±0.05	y	y	y
206	16.43	14.16±0.04	13.63±0.04	13.30±0.05	y	y	y
208	15.30	13.34±0.04	12.79±0.04	12.48±0.03	y	y	y
209	14.09	12.58±0.03	12.05±0.03	11.73±0.03	y	y	y
210	16.63	14.57±0.04	14.08±0.05	13.66±0.05	y	y	y
211	15.99	13.88±0.03	13.38±0.04	13.00±0.04	y	y	y
212	14.90	13.22±0.03	12.69±0.03	12.37±0.03	y	y	y
214	17.14	14.70±0.04	14.10±0.04	13.69±0.05	y	y	y
215	17.10	14.99±0.05	14.47±0.06	14.14±0.06	y	y	y
216	14.76	12.91±0.03	12.26±0.04	11.97±0.03	y	y	y
217	13.85	12.33±0.03	11.83±0.04	11.52±0.03	y	y	y
218	17.38	15.13±0.05	14.57±0.06	14.36±0.09	y	y	y
219	16.14	14.18±0.04	13.74±0.04	13.37±0.05	y	y	y
220	16.46	14.39±0.04	13.83±0.04	13.40±0.04	y	y	y
221	16.38	14.41±0.04	13.93±0.07	13.48±0.05	y	y	y
222	14.96	13.24±0.03	12.71±0.04	12.35±0.03	y	y	y
223	17.15	15.10±0.06	14.61±0.08	14.25±0.09	y	y	y
224	14.71	12.85±0.03	12.39±0.05	12.01±0.04	y	y	y
225	15.54	13.71±0.03	13.26±0.06	12.91±0.03	y	y	y
226	14.86	13.16±0.03	12.64±0.04	12.37±0.03	y	y	y
227	15.44	13.75±0.03	13.27±0.05	12.94±0.03	y	y	y
228	17.37	15.05±0.05	14.53±0.07	14.03±0.07	y	y	y
229	15.40	13.58±0.03	13.03±0.05	12.66±0.04	y	y	y
230	14.18	12.60±0.03	12.06±0.06	11.70±0.04	y	y	y
231	15.10	13.35±0.03	12.85±0.03	12.53±0.03	y	y	y
232	16.02	14.20±0.04	13.79±0.05	13.28±0.04	y	y	y
233	13.70	12.27±0.03	11.73±0.04	11.39±0.03	y	y	y
238	15.59	13.70±0.03	13.11±0.03	12.73±0.03	y	y	y
239	15.19	13.56±0.03	13.06±0.03	12.69±0.03	y	y	y
241	14.26	12.95±0.03	12.39±0.03	12.04±0.03	y	y	y
242	16.11	14.26±0.03	13.73±0.04	13.29±0.04	y	y	y
243	13.97	12.42±0.03	11.86±0.03	11.47±0.03	y	y	y
244	14.29	12.64±0.03	12.11±0.03	11.72±0.02	y	y	y
245	15.07	13.49±0.03	12.96±0.03	12.58±0.03	y	y	y
246	15.16	13.60±0.03	13.03±0.03	12.74±0.04	y	y	y
247	16.90	15.21±0.06	14.57±0.06	14.20±0.07	N	y	y
248	15.16	13.45±0.03	12.95±0.03	12.63±0.04	y	y	y
250	15.49	13.61±0.03	13.17±0.03	12.80±0.04	y	y	y
251	15.57	13.84±0.03	13.29±0.04	12.95±0.04	y	y	y
252	15.38	13.80±0.03	13.29±0.03	12.91±0.04	y	y	y
253	17.24	15.26±0.05	14.63±0.06	14.25±0.07	y	y	y
255	16.07	14.11±0.03	13.64±0.03	13.28±0.04	y	y	y
260	15.73	13.97±0.03	13.48±0.04	13.07±0.04	y	y	y
264	14.98	13.41±0.03	12.86±0.03	12.49±0.04	y	y	y
267	16.41	14.61±0.04	14.06±0.05	13.67±0.05	y	y	y
272	16.53	14.74±0.04	14.14±0.05	13.70±0.06	y	y	y
276	14.01	13.17±0.03	12.84±0.03	12.70±0.04	N	N	N
280	16.75	14.77±0.04	14.32±0.06	13.79±0.06	y	y	y
281	15.79	14.09±0.03	13.55±0.04	13.16±0.04	y	y	y
285	14.97	13.36±0.03	12.83±0.03	12.52±0.04	y	y	y
290	14.71	13.14±0.03	12.60±0.03	12.27±0.03	y	y	y
291	16.40	14.37±0.04	13.83±0.04	13.44±0.05	y	y	y
292	16.73	14.96±0.05	14.46±0.06	14.15±0.07	N	y	y

BPL	I _C	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK
293	17.16	15.17±0.05	14.67±0.07	14.28±0.08	y	y	y
296	16.79	15.07±0.05	14.50±0.06	14.11±0.07	N	y	y
298	14.20	12.66±0.03	12.16±0.03	11.81±0.03	y	y	y
299	16.61	14.74±0.04	14.24±0.05	13.84±0.05	y	y	y
301	15.58	13.85±0.03	13.30±0.03	12.93±0.03	y	y	y
305	14.07	12.62±0.03	12.10±0.03	11.74±0.02	y	y	y
310	14.44	12.90±0.03	12.36±0.04	12.06±0.04	y	y	y
311	14.96	13.41±0.03	12.85±0.03	12.54±0.03	y	y	y
313	16.44	14.55±0.04	13.99±0.05	13.74±0.05	y	y	y
314	13.82	12.65±0.03	12.07±0.03	11.78±0.03	N	y	y
318	13.88	12.61±0.03	12.00±0.03	11.74±0.03	N	y	y
319	13.75	12.87±0.03	12.44±0.03	12.28±0.03	N	N	N
320	15.59	14.02±0.03	13.50±0.03	13.15±0.03	y	y	y
321	14.41	12.91±0.03	12.30±0.03	11.93±0.04	y	y	y
322	15.86	14.19±0.03	13.68±0.03	13.25±0.04	y	y	y
323	16.27	14.38±0.03	13.93±0.04	13.56±0.04	y	y	y
324	14.15	12.77±0.03	12.21±0.03	11.84±0.03	y	y	y
325	17.41	15.21±0.05	14.81±0.07	14.28±0.08	y	y	y
326	16.96	15.06±0.04	14.50±0.05	14.27±0.08	N	y	y
327	18.00	15.52±0.05	14.93±0.07	14.78±0.12	y	y	y
328	16.95	14.72±0.04	14.10±0.04	13.68±0.05	y	y	y
329	14.52	13.05±0.03	12.47±0.03	12.15±0.03	y	y	y
330	15.73	14.16±0.03	13.57±0.03	13.23±0.04	y	y	y
331	17.67	15.41±0.05	14.75±0.06	14.34±0.09	y	y	y
332	16.41	14.59±0.04	14.04±0.05	13.66±0.05	y	y	y
333	14.69	13.02±0.03	12.45±0.03	12.13±0.03	y	y	y
334	17.47	15.27±0.05	14.67±0.06	14.31±0.08	y	y	y
335	15.36	13.55±0.03	13.05±0.04	12.66±0.03	y	y	y
336	17.46	15.30±0.05	14.70±0.06	14.26±0.08	y	y	y
337	16.71	14.69±0.04	14.15±0.04	13.73±0.05	y	y	y
338	16.81	15.07±0.04	14.52±0.05	14.24±0.07	N	y	y
339	13.84	12.43±0.03	11.88±0.03	11.46±0.03	y	y	y

Table A3. Photometry (c2-5), membership (c6-8) and binarity (c9) of RIZ-Pr Praesepe candidates. Membership is from the IK CMD, the JK CMD and the JHK diagram. Refs; (a) Pinfield et al. (1997), (b) Hodgkin et al. (1999). The [†] is also IZ-Pr 40. The [‡] is also IZ-Pr 86. n-d = non-detection

RIZ-Pr	I _C ^a	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK	Bin
1	20.14	17.06±0.05	16.47±0.04	16.14±0.08 ^b	y	y	y	
2 [†]	18.19	15.67±0.05	15.11±0.03	14.77±0.04 ^b	y	y	y	b
3	21.20	19.05±0.17	18.01±0.12	16.77±0.08	y	N	N	
4	18.52	16.04±0.03	15.56±0.03	15.23±0.04 ^b	y	y	y	
5	20.86	19.09±0.17	18.20±0.16	16.94±0.06	y	N	N	
6	21.09	19.10±0.18	18.11±0.16	17.07±0.08	y	N	N	
				17.01±0.07				
				17.32±0.05 ^b				
7	21.24	–	–	18.56±0.29	N	–	–	
				19.15±0.05 ^b				
8	17.81	15.40±0.05	14.89±0.03	14.60±0.04 ^b	y	y	y	
9	21.25	–	–	17.72±0.13	N	–	–	
10	20.54	–	–	18.08±0.07 ^b	N	–	–	
11	19.47	16.66±0.05	16.23±0.04	15.84±0.14 ^b	y	y	y	
12	20.10	–	–	16.91±0.21 ^b	N	–	–	
13	20.24	–	–	19.04±0.50	N	–	–	
				18.49±0.11 ^b				
14	20.59	–	–	17.82±0.06 ^b	N	–	–	
15	21.36	–	–	18.49±0.27	N	–	–	
16	21.10	–	–	18.60±0.31	N	–	–	
17	20.41	18.23±0.10	17.65±0.12	17.28±0.11	N	N	y	
18	19.63	16.40±0.04	15.74±0.03	15.40±0.04 ^b	y	y	N?	b
19	21.42	–	–	18.04±0.18	N	–	–	
20	18.48	16.16±0.04	15.57±0.03	15.19±0.04 ^b	y	y	y	
21 [‡]	18.73	15.89±0.03	15.39±0.03	15.03±0.03 ^b	y	y	y	b
22	21.55	–	–	n-d	N	–	–	
23	19.06	16.63±0.04	16.04±0.03	15.53±0.03	y	y	y	b
				15.98±0.16 ^b				
24	20.43	17.30±0.05	16.68±0.04	16.17±0.06	y	y	y	b
				16.19±0.03 ^b				
25	21.20	–	–	18.62±0.31	N	–	–	
26	18.08	–	–	15.49±0.04 ^b	N	–	–	

Table A4. Photometry (c2-5), membership (c6-8) and binarity (c9) of IZ-Pr Praesepe candidates. Membership is from the IK CMD, the JK CMD and the JHK diagram. Refs; (a) Pinfield etal (2000), (b) 2MASS. The [†] is also RIZ-Pr 2. The [‡] is also RIZ-PR 21.

IZ-Pr	I _C ^a	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK	Bin
1	18.02	15.60±0.03	15.13±0.05	14.72±0.03	y	y	y	b
2	17.25	15.44±0.03	14.94±0.05	14.64±0.03	y	y	y	
3	18.19	15.87±0.03	15.36±0.05	15.09±0.02	y	y	y	
4	18.21	16.56±0.13 ^b	15.76±0.13 ^b	15.31±0.06	y	y	?	
5	18.33	—	—	15.83±0.03	N	—	—	
6	17.26	15.62±0.06 ^b	15.15±0.08 ^b	14.79±0.03	y	y?	?	
7	19.90	—	—	16.52±0.05	N	—	—	
8	17.22	15.57±0.06 ^b	15.16±0.08 ^b	14.72±0.03	y	y?	?	
9	20.37	—	—	17.66±0.12	N	—	—	
10	18.26	15.79±0.03	15.30±0.05	14.95±0.02	y	y	y	b
11	17.47	15.63±0.03	15.10±0.05	14.82±0.03	y	y	y	
12	19.93	—	—	17.36±0.09	N	—	—	
13	19.14	—	—	16.08±0.04	N	—	—	
14	17.84	16.03±0.07 ^b	15.50±0.09 ^b	15.02±0.03	y	y	?	
15	17.94	16.08±0.08 ^b	15.63±0.09 ^b	15.11±0.03	y	y	?	
16	17.18	—	—	14.99±0.03	N	—	—	
17	17.61	15.74±0.03	15.24±0.05	14.86±0.03	y	y	y	
18	18.54	16.18±0.03	15.81±0.03	15.30±0.04	y	y	N	
19	19.93	—	—	16.78±0.06	N	—	—	
20	17.22	15.43±0.03	14.92±0.05	14.58±0.03	y	y	y	
21	19.72	—	—	17.19±0.08	N	—	—	
22	19.65	—	—	16.18±0.04	y	—	—	
23	19.20	16.47±0.03	16.04±0.03	15.57±0.04	y	y	y	
24	17.52	15.47±0.03	14.95±0.05	14.56±0.03	y	y	y	b
25	17.98	15.49±0.03	14.94±0.05	14.64±0.02	y	y	y	b
26	17.38	15.30±0.03	14.86±0.04	14.52±0.03	y	y	y	
27	19.81	—	—	16.77±0.06	N	—	—	
28	17.89	16.22±0.09 ^b	15.50±0.09 ^b	15.15±0.03	y	y	?	
29	19.62	—	—	16.90±0.06	N	—	—	
30	17.55	15.51±0.03	15.09±0.05	14.72±0.03	y	y	y	
31	17.41	15.38±0.03	14.91±0.02	14.54±0.03	y	y	y	
32	19.19	—	—	15.88±0.03	y	?	—	
33	17.30	15.36±0.03	14.90±0.05	14.55±0.03	y	y	y	
34	19.34	—	—	17.49±0.10	N	—	—	
35	19.77	—	—	16.50±0.05	N	—	—	
36	19.41	16.62±0.04	16.11±0.03	15.66±0.02	y	y	y	
37	19.28	—	—	16.40±0.04	N	—	—	
38	18.22	—	—	15.52±0.03	N	—	—	
39	19.44	—	—	16.58±0.05	N	—	—	
40 [†]	18.04	15.74±0.04	15.08±0.03	14.79±0.04	y	y	y	
41	18.02	15.75±0.04	15.28±0.02	14.83±0.03	y	y	y	b
42	20.33	—	—	17.03±0.08	N	—	—	
43	19.47	—	—	16.09±0.10	y	—	—	
44	17.49	15.76±0.06 ^b	15.32±0.08 ^b	14.84±0.04	y	y	?	
45	17.18	—	—	15.46±0.11	N	—	—	
50	18.29	—	—	15.43±0.03	y	—	—	
52	19.21	—	—	16.27±0.04	N	—	—	
54	17.67	15.87±0.08 ^b	15.39±0.10 ^b	15.07±0.14 ^b	y	y?	?	
55	17.40	15.74±0.07 ^b	15.18±0.08 ^b	15.15±0.15 ^b	N	N	?	
56	17.38	15.51±0.06 ^b	14.93±0.07 ^b	14.53±0.09 ^b	y	y	?	
57	17.45	15.62±0.06 ^b	15.01±0.07 ^b	14.62±0.09 ^b	y	y	?	
59	18.47	—	—	15.83±0.04	N	—	—	
60	17.48	15.60±0.06 ^b	15.20±0.09 ^b	14.81±0.11 ^b	y	y?	?	
61	17.59	16.07±0.09 ^b	15.38±0.10 ^b	14.99±0.13 ^b	y	y	?	
62	17.77	16.07±0.09 ^b	15.55±0.11 ^b	15.11±0.14 ^b	y	y	?	
63	17.42	15.72±0.06 ^b	14.99±0.08 ^b	14.51±0.08 ^b	y	y	?	b
64	17.44	15.30±0.05 ^b	14.66±0.06 ^b	14.19±0.06 ^b	y	y	?	b
66	17.26	15.40±0.05 ^b	14.81±0.07 ^b	14.61±0.09 ^b	y	y?	?	
69	17.30	15.45±0.05 ^b	14.89±0.07 ^b	14.53±0.09 ^b	y	y	?	

IZ-Pr	I _C ^a	J _{MKO}	H _{MKO}	K _{MKO}	IK	JK	JHK	Bin
70	17.32	15.40±0.05 ^b	14.74±0.07 ^b	14.53±0.09 ^b	y	y?	?	
72	17.29	15.38±0.05 ^b	14.93±0.07 ^b	14.36±0.09 ^b	y	y	?	b
74	17.87	15.89±0.07 ^b	15.06±0.08 ^b	14.90±0.12 ^b	y	y	?	
77	17.41	15.71±0.06 ^b	15.14±0.08 ^b	14.76±0.10 ^b	y	y	?	
80	17.37	15.55±0.05 ^b	14.81±0.07 ^b	14.52±0.08 ^b	y	y	?	
82	19.24	–	–	16.54±0.04	N	–	–	
84	17.99	15.95±0.07 ^b	15.26±0.08 ^b	15.13±0.14 ^b	y	y?	?	
85	18.92	–	–	17.03±0.05	N	–	–	
86 [‡]	18.63	15.89±0.03	15.39±0.03	15.01±0.03	y	y	y	
88	17.98	16.11±0.09 ^b	15.85±0.15 ^b	15.15±0.15 ^b	y	y	?	
93	19.29	16.76±0.04	16.21±0.03	15.75±0.03	y	y	y	
94	18.25	16.02±0.10 ^b	15.44±0.11 ^b	15.00±0.03	y	y	–	b
95	19.47	–	–	16.78±0.03	N	–	–	
97	19.50	–	–	16.75±0.04	N	–	–	
99	17.86	15.88±0.09 ^b	15.19±0.10 ^b	14.98±0.13 ^b	y	y	?	
100	17.34	15.40±0.06 ^b	14.95±0.09 ^b	14.59±0.08 ^b	y	y?	?	
104	19.50	–	–	16.84±0.06	N	–	–	
105	18.80	–	–	16.15±0.03	N	–	–	
108	17.31	15.47±0.06 ^b	14.83±0.07 ^b	14.57±0.10 ^b	y	y	?	
110	18.68	–	–	16.74±0.05	N	–	–	
112	17.89	16.01±0.09 ^b	15.53±0.12 ^b	15.17±0.15 ^b	y	y?	?	
113	18.44	16.03±0.03	15.58±0.03	15.17±0.03	y	y	y	
117	19.09	16.56±0.04	16.02±0.03	15.70±0.03	y	y	y	
119	17.94	15.77±0.08 ^b	15.00±0.08 ^b	14.66±0.09 ^b	y	y	?	b
120	17.48	15.68±0.07 ^b	15.28±0.09 ^b	14.75±0.11 ^b	y	y	?	
123	19.49	15.79±0.08 ^b	15.41±0.11 ^b	15.26±0.15 ^b	y	N	?	
125	17.48	16.01±0.09 ^b	15.52±0.11 ^b	15.21±0.15 ^b	N	y?	?	
126	18.79	15.94±0.09 ^b	15.57±0.12 ^b	15.22±0.02	y	y?	?	b
127	19.98	16.48±0.16 ^b	16.18±0.23 ^b	15.49±0.19 ^b	y	y	?	
128	17.50	15.57±0.07 ^b	14.89±0.08 ^b	14.62±0.09 ^b	y	y	?	