



## LJMU Research Online

**Conceicao, MS, Traina Chacon-Mikahil, MP, Telles, GD, Libardi, CA, Mendes Junior, EM, Vechin, FC, Lugnani De Andrade, AL, Gaspari, AF, Brum, PC, Cavaglieri, CR, Serag, S, Spiegelman, BM, Hawley, JA and Camera, DM**

**Attenuated PGC-1 alpha Isoforms following Endurance Exercise with Blood Flow Restriction**

<http://researchonline.ljmu.ac.uk/id/eprint/5434/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Conceicao, MS, Traina Chacon-Mikahil, MP, Telles, GD, Libardi, CA, Mendes Junior, EM, Vechin, FC, Lugnani De Andrade, AL, Gaspari, AF, Brum, PC, Cavaglieri, CR, Serag, S, Spiegelman, BM, Hawley, JA and Camera, DM (2016) Attenuated PGC-1 alpha Isoforms following Endurance Exercise with**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>



1 Running Title: PGC-1 $\alpha$  isoform and blood flow restriction

2

3 Attenuated PGC-1 $\alpha$  isoforms following endurance exercise with blood flow restriction

4

5 Miguel S. Conceição<sup>1</sup>; Mara P.T Chacon-Mikahil<sup>1</sup>; Guilherme D. Telles<sup>1</sup>; Cleiton A.

6 Libardi<sup>2</sup>; Edson M. M. Junior<sup>1</sup>; Felipe C. Vechin<sup>3</sup>; André L. Andrade<sup>1</sup>; Arthur F.

7 Gáspari<sup>1</sup>; Patrícia C. Brum<sup>3</sup>; Cláudia R. Cavaglieri<sup>1</sup>; Sara Serag<sup>4</sup>; Bruce M.

8 Spiegelman<sup>4</sup>; John A. Hawley<sup>5,6</sup>; Donny M. Camera<sup>5</sup>.

9

10 <sup>1</sup>Faculty of Physical Education, University of Campinas – Campinas/Brazil; <sup>2</sup>Laboratory

11 of Neuromuscular Adaptations to Resistance Training, Department of Physical

12 Education, Federal University of São Carlos – São Carlos/Brazil; <sup>3</sup>School of Physical

13 Education and Sport, University of São Paulo, São Paulo/Brazil; <sup>4</sup>Department of Cell

14 Biology, Dana-Farber Cancer Institute, Harvard Medical School, Boston/USA; <sup>5</sup>Mary

15 MacKillop Institute for Health Research, Centre for Exercise and Nutrition, Australian

16 Catholic University, Melbourne, Australia; <sup>6</sup>Research Institute for Sport and Exercise

17 Sciences, Liverpool John Moores University, Liverpool, United Kingdom

18

19 Author for correspondence

20 Miguel S. Conceição

21 Faculty of Physical Education, University of Campinas

22 Av. Érico Veríssimo, 701, Cidade Universitária “Zeferino Vaz” Barão Geraldo,

23 Campinas, São Paulo, Brazil. CEP: 13083-851

24 Email: conceicao.miguel0106@gmail.com

25 Phone: + 55 19 35216625

26 **ABSTRACT**

27 Introduction: Exercise performed with blood flow restriction simultaneously enhances  
28 the acute responses to both myogenic and mitochondrial pathways with roles in training  
29 adaptation. We investigated isoform-specific gene expression of the peroxisome  
30 proliferator-activated receptor gamma (PPARG) coactivator 1 and selected target genes  
31 and proteins regulating skeletal muscle training adaptation. Methods: 9 healthy,  
32 untrained males participated in a randomized, counter-balanced, cross-over design in  
33 which each subject completed a bout of low-intensity endurance exercise performed  
34 with blood flow restriction (15 min cycling at 40% of  $VO_{2peak}$ , BFR-EE), endurance  
35 exercise (30 min cycling at 70% of  $VO_{2peak}$ , EE) or resistance exercise (4 x 10  
36 repetitions of leg press at 70% of 1-repetition maximum, RE) separated by at least one-  
37 week recovery. A single resting muscle biopsy (*vastus lateralis*) was obtained two  
38 weeks before the first exercise trial (rest) and 3 h after each bout. Results: Total PGC-1 $\alpha$   
39 mRNA abundance, along with all four isoforms, increased above rest with EE only  
40 ( $P<0.05$ ) being higher than BFR-EE ( $P<0.05$ ). PGC-1 $\alpha$ 1, 2 and 4 were higher after EE  
41 compared to RE ( $P<0.05$ ). EE also increased VEGF, Hif-1 $\alpha$  and MuRF-1 mRNA  
42 abundance above rest ( $P<0.05$ ) while COXIV mRNA expression increased with EE  
43 compared to BFR-EE ( $P<0.05$ ). Conclusion: The attenuated expression of all four PGC-  
44 1 $\alpha$  isoforms when endurance exercise is performed with blood flow restriction suggests  
45 this type of exercise provides an insufficient stimulus to activate the signaling pathways  
46 governing mitochondrial and angiogenesis responses observed with moderate- to high  
47 intensity endurance exercise.

48

49 Key words: mitochondrial biogenesis; cell signalling; skeletal muscle; adaptation;  
50 angiogenesis; high intensity exercise

## 51 **Introduction**

52           Skeletal muscle is a highly malleable tissue that can alter its phenotype  
53 according to the contractile stimulus imposed (39). For instance, moderate-intensity  
54 (i.e., <65% of peak oxygen uptake [ $\dot{V}O_{2peak}$ ]) endurance exercise training enhances  
55 whole-body  $\dot{V}O_{2peak}$  (4), increases the maximal activities of oxidative enzymes, and  
56 shifts patterns of substrate selection from carbohydrate- to fat-based fuels (18). In  
57 contrast, strenuous (80% of one repetition maximum [1-RM]) resistance exercise has  
58 little or no effects on whole-body  $\dot{V}O_{2peak}$  and oxidative enzyme profiles (11) but  
59 increases myofibrillar protein accretion and muscle cross-sectional area (CSA) (30).

60

61           While resistance and endurance exercise could be considered at opposite ends of  
62 the ‘adaptation continuum’ by virtue of their divergent biochemical and morphological  
63 phenotypes, blood flow restriction during low-intensity endurance exercise (BFR-EE)  
64 improves both  $\dot{V}O_{2peak}$ , muscle strength and CSA (1, 2). Abe and co-workers (1)  
65 reported increased isometric muscle strength, muscle CSA and  $\dot{V}O_{2peak}$  following 8  
66 weeks (24 training sessions) of low-intensity cycle exercise (15 min at 40%  $\dot{V}O_{2peak}$ )  
67 performed with BFR-EE compared to same exercise undertaken without BFR. While  
68 these adaptation responses are considerably lower in magnitude relative to conventional  
69 endurance and resistance training performed at higher intensities, the local hypoxia  
70 induced by BFR appears to induce an additive ‘metabolic stressor’ that perturbs cellular  
71 homeostasis (17) and concomitantly enhances both anabolic and oxidative adaptations.

72

73           The cellular mechanisms mediating adaptation responses to exercise are  
74 complex involving the cross talk of several intracellular signaling systems that  
75 ultimately form the basis for specific phenotypic responses with divergent contractile

76 modes (17). The transcriptional co-activator Peroxisome proliferator-activated receptor  
77 gamma (PPARG) coactivator 1 alpha (PGC-1 $\alpha$ ) is a ‘master regulator’ of many  
78 endurance exercise-induced adaptations by virtue of its central role in promoting  
79 mitochondrial biogenesis, angiogenesis, and inflammatory proteins (17). Transcription  
80 of the PGC-1 $\alpha$  gene has been shown to be under the control of several promoter regions  
81 with activation of the alternative PGC-1 $\alpha$ 1 promoter resulting in the transcription of  
82 three additional isoforms: PGC-1 $\alpha$ 2, - $\alpha$ 3 and - $\alpha$ 4. Ruas and colleagues (32) recently  
83 demonstrated a preferential increase in the PGC1- $\alpha$ 4 isoform following resistance  
84 exercise in human skeletal muscle. However, little is known about the regulation of the  
85  $\alpha$ 2 and  $\alpha$ 3 isoforms and, to date, no studies have investigated the expression of all four  
86 PGC-1 $\alpha$  isoforms to diverse contractile stimuli such as resistance and endurance  
87 exercise, or following BFR, in humans. Accordingly, the aim of the present study was  
88 to compare the acute molecular responses mediated by the different PGC-1 $\alpha$  isoforms  
89 following low intensity endurance exercise (BFR-EE), resistance exercise (RE) and  
90 moderate endurance exercise (EE). As BFR-EE can promote both endurance capacity  
91 and muscle hypertrophy responses, we hypothesised EE and RE would selectively  
92 increase the expression of the PGC-1 $\alpha$ 1 and  $\alpha$ 4 isoforms, respectively. In contrast, we  
93 hypothesized that BFR-EE would upregulate a molecular signature involving the  
94 increase of both isoforms and their respective anabolic and mitochondrial gene targets.

95

## 96 **METHODS**

### 97 *Subjects*

98           Nine untrained, healthy male subjects [age  $22.4 \pm 3.0$  yr, body mass (BM)  $73.5 \pm$   
99  $9.7$  kg, height  $1.79 \pm 0.05$  m, maximal oxygen uptake test ( $VO_{2peak}$ )  $36.8 \pm 4.8$  mLkg<sup>-1</sup>  
100  $\cdot$ min<sup>-1</sup>, leg press one repetition maximum (1-RM)  $266 \pm 66$  kg; values are mean  $\pm$  SD]

101 voluntarily participated in this study. The experimental procedures and possible risks  
102 associated with the study were explained to all subjects, who provided written informed  
103 consent before participation. The study was approved by the local University's Ethics  
104 Committee and conducted in conformity with the policy statement regarding the use of  
105 human subjects according to the latest revision of the *Declaration of Helsinki*.

106

### 107 *Experimental Design*

108 The study employed a randomized counter-balanced, cross-over design in which  
109 each subject completed a bout of either resistance exercise (RE), endurance cycling  
110 exercise (EE) or low-intensity cycling exercise combined with blood flow restriction  
111 (BFR-EE). Two weeks prior to the first exercise session, a resting muscle biopsy was  
112 obtained before participants underwent  $VO_{2peak}$  and one-repetition maximum (1-RM)  
113 testing, and exercise familiarization. Exercise trials were separated by a one-week  
114 recovery period during which time subjects maintained their habitual diet and physical  
115 activity patterns.

116

### 117 *Preliminary Testing*

118  $VO_{2peak}$ . Participants performed a maximum graded exercise test on a cycle ergometer  
119 with electromagnetic braking (Quinton modelo: Corival 400, Lode BV, Groningen,  
120 Netherlands) based on a protocol used in previously published paper that investigated  
121 BFR-EE (1). Briefly, after resting on the bike for 5 min, participants commenced the  
122 incremental test protocol. Briefly, subjects commenced cycling at an initial load of 50  
123 W for 1 min and the workload was increased by 15 W/min until a workload of 200 W  
124 was reached, after which further increases were 10 W/min increments. The test  
125 continued until voluntary exhaustion, defined by two of the three following criteria:

126  $\text{VO}_{2\text{peak}}$  plateau ( $< 2.1 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  of variation),  $> 1.10$  respiratory exchange ratio,  
127 and/or heart rate higher than 90% of maximum estimated from age (19). Gas exchange  
128 data were collected continuously using an automated breath-by-breath metabolic system  
129 (CPX, Medical Graphics, St. Paul, Minnesota, USA) and the highest oxygen  
130 consumption value was defined as the peak oxygen consumption ( $\text{VO}_{2\text{peak}}$ ) over any 30  
131 sec period. To confirm the appropriateness of this protocol for this study we performed  
132 a pilot study to verify repeatability in  $\text{VO}_{2\text{peak}}$  measures and observed a strong  
133 repeatability in  $\text{VO}_{2\text{peak}}$  (3.0%), power (1.9%), respiratory exchange ratio (RER) (5.6%),  
134 and time to exhaustion (1.6%) measures.

135

### 136 *Maximal Strength*

137         The one-repetition maximum (1-RM) test was performed on a leg press machine  
138 (45° leg press, G3-PL70; Matrix, São Paulo, Brazil) as previously described (8). Briefly,  
139 participants performed a 5 min warm-up on a cycle ergometer riding at 25 W.  
140 Participants then undertook 1 x 10 repetitions at 50% of their estimated 1-RM, followed  
141 by 1 x 3 repetitions at 70% of the estimated 1-RM with 1-min rest between sets.  
142 Participants then performed a series of single repetitions until the maximum load (1-  
143 RM) lifted was established with fully eccentric-concentric movement with 90° range of  
144 motion. Repetitions were separated by a 3-min recovery and were used to establish the  
145 maximum load/weight that could be moved through the full range of motion once, but  
146 not a second time.

147

### 148 *Diet/Exercise Control*

149         Before each experimental trial (described subsequently), subjects were instructed  
150 to refrain from exercise training and vigorous physical activity, and alcohol and caffeine



151 consumption for a minimum of 48 h. Subjects were provided with standardized  
152 prepacked meals that consisted of 3 g carbohydrate/kg body mass (BM), 0.5 g  
153 protein/kg BM, and 0.3 g fat/kg body mass consumed as the final caloric intake the  
154 evening before reporting for an experimental trial.

155

#### 156 Experimental Testing Sessions

157 On the morning of an experimental trial, subjects reported to the laboratory after a ~10-  
158 h overnight fast. After resting in the supine position for ~15 min and under local  
159 anaesthesia (2–3 mL of 1% Xylocaine), a resting biopsy was obtained from the *vastus*  
160 *lateralis* using a 5-mm Bergstrom needle modified with suction (7). Approximately 100  
161 mg of muscle was removed, dissected free from blood and connective tissue and snap  
162 frozen in liquid nitrogen before being stored at – 80°C until subsequent analyses. Due to  
163 ethical constraints regarding the total number of muscle biopsies allowed, this single  
164 resting biopsy was used as a basal control for all subsequent exercise trials. Two weeks  
165 later participants returned to the laboratory having (after the same pre-trial diet and  
166 exercise control) to undertake the first of three randomly assigned exercise sessions  
167 (described below). Each exercise trial was separated by a one week wash out. Following  
168 the completion of each exercise session, subjects rested for 180 min after which time a  
169 muscle biopsy was obtained. Subsequent incisions were performed 3 cm proximal to  
170 each other. Blood samples were collected before each exercise session and immediately,  
171 1, 2 hr and 3 hr post exercise. Blood samples were immediately placed in microtubes  
172 containing 1% sodium fluoride and then centrifuged at 3000 rpm for 5 min to separate  
173 the plasma before being aliquoted and frozen in liquid nitrogen and stored at -80°C.

174

#### 175 *Resistance Exercise (RE)*

176 After a standardized warm-up on a cycle ergometer consisting of 5 min light  
177 cycling at 25 W, subjects performed 4 sets of 10 repetitions leg press exercise (45° leg  
178 press machine; G3-PL70; Matrix) at 70% of 1-RM. Each set was separated by a 1 min  
179 recovery period during which time subjects remained seated on the leg press machine.  
180 Complete concentric/eccentric movements were performed with 90° of range of motion  
181 and strong verbal encouragement was provided during each set. The volume and  
182 intensity of this session was based on the recommendations of American College Sports  
183 Medicine (ACSM) (3). All participants completed every repetition from each respective  
184 set.

185

#### 186 *Endurance Exercise (EE)*

187 Following a standardized warm up (described previously), subjects performed  
188 30 min of continuous cycling at a power output that elicited ~at 70% of individual  
189  $VO_{2peak}$ . Subjects were fan-cooled and provided visual feedback for pedal frequency,  
190 power output, and elapsed time were provided to subjects. The volume and intensity of  
191 this session were based on the recommendations of ACSM (4). All participants  
192 completed the full 30 min session.

193

#### 194 *Low Intensity Blood Flow Restriction (BFR-EE)*

195 Subjects performed 15 min continuous cycling with a cuff strapped over the  
196 thigh at a power output that elicited at 40% of  $VO_{2peak}$ , as previously reported (1). An  
197 18-cm wide cuff was placed on the proximal portion of the thigh (inguinal fold region)  
198 over the tibial artery and once in position, was inflated until an absence of auditory  
199 blood pulse detected through auscultation with a vascular Doppler probe (DV-600;  
200 Marted, São Paulo, Brazil). Pressure was then slowly released until the first arterial

201 pulse was detected which was considered the systolic pressure at the tibial artery. Cuff  
202 pressure was set at 80% of the maximum tibial arterial pressure and the cuff was  
203 inflated through-out the entire exercise session (22).

204

## 205 *Analytical Procedures*

### 206 *Blood Lactate*

207 Plasma lactate concentration was measured on a spectrophotometer (ELx800,  
208 Biotek, Winooski, USA) using a commercial kit (Biotechnica, Varginha, Brazil)  
209 according to the manufacturer's protocol.

210

### 211 *RNA Extraction and Quantification*

212 Approximately 20 mg of skeletal muscle was homogenized in TRIzol with  
213 chloroform added to form an aqueous RNA phase. This RNA phase was then  
214 precipitated by mixing with ice-cold isopropanol alcohol and the resulting pellet was  
215 washed and re-suspended in 40  $\mu$ l of RNase-free water. Extracted RNA was quantified  
216 using a NanoDrop 1000 spectrophotometer (Nanodrop Technologies, Wilmington,  
217 USA) by measuring absorbance at 260 nm and 280 nm.

218

### 219 *Reverse Transcription*

220 First-strand complementary DNA (cDNA) synthesis was performed using  
221 commercially available TaqMan Reverse Transcription Reagents (Invitrogen,  
222 Melbourne, Australia) in a final reaction volume of 20  $\mu$ L. All RNA and negative  
223 control samples were reverse transcribed to cDNA in a single run from the same reverse  
224 transcription master mix. Serial dilutions of a template human skeletal muscle RNA  
225 (AMBION; Cat No AM7982) was included to ensure efficiency of reverse transcription

226 and for calculation of a standard curve for real-time quantitative polymerase chain  
227 reaction (RT-PCR).

228

### 229 *Real-Time PCR*

230 Quantification (in duplicate) of mRNA was performed using a CFX96 Touch™  
231 Real-Time PCR Detection System (Bio Rad, California, USA). Taqman-FAM-labelled  
232 primer/probes for MuRF-1 (Cat No. Hs00822397\_m1), COXIV (Cat No.  
233 Hs00971639\_m1), IL-6 (Cat No. Hs00985639\_m1), PGC-1α (Cat No.  
234 Hs01016719\_m1), HIF-1α (Cat No. Hs00153153\_m1), Myostatin (Hs00976237\_m1),  
235 IGF-1 (Hs01547656\_m1) and VEGF (Cat No. Hs00900055\_m1) were used in a final  
236 reaction volume of 20 μL. PCR treatments were 2 min at 50 °C for UNG activation, 10  
237 min at 95 °C then 40 cycles of 95 °C for 15 s and 60 °C for 60s. Glyceraldehyde-3-  
238 phosphate dehydrogenase (GAPDH) (Cat No Hs02758991\_g1) was used as a  
239 housekeeping gene and was stably expressed between exercise interventions (data not  
240 shown). The relative amounts of mRNAs were calculated using the relative  
241 quantification ( $\Delta\Delta\text{CT}$ ) method (24). All Taqman-based PCR experiments were  
242 performed in the Centre for Exercise and Nutrition laboratory at the Australian Catholic  
243 University.

244

### 245 *Quantification of PGC-1α isoforms*

246 RNA was extracted from a separate piece of snap frozen muscle (~20 mg) using TRIzol  
247 (Invitrogen) and purified using QIAGEN RNeasy mini-columns. Reverse transcription  
248 was performed using a High Capacity cDNA Reverse Transcription kit (Applied Bio-  
249 systems). Real-Time Quantitative PCR was carried out in a SYBR Green ER PCR  
250 Master Mix (Invitrogen)/ 384-well format using an ABI PRISM 7900HT (Applied

251 Biosystems). Relative mRNA levels were calculated using the comparative CT method  
252 and normalized to cyclophilin mRNA. Primer sequences are as follows: Cyclophilin  
253 (forward: GGAGATGGCACAGGAGGAA; reverse: GCCCGTAGTGC TTCAGTTT),  
254 PGC1 $\alpha$ 1 (forward: ATG GAG TGA CAT CGA GTG TGC T; reverse: GAG TCC ACC  
255 CAG AAA GCT GT), PGC1 $\alpha$ 2 (forward: AGT CCA CCC AGA AAG CTG TCT;  
256 reverse: ATG AAT GAC ACA CAT GTT GGG), PGC1 $\alpha$ 3 (forward: CTG CAC CTA  
257 GGA GGC TTT ATG C; reverse: CAA TCC ACC CAG AAA GCT GTC T), and  
258 PGC1 $\alpha$ 4 (forward: TCA CAC CAA ACC CAC AGA GA; reverse: CTG GAA GAT  
259 ATG GCA CAT). All SYBR Green-based PCR experiments were performed in the  
260 Department of Cell Biology laboratory at the Dana-Farber Cancer Institute, Harvard  
261 Medical School (USA).

262

### 263 *Western Blots*

264       Approximately 30 mg of muscle was homogenized in buffer containing 50 mM  
265 Tris·HCl pH 7.5, 1 mM EDTA, 1 mM EGTA, 10% glycerol, 1% Triton X-100, 50 mM  
266 NaF, 5 mM sodium pyrophosphate, 1 mM DTT, 10%  $\mu$ g/ml trypsin inhibitor, 2  $\mu$ g/ml  
267 aprotinin, 1mM benzamidine, and 1 mM PMSF. After determination of protein  
268 concentration (Pierce, Rockford, IL), lysate was resuspended in Laemmli sample buffer.  
269 Lysate was then re-suspended in Laemmli sample buffer with 40  $\mu$ g of protein loaded  
270 onto 4–20% Mini-PROTEAN TGX Stain-Free™ Gels (Bio Rad, California, USA). Post  
271 electrophoresis gels were activated according to the manufacturer's details (Chemidoc,  
272 Bio-Rad, Gladesville, Australia) and then transferred to polyvinylidene fluoride (PVDF)  
273 membranes. After transfer, a Stain-Free image of the PVDF membranes for total protein  
274 normalization was obtained before membranes were rinsed briefly in distilled water and  
275 blocked with 5% non-fat milk, washed with 10 mM of Tris–HCl, 100 mM of NaCl, and

276 0.02% Tween 20, and incubated with primary antibody (1:1000) overnight at 4 °C.  
277 Membranes were incubated with secondary antibody (1:2,000), and proteins were  
278 detected via chemiluminescence (Amersham Biosciences, Buckinghamshire, UK;  
279 Pierce Biotechnology, Rockford, IL) and quantified by densitometry. All sample time  
280 points for each subject were run on the same gel. Polyclonal anti-phospho-mTOR<sup>Ser2448</sup>  
281 (no. 2971), -p70 S6K<sup>Thr389</sup> (no. 9206), - adenosine monophosphate kinase (AMPK)<sup>Thr172</sup>  
282 (no. 2531), eukaryotic translation initiation factor 4E-binding protein 1 (4E-BP1) 4E-  
283 BP1<sup>Thr37/46</sup> (no. 9459), eEF2 eukaryotic translation elongation factor 2 (eEF2) eEF2<sup>Thr56</sup>  
284 (no. 2331) and p53<sup>Ser15</sup> (no. 9284) were purchased from Cell Signaling Technology  
285 (Danvers, MA, USA). Volume density of each target protein band was normalized to  
286 the total protein loaded into each lane using stain-free technology (15) with data  
287 expressed in arbitrary units. Due to low sample availability, phosphorylated proteins  
288 were unable to be normalized to their respective total protein content and were therefore  
289 also normalized to stain-free protein levels.

290

### 291 **Statistical analysis**

292 Statistical analysis was performed using SAS version 9.3 for Windows (SAS  
293 Institute Inc., Cary, NC, USA). Data normality and variance equality were assessed  
294 through the Shapiro-Wilk and Levene tests. One-way ANOVA with repeated measures  
295 (factor: condition) was performed for gene and protein expression analyses. A mixed  
296 model ANOVA, assuming group and time as fixed factors and subjects as a random  
297 factor, was performed for blood lactate data. Tukey post hoc analysis was used for  
298 multiple comparison purposes when significant F-values were found. The significance  
299 level was set at  $P \leq 0.05$ . Data are presented as Mean  $\pm$  Standard Deviation (SD).

300

301 **Results**

302 **mRNA expression**

303 **Total PGC1- $\alpha$  and isoforms**

304 Total PGC-1 $\alpha$  mRNA (Figure 1A) increased with EE above rest ( $P < 0.0001$ ), RE  
305 ( $P = 0.0013$ ) and BFR-EE ( $P > 0.0001$ ). There was a significant increase in PGC-1 $\alpha$ 1  
306 mRNA with EE above rest ( $P = 0.0450$ ), RE ( $P = 0.0069$ ) and BFR-EE ( $P = 0.0349$ )  
307 (Figure 1B). There was also a significant increase in PGC-1 $\alpha$ 2 mRNA (Figure 1C) with  
308 EE above rest ( $P < 0.0001$ ), RE ( $P = 0.0003$ ) and BFR-EE ( $P < 0.0001$ ). PGC-1 $\alpha$ 3 mRNA  
309 (Figure 1D) increased with EE above rest ( $P = 0.0389$ ). There was also increases PGC-  
310 1 $\alpha$ 4 mRNA (Figure 1E) with EE above rest ( $P = 0.0035$ ), RE ( $P = 0.0469$ ) and BFR-EE  
311 ( $P = 0.0140$ ).

312

313 **\*\*Figure 1 here\*\***

314

315 **VEGF, COXIV, HIF-1a**

316 There was a significant increase in VEGF mRNA (Figure 2A) with EE above  
317 rest ( $P = 0.0180$ ) and RE ( $P = 0.0069$ ). COXIV mRNA expression increased with EE  
318 above BFR-EE ( $P = 0.0550$ ) (Figure 2B). There was a significant increase in HIF-1a  
319 abundance with EE above Rest ( $P = 0.0530$ ) (Figure 2C).

320

321 **\*\*Figure 2 here\*\***

322

323 **IL-6 – IGF-1 - Myostatin - MurRF1**

324 IL-6, IGF-1 and Myostatin mRNA expression were unchanged post-exercise  
325 (Figure 3 A, B, C). There was a post-exercise increase in MuRF1 mRNA abundance

326 with EE above Rest (P=0.0003), RE (P=0.0256) and BFR-EE (P=0.0007) (Figure 3D).

327

328 **\*\*Figure 3 here\*\***

329

### 330 **Cell Signaling**

#### 331 **mTOR -p70S6K -4E-BP1 -eEF2**

332 There were no changes in mTOR<sup>Ser2448</sup>, p70S6K<sup>Thr389</sup>, 4E-BP1<sup>Thr37/46</sup> or eEF2<sup>Thr56</sup>  
333 phosphorylation post-exercise or between exercise groups (Figure 4).

334

335 **\*\*Figure 4 here\*\***

336

#### 337 **AMPK -p53**

338 AMPK<sup>Thr156</sup> and p53<sup>Ser15</sup> phosphorylation were unchanged post-exercise (Figure  
339 5).

340

341 **\*\*Figure 5 here\*\***

342

#### 343 **Plasma lactate concentration**

344 Lactate concentration increased above rest immediately post-exercise for all  
345 interventions (P<.0001 for all comparisons; Table 1). Lactate concentration remained  
346 elevated at 1 h, 2 h and 3 h post-exercise for EE and RE, and 1 h and 2 h for BFR-EE  
347 (P<.0001 for all comparisons).

348



349

**\*\*Table 1 here\*\***

350

## 351 **Discussion**

352           Low intensity (<50% of  $\text{VO}_{2\text{peak}}$ ) endurance training with blood flow restriction  
353 has been shown to concomitantly promote isometric muscle strength, muscle CSA and  
354  $\text{VO}_{2\text{peak}}$  (1, 2). While these enhanced adaptation responses are considerably lower in  
355 magnitude compared to conventional resistance or endurance exercise performed  
356 without any blood flow restriction, the underlying molecular mechanisms mediating  
357 these responses remain largely undefined. For the first time we report that low intensity  
358 endurance cycling exercise performed with blood flow restriction failed to increase  
359 PGC-1 $\alpha$  expression to that commonly observed with ‘conventional’ endurance exercise.  
360 Moreover, we show isoform-specific post-exercise increases in the  $\alpha 4$  isoform along  
361 with Hif-1 $\alpha$  and VEGF mRNA expression following higher intensity endurance  
362 exercise without blood flow restriction. Taken collectively, our novel findings suggest  
363 that cycle exercise undertaken with blood flow restriction is unable to provoke the  
364 perturbations to cellular homeostasis necessary to induce activation of the cell signaling  
365 events regulating mitochondrial biogenesis and angiogenesis that take place with higher  
366 intensity endurance exercise without blood flow restriction.

367

368 A growing body of evidence suggests that exercise undertaken with blood flow  
369 restriction can enhance exercise adaptation. A recent meta-analysis reported both low  
370 load/intensity resistance (20–30% 1 RM) and aerobic walking exercise performed with  
371 blood flow restriction can induce increases in muscle strength and hypertrophy,  
372 although with smaller gains compared to high intensity resistance exercise alone (35).  
373 However, little is known about the molecular mechanisms mediating these responses

374 when low intensity endurance exercise is undertaken with blood flow restriction. As  
375 such, we compared the expression of key gene and protein targets implicated in a range  
376 of exercise adaptation responses such as hypertrophy, mitochondrial biogenesis, muscle  
377 proteolysis, substrate metabolism and angiogenesis between BFR-EE, and conventional  
378 bouts of RE and EE. We particularly focused on the four different full-length PGC-1 $\alpha$   
379 isoforms putatively implicated in anabolic and mitochondrial-related adaptation  
380 responses.

381

382 In agreement with previous studies (5, 23, 29), we observed significant increases in total  
383 PGC-1 $\alpha$  mRNA following continuous endurance exercise performed at 70% of VO<sub>2peak</sub>.  
384 This increase in PGC-1 $\alpha$  mRNA was concomitant with greater abundance of VEGF, a  
385 target of PGC-1 $\alpha$  (37). However, in contrast to our original hypothesis, this response  
386 was absent following a bout of low-intensity endurance exercise (40% VO<sub>2peak</sub>)  
387 performed with blood flow restriction. In an attempt to identify possible mechanisms  
388 responsible for this attenuated PGC-1 $\alpha$  response, we investigated IL-6 expression to  
389 determine whether an increase in the muscular inflammatory program was implicated in  
390 the blunted response. This hypothesis was based on previous data showing an inverse  
391 relationship between skeletal muscle PGC-1 $\alpha$  and IL-6 expression (16). However, IL-6  
392 mRNA expression post-exercise was unchanged in all exercise groups suggesting any  
393 acute increase in muscle inflammation caused by BFR-EE was not responsible for the  
394 reduced PGC-1 $\alpha$  expression observed. We also investigated other cellular markers  
395 implicated in exercise adaptation responses that can regulate PGC-1 $\alpha$  expression.  
396 AMPK is an intracellular 'fuel gauge' that can phosphorylate PGC-1 $\alpha$  and increase its  
397 transcriptional activity (36) while the apoptogenic protein p53 has emerged as another  
398 signaling regulator of skeletal muscle exercise-induced mitochondrial biogenesis and

399 substrate metabolism that can translocate to the nucleus upon activation and induce  
400 PGC-1 $\alpha$  expression (17). Phosphorylation of either of these protein targets was  
401 unaltered post-exercise suggesting other molecular markers and/or physiological  
402 mechanisms may be responsible for the upregulation of PGC-1 $\alpha$  with high intensity  
403 endurance exercise. One plausible explanation for these discrepant findings may be the  
404 level of glycogen utilization between exercise sessions in our untrained subjects. We  
405 (10) and others (6, 31) have shown greater post-exercise PGC-1 $\alpha$  expression with low-  
406 compared to normal or high glycogen concentration and although we did not measure  
407 muscle glycogen use in the current study due to limited muscle tissue availability, the  
408 longer duration and higher intensity exercise bout is likely to have induced greater  
409 glycogen depletion compared to the endurance exercise session performed with blood  
410 flow restriction.

411

412 Another possible explanation for the discrepancy in PGC-1 $\alpha$ 1 expression between the  
413 two endurance-based exercise bouts is the large differences in estimated energy  
414 expenditure. Exercise energy expenditure after BFR-EE was ~4 fold less compared to  
415 the EE protocol with total energy expenditure positively associated with PGC-1 $\alpha$   
416 expression ( $r=0.73$ ,  $P=0.039$ ). Increased PGC-1 $\alpha$  mRNA expression has been observed  
417 after 30 min running compared to bouts of 20 and 10 min (37). Thus, total exercise-  
418 induced energy expenditure may be an overriding determinant of PGC-1 $\alpha$  expression  
419 responses post-exercise.

420

421 Low intensity endurance exercise with BFR was also unable to induce the expression of  
422 PGC-1 $\alpha$ 4 compared to higher intensity endurance exercise without blood flow  
423 restriction. The PGC-1 $\alpha$ 4 isoform has been proposed to promote muscle hypertrophy by

424 inducing IGF-1 expression and reducing the expression of myostatin, a negative  
425 regulator of muscle growth (32). The increase in PGC-1 $\alpha$ 4 mRNA expression with EE  
426 in the current study was mirrored by a small, non-significant, increase and decrease in  
427 IGF1 and myostatin expression, respectively. Ruas and colleagues were the first to  
428 show a selective increase in PGC-1 $\alpha$ 4 expression (concomitant with decreased  
429 myostatin abundance) with resistance compared to endurance exercise in human skeletal  
430 muscle (32). However, this expression pattern was observed following 8 weeks whole-  
431 body resistance training. Thus, a limitation of our study is that we only incorporated a  
432 single bout of isolated leg press suggesting longer training programs/ exercise stimulus  
433 may be required to induce this selective PGC-1 $\alpha$ 4 response. Nonetheless, another recent  
434 publication reported increased truncated and non-truncated PGC-1 $\alpha$  transcripts from  
435 both alternative and proximal promoter sites 2 hours following an acute bout of  
436 resistance exercise that incorporated the same volume and intensity as our study (40).  
437 This indicates the resistance exercise bout performed in our study was likely sufficient  
438 to induce the appropriate signal to increase the expression of this isoform however  
439 potential differences in post-exercise biopsy timing between this study and ours (2 h vs.  
440 3 h) may explain why we did not observe this increase with resistance exercise.

441

442 Increased PGC-1 $\alpha$ 4 and VEGF expression has also been reported in primary myotubes  
443 treated under hypoxic conditions suggesting low oxygen conditions to be favorable for  
444 the activation of this isoform (38). In the current study, the transcription factor Hif-1 $\alpha$ , a  
445 key regulator of angiogenesis in situations of hypoxia (34), was unchanged following  
446 BFR-EE, while RE and EE induced 2-fold higher post-exercise changes in lactate  
447 compared to BFR-EE. While it is possible a greater metabolic and hypoxic stimulus  
448 may be required to increase PGC-1 $\alpha$ 4 signaling, others have reported unchanged blood

449 lactate following aerobic-based exercise with blood flow restriction (26). Moreover, the  
450 same occlusion protocol (15 min cycle at 40%  $\text{VO}_{2\text{peak}}$ ) has been shown to improve  
451 muscle volume and  $\text{VO}_{2\text{peak}}$ , during a chronic training intervention (1). Thus, it is  
452 possible chronic exposure to this occlusion stimulus may be required to elicit increases  
453 in PGC-1 $\alpha$ 4 expression. As this is the first study to investigate changes in Hif-1 $\alpha$   
454 following endurance cycling exercise with BFR it is difficult to compare our results to  
455 those of previous investigations incorporating resistance exercise and BFR. However,  
456 we speculate that when performed with blood flow restriction, the lower contractile  
457 intensity associated with ‘conventional’ endurance compared to resistance (or sprint)  
458 exercise, provides adequate blood flow to the exercising musculature and adjoining  
459 capillary beds in order to prevent tissue de-oxygenation. Further studies comparing  
460 different low intensity endurance exercise protocols with resistance exercise that  
461 incorporate blood flow restriction are required to corroborate this hypothesis.

462

463 Another novel finding from the current study was the post-exercise increases in the  
464 PGC-1 $\alpha$ 2 and 3 isoforms. Similar to the  $\alpha$ 1 and  $\alpha$ 4 isoforms, both PGC-1 $\alpha$ 2 and  $\alpha$ 3  
465 increased above rest with higher intensity endurance exercise and were significantly  
466 elevated compared to resistance exercise. Both isoforms are expressed in skeletal  
467 muscle and brown adipose tissue although little is known about the regulatory targets of  
468 these isoforms and their capacity to mediate exercise adaptation responses (27). Based  
469 on the elevated response following endurance compared to resistance exercise, we  
470 propose these isoforms to mediate physiological processes related to mitochondrial  
471 biogenesis and substrate metabolism.

472

473 Considering low load endurance exercise with BFR can increase muscle strength and  
474 hypertrophy (35), we also investigated markers of translation initiation, elongation and  
475 muscle proteolysis. Previous studies have reported increases in mTOR and p70S6K  
476 phosphorylation that have formed the basis for enhanced rates of muscle protein  
477 synthesis following resistance exercise with blood flow restriction (13, 14).  
478 Nonetheless, the phosphorylation status of these proteins as well as 4E-BP1 and eEF2  
479 were unchanged 3 h post-exercise in the current study. This is in agreement with the  
480 results of Ozaki and colleagues (28) who observed no changes in Akt, mTOR or  
481 p70S6K phosphorylation following 20 min treadmill walking performed with blood  
482 flow restriction despite a higher intensity exercise bout (55%  $\text{VO}_{2\text{peak}}$ ) compared to our  
483 protocol. While our study design was somewhat limited by only having the single post-  
484 exercise biopsy (9), this sampling time-point was specifically chosen based on previous  
485 studies showing significant, and in some cases maximal, increases in PGC-1 $\alpha$  mRNA  
486 expression in response to an exercise challenge (5, 23). Future studies investigating  
487 endurance exercise undertaken with BFR-EE should include a time-course of signaling  
488 responses in order to determine the optimal 'window' for muscle sampling in  
489 subsequent investigations.

490

491 Several other factors including the width and pressure of cuff used during BFR must  
492 also be considered. Previous studies have reported smaller increases in muscle CSA  
493 when lower body resistance training is undertaken with BFR (compared to no BFR) at  
494 the site of the cuff (12, 20). While this indicates a narrow cuff may be advantageous for  
495 promoting anabolic adaptation responses due to compressing less muscle tissue, a recent  
496 study comparing the effects of a wide versus narrow cuff reported similar increases in  
497 maximum strength and muscle cross sectional area following 12 weeks of unilateral

498 elbow flexion performed at 20% of 1RM (21). Also, a recent study showed that there  
499 was no difference in either muscle strength or hypertrophy between different occlusion  
500 pressures (25). Thus, the use of a wider cuff, as used in our protocol, appears unlikely to  
501 attenuate chronic muscle anabolic responses. Regardless, these studies are currently  
502 only limited to BFR with resistance exercise. Future studies comparing these parameters  
503 when endurance exercise is performed with BFR are required. Finally, MuRF-1 mRNA  
504 expression increased post endurance exercise which resulted in a higher expression  
505 above endurance exercise with BFR and resistance exercise. MuRF-1 mediates the  
506 ubiquitin proteasome system by 'labelling' cleaved myofibril segments for degradation  
507 (33). It is unclear whether this increase in expression with high intensity endurance  
508 exercise represents general tissue remodeling, particularly considering our participants  
509 were untrained and the unaccustomed contractile stimulus, or a greater induction of  
510 protein degradation.

511

512 In summary, this is the first study to investigate the molecular mechanisms mediating  
513 muscle adaptation responses to low intensity endurance cycling exercise with blood  
514 flow restriction. The attenuated expression of all four PGC-1 $\alpha$  isoforms when endurance  
515 exercise is performed with blood flow restriction suggests this type of exercise is unable  
516 to induce the appropriate metabolic perturbation capable of activating the cell signaling  
517 machinery responsible for mitochondrial biogenesis and angiogenesis responses with  
518 moderate-to-high intensity endurance exercise. Longer training programs incorporating  
519 endurance exercise with BFR that correlate measurements of these molecular markers  
520 with functional adaptation responses such as changes in  $VO_{2peak}$  and cycle time to  
521 fatigue will yield important information to the efficacy of this training method to

522 enhance training adaptation and subsequently improve health outcomes in populations  
523 that may be unable to perform, prolonged exercise.

524

## 525 **Acknowledgments**

526           The authors would like to express gratitude for the FAPESP (2014/00985-0) and  
527 FAEPEX for financial support. The writing of this publication was also supported by a  
528 Collaborative Research Networks (CRN) grant awarded to J. A. Hawley (2013000443).  
529 The authors declare no declare no conflicts of interest. The results of the present study  
530 do not constitute endorsement by ACSM.

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546



547 **References**

- 548 1. Abe T, Fujita S, Nakajima T et al. Effects of Low-Intensity Cycle Training with  
549 Restricted Leg Blood Flow on Thigh Muscle Volume and VO<sub>2</sub>MAX in Young  
550 Men. *J Sports Sci Med*. 2010;9(3):452-8.
- 551 2. Abe T, Kearns CF, Sato Y. Muscle size and strength are increased following  
552 walk training with restricted venous blood flow from the leg muscle, Kaatsu-  
553 walk training. *J Appl Physiol (1985)*. 2006;100(5):1460-6.
- 554 3. ACSM. American College of Sports Medicine position stand. Progression  
555 models in resistance training for healthy adults. *Med Sci Sports Exerc*.  
556 2009;41(3):687-708.
- 557 4. ACSM. American College of Sports Medicine position stand. Quantity and  
558 quality of exercise for developing and maintaining cardiorespiratory,  
559 musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance  
560 for prescribing exercise. *Med Sci Sports Exerc*. 2011;43(7):1334-59.
- 561 5. Bartlett JD, Hwa Joo C, Jeong TS et al. Matched work high-intensity interval  
562 and continuous running induce similar increases in PGC-1alpha mRNA, AMPK,  
563 p38, and p53 phosphorylation in human skeletal muscle. *J Appl Physiol (1985)*.  
564 2012;112(7):1135-43.
- 565 6. Bartlett JD, Louhelainen J, Iqbal Z et al. Reduced carbohydrate availability  
566 enhances exercise-induced p53 signaling in human skeletal muscle: implications  
567 for mitochondrial biogenesis. *Am J Physiol Regul Integr Comp Physiol*.  
568 2013;304(6):R450-8.
- 569 7. Bergstrom J. Percutaneous needle biopsy of skeletal muscle in physiological and  
570 clinical research. *Scandinavian journal of clinical and laboratory investigation*.  
571 1975;35(7):609-16.

- 572 8. Brown LE, Weir JP. Procedures recommendation I: Accurate assessment of  
573 muscular strength and power. *Journal of Exercise Physiology Online*. 2001;4:1-  
574 21.
- 575 9. Camera DM, Edge J, Short MJ, Hawley JA, Coffey VG. Early time course of  
576 Akt phosphorylation after endurance and resistance exercise. *Med Sci Sports*  
577 *Exerc*. 2010;42(10):1843-52.
- 578 10. Camera DM, Hawley JA, Coffey VG. Resistance exercise with low glycogen  
579 increases p53 phosphorylation and PGC-1alpha mRNA in skeletal muscle. *Eur J*  
580 *Appl Physiol*. 2015;115(6):1185-94.
- 581 11. Campos GE, Luecke TJ, Wendeln HK et al. Muscular adaptations in response to  
582 three different resistance-training regimens: specificity of repetition maximum  
583 training zones. *Eur J Appl Physiol*. 2002;88(1-2):50-60.
- 584 12. Ellefsen S, Hammarstrom D, Strand TA et al. Blood flow-restricted strength  
585 training displays high functional and biological efficacy in women: a within-  
586 subject comparison with high-load strength training. *Am J Physiol Regul Integr*  
587 *Comp Physiol*. 2015;309(7):R767-79.
- 588 13. Fry CS, Glynn EL, Drummond MJ et al. Blood flow restriction exercise  
589 stimulates mTORC1 signaling and muscle protein synthesis in older men. *J Appl*  
590 *Physiol*. 2010;108(5):1199-209.
- 591 14. Gundermann DM, Walker DK, Reidy PT et al. Activation of mTORC1 signaling  
592 and protein synthesis in human muscle following blood flow restriction exercise  
593 is inhibited by rapamycin. *Am J Physiol Endocrinol Metab*.  
594 2014;306(10):E1198-204.
- 595 15. Gurtler A, Kunz N, Gomolka M et al. Stain-Free technology as a normalization  
596 tool in Western blot analysis. *Anal Biochem*. 2013;433(2):105-11.

- 597 16. Handschin C, Choi CS, Chin S et al. Abnormal glucose homeostasis in skeletal  
598 muscle-specific PGC-1alpha knockout mice reveals skeletal muscle-pancreatic  
599 beta cell crosstalk. *The Journal of clinical investigation*. 2007;117(11):3463-74.
- 600 17. Hawley JA, Hargreaves M, Joyner MJ, Zierath JR. Integrative biology of  
601 exercise. *Cell*. 2014;159(4):738-49.
- 602 18. Holloszy JO, Booth FW. Biochemical adaptations to endurance exercise in  
603 muscle. *Annual review of physiology*. 1976;38:273-91.
- 604 19. Howley ET, Bassett DR, Jr., Welch HG. Criteria for maximal oxygen uptake:  
605 review and commentary. *Med Sci Sports Exerc*. 1995;27(9):1292-301.
- 606 20. Kacin A, Strazar K. Frequent low-load ischemic resistance exercise to failure  
607 enhances muscle oxygen delivery and endurance capacity. *Scand J Med Sci*  
608 *Sports*. 2011;21(6):e231-41.
- 609 21. Laurentino GC, Loenneke JP, Teixeira EL, Nakajima E, Iared W, Tricoli V. The  
610 Effect of Cuff Width on Muscle Adaptations after Blood Flow Restriction  
611 Training. *Med Sci Sports Exerc*. 2015.
- 612 22. Laurentino GC, Ugrinowitsch C, Roschel H et al. Strength training with blood  
613 flow restriction diminishes myostatin gene expression. *Med Sci Sports Exerc*.  
614 2012;44(3):406-12.
- 615 23. Little JP, Safdar A, Bishop D, Tarnopolsky MA, Gibala MJ. An acute bout of  
616 high-intensity interval training increases the nuclear abundance of PGC-1alpha  
617 and activates mitochondrial biogenesis in human skeletal muscle. *Am J Physiol*  
618 *Regul Integr Comp Physiol*. 2011;300(6):R1303-10.
- 619 24. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-  
620 time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*.  
621 2001;25(4):402-8.

- 622 25. Lixandrao ME, Ugrinowitsch C, Laurentino G et al. Effects of exercise intensity  
623 and occlusion pressure after 12 weeks of resistance training with blood-flow  
624 restriction. *Eur J Appl Physiol*. 2015.
- 625 26. Loenneke JP, Thrower AD, Balapur A, Barnes JT, Pujol TJ. Blood flow-  
626 restricted walking does not result in an accumulation of metabolites. *Clin*  
627 *Physiol Funct Imaging*. 2012;32(1):80-2.
- 628 27. Martinez-Redondo V, Pettersson AT, Ruas JL. The hitchhiker's guide to PGC-  
629 1alpha isoform structure and biological functions. *Diabetologia*.  
630 2015;58(9):1969-77.
- 631 28. Ozaki H, Kakigi R, Kobayashi H, Loenneke JP, Abe T, Naito H. Effects of  
632 walking combined with restricted leg blood flow on mTOR and MAPK  
633 signalling in young men. *Acta Physiol (Oxf)*. 2014;211(1):97-106.
- 634 29. Perry CG, Lally J, Holloway GP, Heigenhauser GJ, Bonen A, Spriet LL.  
635 Repeated transient mRNA bursts precede increases in transcriptional and  
636 mitochondrial proteins during training in human skeletal muscle. *J Physiol*.  
637 2010;588(Pt 23):4795-810.
- 638 30. Phillips SM. Physiologic and molecular bases of muscle hypertrophy and  
639 atrophy: impact of resistance exercise on human skeletal muscle (protein and  
640 exercise dose effects). *Appl Physiol Nutr Metab*. 2009;34(3):403-10.
- 641 31. Psilander N, Frank P, Flockhart M, Sahlin K. Exercise with low glycogen  
642 increases PGC-1alpha gene expression in human skeletal muscle. *Eur J Appl*  
643 *Physiol*. 2013;113(4):951-63.
- 644 32. Ruas JL, White JP, Rao RR et al. A PGC-1alpha isoform induced by resistance  
645 training regulates skeletal muscle hypertrophy. *Cell*. 2012;151(6):1319-31.

- 646 33. Sanchez AM, Candau RB, Bernardi H. FoxO transcription factors: their roles in  
647 the maintenance of skeletal muscle homeostasis. *Cellular and molecular life*  
648 *sciences : CMLS*. 2014;71(9):1657-71.
- 649 34. Semenza GL. Regulation of physiological responses to continuous and  
650 intermittent hypoxia by hypoxia-inducible factor 1. *Experimental physiology*.  
651 2006;91(5):803-6.
- 652 35. Slys J, Stultz J, Burr JF. The efficacy of blood flow restricted exercise: A  
653 systematic review & meta-analysis. *J Sci Med Sport*. 2015.
- 654 36. Steinberg GR, Kemp BE. AMPK in Health and Disease. *Physiological reviews*.  
655 2009;89(3):1025-78.
- 656 37. Taylor CW, Ingham SA, Ferguson RA. Acute and chronic effect of sprint  
657 interval training combined with postexercise blood-flow restriction in trained  
658 individuals. *Experimental physiology*. 2016;101(1):143-54.
- 659 38. Thom R, Rowe GC, Jang C, Safdar A, Arany Z. Hypoxic induction of vascular  
660 endothelial growth factor (VEGF) and angiogenesis in muscle by truncated  
661 peroxisome proliferator-activated receptor gamma coactivator (PGC)-1alpha. *J*  
662 *Biol Chem*. 2014;289(13):8810-7.
- 663 39. Wilkinson SB, Phillips SM, Atherton PJ et al. Differential effects of resistance  
664 and endurance exercise in the fed state on signalling molecule phosphorylation  
665 and protein synthesis in human muscle. *J Physiol*. 2008;586(Pt 15):3701-17.
- 666 40. Ydfors M, Fischer H, Mascher H, Blomstrand E, Norrbom J, Gustafsson T. The  
667 truncated splice variants, NT-PGC-1alpha and PGC-1alpha4, increase with both  
668 endurance and resistance exercise in human skeletal muscle. *Physiol Rep*.  
669 2013;1(6):e00140.

671 **Figure 1.** (A) Total Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$  (PGC-  
672 1 $\alpha$ ), (B) Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$  (PGC-1 $\alpha$ 1), (C)  
673 Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$ 2 (PGC-1 $\alpha$ 2), (D)  
674 Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$ 3 (PGC-1 $\alpha$ 3) and (E)  
675 Peroxisome proliferator-activated receptor- $\gamma$  coactivator 1 $\alpha$ 4 (PGC-1 $\alpha$ 4) mRNA  
676 abundance at rest and 3 h post-exercise recovery following endurance exercise (EE),  
677 resistance exercise (RE) or low-intensity associated with blood flow restriction (BFR-  
678 EE). Values are expressed relative to GAPDH and presented in arbitrary units (mean  $\pm$   
679 SD, n=9). a= Significant different from Rest ( $P \leq 0.05$ ); b= Significant different from  
680 HI-RT ( $P \leq 0.05$ ); c= Significant different from BFR-EE ( $P \leq 0.05$ ).

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696 **Figure 2.** (A) Vascular endothelial growth factor (VEGF), (B) Cytochrome c oxidase  
697 subunit 4 isoform 1 (COXIV) and (C) hypoxia-inducible factor-1 alpha (HIF-1 $\alpha$ )  
698 mRNA abundance at rest and 3 h post-exercise recovery following endurance exercise  
699 (EE), resistance exercise (RE) or low-intensity associated with blood flow restriction  
700 (BFR-EE). Values are expressed relative to GAPDH and presented in arbitrary units  
701 (mean  $\pm$  SD, n=9). a= Significant different from Rest ( $P \leq 0.05$ ); b= Significant different  
702 from HI-RT ( $P \leq 0.05$ ); c= Significant different from BFR-EE ( $P \leq 0.05$ ).

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721 **Figure 3.** (A) Interleukin 6 (IL-6), (B) Insulin-like growth factor 1(IGF-1), (C) Muscle  
722 RING finger 1 (MURF1) and (D) Myostatin mRNA abundance at rest and 3 h post-  
723 exercise recovery following endurance exercise (EE), resistance exercise (RE) or low-  
724 intensity associated with blood flow restriction (BFR-EE). Values are expressed relative  
725 to GAPDH and presented in arbitrary units (mean  $\pm$  SD, n=9). a= Significant different  
726 from Rest ( $P \leq 0.05$ ); b= Significant different from HI-RT ( $P \leq 0.05$ ); c= Significant  
727 different from BFR-EE ( $P \leq 0.05$ ).

728

729

730

731

732

733

734

735

736

737

738

739



740 **Figure 4.** (A) Mechanistic target of rapamycin (mTOR)<sup>Ser2448</sup> (B) p70S6K<sup>Thr389</sup> (C)  
741 eukaryotic translation initiation factor 4E-binding protein 1 (4E-BP1)<sup>Thr37/46</sup> eukaryotic  
742 elongation factor 2 (eEF2)<sup>Thr56</sup> phosphorylation in skeletal muscle at rest and after 3 h  
743 post-exercise recovery following endurance exercise (EE), resistance exercise (RE) or  
744 low-intensity associated with blood flow restriction (BFR-EE). Values are normalized  
745 to total protein loaded determined by stain free technology in arbitrary units (mean ±  
746 SD, n=9).

747

748

749

750

751

752

753

754

755

756

757

758 **Figure 5.** (A) Adenosine Monophosphate-Activated Protein (AMPK)<sup>Thr172</sup> and (B)  
759 p53<sup>Ser15</sup> phosphorylation in skeletal muscle at rest and after 3 h post-exercise recovery  
760 following endurance exercise (EE), resistance exercise (RE) or low-intensity associated  
761 with blood flow restriction (BFR-EE). Values are normalized to total protein loaded  
762 determined by stain free technology in arbitrary units (mean  $\pm$  SD, n=9).

763

764

765

766

767