

LJMU Research Online

Anderson, L, Naughton, RJ, Close, GL, Di Michele, R, Morgans, R, Drust, B and Morton, JP

Daily Distribution of Macronutrient Intakes of Professional Soccer Players From the English Premier League.

http://researchonline.ljmu.ac.uk/id/eprint/7327/

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Anderson, L, Naughton, RJ, Close, GL, Di Michele, R, Morgans, R, Drust, B and Morton, JP (2017) Daily Distribution of Macronutrient Intakes of Professional Soccer Players From the English Premier League. International Journal of Sport Nutrition & Exercise Metabolism. ISSN 1543-

LJMU has developed LJMU Research Online for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

http://researchonline.ljmu.ac.uk/

1	
2	Daily distribution of macronutrient intakes of professional
3	soccer players from the English Premier League
4	sociel prayers nom the English Premier Longae
5	Liam Anderson ¹ Robert I Naughton ² Graeme L Close ¹ Rocco
6	Di Michele ³ Ryland Morgans ⁴ Barry Drust ¹ and James P
7	Morton ¹
2 8	Worton
Q	
10	¹ Passarah Instituta for Sport and Evaraisa Sajanaas
10	Liverpool John Moores University
17	Tom Reilly Building
12	Purom St Compus
13	Liverpool
15	
16	
10	UK
17	² School of Human and Health Sciences
10	Harold Wilson Building
20	University of Huddersfield
20	Oueensgate
21	Huddersfield
22	HD1 3DH
23	IIK
2 . 25	UK
26	³ Department of Biomedical and Neuromotor Sciences
27	University of Bologna
28	Bologna
29	Italy
30	Turry
31	⁴ Cardiff City Football Club
32	Leckwith Road
33	Cardiff
34	CF11 8AZ
35	UK
36	
37	Running head: Distribution of macronutrient intake
38	Running neuro Distribution of macromations marke
39	Address for correspondence:
40	Dr James Morton
41	Research Institute for Sport and Exercise Sciences
42	Liverpool John Moores University
43	Tom Reilly Building
44	Byrom St Campus
45	Liverpool
46	L3 3AF
47	United Kingdom
48	Email: J.P.Morton@ljmu.ac.uk
49	Tel: +44 151 904 6233
50	

51 Abstract

52 The daily distribution of macronutrient intake can modulate 53 aspects of training adaptations, performance and recovery. We 54 therefore assessed the daily distribution of macronutrient intake 55 (as assessed using food diaries supported by the remote food photographic method and 24 h recalls) of professional soccer 56 57 players (n=6) of the English Premier League during a 7-day 58 period consisting of two match days and five training days. On 59 match days, average carbohydrate (CHO) content of the prematch (<1.5 g.kg⁻¹ body mass) and post-match (1 g.kg⁻¹ body 60 61 mass) meals (in recovery from an evening kick-off) were 62 similar (P>0.05) though such intakes were lower than 63 contemporary guidelines considered optimal for pre-match 64 CHO intake and post-match recovery. On training days, we 65 observed a skewed and hierarchical approach (P<0.05 for all 66 comparisons) to protein feeding such that dinner (0.8 g.kg⁻ ¹)>lunch (0.6 g.kg⁻¹)>breakfast (0.3 g.kg⁻¹)>evening snacks 67 (0.1 g.kg⁻¹). We conclude players may benefit from consuming 68 69 greater amounts of CHO in both the pre-match and post-match 70 meals so as to increase CHO availability and maximize rates of 71 muscle glycogen re-synthesis, respectively. Furthermore, 72 attention should also be given to ensuring even daily distribution of protein intake so as to potentially promote 73 74 components of training adaptation.

75 **Keywords:** glycogen, protein, carbohydrate, soccer,

76 Introduction

77 The elite professional soccer player will typically 78 compete in two games per week as well as partake in three to 79 five daily training sessions (Malone et al., 2014; Morgans et al., 80 2015; Anderson et al., 2015). As such, the fundamental goal of the sport nutritionist is to ensure sufficient energy intake in 81 82 order to promote match day physical performance and recovery 83 (Burke et al., 2011). In relation to professional players of the 84 English Premier League (EPL), we recently observed (in a 85 companion paper) self reported mean daily carbohydrate (CHO) intakes of 4.2 and 6.4 g.kg⁻¹ body mass on training days 86 87 and match days, respectively (Anderson et al., 2017). On this 88 basis, we therefore suggested that elite players potentially under-consume CHO when compared with those guidelines 89 90 that are considered optimal to promote muscle glycogen 91 storage (Burke et al., 2011).

92 Nonetheless, in order to provide more informative 93 dietary guidelines (as opposed to total daily energy intake per 94 se), there is also the definitive need to quantify the daily 95 "distribution" of energy and macronutrient intakes. Such a 96 rationale is well documented for CHO given the relevance of 97 both timing and absolute CHO intake in relation to promoting 98 pre-match loading and post-match muscle glycogen re-99 synthesis (Ivy et al., 1988a; Ivy et al., 1988b). To the authors' 100 knowledge, however, the daily distribution of CHO intake on

both training and match days in elite level soccer players hasnot been reported.

103 In contrast to our previous observations of CHO 104 periodization between training and match days (Anderson et al., 2017), we observed consistent daily protein intakes (e.g. 105 106 200 g), the magnitude of which was higher than previously 107 reported in the literature (Maughan, 1997; Bettonviel et al., 108 2016; Gillen et al., 2016). Similar to daily CHO intakes, 109 however, there is also a requirement to quantify daily 110 distribution of protein intakes (Areta et al., 2013; Mamerow et 111 al., 2014). Indeed, these latter authors demonstrated that the 112 timing and even distribution of daily protein doses may have a 113 more influential role in modulating muscle protein synthesis 114 when compared with the absolute dose of protein intake per se, 115 an effect that is evident in response to both feeding alone 116 (Mamerow et al., 2014) and post-exercise feeding (Areta et al., 117 2013). Such skewed approaches to protein feeding have been 118 previously observed in elite youth UK soccer players 119 (Naughton et al., 2016), adult soccer players of the Dutch 120 league (Bettonviel et al., 2016) and a mixed sex cohort of 121 multisport Dutch athletes (Gillen et al. 2016). However, given 122 that we observed higher absolute daily protein intakes 123 (Anderson et al., 2017) compared with all of the 124 aforementioned studies, there is also a need to further

understand the habitual protein feeding patterns in adultprofessional UK soccer players.

127 Accordingly, the aim of the present study was to 128 therefore quantify the daily distribution of energy and 129 macronutrient intakes of professional soccer players of the 130 EPL. Importantly, we provide distribution data related to both 131 training and match days with practical applications therefore 132 related to promoting training adaptations and match day 133 performance. For analysis of total daily energy intake, daily 134 energy expenditure and training and match load of this cohort, 135 the reader is directed to a previous companion paper (Anderson 136 et al., 2017).

137

138 Methods

139 **Participants**

140 Six male professional soccer players from an EPL first 141 team squad (mean \pm SD; age 27 \pm 3 years, body mass 80.5 \pm 142 8.7 kg, height 180 ± 7 cm, body fat 11.9 ± 1.2 %, fat mass $9.2 \pm$ 143 1.6 kg, lean mass 65.0 ± 6.7 kg) volunteered to take part in the 144 study. Players with different positions on the field took part in 145 the study and included 1 wide defender, 1 central defender, 2 central midfielders (1 defending and 1 attacking), 1 wide 146 147 midfielder and 1 center forward. All six players who took part 148 in the study have represented their respective countries at 149 national level. All players remained injury free for the duration of the study. The study was conducted according to the
Declaration of Helsinki and was approved by the University
Ethics Committee of Liverpool John Moores University.

153

154 Study Design

155 Data collection was conducted during the EPL 2015-156 2016 in-season in the month of November. Players continued 157 with their normal in-season training that was prescribed by the 158 club's coaching staff and were available to perform in two 159 competitive games on days 2 and 5 during data collection. 160 During data collection, game 1 kicked off at 20:05 hours and 161 game 2 kicked off at 16:15 hours, both being home fixtures in 162 European and domestic league competitions, respectively. 163 Before the study commenced all players underwent a whole 164 body fan beam Duel-energy X-ray absorptiometry (DXA) 165 measurement scan (Hologic QDR Series, Discovery A, 166 Bedford, MA, USA) in order to obtain body composition, in 167 accordance with the procedures described by Nana et al. 168 (2015).

169

Dietary Intake

171 Self reported EI was assessed from 7-day food diaries 172 for all players and reported in kilocalories (kcal) and 173 kilocalories per kilogram of lean body mass (kcal/kg LBM). 174 Macronutrient intakes were also analysed and reported in

grams (g) and grams per kilogram of body mass $(g.kg^{-1})$. The 175 176 period of 7 days is considered to provide reasonably accurate 177 estimations of habitual energy and nutrient consumptions 178 whilst reducing variability in coding error (Braakhuis et al., 179 2003). On the day prior to data collection, food diaries were 180 explained to players by the lead researcher and an initial dietary 181 habits questionnaire (24 h food recall) was also performed. 182 These questionnaires were used to establish habitual eating 183 patterns and subsequently allow follow up analysis of food 184 diaries. Additionally, they helped to retrieve any potential 185 information that players' may have missed on their food diary 186 In addition, EI was also cross referenced from the input. remote food photographic method (RFPM) in order to have a 187 188 better understanding of portion size and/ or retrieve any 189 information that players' may have missed on their food diary 190 input. This type of method has been shown to accurately 191 measure the EI of free-living individuals (Martin et al., 2009). 192 To further enhance reliability, and ensure that players missed 193 no food or drink consumption, food diaries and RFPM were 194 reviewed and cross checked using a 24-hour recall by the lead 195 researcher after one day of entries (Thompson & Subar, 2008). 196 As such, the lead researcher used these three sources of energy 197 (i.e. food diaries, 24 h recall and RFPM) intake data in 198 combination to collectively estimate daily energy and 199 macronutrient intake / distribution. To obtain energy and 200 macronutrient composition, the Nutritics professional diet 201 analysis software (Nutritics Ltd, Ireland) was used. Energy and 202 macronutrient intake was further assessed in relation to timing 203 of ingestion. Meals on training days were split into breakfast, 204 morning snack, lunch, afternoon snack, dinner and evening 205 snack. Time and type of consumption was used to distinguish 206 between meals; breakfast (main meal consumed between 6-207 9.30am), morning snack (foods consumed between the 208 breakfast main meal and the lunch), lunch (main meal 209 consumed between 11.30-1.30pm), afternoon snack (foods 210 consumed between lunch and dinner), dinner (main meal 211 consumed between 5-8pm), and evening snack (foods 212 consumed after dinner and prior to sleep).

213 Meals on match days were split into pre-match meal 214 (PMM), pre-match snack (PMS), during match (DM), post-215 match (PM) and post-match recovery meal (PMRM). Timing of 216 events was used to distinguish between meals on match days; 217 PMM (main meal consumed 3 hours prior to kick off), PMS 218 (foods consumed between the PMM and entering the changing 219 rooms after the cessation of the warm up), DM (foods 220 consumed from when the players entered the changing rooms 221 after the warm up until the final whistle or since they were 222 substituted), PM (foods consumed in the changing rooms after 223 the match), PMRM (main meal consumed <3 hours after the 224 end of the match).

225

226 Inter-Researcher Reliability of the Methods

227 To assess inter-researcher reliability, author one, author two 228 and an independent researcher (not included on the authorship) 229 individually assessed energy intake data for one day of one 230 player selected at random. No significant difference was 231 observed (as determined by one-way ANOVA) between 232 researchers for energy (P=0.95), CHO (P=0.99), protein 233 (P=0.95) or fat (P=0.80) intake. Daily totals for researchers 1, 234 2 and 3 were as follows: energy intake = 3174, 3044 and 3013235 kcals; CHO = 347, 353 and 332 g; protein = 208, 201, and 194 236 g and fat = 106, 92 and 101 g, respectively.

237

238 Statistical Analysis

239 All data are presented as the mean \pm standard deviation 240 (SD). Meal distribution data was using linear mixed models 241 with meal as the fixed factor. A random intercept was set for 242 each individual player. When there was a significant (P < 0.05) 243 effect of the fixed factor, Tukey post-hoc pairwise comparisons 244 were performed to identify which categories of the factor 245 differed. This whole analysis was performed separately for 246 training and match days. In the match day's analysis, a fixed 247 factor for day was also included to compare energy intake and 248 distribution of the two different match days. In all the analyses, 249 statistical significance was set at P<0.05. The statistical analysis was carried out with R, version 3.3.1.

251

252

253 Results

254 Energy and Macronutrient Distribution Across Meals on

255 Training Days

256 There were significant differences in the reported 257 absolute and relative energy and macronutrient between meals 258 consumed on training days (P<0.01 for all examined absolute 259 and relative energy intake variables; see Figure 1). 260 Specifically, players consumed higher absolute and relative EI 261 at dinner compared with breakfast, morning, afternoon and 262 evening snacks (P<0.01 for all comparisons). Additionally, 263 absolute and relative EI was also greater at lunch compared 264 with the morning and evening snacks (P<0.01). Absolute and 265 relative CHO intakes were higher at dinner compared with 266 morning snack (both P<0.01), lunch (both P<0.05) and evening 267 snack (both P<0.01), with relative CHO intake also being 268 higher at dinner compared with breakfast (P=0.04).

Protein and relative protein intakes were greater at dinner compared with breakfast, morning snacks, afternoon snacks and evening snacks (P<0.01 for all comparisons). In addition, absolute and relative protein intakes were greater at lunch compared with breakfast, morning snacks and evening snacks (P<0.01 for all comparisons). Both absolute and relative protein intakes were also higher at breakfast compared with evening snack (both P<0.02) and higher at the afternoon snack compared with the evening snack (both P<0.01).

In relation to fat intake, both absolute and relative intakes were higher at dinner compared with the morning, afternoon snacks and evening snacks (P<0.05 for all comparisons). Additionally, fat intake was also higher at lunch compared with the morning snack (P<0.01 for both absolute and relative intakes).

284

285 Energy and Macronutrient Intake Across Meals on Match286 Days

287 There was no significant difference (P>0.05 for all 288 meals; see Figure 2) in absolute and relative energy and 289 macronutrient intake between meals on the two difference 290 match days. However, significant differences were observed 291 between meals consumed on match days for all energy and 292 macronutrient variables (all P<0.05; see Figure 2). The absolute 293 and relative energy and protein intake were higher in the PMM 294 and PM compared with the PMS, DM and PMRM (all P<0.05). 295 Additionally, the absolute and relative CHO intake were also 296 higher in the PMM and PM compared with the PMS and DM 297 (all P<0.05). Fat intake in the PMM and the PM, when 298 expressed in both absolute and relative terms, were higher than

the PMS and DM (all P<0.05), where the PMM was also lowerthan the PMRM (both P<0.05).

301

302

303 Discussion

304 Having previously quantified the daily "total" energy 305 intake and expenditure of the players studied here (Anderson et 306 al., 2017), the aim of the present study was to subsequently 307 quantify the daily distribution of energy and macronutrient 308 intakes on both training and match days. Importantly, we 309 observed that players adopt a skewed approach to feeding on 310 training days such that absolute energy intake, CHO and 311 protein intake are consumed in a hierarchical manner of 312 dinner>lunch>breakfast>snacks. Moreover, we also observed 313 that players tend to under-consume CHO on match days in 314 relation to pre-match and post-match meals, especially in 315 recovery from an evening kick-off time. Taken together, our 316 data highlight the importance of obtaining dietary data related 317 to distribution (as opposed to total daily energy intake per se, 318 Anderson et al., 2017) given the implications related to 319 components of training adaptation, performance and recovery.

In our companion paper (Anderson et al., 2017), we reported that the players studied herein practiced elements of CHO periodization such that total daily CHO intake was greater on match days (i.e. 6.4 g.kg⁻¹ BM) compared with

training days (i.e. 4.2 g.kg⁻¹ BM). Although such CHO 324 325 periodization strategies may be in accordance with the principle 326 of "fuel for the work required" (Impey et al. 2016; Bartlett et 327 al., 2015; Hawley & Morton, 2015), we suggested that players 328 were likely under-consuming CHO in terms of maximizing 329 match day physical performance and recovery. Further 330 evidence highlighting this potential "sub-optimal CHO intake" 331 is also provided by the dietary distribution data provided here. 332 For example, in relation to match day itself, our data suggest 333 that players did not meet current CHO guidelines for which to 334 optimize aspects of physical (Burke et al., 2011), technical (Ali 335 & Williams, 2009) and cognitive (Welsh et al., 2002) performance. Indeed, both the pre-match meal ($< 1.5 \text{ g.kg}^{-1}$ 336 body mass) and CHO feeding during match play (~30 g.h⁻¹; 337 four players consumed <30 g.h⁻¹, see Anderson et al. 2017) 338 339 could be considered sub-optimal in relation to those studies 340 (Wee et al., 2005; Foskett et al., 2008) demonstrating higher CHO intakes (e.g. 2-3 g.kg⁻¹ body mass and 60 g.h⁻¹, 341 342 respectively) induce physiological benefits that are facilitative 343 of improved high-intensity intermittent performance e.g. high 344 pre-exercise glycogen stores, maintenance of plasma 345 glucose/CHO oxidation during exercise and muscle glycogen 346 sparing.

347

348 Given that the present study was conducted during a 349 two game per week schedule, there was the obvious nutritional 350 requirement to maximize muscle glycogen storage in the 24-48 351 h after each game (Krustrup et al., 2006; Bassau et al., 2002). 352 To this end, we also observed CHO intakes that would be 353 considered sub-optimal in relation to maximizing rates of post-354 match muscle glycogen re-synthesis (Jentjens & Jeukendrup, 355 2003). Indeed, in contrast to the well-accepted guidelines of 1.2 g.kg⁻¹ body mass for several hours post-exercise, we 356 observed reported intakes of <1 g.kg⁻¹ in the immediate period 357 358 after match day 1 (i.e. the night-time kick off). Such post-game 359 intakes coupled with the relatively low absolute daily intake 360 (i.e. 4 $g.kg^{-1}$) on the subsequent day (see Anderson et al., 2017) 361 would inevitably ensue that absolute muscle glycogen re-362 synthesis was likely compromised, an effect that may be 363 especially prevalent in type II fibres (Gunnarsson et al., 2013). 364 It is noteworthy, however, that the high absolute protein intakes 365 consumed in the post-match period (i.e. >50 g) would likely 366 potentiate rates of muscle glycogen re-synthesis when 367 consumed in the presence of sub-optimal CHO availability 368 (Van Loon et al., 2000).

369 Despite our observation of CHO periodization during 370 the weekly microcycle, we previously observed (Anderson et 371 al., 2017) consistent daily protein intakes (approximately 200 g 372 per day), the magnitude of which was higher than that typically 373 reported (<150 g/day) previously for adult (Maughan, 1997; 374 Bettonviel et al., 2016) and youth professional male soccer 375 players (Naughton et al., 2016). Similar to CHO intake, 376 however, it is also prudent to consider the daily distribution of 377 protein feeding given that both that skewed and sub-optimal 378 intakes at specific meal times can reduce rates of muscle 379 protein synthesis (Areta et al., 2013; Mamerow et al., 2014). 380 Indeed, recent data suggest that the timing and even distribution of daily protein doses may have a more influential 381 382 role in modulating muscle protein synthesis when compared 383 with the absolute dose of protein intake, an effect that is 384 evident in response to both feeding alone (Mamerow et al., 385 2014) and post-exercise feeding (Areta et al., 2013). In this 386 regard, we observed a skewed pattern of daily protein intake in 387 that absolute protein was consumed in a hierarchical order 388 where dinner>lunch>breakfast>snacks. This finding also 389 agrees with our previous observations on the protein feeding 390 patterns of elite youth soccer players (Naughton et al., 2016) as 391 well as adult players from the Dutch league (Bettonviel et al., 392 2016) and a mixed sex cohort of Dutch athletes (Gillen et al. 393 Nonetheless, given that we observed higher daily 2016). 394 protein intakes (>200 g/day) compared with the previous 395 (typically <150 g/day), studies examination of daily 396 distribution data also allows us to comment on those meals that 397 led to greater absolute protein intake. In this regard, it appears

that an additional absolute intake of approximately 20-25 g at
both lunch and dinner accounted for the greater absolute total
daily intake.

401 Based on recent data suggesting that trained athletes 402 (especially those with higher lean mass) may require protein 403 doses of approximately 40 g (MacNaughton et al., 2016) as 404 well as the importance of protein feeding prior to sleep (Res et 405 al., 2012), our data suggest that breakfast and morning, 406 afternoon and bedtime snacks are key times to improve for the 407 present sample. We acknowledge, however, that protein 408 requirements (both in absolute dosing and timing) should be 409 tailored to the specific population in question in accordance 410 with timing of training sessions, training load and moreover, 411 individualized training goals.

412 Despite the novelty and practical application of the 413 current study, our data are not without limitations, largely a 414 reflection of the practical demands of data collection in an elite 415 football setting. Firstly, this study is reflective of only six 416 players from one team only (albeit reflective of a top EPL 417 team) and hence may not be representative of the customary 418 training and nutritional habits of other teams. Nonetheless, we 419 deliberately recruited players with different playing positions in 420 an attempt to provide a more representative sample of 421 professional soccer players. Secondly, our deliberate choice to 422 study a two game week scenario (as is highly relevant for elite 423 level players) may not be applicable to players of lower 424 standards. Thirdly, as with all dietary analysis studies, our data 425 may be limited by both under-reporting and inter-researcher 426 variability in ability to assess dietary intakes. Indeed, whilst 427 we observed no significant group mean changes in body mass 428 over the data collection period, two of our subjects did appear 429 to under report whereas four of the subjects reported energy 430 intake data that was comparable (within 200 kcal) to energy 431 expenditure data (see Anderson et al. 2017). Finally, both of 432 the games studied here represented home games and hence the 433 nutritional choices are likely to be influenced by the philosophy 434 and service provision of the club coaching and catering staff.

435 In summary, we simultaneously quantified for the first 436 time the daily distribution of energy and macronutrient intakes 437 of EPL soccer players on both training and match days. Our 438 data suggest that players may benefit from consuming greater 439 amounts of CHO in both the pre-match and post-match meals 440 so as to increase CHO availability and maximize rates of 441 muscle glycogen re-synthesis, respectively. Furthermore, we 442 also observed that daily protein intake was consumed in a 443 hierarchical manner such that dinner > lunch > breakfast > 444 snacks. Attention should also be given to therefore ensuring 445 even distribution of daily protein intake so as to potentially 446 promote components of training adaptation.

447

449 Acknowledgements

450

The authors would like to thank all of the participating players
for the cooperation and commitments during all data collection
procedures. We would also like to thank the team's coaches for
cooperation during data collection.

- 455
- 456

457 **References**

458

Ali, A., & Williams, C. (2009). Carbohydrate ingestion and
soccer skill performance during prolonged intermittent
exercise. *J Sports Sci*, *27*, 1499-1508.

462

Anderson, L., Orme, P., Di Michele, R., Close, G.L., Morgans,
R., Drust, B., & Morton, J.P. (2015). Quantification of training
load during one-, two- and three-game week schedules in
professional soccer players from the English Premier League:
implications for carbohydrate periodization. *J Sports Sci, 4*, 1-

- 468 10.
- 469
- 470 Anderson, L., Orme, P., Naughton, R.J., Close, G.L., Milsom,
- 471 J., Rydings, D., O'Boyle, A., Di Michele, R., Louis, J.,
- 472 Morgans, R., Drust, B., & Morton, J.P. (2017). Energy intake

- 476
- Areta, J.L., Burke, L.M., Ross, M.L., Camera, D.M., West,
 D.W., Broad, E.M., Coffey, V.G. (2013). Timing and
 distribution of protein ingestion during prolonged recovery
 from resistance exercise alters myofibrillar protein synthesis. *J Physiol, 591,* 2319-2331.
- 482
- Bartlett, J.D., Hawley, J.A., & Morton, J.P. (2015).
 Carbohydrate availability and exercise training adaptation: Too
 much of a good thing? *Eur J Sport Sci, 15,* 3-12.
- 486
- Bassau, V.A., Fairchild, T.J., Rao, A., Steele, P., & Fournier,
 PA. (2002). Carbohydrate loading in human muscle: an
 improved 1 day protocol. *Eur J Appl Physiol*, *87*, 290-295.
- 490
- Bettonviel, A.EO., Brinkmans, N.Y.J., Russcher, K.,
 Wardenaar, F.C., & Witard, O.C. (2016). Nutritional status and
 daytime pattern of protein intake on match, post-match, rest
 and training days in senior professional and youth elite soccer
 players. *Int J Sport Nutr Exerc Metab, epub ahead of print.*

Braakhuis, A.J., Meredith, K., Cox, G.R., Hopkins, W.G., &
Burke, L.M. (2003). Variability in estimation of self-reported
dietary intake data from elite athletes resulting from coding by
different sports dietitians. *Int J Sport Nutr Exerc Metab, 13,*152-165.

- 502
- 503 Burke, L.M., Hawley, J.A., Wong, S.H., & Jeukendrup, A.E.
- 504 (2011). Carbohydrates for training and competition. *J Sports*505 *Sci*, *29*, 17-27.
- 506
- Foskett, A., Williams, C., Boobis, L., & Tsintzas, K. (2008).
 Carbohydrate availability and energy metabolism during
 intermittent running. *Med Sci Sports Exerc, 40*, 96-103.
- 510
- Gillen, J.B., Trommelen, J., Wardenaar, F.C., Brinkmans,
 N.Y.J., Versteegen, J.J., Jonvik, K.L., Kapp, C., de Vries, J.,
 van den Borne, J.J.G.C., Gibala, M., & van Loon, L.J.C.
 (2016). Dietary protein intake and distribution patterns of welltrained Dutch athletes. *Int J Sport Nutr Exerc Metab, epub ahead of print.*
- 517
- 518 Gunnarsson, T.P., Bendiksen, M., Bischoff, R., Christensen,
 519 P.M., Lesivig, B., Madsen, K., ... Bangsbo, J. (2013). Effect of
- 520 whey protein- and carbohydrate-enriched diet on glycogen

- resynthesis during the first 48 h after a soccer game. *Scand J Med Sci Sports, 23,* 508-515.
- 523
- Hawley, J.A., & Morton, J.P. (2014). Ramping up the signal:
 promoting enurance training adaptation in skeletal muscle by
 nutritional manipulation. *Clin Exp Pharmacol Physiol*, *41*, 608-*613*.
- 528
- 529 Impey, S.G., Hammon, K.M., Shepherd, S.O., Sharples, A.P.,
- 530 Stewart, C., Limb, M., Smith, K., Philp, A., Jeromson, S.,
- 531 Hamilton, D.L., Close, G.L., & Morton, J.P. (2016). Fuel for
- the work required: a practical approach to amalgamating train-
- 533 low paradigms for endurance athletes. *Physiol Rep, 4*, e12803.
- 534
- Ivy, J, L., Katz, A.L., Cutler, C.L., Sherman, W.M., & Coyle,
 E.F. (1988a). Muscle glycogen synthesis after exercise: effect
 of time of carbohydrate ingestion. *J Appl Physiol, 64*, 14801485.
- 539
- 540 Ivy, J.L., Lee, M.C., Brozinick, J.T. Jr., Reed. M.J. (1988b).
- 541 Muscle glycogen storage after different amounts of542 carbohydrate ingestion. *J Appl Physiol*, 65, 2018-2023.
- 543

- Jentjens, R., & Jeukendrup, A. (2003). Determinants of postexercise glycogen synthesis during short-term recovery. *Sports Med*, *33*, 117-144.
- 547
- Krustrup, P., Mohr, M., Steensberg, A., Bencke, J., Kjer, M., &
 Bangsbo, J. (2006). Muscle and blood metabolites during a
 soccer game: Implications for sprint performance. *Med Sci Sports Exerc, 38*, 1165-1174.
- 552
- Malone, J.J., Di Michele, R., Morgans, R., Burgess, D.,
 Morton, J.P., & Drust, B. (2015). Seasonal training-load
 quantification in elite English premier league soccer players. *Int J Sports Physiol Perform, 10,* 489-497.
- 557
- Martin, C.K., Han, H., Coulon, S.M., Allen, H.R., Champagne,
 C.M., & Anton, S.D. (2009). A novel method to remotely
 measure food intake of free-living people in real-time: The
 remote food photography method (RFPM). *B J Nutr, 101,* 446456.
- 563
- Maughan, R.J. (1997). Energy and macronutrient intakes of
 professional football (soccer) players. *Br J Sports Med*, *31*, 4547.
- 567

- Macnaughton, L.S., Wardle, S.L., Witard, O.C., McGlory, C.,
 Hamilton, D.L., Jeromson, S., Lawrence, C.E., Wallis, G.A., &
 Tipton, K.D. (2016). The response of muscle protein synthesis
 following whole-body resistance exercise is greater following
 40 g than 20 g of ingested whey protein. *Physiol Rep, 4*,
 e12893.
- 574
- 575 Mamerow, M.M., Mettler, J.A., English, K.L., Casperson, S.L.,
- 576 Arentson-Lantz, E., Sheffield-Moore, M., Layman, D.K., &
- 577 Paddon-Jones, D. (2014). Dietary protein distribution positively
 578 influences 24-h muscle protein synthesis in healthy adults. J
 579 *Nutri, 144,* 876-880.
- 580
- 581
- 582 Morgans, R., Orme, P., Anderson, L., Drust, B., & Morton, J.P.
- 583 (2014). An intensitve Winter fixture schedule induces a
 584 transient fall in salivary IgA in English premier league soccer
 585 players. *Res Sports Med*, *22*, 346-354.
- 586
- 587 Nana, A., Slater, G.J., Stewart, A.D., & Burke, L.M. (2015).
- 588 Methodology review: using dual-energy X-ray absorptiometry
- 589 (DXA) for the assessment of body composition in athletes and
- active people. Int J Sport Nutr Exerc Metab, 25, 198-215.
- 591

Naughton, R.J., Drust, B., O'Boyle, A., Morgans, R., Abayomi,
J., Davies, L.G., Morton, J.P., & Mahon, E. (2016). Daily
distribution of carbohydrate, protein and fat intake in elite
youth academy soccer players over a 7-day training period. *Int J Sport Nutri Exerc Metab, epub ahead of print.*

597

Res, P.T., Groen, B., Pennings, B., Beelen, M., Wallis, G.A.,
Gijsen, A.P., Senden, J.M., Van Loon, L.J. (2012). Protein
ingestion before sleep improves postexercise overnight
recovery. *Med Sci Sports Exerc, 44,* 1560-1569.

602

Thompson, F., & Subar, A. (2008). Dietary assessment
methodology. In A. M. Coulston & C.J. Boushey (Eds.), *Nutrition in the prevention and treatment of disease* (pp. 3-39).

606 San Diego, CA: Academic Press.

607

Wee, S., Williams, C., Tsintzas, K., & Boobis, L. (2005).
Ingestion of a high-glycemic index meal increases muscle
glycogen storage at rest but augments its utilization during
subsequent exercise. *J App Physiol, 99,* 707-714.

612

Welsh, R.S., Davis, J.M., Burke, J.R., & Williams, H.G.
(2002). Carbohydrates and physical/mental performance during
intermittent exercise to fatigue. *Med Sci Sports Exerc, 34*, 72373.



Figure 1: Energy and macronutrient intakes meal distribution on training days. Figure A=absolute energy expenditure, Figure B=energy expenditure relative to lean body mass, Figure C=absolute carbohydrate, Figure D=relative carbohydrate, Figure E=absolute protein, Figure F=relative protein, Figure G=absolute fat and Figure H=relative fat. a denotes difference from breakfast, b denotes different from morning snack, c denotes difference from lunch, d denotes difference from afternoon snack, e denotes difference from dinner, f denotes difference from evening snack.



Figure 2: Energy and macronutrient intake meal distribution on the two match days during the study period. Black bars=match day 1 and white bars=match day 2. a denotes difference from PMM, b denotes difference from PMRM. PMM=Pre Match Meal, PMS=Pre-Match Snack, DM=During-Match, PM=Post-Match, PMRM=Post-Match Recovery Meal.