DOI 10.1186/s40101-016-0112-8

Journal of Physiological Anthropology

ORIGINAL ARTICLE Open Access

Whole-body patterns of the range of joint motion in young adults: masculine type and feminine type

Keiichi Moromizato[1], Ryosuke Kimura[1*], Hitoshi Fukase[2], Kyoko Yamaguchi[1,3] and Hajime Ishida[1]

Abstract

Background: Understanding the whole-body patterns of joint flexibility and their related biological and physical factors contributes not only to clinical assessments but also to the fields of human factors and ergonomics. In this study, ranges of motion (ROMs) at limb and trunk joints of young adults were analysed to understand covariation patterns of different joint motions and to identify factors associated with the variation in ROM.

Methods: Seventy-eight healthy volunteers (42 males and 36 females) living on Okinawa Island, Japan, were recruited. Passive ROM was measured at multiple joints through the whole body (31 measurements) including the left and right side limbs and trunk.

Results: Comparisons between males and females, dominant and non-dominant sides, and antagonistic motions indicated that body structures influence ROMs. In principal component analysis (PCA) on the ROM data, the first principal component (PC1) represented the sex difference and a similar covariation pattern appeared in the analysis within each sex. Multiple regression analysis showed that this component was associated with sex, age, body fat %, iliospinale height, and leg extension strength.

Conclusions: The present study identified that there is a spectrum of "masculine" and "feminine" types in the whole-body patterns of joint flexibility. This study also suggested that body proportion and composition, muscle mass and strength, and possibly skeletal structures partly explain such patterns. These results would be important to understand individual variation in susceptibility to joint injuries and diseases and in one's suitable and effective postures and motions.

Keywords: Range of motion, Joints, Young adult, Principal component analysis, Multiple regression analysis, Sexual dimorphism, Hand/foot dominance

Introduction

In the field of orthopaedics and rehabilitation medicine, measuring range of motion (ROM) is a clinical procedure to evaluate a mechanical joint problem caused by disorders of the locomotor apparatus. The purpose of ROM measurement is not only to observe the extent of inhibition but also to identify the factors that restrict joint movement and to evaluate the effectiveness of treatment and training. A measurement method of ROM was established by the American Academy of Orthopaedic Surgeons (AAOS) [1], in which the standard anatomical position was defined as the neutral zero starting position, and this method has been used internationally. The AAOS has provided the reference values for normal joint ROM. However, there is great variation in ROM even among healthy individuals, depending on sex, age, physical constitution, daily activities etc. Therefore, in clinical assessments using ROM, it is important to establish an individual standard for each patient. For this purpose, it is indispensable to identify which biological and physical factors affect ROM. In addition, understanding the correlations in ROM of different joints can improve clinical assessments for each individual.

* Correspondence: rkimura@med.u-ryukyu.ac.jp
[1]Department of Human Biology and Anatomy, Graduate School of Medicine, University of the Ryukyus, Uehara 207, Nishihara, Okinawa 903-0215, Japan
Full list of author information is available at the end of the article

BioMed Central

© 2016 The Author(s). **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

UNCORRECTED PROOF

There have been many studies of joint ROM to date. Most of the studies have focused only on upper or lower extremity joint motions, focusing on the effects of age, sex, and/or side dominance and on some special population such as sports athletes and disease patients [2–9]. Even in the studies that have examined six major limb joints of the upper and lower limbs [10–18], correlations among different joints have not been sufficiently discussed. In addition, there have been few studies measuring ROM throughout the whole body, including the trunk joints [19].

Understanding whole-body patterns of joint flexibility and identifying their related biological and physical factors contribute not only to clinical assessments but also to the fields of human factors and ergonomics. Biological and physical factors such as age, sex, physical constitution, and daily activities can affect patterns of joint flexibility. Then, whole-body patterns of joint flexibility can have influences on whole-body motions and eventually can be important to know one's suitable and effective postures and motions.

The aims of this study were to understand the whole-body patterns of joint flexibility and to identify factors associated with variations in ROM. For this purpose, we measured ROMs of the limb and trunk joints of young adults and analysed their association with biological and physical factors. We also compared dominant and non-dominant sides to obtain a cue for related factors. Principal component analysis (PCA) was performed using multiple joint ROM data to identify covariation patterns among different joints. It was found that a major pattern that explains the variation between sexes also appears within each sex, and this pattern appears to be associated with some somatometric and sthenometric measurements.

Materials and methods

Subjects

We recruited volunteers living in Okinawa Island, Japan. Inclusive criteria were healthy males and females between the ages of 20 and 29 who did not have any joint diseases and any history of orthopaedic surgery on joints. As shown in Additional file 1: Table S1, the subjects consisted of 36 females and 42 males. Their ages were concentrated in early twenties and ranged from 20 to 25 years in females (mean 20.8 years, SD 1.2 years) and from 20 to 29 years in males (mean 21.4 years, SD 1.9 years). Hand/arm and foot/leg dominances were determined based on a questionnaire [20, 21]. Most individuals had an experience of sports when they were high school students. There was no significant difference in the frequencies of sports experience between males and females (Fisher's exact test). All subjects provided their written informed consent to participate in this research project.

Measurements

ROM data were collected from the subjects using a goniometer (OG Giken. Co. Ltd., Okayama, Japan). All ROM measurements were performed by four observers after confirming that the inter-observer errors in the measurements were small ($0.97 < ICC < 0.99$). The motions examined are shown in Table 1.

Height, weight, upper limb length, iliospinale height, forearm circumference, forearm minimum circumference, calf circumference, and ankle circumference were measured following the standard anthropometric method [22]. On

Table 1 Summary of somatometry and ROM for each joint motion

Item	All		Female		Male	
	Mean	SD	Mean	SD	Mean	SD
Height [cm]	163.7	9.0	156.9	5.2	168.9	7.7
Weight [kg]	57.0	9.3	51.8	6.4	61.1	9.3
Body fat percentage [%]	20.6	5.2	22.1	3.3	19.4	6.1
Lean body mass [kg]	45.3	8.0	40.3	5.0	49.2	7.8
ROMs [°]						
Shoulder flexion	176.4	5.6	178.4	3.2	174.7	6.5
Shoulder extension	66.5	6.2	67.6	7.1	65.7	5.4
Shoulder abduction	179.7	1.0	179.6	0.9	179.8	1.2
Shoulder external rotation	92.9	9.3	94.5	8.2	91.5	10.1
Shoulder internal rotation	62.5	11.0	67.4	9.2	58.4	10.9
Shoulder horizontal flexion	133.6	9.4	137.1	9.2	130.7	8.6
Shoulder horizontal extension	56.1	8.6	58.3	8.5	54.3	8.3
Elbow flexion	142.6	4.9	144.5	4.0	141.0	5.1
Elbow extension	4.3	5.4	5.6	6.1	3.2	4.6
Wrist extension	81.1	9.1	83.7	7.9	79.0	9.6
Wrist flexion	88.2	10.8	89.6	12.9	87.0	8.5
Fingers V MCP flexion	104.9	12.9	104.6	14.6	105.2	11.4
Fingers V MCP extension	71.8	15.7	73.6	14.1	70.3	16.8
Trunk flexion	35.3	9.5	32.9	9.3	37.3	9.2
Trunk extension	28.2	8.7	28.3	7.6	28.2	9.7
Trunk rotation	48.3	10.8	43.9	11.3	52.2	8.8
Trunk lateral bending	23.2	4.6	21.3	4.2	24.8	4.4
Hip flexion	128.4	6.7	130.4	7.8	126.7	5.1
Hip extension	17.0	3.9	16.1	3.7	17.9	3.9
Hip abduction	33.0	4.8	34.1	5.3	32.1	4.1
Hip adduction	13.8	6.3	13.9	4.1	13.7	7.7
Hip external rotation	43.9	8.4	40.6	8.1	46.7	7.6
Hip internal rotation	42.1	9.6	47.9	9.3	37.1	6.5
Knee flexion	147.9	5.9	148.9	5.0	147.1	6.4
Knee extension	3.5	4.8	5.3	5.4	2.0	3.7
Ankle dorsi flexion	23.0	6.5	23.9	6.2	22.3	6.8
Ankle plantar flexion	51.9	6.6	54.1	6.6	50.1	6.1

The average of dominant and non-dominant is shown

UNCORRECTED PROOF

the digitally scanned images of the left hand, the second and fourth finger lengths were measured using software, Image J (ImageJ, NIH, Bethesda, MD, USA), and the 2D:4D ratio was calculated.

Grip strength was measured using the Smedley Hand Dynamometer (OG Giken. Co. Ltd). In the measurements, the subject was in a standing position with arms at their side, not touching their body. We obtained the average of three trials. As for leg strength, the flexor and extensor muscle strengths at the knee were measured using the Biodex System 3 dynamometer (Sakaimed Co. Ltd., Tokyo, Japan). Isometric knee flexion and extension strengths were tested at 60° of knee flexion. Peak torque was recorded for each motion in three trials and the average peak torque was calculated.

The calliper method was used to calculate body fat percentage (BF%). Subcutaneous fat thickness was measured at the mid-point of the posterior surface of an upper arm, at the inferior angle point of the scapula, and at the side of the lateral point from the umbilicus. Then, BF% and lean body mass (LBM) was calculated using the following formula [23–25]:

$$\text{Body surface area}\left(\text{cm}^2\right): \; S = 72.46 \times H^{0.725} \times W^{0.425},$$

$$\text{Body density}: \; D = 1.0935 - 0.000297 \times T \times S / W / 100,$$

$$\text{BF}\% = (4.570 / D - 4.142) \times 100,$$

$$\text{LBM} = W \times (100 - \text{BF}\%)/100,$$

where W, H, and T are weight (kg), height (cm), and the sum of subcutaneous fat thicknesses (mm), respectively. The results of the somatometric and sthenometric measurements are shown in Table 1 and Additional file 1: Table S2, respectively.

Statistical analyses

Statistical analyses were performed using SPSS® Statistics version 19 (IBM Japan, Tokyo, Japan) and Excel Statistics (Social Survey Research Information Co. Ltd., Tokyo, Japan). Basic summary statistics were calculated for each sex. To identify biological and physical factors associated with each ROM, multiple regression analysis was used. Differences between dominant and non-dominant sides were examined by paired t test, in which we subtracted ROM for dominant side from ROM for non-dominant side (ROM(ND) – ROM(D)). PCA was performed to elucidate whole-body patterns of joint ROM. Correlation coefficient and partial correlation coefficient controlling for sex were calculated between each principal component (PC) and each factor. Finally, to disclose factors associated with the whole-body patterns of ROM, multiple regression analysis was conducted. As for bilateral measurements, the averages of the left and right sides were input into these statistical analyses except for the test of the side difference.

Results

Effects of sex, age, height, BF%, and LBM on each motion

The results of measurements are summarized in Table 1. To concretely explain the factors responsible for the individual variation in ROM, multiple regression analyses were performed for each motion, including sex, age, body height, BF%, and LBM as explanatory variables. The female sex significantly increased ROMs for shoulder flexion, internal rotation and horizontal flexion, elbow flexion and extension, wrist extension, and hip flexion, adduction, and internal rotation, but decreased ROMs for hip extension and external rotation, and trunk flexion and rotation (Table 2). Hip extension versus flexion and hip external rotation versus internal rotation are pairs of antagonistic motions. When the total ranges of these antagonistic motions were compared, no significant sex difference was found (data not shown).

The multiple regression analyses also showed that older age is significantly associated with lower ROMs for shoulder external rotation and horizontal flexion, elbow extension, wrist flexion and extension, and higher ROMs for elbow flexion and trunk flexion and rotation.

A higher LBM was significantly related with lower ROMs for shoulder external rotation and horizontal extension, and with higher ROMs for wrist flexion and hip adduction (Table 2). BF% negatively affected ROMs for shoulder external rotation, shoulder horizontal flexion, and elbow flexion and extension. In contrast, BF% was positively associated with trunk flexion and rotation, and hip extension and external rotation.

ROM differences between dominant and non-dominant sides

Significant ROM differences between dominant and non-dominant sides were detected for several motions. The non-dominant side had higher mobility than the dominant side for shoulder internal rotation, hip abduction, and ankle plantar flexion, whereas the opposite was observed for shoulder external rotation, wrist flexion, and hip adduction (Table 3). In the total range of antagonistic motions, however, there were no significant differences except for ankle dorsiflexion and plantarflexion in all the subjects (Table 4).

Whole-body ROM patterns revealed by PCA

PCA was performed on the ROM data of all subjects, female subjects and male subjects. The first three PCs are shown in Fig. 1. When PCs resulting from female and male sample sets were compared, both sexes demonstrated similar patterns of PC loadings in PC1 (Fig. 2a),

UNCORRECTED PROOF

t2.1 **Table 2** Multiple regression analysis for identifying factors associated with ROM for each joint motion

t2.2 Dependent variable	Sex (F:0, M:1)			Age			Height			Body fat %			Lean body mass					
t2.3	*B*	*β*	*P*	*B*	*β*	*P*	*B*	*β*	*P*	*B*	*β*	*P*	*B*	*β*	*P*	Constant	*P*	R^2
t2.4 Shoulder flexion	−4.68	−0.42	*2.5E−02*	−0.05	−0.02	8.9E−01	−0.03	−0.05	8.0E−01	0.04	0.04	7.5E−01	0.14	0.20	3.6E−01	178.39	*3.8E−13*	0.13
t2.5 Shoulder extension	−4.16	−0.33	7.7E−02	0.73	0.19	1.0E−01	0.00	0.00	9.9E−01	0.22	0.18	1.5E−01	0.23	0.30	1.7E−01	37.97	9.9E−02	0.12
t2.6 Shoulder abduction	0.51	0.24	2.1E−01	0.08	0.13	2.8E−01	−0.02	−0.21	3.4E−01	−0.01	−0.06	6.2E−01	−0.01	−0.07	7.6E−01	182.46	*4.8E−55*	0.06
t2.7 Shoulder external rotation	−0.90	−0.05	7.8E−01	−1.67	−0.29	*8.7E−03*	0.36	0.34	9.5E−02	−0.46	−0.25	*3.6E−02*	−0.66	−0.57	*5.6E−03*	109.72	*9.2E−04*	0.23
t2.8 Shoulder internal rotation	−13.06	−0.59	*1.2E−03*	−0.16	−0.02	8.3E−01	−0.05	−0.04	8.4E−01	0.13	0.06	6.2E−01	0.46	0.33	1.0E−01	57.62	1.3E−01	0.21
t2.9 Shoulder horizontal flexion	−11.41	−0.61	*2.9E−04*	−1.75	−0.30	*3.2E−03*	0.38	0.37	5.2E−02	−0.56	−0.30	*5.9E−03*	−0.13	−0.11	5.4E−01	130.98	*3.1E−05*	0.34
t2.10 Shoulder horizontal extension	−1.14	−0.07	7.1E−01	−0.68	−0.13	2.5E−01	0.42	0.44	*3.9E−02*	0.16	0.10	4.3E−01	−0.62	−0.59	*5.8E−03*	27.60	3.6E−01	0.18
t2.11 Elbow flexion	−5.62	−0.57	*1.4E−03*	0.69	0.22	*3.7E−02*	0.17	0.31	1.3E−01	−0.26	−0.27	*2.2E−02*	−0.14	−0.23	2.6E−01	115.15	*1.5E−09*	0.25
t2.12 Elbow extension	−4.67	−0.43	*1.5E−02*	−0.95	−0.28	*9.4E−03*	0.16	0.27	1.8E−01	−0.34	−0.32	*7.5E−03*	−0.07	−0.10	6.0E−01	9.93	5.9E−01	0.24
t2.13 Wrist extension	−8.09	−0.44	*9.7E−03*	−2.21	−0.39	*3.0E−04*	0.26	0.26	1.9E−01	−0.32	−0.18	1.2E−01	−0.02	−0.01	9.4E−01	96.62	*1.8E−03*	0.29
t2.14 Wrist flexion	−6.92	−0.32	6.0E−02	−2.69	−0.40	*2.3E−04*	−0.12	−0.10	6.3E−01	−0.18	−0.09	4.5E−01	0.57	0.43	*3.2E−02*	145.25	*1.2E−04*	0.27
t2.15 Fingers V MCP flexion	−1.50	−0.06	7.7E−01	−0.88	−0.11	3.6E−01	0.15	0.11	6.4E−01	−0.05	−0.02	8.7E−01	0.03	0.02	9.3E−01	98.48	*4.9E−02*	0.02
t2.16 Fingers V MCP extension	−2.54	−0.08	6.7E−01	0.53	0.06	6.4E−01	−0.56	−0.32	1.6E−01	0.41	0.13	3.0E−01	0.68	0.35	1.1E−01	113.74	5.6E−02	0.06
t2.17 Trunk flexion	8.19	0.43	*1.4E−02*	1.83	0.31	*4.0E−03*	−0.03	−0.03	8.8E−01	0.52	0.28	*1.6E−02*	−0.18	−0.16	4.2E−01	−4.52	8.9E−01	0.25
t2.18 Trunk extension	−1.98	−0.11	5.6E−01	−0.18	−0.03	7.9E−01	0.29	0.30	2.0E−01	−0.17	−0.10	4.5E−01	−0.23	−0.21	3.4E−01	−0.17	1.0E + 00	0.03
t2.19 Trunk rotation	13.78	0.64	*1.9E−04*	1.66	0.25	*1.5E−02*	0.03	0.03	8.9E−01	0.62	0.29	*9.4E−03*	−0.36	−0.27	1.6E−01	4.34	9.0E−01	0.33
t2.20 Trunk lateral bending	3.03	0.33	7.8E−02	−0.11	−0.04	7.3E−01	0.00	0.01	9.7E−01	0.04	0.04	7.3E−01	0.05	0.09	6.7E−01	20.00	2.3E−01	0.15
t2.21 Hip flexion	−6.13	−0.46	*1.7E−02*	0.46	0.11	3.4E−01	0.20	0.27	2.2E−01	−0.10	−0.07	5.6E−01	−0.08	−0.10	6.4E−01	94.77	*2.6E−04*	0.11
t2.22 Hip extension	3.34	0.43	*2.1E−02*	0.41	0.17	1.3E−01	−0.03	−0.07	7.5E−01	0.20	0.27	*3.2E−02*	−0.04	−0.09	6.6E−01	9.24	5.1E−01	0.16
t2.23 Hip abduction	−2.88	−0.30	1.1E−01	−0.46	−0.16	1.8E−01	−0.07	−0.12	5.8E−01	−0.21	−0.22	7.4E−02	0.09	0.16	4.6E−01	55.00	*2.4E−03*	0.13
t2.24 Hip adduction	−4.68	−0.37	*4.6E−02*	0.13	0.03	7.6E−01	−0.04	−0.06	7.8E−01	0.16	0.13	3.0E−01	0.47	0.61	*4.9E−03*	−4.57	8.4E−01	0.16
t2.25 Hip external rotation	10.18	0.61	*4.4E−04*	0.83	0.16	1.2E−01	0.03	0.03	8.6E−01	0.65	0.39	*7.1E−04*	−0.19	−0.19	3.3E−01	11.53	6.7E−01	0.31
t2.26 Hip internal rotation	−11.45	−0.60	*4.4E−04*	0.45	0.08	4.5E−01	−0.03	−0.03	8.7E−01	−0.15	−0.08	4.8E−01	0.02	0.02	9.3E−01	46.37	1.3E−01	0.33
t2.27 Knee flexion	−3.51	−0.30	1.2E−01	0.48	0.13	2.6E−01	0.26	0.41	7.3E−02	−0.12	−0.11	4.0E−01	−0.22	−0.31	1.6E−01	109.06	*4.0E−06*	0.09
t2.28 Knee extension	−3.09	−0.32	8.2E−02	−0.38	−0.13	2.6E−01	−0.14	−0.27	2.1E−01	−0.06	−0.06	6.3E−01	0.14	0.24	2.5E−01	31.16	7.3E−02	0.16
t2.29 Ankle dorsi flexion	0.02	0.00	9.9E−01	0.43	0.11	3.6E−01	0.17	0.24	2.9E−01	0.05	0.04	7.5E−01	−0.34	−0.42	5.7E−02	0.27	9.9E−01	0.09
t2.30 Ankle plantar flexion	−3.64	−0.28	1.4E−01	−0.49	−0.12	3.0E−01	0.16	0.22	3.1E−01	0.07	0.05	6.6E−01	−0.17	−0.21	3.4E−01	43.67	7.5E−02	0.12

t2.31 *P* values less than 0.05 are shown in italics

t2.32 *B* partial regression coefficient, *β* standardized partial regression coefficient

UNCORRECTED PROOF

t3.1 **Table 3** Subtraction of the dominant from the non-dominant side, ROM(ND) – ROM(D) [°]

t3.2 Body part	ROM	All			Female			Male		
t3.3		Mean	SD	*P*	Mean	SD	*P*	Mean	SD	*P*
t3.4 Upper	Shoulder flexion	1.0	5.0	1.0E−01	0.3	2.7	5.4E−01	1.6	6.5	1.4E−01
t3.5	Shoulder extension	0.5	6.9	5.5E−01	0.6	6.5	6.0E−01	0.4	7.3	7.4E−01
t3.6	Shoulder abduction	0.3	1.4	1.0E−01	0.4	1.9	1.8E−01	0.1	0.8	3.2E−01
t3.7	Shoulder external rotation	−1.6	6.3	*3.2E−02*	−1.3	7.0	2.8E−01	−1.9	5.6	*4.6E−02*
t3.8	Shoulder internal rotation	4.3	17.0	*3.5E−02*	5.1	15.9	6.8E−02	3.6	18.1	2.4E−01
t3.9	Shoulder horizontal flexion	−0.5	9.0	6.5E−01	−0.9	7.7	5.1E−01	−0.1	10.1	9.4E−01
t3.10	Shoulder horizontal extension	−0.5	9.8	6.7E−01	−0.3	9.5	8.6E−01	−0.7	10.1	6.9E−01
t3.11	Elbow flexion	1.1	5.6	9.2E−02	1.2	4.9	1.7E−01	1.1	6.1	2.9E−01
t3.12	Elbow extension	0.2	2.6	5.0E−01	−0.3	3.0	5.7E−01	0.7	2.1	5.8E−02
t3.13	Wrist extension	1.2	8.2	2.2E−01	0.4	5.7	6.5E−01	1.9	10.0	2.6E−01
t3.14	Wrist flexion	−2.0	7.7	*3.4E−02*	−1.0	7.8	4.4E−01	−2.8	7.6	*2.9E−02*
t3.15	Fingers V MCP flexion	0.7	10.3	5.6E−01	2.2	9.4	1.8E−01	−0.7	10.9	7.1E−01
t3.16	Fingers V MCP extension	0.6	10.7	6.5E−01	0.6	11.3	7.8E−01	0.6	10.2	7.3E−01
t3.17 Trunk	Trunk rotation	0.3	6.6	7.2E−01	−1.0	5.5	2.9E−01	1.4	7.4	2.4E−01
t3.18	Trunk lateral bending	0.3	4.1	4.9E−01	0.1	4.3	8.4E−01	0.5	4.1	4.4E−01
t3.19 Lower	Hip flexion	−0.5	7.4	5.8E−01	−1.0	7.7	4.4E−01	0.0	7.2	1.0E + 00
t3.20	Hip extension	0.8	3.5	5.1E−02	0.6	3.2	2.9E−01	1.0	3.8	1.0E−01
t3.21	Hip abduction	3.5	8.1	*3.7E−04*	4.3	9.2	*9.7E−03*	2.8	6.9	*1.6E−02*
t3.22	Hip adduction	−3.2	10.6	*1.1E−02*	−2.0	5.3	*3.3E−02*	−4.3	13.7	5.7E−02
t3.23	Hip external rotation	2.0	9.8	8.0E−02	3.4	8.4	*2.1E−02*	0.8	10.8	6.6E−01
t3.24	Hip internal rotation	−0.5	8.7	6.0E−01	−1.4	9.9	4.0E−01	0.3	7.5	8.3E−01
t3.25	Knee flexion	0.7	6.1	3.5E−01	−1.0	6.0	3.3E−01	2.1	5.9	*2.8E−02*
t3.26	Knee extension	0.3	2.5	3.6E−01	0.4	2.8	4.0E−01	0.2	2.3	6.8E−01
t3.27	Ankle dorsi flexion	1.5	7.1	7.8E−02	2.0	5.8	5.1E−02	1.0	8.1	4.4E−01
t3.28	Ankle plantar flexion	2.1	7.0	*1.2E−02*	1.7	6.2	1.1E−01	2.4	7.7	5.8E−02

t3.29 Non-dominant (ND) and dominant (D) sides were determined by hand/arm for upper body and by foot/leg for lower body. *P* values less than 0.05 are shown in italics

t4.1 **Table 4** Subtraction of the dominant from the non-dominant side in the total range of antagonistic motions, ROM(ND) – ROM(D) [°]

t4.2 Body part	Antagonistic motions	All			Female			Male		
t4.3		Mean	SD	*P*	Mean	SD	*P*	Mean	SD	*P*
t4.4 Upper	Shoulder flx + ext	1.5	9.3	1.8E−01	0.9	6.8	4.5E−01	2.0	11.1	2.8E−01
t4.5	Shoulder abd + add	0.3	1.4	1.0E−01	0.4	1.9	1.8E−01	0.1	0.8	3.2E−01
t4.6	Shoulder int rot + ext rot	2.7	18.1	2.1E−01	3.8	15.2	1.5E−01	1.7	20.6	6.2E−01
t4.7	Shoulder hori flx + hori ext	−1.0	12.6	5.1E−01	−1.2	13.5	6.1E−01	−0.8	11.9	6.8E−01
t4.8	Elbow flx + ext	1.3	6.0	6.3E−02	0.9	6.3	4.2E−01	1.8	5.7	6.8E−02
t4.9	Wrist flx + ext	−0.8	10.3	5.3E−01	−0.6	8.2	6.8E−01	−0.9	12.0	6.3E−01
t4.10	Fingers V MCP flx + ext	1.3	15.1	4.8E−01	2.8	15.2	3.0E−01	−0.1	15.0	9.7E−01
t4.11 Lower	Hip flx + ext	0.8	7.7	1.8E−01	0.3	8.1	8.3E−01	1.3	7.5	1.1E−01
t4.12	Hip abd + add	0.3	13.7	9.0E−01	2.0	11.0	3.1E−01	−1.2	15.7	6.1E−01
t4.13	Hip int rot + ext rot	2.3	12.0	8.3E−02	2.6	10.8	1.8E−01	2.1	13.0	2.5E−01
t4.14	Knee flx + ext	0.6	6.7	2.6E−01	−0.8	7.5	5.5E−01	1.9	5.8	*1.7E−02*
t4.15	Ankle dorsi flx + plantar flx	3.5	8.3	*2.9E−04*	3.8	8.0	*1.1E−02*	3.2	8.7	*1.1E−02*

t4.16 Non-dominant (ND) and dominant (D) sides were determined by hand/arm for upper body and by foot/leg for lower body. *P* values less than 0.05 are shown in italics
t4.17 *flx* flexion, *ext* extension, *abd* abduction, *add* adduction, *int rot* internal rotation, *ext rot* external rotation, *hori flx* horizontal flexion, *hori ext* horizontal extension,
t4.18 *dorsi flx* dorsi flexion, *plantar flx* plantar flexion

UNCORRECTED PROOF



f1.1 **Fig. 1** Principal component loadings of the first three PCs. **a** For all samples (females and males). **b** For female samples. **c** For male samples.
f1.2 The average of right and left ROMs was used when the joint motion had right and left data



f2.1 **Fig. 2** Comparison of PC1 in three PCAs using all samples, only female samples, and only male samples. **a** Principal component loadings. The
f2.2 joint motions are sorted in the order of loadings in $PC1_{all}$. *Black bars* indicate motions in which females have a significantly larger ROM than
f2.3 males. *Shaded bars* indicate motions in which males have a significantly larger ROM than females. **b** Correlations between $PC1_{all}$ and $PC1_{females}$
f2.4 and between $PC1_{all}$ and $PC1_{males}$

whereas different patterns were observed in PC2 and PC3. There were strong and significant correlations in the loadings between $PC1_{all}$ and $PC1_{females}$ ($r = 0.75$) and between $PC1_{all}$ and $PC1_{males}$ ($r = 0.96$) (Fig. 2b). Extremely positive values of loadings were observed for wrist extension, shoulder horizontal flexion, and elbow extension, and extremely negative values were observed for trunk rotation, hip external rotation, trunk flexion,

and hip extension in $PC1_{all}$. It is notable that the motions in which females have higher mobility than males showed positive values (black bars in Fig. 2a), whereas the motions in which females have lower mobility than males showed negative values (shaded bars in Fig. 2a). These results indicated that $PC1_{all}$ is a component representing the sex difference, while this component is also observed within each sex.

Factors associated with the whole-body ROM patterns

T5 Table 5 shows correlation coefficients and partial correlation coefficients controlling for sex between PCs and the somatometric/sthenometric measurements. $PC1_{all}$ was significantly correlated with age, BF%, iliospinale height, and leg extension strength after controlling for sex. $PC2_{all}$ was also associated with sex, and significant partial correlations with age, forearm circumference, grip strength, leg extension strength, and leg flexion strength were observed. $PC3_{all}$ was not affected by sex and had significant correlations with weight and forearm, minimum forearm, calf, and ankle circumferences.

To examine the independency of the effect of each factors and to further narrow down the factors that have a direct effect on the whole-body ROM pattern, we subsequently performed multiple regression analysis. Somatometric/sthenometric measurements that were significantly correlated with PCs as mentioned above were included as explanatory variables, and then the variables were chosen thorough stepwise procedures. As a result, sex, age, BF%, iliospinale height, and leg extension strength were associated with $PC1_{all}$ (Table 6), T6 which indicated that these factors have independent effects. $PC2_{all}$ was associated negatively with being male but positively with age and grip strength. Grip strength was likely to represent the whole-body muscle strength since it was correlated with removed factors, leg extension, and flexion strength. The analysis for $PC3_{all}$ suggested that the effects of weight and limb circumferences are not independent of each other, and forearm circumference could best explain the $PC3_{all}$ scores.

Discussion

The results for the sex difference for each ROM (Table 2) were mostly consistent with previous studies; the majority of limb joints had a larger ROM in females than in males, while males were more flexible than females in only four joint motions, including trunk flexion, trunk rotation, hip extension, and hip external rotation [14, 15, 18, 19]. The present study showed that age has negative correlations only with several joint motions. However, since ages of subjects were concentrated in early twenties, careful interpretation should be required for the effect of age. Because most individuals had a sports experience when they were high school students, the period of time after they ceased exercise may have an influence on ROM. Previous observations of a broader range of age groups have disclosed negative effects of age-related changes on the ROM patterns [2, 5, 8, 19]. A previous study of older

t5.1 **Table 5** Correlations of PCs with somatometric/sthenometric measurements

t5.2 Measurement	$PC1_{all}$				$PC2_{all}$				$PC3_{all}$			
t5.3	*R*	*P*	*R**	*P*	*R*	*P*	*R**	*P*	*R*	*P*	*R**	*P*
t5.4 Sex	−0.51	*2.6E−06*	–	–	−0.45	*5.4E−05*	–	–	−0.16	1.7E−01	–	–
t5.5 Age	−0.64	*4.5E−10*	−0.52	*2.7E−06*	0.23	*4.6E−02*	0.32	*7.4E−03*	0.07	5.7E−01	−0.01	9.1E−01
t5.6 Height	−0.31	*6.0E−03*	0.23	5.8E−02	−0.31	*5.7E−03*	0.09	4.4E−01	−0.07	5.6E−01	0.04	7.6E−01
t5.7 Weight	−0.36	*1.5E−03*	0.02	9.0E−01	−0.35	*1.8E−03*	0.12	3.1E−01	−0.23	*4.9E−02*	−0.14	2.5E−01
t5.8 Body fat percentage	−0.10	3.7E−01	−0.48	*1.9E−05*	0.31	*6.4E−03*	0.15	2.1E−01	−0.15	2.0E−01	−0.12	3.2E−01
t5.9 Lean body mass	−0.27	*2.0E−02*	0.23	5.6E−02	−0.40	*3.2E−04*	0.06	6.2E−01	−0.14	2.2E−01	−0.08	4.9E−01
t5.10 Upper limb length	−0.35	*1.8E−03*	0.12	3.1E−01	−0.28	*1.5E−02*	0.12	3.3E−01	−0.04	7.2E−01	0.04	7.2E−01
t5.11 Iliospinale height	−0.17	1.3E−01	0.36	*2.3E−03*	−0.32	*4.2E−03*	−0.05	6.8E−01	−0.08	5.0E−01	0.02	8.9E−01
t5.12 Forearm circumference	−0.48	*1.2E−05*	−0.16	1.9E−01	−0.26	*2.2E−02*	0.28	*1.8E−02*	−0.24	*3.5E−02*	−0.17	1.6E−01
t5.13 Minimum forearm circumference	−0.43	*1.0E−04*	−0.08	4.9E−01	−0.29	*1.1E−02*	0.15	2.0E−01	−0.24	*4.1E−02*	−0.10	4.1E−01
t5.14 Calf circumference	−0.37	*9.6E−04*	−0.15	2.1E−01	−0.23	5.2E−02	0.17	1.5E−01	−0.27	*1.9E−02*	−0.17	1.5E−01
t5.15 Ankle circumference	−0.14	2.3E−01	0.19	1.2E−01	−0.34	*2.7E−03*	−0.01	9.7E−01	−0.26	*2.6E−02*	−0.13	2.9E−01
t5.16 Grip strength	−0.34	*2.8E−03*	0.14	2.4E−01	−0.15	2.0E−01	0.40	*5.4E−04*	−0.14	2.4E−01	−0.01	9.6E−01
t5.17 Leg extension strength	−0.56	*2.1E−07*	−0.36	*1.8E−03*	−0.15	1.9E−01	0.31	*8.7E−03*	−0.20	9.2E−02	−0.13	2.8E−01
t5.18 Leg flexion strength	−0.44	*8.7E−05*	−0.10	4.0E−01	−0.26	*2.6E−02*	0.24	*4.8E−02*	−0.12	3.3E−01	0.11	3.8E−01
t5.19 2D:4D ratio	−0.04	7.5E−01	−0.07	5.8E−01	−0.04	7.4E−01	0.01	9.5E−01	−0.10	3.7E−01	0.02	8.4E−01

t5.20 *P* values less than 0.05 are shown in italics
t5.21 *R* correlation coefficient, *R** partial correlation coefficient controlling for sex

UNCORRECTED PROOF

Q5 **Table 6** Multiple regression analysis for identifying factors that explain PCs

PC	Explanatory variables	*B*	*β*	*P*	Eliminated variables
$PC1_{all}$	Sex (F:0, M:1)	−2.61	−0.60	*2.8E−06*	
	Age	−0.45	−0.28	*2.1E−03*	
	Body fat percentage	−0.14	−0.31	*5.4E−04*	
	Iliospinale height	0.13	0.35	*1.9E−03*	
	Leg extension strength	−0.02	−0.29	*2.0E−02*	
$PC2_{all}$	Sex (F:0, M:1)	−3.20	−0.90	*3.7E−07*	Forearm circumference
	Age	0.35	0.27	*7.7E−03*	Leg extension strength
	Grip strength	0.11	0.58	*5.9E−04*	Leg flexion strength
$PC3_{all}$	Forearm circumference	−0.19	−0.27	1.8E−02	Weight
					Minimum forearm circumference
					Calf circumference
					Ankle circumference

B partial regression coefficient, *β* standardized partial regression coefficient

adults showed that shoulder abduction and hip flexion are associated negatively with age and positively with muscular strength [26]; this may reflect that changes in physical activity due to ageing strongly affect both joint flexibility and muscular strength.

It is worth noting that the BF% and LBM showed not only negative effects on some joint motions but also positive effects on other joint motions. Negative correlations between BF% and several joint motions are likely due to physical obstruction by fat tissue caught between the bones constituting the joint. Shoulder horizontal flexion is a clear example of limitation by fat tissue (Table 2). The results of the multiple regression analyses indicated that the BF% contributes to the limitation of ROM in the upper limb, whereas it increases ROM in trunk flexion and rotation and hip external rotation. Causes of the positive correlation between BF% and ROM need to be further investigated. In the case of shoulder external rotation and horizontal flexion, effects of physical obstruction by muscles and the skeleton can explain the negative associations with LBM. On the other hand, the positive associations of LBM with wrist flexion and hip adduction can result from an indirect association; daily exercise may increase the flexibility of wrist and hip joints, as well as LBM.

Regarding hip joints, females were more flexible than males in flexion, adduction, and internal rotation, and vice versa in extension and external rotation (Table 2). However, no significant sex difference in the total range of antagonistic motions, such as flexion versus extension and external rotation versus internal rotation, suggests that each ROM of the hip joints is affected by skeletal morphology that determines relative positions and angles between bones. Sexual dimorphisms in anteversion of the acetabulum and femoral neck are well known; acetabular anteversion is defined as a forward tilt of the acetabular opening plane with respect to the sagittal plane, and femoral neck anteversion is defined as anterior rotation of femoral neck compared to the axis of the femoral condyles. In general, females have larger femoral neck anteversion than males, which is considered to be a reason for larger hip internal rotation and smaller hip external rotation in females than males [27, 28]. In addition, Nakahara et al. [29] found that a larger acetabulum anteversion in females than in males causes larger hip flexion and hip internal rotation, whereas males have larger ROMs than females in the antagonistic motions that are hip extension and hip external rotation.

As for the trunk, males had a greater ROM of flexion and rotation than females. Females generally have a shorter spinal column and a larger lumbar lordosis than males [30], which is considered to be a reason for females' smaller trunk flexion and rotation. A kinematic analysis of rising from a chair reported that lumbar spine flexion occurs concurrently with hip flexion [31]; this suggests that lumbar spine flexion compensates for inflexibility of the hip joint motion in males.

The data on differences between dominant and non-dominant sides also provide information on factors affecting the variations in ROM. Joint motions that had a larger ROM on the dominant side than on the non-dominant side were shoulder external rotation, wrist flexion, and hip adduction. This result suggests an involvement of daily activity in the variation in ROM. Regarding the asymmetry of shoulder joints, it has been reported that the side of the dominant hand/arm has a significantly larger ROM than the other side, especially among individuals who have experience in sports with overhead-throwing motion [32, 33]. In the present study, we reanalysed only males who had experience in overhead-throwing motion sports, and confirmed an increased difference between the sides in shoulder external rotation ($n = 19$, ROM(ND) – ROMD) = −2.4 ± 4.2, $P = 0.0245$). On the other hand, some joint motions showed a larger ROM on the non-dominant side than on the dominant side. Of these motions, shoulder internal rotation and hip abduction are antagonistic movements of shoulder external rotation and hip adduction, respectively, that showed larger motion on the dominant side. These side differences may be due to off-centred neutral posture because the total ranges of antagonistic motions had no significant difference between the dominant and non-dominant sides. It is well known that side dominance causes asymmetry of posture. In addition,

UNCORRECTED PROOF

previous studies have reported that asymmetric daily posture, such as side sitting, can be related to ROM asymmetry [13, 34]. Alternatively, the difference between dominant and non-dominant sides may be attributed to muscle mass or extension of muscles and tendons; a forced and continuous motion on the dominant side increases its ROM by stretching the muscles antagonistic to the motion. In contrast, the reverse motion is limited by the developed muscles being an obstacle.

As for the relationships in ROM among different joints, Allander et al. [10] reported significant correlations among shoulder, wrist, metacarpophalangeal joint I (MCP I), and hip; in particular, wrist mobility was related to the mobility of the other three joints. However, no study has analysed the covariation patterns of whole-body ROM. On our PCA, $PC1_{all}$ was associated with sex differences, and even when females and males were separately analysed, similar covariation patterns appeared as PC1. These results indicate that not only sexual dimorphism but also other factors, such as body fat, lower limb length, and muscle mass, can be involved in the component. In addition, our study also showed that $PC2_{all}$ and $PC3_{all}$ were significantly associated with muscle strength and limb circumference, respectively. This also indicated that body composition affects the whole-body patterns of joint flexibility.

Based on the results of PCA, we refer to the positive direction of $PC1_{all}$ as "feminine type", and the negative direction as "masculine type" (Fig. 3). Feminine type is characterized by high flexibility of the upper limbs, such as wrist extension and flexion, shoulder horizontal flexion, and elbow extension, while masculine type is characterized by high flexibility of trunk flexion, trunk rotation, hip extension, and hip external rotation. On regression analysis, sex, age, BF%, iliospinale height, and leg extension strength were associated with $PC1_{all}$. BF% had a negative association with the $PC1_{all}$ score, which means that an increased BF% is related to masculine type. Golden et al. [35] have also suggested that an increase of BMI is correlated with a decrease of ROM and that a decreased amount of daily activity leads to both an increased BMI and decreased ROM in the whole body. Iliospinale height is an index of limb length; thus, the positive correlation between iliospinale height and $PC1_{all}$ suggests that the longer the limbs are, the higher the tendency for feminine type the individual has. Leg extension strength, being an index of muscle mass, was associated with a tendency to have masculine type.



Fig. 3 A schematic representation of the whole-body ROM patterns. Being flexible in some motions is characteristic of two contrary types, masculine type (*left*) and feminine type (*right*)

Furthermore, the covariation pattern of $PC1_{all}$ should be strongly affected by acetabulum and femoral neck anteversion because antagonistic motions of hip joints, and hip abduction and adduction, were associated with sex oppositely. In a previous study, it was also reported that an increase in femoral neck anteversion contributed to a decrease in muscle strength of the gluteus medius and vastus medialis [36]. Therefore, the present study suggests that acetabular and femoral neck anteversion and muscle strength, being related to each other in a complicated manner, have an influence on hip joint ROM.

As shown above, multiple factors are likely to be associated with ROM and whole-body patterns of ROM. To understand how cultural differences affect ROM, further global comparisons will be indispensable [3, 10, 37]. In addition, genetic factors associated with joint flexibility still remain to be elucidated. A twin study has reported that the heritability for lumbar flexion is 64 % [38]. It has also been reported that the levels of femoral anteversion are highly correlated between siblings, indicating that this trait is partially heritable [39]. Therefore, the whole-body patterns of joint motions need to be further studied from the various perspectives, including genetic and environmental factors.

Our present study, clarifying the covariation patterns of joint flexibility, will contribute to the prevention of joint injuries and to the evaluation of dysfunction in patients with musculoskeletal diseases. For example, it has been known that anterior cruciate ligament injuries are more frequent in females than in males partly because of joint laxity [40], and therefore, it is possible that the "feminine type" has a higher susceptibility of the knee joint injury than the "masculine type" when they are compared within each sex. Further studies are needed to provide prevention and therapy programmes in consideration of the patterns of joint motions. In addition, it would be essential to know one's type of joint flexibility and one's suitable and effective postures and motions in order to improve the performance in sports and daily activities.

Conclusion

A covariation pattern of ROMs that shows sexual dimorphism was found by PCA. Such covariation pattern was also observed within each sex as a spectrum of "masculine" and "feminine" types and was shown to be partly associated with body proportion and composition and with muscle mass and strength. Comparisons between dominant and non-dominant sides and between antagonistic motions provide suggestions that ordinary posture, daily motions, and skeletal morphology such as acetabular and femoral neck anteversion contribute to individual differences in ROMs. Such knowledge will contribute to the prevention of joint injuries and to improve one's performance in sports and daily activities.

Additional file

Additional file 1: Table S1. Age and hand/arm and foot/leg dominances of the subjects. Table S2. Somatometric and sthenometric measurements in the subjects. (DOCX 23 kb)

Abbreviations

2D:4D: The ratio of the second and fourth finger lengths; AAOS: American Academy of Orthopaedic Surgeons; BF%: Body fat percentage; LBM: Lean body mass; MCP I: Metacarpophalangeal joint I; PCA: Principal component analysis; ROM: Range of motion

Acknowledgements

The authors wish to express their profound gratitude to the volunteers for participating in this study. We are also grateful for the help given by Dr. Akira Kawaguchi.

Funding

This study was supported in part by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (Nos. 22370087, 25251042).

Authors' contributions

KM participated in the design of the study, performed physical measurements and the statistical analysis, and drafted the manuscript. RK conceived of the study, participated in the design of the study, and drafted the manuscript. HF contributed to the interpretation of the statistical analysis and helped to draft the manuscript. KY contributed to the statistical analysis and helped to draft the manuscript. HI conceived of the study, participated in the design of the study, and drafted the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

[1]Department of Human Biology and Anatomy, Graduate School of Medicine, University of the Ryukyus, Uehara 207, Nishihara, Okinawa 903-0215, Japan. [2]Division of Human Evolution Studies, Graduate School of Medicine, Hokkaido University, Kita 15 Nishi 7, Kita-ku, Sapporo, Hokkaido 060-8638, Japan. [3]Present address: School of Natural Sciences and Psychology, Liverpool John Moores University, Byrom Street, Liverpool L3 3AF, UK.

Received: 16 May 2016 Accepted: 15 September 2016

References

1. American Academy of Orthopaedic Surgeons. Joint motion: method of measuring and recording. Chicago, IL: American Academy of Orthopaedic Surgeons; 1965.
2. Chapleau J, Canet F, Petit Y, Sandman E, Laflamme GY, Rouleau DM. Demographic and anthropometric factors affecting elbow range of motion in healthy adults. J Shoulder Elbow Surg. 2013;22(1):88–93.
3. Gunal I, Kose N, Erdogan O, Gokturk E, Seber S. Normal range of motion of the joints of the upper extremity in male subjects, with special reference to side. J Bone Joint Surg Am. 1996;78(9):1401–4.
4. Gupta A, Fernihough B, Bailey G, Bombeck P, Clarke A, Hopper D. An evaluation of differences in hip external rotation strength and range of motion between female dancers and non-dancers. Br J Sports Med. 2004;38(6):778–83.
5. James B, Parker AW. Active and passive mobility of lower limb joints in elderly men and women. Am J Phys Med Rehabil. 1989;68(4):162–7.
6. Leszko F, Hovinga KR, Lerner AL, Komistek RD, Mahfouz MR. In vivo normal knee kinematics: is ethnicity or gender an influencing factor? Clin Orthop Relat Res. 2011;469(1):95–106.
7. Nonaka H, Mita K, Watakabe M, Akataki K, Suzuki N, Okuwa T, et al. Age-related changes in the interactive mobility of the hip and knee joints: a geometrical analysis. Gait Posture. 2002;15(3):236–43.
8. Roach KE, Miles TP. Normal hip and knee active range of motion: the relationship to age. Phys Ther. 1991;71(9):656–65.

UNCORRECTED PROOF

9. Steultjens MP, Dekker J, van Baar ME, Oostendorp RA, Bijlsma JW. Range of joint motion and disability in patients with osteoarthritis of the knee or hip. Rheumatology. 2000;39(9):955–61.
10. Allander E, Bjornsson OJ, Olafsson O, Sigfusson N, Thorsteinsson J. Normal range of joint movements in shoulder, hip, wrist and thumb with special reference to side: a comparison between two populations. Int J Epidemiol. 1974;3(3):253–61.
11. Boone DC, Azen SP. Normal range of motion of joints in male subjects. J Bone Joint Surg Am. 1979;61(5):756–9.
12. Macedo LG, Magee DJ. Differences in range of motion between dominant and nondominant sides of upper and lower extremities. J Manipulative Physiol Ther. 2008;31(8):577–82.
13. Macedo LG, Magee DJ. Effects of age on passive range of motion of selected peripheral joints in healthy adult females. Physiother Theory Pract. 2009;25(2):145–64.
14. Okabe T, Watanabe H, Amano T. The range of joint motions of the extremities in healthy Japanese people—the differences according to the sex. Sogo Rehabilitation. 1980;8(1):41–56 [in Japanese].
15. Soucie JM, Wang C, Forsyth A, Funk S, Denny M, Roach KE, et al. Range of motion measurements: reference values and a database for comparison studies. Hemophilia. 2011;17(3):500–7.
16. Takemasa S, Shimada T, Hidaka M. Normal range of motion of joints in the aged people. Bull Health Sci Kobe. 1997;13:77–82 [in Japanese].
17. Walker JM, Sue D, Miles-Elkousy N, Ford G, Trevelyan H. Active mobility of the extremities in older subjects. Phys Ther. 1984;64(6):919–23.
18. Watanabe H, Ogata K, Amano T, Okabe T. The range of joint motions of the extremities in healthy Japanese people: the difference according to the age. Nihon Seikeigeka Gakkai Zasshi. 1979;53(3):275–91 [in Japanese].
19. Chung MJ, Wang MJ. The effect of age and gender on joint range of motion of worker population in Taiwan. Int J Ind Ergon. 2009;39(4):596–600.
20. Chapman JP, Chapman LJ, Allen JJ. The measurement of foot preference. Neuropsychologia. 1987;25(3):579–84.
21. Hatta T, Nakatsuka Z. Hatta-Nakatsuka handedness test. In Ohno, S. (Ed.) Festschrift to Prof. Ohnishi. Osaka, Japan: Osaka City University Press; 1974 [in Japanese].
22. Knussmann R. Anthropologie: Handbuch der vergleichenden Biologie des Menschen, vol. I. Stuttgart, Germany: Gustav Fischer Verlag; 1988.
23. Brozek J, Kihlberg JK, Taylor HL, Keys A. Skinfold distributions in middle-aged american men: a contribution to norms of leanness-fatness. Ann N Y Acad Sci. 1963;110:492–502.
24. Nagamine S, Suzuki S. Anthropometry and body composition of Japanese young men and women. Hum Biol. 1964;36(1):8–15.
25. Nagamine S. Obesity and weight loss methods. In: Nagamine S, editor. Contemporary sports science: Sport, Energy and Nutrition. Tokyo, Japan: Taishukan; 1979. p. 259–83 [in Japanese].
26. Stathokostas L, McDonald MW, Little RM, Paterson DH. Flexibility of older adults aged 55–86 years and the influence of physical activity. J Aging Res. 2013;743–843
27. Braten M, Terjesen T, Rossvoll I. Femoral anteversion in normal adults: ultrasound measurements in 50 men and 50 women. Acta Orthop Scand. 1992;63(1):29–32.
28. Kozic S, Gulan G, Matovinovic D, Nemec B, Sestan B, Ravlic-Gulan J. Femoral anteversion related to side differences in hip rotation: passive rotation in 1,140 children aged 8-9 years. Acta Orthop Scand. 1997;68(6):533–6.
29. Nakahara I, Takao M, Sakai T, Nishii T, Yoshikawa H, Sugano N. Gender differences in 3D morphology and bony impingement of human hips. J Orthop Res. 2011;29(3):333–9.
30. Amonoo-Kuofi HS. Changes in the lumbosacral angle, sacral inclination and the curvature of the lumbar spine during aging. Acta Anat. 1992;145(4):373–7.
31. Fotoohabadi MR, Tully EA, Galea MP. Kinematics of rising from a chair: image-based analysis of the sagittal hip-spine movement pattern in elderly people who are healthy. Phys Ther. 2010;90(4):561–71.
32. Anloague PA, Spees V, Smith J, Herbenick MA, Rubino LJ. Glenohumeral range of motion and lower extremity flexibility in collegiate-level baseball players. Sports Health. 2012;4(1):25–30.
33. Ruotolo C, Price E, Panchal A. Loss of total arc of motion in collegiate baseball players. J Shoulder Elbow Surg. 2006;15(1):67–71.
34. Wescott DJ, Cunningham DL, Hunt DR. Temporal trends in femoral diaphyseal torsional asymmetry among the Arikara associated with postural behavior. Am J Phys Anthropol. 2014;154(4):512–24.
35. Golden DW, Wojcicki JM, Jhee JT, Gilpin SL, Sawyer JR, Heyman MB. Body mass index and elbow range of motion in a healthy pediatric population: a possible mechanism of overweight in children. J Pediatr Gastroenterol Nutr. 2008;46(2):196–201.
36. Nyland J, Kuzemchek S, Parks M, Caborn DN. Femoral anteversion influences vastus medialis and gluteus medius EMG amplitude: composite hip abductor EMG amplitude ratios during isometric combined hip abduction-external rotation. J Electromyogr Kinesiol. 2004;14(2):255–61.
37. Trudelle-Jackson E, Fleisher LA, Borman N, Morrow Jr JR, Frierson GM. Lumbar spine flexion and extension extremes of motion in women of different age and racial groups: the WIN Study. Spine. 2010;35(16):1539–44.
38. Battie MC, Levalahti E, Videman T, Burton K, Kaprio J. Heritability of lumbar flexibility and the role of disc degeneration and body weight. J Appl Physiol. 2008;104(2):379–85.
39. Upadhyay SS, Burwell RG, Moulton A, Small PG, Wallace WA. Femoral anteversion in healthy children: application of a new method using ultrasound. J Anat. 1990;169:49–61.
40. Arendt E, Dick R. Knee injury patterns among men and women in collegiate basketball and soccer: NCAA data and review of literature. Am J Sports Med. 1995;23(6):694–701.

UNCORRECTED PROOF

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

Author Query Form

Journal: Journal of Physiological Anthropology

Title: Whole-body patterns of the range of joint motion in young adults: masculine type and feminine type

Authors: Keiichi Moromizato, Ryosuke Kimura, Hitoshi Fukase, Kyoko Yamaguchi, Hajime Ishida

Article: 112

Dear Authors,

During production of your paper, the following queries arose. Please respond to these by annotating your proofs with the necessary changes/additions. If you intend to annotate your proof electronically, please refer to the E-annotation guidelines. We recommend that you provide additional clarification of answers to queries by entering your answers on the query sheet, in addition to the text mark-up.

Query No.	Query	Remark
Q1	Author names: Please confirm that the author names are presented accurately and in the correct sequence (given names/initials, family name). Author 1: Given name: Keiichi Family name:Moromizato Author 2: Given name: Ryosuke Family name:Kimura Author 3: Given name: Hitoshi Family name:Fukase Author 4: Given name: Kyoko Family name:Yamaguchi Author 5: Given name: Hajime Family name:Ishida	
Q2	Additional file: As a rule, all supplementary files are to be referred as additional files. Thus, "Supplementary Table S1 and Table S2" was changed to "Additional file 1: Table S1, Additional file 1: Table S2". Moreover, titles inside the additional files were also amended to correspond with their modified citations. Please check and advise if action taken is appropriate.	
Q3	Please check if the edit to the sentence “The female sex significantly increased…” is appropriate. Otherwise, kindly amend as necessary.	
Q4	Please check if the edit to the sentence “When PCs resulting from female and… retains the intended meaning.”	
Q5	Please indicate the significance of the italicized data in Table 6 in the form of a table footnote.	

Query No.	Query	Remark
Q6	Additional file: As per journal requirements, every additional file must have a corresponding caption. In this regard, please be informed that the caption was taken from the additional e-file itself. Please advise if action taken appropriate and amend if necessary.	
Q7	Please check if the article title in Reference [14] is presented correctly.	
Q8	Citation details for Reference [26] are incomplete. Please supply the “volume number” of this reference. Otherwise, kindly advise us on how to proceed.	